



A symplectic resolution for the binary tetrahedral group

Manfred Lehn, Christoph Sorger

► To cite this version:

Manfred Lehn, Christoph Sorger. A symplectic resolution for the binary tetrahedral group. 2008.
hal-00331839v1

HAL Id: hal-00331839

<https://hal.science/hal-00331839v1>

Preprint submitted on 17 Oct 2008 (v1), last revised 21 Mar 2010 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A SYMPLECTIC RESOLUTION FOR THE BINARY TETRAHEDRAL GROUP

MANFRED LEHN & CHRISTOPH SORGER

ABSTRACT. We describe an explicit symplectic resolution for the quotient singularity arising from the four-dimensional symplectic representation of the binary tetrahedral group.

Let G be a finite group with a complex symplectic representation V . The symplectic form σ on V descends to a symplectic form $\bar{\sigma}$ on the open regular part of V/G . A proper morphism $f : Y \rightarrow V/G$ is a symplectic resolution if Y is smooth and if $f^*\bar{\sigma}$ extends to a symplectic form on Y . It turns out that symplectic resolutions of quotient singularities are a rare phenomenon. By a theorem of Verbitsky [9], a necessary condition for the existence of a symplectic resolution is that G be generated by symplectic reflections, i.e. by elements whose fix locus on V is a linear subspace of codimension 2. Given an arbitrary complex representation V_0 of a finite group G , we obtain a symplectic representation on $V_0 \oplus V_0^*$, where V_0^* denotes the contragredient representation of V_0 . In this case, Verbitsky's theorem specialises to an earlier theorem of Kaledin [7]: For $V_0 \oplus V_0^*/G$ to admit a symplectic resolution, the action of G on V_0 should be generated by complex reflections, in other words, V_0/G should be smooth. The complex reflection groups have been classified by Shephard and Todd [8], the symplectic reflection groups by Cohen [2]. The list of Shephard and Todd contains as a sublist the finite Coxeter groups.

The question which of these groups $G \subset \mathrm{Sp}(V)$ admits a symplectic resolution for V/G has been solved for the Coxeter groups by Ginzburg and Kaledin [3] and for arbitrary complex reflection groups most recently by Bellamy [1]. His result is as follows:

Theorem 1. (Bellamy) — *If $G \subset \mathrm{GL}(V_0)$ is a finite complex reflection group, then $V_0 \oplus V_0^*/G$ admits a symplectic resolution if and only if (G, V_0) belongs to the following cases:*

Date: 6 October 2008.

1991 *Mathematics Subject Classification.* Primary 14B05; Secondary 14E15, 13P10.

1. (S_n, \mathfrak{h}) , where the symmetric group S_n acts by permutations on the hyperplane $\mathfrak{h} = \{x \in \mathbb{C}^n \mid \sum_i x_i = 0\}$.
2. $((\mathbb{Z}/m)^n \rtimes S_n, \mathbb{C}^n)$, the action being given by multiplication with m -th roots of unity and permutations of the coordinates.
3. (T, S_1) , where S_1 denotes a two-dimensional representation of the binary tetrahedral group T (see below).

However, the technique of Ginzburg, Kaledin and Bellamy does not provide resolutions beyond the statement of existence. Case 1 corresponds to Coxeter groups of type A and Case 2 with $m = 2$ to Coxeter groups of type B . It is well-known that symplectic resolutions of $\mathfrak{h} \oplus \mathfrak{h}^*/S_n$ and $\mathbb{C}^n \oplus \mathbb{C}^n/(\mathbb{Z}/m)^n \rtimes S_n \cong \text{Sym}^n(\mathbb{C}^2/(\mathbb{Z}/m))$ are given as follows:

For a smooth surface Y the Hilbert scheme $\text{Hilb}^n(Y)$ of generalised n -tuples of points on Y provides a crepant resolution $\text{Hilb}^n(Y) \rightarrow \text{Sym}^n(Y)$. Applied to a minimal resolution of the A_{m-1} -singularity \mathbb{C}^2/G , $G \cong \mathbb{Z}/m$, this construction yields a small resolution $\text{Hilb}^n(\widetilde{\mathbb{C}^2/G}) \rightarrow \text{Sym}^n(\widetilde{\mathbb{C}^2/G}) \rightarrow \text{Sym}^n(\mathbb{C}^2/G)$. Similarly, $(\mathfrak{h} \oplus \mathfrak{h}^*)/S_n$ is the fibre over the origin of the barycentric map $\text{Sym}^n(\mathbb{C}^2) \rightarrow \mathbb{C}^2$. Thus $(\mathfrak{h} \oplus \mathfrak{h}^*)/S_n$ is resolved symplectically by the null-fibre of the morphism $\text{Hilb}^n(\mathbb{C}^2) \rightarrow \text{Sym}^n(\mathbb{C}^2) \rightarrow \mathbb{C}^2$.

It is the purpose of this note to describe an explicit symplectic resolution for the binary tetrahedral group.

1. THE BINARY TETRAHEDRAL GROUP

Let $T_0 \subset \text{SO}(3)$ denote the symmetry group of a regular tetrahedron. The preimage of T_0 under the standard homomorphism $\text{SU}(2) \rightarrow \text{SO}(3)$ is the binary tetrahedral group T . As an abstract group, T is the semidirect product of the quaternion group $Q_8 = \{\pm 1, \pm I, \pm J, \pm K\}$ and the cyclic group $\mathbb{Z}/3$. As a subgroup of $\text{SU}(2)$ it is generated by the elements

$$I = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad \text{and} \quad \tau = -\frac{1}{2} \begin{pmatrix} 1+i & -1+i \\ 1+i & 1-i \end{pmatrix}$$

The binary tetrahedral group has 5 irreducible complex representations: A three-dimensional one arising from the quotient $T \rightarrow T_0 \subset \text{SO}_3$, three one-dimensional representations \mathbb{C}_j arising from the quotient $T \rightarrow \mathbb{Z}/3$ with τ acting by $e^{2\pi ij/3}$, and three two-dimensional representations S_0 , S_1 and S_2 . Here S_0 denotes the standard representation of T arising from the embedding $T \subset \text{SU}_2$. This representation is itself symplectic, its quotient S_0/T being the well-known Klein-DuVal singularity of type E_6 . The two other representations can be written as $S_j = S_0 \otimes \mathbb{C}_j$, $j = 1, 2$. They

are dual to each other, and the diagonal action of T on $S_1 \oplus S_2$ provides the embedding of T to Sp_4 that is of interest in our context. It is as this subgroup of Sp_4 that T appears in the list of Shephard and Todd under the label “No. 4”.

Whereas the action of T on S_0 is symplectic, the action of T on S_1 and S_2 is generated by complex reflections of order 3. Overall, there are 8 elements of order 3 in T or rather 4 pairs of inverse elements, forming 2 conjugacy classes. To these correspond 4 lines in S_1 of points with nontrivial isotropy groups. Let $C_1 \subset S_1$ and $C_2 \subset S_2$ denote the union of these lines in each case. Then $C_1 \times S_2$ and $S_1 \times C_2$ are invariant divisors in $S_1 \oplus S_2$. However, the defining equations are invariant only up to a scalar. Consequently, their images W_1 and W_2 in the quotient $Z = S_1 \oplus S_2/T$ are Weil divisors but not Cartier. The reduced singular locus $\mathrm{sing}(Z)$ is irreducible and off the origin a transversal A_2 singularity. It forms one component of the intersection $W_1 \cap W_2$.

For $j = 1, 2$, let $\alpha_j : Z'_j \rightarrow Z$ denote the blow-up along W_j . Next, let W'_j be the reduced singular locus of Z'_j , and let $\beta_j : Z''_j \rightarrow Z'_j$ denote the blow-up along W'_j .

Theorem 2. — *The morphisms $\sigma_j = \alpha_j \beta_j : Z''_j \rightarrow Z$, $j = 1, 2$, are symplectic resolutions.*

Proof. As all data are explicit, the assertion can be checked by brute calculation. To cope with the computational complexity we use the free computer algebra system SINGULAR¹ [4]. It suffices to treat one of the two cases of the theorem. We indicate the basic steps for $j = 2$. In order to improve the readability of the formulae we write $q = \sqrt{-3}$.

Let $\mathbb{C}[x_1, x_2, x_3, x_4]$ denote the ring of polynomial functions on $S_1 \oplus S_2$. The invariant subring $\mathbb{C}[x_1, x_2, x_3, x_4]^T$ is generated by eight elements, listed in table 1. The kernel I of the corresponding ring homomorphism

$$\mathbb{C}[z_1, \dots, z_8] \rightarrow \mathbb{C}[x_1, x_2, x_3, x_4]^T$$

is generated by nine elements, listed in table 2. The curve C_2 is given by the semiinvariant $x_3^4 + 2qx_3^2x_4^2 + x_4^4$. In order to keep the calculation as simple as possible, the following observation is crucial: Modulo I , the Weil divisor W_2 can be described by 6 equations, listed in table 3. This leads to a comparatively ‘small’ embedding $Z'_2 \rightarrow \mathbb{P}_Z^5$ of Z -varieties. Off the origin, the effect

¹A documented SINGULAR file containing all the calculations is available from the authors upon request.

Table 1: generators for the invariant subring $\mathbb{C}[x_1, x_2, x_3, x_4]^T$:

$$\begin{aligned}
z_1 &= x_1x_3 + x_2x_4, & z_4 &= x_2x_3^3 - qx_1x_3^2x_4 + qx_2x_3x_4^2 - x_1x_4^3, \\
z_2 &= x_3^4 - 2qx_3^2x_4^2 + x_4^4, & z_5 &= x_2^3x_3 - qx_1^2x_2x_3 + qx_1x_2^2x_4 - x_1^3x_4, \\
z_3 &= x_1^4 + 2qx_1^2x_2^2 + x_2^4, & z_6 &= x_1^5x_2 - x_1x_2^5, \\
z_7 &= x_3^5x_4 - x_3x_4^5, & z_8 &= x_1x_2^2x_3^3 - x_2^3x_3^2x_4 - x_1^3x_3x_4^2 + x_1^2x_2x_4^3.
\end{aligned}$$

Table 2: generators for $I = \ker(\mathbb{C}[z_1, \dots, z_8] \rightarrow \mathbb{C}[x_1, \dots, x_4]^T)$.

$$\begin{aligned}
&qz_1^3z_5 - z_1z_3z_4 - 2z_2z_6 - z_5z_8, & z_1z_5^2 + 2z_4z_6 + z_3z_8, \\
&qz_1^3z_4 + z_1z_2z_5 - 2z_3z_7 - z_4z_8, & z_1z_4^2 - 2z_5z_7 - z_2z_8, \\
&-z_1^4 + z_2z_3 - z_4z_5 - 3qz_1z_8, & qz_1^2z_3z_5 - 2z_1^3z_6 - z_3^2z_4 + z_5^3 - 6qz_6z_8, \\
&z_1^2z_4z_5 + qz_1^3z_8 + 4z_6z_7 - z_8^2, & qz_1^2z_2z_4 - 2z_1^3z_7 - z_4^3 + z_2^2z_5 - 6qz_7z_8, \\
&4z_1^2z_4z_5 + qz_3z_4^2 - qz_2z_5^2 + 4z_6z_7 + 8z_8^2
\end{aligned}$$

Table 3: generators for the ideal of the Weil divisor $W_2 \subset Z$.

$$\begin{aligned}
b_1 &= z_3z_7 + 2z_4z_8, & b_2 &= z_2z_4 + 2qz_1z_7, & b_3 &= z_2z_3 - 4qz_1z_8, \\
b_4 &= z_2^3 + 12qz_7^2, & b_5 &= z_1z_2^2 - 6z_4z_7, & b_6 &= z_1^2z_2 - qz_4^2.
\end{aligned}$$

of blowing-up of W_2 is easy to understand even without any calculation: the action of the quaternion normal subgroup $Q_8 \subset T$ on $S_1 \oplus S_2 \setminus \{0\}$ is free. The action of $\mathbb{Z}/3 = T/Q_8$ on $S_1 \oplus S_2/Q_8$ produces transversal A_2 -singularities along a smooth two-dimensional subvariety. Blowing-up along W_1 or W_2 is a partial resolution: it introduces a \mathbb{P}^1 fibre over each singular point, and the total space contains a transversal A_1 -singularity.

The homogeneous ideal $I'_2 \subset \mathbb{C}[z_1, \dots, z_8, b_1, \dots, b_6]$ that describes the subvariety $Z'_2 \subset \mathbb{P}_Z^5$ is generated by I and 39 additional polynomials. In order to understand the nature of the singularities of Z'_2 we consider the six affine charts $U_\ell = \{b_\ell = 1\}$. The result can be summarised like this: The singular locus of Z'_2 is completely contained in $U_2 \cup U_3$, so only these charts are relevant for the discussion of the second blow-up. In fact, the corresponding affine coordinate rings have the following description:

$$R_2 = \mathbb{C}[z_1, b_3, b_4, b_5, b_6]/(b_5b_6 - 2qz_1)^2 + b_4(3qb_3 - b_6^3)$$

is a transversal A_1 -singularity.

$$R_3 = \mathbb{C}[z_1, z_3, z_5, z_6, b_1, b_2, b_6]/J,$$

where J is generated by five elements, listed in table 4. Inspection of these generators shows that $\text{Spec}(R_2)$ is isomorphic to the singularity $(\mathfrak{h}_3 \oplus \mathfrak{h}_3^*)/S_3$,

Table 4: generators for the ideal sheaf J of $Z'_2 \subset \mathbb{C}^7$ in the third chart:

$$\begin{aligned}
&4z_1b_1 + qz_3b_2 + z_5b_6, & z_1z_5 + z_3b_1 + qz_6b_6, \\
&z_1^2b_6 - z_3b_6^2 - 4qb_1^2 - 3z_5b_2, & z_1^2z_3 - z_3^2b_6 - qz_5^2 - 12z_6b_1, \\
&z_1^3 - z_1z_3b_6 + qz_5b_1 + 3qz_6b_2
\end{aligned}$$

the symplectic singularity of Coxeter type A_2 that appears as case 1 in Bellamy's theorem. It is well-known that blowing up the singular locus yields a small resolution. For arbitrary n , this is a theorem of Haiman [5, Prop. 2.6], in our case it is easier to do it directly. Thus blowing-up the reduced singular locus of Z'_2 produces a smooth resolution $Z''_2 \rightarrow Z$.

It remains to check that the morphism $\alpha_2 : Z_2 \rightarrow Z$ is semi-small. For this it suffices to verify that the fibre $E = (\alpha_2^{-1}(0))_{\text{red}}$ over the origin is two-dimensional and not contained in the singular locus of Z'_2 . Indeed, the computer calculation shows that $E \subset \mathbb{P}^5$ is given by the equations $b_1, b_3b_5, b_3b_4, b_5^2 - b_4b_6$ and hence is the union of two irreducible surfaces. The singular locus of Z'_2 is irreducible and two-dimensional and dominates the singular locus of Z . Thus the second requirement is fulfilled, too. \square

Though the theorem admits an almost conceptual formulation the proof does not: in fact, we do explicit calculations that given the complexity of the singularity we were able to carry out only by means of appropriate software. Remark that even in the case of the classical ADE-singularities arising from finite subgroups $G \subset \text{SU}(2)$ the actual resolutions of \mathbb{C}^2/G could only be described by explicit calculations. The difference to our case essentially is one of complexity: The dimension is four instead of two, there are 8 basic invariants satisfying 9 relations instead of Klein's three invariants with a single relation, and the singular locus is itself a complicated singular variety instead of an isolated point. It is only rather recently and after several decades that Nakamura gave us a conceptual tool resolving all Kleinian singularities in one stroke: the G -Hilbert scheme, known to be smooth in dimensions 2 and 3, but which is actually singular in our case of dimension 4, as we will see in the next paragraph.

2. THE EQUIVARIANT HILBERT SCHEME

Following Nakamura, we denote by $T\text{-Hilb}(\mathbb{C}^4)$ the Hilbert scheme of all T -equivariant zero-dimensional subschemes $\xi \subset \mathbb{C}^4 = S_1 \oplus S_2$ with the

property that \mathcal{O}_ξ is isomorphic as a T -representation to the regular representation of T . There is a canonical morphism $T\text{-Hilb}(\mathbb{C}^4) \rightarrow \mathbb{C}^4/T$ that resolves the singularities off the origin.

The two factors of the group $\mathbb{C}^* \times \mathbb{C}^*$ act on $S_1 \oplus S_2$ via dilations on the first and second summand, respectively, and the polynomial ring may accordingly be decomposed into irreducible $T \times \mathbb{C}^* \times \mathbb{C}^*$ -representations. Using this decomposition one can see that $T\text{-Hilb}(\mathbb{C}^4)$ contains a component isomorphic to $\mathbb{P}^2 \times \mathbb{P}^2$ and consisting entirely of subschemes ξ that are supported at the origin of \mathbb{C}^4 . Thus a generic point of this component cannot be deformed to the T -orbit of a general point in \mathbb{C}^4 , and $T\text{-Hilb}(\mathbb{C}^4)$ is not irreducible. Our calculation shows that $T\text{-Hilb}(\mathbb{C}^4)$ has only one more component, namely the closure of the points corresponding to general orbits and which we call for lack of a better name the dynamical Hilbert scheme $T\text{-Hilb}(\mathbb{C}^4)^{\text{dyn}}$. Moreover, $T\text{-Hilb}(\mathbb{C}^4)^{\text{dyn}}$ is smooth and intersects $\mathbb{P}^2 \times \mathbb{P}^2$ transversally, and finally that there are morphisms

$$\begin{array}{ccc}
 & T\text{-Hilb}(\mathbb{C}^4)^{\text{dyn}} & \\
 \swarrow & & \searrow \\
 Z_1'' & & Z_2'' \\
 \searrow & & \swarrow \\
 & Z &
 \end{array}$$

so that the two resolutions of Z discussed above are related by a Mukai-flop. However, the computations on which these assertions are based are far more involved than those referred to above, involving for example the calculation of the versal equivariant deformation spaces of all $\mathbb{C}^* \times \mathbb{C}^*$ -fixed points on $T\text{-Hilb}(\mathbb{C}^4)$. Due to their complexity these calculations might be prone to error, and the last claims are not water-proof.

REFERENCES

- [1] G. Bellamy, On singular Calogero-Moser spaces. July 2007. arXiv:0707.3694
- [2] A. M. Cohen, Finite quaternionic reflection groups. J. Algebra 64 (1980), no. 2, 293–324.
- [3] V. Ginzburg, D. Kaledin, Poisson deformations of symplectic quotient singularities. Adv. Math. 186 (2004), no. 1, 1–57.
- [4] G.-M. Greuel, G. Pfister, and H. Schönemann. SINGULAR 3-0-4. A Computer Algebra System for Polynomial Computations. Centre for Computer Algebra, University of Kaiserslautern (2001). <http://www.singular.uni-kl.de>
- [5] M. Haiman, t, q -Catalan numbers and the Hilbert scheme, Discrete Math. 193 (1998), no. 1-3, 201–224, Selected papers in honor of Adriano Garsia (Taormina, 1994).

- [6] M. Haiman, Hilbert schemes, polygraphs, and the Macdonald positivity conjecture., J. Amer. Math. Soc. 14 (2001), 941–1006.
- [7] D. Kaledin, On crepant resolutions of symplectic quotient singularities. Selecta Math. (N.S.) 9 (2003), no. 4, 529–555.
- [8] G. C. Shephard, J. A Todd, Finite unitary reflection groups. Canadian J. Math. 6, (1954). 274–304.
- [9] M. Verbitsky, Holomorphic symplectic geometry and orbifold singularities. Asian J. Math. 4 (2000), no. 3, 553–563.

MANFRED LEHN, FACHBEREICH MATHEMATIK UND INFORMATIK, JOHANNES GUTENBERG-UNIVERSITÄT MAINZ, D-55099 MAINZ

E-mail address: `lehn@mathematik.uni-mainz.de`

CHRISTOPH SORGER, LABORATOIRE DE MATHÉMATIQUES JEAN LERAY (UMR 6629 DU CNRS) & INSTITUT UNIVERSITAIRE DE FRANCE, UNIVERSITÉ DE NANTES, BP 92208, 2, RUE DE LA HOUSINIÈRE, F-44322 NANTES CEDEX 03, FRANCE

E-mail address: `christoph.sorger@univ-nantes.fr`