



**HAL**  
open science

## Excited Brownian Motions

Olivier Raimond, Bruno Schapira

► **To cite this version:**

| Olivier Raimond, Bruno Schapira. Excited Brownian Motions. 2008. hal-00332059v2

**HAL Id: hal-00332059**

**<https://hal.science/hal-00332059v2>**

Preprint submitted on 20 Oct 2008 (v2), last revised 17 Oct 2010 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# EXCITED BROWNIAN MOTIONS

OLIVIER RAIMOND AND BRUNO SCHAPIRA

ABSTRACT. We introduce and study a natural continuous time version of excited random walks. In the case of nonnegative drift, we obtain a necessary and sufficient condition for recurrence. This result is analogous to Zerner's result [Zer1] for excited (or cookie) random walks. We use similar arguments.

## 1. INTRODUCTION

Random processes that interact with their past trajectory have been studied a lot these past years. Reinforced random walks were introduced by Coppersmith and Diaconis, and then studied by Pemantle, Davis and many other authors (see the recent survey [Pem]). Some examples of time and space-continuous processes defined by a stochastic differential equation have also been studied. For example the self-interacting (or attracting) diffusions studied by Benaïm, Ledoux and Raimond (see [BeLR, BeR1, BeR2]) and also by Cranston, Le Jan, Herrmann, Kurtzmann and Roynette (see [CLJ, HR, Ku]), are processes defined by a stochastic differential equation for which the drift term is a function of the present position and of the occupation measure of the past process. Another example which is not solution of a stochastic differential equation was studied by Tóth and Werner [TW].

Carmona, Petit and Yor [CPY], Davis [D2], and Perman and Werner [PW] (see also other references therein) studied what they called a perturbed Brownian motion, which is the real valued process  $X$  defined by

$$X_t = B_t + \alpha \sup_{s \leq t} X_s + \beta \inf_{s \leq t} X_s,$$

where  $B$  is a Brownian motion. This process can be viewed as a weak limit of once edge-reinforced random walks on  $\mathbb{Z}$  (see in particular [D1, D2, W]).

More recently, excited (or cookie) random walks were introduced by Benjamini and Wilson [BW], and then further studied first on  $\mathbb{Z}^d$  [ABK, BaS1, BaS2, KZer, Ko1, Ko2, Zer1, Zer2], but also on trees [BaS3, V]. In this class of walks, the transition probabilities depend on the number of times the walk has visited the present site. In particular Kosygina and Zerner [KZer] showed that on  $\mathbb{Z}$  and if  $p_i$  is the probability to go from  $x$  to  $x + 1$  after the  $i$ -th visit of  $x$ , then the walk is a.s. recurrent if, and only if,

$$\sum_i (2p_i - 1) \in [-1, 1]$$

and it is a.s. transient otherwise.

---

2000 *Mathematics Subject Classification.* 60F20; 60K35.

*Key words and phrases.* Reinforced process; Excited process; Self-interacting process.

We present what could be a continuous time version of this. Excited Brownian motions considered here are defined by the stochastic differential equation

$$dX_t = dB_t + \varphi(X_s, L_s^{X_s}) ds,$$

for some bounded measurable  $\varphi$ , and where  $B$  is a Brownian motion and  $L_s^x$  is the local time in  $x$  at time  $s$  of  $X$ . We prove here a similar result to the one of Zerner [Zer1] for excited random walks: when  $\varphi(x, l)$  is a nonnegative function constant in  $x$ , recurrence is equivalent to

$$\int_0^\infty \varphi(0, l) dl \leq 1.$$

In particular when  $\varphi(x, l) = b1_{\{l \leq L\}}$ , recurrence is equivalent to  $bL \in [-1, 1]$ . This last case corresponds to the case the drift is  $b$  when the local time is less than  $L$  and 0 after. This is similar with the excited random walk with  $M$  cookies per sites: when a cookie (or more) is present at site  $x$  there is a probability  $p$  to go from  $x$  to  $x + 1$  (and after this visit the cookie is eaten), and when there is no cookie this probability is  $1/2$ , in which case recurrence is equivalent to  $(2p - 1)M \in [-1, 1]$ .

The paper is organized as follows. In the next section we define excited Brownian motions and give some elementary properties. In section 3 we describe the law of the excursions of our process above or below some level. In section 4 we study the property of recurrence and prove a general 0-1 law. In section 5 we study the particular case of nonnegative  $\varphi$  and obtain a necessary and sufficient criterion for recurrence or transience. In the last section, we add some remarks for general  $\varphi$  and prove in particular a necessary condition for recurrence. It is natural to believe that this condition is also sufficient, as it is the case for excited random walks [KZer]. But a proof of this is out of reach with the techniques of the present paper. Our arguments are similar to the ones of [Zer1].

*Acknowledgments:* We would like to thank Itai Benjamini for his suggestion to study the processes introduced in this paper.

## 2. DEFINITIONS AND FIRST PROPERTIES

Denote by  $(\Omega, \mathcal{F}, \mathbb{Q}_x)$  the Wiener space, where  $\mathbb{Q}_x$  is the law of a real Brownian motion started at  $x$ . Define  $X_t(\omega) = \omega(t)$  for all  $t \geq 0$  and  $(\mathcal{F}_t, t \geq 0)$  the filtration associated to  $X$ . In the following,  $L_t^y$  denotes the local time process of  $X$  at level  $y$  and at time  $t$ .

Let  $\Lambda$  be the set of measurable bounded functions  $\varphi : \mathbb{R} \times \mathbb{R}^+ \rightarrow \mathbb{R}$ . The subset of  $\Lambda$  of nonnegative functions will be denoted by  $\Lambda^+$ . We will denote by  $\Lambda_c$  and  $\Lambda_c^+$  the sets of functions  $\varphi$  in  $\Lambda$  (resp. in  $\Lambda^+$ ) such that  $\varphi(x, l)$  is a constant function of  $x$ .

For  $\varphi \in \Lambda$ , set

$$M_t^\varphi := \exp \left( \int_0^t \varphi(X_s, L_s^{X_s}) dX_s - \frac{1}{2} \int_0^t \varphi^2(X_s, L_s^{X_s}) ds \right).$$

Then  $(M_t^\varphi, t \geq 0)$  is an  $(\mathcal{F}_t, t \geq 0)$  martingale. Define  $\mathbb{P}_{x,t}^\varphi$  as the probability measure on  $(\Omega, \mathcal{F}_t)$  having a density  $M_t^\varphi$  with respect to  $\mathbb{Q}_x$  restricted to  $\mathcal{F}_t$ . By consistency, it is possible to construct a (unique) probability measure  $\mathbb{P}_x^\varphi$  on  $\Omega$ , such that  $\mathbb{P}_x^\varphi$  restricted to  $\mathcal{F}_t$  is  $\mathbb{P}_{x,t}^\varphi$ . By the transformation of drift formula (Girsanov

Theorem), one proves that under  $\mathbb{P}_x^\varphi$ ,

$$B_t = X_t - x - \int_0^t \varphi(X_s, L_s^{X_s}) ds$$

is a Brownian motion started at 0.

This proves (the uniqueness follows by a similar argument: start with  $\mathbb{P}_x^\varphi$  the law of a solution and then construct  $\mathbb{P}_{x,t}^\varphi$ ) the following

**Proposition 2.1.** *Let  $(x, \varphi) \in \mathbb{R} \times \Lambda$ . Then there is a unique solution  $(X, B)$  to the equation*

$$(1) \quad X_t = x + B_t + \int_0^t \varphi(X_s, L_s^{X_s}) ds$$

with  $L_t^y$  the local time of  $X$  at level  $y$  and at time  $t$ , and such that  $B$  is a Brownian motion started at 0.

For clarity, we will sometimes write  $\mathbb{Q}$  for  $\mathbb{Q}_0$  and  $\mathbb{P}$  for  $\mathbb{P}_0^\varphi$ , if there is no ambiguity on  $\varphi$ . We will use the notation  $\mathbb{E}$  and  $\mathbb{E}_x^\varphi$  for the expectations with respect to  $\mathbb{P}$  and  $\mathbb{P}_x^\varphi$ . For other probability measures  $\mu$ , the expectation of a random variable  $Z$  will simply be denoted by  $\mu(Z)$ .

Let  $(x, \varphi) \in \mathbb{R} \times \Lambda$ . Under  $\mathbb{P}_x^\varphi$ , for all stopping times  $T$ , on the event  $\{T < \infty\}$ ,

$$(2) \quad \text{the law of } (X_{t+T}, t \geq 0) \text{ given } \mathcal{F}_T \text{ is } \mathbb{P}_{X_T}^{\varphi_T},$$

where  $\varphi_T \in \Lambda$  is defined by

$$\varphi_T(y, l) = \varphi(y, L_T^y + l).$$

Note that  $(X_t, \varphi_t)$  is a Markov process and that (2) is just the strong Markov property for this process.

We denote by  $D_t$  the drift accumulated at time  $t$  :

$$D_t = \int_0^t \varphi(X_s, L_s^{X_s}) ds.$$

**Lemma 2.2.** *Set  $h(x, l) = \int_0^l \varphi(x, u) du$ . The drift term  $D_t$  is also equal to*

$$\int_{\mathbb{R}} h(x, L_t^x) dx.$$

*Proof.* This follows from the occupation times formula given in exercise (1.15) in the chapter VI of Revuz–Yor [RY].  $\square$

In the following, we set for any Borel set  $A$  of  $\mathbb{R}$ ,

$$D_t^A = \int_A h(x, L_t^x) dx.$$

We will use also the notation  $D_t^+$ ,  $D_t^-$  and  $D_t^k$ ,  $k \in \mathbb{Z}$ , respectively for  $D_t^{\mathbb{R}^+}$ ,  $D_t^{\mathbb{R}^-}$  and  $D_t^{(k, k+1)}$ . Note that (this is still a consequence of Exercise (1.15) Chapter VI in [RY])

$$D_t^A = \int_0^t \varphi(X_s, L_s^{X_s}) 1_A(X_s) ds.$$

**Lemma 2.3.** *Let  $(\varphi_n)_{n \geq 0} \in \Lambda$  be a sequence of functions, such that  $\varphi_n(x, l)$  converges toward  $\varphi(x, l)$ , when  $n \rightarrow +\infty$ , for a.e.  $x$  and  $l$ , and such that*

$$\sup_{x, n, u} |\varphi_n(x, u)| < +\infty.$$

*Then  $\mathbb{P}_x^{\varphi_n}$  converges weakly on  $\Omega$  toward  $\mathbb{P}_x^\varphi$  for all  $x$ .*

*Proof.* Let  $Z$  be a bounded continuous  $\mathcal{F}_t$ -measurable random variable. We have to prove that  $\mathbb{E}_x^{\varphi_n}(Z)$  converges towards  $\mathbb{E}_x^\varphi(Z)$ . Since  $\mathbb{E}_x^{\varphi_n}(Z)$  and  $\mathbb{E}_x^\varphi(Z)$  are respectively equal to  $\mathbb{Q}_x(ZM_t^{\varphi_n})$  and to  $\mathbb{Q}_x(ZM_t^\varphi)$  it suffices to prove that  $M_t^{\varphi_n}$  converges in  $L^2$  towards  $M_t^\varphi$ . Using Itô calculus, we prove that

$$\frac{d}{dt} \mathbb{Q}_x[(M_t^{\varphi_n} - M_t^\varphi)^2] \leq 2\mathbb{Q}_x[(\varphi_n - \varphi)^2(X_t, L_t^{X_t})(M_t^\varphi)^2] + C\mathbb{Q}_x[(M_t^{\varphi_n} - M_t^\varphi)^2],$$

for some constant  $C > 0$ . By dominated convergence, the first term of the right hand term converges to 0. We conclude by using Gronwall's lemma.  $\square$

### 3. CONSTRUCTION WITH EXCURSIONS

Define the processes  $A^+$  and  $A^-$  as follows:

$$A_t^+ = \int_0^t 1_{\{X_s > 0\}} ds \quad \text{and} \quad A_t^- = \int_0^t 1_{\{X_s < 0\}} ds.$$

Define the right-continuous inverses of  $A^+$  and  $A^-$  as

$$\kappa^+(t) = \inf\{u > 0 \mid A_u^+ > t\} \quad \text{and} \quad \kappa^-(t) = \inf\{u > 0 \mid A_u^- > t\}.$$

Define the two processes  $X^+$  and  $X^-$  by

$$X_t^+ = X_{\kappa^+(t)} \quad \text{and} \quad X_t^- = X_{\kappa^-(t)}.$$

Denote by  $\mathbb{Q}^+$  and  $\mathbb{Q}^-$  the laws respectively of  $X^+$  and  $X^-$  under  $\mathbb{Q}$ , and let  $\mathbb{Q}_t^+$  and  $\mathbb{Q}_t^-$  respectively be their restrictions to  $\mathcal{F}_t$  (then  $\mathbb{Q}_t^\pm$  is the law of  $(X_s^\pm; s \leq t)$ ). It is known that  $\mathbb{Q}^+$  (resp.  $\mathbb{Q}^-$ ) is the law of a Brownian motion reflected above 0 (resp. below 0) and started at 0. The process  $\beta$ , defined by

$$\beta_t := X_t - L_t^0 \quad (\text{resp. } \beta_t := X_t + L_t^0)$$

(recall that  $L^0$  is the local time process in 0 of  $X$ ) is a Brownian motion under  $\mathbb{Q}^+$  (resp. under  $\mathbb{Q}^-$ ). Denote by  $N_t^\varphi$  the martingale on  $(\Omega, \mathcal{F}, \mathbb{Q}^\pm)$  defined by

$$N_t^\varphi := \exp\left(\int_0^t \varphi(X_s, L_s^{X_s}) d\beta_s - \frac{1}{2} \int_0^t \varphi^2(X_s, L_s^{X_s}) ds\right).$$

Let  $\mathbb{P}^{\varphi, \pm}$  be the measures whose restrictions  $\mathbb{P}_t^{\varphi, \pm}$  to  $\mathcal{F}_t$  are defined by

$$\mathbb{P}_t^{\varphi, \pm} := N_t^\varphi \cdot \mathbb{Q}_t^\pm \quad \forall t \geq 0.$$

Note that, by using Girsanov Theorem, on the space  $(\Omega, \mathcal{F}, \mathbb{P}^{\varphi, \pm})$ ,

$$(3) \quad X_t = \beta_t^\pm \pm L_t + \int_0^t \varphi(X_s, L_s^{X_s}) ds,$$

with  $\beta_t^\pm$  a Brownian motion.

Set  $\tilde{\mathbb{P}}^\varphi := \mathbb{P}^{\varphi, +} \otimes \mathbb{P}^{\varphi, -}$  and let  $(X^1, X^2)$  be the canonical process of law  $\tilde{\mathbb{P}}^\varphi$ . Then  $X^1$  and  $X^2$  are independent and respectively distributed like  $\mathbb{P}^{\varphi, +}$  and  $\mathbb{P}^{\varphi, -}$ .

Denote by  $L^{(1)}$  and  $L^{(2)}$  the local time processes in 0 of  $X^1$  and  $X^2$ , and define their right continuous inverses  $\tau^1$  and  $\tau^2$  by

$$\tau_s^i = \inf\{t \mid L_t^{(i)} > s\} \quad \text{for } i \in \{1, 2\}.$$

Denote by  $e^1$  and  $e^2$  their excursion processes out of 0: for  $s \leq L_\infty^{(i)}$ ,

$$e_s^i(u) = X_{\tau_{s-}^i + u}^i \quad \text{for all } u \in (0, \tau_s^i - \tau_{s-}^i),$$

if  $\tau_s^i - \tau_{s-}^i > 0$ , and  $e_s^i = 0$  otherwise. Let now  $e$  be the excursion process obtained by adding  $e^1$  and  $e^2$ .

Denote by  $\Xi$  the measurable transformation that reconstructs a process out of its excursion process (see [RY] Proposition (2.5) p.482). Note that  $\Xi$  is not a one to one map, it is only surjective (think of processes having an infinite excursion out of 0, in which case  $L_\infty^0 < \infty$ ).

**Proposition 3.1.** *The following hold*

- (i) *The law of the process  $\Xi e$  is  $\mathbb{P}$ .*
- (ii)  $((\Xi e)_t^+, t \leq L_\infty^{(2)}) = (X_t^1, t \leq L_\infty^{(2)})$ .
- (iii)  $((\Xi e)_t^-, t \leq L_\infty^{(1)}) = (X_t^2, t \leq L_\infty^{(1)})$ .
- (iv) *Denote by  $L$  the local time in 0 of  $\Xi e$ . Then for all  $t \in \mathbb{R}^+ \cup \{\infty\}$ ,  $L_t = L_t^{(1)} \wedge L_t^{(2)}$ .*

*Proof.* Assume first that  $\varphi(x, l) = 0$  if  $x \in (-c, c)$ , for some constant  $c > 0$ . For any  $\epsilon \in (-c, c)$ , define a process  $X^\epsilon$  as follows: set  $T_0^\epsilon = 0$  and for  $n \geq 1$ ,

$$S_n^\epsilon = \inf\{t \geq T_{n-1}^\epsilon \mid X_t \in \{-\epsilon, \epsilon\}\},$$

$$T_n^\epsilon = \inf\{t \geq S_n^\epsilon \mid X_t = 0\}.$$

Define also

$$A_\epsilon(t) = \sum_{n \geq 1} (T_n^\epsilon \wedge t - S_n^\epsilon \wedge t),$$

and let  $\kappa_\epsilon(t)$  be the right-continuous inverse of  $A_\epsilon^+$ . Then set

$$X_t^\epsilon := X_{\kappa_\epsilon(t)}.$$

Now observe that during each time-interval  $(S_n^\epsilon, T_n^\epsilon)$ , the local time in 0 of  $X$  cannot increase. And for  $t \in (A_\epsilon(S_n^\epsilon), A_\epsilon(T_n^\epsilon))$ ,  $\kappa_\epsilon(t) = S_n^\epsilon + (t - A_\epsilon(S_n^\epsilon))$ . So by (3), during the intervals  $(A_\epsilon(S_n^\epsilon), A_\epsilon(T_n^\epsilon))$ , if  $X$  follows the law of  $\Xi e$ , then  $X^\epsilon$  is solution of the SDE

$$dX_t^\epsilon = dR_t^\epsilon + \varphi\left(X_t^\epsilon, L_{\kappa_\epsilon(t)}^{X_t^\epsilon}\right) dt,$$

where  $L$  is the local time of  $X$ , and  $R^\epsilon$  is the Brownian motion defined by

$$R_t^\epsilon = \sum_{k=1}^{n-1} (B_{T_k^\epsilon} - B_{S_k^\epsilon}) + (B_{\kappa_\epsilon(t)} - B_{S_n^\epsilon})$$

for  $t \in (A_\epsilon(S_n^\epsilon), A_\epsilon(T_n^\epsilon))$ .

Denote by  $L^{\epsilon, \cdot}$  the local time process of  $X^\epsilon$ . Then

$$L_{\kappa_\epsilon(t)}^x = L_t^{\epsilon, x} \quad \forall t \geq 0 \quad \forall x \notin (-c, c).$$

Since  $\varphi(x, l) = 0$  when  $x \in (-c, c)$ ,

$$\varphi(x, L_{\kappa_\epsilon(t)}^x) = \varphi(x, L_t^{\epsilon, x}) \quad \forall t \geq 0 \quad \forall x \in \mathbb{R}.$$

Thus  $X^\epsilon$  satisfies in fact the SDE:

$$(4) \quad dX_t^\epsilon = dR_t^\epsilon + \varphi\left(X_t^\epsilon, L_t^{\epsilon, X_t^\epsilon}\right) dt,$$

up to the first time it hits 0. And when it hits 0, it jumps instantaneously to  $\epsilon$  or to  $-\epsilon$  with probability 1/2, independently of its past trajectory. Note that this determines the law of  $X^\epsilon$ .

But it follows also from (1), that if  $X$  has law  $\mathbb{P}$ , then  $X^\epsilon$  solves as well the SDE (4) up to the first time it hits 0, and then jump to  $\epsilon$  or  $-\epsilon$  with probability 1/2 (since there is no drift in  $(-c, c)$ ). Since moreover  $X^\epsilon$  converges to  $X$ , when  $\epsilon$  goes to 0, we conclude that the law of  $\Xi e$  is  $\mathbb{P}$ .

To finish the proof of (i), for  $c > 0$ , define  $\varphi_c$  by

$$\varphi_c(x, l) = \varphi(x, l)1(x \notin [-c, c]).$$

By Lemma 2.3 we know that  $\mathbb{P}_0^{\varphi_c}$  converges toward  $\mathbb{P}_0^\varphi$ , when  $c \rightarrow 0$ . It can be also seen that  $\mathbb{P}^{\varphi_c, +}$  and  $\mathbb{P}^{\varphi_c, -}$  converge respectively towards  $\mathbb{P}^{\varphi, +}$  and  $\mathbb{P}^{\varphi, -}$ . Since  $\Xi e$  is a measurable transformation of  $(X^1, X^2)$ , we can conclude.

Assertions (ii), (iii) and (iv) are immediate: in the construction of  $\Xi e$ , one needs only to know  $e_s$  for  $s \leq L_\infty = L_\infty^{(1)} \wedge L_\infty^{(2)}$ . So  $(\Xi e)^+$  and  $(\Xi e)^-$  can be respectively reconstructed with the positive and negative excursions of  $(e_s, s \leq L_\infty^{(1)} \wedge L_\infty^{(2)})$ .  $\square$

#### 4. RECURRENCE, TRANSCIENCE AND A 0 – 1 LAW

Let  $\varphi \in \Lambda$ . Consider the events  $R_a := \{L_\infty^a = +\infty\}$ ,  $a \in \mathbb{R}$ . Using conditional Borel-Cantelli lemma, one can prove that for all  $x, y, z$ , and all  $\varphi \in \Lambda$ ,  $\mathbb{P}_x^\varphi$ -a.s.,  $R_y = R_z$ . In the following, we will denote by  $R$  the event of recurrence ( $= R_a$  for all  $a$ ).

We will first study the question of recurrence and transience for the processes  $X^1$  and  $X^2$  separately, where  $X^1$  and  $X^2$  are independent respectively of law  $\mathbb{P}^{\varphi, +}$  and  $\mathbb{P}^{\varphi, -}$ . Note that we still have for all  $x \geq 0$  (resp.  $x \leq 0$ ), and all  $\varphi \in \Lambda$ ,  $\mathbb{P}^{\varphi, +}$ -a.s. (resp.  $\mathbb{P}^{\varphi, -}$ -a.s.),  $R_x = R_0 (= R)$ . So in all cases,  $R$  is the event  $\{L_\infty^0 = \infty\}$ .

Fix  $x \in \mathbb{R}$  and denote by  $\varphi_x$  the function in  $\Lambda$  such that for all  $(y, l) \in \mathbb{R} \times \mathbb{R}^+$ ,

$$\varphi_x(y, l) = \varphi(x + y, l).$$

In the following,  $\mathbb{P}_x^{\varphi, \pm}$  will denote  $\mathbb{P}^{\varphi_x, \pm}$ .

**Proposition 4.1.** *For all  $x \in \mathbb{R}$  and all  $\varphi \in \Lambda$ , the following holds*

$$(5) \quad \mathbb{P}_x^\varphi(R) = \mathbb{P}_x^{\varphi, +}(R) \times \mathbb{P}_x^{\varphi, -}(R).$$

*Proof.* This is a straightforward application of Proposition 3.1 since

$$\begin{aligned} \mathbb{P}_x^\varphi(R) &= \tilde{\mathbb{P}}^{\varphi_x}(L_\infty = \infty) \\ &= \tilde{\mathbb{P}}^{\varphi_x}(L_\infty^{(1)} = L_\infty^{(2)} = \infty). \end{aligned}$$

and we conclude since  $L_\infty^{(1)}$  and  $L_\infty^{(2)}$  are independent.  $\square$

For  $t > 0$  and  $a \in \mathbb{R}$ , set  $\sigma_t^a = \inf\{s > 0 \mid \int_0^s 1_{\{X_u \geq a\}} du > t\}$ . Then we have

**Lemma 4.2.** *Let  $a \in \mathbb{R}$ ,  $x, y \leq a$  and  $\varphi, \psi \in \Lambda$  be such that  $\varphi(z, l) = \psi(z, l)$  for all  $z \geq a$  and all  $l \geq 0$ . Then  $(X_{\sigma_t^a} - a, t \geq 0)$  has the same distribution under  $\mathbb{P}_x^{\varphi, +}$  and under  $\mathbb{P}_y^{\psi, +}$ . In particular,*

$$\mathbb{P}_x^{\varphi, +}(L_\infty^a = \infty) = \mathbb{P}_y^{\psi, +}(L_\infty^a = \infty).$$

*Proof.* The lemma follows immediately from Proposition 3.1.  $\square$

Note that a similar proposition holds for  $\mathbb{P}^{\varphi,-}$ . A direct consequence of this lemma is that

**Proposition 4.3.** *For all  $x \in \mathbb{R}$ ,  $\mathbb{P}_x^{\varphi,\pm}(R) = \mathbb{P}^{\varphi,\pm}(R)$ .*

With this in hands, we can prove the

**Proposition 4.4** (Zero-one law). *Let  $\varphi \in \Lambda$ . Then  $\mathbb{P}^{\varphi,+}(R)$  and  $\mathbb{P}^{\varphi,-}(R)$  both belong to  $\{0, 1\}$ .*

*Proof.* Assume that  $\mathbb{P}^{\varphi,+}(R) > 0$ . By martingale convergence theorem,  $\mathbb{P}^{\varphi,+}$ -a.s.

$$\begin{aligned} 1_R &= \lim_{z \rightarrow +\infty} \mathbb{P}_x^{\varphi,+}(R \mid \mathcal{F}_{T_z}) \\ &= \lim_{z \rightarrow +\infty} \mathbb{P}_z^{\varphi T_z,+}(R) \\ &= \mathbb{P}^{\varphi,+}(R), \end{aligned}$$

by application of Lemma 4.2 and Proposition 4.3. Thus  $\mathbb{P}^{\varphi,+}(R) = 1$ , which proves the proposition.  $\square$

Denote by  $T$  the event of transience. Then  $T = R^c = \{L_\infty^0 < \infty\}$ . Note that  $\mathbb{P}^{\varphi,\pm}$ -a.s.,  $T = \{X \rightarrow \pm\infty\}$ , and that  $\mathbb{P}_x^\varphi$ -a.s.,  $T = \{X \rightarrow +\infty\} \cup \{X \rightarrow -\infty\}$  and that  $\tilde{\mathbb{P}}^\varphi$ -a.s.,  $\{\exists e \text{ is transient}\} = \{X^1 \rightarrow \infty\} \cup \{X^2 \rightarrow -\infty\}$ . Then we have the criterion

**Proposition 4.5.** *Let  $\varphi \in \Lambda$ .*

- (i)  $\mathbb{P}_x^\varphi(R) = 1$  for all  $x$  if, and only if,  $\mathbb{P}^{\varphi,+}(R) = \mathbb{P}^{\varphi,-}(R) = 1$ .
- (ii)  $\mathbb{P}_x^\varphi(R) = 0$  for all  $x$  if, and only if,  $\mathbb{P}^{\varphi,+}(R) = 0$  or  $\mathbb{P}^{\varphi,-}(R) = 0$ .

*Proof.* This is a straightforward consequence of Propositions 4.1, 4.3 and 4.4.  $\square$

## 5. CASE $\varphi \geq 0$

In this section, we will consider functions belonging to  $\Lambda^+$ . In this case, it is obvious that  $\mathbb{P}^{\varphi,-}(R) = 1$ . Thus the recurrence property only depends on  $\mathbb{P}^{\varphi,+}(R)$ .

Set  $T_a = \inf\{t > 0 \mid X_t = a\}$ .

**Lemma 5.1.** *Let  $\varphi \in \Lambda^+$ , for which there exists  $x_0$  such that for all  $x \leq x_0$  and all  $l \geq 0$ ,  $\varphi(x, l) = \varphi(x_0, l)$ , and where  $\varphi(x_0, l)$  is non zero function of  $l$ . Then, for all  $a \geq x$ ,*

$$\mathbb{E}_x^\varphi(D_{T_a}) = a - x.$$

*Proof.* Without losing generality, we prove this result for  $x = 0$ . Take  $a > 0$ . We have for all  $n$ ,

$$0 = \mathbb{E}_0^\varphi(B_{T_a \wedge n}) = \mathbb{E}_0^\varphi(X_{T_a \wedge n}) - \mathbb{E}_0^\varphi(D_{T_a \wedge n}).$$

By monotone convergence,  $\lim_{n \rightarrow \infty} \mathbb{E}_0^\varphi(D_{T_a \wedge n}) = \mathbb{E}_0^\varphi(D_{T_a})$ . Thus, if we can prove that  $\lim_{n \rightarrow \infty} \mathbb{E}_0^\varphi(X_{T_a \wedge n}) = a$ , the lemma will follow. Since  $\mathbb{P}_0^\varphi$ -a.s.,  $X_{T_a} = a$  and  $T_a < \infty$ , this is equivalent to prove that

$$\lim_{n \rightarrow \infty} \mathbb{E}_0^\varphi[X_n 1_{\{n < T_a\}}] = 0.$$

For all  $n$ ,

$$\min_{t < T_a} X_t \leq X_n 1_{\{n < T_a\}} \leq a \quad \mathbb{P}_0^\varphi - a.s.$$



If one proves that  $\min_{t < T_a} X_t$  is integrable, we can conclude by dominated convergence. One has for all  $c > 0$ ,

$$\begin{aligned} \mathbb{E}_0^\varphi \left[ -\min_{t < T_a} X_t \right] &\leq \mathbb{E}_0^\varphi [D_{T_a}/c] \\ &\quad + \sum_{i \geq 1} i \times \mathbb{P}_0^\varphi \left[ -\min_{t < T_a} X_t \in [i-1, i], D_{T_a} < ci \right]. \end{aligned}$$

Clearly  $\mathbb{E}_0^\varphi [D_{T_a}/c] \leq a/c < +\infty$ . Moreover for all  $i \geq 1$ ,

$$(6) \quad \mathbb{P}_0^\varphi \left[ -\min_{t < T_a} X_t \in [i-1, i], D_{T_a} < ci \right] \leq \mathbb{P}_0^\varphi [T_{-i} < T_a, D_{T_a} < ci].$$

Now on the event  $\{T_{-i} < T_a\}$ ,

$$D_{T_a} \geq \sum_{k=1}^i \left( \int_{-k}^{-k+1} h(y, L_{T_{-k}}^y) dy \right) \mathbf{1}_{\{T_{-k} < \infty\}}.$$

Set

$$\alpha_k := \left( \int_{-k}^{-k+1} h(y, L_{T_{-k}}^y) dy \right) \mathbf{1}_{\{T_{-k} < \infty\}}.$$

We will prove that there exists a constant  $\alpha > 0$ , such that for all integer  $k$  greater than  $-x_0$ ,

$$(7) \quad \mathbb{E}_0^\varphi [\alpha_{k+1} \mid \mathcal{F}_{T_{-k}}] \geq \alpha \mathbf{1}_{\{T_{-k} < +\infty\}}.$$

By (2) we have

$$\begin{aligned} \mathbb{E}_0^\varphi [\alpha_{k+1} \mid \mathcal{F}_{T_{-k}}] &= \mathbb{E}_{-k}^{\varphi^{T_{-k}}} [\alpha_{k+1}] \mathbf{1}_{\{T_{-k} < \infty\}} \\ &= \mathbb{E}_0^{\psi_k} [\alpha_1] \mathbf{1}_{\{T_{-k} < \infty\}} \end{aligned}$$

with  $\psi_k(x, l) = \varphi(x_0, l)$  for  $x < 0$  and  $\psi_k(x, l) = \varphi(x - k, L_{T_{-k}}^{x-k})$  for  $x \geq 0$ . Note that by Proposition 3.1,  $\Xi e$  will reach level  $-1$  in finite time if, and only if,

$$L_\infty^{(1)} \geq L_{T_{-1}}^{(2)},$$

where  $T_{-1}$  denotes also the hitting time of  $-1$  for  $X^2$ . Thus

$$\mathbb{E}_0^{\psi_k} [\alpha_1] = \tilde{\mathbb{E}}^{\psi_k} \left[ F(X_t^2, t \leq T_{-1}) \mathbf{1}_{\{L_\infty^{(1)} \geq L_{T_{-1}}^{(2)}\}} \right],$$

with

$$F(X_t^2, t \leq T_{-1}) = \int_{-1}^0 h(y, L_{T_{-1}}^{(2),y}) dy,$$

where  $L^{(2),y}$  is the local time in  $y$  of  $X^2$ . It is possible to couple  $X^1$  with a Brownian motion  $Y^+$  reflected at 0, started at 0, with drift  $\|\varphi\|_\infty$ , and such that  $X_t^1 \leq Y_t^+$  for all  $t \geq 0$ . Note that the law of  $Y^+$  is  $\mathbb{P}^{\psi,+}$ , where  $\psi(x, l) = \|\varphi\|_\infty$  if  $x \geq 0$  and  $\psi(x, l) = \varphi(x_0, l)$  if  $x < 0$ . Moreover,

$$L_\infty^{(1)} \geq L_\infty^+,$$

with  $L^+$  the local time in 0 of  $Y^+$ . These properties imply that

$$\mathbb{E}_0^{\psi_k} [\alpha_1] \geq \alpha := \mathbb{E}_0^\psi [\alpha_1].$$

It remains to see that  $\alpha$  is positive. For any  $t > 0$ , it is larger than

$$\mathbb{E}_0^\psi [\alpha_1 \mathbf{1}_{\{T_{-1} \leq t\}}].$$

By using Girsanov's transform it suffices to prove that for some  $t > 0$ ,

$$\mathbb{Q} [\alpha_1 1_{\{T_{-1} \leq t\}}] > 0.$$

But

$$\mathbb{Q} [\alpha_1 1_{\{T_{-1} \leq t\}}] = \int_{-1}^0 \mathbb{Q} [h(y, L_{T_{-1}}^y) 1_{\{T_{-1} \leq t\}}] dy.$$

By continuity of  $h$  and  $L$  it suffices to prove that for some  $y \in (-1, 0)$  and  $t > 0$ ,

$$\mathbb{Q} [h(y, L_{T_{-1}}^y) 1_{\{T_{-1} \leq t\}}] > 0.$$

But this is clear for  $y = 1/2$  for instance, since for any  $M > 0$ ,  $\mathbb{Q}[L_{T_{-1}}^{1/2} > M] > 0$ .

Next (7) shows that the process  $(M_i, i \geq 0)$  defined by

$$M_i = \sum_{k=1}^i (\alpha_k - \alpha 1_{\{T_{-k} < +\infty\}}),$$

is a sub-martingale with respect to the filtration  $(\mathcal{F}_{T_{-i}})_{i \geq 1}$ . Moreover on the event  $\{T_{-i} < T_a\}$ , one has

$$M_i = \sum_{k=1}^i (\alpha_k - \alpha).$$

So by taking  $c = \alpha/2$  in (6) one gets

$$\mathbb{P} [T_{-i} < T_a, D_{T_a} < c(i+1)] \leq \mathbb{P}[M_i \leq -\alpha i/2],$$

for all  $i \geq 0$ . We conclude now that  $\mathbb{E}[-\min_{t < T_a} X_t]$  is finite by using standard results on martingales.  $\square$

**Remark 5.2.** As noticed also by Zerner in [Zer1], the condition  $\varphi(x_0, \cdot) \neq 0$  is necessary (the result being obviously false if  $\varphi = 0$ ).

**Remark 5.3.** This lemma can be extended to the case where  $\varphi$  is random and stationary in the sense that for all  $z$ ,  $\varphi^z$  and  $\varphi$  have the same law, where  $\varphi^z$  is defined by  $\varphi^z(x, l) = \varphi(x+z, l)$ .

**Lemma 5.4.** Let  $\varphi \in \Lambda_c^+$ . Then

$$\mathbb{E}_0^\varphi(D_\infty^k) \leq 1,$$

for all  $k \geq 0$ .

*Proof.* It is the same proof than for Lemma 11 in Zerner [Zer1]. We reproduce it here for completeness. Note first that  $\mathbb{E}_0^\varphi[D_\infty^k] = \mathbb{P}_k^{\varphi,+}(D_\infty^0)$ , which does not depend on  $k$  since  $\varphi \in \Lambda_c$ . For  $K \geq 1$  and  $i \leq K-1$ ,

$$D_{T_K} \geq D_{T_K}^+ = \sum_{j=0}^{K-1} D_{T_K}^j \geq \sum_{j=0}^{K-1-i} D_{T_K}^j \geq \sum_{j=0}^{K-1-i} D_{T_{j+i}}^j.$$

Using Lemma 5.1,

$$K = \mathbb{E}_0^\varphi[D_{T_K}] \geq \sum_{j=0}^{K-1-i} \mathbb{E}_0^\varphi[D_{T_{j+i}}^j] \geq (K-i) \mathbb{E}_0^\varphi[D_{T_i}^0].$$

Letting  $i \rightarrow \infty$ , we conclude.  $\square$

**Lemma 5.5.** *Let  $\varphi \in \Lambda^+$  be such that  $\mathbb{P}^{\varphi,+}(R) = 0$ . Then*

$$\lim_{z \rightarrow \infty} \mathbb{E}_0^\varphi(D_{T_z}^+)/z = 1.$$

*If moreover  $\varphi \in \Lambda_c^+$ , then*

$$\mathbb{E}_0^\varphi(D_\infty^0) = 1.$$

*Proof.* The proof of the first part follows the proof of Lemma 6 in Zerner [Zer1]. We write it here for completeness. First, since  $\mathbb{E}^{0,\varphi}(D_{T_z}^+) = \mathbb{P}^{\varphi,+}(D_{T_z})$  and  $\mathbb{P}^{\varphi,+}(R)$  is a function of  $(\varphi(x, \cdot))_{x \geq 0}$ , we can assume that  $\varphi(x, l) = 1$  for all  $x \leq 0$  and all  $l \geq 0$ . Thus Lemma 5.1 can be applied:  $\mathbb{E}_0^\varphi(D_{T_z}) = z$ . So, it suffices to prove that  $\lim_{z \rightarrow \infty} \mathbb{E}_0^\varphi(D_{T_z}^-)/z = 0$ .

For  $i \geq 1$ , let  $\sigma_i = \inf\{j \geq T_i \mid X_j = 0\}$ . We have, for  $z$  an integer,

$$\mathbb{E}_0^\varphi[D_{T_z}^-] = \sum_{i=0}^{z-1} \mathbb{E}_0^\varphi[D_{T_{i+1}}^- - D_{T_i}^-].$$

Note that

$$\begin{aligned} \mathbb{E}_0^\varphi[D_{T_{i+1}}^- - D_{T_i}^-] &= \mathbb{E}_0^\varphi \left[ 1_{\{\sigma_i < T_{i+1}\}} (D_{T_{i+1}}^- - D_{T_i}^-) \right] \\ &= \mathbb{E}_0^\varphi \left[ 1_{\{\sigma_i < T_{i+1}\}} \mathbb{E}_0^{\varphi_{\sigma_i}}(D_{T_{i+1}}^-) \right] \\ &\leq (i+1) \mathbb{P}_0^\varphi(\sigma_i < T_{i+1}) \end{aligned}$$

using again Lemma 5.1. Thus it remains to prove that

$$\lim_{z \rightarrow \infty} \frac{1}{z} \sum_{i=1}^{z+1} i \mathbb{P}_0^\varphi(\sigma_i < T_{i+1}) = 0.$$

Let  $Y_i = \mathbb{P}_0^\varphi[\sigma_i < T_{i+1} \mid \mathcal{F}_{T_i}]$ . Since  $\mathbb{P}_0^\varphi(R) = 0$ , the conditional Borel-Cantelli lemma implies that  $\mathbb{P}_0^\varphi$ -a.s.,  $\sum_i Y_i < +\infty$ . Since  $Y_i \leq 1/i$  ( $X$  being greater than a Brownian motion), for all positive  $\epsilon$ ,

$$i \times \mathbb{P}_0^\varphi(\sigma_i < T_{i+1}) \leq \epsilon + \mathbb{P}_0^\varphi(Y_i \geq \epsilon/i).$$

This implies that

$$\frac{1}{z} \sum_{i=1}^{z+1} i \mathbb{P}_0^\varphi(\sigma_i < T_{i+1}) \leq \epsilon + \frac{1}{z} \mathbb{E}_0^\varphi \left[ \sum_{i=1}^{z+1} 1_{\{Y_i \geq \epsilon/i\}} \right].$$

But since  $\sum_i Y_i < \infty$  a.s., the density of the  $i \leq z$  such that  $Y_i \leq \epsilon/i$  tends to 0 when  $z$  tends to  $\infty$ . Thus the preceding sum converges to 0. This concludes the proof of the first part.

The second part is immediate (see Zerner [Zer1] Theorem 12). Since  $\varphi \in \Lambda_c^+$ , for all  $K \geq 0$ ,

$$\mathbb{E}_0^\varphi[D_\infty^0] = \frac{1}{K} \sum_{k=0}^{K-1} \mathbb{E}_0^\varphi[D_\infty^k] \geq \frac{1}{K} \mathbb{E}_0^\varphi[D_{T_K}^+].$$

We conclude using the first part of the lemma.  $\square$

The next result gives a sufficient criterion for recurrence, when we only know that  $\varphi \in \Lambda^+$ . For  $\varphi \in \Lambda_c^+$ , we will obtain a necessary and sufficient condition (see Theorem 5.10 below).

**Corollary 5.6.** Let  $\varphi \in \Lambda^+$ . For  $x \in \mathbb{R}$ , set  $\delta^x(\varphi) = \int_0^\infty \varphi(x, u) du$ . If

$$\liminf_{z \rightarrow +\infty} \frac{1}{z} \int_0^z \delta^x(\varphi) dx < 1,$$

then  $\mathbf{P}^{\varphi,+}(R) = 1$ .

*Proof.* Since P-a.s.  $D_{T_z}^+ \leq \int_0^z \delta^x(\varphi) dx$ , if  $\liminf \frac{1}{z} \int_0^z \delta^x(\varphi) dx < 1$ , then

$$\liminf \mathbf{E}_0^\varphi(D_{T_z}^+)/z < 1.$$

We conclude by using Lemma 5.5.  $\square$

**Lemma 5.7.** Let  $\varphi \in \Lambda^+$  be such that  $\mathbf{P}^{\varphi,+}(R) = 1$ . Then

$$\mathbf{P}_0^\varphi(T_{-1} = +\infty) > 0.$$

*Proof.* By using Proposition 3.1,

$$\mathbf{P}_0^\varphi(T_{-1} = +\infty) = \tilde{\mathbf{P}}^\varphi(L_\infty^{(1)} < L_{T_{-1}}^{(2)}),$$

where  $T_{-1}$  denotes also the hitting time of  $-1$  for  $X^2$ . Since  $X^1$  and  $X^2$  are independent and since  $\mathbf{P}^{\varphi,+}(R) = 0$  implies that  $\tilde{\mathbf{P}}^{\varphi}$ -a.s.,  $L_\infty^{(1)} < +\infty$ , it suffices to prove that for any  $l > 0$ ,

$$\mathbf{P}^{\varphi,-}(L_{T_{-1}}^0 > l) > 0.$$

Equivalently it suffices to prove that for any  $l > 0$ , there exists  $t > 0$  such that

$$\mathbf{P}^{\varphi,-}(L_{T_{-1}}^0 > l \text{ and } T_{-1} \leq t) > 0.$$

By absolute continuity of  $\mathbf{P}_{|\mathcal{F}_t}^{\varphi,-}$  and  $\mathbf{Q}_{|\mathcal{F}_t}^-$  this is equivalent to

$$\mathbf{Q}^-(L_{T_{-1}} > l \text{ and } T_{-1} \leq t) > 0.$$

But this is well known. Thus the lemma is proved.  $\square$

**Lemma 5.8.** Let  $\varphi \in \Lambda^+$  be such that  $\mathbf{P}^{\varphi,+}(R) = 0$ . Then for any  $M > 0$ ,

$$\mathbf{P}_0^\varphi[L_\infty^0 < M] > 0.$$

*Proof.* We have

$$\begin{aligned} \mathbf{P}_0^\varphi[L_\infty^0 < M] &\geq \mathbf{P}_0^\varphi[L_{T_1}^0 < M \text{ and } X_t > 0 \quad \forall t > T_1] \\ &\geq \mathbf{E}_0^\varphi \left[ \mathbf{1}_{\{L_{T_1}^0 < M\}} \mathbf{P}_1^{\varphi T_1}(T_0 = \infty) \right]. \end{aligned}$$

But Lemma 5.7 implies that a.s.,  $\mathbf{P}_1^{\varphi T_1}(T_0 = \infty) > 0$ . So it remains to prove that  $\mathbf{P}_0^\varphi(L_{T_1}^0 < M) > 0$ . Like in the previous lemma, by absolute continuity, it suffices to prove that

$$\mathbf{Q}(L_{T_1}^0 < M) > 0.$$

But again this is well known. Thus the lemma is proved.  $\square$

**Remark 5.9.** This lemma holds as well for any  $\varphi \in \Lambda$ , such that  $\mathbf{P}_0^\varphi(R) = 0$ .

Finally we obtain the

**Theorem 5.10.** Let  $\varphi \in \Lambda_c^+$ . Then

$$\mathbf{P}^{\varphi,+}(R) = 1 \iff \int_0^\infty \varphi(0, u) du \leq 1.$$

*Proof.* We prove this for  $\varphi$  such that  $\varphi(x, l) = \varphi(0, l)$  for all  $x$ . By Lemma 5.4,  $\mathbf{E}_0^\varphi(D_\infty^0) \leq 1$ . But if  $\mathbf{P}^{\varphi,+}(R) = 1$ , then  $\mathbf{P}_0^\varphi$ -a.s.  $L_\infty^x = +\infty$ , for all  $x$ . So, by using the occupation time formula (see Lemma 2.2) we have  $\mathbf{E}_0^\varphi(D_\infty^0) = \int_0^\infty \varphi(0, u) du$ . This gives the necessary condition. Reciprocally, if  $\mathbf{P}^{\varphi,+}(R) = 0$ , we saw in Lemma 5.5 that  $\mathbf{E}_0^\varphi(D_\infty^0) = 1$ . But by Lemma 5.8, we have  $\mathbf{E}_0^\varphi(D_\infty^0) = \mathbf{E}_0^\varphi(h(0, L_\infty^0)) < \int_0^\infty \varphi(0, u) du$ , which gives the sufficient condition and concludes the proof of the theorem.  $\square$

Note that if  $\varphi \in \Lambda$  is such that for some  $a \in \mathbb{R}$ ,  $\varphi(x, l) = \varphi(a, l) \geq 0$  for all  $x \geq a$  and all  $l \geq 0$ , then

$$\mathbf{P}^{\varphi,+}(R) = 1 \iff \int_0^\infty \varphi(a, u) du \leq 1.$$

This can be proved using the fact that  $\mathbf{P}^{\varphi,+}(R) = \mathbf{P}^{\varphi_{a,+}}(R)$  which does not depend on  $\varphi(x, \cdot)$ , for  $x < a$ .

## 6. ADDITIONAL REMARKS FOR GENERAL $\varphi$

Our first remark concerns the question of transience. More precisely we could ask if it would be possible to have  $\mathbf{P}(X \rightarrow +\infty) = 1 - \mathbf{P}(X \rightarrow -\infty) \in (0, 1)$ . The answer is yes. For instance take  $\varphi \in \Lambda$  such that  $\mathbf{P}^{\varphi,+}(R) = 0$ , i.e. one has  $\mathbf{P}^{\varphi,+}$ -a.s.  $X \rightarrow +\infty$ . Then define  $\psi \in \Lambda$  by  $\psi(x, u) = \varphi(x, u)$  if  $x \geq 0$  and  $\psi(x, u) = -\varphi(x, u)$  if  $x < 0$ . Then by symmetry we have

$$\mathbf{P}_0^\psi(X \rightarrow +\infty) = \mathbf{P}_0^\psi(X \rightarrow -\infty) = 1/2.$$

More generally,

$$\mathbf{P}_0^\varphi(X \rightarrow +\infty) = \tilde{\mathbf{P}}^\varphi(L_\infty^1 < L_\infty^2),$$

and

$$\mathbf{P}_0^\varphi(X \rightarrow -\infty) = \tilde{\mathbf{P}}^\varphi(L_\infty^1 > L_\infty^2).$$

So, if  $L_\infty^1$  and  $L_\infty^2$  are finite random variables, since they are independent with support  $\mathbb{R}^+$ , these two probabilities are positive. Note that the previous section gives a criterion for these random variables to be finite when  $\varphi$  is nonnegative or nonpositive at infinity.

To summarize:  $\mathbf{P}_0^\varphi(R) = 1$  if there exists  $a > 0$  such that

- $\varphi(x, l) \leq 0$  for  $x \geq a$  and  $\varphi(x, l) \geq 0$  for  $x \leq -a$  or that;
- $\varphi(x, l) = \varphi(a, l) \geq 0$  for  $x \geq a$  and  $\varphi(x, l) \geq 0$  for  $x \leq -a$ , and  $\int_0^\infty \varphi(a, u) du \leq 1$ , or that;
- $\varphi(x, l) = \varphi(a, l) \geq 0$  for  $x \geq a$ ,  $\varphi(x, l) = \varphi(-a, l) \leq 0$  for  $x \leq -a$ ,  $\int_0^\infty \varphi(a, u) du \leq 1$  and  $\int_0^\infty \varphi(-a, u) du \geq -1$  or that;
- $\varphi(x, l) \leq 0$  for  $x \leq -a$  and  $\varphi(x, l) = \varphi(-a, l) \leq 0$  for  $x \leq -a$  and  $\int_0^\infty \varphi(0, u) du \geq -1$ .

Similarly  $\mathbf{P}_0^\varphi(R) = 0$  if there exists  $a > 0$  such that

- $\varphi(x, l) = \varphi(a, l) \geq 0$  for  $x \geq a$  with  $\int_0^\infty \varphi(a, u) du > 1$  or that;
- $\varphi(x, l) = \varphi(-a, l) \leq 0$  for  $x \leq -a$  with  $\int_0^\infty \varphi(-a, u) du < -1$ .

Our second remark gives a general necessary condition for recurrence.

**Proposition 6.1.** *Let  $\varphi \in \Lambda_c$ . If*

$$(8) \quad \int_0^{+\infty} |\varphi(0, u)| du < +\infty,$$

then

$$P_0^\varphi(R) = 1 \implies \int_0^\infty \varphi(0, u) \, du \in [-1, 1].$$

*Proof.* Let

$$D_t^{(1)} := \int_0^t \varphi \left( X_s^1, L_s^{(1), X_s^{(1)}} \right) \, ds = \int_0^\infty h(x, L_t^{(1), x}) \, dx,$$

be the total drift accumulated by  $X^{(1)}$  up to time  $t$ . By dominated convergence theorem, that we can apply by using (8), one get for all  $z \geq 0$ ,

$$z = \tilde{E}^\varphi [X_{T_z}^1] = \tilde{E}^\varphi [L_{T_z}^{(1)}] + \tilde{E}^\varphi [D_{T_z}^{(1)}].$$

So for all  $z \geq 0$ ,

$$\tilde{E}^\varphi [D_{T_z}^{(1)}] \leq z.$$

Since  $\varphi \in \Lambda_c$ , this implies

$$\tilde{E}^\varphi [D_\infty^{(1), 0}] \leq 1,$$

with evident notation. But if  $X$  is recurrent, then  $X^1$  is also recurrent and

$$D_\infty^{(1), 0} = \int_0^\infty \varphi(0, u) \, du,$$

which proves that

$$\int_0^\infty \varphi(0, u) \, du \leq 1.$$

The other inequality is obtained similarly by using the recurrence of  $X^2$ . This concludes the proof of the proposition.  $\square$

#### REFERENCES

- [ABK] **Amir G., Benjamini I, Kozma G.:** *Excited random walk against a wall*, Probab. Theory and Related Fields 140, (2008), 83–102.
- [BaS1] **Basdevant A.-L., Singh A.:** *On the speed of a cookie random walk*, Probab. Theory Related Fields 141, (2008), 625–645.
- [BaS2] **Basdevant A.-L., Singh A.:** *Rate of growth of a transient cookie random walk*, Electron. J. Probab. 13, (2008), 811–851.
- [BaS3] **Basdevant A.-L., Singh A.:** *Recurrence and transience of a multi-excited random walk on a regular tree*, arXiv:0803.3284.
- [BeLR] **Benaïm M., Ledoux, M., Raimond, O.:** *Self-interacting diffusions*, Probab. Theory Relat. Fields 122(1), 1–41 (2002).
- [BeR1] **Benaïm, M., Raimond, O.:** *Self-interacting diffusions II: convergence in law*, Ann. Inst. H. Poincaré Probab. Statist. 39(6), 1043–1055 (2003).
- [BeR2] **Benaïm, M., Raimond, O.:** *Self-interacting diffusions III: the symmetric case*, Ann. Probab. 33(5), 1717–1759 (2005).
- [BW] **Benjamini I., Wilson D. B.:** *Excited random walk*, Electron. Comm. Probab. 8 (electronic), (2003), 86–92.
- [CPY] **Carmona P., Petit F., Yor M.:** *Beta variables as times spent in  $[0, \infty[$  by certain perturbed Brownian motions*, J. London Math. Soc. (2) 58, (1998), 239–256.
- [CLJ] **Cranston, M., Le Jan, Y.:** *Self-attracting diffusions: two case studies*, Math. Ann. 303, 87–93 (1995).
- [D1] **Davis B.:** *Reinforced random walk*, Probab. Theory Related Fields 84, (1990), 203–229.
- [D2] **Davis B.:** *Weak limits of perturbed Brownian motion and the equation  $Y_t = B_t + \alpha \sup\{Y_s : s \leq t\} + \beta \inf\{Y_s : s \leq t\}$* , Ann. Probab. 24, (1996), 2007–2023.
- [HR] **Herrmann, S., Roynette, B.:** *Boundedness and convergence of some self-attracting diffusions*, Math. Ann. 325(1), 81–96 (2003).

- [KZer] **Kosygina E., Zerner M. P. W.:** *Positively and negatively excited random walks on integers, with branching processes*, preprint, arXiv:0801.1924.
- [Ko1] **Kozma G:** *Excited random walk in three dimensions has positive speed*, arXiv:math/0310305.
- [Ko2] **Kozma G:** *Excited random walk in two dimensions has linear speed*, arXiv:math/0512535.
- [Ku] **Kurtzmann A.:** *Asymptotic behavior of Self-interacting Diffusions on  $R^d$* , Ph.D. thesis, University of Neuchâtel, (2007).
- [Pem] **Pemantle R.:** *A survey of random processes with reinforcement*, Probab. Surv. 4 (electronic), (2007), 1–79.
- [PW] **Perman, M., Werner W.:** *Perturbed Brownian motions*, Probab. Theory Related Fields 108, (1997), 357–383.
- [RY] **Revuz D., Yor M.:** *Continuous martingales and Brownian motion*, Springer-Verlag, third ed. (1999).
- [TW] **Tóth B., Werner W.:** *The true self-repelling motion*, Probab. Theory Related Fields 111, (1998), 375–452.
- [V] **Volkov S.:** *Excited random walk on trees*, Electron. J. Probab. 8, (2003), 15 pp. (electronic).
- [W] **Werner W.:** *Some remarks on perturbed reflecting Brownian motion*, Sémin. Probab. XXIX, LNM 1613, Springer, Berlin, (1995), 37–43.
- [Zer1] **Zerner M. P. W.:** *Multi-excited random walks on integers*, Probab. Theory Relat. Fields 133, (2005), 98–122.
- [Zer2] **Zerner M. P. W.:** *Recurrence and transience of excited random walks on  $\mathbb{Z}_d$  and strips*, Electron. Comm. Probab. 11, (2006), 118–128 (electronic).

DÉPARTEMENT DE MATHÉMATIQUES, BÂT. 425, UNIVERSITÉ PARIS-SUD 11, F-91405 ORSAY, CEDEX, FRANCE.

*E-mail address:* `olivier.raimond@math.u-psud.fr`

*E-mail address:* `bruno.schapira@math.u-psud.fr`