



HAL
open science

Scaling approach to existence of long cycles in Casimir boxes

Mathieu Beau

► **To cite this version:**

| Mathieu Beau. Scaling approach to existence of long cycles in Casimir boxes. 2009. hal-00332745v2

HAL Id: hal-00332745

<https://hal.science/hal-00332745v2>

Preprint submitted on 25 Feb 2009 (v2), last revised 27 Apr 2009 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Scaling approach to existence of long cycles in Casimir boxes

Mathieu BEAU¹

Université de la Méditerranée and Centre de Physique Théorique - UMR
6207 ²,
Luminy - Case 907, 13288 Marseille, Cedex 09, France

Abstract

We analyse the concept of generalized Bose-Einstein condensation (g-BEC) known since 1982 for the perfect Bose gas (PBG) in Casimir boxes (anisotropic boxes) via the relation between concepts of long cycles and Off-Diagonal-Long-Range-Order (ODLRO) usually considered as an adequate way to describe the standard BEC on the ground state for the cubic box. First we show that these three criteria are equivalent in this case. Then we revise definitions of these concepts which are based on a scaling approach to prove that the classification of the g-BEC (in three types I,II,III) implies a hierarchy of long cycles (depending on their size) and a hierarchy of ODLRO (depending on the coherence length of the condensate).

Keywords: Generalized Bose-Einstein condensation, (Long) Cycles, Off-Diagonal-Long-Range-Order (ODLRO), perfect Bose gas, Casimir boxes, Scaling.

PACS: 05.30.Jp, 03.75.Hh

¹E-mail: mathieubeau@hotmail.fr

²Université de Provence - Aix-Marseille I, Université de la Méditerranée - Aix-Marseille II, Université du Sud - Toulon - Var, FRUMAM (FR 2291)

Contents

1	Introduction	2
2	BEC in Casimir boxes	5
2.1	About the concept of g-BEC condensation and the ODLRO	5
2.2	Feynman theory of cycles and existence of long cycles	8
3	Generalized BEC concept: revisited	10
3.1	Generalized condensation and scaled condensation	10
3.2	Scaling approach of the short/long cycles	12
3.3	Hierarchy of s-long cycles	13
4	Does generalized BEC of the type I, II, III imply a hierarchy of long cycles?	14
4.1	Generalized BEC in the case : $\alpha_1 < 1/2$	14
4.2	Generalized BEC in the case : $\alpha_1 = 1/2$	15
4.3	Generalized BEC in the case : $\alpha_1 > 1/2$	17
5	Does generalized BEC I, II, III imply a hierarchy of ODLRO?	19
5.1	Scaling approach to ODLRO	19
5.2	Hierarchy and anisotropy of ODLRO, coherence of the condensate	21
6	Concluding remarks	24

1 Introduction

About the Bose-Einstein condensation

The Bose-Einstein condensation known since 1925 and discovered in 4He in 1975 (see [Z-B] for historical remarks) attracts now a lot of attention of theoretical and mathematical physicists motivated by experiments with ultracold gases in traps [L-S-S-Y]. Recently experimentalists discovered a Bose-Einstein condensation in very anisotropic traps [M-H-U-B]. This may to imply in certain cases a *fragmentation* of the condensate. In fact this phenomenon was predicted long time ago by Casimir [C]. Then it was carefully studied by mathematical physicists and now it is known under the name of *generalized* Bose-Einstein condensation (g-BEC) *à la van den Berg-Lewis-Pulé* [Z-B]. After the first publication by van den Berg and Lewis in 1982 [vdB-L], a set of articles treated different cases of the Casimir box anisotropy, boundary conditions and external potentials, [vdB], [vdB-L], [vdB-L-P], [vdB-L-L]. They classified the Bose-Einstein condensation into three types. If a finite/infinite number of one-particle quantum states are macroscopically occupied, then the Bose gas manifests BEC of type I/II. If there are no states macroscopically occupied, although in the thermodynamical limit a macroscopic number of particles is accumulated in the ground (zero-mode) state, then the Bose gas manifests BEC of type III.

It is worthy to note that not only anisotropy, but also interaction between particles is able to modify the type of the condensate [Br-Z], [M-V].

In 1953 [F], R.Feynman introduced a concept of *cycles* by rewriting the partition function of the boson gas using the Bose-statistic. In the Feynman-Kac representation, cycles are close trajectories in the Wiener space. This is a mathematical framework for the path-integral formulation of the quantum statistical mechanics. Heuristically at low temperature (or at high density), bosons belong to cycles of different sizes and below some critical point, *infinite* cycles appear in the Feynman-Kac representation. Since Feynman it is a common wisdom to count this equivalent to the (type I) Bose-Einstein condensation [P-O]. For the perfect Bose gas it was proven in [S] and for some weakly interacting Bose gases only recently in [D-M-P], [U].

In 1956, L.Penrose and L.Onsager [P-O] introduced via a reduced density matrix, the concept of Off-Diagonal-Long-Range-Order (ODLRO) to give an alternative description of the Bose-Einstein condensation in ideal or interacting Bose-gas. They worked out convincing arguments that existence of the ODLRO is equivalent to the Bose-Einstein condensation. In fact the ODLRO measures in boson systems a correlation between two infinitely distant points. Recently a contact between the ODLRO and the existence of the long cycles studied in [U] and [U2].

Conceptual and physical problems

In spite of the ODLRO criterion is usually considered as criterion of (type I) BEC, it is not evident that ODLRO is equivalent to generalized BEC. Moreover, it is not evident that the presence of long cycles is equivalent to generalized BEC. Therefore, the present paper is motivated by two questions:

Are the different criteria of Bose-Einstein condensation (generalized BEC, long cycles and ODLRO) equivalent for the *perfect* Bose-gas in anisotropic (Casimir) boxes (Section 2)? Hence, the purpose of the Section 2 is an extension of known results on the long cycles, see A.Suto [S], T.Dorlas, P.Martin and J.Pulé [D-M-P], D.Ueltschi [U], to the Casimir boxes, where the concept of the generalized Bose-Einstein condensation can be explicitly verified.

The second question is how we can classify different types of generalized Bose-Einstein condensation with the help of the concept of cycles and the ODLRO (Sections 3, 4 and 5)?

The aim of this paper is to relate the classification of generalized Bose-Einstein condensation (type I, II and III) in anisotropic boxes to hierarchy of long cycles and the corresponding hierarchy of the ODLRO. More precisely, we would like to know, what is the scale of different sizes of the long cycles (macroscopic or not) and correlations (the coherence length) of the condensate? Our arguments are based on the scaling approach. To this end, we propose a scaling formulation for the condensate density and for the notion of long cycles (Section 3), as well as for the reduced density matrix and the ODLRO (Section 5).

Since the seventies [dG] scaling concepts used in polymer physics. In the present paper we adapt this scaling approach to the cycles (and two-point correlation function) because there is a deep analogy between cycle representation of boson systems and polymers

[C-S]. Below we give a mathematical presentation of scaling concepts concerning the Bose-Einstein condensation of Ideal Bose-Gas in Casimir boxes.

It is relevant in physics of coherent states, since it relates the box geometry constrains to the coherence shape of condensate clouds and to "geometry" of the boson cycles (polymers). Heuristically, there is a scaling relation between the coherence length r and the size of long cycles involved n bosons, which is given by $r = \lambda_\beta n^{1/2}$. Here $\lambda_\beta = \hbar\sqrt{2\pi\beta/m}$ is the thermal de Broglie length, provided correlation function has the form (5.52). This relation is analogous to the scaling law for the ideal polymer chain [dG], where the size of the chain R is proportional to the number of monomers N : $R = l_0 N^{1/2}$, where l_0 is the effective length of a monomer.

Results

This paper contains two kind of results interesting for physical properties of the perfect Bose-gas in Casimir boxes: $\Lambda = L_1 \times L_2 \times L_3$, with $L_\nu = V^{\alpha_\nu}$, $\nu = 1, 2, 3$, $\alpha_1 + \alpha_2 + \alpha_3 = 1$). These results establish a contact between generalized van den Berg-Lewis-Pulé condensate and experimental data concerning the fragmented BEC [M-H-U-B].

The first result is that the number of particles N_0 in the condensate for finite size system (N particles) is (see section 3.2):

$$N_0 = n_1 + n_2 + \dots + n_M,$$

where n_i are the numbers of particles in the condensate states, with $n_i = O(N)$ and $M = O(1)$, if $\alpha_1 \leq 1/2$ (see Theorem 4.1 and 4.2) or $n_i = O(N^\delta)$ and $M = O(N^{1-\delta})$ (such that $N_0 = O(N)$), $\delta = 2(1 - \alpha_1) < 1$, if $\alpha_1 > 1/2$ (see Theorem 4.3). This result can be related to the fragmentation theory of Bose-Einstein condensation [M-H-U-B]. The second part of this result (see section 4) is that the order of the size of long cycles is macroscopic (of the order $O(N)$) if we have generalized BEC of type I or II (i.e. $M = O(N)$) and that the order of the size of long cycles is microscopic size (of the order $O(N^\delta)$), see Theorem 4.1, 4.2 and 4.3.

The second result is that by virtue of Theorem 5.1 the two points correlation function (at different scales) has the form:

$$\begin{aligned} \lim_{V \uparrow \infty} \sigma_\Lambda(x - x') &= \rho_0(\beta), \text{ for } \|x - x'\| = O(V^{\alpha_1}), \text{ if } \alpha_1 < 1/2, \\ &= \sum_{n_1 \in \mathbb{Z}^1} \frac{\cos(\pi n y)}{\pi \lambda_\beta^2 n^2 + B}, \text{ for } \|x - x'\| = yV^{\alpha_1}, \text{ if } \alpha_1 = 1/2, \\ &= \rho_0(\beta) e^{-2y\sqrt{\pi C}/\lambda_\beta}, \text{ for } \|x - x'\| = yV^{\delta/2}, \text{ if } \alpha_1 > 1/2, \end{aligned}$$

where $\rho_0(\beta)$ is the density of particles in the condensate, B and C are two positive constants respectively given by (2.8) and (2.9). This form implies that the order of the *condensate coherence length* coincides with the size of the box in the case of generalized BEC of type I or II (i.e. long cycles of macroscopic size), but it is smaller if we have generalized BEC of type III (i.e. long cycles of microscopic size). The last case shows decreasing of the coherence length for the elongated condensate. Notice that existence of this phenomenon is indicated in physical literature [P-S-W].

To make a contact with experiments involved the cold atoms confined in magnetic traps one should extend these results to the Bose-gas in a weak harmonic potential. So, an important question is theoretical and experimental study of very anisotropic systems (like quasi-2D or quasi-1D systems) to understand coherence properties of condensate, which mimics the Casimir case.

2 BEC in Casimir boxes

In this section we give a synthetic review of the different concepts around Bose-Einstein condensation such as occupation number, generalized BEC (g-BEC), cycles, long cycles, reduce density matrix, Off-Diagonal-Long-Range-Order (ODLRO) and the link between the different criterions of BEC (g-BEC, long cycles and ODLRO).

2.1 About the concept of g-BEC condensation and the ODLRO

Basic notions

Let us consider the grand-canonical (β, μ) perfect Bose-gas (PBG) in Casimir boxes $\Lambda = L_1 \times L_2 \times L_3 \in \mathbb{R}^{d=3}$, of volume $|\Lambda| = V$ with sides length $L_\nu = V^{\alpha_\nu}$, $\nu = 1, 2, 3$, where $\alpha_1 \geq \alpha_2 \geq \alpha_3 > 0$, $\alpha_1 + \alpha_2 + \alpha_3 = 1$. For the single-particle hamiltonian $H_\Lambda^{(N=1)} = T_\Lambda^{(1)} := -(\hbar^2/2m)\Delta$ with periodic boundary conditions, we get the dual vector-spaces Λ^* defined by:

$$\Lambda^* = \left\{ k \in \mathbb{R}^3 : k = \left(\frac{2\pi n_1}{V^{\alpha_1}}, \frac{2\pi n_2}{V^{\alpha_2}}, \frac{2\pi n_3}{V^{\alpha_3}} \right); n_\nu \in \mathbb{Z}^1 \right\}. \quad (2.1)$$

In the grand-canonical ensemble, the mean values of the k -mode particle densities $\{\rho_\Lambda(k)\}_{k \in \Lambda^*}$ are:

$$\rho_\Lambda(k) := \frac{1}{V} \langle N_\Lambda(k) \rangle_\Lambda(\beta, \mu) = \frac{1}{V} \frac{1}{e^{\beta(\epsilon_\Lambda(k) - \mu)} - 1}, \quad (2.2)$$

where $\epsilon_\Lambda(k) = \hbar^2 k^2 / 2m$, $k \in \Lambda^*$ are the eigenvalue of the Laplacian with periodic boundary conditions, $\langle N_\Lambda(k) \rangle_\Lambda(\beta, \mu)$ is the Gibbs expectation of the particles number operator $N_\Lambda(k)$ in the mode k . The total density of particles is $\rho_\Lambda(\beta, \mu) := \sum_{k \in \Lambda^*} \rho_\Lambda(k)$

Let us recall that the eigenfunctions $\{\Psi_{\Lambda,k}^{(N=1)}(x)\}_{k \in \Lambda^*}$ of the single particle hamiltonian $T_\Lambda^{(1)}$ are:

$$\Psi_{\Lambda,k}^{(1)}(x) = \frac{1}{\sqrt{V}} e^{ik \cdot x}. \quad (2.3)$$

London scaling approach and g-BEC

In fact it was F.London [L] who for the first time used implicitly the *scaling* approach to solve the controversy between the G.Uhlenbeck mathematical arguments against condensation of the perfect Bose-gas for high densities and Einstein's intuitive reasoning in favour of this phenomenon. His line of reasoning was based on the following observations: since the explicit formula for the total particle density $\rho_\Lambda(\beta, \mu)$ in the box Λ is known only in grand-canonical ensemble (β, μ) , to ensure fixed density ρ in this box one has first to solve equation $\rho = \rho_\Lambda(\beta, \mu)$ which determine the corresponding value of the chemical potential $\bar{\mu}_\Lambda := \bar{\mu}_\Lambda(\beta, \rho)$.

Then the van den Berg-Lewis-Pulé formulation of the g-BEC concept in Casimir boxes gets the form:

Definition 2.1 *We say that for the grand-canonical PBG manifests the g-BEC for a fixed total density of particle ρ , if one has:*

$$\rho_0(\beta, \rho) := \lim_{\epsilon \downarrow 0} \lim_{V \uparrow \infty} \sum_{\{k \in \Lambda^* : \|k\| \leq \epsilon\}} \rho_\Lambda(k) > 0, \quad (2.4)$$

where $\rho_\Lambda(k)$ are defined by (2.2) for $\mu = \bar{\mu}_\Lambda(\beta, \rho)$.

This motivates the following classification:

Definition 2.2 • *One gets g-BEC of type I if a finite number of the single-particle states are macroscopically occupied.*

• *There is g-BEC of type II is an infinite (countable) number of the single-particle states are macroscopically occupied.*

• *The g-BEC is called type III, if none of the single-particle state is macroscopically occupied, but $\rho_0(\beta, \rho) > 0$.*

As usually one introduces for the PBG the critical density, $\rho_c(\beta)$ defined by:

$$\rho_c(\beta) := \sup_{\mu < 0} \lim_{V \uparrow \infty} \rho_\Lambda(\beta, \mu) = g_{3/2}(1)/\lambda_\beta^3, \quad (2.5)$$

where $g_s(z) := \sum_{j=1}^{\infty} z^j/j^s$ is related to the Riemann zeta-function $\zeta(s) := g_s(1)$. Here $\lambda_\beta = \hbar\sqrt{2\pi\beta/m}$ is the thermal de Broglie length.

The following proposition is due to [vdB-L-P]:

Proposition 2.1 *For particles densities $\rho < \rho_c(\beta)$ there is no g-BEC of the PBG in Casimir boxes and the chemical potential $\mu = \bar{\mu}(\beta, \rho)$ is a unique solution of equation:*

$$\rho = g_{3/2}(e^{\beta\mu})/\lambda_\beta^3 \quad (2.6)$$

Let $1/2 > \alpha_1$, then for a fixed particles density $\rho > \rho_c(\beta)$ the chemical potential $\bar{\mu}_\Lambda = -A/\beta V + o(1/V)$, with $A > 0$ and there is g-BEC of the type I in the single zero-mode $k = 0$. Here A is a solution of equation :

$$\rho - \rho_c(\beta) = \frac{1}{A}. \quad (2.7)$$

If $1/2 = \alpha_1$, then the chemical potential $\bar{\mu}_\Lambda = -B/\beta V + o(1/V)$, with $B > 0$ and one gets the g-BEC of type II in the infinite number of modes:

$$\begin{aligned} \lim_{V \uparrow \infty} \rho_\Lambda(k) &= \frac{1}{B + \pi\lambda_\beta^2 n_1^2}, \text{ for } k = (2\pi n_1/V^{\alpha_1}, 0, 0), \ n_1 \in \mathbb{Z}^1, \\ &= 0, \text{ for } k \neq (2\pi n_1/V^{\alpha_1}, 0, 0), \ n_1 \in \mathbb{Z}^1. \end{aligned}$$

Here B is a solution of equation:

$$\rho - \rho_c(\beta) = \sum_{n_1 \in \mathbb{Z}^1} \frac{1}{B + \pi \lambda_\beta^2 n_1^2} . \quad (2.8)$$

If $1/2 < \alpha_1$, then the chemical potential $\bar{\mu}_\Lambda = -C/\beta V^\delta + o(1/V^\delta)$, with $\delta = 2(1 - \alpha_1)$ and $C > 0$. The corresponding g-BEC is of the type III: for all $k \in \Lambda^*$ we have $\lim_{V \uparrow \infty} \rho_\Lambda(k) = 0$, although $\rho_0(\beta, \rho) > 0$ and C is a solution of equation:

$$\rho - \rho_c(\beta) = \frac{\sqrt{\pi}}{\lambda_\beta C^{1/2}} . \quad (2.9)$$

Reduce density matrix and ODLRO

Definition 2.3 We say that the PBG manifests for a fixed total density of particles ρ an ODLRO, if gets a nontrivial limit:

$$\sigma(\beta, \rho) := \lim_{\|x-x'\| \uparrow \infty} \lim_{V \uparrow \infty} \sigma_\Lambda(\beta, \rho; x, x') > 0 , \quad (2.10)$$

where $\sigma(\beta, \rho; x, x') := \lim_{V \uparrow \infty} \sigma_\Lambda(\beta, \rho; x, x')$ is the two-point correlation function, defined by:

$$\sigma_\Lambda(\beta, \rho; x, x') := \sum_{k \in \Lambda^*} \langle N_\Lambda(k) \rangle_\Lambda(\beta, \rho) \Psi_{\Lambda, k}^{(1)}(x) \Psi_{\Lambda, k}^{(1)}(x')^* = \sum_{k \in \Lambda^*} \rho_\Lambda(k) e^{ik \cdot (x-x')} , \quad (2.11)$$

for periodic boundary conditions. Here $\rho_\Lambda(k)$ is defined by (2.2).

Recall that according to Definition 2.1 for a non-zero BEC of types I and II we obtain in the thermodynamical limit a nontrivial particle density in the mode null, i.e. particle density with de Broglie wave-length equals infinity. This allows a *communication* in the condensate through the whole space (in \mathbb{R}^3). It is less evident for the g-BEC of the type III, but the heuristic arguments can be checked showing that the g-BEC and the ODLRO are equivalent for the PBG in Casimir boxes:

Theorem 2.1 Consider the PBG in Casimir boxes Λ . Then:

$$\begin{aligned} \sigma(\beta, \rho) &= 0, \text{ for } \rho < \rho_c, \\ &= \rho - \rho_c(\beta), \text{ for } \rho > \rho_c, \end{aligned} \quad (2.12)$$

i.e. the ODLRO is non-zero if and only if there is g-BEC.

Proof

We split the correlation function in two parts, the first one is the correlation due to the (future) condensate and the second one corresponds to the particles outside the condensate:

$$\begin{aligned} \sigma(\beta, \rho; x, x') &:= \lim_{\epsilon \downarrow 0} \lim_{V \uparrow \infty} \sum_{k \in \{\Lambda^*: \|k\| \leq \epsilon\}} \rho_\Lambda(k) e^{ik \cdot (x-x')} + \lim_{\epsilon \downarrow 0} \lim_{V \uparrow \infty} \sum_{k \in \{\Lambda^*: \|k\| > \epsilon\}} \rho_\Lambda(k) e^{ik \cdot (x-x')} \\ &= \lim_{\epsilon \downarrow 0} \lim_{V \uparrow \infty} \sum_{k \in \{\Lambda^*: \|k\| \leq \epsilon\}} \rho_\Lambda(k) + \frac{1}{(2\pi)^3} \int_{k \in \mathbb{R}^3} dk \frac{e^{ik \cdot (x-x')}}{e^{\beta(\hbar^2 k^2 / 2m - \bar{\mu}(\beta, \rho))} - 1}, \end{aligned}$$

since $\forall k \in \Lambda^* : \|k\| \leq \epsilon, k \cdot (x - x') \rightarrow 0$ when $\epsilon \rightarrow 0$. Here $\bar{\mu}(\beta, \rho) := \lim_{V \uparrow \infty} \bar{\mu}_\Lambda(\beta, \rho)$.

Then by virtue of Proposition 2.1 and by the Riemann-Lebesgue Theorem we obtain the result. \square

Since Penrose and Onsager [P-O], the ODLRO is known as the most relevant criterion of condensation because it is valid with or without interactions between particles. Here we established that ODLRO is not equivalent to the usual criterion of BEC (macroscopic occupation of the ground state) but to the generalized BEC. Thus for the PBG a *true* criterion of condensation is the generalized BEC. It is natural to suppose the same interacting Bose-gases.

2.2 Feynman theory of cycles and existence of long cycles

Feynman concept of cycles

Here we recall the Feynman concept of cycles introduced in 1953 [F]. It is related to the Bose statistic and the Feynman-Kac representation for partition functions.

Recall that for the PBG in boxes $\Lambda = L_1 \times L_2 \times L_3 \subset \mathbb{R}^3$ with *periodic* boundary conditions, (i.e. dual vector-spaces $\Lambda^* := \{k \in \mathbb{R}^3 : k = (2\pi n_1/L_1, 2\pi n_2/L_2, 2\pi n_3/L_3), n_\nu \in \mathbb{Z}^1\}$, see (2.1)) the grand-canonical pressure has the form:

$$\begin{aligned} p_\Lambda(\beta, \mu) &= \frac{1}{\beta|\Lambda|} \log(\Xi_\Lambda^0(\beta, \mu)) = \frac{1}{\beta|\Lambda|} \sum_{k \in \Lambda^*} \ln \left[(1 - e^{-\beta(\epsilon_\Lambda(k) - \mu)})^{-1} \right] \\ &= \frac{1}{\beta|\Lambda|} \sum_{j=1}^{\infty} \frac{1}{j} e^{j\beta\mu} \text{Tr}_{\mathcal{H}_\Lambda^{(1)}} \left(e^{-j\beta T_\Lambda^{(1)}} \right) , \end{aligned} \quad (2.13)$$

where $\Xi_\Lambda^0(\beta, \mu)$ is the PBG grand-canonical partition function and we used $\text{Tr}_{\mathcal{H}_{\Lambda,1}} \left(e^{-j\beta T_\Lambda^{(1)}} \right) = \sum_{k \in \Lambda^*} e^{-j\beta \epsilon_\Lambda(k)}$.

The Feynman-Kac representation naturally appears from (2.13) if we consider the representation of the trace of the *Gibbs semigroup*: $\left\{ e^{-\beta T_\Lambda^{(1)}} \right\}_{\beta > 0}$, via its kernel $K_\Lambda(\beta; x, x') = \left(e^{-\beta T_\Lambda^{(1)}} \right) (x, x')$:

$$\text{Tr}_{\mathcal{H}_\Lambda^{(1)}} \left(e^{-\beta T_\Lambda^{(1)}} \right) = \int_\Lambda dx K_\Lambda(\beta; x, x) . \quad (2.14)$$

It is known [G] that the kernel $K_\Lambda(\beta; x, x')$ can be represented as a path integral over Wiener trajectories starting at point x and finishing at point x' . Thus $K_\Lambda(\beta; x, x)$ can be represent as a Wiener integral over close trajectories (loops) starting and finishing at the same point. The order of the size of the trajectories coincides with the size of the quantum fluctuations λ_β , known as the thermal de Broglie length [M].

By virtue of the Gibbs semigroup properties and by expressions (2.13), (2.14), we get:

$$p_\Lambda(\beta, \mu) = \sum_{j=1}^{\infty} \frac{1}{j} e^{j\beta\mu} \int_\Lambda dx_1 \int_\Lambda dx_2 \dots \int_\Lambda dx_j K_\Lambda(\beta; x_1, x_2) K_\Lambda(\beta; x_2, x_3) \dots K_\Lambda(\beta; x_j, x_1) , \quad (2.15)$$

In (2.15) each integral over $(x_1, \dots, x_j) \in \Lambda^j$ correspond to the impact of the Wiener's loops of the length $j\lambda_\beta$ [G], [M].

Notice that the total density of particles in the grand-canonical ensemble is given by:

$$\rho_\Lambda(\beta, \mu) := \frac{\langle N_\Lambda \rangle_\Lambda(\beta, \mu)}{V} = \partial_\mu p_\Lambda(\beta, \mu) = \frac{1}{V} \sum_{j=1}^{\infty} e^{j\beta\mu} \text{Tr}_{\mathcal{H}_\Lambda^{(1)}} e^{-j\beta T_\Lambda^{(1)}} . \quad (2.16)$$

One can use this representation to identify the *repartition* of the total density (2.16) over densities of particles involved into loops of the length $j\lambda_\beta$:

$$\rho_{\Lambda,j}(\beta, \mu) := \frac{1}{V} e^{j\beta\mu} \text{Tr}_{\mathcal{H}_\Lambda^{(1)}} e^{-j\beta T_\Lambda^{(1)}} . \quad (2.17)$$

The j -loops particles density (2.17) and the representation (2.15) are the key notions for the concept of the *short/long* cycles. Indeed, after the thermodynamical limit one can obtain loops of *finite* sizes or *infinite* sizes, i.e. one can relate the BEC to appearance of loops of the infinite size as an explanation of the long-range order and the macroscopic size of the quantum fluctuations [F], [P-O], [U]. The mathematical basis of the Feynman's cycles approach is related to the boson permutation group, see [D-M-P],[U], [S] and [M].

BEC and the concept of short/long cycles

Definition 2.4 *We say that the representation (2.16) for the grand-canonical PBG contains only short cycles if:*

$$\rho_{\text{short}}(\beta, \mu) := \lim_{M \rightarrow \infty} \left\{ \lim_{V \uparrow \infty} \sum_{j=1}^M \rho_{\Lambda,j}(\beta, \mu) \right\} = \rho(\beta, \mu) := \lim_{V \uparrow \infty} \rho_\Lambda(\beta, \mu) , \quad (2.18)$$

i.e. it coincides with the total particle density. Since in general the limits in (2.18) are not interchangeable, we say that for a given total particle density ρ the representation (2.16) contains macroscopic number of long cycles, if $\rho > \rho_{\text{short}}(\beta, \bar{\mu}(\beta, \rho))$, or equivalently if:

$$\rho_{\text{long}}(\beta, \rho) := \lim_{M \rightarrow \infty} \left\{ \lim_{V \uparrow \infty} \sum_{j=M+1}^{\infty} \rho_{\Lambda,j}(\beta, \bar{\mu}_\Lambda(\beta, \rho)) \right\} > 0 . \quad (2.19)$$

Since Feynman [F] the presence of the non-zero density of the long cycles is usually connected with the existence of zero-mode BEC, but a rigorous proof of this conjecture for a certain class of models has been obtained only recently. There we noticed that even for the PBG type I BEC the mathematical proof of this connection was not straightforward and appealed to a non-trivial analysis, see [S], [D-M-P],[U].

Theorem 2.2 *Let us consider a PBG in Casimir boxes Λ . Then we have:*

$$\begin{aligned} \rho_{\text{long}}(\beta, \rho) &= 0, \text{ if } \rho < \rho_c(\beta) , \\ &= \rho - \rho_c(\beta), \text{ if } \rho > \rho_c(\beta) , \end{aligned} \quad (2.20)$$

where $\rho_c(\beta)$ is the critical density defined by (2.5).

Proof: By virtue of (rhoj0) and (2.18) we have:

$$\begin{aligned}\rho_{\text{short}}(\beta, \mu) &= \lim_{M \rightarrow \infty} \left\{ \lim_{V \uparrow \infty} \sum_{j=1}^M e^{j\beta\mu} \prod_{\nu=1}^3 \frac{1}{V^{\alpha_\nu}} \sum_{n_\nu \in \mathbb{Z}^1} e^{-j\beta\pi\lambda_\beta^2 (n_\nu/V^{\alpha_\nu})^2} \right\} \\ &= \sum_{j=1}^{\infty} e^{j\beta\mu} \prod_{\nu=1}^3 \int_{\mathbb{R}} d\xi_\nu e^{-j\beta\pi\lambda_\beta^2 \xi_\nu^2} = \sum_{j=1}^{\infty} \frac{e^{\beta\mu}}{\lambda_\beta^3 j^{3/2}} =: \frac{g_{3/2}(e^{\beta\mu})}{\lambda_\beta^3},\end{aligned}$$

c.f. definition of $g_s(z)$. By virtue of Proposition 2.1, if $\rho < \rho_c(\beta)$ we have $\rho_{\text{short}}(\beta, \rho) = \rho$ and if $\rho > \rho_c(\beta)$ we have $\rho_{\text{short}}(\beta, \rho) = \rho_c(\beta)$. So, by Definition 2.4 we conclude the proof. \square

Intuitively the size of long cycles for usual BEC (ground state macroscopic occupation) are of the order of the total particle number ($j = O(N)$). What happens if the condensate is fragmented, i.e. of the type II or III? Below we apply a scaling approach to study these cases and we prove in Theorem 4.3 that the order of the size of long cycles is smaller than the number of particles ($O(N^\delta)$, $\delta < 1$).

3 Generalized BEC concept: revisited

In this section we propose a modification of the concept of generalized BEC, which we call *scaled* BEC (s-BEC). This implies the corresponding modification of the concept for cycles (*scaled* short/long cycles denoted s-short/s-long cycles) via similar scaling arguments. Moreover, we introduce a classification of the s-BEC (type I, II, III) and the *hierarchy* of the s-long cycles, which distinguishes long-microscopic/macrosopic cycles.

3.1 Generalized condensation and scaled condensation

The original van den Berg-Lewis-Pulé concept of the g-BEC [vdB-L-P] was not explicitly addressed to detect a *fine* structure of the condensate: *a priori* it does not allow to make the distinction between generalized BEC of *types* I, II or III. In fact one can do this analysis, as it was done for the first time in [vdB-L] for the case of the Casimir boxes.

To make this facet more evident we introduce here a *new* definition of the *generalized* BEC, called a *scaled* BEC (s-BEC). Take for simplicity the PBG in Casimir boxes Λ with *periodic* boundary conditions, i.e. with the dual vector-spaces Λ^* defined by (2.1), and with the k -modes mean particle densities $\{\rho_\Lambda(k)\}_{k \in \Lambda}$ defined by (2.2).

Definition 3.1 *We say that for a fixed total density ρ the (perfect) Bose gas manifests a s-BEC in boxes Λ , if there exists a positive decreasing function $\eta : V \mapsto \mathbb{R}_+$, such that $\lim_{V \uparrow \infty} \eta(V) = 0$ and we have:*

$$\rho_\eta(\beta, \rho) := \liminf_{V \uparrow \infty} \sum_{\{k \in \Lambda^* : \|k\| \leq \eta(V)\}} \rho_\Lambda(k) > 0. \quad (3.21)$$

Remark 3.1 *Recall that the van den Berg-Lewis-Pulé definition of g-BEC is formulated as:*

$$\rho_0(\beta, \rho) := \lim_{\epsilon \downarrow 0} \lim_{V \uparrow \infty} \sum_{\{k \in \Lambda^* : \|k\| \leq \epsilon\}} \rho_\Lambda(k) > 0. \quad (3.22)$$

Hence, the two Definitions 2.1 and 3.1 are evidently not equivalent. Moreover, we show that our Definition 3.1 allows also to connect a fine mode-structure of the condensate of types I, II or III with the long-cycles hierarchy, and to show that there is a relation between the structure of the condensate and the size of cycles.

The following statement is an evident consequence of Definitions 2.1 and 3.1:

Lemma 3.1 For any function $\eta(V)$ we have:

$$0 \leq \rho_\eta(\beta, \rho) \leq \rho_0(\beta, \rho) . \quad (3.23)$$

This lemma means that s-BEC implies g-BEC and (i.e. that if there is no g-BEC, then there is no s-BEC).

A simple example of application of the s-BEC approach is the possibility to distinguish the type I, II or III condensations of the PBG in Casimir boxes Λ with *periodic* boundary conditions.

Proposition 3.1 The rate $\eta(V) = O(1/V^{1/2})$ is an important threshold to refine a discrimination between different types of g-BEC. If for example, one takes $\eta_\delta(V) = 2\pi/V^{(1/2-\delta)}$ such that $\delta > 0$, then we obtain:

$$\begin{aligned} \rho_{\eta_\delta}(\beta, \rho) &= \rho_0(\beta, \rho) , & \text{for } \alpha_1 \leq 1/2 , \\ \rho_{\eta_\delta}(\beta, \rho) &= 0 , & \text{for } \alpha_1 > 1/2 + \delta , \\ \rho_{\eta_\delta}(\beta, \rho) &= \rho_0(\beta, \rho) , & \text{for } 1/2 + \delta > \alpha_1 > 1/2 . \end{aligned}$$

On the other hand, for $\alpha_1 = 1/2$ and $\eta_\Gamma(V) := 2\pi\Gamma/V^{1/2}$ one gets a modification of the density of the type II condensation:

$$\rho_{\eta_\Gamma}(\beta, \rho) = \sum_{|n_1| \leq \Gamma} \frac{1}{\lambda_\beta^2 n_1^2 + B} < \rho_0(\beta, \rho) .$$

For $\alpha_1 > 1/2$ and $\eta_{\Gamma'}(V) := 2\pi\Gamma'/V^{1-\alpha_1}$ one gets a modification of the density of the type III condensation:

$$\rho_{\eta_{\Gamma'}}(\beta, \rho) = \int_{\mathbb{R}^+} d\zeta \frac{e^{-C\zeta}}{\zeta^{1/2} \lambda_\beta} \text{erf}(\Gamma' \lambda_\beta \zeta / \sqrt{4\pi}) < \rho_0(\beta, \rho) ,$$

where $\text{erf}(\cdot)$ stands for error function and where C is the unique solution of equation (2.9).

Proof: By virtue of the proof of the Theorems 4.1, 4.2 and 4.3, with the different choices of $\eta(V)$, we can obtain the results. \square

Note that the scaling criterion of condensation (Definition 3.1) is more adapted for physical description of BEC because a judicious choice of the function $\eta(V)$ may give a support of momentum distribution of the condensate. It also could be useful for numerical analysis of experimentally observed *fragmented* condensates [M-H-U-B].

If N_0 is the number of particle in the condensate then one can say that there is fragmentation of first type if $N_0 = n_1 + n_2 + \dots + n_M$ with $n_i = O(N)$ and $M = O(1)$, (where n_i are the numbers of particles in the condensate states), or of the second type if $n_i = O(N^\delta)$ and $M = O(N^{1-\delta})$, $\delta < 1$ (such that $N_0 = O(N)$).

3.2 Scaling approach of the short/long cycles

First we introduce the concepts of *scaled* short/long cycles (s-short/long cycles):

Definition 3.2 *We say that Bose-gas manifests s-long cycles, if there exists a positive increasing function of the volume $\lambda : \mathbb{R}^+ \rightarrow \mathbb{N}^+$, such that $\lim_{V \uparrow \infty} \lambda(V) = \infty$ and:*

$$\rho_{\text{long},\lambda}(\beta, \rho) := \liminf_{V \uparrow \infty} \sum_{j \geq \lambda(V)} \rho_{\Lambda,j}(\beta, \bar{\mu}_{\Lambda}(\beta, \rho)) > 0, \quad (3.24)$$

where $\rho_{\Lambda,j}(\beta, \mu)$ is given by (2.17).

The following statement is an evident consequence of Definitions 2.4 and 3.2:

Lemma 3.2 *In the particular case of PBG in Casimir boxes we have:*

$$0 \leq \rho_{\text{long},\lambda}(\beta, \rho) \leq \rho_{\text{long}}(\beta, \rho), \quad (3.25)$$

for any function $\lambda(V)$. Here $\rho_{\text{long}}(\beta, \rho)$ is given by (2.19).

This lemma implies that the presence of s-long cycles imply the presence of the long cycles (c.f. Definition 2.4).

A simple example of application of the s-long cycles approach is a possibility to distinguish the type I, II or III condensations of the PBG in Casimir boxes Λ with *periodic* boundary conditions.

Proposition 3.2 *If $\lambda(V) = V^\delta$ then for $\rho \geq \rho_c(\beta)$ we obtain:*

$$\begin{aligned} \rho_{\text{long},\lambda}(\beta, \rho) &= 0, \text{ for } \delta > 1, \\ \rho_{\text{long},\lambda}(\beta, \rho) &= \rho_0(\beta, \rho), \text{ for } \alpha_1 \leq 1/2, \ 0 < \delta < 1, \\ \rho_{\text{long},\lambda}(\beta, \rho) &= 0, \text{ for } \alpha_1 > 1/2, \ 2(1 - \alpha_1) < \delta. \\ \rho_{\text{long},\lambda}(\beta, \rho) &= \rho_0(\beta, \rho), \text{ for } \alpha_1 > 1/2, \ 0 < \delta < 2(1 - \alpha_1), \end{aligned}$$

Proof: Adapting the proof of the Theorems 4.1, 4.2 and 4.3 with the different choices of δ for $\lambda(V) = V^\delta$ we can obtain the results. \square

Definition 3.3 *If $j : \mathbb{R}^+ \rightarrow \mathbb{N}^+$ is a bounded positive increasing function of the volume, i.e. $\lim_{V \uparrow \infty} j(V) < \infty$, then $\rho_{\Lambda,j(V)}(\beta, \bar{\mu}_{\Lambda}(\beta, \rho))$ is the density of particles in the s-short-cycles of size $j(V)$.*

Definition 3.4 *If $j : \mathbb{R}^+ \rightarrow \mathbb{N}^+$ is a positive increasing function of the volume such as $\lim_{V \uparrow \infty} j(V) = \infty$, then $\rho_{\Lambda,j(V)}(\beta, \bar{\mu}_{\Lambda}(\beta, \rho))$ is the density of particles in the s-long-cycles of size $j(V)$.*

There is a natural classification of s-long-cycles:

- if $\lim_{V \uparrow \infty} (j(V)/V) = 0$, we say that $\rho_{\Lambda,j(V)}(\beta, \bar{\mu}_{\Lambda}(\beta, \rho))$ is the density of particles in the microscopic-long-cycles of size $j(V)$,
- if $0 < \lim_{V \uparrow \infty} (j(V)/V) < \infty$, we say that $\rho_{\Lambda,j(V)}(\beta, \bar{\mu}_{\Lambda}(\beta, \rho))$ is the density of particles in the macroscopic-cycles of size $j(V)$
- if $\lim_{V \uparrow \infty} (j(V)/V) = \infty$, we say that $\rho_{\Lambda,j(V)}(\beta, \bar{\mu}_{\Lambda}(\beta, \rho))$ is the density of particles in the large-cycles of size $j(V)$.

To clarify Definition 3.4, we make the following remark:

Remark 3.2 *We say that if $j(V)$ in (2.17) is of the order V^α if $0 < \lim_{V \uparrow \infty} (j(V)/V^\alpha) < \infty$, for example $j(V) = xV^\alpha$, $x > 0$. If $\alpha < 1$, we are in the first case of the classification in Definition 3.4. This case is important because there is a question: can we have a macroscopic quantity of particles in the microscopic-long-cycles? If $\alpha = 1$ we are in the second case of the classification of s-long-cycles and if $\alpha > 1$ we are in the third case. Of course, one can take above any increasing function $j(V)$ including e.g. $\ln(V)$.*

3.3 Hierarchy of s-long cycles

Definitions 3.3, 3.4 and the Remark 3.2 allow to give a natural classification of scaled-long (s-long) cycles. We call this classification a *hierarchy* of cycles, ordered according their size.

In general there are long cycles of any size:

Definition 3.5 *We say that the Bose gas manifests the s-long cycles of the order $\lambda(V)$ where $\lambda : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a positive increasing function of the volume, if there exist two positive real numbers x and y such as the s-long cycles particle density:*

$$\lim_{V \uparrow \infty} \sum_{j=x\lambda(V)}^{y\lambda(V)} \rho_{\Lambda,j}(\beta, \bar{\mu}_\Lambda(\beta, \rho)) > 0.$$

Then the total density of particles in cycles of size of the order $\lambda(V)$ is:

$$\rho_{\text{long}}(\beta, \rho | \lambda) := \lim_{x \downarrow 0; y \uparrow \infty} \lim_{V \uparrow \infty} \sum_{j=x\lambda(V)}^{y\lambda(V)} \rho_{\Lambda,j}(\beta, \bar{\mu}_\Lambda(\beta, \rho)).$$

We introduce a classification of s-long cycles:

Definition 3.6 - *We say that the Bose-gas manifests macroscopic-cycles, if the Bose gas manifests the s-long cycles of the order V .*

- *We say that the Bose-gas manifests long-microscopic-cycles, if the Bose gas manifests the s-long cycles of orders smaller than V .*

In the next section we see that in the particular case of the PBG in Casimir boxes, the classification of s-BEC induces a hierarchy in classification of s-long cycles. By virtue of Theorem 4.1, 4.2 and 4.3 we obtain:

Proposition 3.3 *If x and y are two positive real numbers, then we have:*

$$\begin{aligned} \lim_{\Lambda} \sum_{j=xV^\delta}^{yV^\delta} \rho_{\Lambda,j}(\beta, \bar{\mu}_\Lambda(\beta, \rho)) &= (e^{-xA} - e^{-yA})\rho_0(\beta, \rho), \text{ for } \alpha_1 < 1/2, \delta = 1, \\ &= (e^{-xB} - e^{-yB})\rho_0(\beta, \rho), \text{ for } \alpha_1 = 1/2, \delta = 1, \\ &= (e^{-xC} - e^{-yC})\rho_0(\beta, \rho) \text{ for } \alpha_1 > 1/2, \delta = 2(1 - \alpha_1). \end{aligned}$$

where A is the unique solution of the equation (2.7), B is the unique solution of the equation (2.8) and C is the unique solution of the equation (2.9).

This proposition gives an illustration of the hierarchy of cycles, in the first case ($\alpha_1 < 1/2$) and second case ($\alpha_1 = 1/2$) the Bose gas manifests presence of macroscopic-cycles, in the third case ($\alpha_1 > 1/2$), where the Bose gas manifests presence of long-microscopic-cycles of the order $V^{2(1-\alpha_1)}$. A link with the results given by Proposition 2.1 we are going to discuss in details in the next Section 3.

4 Does generalized BEC of the type I, II, III imply a hierarchy of long cycles?

The aim of this section is to relate a fine structure of g-BEC and s-BEC (type I, II or III) with hierarchy of long cycles. Recall that we deal with Casimir boxes $\Lambda = V^{\alpha_1} \times V^{\alpha_2} \times V^{\alpha_3}$, $\alpha_1 \geq \alpha_2 \geq \alpha_3 > 0$, $|\Lambda| = V$, ($\alpha_1 + \alpha_2 + \alpha_3 = 1$).

4.1 Generalized BEC in the case : $\alpha_1 < 1/2$

In this case the geometry is close to the usual cubic box. Our main result here is that the g-BEC of type I implies the macroscopic-cycles in the fundamental state.

Theorem 4.1 *If one takes the Casimir boxes Λ with $1/2 > \alpha_1$, then for a fixed density of particles $\rho > \rho_c(\beta)$ the chemical potential is $\bar{\mu}_\Lambda := \bar{\mu}_\Lambda(\beta, \rho) = -A/\beta V + o(1/V)$, with $A > 0$. This implies the s-BEC as well as the g-BEC of type I in the zero mode (ground state) together only with macroscopic-cycles in this mode. Here A is the unique solution of equation (2.7).*

Proof: Taking into account (2.1) we denote the family of Casimir boxes by Λ_I with the corresponding dual space:

$$\Lambda_I^* := \left\{ k \in \mathbb{R}^3 : k = \left(\frac{2\pi n_1}{V^{\alpha_1}}, \frac{2\pi n_2}{V^{\alpha_2}}, \frac{2\pi n_3}{V^{\alpha_3}} \right); n_\nu \in \mathbb{Z}^1; 1/2 > \alpha_1 \right\}. \quad (4.26)$$

Let $\Lambda_{0,I}^*$ be a subset of Λ_I^* defined by:

$$\Lambda_{0,I}^* = \{k \in \Lambda_I^* : \|k\| \leq \eta_I(V)\}, \quad (4.27)$$

where $\eta_I(V) = 1/V$. Then we have $\Lambda_{0,I}^* = \{k = 0\}$.

Let us consider the total density of particles:

$$\rho := \lim_{V \uparrow \infty} \rho_\Lambda(\beta, \bar{\mu}_\Lambda) = \rho_{\text{short}}(\beta, \rho) + \rho_{\text{long}}(\beta, \rho), \quad (4.28)$$

c.f. (2.18), (2.19).

We can decompose the density of particles in long-cycles into two parts defined by:

$$\rho_{\text{long}}(\beta, \rho) := \rho_{\text{long}}(\Lambda_I^* \setminus \Lambda_{0,I}^*) + \rho_{\text{long}}(\Lambda_{0,I}^*), \quad (4.29)$$

where $\rho_{\text{long}}(\Lambda_I^* \setminus \Lambda_{0,I}^*)$ is the limiting density of particles in long-cycles outside $\Lambda_{0,I}^*$:

$$\rho_{\text{long}}(\Lambda_I^* \setminus \Lambda_{0,I}^*) := \lim_{M \rightarrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_I^* \setminus \Lambda_{0,I}^*} \sum_{j=M}^{\infty} \rho_{\Lambda,j}(k) \right), \quad (4.30)$$

where the spectral repartition of particles density in j -cycles is:

$$\rho_{\Lambda,j}(k) := \frac{1}{V} e^{j\beta\bar{\mu}_{\Lambda}} e^{-j\beta\epsilon_{\Lambda}(k)}, \quad (4.31)$$

and $\bar{\mu}_{\Lambda} := \bar{\mu}_{\Lambda}(\beta, \mu)$ is the solution of equation $\rho = \rho_{\Lambda}(\beta, \mu)$.

First we shall estimate the density of particles in long-cycles of $\Lambda_I^* \setminus \Lambda_{0,I}^*$ by (4.30) and asymptotic for $\bar{\mu}_{\Lambda}$ we get:

$$\begin{aligned} \rho_{\text{long}}(\Lambda_I^* \setminus \Lambda_{0,I}^*) &= \lim_{M \rightarrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_I^* \setminus \Lambda_{0,I}^*} \sum_{j=M}^{\infty} \frac{1}{V} e^{j\beta\bar{\mu}_{\Lambda}} e^{-j\beta\epsilon_{\Lambda}(k)} \right), \\ &= \lim_{M \rightarrow \infty} \sum_{j=M}^{\infty} \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} dk e^{-j\pi\lambda_{\beta}^2 k^2}, \\ &= 0. \end{aligned}$$

Consequently there is no long cycles in $\Lambda_I^* \setminus \Lambda_{0,I}^*$ and since our last estimate is valid for any $M \rightarrow \infty$ we conclude that there are no s-long cycles in $\Lambda_I^* \setminus \Lambda_{0,I}^*$ (see Definition 3.2).

Now Let us consider the modes in $\Lambda_{0,I}^*$, we would like to prove that the PBG manifests s-long cycles of the order $O(V)$, i.e. macroscopic-cycles (see Definition 3.6).

Since $\bar{\mu}_{\Lambda} = -A/\beta V + o(1/V)$, with $A > 0$ we have:

$$\begin{aligned} \rho_{\text{long}}(\Lambda_{0,I}^* | \text{macro}) &:= \lim_{x \downarrow 0; y \uparrow \infty} \lim_{V \uparrow \infty} \sum_{j=xV}^{yV} \frac{1}{V} e^{j\beta\bar{\mu}_{\Lambda}} \\ &= \lim_{x \downarrow 0; y \uparrow \infty} \lim_{V \uparrow \infty} \frac{(e^{-xA+O(1/V)} - e^{-yA+O(1/V)})}{e^{-\beta\bar{\mu}_{\Lambda}} - 1} \\ &= \lim_{V \uparrow \infty} \frac{1}{e^{-\beta\bar{\mu}_{\Lambda}} - 1} = \lim_{V \uparrow \infty} \rho_{\Lambda}(\Lambda_{0,I}^*), \end{aligned} \quad (4.32)$$

where $\rho_{\Lambda}(\Lambda_{0,I}^*) := \sum_{k \in \Lambda_{0,I}^*} \rho_{\Lambda}(k)$ is the density of particles in $\Lambda_{0,I}^*$.

We can easily calculate $\rho_{\Lambda}(\Lambda_{0,I}^*)$:

$$\rho_{\Lambda}(\Lambda_{0,I}^*) = \frac{1}{V} \frac{1}{e^{-\beta\bar{\mu}_{\Lambda}} - 1} = \frac{1}{V} \frac{1}{e^{\beta \frac{A}{\beta V} + o(\frac{1}{V})} - 1} = \frac{1}{A} + o\left(\frac{1}{V}\right). \quad (4.33)$$

So by virtue of (4.29), (4.32) and (4.33):

$$\rho_{\text{long}}(\Lambda_{0,I}^* | \text{macro}) = \rho_{\text{long}}(\beta, \rho) = \frac{1}{A}, \quad (4.34)$$

We know by Theorem 2.2 that the density of particles in short-cycles is equal to the critical density. Consequently by virtue of (4.28), (4.34) we can conclude the proof of the theorem. \square

4.2 Generalized BEC in the case : $\alpha_1 = 1/2$

Our main result here is a theorem about g-BEC of type II due to the presence of macroscopic-cycles in an infinite (in the thermodynamical limit) number of modes.

Theorem 4.2 For the Casimir boxes with $1/2 = \alpha_1$ and a fixed particle density $\rho > \rho_c(\beta)$ the chemical potential is $\bar{\mu}_\Lambda = -B/\beta V + o(1/V)$, with $B > 0$. This implies the *s*-BEC as well as the *g*-BEC of type II in an infinite (in thermodynamical limit) number of modes and simultaneously macroscopic-cycles only in these modes. Here B is a unique solution of equation (2.8).

Proof: Taking into account (2.1) we denote the family of Casimir boxes by Λ_{II} and the dual space is:

$$\Lambda_{II}^* = \left\{ k \in \mathbb{R}^3 : k = \left(\frac{2\pi n_1}{V^{\alpha_1}}, \frac{2\pi n_2}{V^{\alpha_2}}, \frac{2\pi n_3}{V^{\alpha_3}} \right); n_\nu \in \mathbb{Z}^1; 1/2 = \alpha_1 \right\}, \quad (4.35)$$

Let $\Lambda_{0,II,\Gamma}^*$ be a subset of Λ_{II}^*

$$\Lambda_{0,II,\Gamma}^* = \{ k \in \Lambda_{II}^* : \|k\| \leq \eta_{II}^\Gamma(V) \}, \quad (4.36)$$

where:

$$\eta_{II}^\Gamma(V) := \frac{2\pi\Gamma}{V^{1/2}}, \Gamma \in \mathbb{N}^*, \quad (4.37)$$

Notice that this set contains the whole value of the condensate as well as particles involved in the long-cycles for $\Gamma \rightarrow \infty$ after thermodynamical limit.

Again we decompose the density of particles in long cycles into two parts:

$$\rho_{\text{long}}(\beta, \rho) = \lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{II}^* \setminus \Lambda_{0,II,\Gamma}^*) + \lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{0,II,\Gamma}^*). \quad (4.38)$$

We take the limit $\Gamma \rightarrow \infty$ to have the totality of the condensate in the first part.

The second part of 4.38 is:

$$\begin{aligned} \lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{II}^* \setminus \Lambda_{0,II,\Gamma}^*) &= \lim_{\Gamma \uparrow \infty} \lim_{M \uparrow \infty} \lim_{\Lambda} \left(\sum_{j=M}^{\infty} \frac{1}{V} e^{-j\beta C/V} \sum_{\|k\| > 2\pi\Gamma/\sqrt{V}} e^{-j\beta\epsilon_\Lambda(k)} \right) \\ &= \lim_{\Gamma \uparrow \infty} \lim_{M \uparrow \infty} \lim_{\Lambda} \left(\sum_{j=M}^{\infty} \frac{1}{V} e^{-j\beta C/V} \sum_{1/V^{1/2-\epsilon} > \|k\| > 2\pi\Gamma/\sqrt{V}} e^{-j\beta\epsilon_\Lambda(k)} \right) \\ &+ \lim_{\Gamma \uparrow \infty} \lim_{M \uparrow \infty} \lim_{\Lambda} \left(\sum_{j=M}^{\infty} \frac{1}{V} e^{-j\beta C/V} \sum_{\|k\| > 1/V^{1/2-\epsilon}} e^{-j\beta\epsilon_\Lambda(k)} \right), \quad (4.39) \end{aligned}$$

with $1/2 - \epsilon > \alpha_2$. Then we calculate the second term of 4.39:

$$\lim_{\Gamma \uparrow \infty} \lim_{M \uparrow \infty} \lim_{\Lambda} \left(\sum_{j=M}^{\infty} \frac{1}{V} e^{-j\beta C/V} \sum_{\|k\| > 1/V^{1/2-\epsilon}} e^{-j\beta\epsilon_\Lambda(k)} \right) = \lim_{M \uparrow \infty} \sum_{j=M}^{\infty} \frac{1}{j^{3/2} \lambda_\beta^3} = 0.$$

The first term of 4.39 have an upper bound:

$$\lim_{\Gamma \uparrow \infty} \lim_{\Lambda} \left(\sum_{j=1}^{\infty} \frac{1}{V} e^{-j\beta C/V} \sum_{1/V^{1/2-\epsilon} > \|k\| > 2\pi\Gamma/\sqrt{V}} e^{-j\beta\epsilon_\Lambda(k)} \right) = \lim_{\Gamma \uparrow \infty} \sum_{|n_1| > \Gamma} \frac{1}{\pi \lambda_\beta^2 n_1^2 + B} = 0,$$

consequently the first term of (4.39) is null so $\lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{II}^* \setminus \Lambda_{0,II,\Gamma}^*) = 0$.

Now let us consider the modes in $\Lambda_{0,II,\Gamma}^*$. We would like to apply the same strategy as the proof of the Theorem 4.1 to prove that the PBG manifests s-long cycles of the order $O(V)$, i.e. macroscopic-cycles (see Definition 3.6).

Since $\bar{\mu}_\Lambda = -B/\beta V + o(1/V)$, with $B > 0$ we have:

$$\begin{aligned}
\lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{0,II,\Gamma}^* | \text{macro}) &:= \lim_{\Gamma \uparrow \infty} \lim_{x \downarrow 0; y \uparrow \infty} \lim_{V \uparrow \infty} \frac{1}{V} \sum_{k \in \Lambda_{0,II}^*} \sum_{j=xV}^{yV} \rho_{\Lambda,j}(k) \\
&= \lim_{\Gamma \uparrow \infty} \lim_{x \downarrow 0; y \uparrow \infty} \lim_{V \uparrow \infty} \sum_{k \in \Lambda_{0,II,\Gamma}^*} \frac{(e^{-xB+O(1/V)} - e^{-yB+O(1/V)})}{e^{\beta(\epsilon_\Lambda(k) - \bar{\mu}_\Lambda)} - 1} \\
&= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \sum_{k \in \Lambda_{0,II,\Gamma}^*} \frac{1}{e^{\beta(\epsilon_\Lambda(k) - \bar{\mu}_\Lambda)} - 1} \\
&= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{\mathbf{k} \in \Lambda_{0,II,\Gamma}^*} \rho_\Lambda(\mathbf{k}) \right). \tag{4.40}
\end{aligned}$$

We can easily calculate:

$$\begin{aligned}
\lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{\mathbf{k} \in \Lambda_{0,II,\Gamma}^*} \rho_\Lambda(\mathbf{k}) \right) &= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \sum_{n_1=-\Gamma}^{\Gamma} \frac{1}{V} \frac{1}{e^{\beta(\pi \lambda_\beta^2 n_1^2 / V + B/V + O(1/V))} - 1} \\
&= \sum_{n_1 \in \mathbb{Z}^1} \frac{1}{B + \pi \lambda_\beta^2 n_1^2}. \tag{4.41}
\end{aligned}$$

So by virtue of (4.38), (4.40) and (4.41):

$$\lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{0,II,\Gamma}^* | \text{macro}) = \rho_{\text{long}}(\beta, \rho) = \sum_{n_1 \in \mathbb{Z}^1} \frac{1}{B + \pi \lambda_\beta^2 n_1^2}, \tag{4.42}$$

We know by Theorem 2.2 that the density of particles in short-cycles is equal to the critical density. Consequently by virtue of (4.28), (4.42) we can conclude the proof of the theorem. \square

4.3 Generalized BEC in the case : $\alpha_1 > 1/2$

Our main result is the theorem about g-BEC of type III due to the presence of *long-microscopic* cycles in infinite (in the thermodynamical limit) number of modes.

Theorem 4.3 *If one takes the Casimir boxes $\Lambda = V^{\alpha_1} \times V^{\alpha_2} \times V^{\alpha_3}$ with $1/2 > \alpha_1$, then for a fixed density of particles $\rho > \rho_c(\beta)$ the chemical potential is $\bar{\mu}_\Lambda(\beta, \rho) = -C/\beta V^\delta + o(1/V^\delta)$, with $\delta = 2(1 - \alpha_1)$ and $C > 0$. This implies the s-BEC as well as the g-BEC of the type III in infinite (in the thermodynamical limit) number of modes, together with long-microscopic cycles of the order V^δ , but only in these modes. Here C is a unique solution of equation (2.9).*

Proof: Taking into account (2.1) we denote the family of Casimir boxes Λ_{III} and the dual spaces are defined by:

$$\Lambda_{III}^* = \left\{ k \in \mathbb{R}^3 : k = \left(\frac{2\pi n_1}{V^{\alpha_1}}, \frac{2\pi n_2}{V^{\alpha_2}}, \frac{2\pi n_3}{V^{\alpha_3}} \right); n_i \in \mathbb{Z}^1; \alpha_1 > 1/2 \right\}. \quad (4.43)$$

Let $\Lambda_{0,III,\Gamma}^*$ be a subset of Λ_{III}^* :

$$\Lambda_{0,III,\Gamma}^* = \{ k \in \Lambda_{III}^* : \|k\| \leq \eta_{III}^\Gamma(V) \}, \quad (4.44)$$

where:

$$\eta_{III}^\Gamma(V) := \frac{2\pi\Gamma}{V^{\delta/2}}, \quad \Gamma \in \mathbb{N}^*, \quad (4.45)$$

here $\delta = 2(1 - \alpha_1) < 1$.

We show that $\Lambda_{0,III,\Gamma}^*$ contains the whole value of the condensate as well as the particles involved into long cycles in the limit $\Gamma \rightarrow \infty$.

Before to present formal arguments, let us make a remark about *qualitative* difference between the cases $\alpha_1 > 1/2$ and $\alpha_1 \leq 1/2$. With the definition of $\eta_{III}^\Gamma(V)$ (4.45), we see that the number of states in $\Lambda_{0,III,\Gamma}^*$ is of the order $O(V^{2\alpha_1-1})$ that goes to infinity, with increasing volume, there are much more states in the condensate than in $\Lambda_{0,II,\Gamma}^*$ (defined by 4.36). Heuristically one can say that $\Lambda_{0,III,\Gamma}^*$ contains long cycles of sizes of the order $O(V^\delta)$ in a number of modes of the order $O(V^{2\alpha_1-1})$. Thus the number of particles in these s-long cycles is of the order $O(V^\delta)O(V^{2\alpha_1-1}) = O(V)$, which is macroscopic. For this reason there is a macroscopic condensate (the order of the number of particles is $O(V)$) because there is an accumulation of microscopic condensates (the order of the number of particles is $O(V^{2\alpha_1-1})$ which is smaller than $O(V)$) as well as an accumulation of long-microscopic cycles in $\Lambda_{0,III,\Gamma}^*$ at each mode of the condensate.

Again we decompose the density of particles in long cycles into two parts:

$$\rho_{\text{long}}(\beta, \rho) = \lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{III}^* \setminus \Lambda_{0,III,\Gamma}^*) + \lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{0,III,\Gamma}^*). \quad (4.46)$$

By the same argument as in the proof of Theorem 4.2, one finds that the first term of 4.46 is null.

Now let us consider the modes in $\Lambda_{0,III,\Gamma}^*$. In this case, we would like to prove that the PBG manifests s-long cycles of the order $O(V^\delta)$, $\delta = 2(1 - \alpha_1) < 1$, i.e. microscopic-cycles (see Definition 3.5 and 3.6).

Since $\bar{\mu}_\Lambda = -C/\beta V^\delta + o(1/V^\delta)$ with $\delta = 2(1 - \alpha_1)$ and $C > 0$ we obtain:

$$\begin{aligned} \lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{0,III,\Gamma}^* | \text{micro}) &:= \lim_{\Gamma \uparrow \infty} \lim_{x \downarrow 0; y \uparrow \infty} \lim_{V \uparrow \infty} \frac{1}{V} \sum_{k \in \Lambda_{0,III,\Gamma}^*} \sum_{j=xV^\delta}^{yV^\delta} \frac{1}{V} e^{j\beta\bar{\mu}_\Lambda} e^{-j\beta\epsilon_\Lambda(k)} \\ &= \lim_{\Gamma \uparrow \infty} \lim_{x \downarrow 0; y \uparrow \infty} \lim_{V \uparrow \infty} \sum_{k \in \Lambda_{0,III,\Gamma}^*} \frac{(e^{-xC+O(1/V)} - e^{-yC+O(1/V)})}{e^{\beta(\epsilon_\Lambda(k) - \bar{\mu}_\Lambda)} - 1} \\ &= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{\mathbf{k} \in \Lambda_{0,III,\Gamma}^*} \rho_\Lambda(\mathbf{k}) \right). \end{aligned} \quad (4.47)$$

We can easily calculate:

$$\lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{\mathbf{k} \in \Lambda_{0,III,\Gamma}^*} \rho_{\Lambda}(\mathbf{k}) \right) = \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\frac{1}{V^{\delta}} \frac{1}{V^{2\alpha_1-1}} \sum_{j=1}^{\infty} e^{-(j/V^{\delta})C} \sum_{n_1=-\Gamma}^{\Gamma} e^{-\pi \lambda_{\beta}^2 (j/V^{\delta})(n_1/V^{2\alpha_1-1})^2} \right).$$

Since this expression is nothing but the limit of double Darboux-Riemann sums, in thermodynamical limit we obtain a double integral:

$$\lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{\mathbf{k} \in \Lambda_{0,III,\Gamma}^*} \rho_{\Lambda}(\mathbf{k}) \right) = \lim_{\Gamma \uparrow \infty} \int_{\mathbb{R}^+} d\zeta e^{-\zeta C} \int_{-\Gamma}^{\Gamma} d\xi e^{-\zeta \pi \lambda_{\beta}^2 \xi^2} = \int_{\mathbb{R}^+} d\zeta \frac{e^{-\zeta C}}{\sqrt{\zeta} \lambda_{\beta}} = \frac{\sqrt{\pi}}{C^{1/2} \lambda_{\beta}} \quad (4.48)$$

Hence:

$$\lim_{\Gamma \uparrow \infty} \rho_{\text{long}}(\Lambda_{0,III,\Gamma}^* | \text{micro}) = \rho_{\text{long}}(\beta, \rho) = \frac{\sqrt{\pi}}{C^{1/2} \lambda_{\beta}}, \quad (4.49)$$

by virtue of (4.46), (4.47) and (4.48).

We know by Theorem 2.2 that the density of particles in short-cycles is equal to the critical density. Consequently by virtue of (4.28), (4.49) we can conclude the proof of the theorem. \square

5 Does generalized BEC I, II, III imply a hierarchy of ODLRO?

In the Introduction we presented three concepts related to the BEC: g-BEC, long cycles and ODLRO. We present in Sections 2 and 3 two new concepts: *scaled* BEC (s-BEC) and *scaled* short/long cycles (s-short/long cycles) associated with g-BEC and short/long cycles. Thus it seems consistent to introduce a concept of *scaled* ODLRO (s-ODLRO) via our scaling approach to study the hierarchy of ODLRO.

This part gives a physical meaning of the scaling approach because it allows the study of the coherence of the condensate at large scale. We show that for very anisotropic cases ($\alpha_1 > 1/2$) the coherence length (maximal length of correlation) is not equal to the size of the box (see Theorem 5.1).

5.1 Scaling approach to ODLRO

Recall that the generalized criterion of ODLRO is that there is ODLRO if and only if there is g-BEC, see Theorem 2.1. The standard definition of ODLRO is formulated in Definition 2.3:

$$\sigma(\beta, \rho) := \lim_{\|x-x'\| \uparrow \infty} \sigma(\beta, \rho; x, x'),$$

where $\sigma(\beta, \rho; x, x')$ is the two-point correlation function between two points x and x' after thermodynamical limit. Notice that this definition can not be satisfactory when we use the definition of s-BEC, since we do not precise yet what are the scales of large correlations. It seems to be interesting to take thermodynamical limit at the same time as we take the two points x and x' at increasing distance.

A natural question is whether we are able to detect different types of s-BEC (as well as g-BEC) with the help of a generalized criterion of ODLRO based on our scaling approach? We call it a *scaled* ODLRO (s-ODLRO).

Definition 5.1 *The PBG manifests a s-ODLRO if there exists a vector-valued function of volume $X : V \mapsto X(V) \in \Lambda$ such that $\lim_{V \uparrow \infty} |X_\nu(V)| = \infty$, $\nu = 1, 2, 3$ and:*

$$\sigma_X(\beta, \rho) := \lim_{V \uparrow \infty} (\sigma_{\Lambda, X})(V) > 0, \quad (5.50)$$

where $(\sigma_{\Lambda, X})(V)$ is the two-point scaled-correlation function (two-point s-correlation function) for $x(V), x'(V) \in \Lambda$, see (2.11):

$$(\sigma_{\Lambda, X})(V) := \sigma_\Lambda(\beta, \rho; x(V) - x'(V)) = \sum_{k \in \Lambda^*} \rho_\Lambda(k) e^{ik \cdot X(V)}, \quad (5.51)$$

here $X(V) = (x - x')(V) \in \Lambda$.

Remark 5.1 *By (2.1) and (2.2) one can write (2.11) like:*

$$\sigma(\beta, \rho; x, x') = \sum_{j=1}^{\infty} e^{j\beta\bar{\mu}_\Lambda} \prod_{\nu=1}^3 \theta_3\left(\frac{\pi}{V^{2\alpha_\nu}}(x_\nu - x'_\nu), e^{-j\pi \frac{\lambda_\beta^2}{V^{2\alpha_\nu}}}\right), \quad (5.52)$$

where $\theta_3(u, q) := \sum_{n \in \mathbb{Z}^1} q^{n^2} e^{2inu}$ is the elliptic theta-function.

This implies the following proposition:

Proposition 5.1 *By (5.51) the two-point correlation function as well as the two-point s-correlation function (see Definition 5.1) are non-negative symmetric and L_ν -periodic functions of $x_\nu - x'_\nu$, $\nu = 1, 2, 3$ on \mathbb{R} , and decreasing/increasing on $[nL_\nu, nL_\nu + L_\nu/2] \subset \mathbb{R}^+$, $n \in \mathbb{N}$, respectively on $[nL_\nu + L_\nu/2, (n+1)L_\nu] \subset \mathbb{R}^+$, $n \in \mathbb{N}$ (i.e. monotone on the semi-periods).*

Proof: These properties follow from the properties of the elliptic theta-function [A-S].
□

The following statement is a direct consequence of Definitions 2.3 and Remark 5.1:

Lemma 5.1 *For any vector-valued $X(V)$ we have:*

$$0 \leq \sigma_X(\beta, \rho) \leq \sigma(\beta, \rho). \quad (5.53)$$

This Lemma means that the s-ODLRO implies standard ODLRO.

5.2 Hierarchy and anisotropy of ODLRO, coherence of the condensate

Here we use Definition 5.1 to analyze the s-BEC and the s-long cycles in the Casimir boxes. Notice that the usual criterion of ODLRO is such that we have no indication of the scale of long correlations because we study their correlations after thermodynamical limit.

We introduce a classification of the s-OLDRO which is formally defined by:

Definition 5.2 *The PBG manifests the macroscopic-ODLRO in the direction x_ν , if there exist a vector $X(V) = (X_1(V), X_2(V), X_3(V)) \in \Lambda$ such that: $\lim_{V \uparrow \infty} |X_\nu(V)|/V^{\alpha_\nu} > 0$ and $\sigma_X(\beta, \rho) > 0$.*

Definition 5.3 *If the PBG does not manifests the macroscopic-ODLRO in the direction x_ν , although the PBG manifests the s-ODLRO, then it manifests the microscopic-ODLRO in the direction x_ν .*

With the periodic boundary conditions the system is homogeneous and so there is no localization of the condensate in the space contrary to the case of the Dirichlet boundary conditions. However the coherence length of the condensate could be studied by virtue of the precedent Definitions 5.2 and 5.3.

Theorem 5.1 *Let us consider the grand-canonical PBG in Casimir boxes $\Lambda = V^{\alpha_1} \times V^{\alpha_2} \times V^{\alpha_3}$ with Dirichlet boundary conditions and a fixed density of particles ρ . Let $X : V \in \mathbb{R}^+ \mapsto X(V) = (X_1(V), X_2(V), X_3(V)) \in \Lambda$, $\lim_{V \uparrow \infty} X_\nu(V) = \infty$, $0 < X_\nu(V) \leq V^{\alpha_\nu}/2$, $\nu = 1, 2, 3$. Then we have the following results concerning the s-ODLRO, see (5.50):*

$$\sigma_X(\beta, \rho) = 0, \text{ for } \rho < \rho_c(\beta), \quad (5.54)$$

Whereas for $\rho > \rho_c(\beta)$ we get:
for $\alpha_1 < 1/2$:

$$\sigma_X(\beta, \rho) = \rho_0(\beta), \quad (5.55)$$

for $\alpha_1 = 1/2$:

$$\sigma_X(\beta, \rho) = \rho_0(\beta), \text{ for } \lim_{V \uparrow \infty} (X_1(V)/V^{\alpha_1}) = 0, \quad (5.56)$$

$$= \sum_{n_1 \in \mathbb{Z}^1} \frac{\cos \pi n_1 x}{\pi \lambda_\beta^2 n_1^2 + B} < \rho_0(\beta), \text{ for } X_1(V) = xV^{\alpha_1}/2, \text{ } 0 < x < 1, \text{ } , \quad (5.57)$$

for $\alpha_1 > 1/2$:

$$\sigma_X(\beta, \rho) = \rho_0(\beta), \text{ for } \lim_{V \uparrow \infty} (X_1(V)/V^\delta) = 0, \text{ } \delta = 2(1 - \alpha_1), \quad (5.58)$$

$$= \rho_0(\beta) e^{-2x\sqrt{\pi c}/\lambda_\beta} < \rho_0(\beta), \text{ for } X_1(V) = xV^{\delta/2}, \text{ } x > 0, \quad (5.59)$$

$$= 0, \text{ for } \lim_{V \uparrow \infty} (X_1(V)/V^{\delta/2}) = 0. \quad (5.60)$$

Proof: To ensure that the correlation function is monotonously decreasing for periodic boundary conditions, we choose $0 < X_\nu \leq \frac{1}{2}V^{\alpha_\nu}$, $\nu = 1, 2, 3$, see Proposition 5.1.

The first step of the proof is to study the case $\rho < \rho_c(\beta)$:

Since $\sigma(\beta, \rho) = 0$ (Theorem 2.1), by Lemma 5.1 we get $\sigma_X(\beta, \rho) = 0$ for any vector $X(V) \in \Lambda$.

The second step is to study the case $\rho > \rho_c(\beta)$:

For $\alpha_1 < 1/2$ by Definition 5.1 we have:

$$\begin{aligned} \sigma_X(\beta, \rho) &= \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{0,I}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) + \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_I^* \setminus \Lambda_{0,I}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) \\ &= \lim_{V \uparrow \infty} \left(\frac{1}{V} \sum_{j=1}^{\infty} e^{-Aj/V} \right) + \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_I^* \setminus \Lambda_{0,I}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) \end{aligned} \quad (5.61)$$

where Λ_I^* is the dual vector space given by equation (4.26) and $\Lambda_{0,I}^* = \{k = 0\}$ is the subset corresponding to the condensate defined by (4.27). The first term of (5.61) is equal to $\rho_0(\beta, \rho)$ by virtue of Theorem 4.1, so given that $\sigma(\beta, \rho) = \rho_0(\beta, \rho)$ (Theorem 2.1) and by Lemma 5.1 the second term of (5.61) have to be null and we get the result.

For $\alpha_1 = 1/2$ by Definition 5.1 we obtain:

$$\begin{aligned} \sigma_X(\beta, \rho) &= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{0,II,\Gamma}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) + \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{II}^* \setminus \Lambda_{0,II,\Gamma}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) \\ &= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\frac{1}{V} \sum_{j=1}^{\infty} e^{-Bj/V} \sum_{n_1=-\Gamma}^{\Gamma} e^{-\pi \lambda_\beta^2 n_1^2 (j/V)} e^{iX_1(V)(n_1/V)} \right) \\ &+ \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{II}^* \setminus \Lambda_{0,II,\Gamma}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) \end{aligned} \quad (5.62)$$

where Λ_{II}^* is the dual vector space given by equation (4.35) and $\Lambda_{0,II,\Gamma}^*$ is the subset corresponding to the condensate defined by (4.36).

If $\lim_{V \uparrow \infty} (X_1(V)/V^{\alpha_1}) = 0$, the first term in the right-hand side of (5.62) is equal to $\rho_0(\beta, \rho)$ by virtue of Theorem 4.2. Given that $\sigma(\beta, \rho) = \rho_0(\beta, \rho)$ (Theorem 2.1) and by Lemma 5.1, the second term in (5.62) is null and we obtain the result.

Let $\lim_{V \uparrow \infty} (X_1(V)/V^{\alpha_1}) = x$, $0 < x \leq 1/2$. Since the sum inside the limit in the first term of the right-hand side of (5.62) is a Darboux-Riemann sum, one get:

$$\int_{\mathbb{R}^+} d\chi e^{-B\chi} \sum_{n_1 \in \mathbb{Z}^1} e^{-\pi \lambda_\beta^2 n_1^2 \chi} e^{ixn_1} = \sum_{n_1 \in \mathbb{Z}^1} \frac{\cos \pi n_1 x}{\lambda_\beta^2 n_1^2 + B}.$$

The second term in the right hand side of (5.62) is null because the phase implies that it is smaller than the density of particles in $\Lambda_{II}^* \setminus \Lambda_{II,0,\Gamma}^*$ which is null in the limit $\Gamma \rightarrow \infty$.

For $\alpha_1 > 1/2$ by Definition 5.1 and by virtue of (4.48) we have:

$$\begin{aligned}
\sigma_X(\beta, \rho) &= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{0,III,\Gamma}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) + \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{III}^* \setminus \Lambda_{0,III,\Gamma}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) \\
&= \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\frac{1}{V^\delta} \frac{1}{V^{2\alpha_1-1}} \sum_{j=1}^{\infty} e^{-Cj/V^\delta} \sum_{n_1=-\Gamma}^{\Gamma} e^{-\pi\lambda_\beta^2(j/V^\delta)(n_1/V^{2\alpha_1-1})^2} e^{iX_1(V)n_1/V^{\alpha_1}} \right) \\
&+ \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{III}^* \setminus \Lambda_{0,III,\Gamma}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right) \tag{5.63}
\end{aligned}$$

where Λ_{III}^* is the dual vector space given by equation (4.43) and $\Lambda_{0,III}^*$ is the subset corresponding to the condensate defined by (4.44).

If $\lim_{V \uparrow \infty} (X_1(V)/V^{\delta/2}) = 0$, the right-hand side of (5.63) is equal to $\rho_0(\beta, \rho)$ by virtue of Theorem 4.3. Since $\sigma(\beta, \rho) = \rho_0(\beta, \rho)$ (Theorem 2.1), by Lemma 5.1 the second term of (5.63) have to be null and we get the result.

Let $\lim_{V \uparrow \infty} (X_1(V)/V^{\delta/2}) = x$, $x > 0$, $\delta = 2(1 - \alpha_1)$. Then the sum inside the limit in the first term of (5.63) is a double Darboux-Riemann sum, which implies:

$$\int_{\mathbb{R}^+} d\xi e^{-C\xi} \int_{\chi \in \mathbb{R}} d\chi e^{-\pi\lambda_\beta^2 \xi \chi^2} e^{2\pi i x \chi} = \frac{\sqrt{\pi}}{\lambda_\beta \sqrt{C}} e^{-2x\sqrt{\pi C}/\lambda_\beta}.$$

By the same argument as in the case $\alpha_1 = 1/2$ the second part of (5.63) is null (the precedent expression is a decreasing function of x for $x > 0$) and thus we obtain the result.

Let $\lim_{V \uparrow \infty} (X_1(V)/V^{\delta/2}) = \infty$. Since the correlation function is a decreasing function for $0 < X_\nu \leq V^{\alpha_\nu}/2$ (see Proposition 5.1), it is uniformly bounded by the above estimate with $X(V) = xV^\delta$, $x > 0$:

$$\int_{\mathbb{R}^+} d\xi e^{-C\xi} \int_{\chi \in \mathbb{R}} d\chi e^{-\pi\lambda_\beta^2 \xi \chi^2} e^{2\pi i x \chi} + \lim_{\Gamma \uparrow \infty} \lim_{V \uparrow \infty} \left(\sum_{k \in \Lambda_{III}^* \setminus \Lambda_{0,III,\Gamma}^*} \rho_\Lambda(k) e^{ik \cdot X(V)} \right).$$

When x tends to infinity, the first part goes to zero (by the Riemann-Lesbegue theorem) then the precedent arguments show that the second part is also null. This concludes the proof. \square

We give here a classification of the s-ODLRO for three cases of Casimir boxes:

Theorem 5.2 *If one takes the Casimir boxes with $\alpha \leq 1/2$ then for a fixed density $\rho > \rho_c(\beta)$ the PBG manifests macroscopic-ODLRO in three directions. If $\alpha > 1/2$ then for a fixed density $\rho > \rho_c(\beta)$ the PBG manifests microscopic-ODLRO in direction x_ν and macroscopic-ODLRO in other directions.*

Proof: Definitions 5.2, 5.3 and Theorem 5.1 give the proof of the theorem. \square

It is remarkable that for type I and II g-BEC in Casimir boxes corresponding to $\alpha_1 \leq 1/2$ the condensate is spatially macroscopic whereas for the case $\alpha_1 > 1/2$ the condensate is spatially macroscopic in two directions but microscopic in the most anisotropic direction

x_1 . It is naturally to guess that there is a link between the size of s-long cycles and coherence length of the condensate. We can see this explicitly in [U] where the competition between the size of correlation X and the size of cycle j indicates that the coherence length is of the order of the square root of the size of the s-long cycles (e.g. $V^{\delta/2}$, $\delta = 2(1 - \alpha_1)$ in the case of the PBG in Casimir boxes with $\alpha_1 > 1/2$).

6 Concluding remarks

Some technical remarks

In this paper we introduce a new concept of BEC which is called the scaled BEC (s-BEC) to adapt the London scaling approach to the problem of g-BEC. It implies the van den Berg-Lewis-Pulé classification of BEC in three types (I,II,III) illustrated for the particular case of the PBG in Casimir boxes. This is a first formal step necessary before to study more carefully the different case of g-BEC for the PBG in Casimir boxes. One can see this by virtue of Proposition and 3.1.

The fundamental question that we study in this paper is the relation of different types of g-BEC (I,II,III) with the long cycles and with the ODLRO. Our results concerning the PBG in Casimir boxes can be summarized as follows:

- we introduced new concept of short/long cycles called scaled short/long cycles (s-short/long cycles, see Definition 3.2) to distinguish different types of g-BEC, see Theorems 4.1, 4.2 4.3 and Remark 3.2 and Proposition 3.3. This paper is based on the estimation of the size of s-long cycles in the condensate, see Definitions 3.3 at 3.5. If the size of s-long cycles is macroscopic, then the g-BEC is of the type I or II but if the s-long cycles are microscopic, then the g-BEC is of type III.

- we introduced new concept of ODLRO, called scaled ODLRO (s-ODLRO, see Definition 5.1), to distinguish the different types of g-BEC, see Theorems 5.1, 5.2. Our arguments are based on the estimate of the coherence length of the condensate, see Definitions 5.2, 5.3. If the coherence length is macroscopic in the three directions, then the g-BEC is of type I or II, and if in one of the three dimensions the coherence length is microscopic, then the g-BEC is of type III.

One can see that the proof of the Theorem 4.1, 4.2 and 4.3 is based on an analysis of geometric series easily done via s-long cycles. For this reason we can say that s-long cycles is a well adapt technique to study classification of BEC. Another reason is that the concept of cycles is independent of the representation of the gas (Feynman-Kac versus spectral representation). So cycles could be useful to study generalized BEC for an interacting Bose gas.

For simplicity we consider here the PBG with periodic boundary conditions but one can adapt present paper to Dirichlet or Neumann boundary conditions for which we can characterize the geometric form of the condensate cloud via the concept of scaled local density. In these cases the results concerning the hierarchies of cycles and ODLRO do not change. However, if one takes attractive boundary conditions, see [V-V-Z], we guess that the result should be different, since in this case the condensate is localized in two modes and it is not homogeneous. One can suppose that the s-long cycles are macroscopic but the most interesting is the coherence length of the condensate and its geometric form.

The last technical remark concerns the validity of the results in the canonical ensemble. It is known [P-Z] that in the canonical ensemble for the PBG in Casimir boxes there is the same type of generalized BEC as in the grand-canonical ensemble for the equivalent geometry. Thus we believe that the principle conclusions concerning the hierarchy of long cycles and of ODLRO should not change. However given that the amount of condensate in individual states are different [P-Z] from the grand-canonical case and some results (formulae in the Propositions 3.1 and 3.3 and Theorem 5.1) should be different.

Conceptual and physical remarks

First notice that ODLRO, long cycles and generalized BEC are equivalent criteria for PBG in Casimir boxes. Consequently generalized BEC is more relevant than usual BEC (macroscopical occupation of ground state) because it contains all cases of BEC (fragmented or not).

In this article we present a scaling approach of Bose-Einstein condensation. It allows classification of different types of condensate via scaling size of long cycles in relation with scaling size of large correlations. The interest of the study of long cycles using scaling approach is the knowledge on coherence properties of condensate by virtue of the rule of Bose statistics in the two points correlation function. Heuristically the order of the size of large correlation is the square root of the order of the size of long cycles.

Scaling approach should be useful for interpretation of thermodynamical results for large particles number systems. It is interesting because one can study the effect of the geometry of the box on geometry and coherence properties of the condensate. In this paper, we show that a condensate is like a finite or infinite number of macroscopic particle (type I or II) or infinite number of microscopic particles (type III, analogous to quasi-condensate [M-H-U-B], [P-S-W]) formed by "closed polymers chain" (cycles) of macroscopic or microscopic size related to coherence length (square root of size of cycles). Physically it could be interesting to study the correlation function of a condensate in an harmonic trap with pulsations $\omega_x, \omega_y, \omega_z$ using our scaling approach (e.g. for very anisotropic traps).

Perspectives

Here we took Casimir boxes to illustrate our concepts. But we can use *van den Berg* boxes which is a generalization of Casimir boxes ($\Lambda_L = L_1(L) \times L_2(L) \times L_3(L)$ with $V_L := |\Lambda_L| = L_1(L)L_2(L)L_3(L)$ where $L_i(L)$ are functions of a parameter L such as $\lim_{L \rightarrow \infty} L_i(L) = \infty$). These boxes are very interesting because with particular choice of the functions $L_i(L)$, e.g. (see[vdB]) $L_1(L) = L_2(L) = e^L, L_3(L) = L$ with Dirichlet boundary conditions, for $\rho_c(\beta) < \rho < \rho_m(\beta)$ the g-BEC is of type III and for $\rho > \rho_m(\beta)$ it seems there is the *coexistence* of g-BEC of type I and type III, where $\rho_c(\beta)$ is the critical density defined by (2.5) and $\rho_m(\beta)$ is a critical density defined in this particular case by $\rho_m(\beta) = \rho_c(\beta) + 1/\beta\pi$. In fact $\rho_c(\beta)$ separates regimes condensate - non condensate and $\rho_m(\beta)$ separates the different types of generalized BEC. It seems to be analogous with quasi-condensate/condensate transition [P-S-W]. It is natural now to study this curious phenomena using our approach, which will be a subject for another article.

Whether the scaling approach for the interacting gas problem is still relevant? One can see that the concept of g-BEC is well formulated for the PBG but not for the interacting Bose gas. Then how to study the classification of BEC for the interacting Bose gas if we have no more indications about the spectral properties of the gas? This problem could be solved using our scaling approach (of long cycles or ODLRO). Thus the next step will be the application of these new methods for some models of interacting Bose gas (especially for weakly interacting Bose gas at first time, take e.g. an interaction $U = \sum_{k \in \Lambda^*} g N_k^2 / 2V$, $g > 0$, see [Br-Z])). For strongly interacting bosons in *quantum crystals*, there is formation of infinite cycles [U2]. It could be interesting to study them using scaling approach of long cycles.

Another interesting problem is the characterization of quantum fluctuations in many body systems. Scaling approach could be also relevant to study finite size scaling effect [B-D-T] appearing in some systems, like Casimir effect in quantum liquids [M-Z], [Z-al].

Acknowledgments

I would like to thank my promotor Valentin Zagrebnov for introduction me into this subject, encouragements, helpful remarks and discussions. I benefited a lot by the referees constructive criticism and suggestions to improve the first version of the manuscript.

References

- [A-S] M.Abramowitz, I.A.Stegun, eds (1972), Handbook of Mathematical functions with formulas, graphs and mathematical tables, New York: Dover Publications.
- [B-D-T] J.G.Brankov, D.M.Danchev, N.S.Tonchev, World Scientific Publishing Co. Pte. Ltd, Theory of critical phenomena in finite-size systems: scaling and quantum effects, Series on modern condensed matter physics; vol. 9, (2000).
- [Br-Z] J.B.Bru, V.A.Zagrebnov, Exactly soluble model with two kinds of Bose-Einstein condensations, *Physica A* **268**, 309-325, (1999).
- [C] H.B.G.Casimir, On Bose-Einstein condensation, *Fundamental Problems in Statistical Mechanics III*, ed E.G.D.Cohen, p. 188-196, (1968)
- [C-S] J.A.Cuesta, R.P.Sear Phase transition analogous to Bose-Einstein condensation in systems of noninteracting surfactant aggregates, *Physical Review E* **65**, 031406, (2002).
- [D-M-P] T.C.Dorlas, Ph.A.Martin, J.V.Pulé, Long Cycles in a Perturbed Mean Field Model of a Boson Gas, *Journal of Statistical Physics* **121**, 433-461 (2005).
- [F] R.P.Feynman, Atomic Theory of the λ -Transition in Helium, *The Physical Review* **91**, 1291 (1953).
- [dG] P-G.de Gennes, Scaling concepts in polymer physics, Cornell University (1979).

- [G] J.Ginibre, "Some applications of functional integration in statistical mechanics" Statistical Mechanics and Quantum Field Theory, C. DeWitt, R.Stora, eds, Les Houches, Gordon and Breach, 1971, p. 327.
- [L] F.London On the Bose-Einstein condensation, *Phys. Rev.* **54**, 947-954 (1938).
- [L-S-S-Y] The Mathematics of the Bose Gas and its Condensation, Series: Oberwolfach Seminars , Vol. 34 E.H.Lieb, R.Seiringer, J.P.Solovej, J.Yngvason, 2005, VIII, p.208.
- [M] Ph.A.Martin, Quantum mayer graph: application to Bose and Coulomb gases, *Acta Physica Polonica B* **34**, 3629 (2003).
- [M-H-U-B] E. Mueller, T. Ho, M. Ueda, G. Baym, Fragmentation of Bose-Einstein condensates, *Physical Review A* **74**, 063623 (2006).
- [M-V] T.Michoel, A.Verbeure, Nonextensive Bose-Einstein condensation model, *J.Math.Phys.* **40**, 1668-1679, (1999).
- [M-Z] Ph.A.Martin, V.A.Zagrebnov, The Casimir effect for the Bose-gas in slabs, *Europhysics Letters* **73**(1), 15-20 (2006).
- [P-O] O.Penrose, L.Onsager Bose-Einstein Condensation and Liquid Helium, *Physical Review* **104**, 576 (1956).
- [P-S-W] D.S.Petrov, G.V.Shlyapnikov, J.T.M.Walraven Phase-Fluctuating 3D Bose-Einstein Condensates in Elongated Traps, *Phys. Rev. Lett.* **87**, 50404 (2001).
- [P-Z] J.Pulé, V.A.Zagrebnov The canonical perfect Bose gas in Casimir boxes, *J. Math. Phys.* **45**, 3565 (2004).
- [S] A. Sütö, Percolation transition in the Bose gas: II, *J.Phys.A:Math.Gen* **35**, 6995-7002 (2002).
- [U] D. Ueltschi, Feynman cycles in the Bose gas, *J. Math. Phys.* **47**, 123303 (2006).
- [U2] D. Ueltschi, Relation between Feynman Cycles and Off-Diagonal Long-Range Order, *Physical Review Letters* **97**, 170601, (2006).
- [vdB] M. van den Berg On Condensation in the Free-Boson Gas and the Spectrum of the Laplacian, *Journal of Statistical Physics* **31**, 623-637, (1983).
- [vdB-L] M. van den Berg, J.T.Lewis, On generalized condensation in the free boson gas, *Physica 110A*, 550-564 (1982).
- [vdB-L-L] M. van den Berg, J.T.Lewis and M.Lunn On the general theory of Bose-Einstein condensation and the state of the free boson gas, *Helv.Phys.Acta* **59**, 1289-1310 (1986).
- [vdB-L-P] M. van den Berg, J.T.Lewis and J.V.Pulé A general theory of Bose-Einstein condensation, *Helv.Phys.Acta* **59**, 1271-1288 (1986).

- [V-V-Z] L.Vandedevenne, A.Verbeure and V.A.Zagrebnov, Equilibrium states for the Bose gas, *J.Math.Phys.* **45**, 1606–1622, (2004).
- [Z-al] R. Zandi et al., Thinning of superfluidity films below the critical point, *Physical Review E* **76**, 030601 (2007).
- [Z-B] V.A.Zagrebnov, and J.B.Bru, The Bogoliubov model of weakly imperfect Bose gas, *Phys. Rep.* **350**, no. 5-6, 291434, (2001)