HAL
open science

# Quasi-energy function for diffeomorphisms with wild separatrices

Viatcheslav Grines, Francois Laudenbach, Olga Pochinka

## To cite this version:

Viatcheslav Grines, Francois Laudenbach, Olga Pochinka. Quasi-energy function for diffeomorphisms with wild separatrices. Mathematical Notes (transl. of Mat. Zametki), 2009, 86 (2), pp.163-170. hal-00333484

**HAL Id: hal-00333484**

**https://hal.science/hal-00333484v1**

Submitted on 23 Oct 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Quasi-energy function for diffeomorphisms with wild separatrices

V. Grines* F. Laudenbach† O. Pochinka‡

October 23, 2008

**Abstract**

According to Pixton [8] there are Morse-Smale diffeomorphisms of $\mathbb{S}^3$ which have no energy function, that is a Lyapunov function whose critical points are all periodic points of the diffeomorphism. We introduce the concept of quasi-energy function for a Morse-Smale diffeomorphism as a Lyapunov function with the least number of critical points and construct a quasi-energy function for any diffeomorphism from some class of Morse-Smale diffeomorphisms on $\mathbb{S}^3$.

*Mathematics Subject Classification:* 37B25, 37D15, 57M30.
*Keywords:* Morse-Smale diffeomorphism, Lyapunov function, Morse theory.

First and third authors thank grant RFBR No 08-01-00547 of Russian Academy for partial financial support.

## 1 Formulation of results

According to [3], given a closed smooth $n$-manifold $M^n$ and a Morse function $\varphi : M^n \to \mathbb{R}$ is called a *Morse-Lyapunov function* for Morse-Smale diffeomorphism $f : M^n \to M^n$ if:

1) $\varphi(f(x)) < \varphi(x)$ if $x \notin Per(f)$ and $\varphi(f(x)) = \varphi(x)$ if $x \in Per(f)$, where $Per(f)$ is the set of periodic points of $f$;

2) any point $p \in Per(f)$ is a non-degenerate maximum of $\varphi|_{W^u(p)}$ and a non-degenerate minimum of $\varphi|_{W^s(p)}$.

**Definition 1.1** *Given a Morse-Smale diffeomorphism $f : M^n \to M^n$, a function $\varphi : M^n \to \mathbb{R}$ is a quasi-energy function for $f$ if $\varphi$ is a Morse-Lyapunov function for $f$ and has the least possible number of critical points among all Morse-Lyapunov functions for $f$.*

*N. Novgorod State University, Gagarina 23, N. Novgorod, 603950 Russia, grines@vmk.unn.ru.

†Laboratoire de mathématiques Jean Leray, UMR 6629 du CNRS, Faculté des Sciences et Techniques, Université de Nantes, 2, rue de la Houssinière, F-44322 Nantes cedex 3, France, francois.laudenbach@univ-nantes.fr.

‡N. Novgorod State University, Gagarina 23, N. Novgorod, 603950 Russia, olga-pochinka@yandex.ru.

In this paper we consider the class $G_4$ of Morse-Smale diffeomorphisms $f : \mathbb{S}^3 \to \mathbb{S}^3$ whose nonwandering set consists of exactly four fixed points: one source $\alpha$, one saddle $\sigma$ and two sinks $\omega_1$ and $\omega_2$. It follows from [9] (theorem 2.3), that the closure of each connected component (separatrix) of the one-dimensional manifold $W^u(\sigma) \setminus \sigma$ is homeomorphic to a segment which consists of this separatrix and two points: $\sigma$ and some sink. Denote by $\ell_1, \ell_2$ the one-dimensional separatrices containing the respective sinks $\omega_1, \omega_2$ in their closures. According to [9], $\bar{\ell}_i, i = 1, 2$ is everywhere smooth except, maybe, at $\omega_i$. So the topological embedding of $\bar{\ell}_i$ may be complicated in a neighborhood of the sink.

According to [1], $\ell_i$ is called *tame* (or *tamely embedded*) if there is a homeomorphism $\psi_i : W^s(\omega_i) \to \mathbb{R}^n$ such that $\psi_i(\omega_i) = O$, where $O$ is the origin and $\psi_i(\bar{\ell}_i \setminus \sigma)$ is a ray starting from $O$. In the opposite case $\ell_i$ is called *wild*. It follows from a criterion in [4] that the tameness of $\ell_i$ is equivalent to the existence of a smooth 3-ball $B_i$ around $\omega_i$ in any neighborhood of $\omega_i$ such that $\ell_i \cap \partial B_i$ consists of exactly one point. Using lemma 4.1 from [3] it is possible to make this criterion more precise in our dynamical setting: $\ell_i$ is tame if and only if there is 3-ball $B_{\omega_i}$ such that $\omega_i \in f(B_{\omega_i}) \subset int\ B_{\omega_i} \subset W^s(\omega_i)$ and $\ell_i \cap \partial B_{\omega_i}$ consists of exactly one point.

It was proved in [2] that, for every diffeomorphism $f \in \mathcal{G}_4$, at least one separatrix ($\ell_1$ say) is tame. It was also shown that the topological classification of diffeomorphisms from $\mathcal{G}_4$ is reduced to the embedding classifications of the separatrix $\ell_2$; hence there are infinitely many diffeomorphisms from $\mathcal{G}_4$ which are not topologically conjugate.

To characterize a type of embedding of $\ell_2$ we introduce some special Heegaard splitting of $\mathbb{S}^3$. Let us recall that a three-dimensional orientable manifold is *a handlebody of genus* $g \geq 0$ if it is obtained from a 3-ball by an orientation reversing identification of $g$ pairs of pairwise disjoint 2-discs in its boundary. The boundary of such a handlebody is an orientable surface of genus $g$.

Let $P^+ \subset \mathbb{S}^3$ be a handlebody of genus $g$ such that $P^- = \mathbb{S}^3 \setminus intP^+$ is a handlebody (necessarily of the same genus as $P^+$). Then the pair $(P^+, P^-)$ is a Heegaard splitting of genus $g$ of $\mathbb{S}^3$ with Heegaard surface $S = \partial P^+ = \partial P^-$.

**Definition 1.2** *A Heegaard splitting $(P^+, P^-)$ of $\mathbb{S}^3$ is said to be adapted to $f \in G_4$, or $f$-adapted, if:*

*a) $\overline{W^u(\sigma)} \subset f(P^+) \subset int\ P^+$;*

*b) $W^s(\sigma)$ intersects $\partial P^+$ transversally and $W^s(\sigma) \cap P^+$ consists of a unique 2-disc.*

*An $f$-adapted Heegaard splitting $\mathbb{S}^3 = P^+ \cup P^-$ is said to be minimal if its genus is minimal among all $f$-adapted splittings.*

For each integer $k \geq 0$ we denote by $G_{4,k}$ the set of diffeomorphisms $f \in G_4$ for which the minimal $f$-adapted Heegaard splitting has genus $k$. It is easily seen that, for each $f \in G_{4,0}$, $\ell_2$ is tame and, according to [3], $f$ possesses an energy function. Conversely any diffeomorphism in $G_{4,k}$, $k > 0$, has no energy function (see [8]). Figure 1 shows the phase portrait of a diffeomorphism $G_{4,1}$. The main result of this paper is the following.

**Theorem 1** *Every quasi-energy function for a diffeomorphism $f \in G_{4,1}$ has exactly six critical points.*




Figure 1: A diffeomorphism from the class $G_{4,1}$

## 2 Recollection of Morse theory

According to Milnor ([6], section 3), we use the following definitions.

A compact $(n+1)$-dimensional *cobordism* is a triad $(W, L_0, L_1)$ where $L_0$ and $L_1$ are closed manifolds of dimension $n$ and $W$ is a compact $(n+1)$-dimensional manifold whose boundary consists of the disjoint union $L_0 \cup L_1$. It is an *elementary* cobordism when it possesses a Morse function $\varphi : W \to [0,1]$ with only one critical point and such that $\varphi^{-1}(i) = L_i$ for $i = 0, 1$. When the index of the unique critical point is $r$, one speaks of an elementary cobordism of index $r$.

In this situation, $L_1$ is obtained from $L_0$ by a *surgery* of index $r$, that is: there is an embedding $h : \mathbb{S}^{r-1} \times \mathbb{D}^{n-r+1} \to L_0$ such that $L_1$ is diffeomorphic to the manifold obtained from $L_0$ by removing the interior of the image of $h$ and gluing $\mathbb{D}^r \times \mathbb{S}^{n-r}$, or

$$L_1 \cong \mathbb{D}^r \times \mathbb{S}^{n-r} \bigcup_{h|_{\mathbb{S}^{r-1}\times\mathbb{S}^{n-r}}} L_0 \setminus int\ (h(\mathbb{S}^{r-1} \times \mathbb{D}^{n-r+1}))\,.$$

Conversely, the following statement holds (see [6], Theorem 3.12):

**Statement 2.1** *If $L_1$ is obtained from $L_0$ by a surgery of index $r$, then there exists an elementary cobordism $(W, L_0, L_1)$ of index $r$.*

On figure 2 it is seen a surgery of index 1 from the 2-sphere to the 2-torus with some level sets of a Morse function on the corresponding elementary cobordism.

Finally, we recall the weak Morse inequalities (see [5], Theorem 5.2).




Figure 2: An elementary cobordism

**Statement 2.2** *Let $M^n$ be a closed manifold, $\varphi : M^n \to \mathbb{R}$ be a Morse function, $C_q$ be the number of critical points of index $q$ and $\beta_q(M^n)$ be the $q$-th Betti number of the manifold $M^n$. Then $\beta_q(M^n) \leq C_q$ and the Euler characteristic $\chi(M^n) := \sum\limits_{q=0}^{n}(-1)^q\beta_q(M^n)$ equals $\sum\limits_{q=0}^{n}(-1)^q C_q$.*

# 3 Proof of Theorem 1

Let $f$ be a Morse-Smale diffeomorphism of the 3-sphere belonging to $G_{4,1}$. As the number of critical points of any Morse function on a closed 3-manifold is even (it follows from statement 2.2) and greater than four (as $Per(f) \subset Cr(\varphi)$ and $\ell_2$ is wild) then, for proving theorem 1, it is enough to construct a Lyapunov function with six critical points.

## 3.1 Auxiliary statements

For the proof of the following statements 3.1 and 3.2 we refer to [3], lemma 2.2 and lemma 4.2.

**Statement 3.1** *Let $p$ be a fixed point of a Morse-Smale diffeomorphism $f : M^n \to M^n$ such that $\dim W^u(p) = q$. Then, in some neighborhood $U_p$ of $p$, there exist local coordinates $x_1, \ldots, x_n$ vanishing at $p$ and an energy function $\varphi_p : U_p \to \mathbb{R}$ such that*

$$\varphi_p(x_1, \ldots, x_n) = q - x_1^2 - \ldots - x_q^2 + x_{q+1}^2 + \ldots + x_n^2$$

*and $(TW^u(p) \cap U_p) \subset Ox_1 \ldots x_q$, $(TW^s(p) \cap U_p) \subset Ox_{q+1} \ldots x_n$.*

**Statement 3.2** *Let $\omega$ be a fixed sink of a Morse-Smale diffeomorphism $f : M^3 \to M^3$ and $B_\omega$ be a 3-ball with boundary $S_\omega$ such that $\omega \in f(B_\omega) \subset int\ B_\omega \subset W^s(\omega)$. Then there exists an energy function $\varphi_{B_\omega} : B_\omega \to \mathbb{R}$ for $f$ having $S_\omega$ as a level set.*

**Lemma 3.3** *Let $\omega$ be a fixed sink of a Morse-Smale diffeomorphism $f : M^3 \to M^3$ and $Q_\omega$ be a solid torus such that $\omega \in f(Q_\omega) \subset int\ Q_\omega \subset W^s(\omega)$. Then there exists a 3-ball $B_\omega$ such that $f(Q_\omega) \subset B_\omega \subset int\ Q_\omega$.*

**Proof:** Let $D_0$ be a meridian disk in $Q_\omega$ such that $\omega \notin D_0$. As $Q_\omega \subset W^s(\omega)$ there is an integer $N$ such that $f^n(Q_\omega) \cap D_0 = \emptyset$ for every $n > N$. We may also assume that $D_0$ is transversal to $G = \bigcup\limits_{n\in\mathbb{Z}} f^n(\partial Q_\omega)$, and hence $G \cap int\ D_0$ consists of a finite family $\mathcal{C}_{D_0}$ of intersection curves. Each intersection curve $c \in \mathcal{C}_{D_0}$ belongs to $f^k(\partial Q_\omega)$ for some integer $k \in \{1, \dots, N\}$. There are two cases: (1) $c$ bounds a disk on $f^k(\partial Q_\omega)$; (2) $c$ does not bound a disk on $f^k(\partial Q_\omega)$. Let us decompose $\mathcal{C}_{D_0}$ as union of two pairwise disjoint parts $\mathcal{C}^1_{D_0}$ and $\mathcal{C}^2_{D_0}$ consisting of curves with property (1) or (2), accordingly.

Let us show that there is a meridian disk $D_1$ in $Q_\omega$ such that $D_1$ is transversal to $G$ and $G \cap int\ D_1$ consists of family $\mathcal{C}_{D_1} = \mathcal{C}^2_{D_0}$ of intersection curves. If $\mathcal{C}^1_{D_0} = \emptyset$ then $D_1 = D_0$. In the opposite case for any curve $c \in \mathcal{C}^1_{D_0}$ denote by $d_c$ the disk on $f^k(\partial Q_\omega)$ such that $\partial d_c = c$. Notice that $d_c$ does not contain a curve from the family $\mathcal{C}^2_{D_0}$. Then there is $c \in \mathcal{C}_{D_1}$ which is innermost on $f^k(\partial Q_\omega)$ in the sense that the interior of $d_c$ contains no intersection curves from $\mathcal{C}_{D_0}$. For such a curve $c$ denote $e_c$ the disk on $D_0$ such that $\partial e_c = c$. As $int\ Q_\omega \setminus D_0$ is an open 3-ball then $e_c \cup d_c$ bounds a unique 3-ball $b_c \subset int\ Q_\omega$. Set $D'_c = (D_0 \setminus e_c) \cup d_c$. There is a smooth approximation $D_c$ of $D'_c$ such that $D_c$ is a meridian disk on $Q_\omega$, $D_c$ is transversal to $G$. Moreover $G \cap int\ D_c$ consists of a family $\mathcal{C}_{D_c}$ of intersection curves having less elements than $\mathcal{C}_{D_0}$; indeed, $c$ disappeared and also all curves from $\mathcal{C}_{D_0}$ lying in $int\ e_c$. We will repeat this process until getting a meridian disk $D_1$ with the required property.

Now let $c \in \mathcal{C}_{D_1}$, $c \in f^k(\partial Q_\omega)$. Denote $e_c$ the disk that $c$ bounds in $D_1$. Let us choose $c$ innermost in $D_1$ in the sense that the interior of $e_c$ contains no intersection curves from $\mathcal{C}_{D_1}$. There are two cases: (a) $e_c \subset f^k(Q_\omega)$ and (b) $int\ e_c \cap f^k(Q_\omega) = \emptyset$.

In case (a) $e_c$ is a meridian disk of $f^k(Q_\omega)$ and $D = f^{-k}(e_c)$ is a meridian disks in $Q_\omega$ such that $f(Q_\omega) \cap D = \emptyset$. Indeed, by construction $int\ e_c \cap G = \emptyset$, hence $int\ D \cap G = \emptyset$. Thus we can find the required 3-ball $B_\omega$ inside $int\ Q_\omega \setminus D_1$.

In case (b) there is a tubular neighborhood $V(e_c) \subset int\ Q_\omega$ of the disk $e_c$ such that $G \cap int\ V(e_c) = \emptyset$ and $B_k = f^k(Q_\omega) \cup V(e_c)$ is 3-ball. Then $f^k(Q_\omega) \subset B_k \subset int\ f^{k-1}(Q_\omega)$. Thus $B_\omega = f^{1-k}(B_k)$ is the required 3-ball. $\diamond$

## 3.2 Construction of a quasi-energy function for a diffeomorphism $f \in G_{4,1}$

As a similar construction was done in section 4.3 of [3], we only give a sketch of it below.

1. Construct an energy function $\varphi_p : U_p \to \mathbb{R}$ near each fixed point $p$ of $f$ as in statement 3.1.




Figure 3: Heegaard decomposition $(Q^+, Q^-)$

2. By definition of the class $G_{4,1}$, for each $f \in G_{4,1}$ there is a solid torus $P^+$ belonging to a Heegaard splitting $(P^+, P^-)$ of $\mathbb{S}^3$ and such that:

   a) $\overline{W^u(\sigma)} \subset f(P^+) \subset int\ P^+$;

   b) $W^s(\sigma)$ intersects $\partial P^+$ transversally and $W^s(\sigma) \cap P^+$ consists of a unique 2-disk.

   As $\mathbb{S}^3 \setminus \overline{W^s(\sigma)}$ is the disjoint union $W^s(\omega_1) \cup W^s(\omega_2)$, then by property b), the disk $P^+ \cap W^s(\sigma)$ is separating in $P^+$. Moreover there exists a neighborhood of $P^+ \cap W^s(\sigma)$, such that after removing it from $P^+$ we get a 3-ball $P_{\omega_1}$ and solid torus $P_{\omega_2}$ with the following properties for each $i = 1, 2$:

   i) $\omega_i \in f(P_{\omega_i}) \subset int\ P_{\omega_i} \subset W^s(\omega_i)$;

   ii) $\partial P_{\omega_i}$ is a Heegaard surface and $\ell_i \cap \partial P_{\omega_i}$ consists of exactly one point.

   Due to the $\lambda$-lemma[1] (see, for example,[7]), replacing $P_{\omega_i}$ by $f^{-n}(P_{\omega_i})$ for some $n > 0$ if necessary, we may assume that $\partial P_{\omega_i}$ is transversal to the regular part of the critical level set $C := \varphi_\sigma^{-1}(1)$ of the function $\varphi_\sigma$ and the intersections $C \cap \partial P_{\omega_i}$ consist of exactly one circle. For $\varepsilon \in (0, \frac{1}{2})$ define $H_\varepsilon^+$ as the closure of $\{x \in U_\sigma \mid x \notin (P_{\omega_1} \cup P_{\omega_2}),\ \varphi_\sigma(x) \leq 1+\varepsilon\}$ and set $P_\varepsilon^+ = P_{\omega_1} \cup P_{\omega_2} \cup H_\varepsilon^+$. In the same way as in [3] it is possible to choose $\varepsilon > 0$ such that $\partial P_{\omega_i}$ intersects transversally each level set with value in $[1-\varepsilon,\ 1+\varepsilon]$; this intersection consists of one circle. Taking a smoothing $Q^+$ of $P_\varepsilon^+$ we have $f(Q^+) \subset int\ Q^+$ and $\Sigma := \partial Q^+$ is a Heegaard surface of genus 1. Let $Q^-$ be the closure of $\mathbb{S}^3 \setminus int\ Q^+$ (see

[1]The $\lambda$-lemma claims that $f^{-n}(S_{\omega_i}) \cap U_\sigma$ tends to $\{x_1 = 0\} \cap U_\sigma$ in the $C^1$ topology when $n$ goes to $+\infty$.

figure 3). It is easy to check that $Q^+$ is isotopic to $P^+$. Therefore, the pair $(Q^+, Q^-)$ is an $f$-adapted Heegaard splitting with the property that the disk $Q^+ \cap W^s(\sigma)$ lies in $U_\sigma$.

3. For each $i = 1, 2$, let $\tilde{P}_{\omega_i}$ be a handlebody of genus $i-1$ such that $f(P_{\omega_i}) \subset \tilde{P}_{\omega_i} \subset int\ P_{\omega_i}$, $\partial\tilde{P}_{\omega_i}$ intersects transversally each level set with value in $[1-\varepsilon, 1+\varepsilon]$ along one circle and $P_{\omega_i} \setminus int\ \tilde{P}_{\omega_i}$ is diffeomorphic to $\partial P_{\omega_i} \times [0,1]$. Define $d_i$ as the closure of $\{x \in U_\sigma \mid x \in (W^s(\omega_i) \setminus \tilde{P}_{\omega_i}),\ \varphi_\sigma(x) = 1-\varepsilon\}$. By construction $d_i$ is a disk whose boundary curve bounds a disk $D_i$ in $\partial\tilde{P}_{\omega_i}$. We form $S_i$ by removing the interior of $D_i$ from $\partial\tilde{P}_{\omega_i}$ and gluing the $d_i$. Denote $P(S_i)$ the handlebody of genus $i-1$ bounded by $S_i$ and containing $\omega_i$. As in [3] it is possible to choose $\varepsilon$ such that $f(P(S_i)) \subset int\ P(S_i)$.

   Let $K$ be the domain between $\partial Q^+$ and $S_1 \cup S_2$. We introduce $T^+$, the closure of $\{x \in \mathbb{S}^3 \mid x \notin (P_{\omega_1} \cup P_{\omega_2}),\ 1-\varepsilon \leq \varphi_\sigma(x) \leq 1+\varepsilon\}$; observe $T^+ \subset U_\sigma$. We define a function $\varphi_K : K \to \mathbb{R}$ whose value is $1+\varepsilon$ on $\partial Q^+$, $1-\varepsilon$ on $S_1 \cup S_2$, coinciding with $\varphi_\sigma$ on $K \cap T^+$ and without critical points outside $T^+$. This last condition is easy to satisfy as the domain in question is a product cobordism. In a similar way to [3], section 4.3, one can check that $\varphi_K$ is a Morse-Lyapunov function.

4. As $P(S_1)$ is a 3-ball such that $\omega_1 \in f(P(S_1)) \subset int\ P(S_1) \subset W^s(\omega_1)$, then by statement 3.2 there is an energy function $\varphi_{P(S_1)} : P(S_1) \to \mathbb{R}$ for $f$ with $S_1$ as a level set with value $1-\varepsilon$.

5. As $P(S_2)$ is a solid torus such that $\omega_2 \in f(P(S_2)) \subset int\ P(S_2) \subset W^s(\omega_2)$, then according to lemma 3.3 there is a 3-ball $B_{\omega_2}$ such that $f(P(S_2)) \subset B_{\omega_2} \subset int\ P(S_2)$. As in the previous item, there is an energy function $\varphi_{B_{\omega_2}} : B_{\omega_2} \to \mathbb{R}$ for $f$ with $\partial B_{\omega_2}$ as a level set with value $\frac{1}{2}$.

6. As $P(S_2)$ is a solid torus, it is obtained from a 3-ball by an orientation reversing identification of a pair of disjoint 2-discs in its boundary; hence the solid torus is the union of a 3-ball and an elementary cobordism of index 1. Since, up to isotopy, there is only one 3-ball in the interior of a solid torus, then $(W_{\omega_2}, \partial B_{\omega_2}, S_2)$ is an elementary cobordism of index 1, where $W_{\omega_2} = P(S_2) \setminus int\ B_{\omega_2}$. Hence $W_{\omega_2}$ possesses a Morse function $\varphi_{W_{\omega_2}}$ with only one critical point of index 1 and such that $\varphi_{W_{\omega_2}}(\partial B_{\omega_2}) = \frac{1}{2}$, $\varphi_{W_{\omega_2}}(S_2) = 1-\varepsilon$.

7. Define the smooth function $\varphi^+ : Q^+ \to \mathbb{R}$ by the formula

$$\varphi^+(x) = \begin{cases} \varphi_K(x), x \in K; \\ \varphi_{P(S_1)}(x), x \in P(S_1); \\ \varphi_{B_{\omega_2}}(x), x \in B_{\omega_2}; \\ \varphi_{W_{\omega_2}}(x), x \in W_{\omega_2}. \end{cases}$$

   Then $\varphi^+$ is a Morse-Lyapunov function for $f|_{Q^+}$ with one additional critical point.

8. By the construction $Q^-$ is a solid torus such that $\alpha \in f^{-1}(Q^-) \subset int\ Q^- \subset W^u(\alpha)$. Since $\alpha$ is a sink for $f^{-1}$ then, as in item 4, there is a 3-ball $B_\alpha$ such that $f^{-1}(Q^-) \subset B_\alpha \subset int\ Q^-$ and an energy function $\varphi_{B_\alpha} : B_\alpha \to \mathbb{R}$ for $f^{-1}$ with $\partial B_\alpha$ as a level set of value $\frac{1}{2}$.

9. Similarly to item 5, $\partial Q^-$ is obtained from $\partial B_\alpha$ by a surgery of index 1. Therefore $(W_\alpha, \partial Q^-, \partial B_\alpha)$ is an elementary cobordism of index 1, where $W_\alpha = Q^- \setminus int\ B_\alpha$. Hence, $W_\alpha$ possesses a Morse function $\varphi_{W_\alpha}$ with only one critical point of index 1. We may choose $\varphi_{W_\alpha}(\partial B_\alpha) = \frac{1}{2}$, $\varphi_{W_\alpha}(\partial Q^-) = 2 - \varepsilon$.

10. Define the smooth function $\varphi^- : Q^- \to \mathbb{R}$ by the formula

$$\varphi^-(x) = \begin{cases} 3 - \varphi_{B_\alpha}(x), x \in \varphi_{B_\alpha}; \\ 3 - \varphi_{W_\alpha}(x), x \in \varphi_{W_\alpha}. \end{cases}$$

Then $\varphi^-$ is a Morse-Lyapunov function for $f|_{Q^-}$ with one additional critical point.

11. The function $\varphi : \mathbb{S}^3 \to \mathbb{R}$ defined by $\varphi|_{Q^+} = \varphi^+$ and $\varphi|_{Q^-} = \varphi^-$ is the required Morse-Lyapunov function for the diffeomorphism $f$ with exactly six critical points.

# References


[1] E. Artin, R. Fox, *Some wild cells and spheres in three-dimensional space*, Annals of Math. (1948) 49, 979-990.

[2] Ch. Bonatti, V. Grines, *Knots as topological invariant for gradient-like diffeomorphisms of the sphere $S^3$*, Journal of Dynamical and Control Systems (2000) 6, 579-602.

[3] V. Grines, F. Laudenbach, O. Pochinka, *Self-indexing energy function for Morse-Smale diffeomorphisms on 3-manifolds*, submitted to published.

[4] O.G. Harrold, H.C. Griffith, E.E. Posey, *A characterization of tame curves in three-space,* Trans. Amer. Math. Soc. (1955) 79, 12-34.

[5] J. Milnor, *Morse theory,* Princeton University Press, 1963.

[6] J. Milnor, *Lectures on the h-cobordism Theorem,* Princeton University Press, 1965.

[7] J. Palis, *On Morse-Smale dynamical systems,* Topology (1969) 8, 385-404.

[8] D. Pixton, *Wild unstable manifolds,* Topology (1977) 16, 167-172.

[9] S. Smale, *Differentiable dynamical systems,* Bull. Amer. Math. Soc. (1967) 73, 747-817.