



HAL
open science

Comparative bioaccumulation of trace elements between Nautilus pompilius and N. macromphalus (Cephalopoda: Nautiloidea) from Vanuatu and New-Caledonia

Mathieu Pernice, Julien Boucher, Renata Boucher-Rodoni, Pascale Joannot, Paco Bustamante

► To cite this version:

Mathieu Pernice, Julien Boucher, Renata Boucher-Rodoni, Pascale Joannot, Paco Bustamante. Comparative bioaccumulation of trace elements between *Nautilus pompilius* and *N. macromphalus* (Cephalopoda: Nautiloidea) from Vanuatu and New-Caledonia. *Ecotoxicology and Environmental Safety*, 2009, 72 (2), pp.365-371. <10.1016/j.ecoenv.2008.04.019>. <hal-00336127>

HAL Id: hal-00336127

<https://hal.science/hal-00336127v1>

Submitted on 1 Nov 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **Comparative bioaccumulation of trace elements between *Nautilus pompilius* and *N.***
2 ***macromphalus* (Cephalopoda: Nautiloidea) from Vanuatu and New-Caledonia**

3

4 Running title: influence of the environment on bioaccumulation of trace elements in *Nautilus*

5

6 PERNICE Mathieu¹, BOUCHER Julien², BOUCHER-RODONI Renata¹, JOANNOT
7 Pascale³, BUSTAMANTE Paco^{4#}

8

9 ¹ UMR 5178 Biologie des Organismes Marins et Ecosystèmes, DMPA, Muséum National
10 d'Histoire Naturelle, 55 rue Buffon, 75005 Paris, France.

11 ² Ecole Polytechnique Fédérale de Lausanne, LGCB-ISIC-SB-EPFL1015 Lausanne,
12 Switzerland.

13 ³ Direction des Collections, Muséum National d'Histoire Naturelle, Cour Fagon, 57 rue
14 Cuvier, 75005 Paris, France.

15 ⁴ Littoral Environnement et Sociétés (LIENSs), UMR 6250 CNRS-Université La Rochelle, 2
16 Rue Olympe de Gouges, 17042 La Rochelle Cedex 01, France

17

18 Correspondence to: Dr. Paco BUSTAMANTE
19 Littoral Environnement et Sociétés (LIENSs),
20 2 Rue Olympe de Gouges
21 F-17042 La Rochelle Cedex 01
22 France
23 Phone : +33 5 465 076 25
24 Fax : +33 5 464 582 64
25 E-mail : pbustama@univ-lr.fr

26

27 **ABSTRACT:** The concentrations of sixteen trace elements were investigated and compared
28 for the first time in the digestive and excreting tissues of two *Nautilus* species (Cephalopoda:
29 Nautiloidea) from two geologically contrasted areas: 1) *N. macromphalus* from New
30 Caledonia, a region characterised by its richness in nickel ores and its lack of tectonic
31 activities and 2) *N. pompilius* from the Vanuatu archipelago showing high volcanic and
32 tectonic activities. In both *Nautilus* species, results clearly highlighted that the digestive gland
33 played a key role in the bioaccumulation and storage of Ag, Cd, Ce, Co, Cu, Fe, La, Nd, V
34 and Zn whereas As, Cr, Mn, Ni, Pb and Se were accumulated in a greater extent in the
35 excreting tissues (i.e. pericardial and renal appendages). Despite contrasting environments, no
36 significant difference ($P < 0.05$) was found between the two *Nautilus* species in the
37 concentrations of most of the essential and non essential elements, including Ni and
38 associated metals in Ni ores (i.e. Co and Mn). As nautilus lives on the outer shelf of barrier
39 reefs, these results strongly support the hypothesis that the New Caledonian lagoon traps the
40 major amount of the trace elements derived from natural erosion and the intense mining
41 activities conducted on land. In contrast, the concentrations of the rare earth elements (Ce, La,
42 and Nd) were significantly higher in *N. pompilius* than in *N. macromphalus*, probably as a
43 result of the local enrichment of Vanuatu waters by specific environmental processes, such as
44 volcanism or upwelling.

45

46 **Keywords:** metals, metalloids, cephalopods, detoxification, excretion, South Pacific

47 1. INTRODUCTION

48

49 Marine organisms are exposed to trace elements that are present in their diet and dissolved in
50 seawater. This double exposure results in element accumulation in their different tissues
51 whether or not these elements are essential to the metabolism (Rainbow, 2002). This process,
52 widely described as bioaccumulation can vary greatly according to the specificities of the
53 organisms such as age/size, sex and lifestyle, and according to the relative bioavailability of
54 the metal in diet and seawater (Rainbow and Wang, 2001). Comparative analysis of trace
55 element concentrations between closely related species living in different geographical areas
56 can therefore allow at assessing the influence of the environment on their bioaccumulation.

57 Among the marine species, cephalopods belong to a molluscan group comprising ca. seven
58 hundred species (Boyle and Rodhouse, 2005), in which high concentrations of trace elements
59 such as Ag, Cd, Cu or Zn have generally been recorded (Martin and Flegal, 1975; Finger and
60 Smith, 1987; Miramand and Bentley, 1992; Yamada et al., 1997; Bustamante et al., 1998;
61 Dorneles et al., 2007). Indeed, several studies have demonstrated that the digestive and the
62 excretory organs of cephalopods play a major role in the bioaccumulation of trace elements,
63 as these organs are deeply involved in the assimilation processes, detoxification and storage
64 of both essential and non-essential metals (Miramand and Guary, 1980; Miramand and
65 Bentley, 1992; Bustamante et al., 2000, 2002a,b, 2004).

66 In the present work, we focused on the bioaccumulation of trace elements in the *Nautilus*
67 genus. Nautiluses are scavenger cephalopods, inhabiting the outer shelf of barrier reefs in the
68 Indo-Pacific area, generally at a depth of 300 to 500 meters (O'Dor et al., 1993; Norman,
69 2000). In spite of its evolutionary interest (*Nautilus* is the last representative of ectocochleate
70 cephalopods, i.e. considered as a "living fossil"; Boyle and Rodhouse, 2005) and its
71 physiological specificities (slow digestion and most of the excretory processes taking place in

72 the pericardial appendages; Schipp et al., 1985; Westermann et al., 2002), only very few
73 studies deal with the bioaccumulation of trace elements in this ancestral genus. High
74 concentrations of several metals and metalloids were reported in the digestive and excretory
75 tissues of *Nautilus macromphalus* (Bustamante et al., 2000), a species endemic to the New
76 Caledonian waters. These authors suggested that the high metal concentrations may be due to
77 a high exposure in the metal-rich conditions in the surrounding New Caledonian waters.
78 Indeed, the largest resources of Ni as laterites in the world are present in New Caledonia
79 where open-cast mining is conducted since the 20th century, leading to a dramatic increase of
80 metal concentrations in coastal waters and, subsequently, the metal burdens in marine biota
81 (Labrosse et al., 2000; Metian et al., 2008). However, it is not clear whether metal
82 accumulation observed in *N. macromphalus* reflected the impact of metal contamination
83 through human activities on the outer shelf waters of New Caledonia or either if it is more due
84 to a natural progressive bioaccumulation during nautilus long life span (*viz.* 10-15 years).

85
86 The aim of the present study was therefore to determine the bioaccumulation of a large range
87 of elements in the genus *Nautilus* comparing two species from geologically contrasted areas:
88 1) *N. macromphalus* from New Caledonia which is characterised by its richness in nickel ores
89 and its lack of tectonic activities and 2) *N. pompilius* from the Vanuatu archipelago which
90 shows high volcanic and tectonic activities (Harrison et al, 1996). The analysed elements
91 comprised 11 metals (Ag, Cd, Co, Cr, Cu, Fe, Mn, Ni, Pb, V and Zn), 2 metalloids (As and
92 Se) and 3 rare earth elements (Ce, La and Nd). These elements were selected for comparison
93 with other species and other published works concerning Nautilus (e.g. Miramand et al. 1992,
94 Bustamante et al. 2000, Ichihashi et al. 2001). The concentrations and relative contents of
95 these different trace elements were determined in the digestive and the excretory tissues (i.e.
96 pericardial and renal appendages) and compared between both nautilus species. The

97 specificities and the potential influence of the environment on bioaccumulation of these trace
98 elements in the *Nautilus* genus are discussed.

99

100 **2. MATERIALS AND METHODS**

101 **a) Biological material**

102 Three *Nautilus pompilius* and five *Nautilus macromphalus* specimens were collected
103 respectively off Santo island coast (Vanuatu, 15° 40' 522 S, 167° 02' 838 E, depth: 334m) and
104 on the outer shelf of New-Caledonia (22° 20' 500 S, 166° 13' 400 E, depth: 300m) in May
105 2006 (Figure 1). Each individual was measured and weighted prior to dissection. The main
106 features of the eight specimens sampled are reported in Table 1.

107 All specimens were dissected aseptically and for each specimen one pericardial appendage,
108 one renal appendage and a fraction of the digestive gland were immediately frozen in liquid
109 nitrogen and stored at -80°C for metal analyses.

110

111 **b) Analytical procedure**

112 Tissue samples were freeze-dried. Aliquots of the samples ranging from 20 to 300 mg were
113 digested using a 3:1 v:v nitric-hydrochloric acid mixture with 65% ultrapure HNO₃ and
114 ultrapure 70% HCl. The acidic digestion was performed overnight under ambient temperature
115 and then heated in a microwave during 30 minutes, with increasing temperature until 105°C,
116 and 15 minutes at 105°C (1200 Watts). After the mineralization process, each sample was
117 diluted to 30 ml or 50 ml with milli-Q quality water, according to the volume of acid added to
118 the mineralization (3 ml and 4.5 ml. respectively).

119 Ag, As, Cd, Ce, Co, Cr, Cu, Fe, La, Mn, Nd, Ni, Pb, Se, V and Zn were analysed either by
120 ICP-OES (Varian[®] Vista-Pro) and ICP-MS (Varian[®] Ultra Mass 700). Reference tissues,
121 dogfish liver DOLT-3 (NRCC), lobster hepatopancreas TORT-2 (NRCC) and Bush Branches

122 and Leaves (GBW 07602) were treated and analysed in the same way as the samples. Results
123 were in good agreement with the certified values, and the standard deviations were low,
124 proving good repeatability of the method. The results for standard reference materials
125 displayed recoveries of the elements ranging from 81% to 119% (n=10). For each set of
126 analyses, blanks were included in each analytical batch. The detection limits ($\mu\text{g.g}^{-1}$ dry wt)
127 for ICP-OES were 8.3 (As, Fe, Zn), 3.3 (Ag, Se), 1.67 (Pb, V), 0.83 (Cd, Co, Cr, Cu, Mn, Ni),
128 and for ICP-MS, they were 0.150 (Ni, V), 0.065 (Cd, Co, Cr, Cu, Mn, Pb), 0.033 (Ag), 0.017
129 (Nd), 0.008 (Ce, La). Average element concentrations are presented with standard deviations.
130 All trace element concentrations are given on a dry weight basis ($\mu\text{g.g}^{-1}$ dry wt).

131

132 *Data analyses*

133 Statistical analyses were performed using MINITAB 13.2 Software. Comparison of trace
134 element concentrations in nautilus tissues between sites was tested by 1-way ANOVA.
135 Hypothesis of normal distribution was tested using the Anderson-Darling test and equality of
136 variance by the Bartlett test. Pearson coefficients were calculated between trace elements in
137 each tissue. The significance level for statistical analyses was always set at $\alpha = 0.05$.

138

139 **3. RESULTS**

140

141 *Trace element concentrations in the tissues*

142 Metal concentrations in the digestive gland, the renal appendages and the pericardial
143 appendages of *N. macromphalus* and *N. pompilius* are reported in Table 2. La, Ce, and Nd
144 concentrations fell below the detection limit in the renal appendages. Among the three tissues
145 analysed, the digestive gland was the major site of accumulation for Ag, Cd, Ce, Co, Cu, Fe,
146 La, Nd, V and Zn (Table 2). The digestive gland also concentrated As and Ni at

147 concentrations close to the highest ones recorded in the other tissues. As, Cr, Mn, Ni, Pb, and
148 Se concentrations were remarkable in excreting tissues. More specifically, the renal
149 appendages exhibited the highest Mn and Se concentrations, whereas As, Ni, and Pb were
150 concentrated at similar concentrations within both renal and pericardial appendages. Finally,
151 the pericardial appendages were the major site of accumulation for Cr.

152

153 *Comparison between species*

154 Generally, the pattern of trace element concentrations was the same in both *N. macromphalus*
155 and *N. pompilius* tissues. However, there were few exceptions and more particularly in the
156 digestive gland: firstly, Ce, Fe, La, and Nd concentrations were significantly ($P < 0.05$) higher
157 in *N. pompilius*; secondly, Cd and Cr concentrations were significantly ($P < 0.05$) higher in *N.*
158 *macromphalus* (Table 3).

159 In the case of the pericardial appendages, As, La, and Nd concentrations were significantly
160 higher ($P < 0.05$) in *N. pompilius* (Table 3). For the renal appendages, Cu was the only element
161 for which a significant difference was detected, *N. pompilius* having significantly higher
162 concentrations than *N. macromphalus* (Table 3).

163

164 *Relative proportion of trace elements between tissues*

165 Among the three tissues considered, the digestive gland contained the largest proportions of
166 most of the trace elements with 73 to 81% of Ag, 90 to 91% of Cd, 61 to 79% of Ce, 80 to
167 86% of Co, 67 to 72% of Cu, 90 to 93% of Fe, 72 to 81% of La, 55 to 75% of Nd, 56 to 67 of
168 V, 81% to 82% of Zn in both *Nautilus* species (Fig. 2). The only exception were As, Cr, Mn,
169 Ni, Se, and Pb which were mainly contained in the excreting tissues, i.e. renal and pericardial
170 appendages (Fig. 2). The renal appendages were the major site of accumulation of Mn, Ni,
171 and Se (i.e. 74 to 89% for Mn, 49 to 54% for Ni; 61 to 73% for Se) while the pericardial

172 appendages accumulated the highest relative proportion of Cr (i.e. 63 to 74%). Although the
173 concentrations of some of the trace elements were high in the excretory organs, these tissues
174 contained in fact low amounts of metals because of their small masses.

175

176 *Correlations between metals*

177 Table 4 shows the correlation between trace elements within the three tissues. It is noteworthy
178 that the rare earth elements were correlated between each other within the digestive gland ($r >$
179 0.985 and $P < 0.001$ in all cases) and the pericardial appendages ($r > 0.922$ and $P < 0.003$ in all
180 cases). In the digestive gland, rare earth elements varied positively with Fe and Mn ($r > 0.918$
181 and $r > 0.807$ and $P < 0.004$ and $P < 0.028$, respectively) but negatively with As ($r > -0.726$ and
182 $P = 0.047$). Cu correlated positively with Zn ($r = 0.834$ and $P = 0.020$ in all cases) but negatively
183 with Se in this tissue ($r = -0.809$ and $P = 0.028$ in all cases).

184 In the renal appendages, correlations were weak, Mn varied positively with Cd ($r = 0.833$,
185 $P = 0.010$) and Zn ($r = 0.735$, $P = 0.038$), Fe with Zn ($r = 0.833$, $P = 0.010$) and Pb with Cr
186 ($r = 0.936$, $P = 0.001$) but negatively with V ($r = -0.753$, $P = 0.031$; Table 3).

187 In the pericardial appendages, Cd concentrations correlated positively with Cr ($r = 0.897$,
188 $P = 0.006$), Ni ($r = 0.965$, $P < 0.001$), and V ($r = 0.931$, $P = 0.002$).

189

190 **4. DISCUSSION**

191

192 The present extensive analysis of sixteen trace element concentrations in *N. macromphalus*
193 and *N. pompilius* digestive and excreting tissues clearly highlighted that these organs
194 constitute important tissues for the bioaccumulation of rare earth elements, metals and
195 metalloids. Such results support the hypothesis of efficient regulation processes for non

196 essential metals in Nautilidae (Bustamante et al., 2000) even if, among the eight studied
197 specimens, the variability of metal concentrations was relatively high (Table 2).

198 In cephalopods, the digestive system plays a major role in both the digestive and the
199 detoxification processes: on the one hand, the digestive gland synthesizes the digestive
200 enzymes and is a place of absorption and assimilation of nutrients (Boucaud-Camou and
201 Boucher-Rodoni, 1983; Boucher-Rodoni and Boucaud-Camou, 1987); on the other hand, this
202 organ is involved in the storage and detoxification of trace elements (Bustamante et al.,
203 2002ab, 2004). Our investigations on trace element concentrations also confirmed the key role
204 of the digestive gland of both *N. macromphalus* and *N. pompilius* in the metabolism of most
205 of the analysed trace elements (i.e. Ag, Cd, Ce, Co, Cu, Fe, La, Nd, V and Zn), as it is also the
206 case in more recent cephalopods, i.e. belonging to the Coleoidae (e.g. Ghiretti-Magaldi et al.,
207 1958; Martin and Flegal, 1975; Miramand and Guary, 1980; Smith et al., 1984; Finger and
208 Smith, 1987; Miramand and Bentley, 1992; Seixas et al., 2005a; Pierce et al., 2008).

209 Our results also stressed the potential of excreting tissues (i.e. renal and pericardial
210 appendages) to accumulate several elements such as As, Cr, Mn, Ni, Pb, and Se. In contrast to
211 the Coleoidae, in nautilus the renal appendages are specialized in the storage of calcium and
212 the pericardial appendages play a major role in the evacuation of the metabolic wastes
213 (Schipf and Martin, 1987). These organs are known to harbour a dual bacterial symbiosis
214 (Pernice et al., 2007) involving spirochaetes and betaproteobacteria. The exact mechanisms
215 by which these bacterial symbionts are involved in nautilus excretion are still enigmatic.
216 Nevertheless, it is widely admitted that micro-organisms can play an important role in the
217 incorporation of trace elements (Worms et al., 2006). Accumulation of Co, Fe and Ni by
218 betaproteobacterial symbionts of the pericardial appendages of *N. macromphalus* has already
219 been observed by electronic microscopy (Pernice et al., in preparation). However, it remains
220 unclear if this accumulative process is only passive (viz. through adsorption processes) or

221 active (viz. effective absorption of trace elements). Indeed, the betaproteobacterial symbionts
222 present in nautilus tissues could play a role in the detoxification and storage of some metals
223 such as Co, Fe and Ni (Table 2). Such processes could involve enzymatic reduction and/or
224 complexation with organic compounds (Pernice et al., 2006), leading to the storage of these
225 metals at high concentrations in the excreting tissues, independently of the nautilus species.
226 The role of this bacterial symbiosis on trace element bioaccumulation and detoxification in
227 nautilus excreting tissues clearly deserves further investigations.

228 Although Nautilidae have been previously reported for their special abilities to accumulate
229 heavy metals at relatively elevated concentrations in their tissues (Bustamante et al., 2000),
230 this study is the first comparative approach aiming to compare trace element concentrations in
231 two *Nautilus* species from two geographically close areas (viz. distant from less than 1000
232 kms). Both species potentially share common trophic features but, at the same time, they live
233 in geologically different environments (richness in nickel ores and lack of tectonic activities
234 in New Caledonia and high volcanic and tectonic activities in Vanuatu), potentially resulting
235 in a different degree of “natural” contamination. New Caledonia is currently the third largest
236 producer of Ni in the world (Dalvi et al., 2004) and mining activities associated to a strong
237 natural erosion have resulted in the enrichment of Ni and associated metals in Ni ores (i.e. Co,
238 Cr, Cu, Fe and Mn) and some other extracted metals (i.e. Ag and Pb) in the coastal waters and
239 consequently in the lagoon food webs (Monniot et al., 1994; Fernandez et al., 2006; Hédouin
240 et al., 2006; Metian et al., 2008). Conversely, it is likely that the seawater concentrations of
241 most of these metals are very low in Santo islands waters and around Vanuatu in general
242 according to the absence of industrial and mining activities in this area (Harrison et al., 1996).
243 Surprisingly, in the present study, very little difference appeared in the sixteen trace element
244 concentrations between *N. macromphalus* from New Caledonia and *N. pompilius* from
245 Vanuatu. Interestingly, there was no significant difference between both *Nautilus* species for

246 Ag, Co, Mn, Ni, Pb, Se, V and Zn (Table 3) indicating that the geographical origin does not
247 seem to be a major factor contributing to interspecific variations of these trace element
248 concentrations. As nautilus lives on the outer shelf of barrier reefs and rarely penetrates into
249 lagoons (Saunders and Ward, 1987; Norman, 2000), our results strongly supported the
250 hypothesis that the New Caledonian lagoon traps the major amount of the trace elements
251 derived from the island (Ambatsian et al., 1997). To the best of our knowledge, there are no
252 published data in the open literature concerning trace elements in organisms from the oceanic
253 waters off New Caledonia, except for two marine mammal species (Bustamante et al., 2003).
254 These authors reported very low concentrations of Co, Cr and Ni in the marine mammal
255 tissues which suggested that mining activity in New Caledonia does not seem to be a
256 significant source of contamination in the oceanic domain. Overall, the global geologic
257 contagion due to the mining activity in New Caledonia seems to be limited to the shoreline
258 waters and to the inner lagoon (Fernandez et al., 2006).
259 More interestingly, the concentrations of rare earth elements were significantly different
260 between the digestive gland and the pericardial appendages of *N. macromphalus* and *N.*
261 *pompilius*. F-transition elements of rare earths have similar physicochemical properties, and it
262 is therefore not surprising that they correlate to each other (Table 4). Our results suggested
263 that rare earth elements would reflect the characteristics of the surrounding environment.
264 Very little is known about the concentrations of Ce, La and Nd in cephalopods, which varied
265 from 0.045 to 1.95 $\mu\text{g}\cdot\text{g}^{-1}$ dwt in the digestive gland of the squid *Stenoteuthis oualaniensis*
266 (Ichihashi et al., 2001). In fact, rare earth elements have chemistry and sorption/complexation
267 characteristics analogous to transuranic elements such as americium or plutonium (Choppin
268 Torres and Chopin, 1984; Kim et al., 1989; Buckau et al., 1992). In Coleoid cephalopods,
269 transuranic elements also accumulate in the digestive and excreting tissues, in particular in the
270 branchial heart and their appendages (Guary et al., 1981; Guary and Fowler, 1982;

271 Bustamante et al., 2006). It is therefore not surprising that they accumulate at relatively high
272 concentrations in nautilus digestive gland and pericardial appendage.

273 Besides Ce, La and Nd, the concentrations of As, Cu and Fe were also higher in *N. pompilius*
274 than in *N. macromphalus*. It is difficult to ascertain the reasons of such a difference but this
275 might also be linked to geological features reflecting a potential enrichment of Vanuatu
276 waters by local environmental processes, such as volcanism or upwelling (Malinovsky and
277 Markevich, 2007). In fact, only two trace elements, Cd and Cr, displayed significantly higher
278 concentrations in *N. macromphalus* than in *N. pompilius* tissues (Table 2). Therefore, the
279 reasons of such interspecific differences in bioaccumulation, i.e. higher concentrations of
280 trace elements in one species, remain unclear whereas an absence of difference could be
281 explained by the similar physiology of these two species (Ward, 1988; Norman, 2000).

282 As for other marine organisms, in cephalopods, trace element concentrations may vary with
283 biological and environmental factors such as temperature, salinity, season, age (size), sex,
284 lifestyle and geographical origin (e.g. Monteiro et al., 1992; Bustamante et al., 1998;
285 Raimundo et al., 2004; Seixas et al., 2005b; Pierce et al., 2008). Further investigations are
286 needed to understand how exactly these factors affect their concentrations in the studied
287 species. However, among these factors, individual size/age is known to be of primary
288 importance in invertebrates (e.g. Warnau et al., 1995; Bustamante and Miramand, 2005;
289 Hédouin et al., 2007). In the present study, the size as well as the two others main features
290 measured (sex and weight) were globally homogenous among the eight studied specimens.
291 Because nautilus are long lived animals, it would be striking to determine the influence of
292 the size/age on trace elements bioaccumulation.

293 Sources of trace elements to nautilus can be (1) seawater, as it passes through the skin and
294 through the gills during respiration, and (2) food, which is now considered as a major route
295 for metal accumulation in marine animals, including both invertebrates and fish (Koyama et

296 al., 2000; Wang, 2002; Bustamante et al., 2002a, 2004). Knowledge of trace element
297 speciation in the direct environment of nautilus is a key factor to ascertain the relative
298 importance of each uptake pathways. Concerning the first pathway (i.e. seawater), further
299 aquatic toxicity studies are needed to determine the fraction of total metal dissolved in
300 seawater (the 'toxic fraction') that react with, and is transported across biological membranes
301 such as nautilus gills. Concerning the second pathway (i.e. food), our comparative analysis
302 suggested that the metal discharges due to intense mining activities conducted on land in New
303 Caledonia are not easily transferred in the food web of nautilus. Therefore, to assess the
304 impact of mining activities on cephalopods in the region of New Caledonia, further studies
305 should be conducted on more coastal species known to be resident in the impacted areas, such
306 as Sepiidae, Sepiolidae or Octopodidae spending all of their life in the waters of the New
307 Caledonian lagoon.

308

309 **5. ACKNOWLEDGMENTS**

310

311 We thank the Aquarium and the centre IRD of Nouméa (New Caledonia) as well as the
312 Maritime college of Luganville (Vanuatu) for their help in providing *Nautilus macromphalus*
313 and *N. pompilius* specimens, respectively. This work was supported by the Muséum National
314 d'Histoire Naturelle (pluriformations programme: “Ecologie fonctionnelle et développement
315 durable des écosystèmes naturels et anthropisés”), the University Pierre-et-Marie Curie, the
316 CNRS (UMR 5178 and UMR 6250 LIENSs) and a research grant from the Government of
317 New Caledonia awarded by the Pacific Fund.

318

319 **6. REFERENCES**

320

321 Ambatsian, P., Fernex, F., Bernat, M., Parron, C., Lecolle, J. 1997. High metal inputs to
322 closed seas: the New Caledonian lagoon. *J. Geochem. Explor.* 59, 59-74.
323

324 Boucaud-Camou, E., Boucher-Rodoni, R. 1983. Feeding and digestion in Cephalopods. In:
325 *The Mollusca, Part 2, physiology* (Wilbur, K M ed). Academic press, New York, 1983, pp.
326 149-180.
327

328 Boucher-Rodoni, R., Boucaud-Camou, E. 1987. Fine structure and absorption of ferritin in the
329 digestive organs of *Loligo vulgaris* and *L. forbesi* (Cephalopoda, Teuthoidea). *J. Morph.* 193,
330 173-184.
331

332 Boyle, P., Rodhouse, P. 2005. *Cephalopods: Ecology and Fisheries* Blackwell publishing,
333 Ames. 452 pp.
334

335 Buckeau, G., Kim, J.I., Klenze, R., Rhee, D.S., Wimmer, H. 1992. A comparative study of the
336 fulvate complexation of trivalent transuranic ions. *Radiochim. Acta* 57, 105-112.
337

338 Bustamante, P., Cherel, Y., Caurant, F., Miramand, P. 1998. Cadmium, copper and zinc in
339 octopuses from Kerguelen Islands, southern Indian Ocean. *Polar Biol.* 19, 264-271.
340

341 Bustamante, P., Grigioni, S., Boucher-Rodoni, R., Caurant, F., Miramand, P. 2000.
342 Bioaccumulation of 12 trace elements in the tissues of the nautilus *Nautilus macromphalus*
343 from New Caledonia. *Mar. Pollut. Bull.* 40, 688-696.
344

345 Bustamante, P., Cosson, R.P., Gallien, I., Caurant, F., Miramand, P. 2002a. Cadmium
346 detoxification processes in the digestive gland of cephalopods in relation to accumulated
347 cadmium concentrations. *Mar. Environ. Res.* 53, 227-241.

348

349 Bustamante, P., Teyssié, J-L., Fowler, S.W., Cotret, O., Danis, B., Miramand, P., Warnau, M.
350 2002b. Biokinetics of zinc and cadmium accumulation and depuration at different stages in
351 the life cycle of the cuttlefish *Sepia officinalis*. *Mar. Ecol. Prog. Ser.* 231, 167–177.

352

353 Bustamante, P., Garrigue, C., Breau, L., Caurant, F., Dabin, W., Greaves, J., Dodemont, R.
354 2003. Trace elements in two odontocetes species (*Kogia breviceps* and *Globicephala*
355 *macrorhynchus*) stranded in New Caledonia (South Pacific). *Environ. Pollut.* 124, 263-271.

356

357 Bustamante, P., Teyssié, J-L., Danis, B., Fowler, S.W., Miramand, P., Cotret, O., Miramand,
358 P., Warnau, M. 2004. Uptake, transfer and distribution of silver and cobalt in tissues of the
359 common cuttlefish *Sepia officinalis* at different stages of its life cycle. *Mar. Ecol. Prog. Ser.*
360 269, 185-195.

361

362 Bustamante, P., Miramand, P. 2005. Evaluation of the variegated scallop *Chlamys varia* as a
363 biomonitor of temporal trends of Cd, Cu, and Zn in the field. *Environ. Pollut.* 138, 109-120.

364

365 Bustamante, P., Teyssié, J-L., Fowler, S.W., Warnau, M. 2006. Assessment of the exposure
366 pathway in the uptake and distribution of americium and cesium in cuttlefish (*Sepia*
367 *officinalis*) at different stages of its life cycle. *J. Exp. Mar. Biol. Ecol.* 331, 198-207.

368

369 Choppin Torres, R.A., Choppin, G.R. 1984. Europium (III) and americium (III) stability
370 constants with humic acid. *Radiochim. Acta* 34, 143-148.

371

372 Dalvi, A.D., Bacon, W.G., Osborne, R.C. 2004. The past and the future of nickel laterites.
373 PDAC International Convention, Trade Show and Investors Exchange. March 7–10, 27 pp.

374

375 Dorneles, P.R., Lailson-Brito, J., Aguiar dos Santos, R., Silva da Costa, P.A., Malm, O.,
376 Freitas Azevedo, A., Machado Torres, J.P. 2007. Cephalopods and cetaceans as indicators of
377 offshore bioavailability of cadmium off Central South Brazil Bight. *Environ. Pollut.* 148, 352–
378 359.

379

380 Eisler, R. 1981. Trace metals concentration in marine organisms. Pergamon Press, New York,
381 685 pp.

382

383 Fernandez, J.M., Ouillon, S., Chevillon, C., Douillet, P., Fichez, R., Legendre, R. 2006. A
384 combined modelling and geochemical study of the fate of terrigenous inputs from mixed
385 natural and mining sources in a coral reef lagoon (New Caledonia) *Mar. Pollut. Bull.* 52, 320–
386 331.

387

388 Finger, J.M., Smith, J.D. 1987. Molecular association of Cu, Zn, Cd and ²¹⁰Po in the digestive
389 gland of the squid *Nototodarus gouldi*. *Mar. Biol.* 95, 87-91.

390

391 Ghiretti-Magaldi, A., Giuditta, A., Ghiretti, F. 1958. Pathways of terminal respiration in
392 marine invertebrates. I. The respiratory system in cephalopods. *J. Cell. Comp. Physiol.* 52,
393 389-429.

394

395 Guary, J-C., Higgo, J.J.W., Cherry, R.D., Heyraud, M. 1981. High concentrations of
396 transuranic and natural radioactive elements in the branchial hearts of the cephalopods
397 *Octopus vulgaris*. Mar. Ecol. Prog. Ser. 4, 123-126.
398

399 Guary, J-C., Fowler, S.W. 1982. Experimental studies on the biokinetics of plutonium and
400 americium in the cephalopod *Octopus vulgaris*. Mar. Ecol. Prog. Ser. 7, 327-335.
401

402 Harrison, N., Gangaiya, P., Morrison, R.J. 1996. Organochlorines in the Coastal Marine
403 Environment of Vanuatu and Tonga. Mar. Pollut. Bull. 32, 575-579.
404

405 Hédouin, L., Metian, M., Teyssié, J-L., Fowler, S.W., Fichez, R., Warnau, M. 2006.
406 Allometric relationships in the bioconcentration of heavy metals by the edible tropical clam
407 *Gafrarium tumidum*. Sci. Total Environ. 366, 154–163.
408

409 Hédouin, L., Pringault, O., Metian, M., Bustamante, P., Warnau, M. 2007. Nickel
410 bioaccumulation in bivalves from the New Caledonia lagoon: Seawater and food exposure.
411 Chemosphere 66, 1449-1457.
412

413 Hirano, S., Suzuki, K.T. 1996. Exposure, metabolism, and toxicity of rare earth and related
414 compounds. Environ. Health Perspect. 104, 85-95.
415

416 Kim, J.I., Buckeau, G., Bryant, E., Klenze, R. 1989. Complexation of americium (III) with
417 humic acid. Radiochim. Acta 48, 135-143.
418

419 Koyama, J., Nanamori, N., Segawa, S. 2000. Bioaccumulation of waterborne and dietary
420 cadmium by oval squid *Sepioteuthis lessoniana*, and its distribution among organs. Mar.
421 Pollut. Bull. 40, 961-996.
422

423 Ichihashi, H., Kohno, H., Kannan, K., Tsumura, A., Yamasaki, S. 2001. Multielemental
424 analysis of purpleback flying squid using high resolution inductively coupled plasma-mass
425 spectrometry (HR ICP-MS). Environ. Sci. Technol. 35, 3103-3108.
426

427 Labrosse, P., Letourneur, Y., Kulbicki, M., Paddon, J.R. 2000. Fish stock assessment of the
428 Northern New Caledonian lagoons: fishing pressure, potential yields and impact on
429 management options. Aquat. Living Res. 13, 91-98.
430

431 Norman, M.D. 2000. *Cephalopods: a world guide*. IKAN publishing, Frankfurt, 320 pp.
432

433 Malinovsky, A.I., Markevich, P.V. 2007. Heavy Clastic Minerals from the Far East Island-
434 Arc Complexes. Russian J. Pacific Geol. 1, 71-81.
435

436 Martin, J.H., Flegal, A.R. 1975. High copper concentrations in squid livers in association with
437 elevated levels of silver, cadmium, and zinc. Mar. Biol. 30, 51-55.
438

439 Metian, M., Bustamante, P., Hédouin, L., Warnau, M. 2008. Accumulation of trace elements
440 in the tropical scallop *Comptopallium radula* from coral reefs in New Caledonia. Environ.
441 Pollut. 152: 543-552.
442

443 Miramand, P., Guary, J-C. 1980. High concentrations of some heavy metals in tissues of the
444 Mediterranean octopus. Bull. Environ. Contam. Toxicol. 24, 783-788.
445

446 Miramand, P., Bentley, D. 1992. Concentration and distribution of heavy metals in tissues of
447 two cephalopods, *Eledone cirrhosa* and *Sepia officinalis*, from the French coast of the English
448 Channel. Mar. Biol. 114, 407-414.
449

450 Monniot, F., Martoja, R., Monniot, C. 1994. Cellular sites of iron and nickel accumulation in
451 ascidians related to the naturally and anthropic enriched New Caledonian environment.
452 Annal. Inst. Océanogr. 70, 205-216.
453

454 Monteiro, L.R., Porteiro, F.M., Gonçalves, J.M. 1992. Inter- and intra-specific variation of
455 mercury levels in muscle of cephalopods from the Azores. Arquipelago 10, 13-22.
456

457 O'Dor, R.K., Forsythe, J., Webber, D.M., Wells, J., Wells, M.J. 1993. Activity levels of
458 *Nautilus* in the wild. Nature 362, 626-627.
459

460 Pernice, M., Wetzel, S., Gros, O., Boucher-Rodoni, R., Dubilier, N. 2007. Enigmatic dual
461 symbiosis in the excretory organs of *Nautilus macromphalus* (Cephalopoda: Nautiloidea).
462 Proc. Royal Soc. Lond. Ser. Biol. Sci. 274, 1143–1152.
463

464 Pernice, M., Boucher, J., Joannot, P., Boucher-Rodoni, R. 2006. Comparative study of
465 bacteria diversity and bioaccumulation of trace elements in the excretory organs of *Nautilus*.
466 In: CIAC symposium 2006, February 6-10, Hobart, Australia, 120 pp.
467

468 Pierce, G.J., Stowasser, G., Hastie, L.C., Bustamante, P. 2008. Geographic, seasonal and
469 ontogenetic variation in cadmium and mercury concentrations in squid (Cephalopoda:

470 Teuthoidea) from UK waters. *Ecotox. Environ. Safety* in press
471 ([doi:10.1016/j.ecoenv.2007.07.007](https://doi.org/10.1016/j.ecoenv.2007.07.007)).

472

473 Rainbow, P.S., Wang, W-X. 2001. Comparative assimilation of Cr, Cr, Se, and Zn by the
474 barnacle *Elminius modestus* from phytoplankton and zooplankton diets. *Mar. Ecol. Prog. Ser.*
475 218, 239–248.

476

477 Rainbow, P.S. 2002. Trace metal concentrations in aquatic invertebrates: why and so what?
478 *Environ. Pollut.* 120, 497–507.

479

480 Raimundo, J., Caetano, M., Vale, C. 2004. Geographical variation and partition of metals in
481 tissues of *Octopus vulgaris* along the Portuguese coast. *Sci. Total Environ.* 325, 71-81.

482

483 Saunders, W., Ward, P.D. 1987. Ecology, distribution and population characteristics of
484 Nautilus. In: *Nautilus, the biology and paleobiology of a living fossil* Saunders, WB &
485 Landman, NH. (eds). New-York & London, Plenum press, 137-162 pp.

486

487 Schipp, R., Martin, A.W., Liebermann, H., Magnier, Y. 1985. Cytomorphology and function
488 of pericardial appendages of Nautilus (Cephalopoda, Tetrabranchiata). *Zoomorph.* 105, 16-29.

489

490 Schipp, R., Martin, A.W. 1987. The excretory system of Nautilus. In: *Nautilus, the biology*
491 *and Paleobiology of a living fossil* (WB Sauders and NH. Landman, eds). Plenum Press, New
492 York, 281-304 pp.

493

494 Seixas, S., Bustamante, P., Pierce, G.J. 2005a. Interannual patterns of variation in
495 concentrations of trace elements in arms of *Octopus vulgaris*. *Chemosphere* 59, 1113-1124.
496

497 Seixas, S., Bustamante, P., Pierce, G.J. 2005b. Accumulation of mercury in the tissues of the
498 common octopus *Octopus vulgaris* (L.) in two localities on the Portuguese coast. *Sci. Total*
499 *Environ.* 340, 113-122.
500

501 Smith, J.D., Plues, L., Heyraud, M., Cherry, R.D. 1984. Concentrations of the elements Ag,
502 Al, Ca, Cd, Cu, Fe, Mg,, Pb and Zn, and the radionuclides ²¹⁰Pb and ²¹⁰Po in the digestive
503 gland of the squid *Nototodarus gouldi*. *Mar. Environ. Res.* 13, 55-68.
504

505 Wang, W-X. 2002. Cd and Se aqueous uptake and exposure of green mussels *Perna viridis*:
506 influences of seston quantity. *Mar. Ecol. Prog. Ser.* 226, 211–221.
507

508 Ward, P.D. 1988. In search of Nautilus “A New York Academy of Sciences book”, Simon &
509 Schuster eds, New-York.
510

511 Warnau, M., Ledent, G., Temara, A. 1995. Allometry of heavy metal bioconcentration in the
512 echinoid *Paracentrotus lividus*. *Arch. Environ. Contam. Toxicol.* 29, 393–399.
513

514 Westermann, B., Ruth, P., Litzbauer, H.D., Beck, I., Beuerlin, K., Schimdtberg, H., Kleta,
515 E.F., Schipp, R. 2002. The digestive tract of *Nautilus pompilius* (Cephalopoda,
516 Tetrabranchiata): an X-ray analytical and computational tomography study on the living
517 animal. *J. Exp. Biol.* 205, 1617-1624.
518

519 Worms, I., Simona, D.F., Hasslera, C.S., Wilkinson, K.J. 2006. Bioavailability of trace metals
520 to aquatic microorganisms: importance of chemical, biological and physical processes on
521 biouptake. *Biochimie* 88, 1721–1731.

522

523 Yamada, H., Takayanagi, K., Tateishi, M., Tagata, H., Ikeda, K. 1997. Organotin compounds
524 and polychlorinated biphenyls of livers in squid collected from coastal waters and open ocean.
525 *Environ. Pollut.* 96(2), 217-226.

526 Table 1. Characteristics of the sampled specimens of *Nautilus pompilius* and *N.*
527 *macromphalus*.

528

Species	Sex	Length (mm)	Weight (g)
<i>N. pompilius</i>			
1	Male	177	825
2	Male	151	660
3	Male	171	796
<i>N. macromphalus</i>			
1	Male	164	766
2	Female	162	676
3	Male	173	870
4	Male	160	758
5	Male	140	480

529 Table 2: Trace elements concentrations (Mean \pm SE; $\mu\text{g g}^{-1}$ dry wt) in the tissues of Nautiloidae from this study and from the literature. The
 530 highest values are in bold. dl: detection limit; NA: not analysed; # only one value.

531

Tissue	Rare earth elements			Non essential metals			Essential metals and metalloids									
	La	Ce	Nd	Ag	Cd	Pb	As	Co	Cr	Cu	Fe	Mn	Ni	Se	V	Zn
<i>Nautilus macromphalus</i> (Bustamante et al. 2000)																
Digestive gland	NA	NA	NA	4.45 \pm 2.68	45.1\pm13.2	0.46 \pm 0.17	166 \pm 26	7.8\pm3.6	4.4 \pm 1.1	106\pm46	666\pm231	10.1 \pm 1.7	16.3 \pm 7.8	NA	8.8\pm2.0	672\pm208
Pericardial appendages	NA	NA	NA	20.3\pm6.2	2.16 \pm 1.63	1.00 \pm 0.82	260\pm111	1.31 \pm 0.80	8.25\pm1.55	27.2 \pm 33.4	78 \pm 46	25.1 \pm 23.0	5.64 \pm 3.54	NA	0.18 \pm 0.09	22.4 \pm 26.6
Renal appendages	NA	NA	NA	1.80 \pm 2.22	1.87 \pm 1.23	1.06\pm0.88	112 \pm 65	0.43 \pm 0.37	2.54 \pm 2.88	7.1 \pm 10.2	343 \pm 160	77.1\pm54.1	22.7\pm13.9	NA	1.02 \pm 0.72	11.0 \pm 16.8
<i>N. macromphalus</i> (Present study)																
Digestive gland	0.30\pm0.07	0.25\pm0.11	0.19\pm0.07	9.42\pm3.91	51.5\pm3.5	0.98 \pm 0.15	257\pm19	4.69\pm1.51	2.95 \pm 0.35	263\pm78	353\pm63	10.9 \pm 1.3	14.0 \pm 1.04	2.65 \pm 0.90	14.8\pm2.28	388\pm61
Pericardial appendages	0.11 \pm 0.02	0.17 \pm 0.05	0.15 \pm 0.05	1.82 \pm 0.54	1.93 \pm 0.76	1.57 \pm 0.11	169 \pm 19	0.65 \pm 0.09	9.59\pm3.35	73.7 \pm 9.9	19 \pm 1	12.7 \pm 2.9	6.93 \pm 1.37	3.72 #	5.62 \pm 2.26	56.5 \pm 1.8
Renal appendages	<dl	<dl	<dl	0.44 \pm 0.12	3.39 \pm 1.16	2.11\pm0.51	152 \pm 17	0.18 \pm 0.03	2.57 \pm 0.36	27.6 \pm 4.2	20 \pm 3	68.2\pm6.2	19.7\pm2.39	9.84\pm0.94	1.54 \pm 0.13	35.2 \pm 2.8
<i>N. pompilius</i> (Present study)																
Digestive gland	1.00\pm0.14	1.58\pm0.32	0.92\pm0.10	5.65\pm1.04	74.3\pm6.8	0.62 \pm 0.07	210 \pm 16	3.85\pm1.04	1.37 \pm 0.24	311\pm61	625\pm51	15.4 \pm 1.7	9.77 \pm 0.07	2.35 \pm 0.84	8.42\pm0.57	470\pm52
Pericardial appendages	0.24 \pm 0.03	0.42 \pm 0.05	0.31 \pm 0.01	1.75 \pm 0.07	2.39 \pm 0.37	1.24\pm0.24	323\pm53	0.79 \pm 0.07	9.30\pm3.87	99.7 \pm 2.4	21 \pm 3	24.1 \pm 4.9	7.02 \pm 0.79	2.49 #	4.94 \pm 0.74	58.2 \pm 2.7
Renal appendages	<dl	<dl	<dl	0.35 \pm 0.03	5.93 \pm 0.31	0.99 \pm 0.21	292 \pm 80	0.18 \pm 0.03	1.98 \pm 0.08	53.3 \pm 4.6	24 \pm 3	307\pm83	19.6\pm0.97	13.3\pm6.12	1.82 \pm 0.02	42.7 \pm 1.5

532 Table 3. One-way ANOVA comparison of trace element concentrations in *Nautilus pompilius*

533 (Np) and in *N. macromphalus* (Nm). NS: not significant.

534

Element	Digestive gland	Pericardial appendages	Renal appendages
Rare earth elements			
Ce	Np > Nm (F= 19.58 ; p= 0.007)	NS	-
La	Np > Nm (F= 23.40; p= 0.005)	Np > Nm (F= 0.609; p= 0.025)	-
Nd	Np > Nm (F= 25.13 ; p= 0.004)	Np > Nm (F= 6.54 ; p= 0.051)	-
Essential elements			
As	NS	Np > Nm (F= 9.58 ; p= 0.027)	NS
Co	NS	NS	NS
Cr	Nm > Np (F= 11.34 ; p= 0.020)	NS	NS
Cu	NS	NS	Np > Nm (F= 15.58; p=0.008)
Fe	Np > Nm (F= 10.09 ; p= 0.025)	NS	NS
Mn	NS	NS	NS
Ni	NS	NS	NS
Se	NS	NS	NS
V	NS	NS	NS
Zn	NS	NS	NS
Non essential elements			
Ag	NS	NS	NS
Cd	Nm > Np (F= 0.740 ; p= 0.023)	NS	NS
Pb	NS	NS	NS

535 Table 4: Correlations between trace elements within the different tissues of nautilus.

536 Not underlined: p<0.05; underlined: p<0.001.

537

Elements	Digestive gland	Renal appendages	Pericardial appendages
La	<u>+ Ce, + Nd, - As, + Fe, + Mn</u>	<dl	<u>+ Ce, + Nd, + Mn</u>
Ce	<u>+ Nd, - As, + Fe, + Mn,</u>	<dl	<u>+ Nd, + Co, + La</u>
Nd	<u>+ Fe, + Mn</u>	<dl	<u>+ Ce, + Co, + La</u>
Cu	+ Zn, -Se	-	-
Fe	+ Mn, -V, + <u>La, + Ce, + Nd</u>	+ Zn	+ Ag, <u>+ Mn</u> , + Cr
Pb	+ V	- V, <u>+ Cr</u>	-
Mn	+ La, + Ce, + Nd, + Fe	+ Cd, +Zn	+ La, <u>+ Fe</u>
Cd	-	<u>+Mn</u>	<u>+ Ni, + V, + Cr</u>

538

539 <dl: concentrations below the detection limit.

540 **Caption to figures**

541

542 Figure 1: Location of *Nautilus pompilius* and *N. macromphalus* specimens' collection.

543

544 Figure 2: Relative distribution of the trace elements between the digestive gland (DG),

545 pericardial appendages (PA) and renal appendages (RA) of *Nautilus pompilius* from Vanuatu


546 (black) and *N. macromphalus* from New Caledonia (white). Scale bars represent 1 standard

547 error.

548

549

550



551

552

553

Figure 1.

554

