



HAL
open science

The non-symmetric operad pre-Lie is free

Nantel Bergeron, Muriel Livernet

► **To cite this version:**

| Nantel Bergeron, Muriel Livernet. The non-symmetric operad pre-Lie is free. 2008. hal-00337068v1

HAL Id: hal-00337068

<https://hal.science/hal-00337068v1>

Preprint submitted on 5 Nov 2008 (v1), last revised 15 Jul 2009 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE NON-SYMMETRIC OPERAD PRE-LIE IS FREE

NANTEL BERGERON AND MURIEL LIVERNET

ABSTRACT. We prove that the pre-Lie operad is a free non-symmetric operad.

CONTENTS

Introduction	1
1. The pre-Lie operad and rooted trees	2
2. A gradation on labelled rooted trees	5
3. The operad pre-Lie is free as a non-symmetric operad	7
References	10

INTRODUCTION

Operads are a specific tool for encoding type of algebras. For instance there are operads encoding associative algebras, commutative and associative algebras, Lie algebras, pre-Lie algebras, dendriform algebras, Poisson algebras and so on. A usual way of describing a type of algebras is by giving the generators and relations. For instance a Lie algebra L is a vector space together with a bilinear product, the bracket (the generator) satisfying the relations $[x, y] = -[y, x]$ and $[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0$ for all $x, y, z \in L$. The vector space of all operations one can perform on n distinct variables in a Lie algebra is $\mathcal{L}ie(n)$, the building block of the symmetric operad $\mathcal{L}ie$. Composition in the operad corresponds to composition of operations. The vector space $\mathcal{L}ie(n)$ has a natural action of the symmetric group, so it is a symmetric operad. The case of associative algebras can be considered in two different ways. An associative algebra A is a vector space together with a product satisfying the relation $(xy)z = x(yz)$. The vector space of all operations one can perform on n distinct variables in an associative algebra is $\mathcal{A}s(n)$, the building block of the symmetric operad $\mathcal{A}s$. The vector space $\mathcal{A}s(n)$ has for basis the symmetric group S_n . But, in view of the relation, one can look also at the vector space of all ordered-preserving operations one can perform on n distinct ordered variables in an associative algebra: this is a vector space of dimension 1 generated by the only operation $x_1 \cdots x_n$. So the non-symmetric operad $\widetilde{\mathcal{A}s}$ describing associative algebras is 1-dimensional for each n : this is the terminal object in the category of non-symmetric operads.

2000 *Mathematics Subject Classification.* 18D, 05E, 17B.

Key words and phrases. rooted tree, pre-Lie algebras.

Livernet supported by the Clay Mathematical Institute and hosted by MIT.

Here is the picture between symmetric and non-symmetric operads. A symmetric operad \mathcal{P} starts with a graded vector space $(\mathcal{P}(n))_{n \geq 0}$ together with an action of the symmetric group S_n on $\mathcal{P}(n)$ for each n . This data is called a symmetric sequence or an \mathbb{S} -module or a vector species. There is a forgetful functor from vector species to graded vector spaces, forgetting the action of the symmetric group. This functor has a left adjoint \mathcal{S} which corresponds to tensoring by the regular representation of the symmetric group. A symmetric operad is an operad in the category of vector species and a non-symmetric operad is an operad in the category of graded vector spaces. The symmetric operad $\mathcal{A}s$ is the image by \mathcal{S} of \star . It is clear that $\mathcal{L}ie$ is not in the image of \mathcal{S} since the Jacobi relation does not respect the order of the variables $x < y < z$ nor the anti-symmetry relation. Still one can regard $\mathcal{L}ie$ as a non-symmetric operad applying the forgetful functor. Salvatore and Tauraso proved in [6] that the operad $\mathcal{L}ie$ is a free non-symmetric operad.

A free non-symmetric operad describes type of algebras which have a set of generating operations and no relations between them. For instance, magmatic algebras are vector spaces together with a bilinear product. There is a well known free non-symmetric operad, built on Stasheff polytopes, see e.g. [7]. Algebras over this operad are vector spaces V together with an n -linear product: $V^{\otimes n} \rightarrow V$ for each n . In a homotopy theory point of view, the category of operads is a Quillen category and free operads play an essential role in the homotopy category. One wants to replace an operad \mathcal{P} by a quasi-free resolution, that is, a free operad together with a differential such that its homology gives \mathcal{P} . For instance, a quasi-free resolution of $\widetilde{\mathcal{A}s}$, the non-symmetric operad for associative algebras is given by the Stasheff polytopes and algebras over it are A_∞ -algebras (associative algebras up to homotopy). This gives us the motivation for studying whether a given symmetric operad is free as a non-symmetric operad or not.

In this paper we prove that the operad pre-Lie is a free non-symmetric operad. Pre-Lie algebras are vector spaces together with a bilinear product satisfying the relation $(x * y) * z - x * (y * z) = (x * z) * y - x * (z * y)$. The operad pre-Lie is based on labelled rooted trees which are of combinatorial interest. In the process of proving the main result, we describe another operad denoted \mathcal{T}_{Max} also based on rooted trees and having the advantage to be a set operad. We prove that it is a free non-symmetric operad. The link between the two operads is made via a gradation on labelled rooted trees.

1. THE PRE-LIE OPERAD AND ROOTED TREES

We first recall the definition of the pre-Lie operad based on labelled rooted trees as in [3]. For $n \in \mathbb{N}^*$, the set $\{1, \dots, n\}$ is denoted by $[n]$ and $[0]$ denotes the empty set. The symmetric group on k letters is denoted by S_k . There are many equivalent definitions of operads and we refer to [5] for basics on operads. Here are the definitions needed for the sequel.

Definition 1.1. A *non-symmetric operad* is a graded vector space $(\mathcal{P}(n))_{n \geq 1}$ together with compositions $\circ_i : \mathcal{P}(n) \otimes \mathcal{P}(m) \rightarrow \mathcal{P}(n + m - 1)$ for $1 \leq i \leq n$ satisfying the

following relations: for $a \in \mathcal{P}(n)$, $b \in \mathcal{P}(m)$ and $c \in \mathcal{P}(\ell)$

$$\begin{aligned} (a \circ_i b) \circ_{j+i-1} c &= a \circ_i (b \circ_j c), & \text{for } 1 \leq j \leq m, \\ (a \circ_i b) \circ_j c &= (a \circ_j c) \circ_{i+\ell-1} b, & \text{for } j < i. \end{aligned}$$

If in addition each $\mathcal{P}(n)$ is acted on the right by the symmetric group S_n and the compositions are equivariant with respect to this action, then the collection $(\mathcal{P}(n))_n$ forms a *symmetric operad*. An *algebra* over an operad \mathcal{P} is a vector space X endowed with evaluation maps

$$\begin{aligned} ev_n : \quad \mathcal{P}(n) \otimes X^{\otimes n} &\rightarrow X \\ p \otimes x_1 \otimes \dots \otimes x_n &\mapsto p(x_1, \dots, x_n) \end{aligned}$$

compatible with the compositions \circ_i : for $p \in \mathcal{P}(n)$, $q \in \mathcal{P}(m)$, $x'_i \in X$ one has

$$(p \circ_i q)(x_1, \dots, x_{n+m-1}) = p(x_1, \dots, x_{i-1}, q(x_i, \dots, x_{i+m-1}), x_{i+m}, \dots, x_{n+m-1}).$$


If the operad is symmetric the evaluation maps are required to be equivariant with respect to the action of the symmetric group as follows:

$$(p \cdot \sigma)(x_1, \dots, x_n) = p(x_{\sigma^{-1}(1)}, \dots, x_{\sigma^{-1}(n)}).$$


Remark 1.2. *Planar tree of operations:* a convenient way to uniquely represent composition of operations in a non-symmetric operad \mathcal{P} is to use a planar tree. An element $a \in \mathcal{P}(n)$ is represented by a planar tree with a single trunk (outgoing edge), one vertex labelled by a and n leaves (incoming edges):



The n leaves are counted from left to right as $1, 2, \dots, n$. Now if we have $a \in \mathcal{P}(n)$, $b \in \mathcal{P}(m)$ and $1 \leq i \leq n$ we represent the composition $a \circ_i b$ by the planar tree



The resulting tree has $n + m - 1$ leaves (counted from left to right) and represents an element of $\mathcal{P}(n + m - 1)$. The two relations in Definition 1.1 corresponds to the following two trees: for $a \in \mathcal{P}(n)$, $b \in \mathcal{P}(m)$ and $c \in \mathcal{P}(\ell)$ we can have



Each relation is obtained by writing down the two ways of interpreting the tree as a composition of operations. In general a planar tree $\mathbb{T}(a_1, a_2, \dots, a_k)$ with k vertices

labelled by elements $a_i \in \mathcal{P}(n_i)$ where n_i is the number of incoming edges at the i th vertex, corresponds to a unique composition of operations in \mathcal{P} independent of any relations. Any full subtree of $\mathbb{T}(a_1, a_2, \dots, a_k)$ is completely determined by the position of its leaves; they form an interval $[p, q]$ where $1 \leq p \leq q \leq n_1 + n_2 + \dots + n_k - k + 1$.

Definition 1.3. Let S be a set. An S -labelled rooted tree is a non planar rooted tree whose vertices are in bijection with S . If $S = [n]$, then we talk about n -labelled rooted trees and denote by $\mathcal{T}(n)$ the set of those trees. It is acted on by the symmetric group by permuting the labels.

The set $\mathcal{T}(3)$ has for elements:

$$(1.1) \quad \begin{array}{ccccccc} \begin{array}{c} 2 \quad 3 \\ \diagdown \quad / \\ \circ \\ / \quad \backslash \\ 1 \end{array} & \begin{array}{c} 1 \quad 3 \\ \diagdown \quad / \\ \circ \\ / \quad \backslash \\ 2 \end{array} & \begin{array}{c} 1 \quad 2 \\ \diagdown \quad / \\ \circ \\ / \quad \backslash \\ 3 \end{array} & \begin{array}{c} 3 \\ | \\ 2 \\ | \\ 1 \\ \circ \end{array} & \begin{array}{c} 2 \\ | \\ 3 \\ | \\ 1 \\ \circ \end{array} & \begin{array}{c} 1 \\ | \\ 3 \\ | \\ 2 \\ \circ \end{array} & \begin{array}{c} 3 \\ | \\ 1 \\ | \\ 2 \\ \circ \end{array} & \begin{array}{c} 2 \\ | \\ 1 \\ | \\ 3 \\ \circ \end{array} & \begin{array}{c} 1 \\ | \\ 1 \\ | \\ 2 \\ \circ \end{array} & \begin{array}{c} 1 \\ | \\ 2 \\ | \\ 3 \\ \circ \end{array} \end{array}$$

In general $\mathcal{T}(n)$ has n^{n-1} elements (see [1] for more details).

We denote by $k\mathcal{T}(n)$ the k -vector space spanned by $\mathcal{T}(n)$.

Theorem 1.4. [3, theorem 1.9] *The collection $(k\mathcal{T}(n))_{n \geq 1}$ forms a symmetric operad, the operad pre-Lie denoted by \mathcal{PL} . Algebras over this operad are pre-Lie algebras, that is, vector spaces L together with a product $*$ satisfying the relation*

$$(x * y) * z - x * (y * z) = (x * z) * y - x * (z * y), \quad \forall x, y, z \in L.$$

We recall the operad structure of \mathcal{PL} as explained in [3]. A rooted tree is naturally oriented from the leaves to the root, so it makes sense to define the set of incoming vertices of a vertex i denoted by $\text{In}(T, i)$. There is also at most one outgoing vertex of a vertex i depending whether i is the root of T or not. For $T \in \mathcal{T}(n)$ and $S \in \mathcal{T}(m)$, we define

$$T \circ_i S = \sum_{f: \text{In}(T, i) \rightarrow [m]} T \circ_i^f S,$$

where $T \circ_i^f S$ is the rooted tree obtained by substituting the tree S for the vertex i in T . The outgoing vertex of i , if it exists, becomes the outgoing vertex of the root of S , whereas the incoming vertices of i are grafted on the vertices of S according to the map f . The root of $T \circ_i^f S$ is the root of T or the root of S in case i is the root of T . There is also a relabelling of the vertices of T and S in $T \circ_i^f S$: we add $i - 1$ to the labels of S and $m - 1$ to the ones of T which are greater than i . Here is an example:

$$(1.2) \quad \begin{array}{c} \begin{array}{c} 1 \quad 3 \\ \diagdown \quad / \\ \circ \\ / \quad \backslash \\ 2 \end{array} \circ_2 \begin{array}{c} 1 \\ | \\ 2 \\ \circ \end{array} = \begin{array}{c} 1 \quad 4 \\ \diagdown \quad / \\ \circ \\ / \quad \backslash \\ 2 \end{array} \circ_2 \begin{array}{c} 2 \\ | \\ 3 \\ \circ \end{array} = \begin{array}{c} 1 \quad 4 \\ \diagdown \quad / \\ \circ \\ | \\ 2 \\ | \\ 3 \\ \circ \end{array} + \begin{array}{c} 1 \\ | \\ 4 \\ | \\ 2 \\ | \\ 3 \\ \circ \end{array} + \begin{array}{c} 4 \\ | \\ 1 \\ | \\ 2 \\ | \\ 3 \\ \circ \end{array} + \begin{array}{c} 1 \quad 2 \quad 4 \\ \diagdown \quad / \quad / \\ \circ \\ | \\ 3 \\ \circ \end{array} \end{array}$$



2. A GRADATION ON LABELLED ROOTED TREES

We introduce a gradation on labelled rooted trees. We prove that in the expansion of the composition of two rooted trees in the operad pre-Lie there is a unique rooted tree of maximal degree and a unique tree of minimal degree, yielding new non-symmetric operad structures on labelled rooted trees.

Definition 2.1. Let T be an n -labelled rooted tree. Let $\{a, b\}$ denote a pair of two adjacent vertices labelled by a and b . The degree of $\{a, b\}$ is $|a - b|$. The degree of T denoted by $\deg(T)$ is the sum of the degrees of its pairs of adjacent vertices. For instance

$$\deg(\begin{array}{c} 1 \quad 3 \\ \diagdown \quad / \\ \circ \\ / \quad \backslash \\ 2 \end{array}) = 2, \quad \deg(\begin{array}{c} 1 \quad 4 \\ \diagdown \quad / \\ \circ \\ | \\ \circ \\ | \\ 3 \end{array}) = 4, \quad \deg(\begin{array}{c} 4 \\ | \\ \circ \\ / \quad \backslash \\ 2 \quad 1 \\ | \\ \circ \\ | \\ 3 \end{array}) = 5, \quad \deg(\begin{array}{c} 1 \\ | \\ \circ \\ / \quad \backslash \\ 2 \quad 4 \\ | \\ \circ \\ | \\ 3 \end{array}) = 3$$

Proposition 2.2. *In the expansion of $T \circ_i S$ in the operad pre-Lie, there is a unique tree of minimal degree and a unique tree of maximal degree.*

For instance, in the equation (1.2) the rooted tree of minimal degree 3 is  and the one of maximal degree 5 is . The other ones are of degree 4.

Proof– Any tree in the expansion of $T \circ_i S$ writes $U_f := T \circ_i^f S$ for some $f : \text{In}(T, i) \rightarrow [m]$. To compute the degree of U_f , we compute the degree of a pair of two adjacent vertices $\{a, b\}$ in U_f . There are 4 cases to consider: a) the pair was previously in S or b) it was previously in T and each vertex was different from i , or c) it was in T of the form $\{i, j\}$ for $j \in \text{In}(T, i)$ or d) if i is not the root of T it was of the form $\{i, k\}$ where k is the outgoing vertex of i .

In case a) the degree of the pair in U_f is the same as it was in S .

In case b), let $\{a', b'\}$ be the corresponding pair in T before relabelling. The degree d of the pair $\{a, b\}$ in U_f is the same as the degree d' of $\{a', b'\}$ except if $a' < i < b'$ or $b' < i < a'$, where $d = d' + m - 1$. Let $\text{gap}(T, i)$ be the number of adjacent pairs of vertices in T satisfying the latter condition.

In case c), let $\{i, j\}$ be the pair in T which gives the pair $\{a, b\}$ in U_f . Let d' be the degree of $\{i, j\}$. If $j < i$ then $\{a, b\} = \{f(j) + i - 1, j\}$. Its degree d is minimal and equals d' if $f(j) = 1$. It is maximal and equals $d' + m - 1$ if $f(j) = m$. If $j > i$ then $\{a, b\} = \{f(j) + i - 1, j + m - 1\}$. Its degree d is minimal and equals d' if $f(j) = m$. It is maximal and equals $d' + m - 1$ if $f(j) = 1$.

In case d), let d' be the degree of $\{i, k\}$. If $k < i$ then $\{a, b\} = \{s + i - 1, k\}$ where s is the label of the root of S . It has degree $d' + s - 1$. If $k > i$, then $\{a, b\} = \{s + i - 1, k + m - 1\}$ and has degree $(m - s) + d'$. Let $\epsilon(T, i, s)$ be $0, s - 1, m - s$ according to the different situations, 0 corresponding to the one where i is the root of T .

As a conclusion

$$\begin{aligned} \deg(T) + \deg(S) + \text{gap}(T, i)(m-1) + \epsilon(T, i, s) &\leq \deg(U_f) \leq \\ \deg(T) + \deg(S) + \text{gap}(T, i)(m-1) + \epsilon(T, i, s) + |\text{In}(T, i)|(m-1). \end{aligned}$$

There is a unique f_{Min} such that $\deg(U_{f_{\text{Min}}})$ is minimal and there is a unique f_{Max} such that $\deg(U_{f_{\text{Max}}})$ is maximal:

$$(2.1) \quad f_{\text{Min}}(k) = \begin{cases} 1 & \text{if } k < i, \\ m & \text{if } k > i, \end{cases}$$

$$(2.2) \quad f_{\text{Max}}(k) = \begin{cases} m & \text{if } k < i, \\ 1 & \text{if } k > i, \end{cases}$$

which ends the proof. \square

Theorem 2.3. *There are two different non-symmetric operad structures on the collection $(k\mathcal{T}(n))_{n \geq 1}$ given by the compositions $T \circ_i^{f_{\text{Min}}} S$ on the one hand and $T \circ_i^{f_{\text{Max}}} S$ on the other hand where f_{Min} and f_{Max} were defined in equations (2.1) and (2.2).*

Proof– For a rooted tree T , let E_T be the set of the oriented edges of the tree: an edge is denoted (a, b) where a, b are the labels of its extremal vertices, and the orientation of the tree is from the leaves to the root. For an integer $a \neq i$ we denote by \tilde{a}_i^m the integer a if $a < i$ or $a + m - 1$ if $a > i$. Given a map $f : \text{In}(T, i) \rightarrow [m]$, the set $E_{T \circ_i^f S}$ has different type of elements:

- $(a + i - 1, b + i - 1)$ for $(a, b) \in E_S$;
- $(\tilde{a}_i^m, \tilde{b}_i^m)$ for $(a, b) \in E_T$ and $a, b \neq i$;
- $(\tilde{a}_i^m, f(a) + i - 1)$ for $(a, i) \in E_T$;
- $(i + s - 1, \tilde{b}_i^m)$ for $(i, b) \in E_T$.

Let $T \in \mathcal{T}(n), S \in \mathcal{T}(m)$ and $U \in \mathcal{T}(p)$. In order to avoid confusion, we denote by $f_{\text{Max}}^{i,p}$ the map sending $k < i$ to p and $l > i$ to 1. We would like to compare the trees

$$V_1 = (T \circ_i^{f_{\text{Max}}^{i,m}} S) \circ_{j+i-1}^{f_{\text{Max}}^{j+i-1,p}} U \quad \text{and} \quad V_2 = T \circ_i^{f_{\text{Max}}^{i,m+p-1}} (S \circ_j^{f_{\text{Max}}^{j,p}} U) :$$

- In V_1 and V_2 , any $(a, b) \in E_U$ converts to $(a + j + i - 2, b + j + i - 2)$.
- In V_1 and V_2 , any $(a, b) \in E_S$ converts to $(\tilde{a}_j^p + i - 1, \tilde{b}_j^p + i - 1)$ if $a, b \neq j$, or converts to $(\tilde{a}_j^p + i - 1, f_{\text{Max}}^{j,p}(a) + i + j - 2)$ if $b = j$ or converts to $(j + i - 1 + u - 1, \tilde{b}_j^p + i - 1)$ if $a = j$.
- In V_1 and V_2 , any $(a, b) \in E_T$ with $a, b \neq i$ converts to $(\tilde{a}_i^{p+m-1}, \tilde{b}_i^{p+m-1})$.
- In V_1 and V_2 , any $(a, i) \in E_T$ converts to $(\tilde{a}_i^{p+m-1}, f_{\text{Max}}^{i,m+p-1}(a) + i - 1)$.

- In V_1 and V_2 , any $(i, b) \in E_T$ converts to $(i - 1 + \text{root}(S \circ_j U), \tilde{b}_i^{m+p-1})$, where $\text{root}(S \circ_j U)$ is the root of $S \circ_j U$. More precisely

$$\text{root}(S \circ_j U) = \begin{cases} s & \text{if } s < j \\ u + j - 1 & \text{if } s = j \\ s + p - 1 & \text{if } s > j. \end{cases}$$

The proof of

$$(T \circ_i^{f_{\text{Max}}^{i,m}} S) \circ_j^{f_{\text{Max}}^{j,p}} U = (T \circ_j^{f_{\text{Max}}^{j,p}} U) \circ_{i+p-1}^{f_{\text{Max}}^{i+p-1,m}} S, \text{ for } j < i$$

is similar and left to the reader. So is the proof with f_{Min} instead of f_{Max} . \square

The two operads on labelled rooted trees defined by the theorem are denoted by \mathcal{T}_{Max} and \mathcal{T}_{Min} . Note that they are set operads, that is, the composition is defined at the level of the sets $\mathcal{T}(n)$ and not only at the level of the vector spaces $k\mathcal{T}(n)$. There is another set operad built on rooted trees: the operad NAP encoding non-associative permutative algebras in [4], in which f_{NAP} is the constant map with value the root of S . This operad has the advantage to be a symmetric operad.

3. THE OPERAD PRE-LIE IS FREE AS A NON-SYMMETRIC OPERAD

We show that \mathcal{T}_{Max} is a free non-symmetric operad. Using Proposition 2.2, we conclude that the operad pre-Lie is free as a non-symmetric operad. To this end we need to introduce some notation on rooted trees.

Definition 3.1. Given two ordered sets S and T , an ordered preserving bijection $\phi : S \rightarrow T$ induces a natural bijection between the set of S -labelled rooted trees and the set of T -labelled rooted trees also denoted by ϕ . A T -labelled rooted tree X is *isomorphic* to an S -labelled rooted tree Y if $X = \phi(Y)$.

Given a rooted tree $T \in \mathcal{T}(n)$ and a subset $K \subseteq [n]$, we denote by $T|_K$ the graph obtained from T by keeping only the vertices of T that are labelled by elements of K and only the edges of T that have two vertices labelled in K . Remark that each connected component of $T|_K$ is a rooted tree itself where the root is given by the unique vertex closest to the root of T in the component. Also, for $c \in [n]$ we denote by $T^{(c)}$ the full subtree of T derived from the vertex labelled by c . For example if $K = \{2, 3, 4, 5, 6\} \subset [7]$ and

$$T = \begin{array}{c} \begin{array}{ccc} 2 & & 7 \\ & \diagdown & / \\ & 6 & \\ & / & \diagdown \\ 5 & & 4 \\ & \diagup & \diagdown \\ & 1 & \\ & / & \diagdown \\ & 3 & \end{array} \end{array}, \quad \text{we have } T|_K = \begin{array}{c} \begin{array}{ccc} 2 & & \\ & \diagdown & / \\ & 6 & \\ & / & \diagdown \\ 5 & & 4 \\ & \diagup & \diagdown \\ & 3 & \end{array} \end{array} \quad \text{and } T^{(1)} = \begin{array}{c} \begin{array}{ccc} 2 & & 7 \\ & \diagdown & / \\ & 6 & \\ & / & \diagdown \\ & 1 & \end{array} \end{array}.$$

For $1 \leq a < b \leq n$, $T \in \mathcal{T}_{\text{Max}}(n - b + a)$ and $S \in \mathcal{T}_{\text{Max}}(b - a + 1)$, let $X = T \circ_a S$. Consider the interval $[a, b] = \{a, a + 1, \dots, b\}$, clearly $X|_{[a,b]}$ is isomorphic to S under the unique ordered bijection $[1, b - a + 1] \rightarrow [a, b]$. Let $a \leq c \leq b$ be the label of the root of $X|_{[a,b]}$. Remark that $X^{(c)}$ and $X|_{[a,b]}$ differ only at a and b where there could be subtrees of X that do not belong to $X|_{[a,b]}$. We can then characterize trees X that are obtained from a nontrivial operation $T \circ_a S$ as follows:

Definition 3.2. A tree $X \in \mathcal{T}_{\text{Max}}(n)$ is called *decomposable* if there exists $1 \leq a < b \leq n$ with $(a, b) \neq (1, n)$ such that

- (i) $X|_{[a,b]}$ is a rooted tree. Let c be the label of its root. One has $a \leq c \leq b$.
- (ii) The subtree $X^{(c)}$ contains $X|_{[a,b]}$ and differs from it only at a and b .
- (iii) All subtrees in $X^{(c)} - X|_{[a,b]}$ attached at a have their root labelled in $[b+1, n]$.
- (iv) All subtrees in $X^{(c)} - X|_{[a,b]}$ attached at b have their root labelled in $[1, a-1]$.

It is clear from the discussion above and the definition of the operad \mathcal{T}_{Max} that X is decomposable if and only if it is the result of a non-trivial composition. Consequently, we say that X is *indecomposable* if it is not decomposable. That is there is no $1 \leq a < b \leq n$ such that (i)–(iv) are satisfied. For example let

$$X = \begin{array}{c} \begin{array}{ccc} 7 & & 4 \\ & \diagdown & / \\ 1 & 3 & 2 \\ & | & \\ 8 & 5 & \\ & | & \\ & 6 & \end{array} , & X|_{[3,5]} = \begin{array}{c} 4 \\ / \\ 3 \\ | \\ 5 \end{array} & \text{and} & X^{(5)} = \begin{array}{c} \begin{array}{ccc} 7 & & 4 \\ & \diagdown & / \\ 1 & 3 & 2 \\ & | & \\ 8 & 5 & \\ & | & \\ & 6 & \end{array} \end{array} ,$$

This tree X is decomposable since for $1 \leq 3 < 5 \leq 8$ we have that $X|_{[3,5]}$ is a single tree and the subtree of $X^{(5)} - X|_{[3,5]}$ are attached at 3 and 5 only. Moreover, the subtree attached at 3 has root labelled by $7 \in [6, 8]$ and the subtrees attached at 5 have roots labelled by $1, 2 \in [1, 2]$. Indeed, in \mathcal{T}_{Max} we have

$$X = \begin{array}{c} \begin{array}{ccc} 1 & 5 & 2 \\ & \diagdown & / \\ 6 & 3 & \\ & | & \\ & 4 & \end{array} \circ_3 \begin{array}{c} 2 \\ | \\ 1 \\ | \\ 3 \end{array} .$$

The reader may check that the following are all the indecomposable trees of \mathcal{T}_{Max} up to arity 3:

$$\begin{array}{c} 2 \\ | \\ 1 \end{array} , \quad \begin{array}{c} 1 \\ | \\ 2 \end{array} \quad \text{and} \quad \begin{array}{ccc} 1 & & 3 \\ & \diagdown & / \\ & 2 & \end{array} .$$

Theorem 3.3. *The non-symmetric operad \mathcal{T}_{Max} is a free non-symmetric operad.*

Proof. If \mathcal{T}_{Max} is not free, then for some n there is a tree $X \in \mathcal{T}_{\text{Max}}(n)$ with two distinct constructions from indecomposables. In Remark 1.2, a composition of operations is completely determined by a unique planar rooted tree. We then have that $X = \mathbb{T}(T_1, T_2, \dots, T_r) = \mathbb{Y}(S_1, S_2, \dots, S_k)$ where $T_1, \dots, T_r, S_1, \dots, S_k$ are indecomposables and $\mathbb{T}(T_1, T_2, \dots, T_r)$ and $\mathbb{Y}(S_1, S_2, \dots, S_k)$ are two distinct trees of operations in \mathcal{T}_{Max} with $r, s > 1$.

Since X is decomposable, we can find $1 \leq a < b \leq n$, such that $X|_{[a,b]}$ is isomorphic to a single T_i in position $[a, b]$ in $\mathbb{T}(T_1, T_2, \dots, T_r)$. Moreover $X|_{[a,b]}$ satisfies (i)–(iv) of Definition 3.2.

If $X|_{[a,b]}$ is also isomorphic to a tree S_j in position $[a, b]$ in $\mathbb{Y}(S_1, S_2, \dots, S_k)$, then we replace X by the smaller tree in $\mathcal{T}_{\text{Max}}(n - b + a)$ that we obtain by removing T_i in $\mathbb{T}(T_1, T_2, \dots, T_r)$ and removing S_j in $\mathbb{Y}(S_1, S_2, \dots, S_k)$. Clearly, this new smaller X has two distinct constructions from indecomposables. We can thus assume that $X|_{[a,b]}$ is not isomorphic to a single S_j in position $[a, b]$ in $\mathbb{Y}(S_1, S_2, \dots, S_k)$.

We now study how $X|_{[a,b]}$ overlap in the position $[a, b]$ of $\mathbb{Y}(S_1, S_2, \dots, S_k)$. Remark first that since all S_j are indecomposables, the interval $[a, b]$ cannot be part of a single S_j of $\mathbb{Y}(S_1, S_2, \dots, S_k)$. Indeed, that would imply that S_j would contain a subtree satisfying Definition 3.2 which would be a contradiction.

We may assume that $a > 1$. To see this, assume that the only sub-interval $[a, b] \subset [1, n]$ such that $X|_{[a,b]}$ is isomorphic to a single T_i in position $[a, b]$ in $\mathbb{T}(T_1, T_2, \dots, T_r)$ is such that $a = 1$. Assume moreover that the only sub-interval $[a', b'] \subset [1, n]$ such that $X|_{[a',b']}$ is isomorphic to a single S_j in position $[a', b']$ in $\mathbb{Y}(S_1, S_2, \dots, S_k)$ is such that $a' = 1$. Since S_j is indecomposable, we must have $b > b'$. Similarly, since T_i is indecomposable, we must have $b < b'$. This implies that $b = b'$ and $T_i = S_j$. This possibility was excluded above. So we must have $a > 1$ or $a' > 1$. In the case where $a = 1$ and $a' > 1$ we could just interchange the role of $\mathbb{T}(T_1, T_2, \dots, T_r)$ and $\mathbb{Y}(S_1, S_2, \dots, S_k)$ and assume that we have $a > 1$.

Now, since T_j is indecomposable, there is no subinterval $[c, d] \subseteq [a, b]$ such that $X|_{[c,d]}$ is isomorphic to a full subtree of operations $\mathbb{Y}'(S_{j_1}, S_{j_2}, \dots, S_{j_\ell})$. Hence we can find $c < a \leq d < b$ such that $X|_{[c,d]} \cong \mathbb{Y}'(S_{j_1}, S_{j_2}, \dots, S_{j_\ell})$ satisfies the Definition 3.2.

The graph $X|_{[a,d]}$ belongs to both $X|_{[a,b]}$ and $X|_{[c,d]}$. Let e be the label of the root of $X|_{[a,b]}$ and f the root of $X|_{[c,d]}$. The two full subtrees $X^{(e)}$ and $X^{(f)}$ both contain $X|_{[a,d]}$. This implies that either $X^{(f)}$ is fully contained in $X^{(e)}$, or $X^{(e)}$ is fully contained in $X^{(f)}$.

Let us assume that $X^{(f)}$ is fully contained in $X^{(e)}$, that means $X|_{[a,b]}$ and $X|_{[c,d]}$ are both subtrees of $X^{(e)}$. From Definition 3.2, we know that the difference between $X^{(e)}$ and $X|_{[a,b]}$ is only some subtrees at a and b . The subtree $X|_{[c,d]}$ contains a but does not contain b . Moreover, the point c is not in $X|_{[a,b]}$, so it must be in a subtree of $X^{(e)}$ attached to a . The unique shortest path between a and c in $X|_{[c,d]}$ belongs to that subtree of $X^{(e)}$. Since it is attached at a and is not in $X|_{[a,b]}$, by (iii) of Definition 3.2, it must have a root $r \in [b + 1, n]$. This is a contradiction, the root r is part of any path joining a and c and $r \notin [c, d]$, hence not in $X|_{[c,d]}$. The case where $X^{(e)}$ is fully contained in $X^{(f)}$ is argued similarly, using condition (iv) of Definition 3.2, and leads to a contradiction as well. We must conclude that \mathcal{T}_{Max} is free. \square

Remark 3.4. The non-symmetric operads \mathcal{T}_{Min} and NAP are not free. Indeed, in the operad \mathcal{T}_{Min} one has the following relation:

$$\begin{array}{c} 2 \\ \bullet \\ | \\ \bullet \\ 1 \end{array} \circ_1 \begin{array}{c} 2 \\ \bullet \\ | \\ \bullet \\ 1 \end{array} = \begin{array}{c} 2 \\ \bullet \\ | \\ \bullet \\ 1 \end{array} \circ_2 \begin{array}{c} 2 \\ \bullet \\ | \\ \bullet \\ 1 \end{array} = \begin{array}{c} 3 \\ \bullet \\ | \\ \bullet \\ 2 \\ | \\ \bullet \\ 1 \end{array}$$

And in the operad NAP one has the following relation

$$\begin{array}{c} 2 \\ \bullet \\ | \\ \bullet \\ 1 \end{array} \circ_1 \begin{array}{c} 1 \\ \bullet \\ | \\ \bullet \\ 2 \end{array} = \begin{array}{c} 1 \\ \bullet \\ | \\ \bullet \\ 2 \end{array} \circ_2 \begin{array}{c} 2 \\ \bullet \\ | \\ \bullet \\ 1 \end{array} = \begin{array}{c} 1 \quad 3 \\ \bullet \quad \bullet \\ \diagdown \quad / \\ \bullet \\ 2 \end{array}$$

Remark 3.5. Let $k\mathcal{T}_{\text{Max}}^0(n)$ denote the k -vector space spanned by the indecomposables of $\mathcal{T}_{\text{Max}}(n)$ and let β_n be its dimension. Let $\alpha(x) = \sum_{n \geq 1} \alpha_n x^n$ be the Hilbert serie associated to the free non-symmetric operad generated by the vector spaces $k\mathcal{T}_{\text{Max}}^0(n)$. It is well known (see e.g. [6]) that one has the identity

$$\beta(\alpha(x)) + x = \alpha(x),$$

where $\beta(x) = \sum_{n \geq 2} \beta_n x^n$. Theorem 3.3 implies that $\alpha_n = n^{n-1}$. As a consequence, we get that the Hilbert series for indecomposable of \mathcal{T}_{Max} is

$$\begin{aligned} \mathcal{H}_{\mathcal{T}_{\text{Max}}^0}(x) &= \sum_{n \geq 0} \dim(k\mathcal{T}_{\text{Max}}^0(n)) x^n = 2x^2 + x^3 + 14x^4 + 146x^5 + \\ &+ 1994x^6 + 32853x^7 + 630320x^8 + 13759430x^9 + \dots \end{aligned}$$

Corollary 3.6. *The non-symmetric operad pre-Lie is a free non-symmetric operad.*

Proof. The natural basis of the vector spaces $\mathcal{PL}(n)$ and $\mathcal{T}_{\text{Max}}(n)$ are the same and is given by the set of n -labelled rooted trees $\mathcal{T}(n)$. However the operad compositions differ. Following Section 2, we say that a tree in \mathcal{PL} is indecomposable if it is not the element of maximal degree in any non-trivial composition of trees in \mathcal{PL} . That is to say it is indecomposable in \mathcal{T}_{Max} . From Theorem 3.3, any rooted tree X is constructed from indecomposables in \mathcal{T}_{Max} in a unique way $\mathbb{T}(T_1, T_2, \dots, T_r)$. If we consider the sequence of operations $\mathbb{T}(T_1, T_2, \dots, T_r)$ in \mathcal{PL} , then X would be the unique maximal degree element in the result, by Proposition 2.2. By triangularity, we have that the set (in \mathcal{PL})

$$\{\mathbb{T}(T_1, T_2, \dots, T_r) : T_i \text{ indecomposable and } \mathbb{T} \text{ any planar tree of operations}\}$$

is linearly independent and span \mathcal{PL} . This means \mathcal{PL} is free. \square

Remark 3.7. The Hilbert Series for indecomposables and the operad \mathcal{PL} are the same as in Remark 3.5.

Remark 3.8. In the previous work [2] we have shown that as symmetric operads we have that $\mathcal{PL} = \mathcal{Lie} \circ F$ for some graded vector space F . We have also shown that F is naturally equipped with a structure of non-symmetric operad. One can readily adapt the method developed here to show that F is also free as a non-symmetric operad.

REFERENCES

- [1] François Bergeron, Gilbert Labelle, and Pierre Leroux. *Combinatorial species and tree-like structures*, volume 67 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1998. Translated from the 1994 French original by Margaret Readdy, With a foreword by Gian-Carlo Rota.
- [2] Nantel Bergeron and Muriel Livernet. A combinatorial basis for the free Lie algebra of the labelled rooted trees. preprint, arXiv:0707.4460, 2007.
- [3] Frédéric Chapoton and Muriel Livernet. Pre-Lie algebras and the rooted trees operad. *Internat. Math. Res. Notices*, 8:395–408, 2001.
- [4] Muriel Livernet. A rigidity theorem for pre-Lie algebras. *J. Pure Appl. Algebra*, 207(1):1–18, 2006.

- [5] Martin Markl, Steve Shnider, and Jim Stasheff. *Operads in algebra, topology and physics*, volume 96 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2002.
- [6] Paolo Salvatore and Roberto Tauraso. The operad Lie is free. arXiv:0802.3010, 2008.
- [7] James Dillon Stasheff. Homotopy associativity of H -spaces. I, II. *Trans. Amer. Math. Soc.* 108 (1963), 275-292; *ibid.*, 108:293-312, 1963.

(Nantel Bergeron) DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO M3J 1P3, CANADA

E-mail address: `bergeron@mathstat.yorku.ca`

URL, Nantel Bergeron: <http://www.math.yorku.ca/bergeron>

(Muriel Livernet) UNIVERSITÉ PARIS13, CNRS, UMR 7539 LAGA, 99, AVENUE JEAN-BAPTISTE CLÉMENT, 93430 VILLETANEUSE, FRANCE

E-mail address: `livernet@math.univ-paris13.fr`

URL, Muriel Livernet: <http://www.math.univ-paris13.fr/livernet>