



HAL
open science

Medium-modified average multiplicity and multiplicity fluctuations in jets

Redamy Perez Ramos

► **To cite this version:**

Redamy Perez Ramos. Medium-modified average multiplicity and multiplicity fluctuations in jets. 2009. hal-00338947v2

HAL Id: hal-00338947

<https://hal.science/hal-00338947v2>

Preprint submitted on 19 May 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Medium-modified average multiplicity and multiplicity fluctuations in jets

Redamy Pérez-Ramos¹

II. Institut für Theoretische Physik, Universität Hamburg
Luruper Chaussee 149, D-22761 Hamburg, Germany

Abstract: The energy evolution of average multiplicities and multiplicity fluctuations in jets produced in heavy-ion collisions is investigated from a toy QCD-inspired model. In this model, we use modified splitting functions accounting for medium-enhanced radiation of gluons by a fast parton which propagates through the quark gluon plasma. The leading contribution of the standard production of soft hadrons is enhanced by a factor $\sqrt{N_s}$ while next-to-leading order (NLO) corrections are suppressed by $1/\sqrt{N_s}$, where the parameter $N_s > 1$ accounts for the induced-soft gluons in the medium. Our results for such global observables are cross-checked and compared with their limits in the vacuum.

¹E-mail: redamy@mail.desy.de

Recent experiments at the Relativistic Heavy Ion Collider (RHIC) have established a phenomenon of strong high-transverse momentum hadron suppression [1], which supports the picture that hard partons going through dense matter suffer a significant energy loss prior to hadronization in the vacuum (for recent review see [2]).

Predictions concerning multi-particle production in nucleus-nucleus collisions can be carried out by using a toy QCD-inspired model introduced by Borghini and Wiedemann in [3]; it allows for analytical computations and may capture some important features of a more complete QCD description. In this model, the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) splitting functions $q \rightarrow g\bar{q}$ and $g \rightarrow gg$ [4] of the QCD evolution equations were distorted so that the role of soft emissions was enhanced by multiplying the infra-red singular terms by the medium factor N_s . The model [3] was further discussed and used on the description of final states hadrons produced in heavy-ion collisions [5].

Within the model, we make predictions for the medium-modified average multiplicity N_A in quark and gluon jets ($A = q, g$) produced in such reactions, for the ratio $r = N_g/N_q$ and finally for the second multiplicity correlators $\langle N_A(N_A - 1) \rangle / N_A^2$, which determines the width of the multiplicity distribution. The starting point of our analysis is the NLO or Modified-Leading-Logarithmic-Approximation (MLLA) master evolution equation for the generating functional [4] which determine the jet properties at all energies together with the initial conditions at threshold at small x , where x is the fraction of the outgoing jet energy carried away by a single gluon. Their solutions with medium-modified splitting functions can be resummed in powers of $\sqrt{\alpha_s/N_s}$ and the leading contribution can be represented as an exponential of the medium-modified anomalous dimension which takes into account the N_s -dependence:

$$N_A \simeq \exp \left\{ \int^Y \gamma_{\text{med}}(\alpha_s(Y)) dY \right\}, \quad (1)$$

where $\gamma_{\text{med}}(\alpha_s)$ can be expressed as a power series of $\sqrt{\alpha_s/N_s}$ in the symbolic form:

$$\gamma_{\text{med}}(\alpha_s) \simeq \sqrt{N_s} \times \sqrt{\alpha_s} \left(1 + \sqrt{\frac{\alpha_s}{N_s}} + \mathcal{O}\left(\frac{\alpha_s}{N_s}\right) \right).$$

Within this logic, the leading double logarithmic approximation (DLA, $\mathcal{O}(\sqrt{N_s\alpha_s})$), which resums both soft and collinear gluons, and NLO (MLLA, $\mathcal{O}(\alpha_s)$), which resums hard collinear partons and accounts for the running of the coupling constant α_s , are complete. The choice $dY = d\Theta/\Theta$, where $\Theta \ll 1$ is the angle between outgoing couples of partons in independent partonic emissions, follows from Angular Ordering (AO) in intra-jet cascades [4]. In order to obtain the hadronic spectra, we advocate for the Local Parton Hadron Duality (LPHD) hypothesis [6]: global and differential partonic observables can be normalized to the corresponding hadronic observables via a certain constant \mathcal{K} that can be fitted to the data, i.e. $N_{g,q}^h = \mathcal{K} \times N_{g,q}$.

The evolution of a jet of energy E and half-opening angle Θ involves the DLA anomalous dimension γ_0 related to the coupling constant α_s through $\gamma_0^2 = 2N_c\alpha_s/\pi$, with $\alpha_s = 2\pi/4N_c\beta_0(Y + \lambda)$, where $Y = \ln(Q/Q_0)$ ($Q = E\Theta$ is the hardness or maximum transverse momentum of the jet), $\lambda = \ln(Q_0/\Lambda_{\text{QCD}})$ is a parameter associated with hadronization (Q_0 is the collinear cut-off parameter, $k_T > Q_0$, and Λ_{QCD} is the intrinsic QCD scale) and $\beta_0 = \frac{1}{4N_c} \left(\frac{11}{3}N_c - \frac{4}{3}T_R \right)$, where $T_R = n_f/2$, n_f being the number of active flavors. At MLLA, as a consequence of angular ordering in parton cascading, the average multiplicity inside a gluon and a quark jet, $N_{g,q}$, obey the system of two-coupled evolution equations [7]

(the subscript Y denotes d/dY)

$$N_{gY} = \int_0^1 dx \gamma_0^2 [\Phi_g^g(N_g(x) + N_g(1-x) - N_g) + n_f \Phi_g^q(N_q(x) + N_q(1-x) - N_q)], \quad (2)$$

$$N_{qY} = \int_0^1 dx \gamma_0^2 [\Phi_q^g(N_g(x) + N_q(1-x) - N_g)], \quad (3)$$

which follow from the MLLA master evolution equation for the generating functional; $N_{g,q} \equiv N_{g,q}(Y)$, $N_{g,q}(x) \equiv N_{g,q}(Y + \ln x)$, $N_{g,q}(1-x) \equiv N_{g,q}(Y + \ln(1-x))$, Φ_A^B denotes the medium-modified DGLAP splitting functions:

$$\begin{aligned} \Phi_g^g(x) &= \frac{N_s}{x} - (1-x)[2 - x(1-x)], \\ \Phi_g^q(x) &= \frac{1}{4N_c}[x^2 + (1-x)^2], \\ \Phi_q^g(x) &= \frac{C_F}{N_c} \left(\frac{N_s}{x} - 1 + \frac{x}{2} \right), \end{aligned} \quad (4)$$

which accounts for parton energy loss in the medium by enhancing the singular terms like $\Phi \approx N_s/x$ as $x \ll 1$ as proposed in the Borghini-Wiedemann model [3]. Thus, when N_s increases the DLA becomes dominant and energy-momentum conservation plays a less important role.

For $Y \gg \ln x \sim \ln(1-x)$, $N_{g,q}(x)$ ($N_{g,q}(1-x)$) can be replaced by $N_{g,q}$ in the hard partonic splitting region $x \sim 1-x \sim 1$ (non-singular or regular parts of the splitting functions), while the dependence at small $x \ll 1$ is kept in the singular term $\Phi(x) \approx N_s/x$ as done in the vacuum. Furthermore, the integration over x can be replaced by the integration over $Y(x) = \ln\left(\frac{xE\Theta}{Q_0}\right)$. Thus, one is left with the approximate system of two-coupled equations,

$$\frac{d^2}{dY^2} N_g(Y) = \gamma_0^2 \left(N_s - a_1 \frac{d}{dY} \right) N_g(Y), \quad (5)$$

$$\frac{d^2}{dY^2} N_q(Y) = \frac{C_F}{N_c} \gamma_0^2 \left(N_s - \tilde{a}_1 \frac{d}{dY} \right) N_q(Y), \quad (6)$$

with the initial conditions at threshold $N_A(0) = 1$ and $N_A'(0) = 0$ and the hard constants:

$$a_1 = \frac{1}{4N_c} \left[\frac{11}{3} N_c + \frac{4}{3} T_R \left(1 - 2 \frac{C_F}{N_c} \right) \right], \quad \tilde{a}_1 = 3/4.$$

The quantum corrections $\propto a_1, \tilde{a}_1$ in (5,6) arise from the integration over the regular part of the splitting functions, they are $\mathcal{O}(\sqrt{\alpha_s/N_s})$ suppressed and *partially* account for energy conservation as happens in the vacuum.

These equations can be solved by applying the inverse Mellin transform to the self-contained gluonic equation (5), which leads to

$$N_g^h(Y) \simeq \mathcal{K} \times \int_C \frac{d\omega}{2\pi i} \omega^{\frac{a_1}{\beta_0} - 2} \exp \left[\omega(Y + \lambda) + \frac{N_s}{\beta_0 \omega} \right], \quad (7)$$

where the contour C lies to the right of all singularities of $N_g(\omega)$ in the complex plane. Since we are concerned with the asymptotic solution of the equation as $Y \gg 1$ ($E\Theta \gg Q_0$), that is the high-energy limit, the inverse Mellin transform (7) can be estimated by the steepest descent method. Indeed, the large




Figure 1: MLLA (8) medium-modified average multiplicity as a function of $Q = E\Theta$ in the vacuum ($N_s = 1$) and in the medium ($N_s = 1.6$ and $N_s = 1.8$) for $n_f = 3$.

parameter is Y and the function in the exponent presents a saddle point at $\omega_0 = \sqrt{N_s/\beta_0(Y + \lambda)}$, such that the asymptotic solution reads

$$N_g^h(Y) \simeq \mathcal{K} \times (Y + \lambda)^{-\frac{\sigma_1}{\beta_0}} \exp \sqrt{\frac{4N_s}{\beta_0}(Y + \lambda)}, \quad (8)$$

where $\sigma_1 = \frac{a_1}{2} - \frac{\beta_0}{4}$. The constant σ_1 is N_s -independent because it resums vacuum corrections. Therefore, the production of soft gluons in the medium becomes $\exp \left[2(\sqrt{N_s} - 1)\sqrt{(Y + \lambda)/\beta_0} \right]$ higher than the standard production of soft gluons in the vacuum [4]. From (1) and (8) one obtains the medium-modified MLLA anomalous dimension $\gamma_{\text{med}} = \frac{1}{N_g} \frac{dN_g}{dY} = \sqrt{N_s}\gamma_0 - \sigma_1\gamma_0^2$, which is nothing but the MLLA rate of multi-particle production with respect to the *evolution-time* variable Y in the dense medium. In Fig. 1, we display the medium-modified average multiplicity (8) with predictions in the vacuum ($N_s = 1$) in the range $10 \leq Q(\text{GeV}) \leq 200$; we set $Q_0 = \Lambda_{\text{QCD}} = 0.23$ GeV in the limiting spectrum approximation [7], and take $\mathcal{K} = 0.2$ from [7]. The values $N_s = 1.6$ and $N_s = 1.8$ in the medium may be realistic for RHIC and LHC phenomenology [3, 5]; the jet energy subrange $10 \leq Q(\text{GeV}) \leq 50$ displayed in Fig. 1 has been recently considered by the STAR collaboration, which reported the first measurements of charged hadrons and particle-identified fragmentation functions from p+p collisions [8] at $\sqrt{s_{\text{NN}}} = 200$ GeV. Finally, the whole jet energy range in the same figure, in particular for those values at $Q \geq 50$ GeV, will be reached at the LHC, i.e $Q = 100$ GeV is an accessible value in this experiment (see [3] and references therein).

We find, as expected, that the production of soft hadrons increases as $N_s > 1$: the available phase space for the production of harder collinear hadrons is restricted as the model itself states. The medium-modified MLLA gluon to quark average multiplicity ratio, $r = N_g/N_q = N_g^h/N_q^h$, following from (8) and (3) reads

$$r = r_0 \left[1 - r_1 \frac{\gamma_0}{\sqrt{N_s}} + \mathcal{O} \left(\frac{\gamma_0^2}{N_s} \right) \right], \quad r_0 = \frac{N_c}{C_F}, \quad (9)$$

where we introduced the coefficient $r_1 = a_1 - \tilde{a}_1$ in the term suppressed by $\gamma_0/\sqrt{N_s}$ as $N_s > 1$. Therefore, if compared with its behavior at $N_s = 1$, we check, as expected from the model [3], that r becomes closer to its asymptotic DLA limit $r_0 = N_c/C_F = 9/4$, as depicted in Fig. 2. Setting $N_s = 1$ in (9), one recovers the appropriate limits in the vacuum [4, 9, 10]. Finally, the gluon jets are still more




Figure 2: MLLA ratio r (9) as a function of $Q = E\Theta$ in the vacuum ($N_s = 1$) and in the medium ($N_s = 1.6$ and $N_s = 1.8$) for $n_f = 3$.

active than the quark jets in producing secondary particles and the shape of the curves are roughly the same.

The normalized second multiplicity correlator $A_2 = \langle N_A(N_A - 1) \rangle / N_A^2$ defines the width of the multiplicity distribution and is related to its dispersion by the formula $D_A^2 = (A_2 - 1)N_A^2 + N_A$ [9]. These moments, which are less inclusive than the average multiplicity, prove to be \mathcal{K} -independent and therefore provide a pure test of multiparticle production. The medium-modified system of two-coupled evolution equations for this observable follows from the MLLA master equation for the azimuthally averaged generating functional [4] and can be written in the convenient form

$$\begin{aligned} \frac{d}{dY}(N_g^{(2)} - N_g^2) &= \int_0^1 dx \gamma_0^2 \Phi_g^g \left[N_g^{(2)}(Y + \ln x) + \left(N_g^{(2)}(Y + \ln(1-x)) - N_g^{(2)}(Y) \right) \right. \\ &\quad \left. + \left(N_g(Y + \ln x) - N_g(Y) \right) \left(N_g(Y + \ln(1-x)) - N_g(Y) \right) \right] \\ &\quad + n_f \int_0^1 dx \gamma_0^2 \Phi_g^q \left[2 \left(N_q^{(2)}(Y + \ln x) - N_q^2(Y + \ln x) \right) - \left(N_g^{(2)}(Y) - N_g^2(Y) \right) \right. \\ &\quad \left. + \left(2N_q(Y + \ln x) - N_g(Y) \right) \left(2N_q(Y + \ln(1-x)) - N_g(Y) \right) \right], \end{aligned} \quad (10)$$

$$\begin{aligned} \frac{d}{dY}(N_q^{(2)} - N_q^2) &= \int_0^1 dx \gamma_0^2 \Phi_q^g \left[N_g^{(2)}(Y + \ln x) + \left(N_q^{(2)}(Y + \ln(1-x)) - N_q^{(2)}(Y) \right) \right. \\ &\quad \left. + 2 \left(N_g(Y + \ln x) - N_q(Y) \right) \left(N_q(Y + \ln(1-x)) - N_q(Y) \right) \right], \end{aligned} \quad (11)$$

which proves to be more suitable for obtaining analytical solutions in the following. We use a new method to compute solutions at MLLA by replacing $N_A^{(2)} = A_2 N_A^2$ on both sides of the expanded equations at $x \sim 1 - x \sim 1$. The notations in (10,11) follow the same logic as those in (5,6). Applying the analysis that led to the system (5,6), we obtain from (10,11)

$$\frac{d^2}{dY^2} (N_g^{(2)} - N_g^2) = \gamma_0^2 \left(N_s - a_1 \frac{d}{dY} \right) N_g^{(2)} + (a_1 - b_1) \gamma_0^2 \frac{d}{dY} N_g^2, \quad (12)$$

$$\frac{d^2}{dY^2} (N_q^{(2)} - N_q^2) = \frac{C_F}{N_c} \gamma_0^2 \left(N_s - \tilde{a}_1 \frac{d}{dY} \right) N_q^{(2)}, \quad (13)$$

where

$$b_1 = \frac{1}{4N_c} \left[\frac{11}{3} N_c - 4 \frac{T_R}{N_c} \left(1 - 2 \frac{C_F}{N_c} \right)^2 \right].$$




Figure 3: MLLA second multiplicity correlator inside a gluon jet (15) as a function of $Q = E\Theta$ in the vacuum ($N_s = 1$) and in the medium ($N_s = 1.6$ and $N_s = 1.8$ for $n_f = 3$).

The constant N_s only affects the leading double logarithmic term of the equations. The terms proportional to a_1 , $(a_1 - b_1)$ and \tilde{a}_1 are hard vacuum corrections, which *partially* account for energy conservation, indeed $\gamma_0^2 \frac{dN}{dY} \approx \sqrt{N_s} \gamma_0^3$ and the relative correction to DLA is $\mathcal{O}(\sqrt{\alpha_s/N_s})$.

Setting $N_g^{(2)} = G_2 N_g^2$ in (12) and making use of (8), the system can be solved iteratively by taking terms up to $\mathcal{O}(\alpha_s)$ into consideration. The analytical solution reads,

$$G_2 - 1 = \frac{1 - \left(\frac{2}{3}a_1 + 2b_1\right) \frac{\gamma_0}{\sqrt{N_s}} + \mathcal{O}\left(\frac{\gamma_0^2}{N_s}\right)}{3 - (4a_1 - \beta_0) \frac{\gamma_0}{\sqrt{N_s}} + \mathcal{O}\left(\frac{\gamma_0^2}{N_s}\right)}, \quad (14)$$

while its expansion in the form $1 + \gamma_0/\sqrt{N_s}$ leads to

$$G_2 - 1 \approx \frac{1}{3} - c_1 \frac{\gamma_0}{\sqrt{N_s}} + \mathcal{O}\left(\frac{\gamma_0^2}{N_s}\right), \quad (15)$$

where the linear combination of color factors can be written in the form

$$c_1 = \frac{1}{4N_c} \left(\frac{55}{9} - 4 \frac{T_R}{N_c} + \frac{112 T_R C_F}{9 N_c N_c} - \frac{32 T_R C_F^2}{3 N_c N_c^2} \right). \quad (16)$$

We use (15) and (9) and substitute $N_q^{(2)} = Q_2 N_q^2$ into (13) such that the solution reads

$$Q_2 - 1 \approx \frac{N_c}{C_F} \left(\frac{1}{3} - \tilde{c}_1 \frac{\gamma_0}{\sqrt{N_s}} \right) + \mathcal{O}\left(\frac{\gamma_0^2}{N_s}\right), \quad (17)$$

where we obtain the combination of color factors

$$\tilde{c}_1 = \frac{1}{4N_c} \left(\frac{55}{9} + \frac{4 T_R C_F}{9 N_c N_c} - \frac{8 T_R C_F^2}{3 N_c N_c^2} \right). \quad (18)$$

Setting $N_s = 1$ in (15) and (17) we get a perfect agreement with the vacuum results [9]. In Fig. 3 and Fig. 4, we compare our results for the medium-modified second multiplicity correlators (15) and (17) with predictions in the vacuum ($N_s = 1$) [9] in the limiting spectrum approximation inside the typical range $10 \leq Q(\text{GeV}) \leq 200$ for RHIC and LHC phenomenology. Similarly to the MLLA ratio $r(N_s)$, Eq. (9), the hard corrections $\mathcal{O}(\gamma_0)$ are suppressed by a factor $1/\sqrt{N_s}$. As expected from the model,




Figure 4: MLLA second multiplicity correlator inside a quark jet (17).

we check that these results approach their DLA limits when N_s increases; moreover, the multiplicity fluctuations of individual events must be larger for quark jets as compared to gluon jets just like in the vacuum [9]. Another interesting feature of these observables concerns the shape of the curves. They are roughly identical and prove not to depend on the medium parameter N_s . Moreover, there exists evidence for a flattening of the slopes as the hardness of the jet $Q = E\Theta$ increases for $N_s \geq 1$ (vacuum and medium). This kind of scaling behavior is known as the Koba-Nielsen-Olsen (KNO) scaling [11]: it was discovered by Polyakov in quantum field theory [12] and experimentally confirmed by e^+e^- measurements [13] for the second and higher order multiplicity correlators.

In this paper we have dealt with the medium-modified average multiplicity and the medium-modified second multiplicity correlator in quark and gluon jets at RHIC and LHC energy scales. The starting point of our calculations is based on the Borghini-Wiedemann work [3], which models parton energy loss in a nuclear medium. The average multiplicity is found to be enhanced by the factor $\sqrt{N_s}$ acting on the exponential leading contribution (8); this leads in particular to the rescaling of the anomalous dimension γ_{med} ($\gamma \rightarrow \gamma_{\text{med}} \approx \sqrt{N_s}\gamma_0$) or equivalently, to the enhancement of the in medium coupling constant. Since hard corrections are suppressed by the extra factor $1/\sqrt{N_s}$, it is straightforward to check that r , G_2 and Q_2 approach the asymptotic DLA limits $r_0 = N_c/C_F$, $G_2 = 4/3$ and $Q_2 = 1 + N_c/3C_F$ [4] when N_s increases. The previously mentioned KNO-scaling experienced by G_2 and Q_2 proves no special sensibility to the model and should normally hold like in the vacuum.

Finally, since these results are model-dependent, they may still be improved in the future, specially after the N_s -dependence of the non-singular parts of the splitting functions (4) has been exactly computed.

Perspective: Many experimental characterizations of the medium-modified intrajet structure in heavy-ion collisions at RHIC and at the LHC require a soft momentum cut-off p_T^{cut} , with $Q > p_T^{\text{cut}}$ to remove the effects of the high multiplicity background. In [3], the soft background was subtracted by integrating the single inclusive differential distribution $\frac{dN}{d \ln p_T}$ (“hump-backed plateau”) over the range $Q \geq p_T \geq p_T^{\text{cut}}$, with $p_T^{\text{cut}} > \Lambda_{QCD}$. Accordingly, the equivalent computation should be performed for the second multiplicity correlator by integrating the double differential inclusive distribution (two-particle correlation) $\frac{d^2N}{d \ln p_{1,T} d \ln p_{2,T}}$ over $p_{i,T}$, with the lower bounds of integration $p_{i,T}^{\text{cut}} > \Lambda_{QCD}$ ($i = 1, 2$). Imposing such a cut-off in our calculations will affect the normalization rather than the behavior and the shape of these observables as a function of N_s and the jet energy scale of the process Q [14].

References

- [1] K. Adcox et al. (PHENIX Collab.), Phys. Rev. Lett. **88** (2002) 022301;
S.S. Adler et al. (PHENIX Collab.), Phys. Rev. Lett. **91** (2003) 072301;
C. Adler et al. (STAR Collab.), Phys. Rev. Lett **89** (2002) 202301.
- [2] F. Arleo, hep-ph/0810.1193 and references therein;
S. Peigné & A.V. Smilga, hep-ph/0810.5702.
- [3] N. Borghini & U.A. Wiedemann, hep-ph/0506218.
- [4] Yu.L. Dokshitzer, V.A. Khoze, A.H. Mueller & S.I. Troyan, Basics of Perturbative QCD, Editions Frontières, Paris (1991).
- [5] S.Sapeta & U.A Wiedemann, Eur. Phys. J. **C55** (2008) 293.
- [6] Ya.I. Azimov, Yu.L. Dokshitzer, V.A. Khoze & S.I. Troian, Z. Phys. **C 27** (1985) 65; Yu.L. Dokshitzer, V.A. Khoze & S.I. Troian, J. Phys. **G 17** (1991) 1585.
- [7] I.M. Dremin, V.A. Nechitailo, Mod. Phys. Lett. **A 9** (1994) 1471; JETP Lett. **58** (1993) 945.
- [8] M. Heinz for the STAR Collaboration, nucl-exp/0809.3769.
- [9] E.D. Malaza & B.R. Webber, Phys. Lett. **B 149** (1984) 501; E.D. Malaza & B.R. Webber, Nucl. Phys. **B 267** (1986) 702.
- [10] V.A. Khoze & W. Ochs, Int. J. Mod. Phys. **A 12** (1997) 2949.
- [11] Yu.L. Dokshitzer, Phys. Lett. **B 305** (1993) 295.
- [12] A.M. Polyakov, Sov. Phys. JETP **32** (1971) 296.
- [13] HRS Coll., Phys. Rev. **D 34** (1986) 3304; AMY Coll., Phys. Rev. **D 42** (1990) 737; DELPHI Coll., Z. Phys. **C-Particles and Fields 50** (1991) 185.
- [14] N. Borghini & R. Perez-Ramos, in preparation.