



HAL
open science

Photoionisation loading of large Sr⁺ ion clouds with ultrafast pulses

Sébastien Removille, Romain Dubessy, Quentin Glorieux, Samuel Guibal,
Thomas Coudreau, Luca Guidoni, Jean-Pierre Likforman

► **To cite this version:**

Sébastien Removille, Romain Dubessy, Quentin Glorieux, Samuel Guibal, Thomas Coudreau, et al..
Photoionisation loading of large Sr⁺ ion clouds with ultrafast pulses. 2008. hal-00340119v1

HAL Id: hal-00340119

<https://hal.science/hal-00340119v1>

Preprint submitted on 20 Nov 2008 (v1), last revised 5 Dec 2008 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Photoionisation loading of large Sr^+ ion clouds with ultrafast pulses.

S. Removille, R. Dubessy, Q. Glorieux, S. Guibal,

T. Coudreau, L. Guidoni, and J.-P. Likforman*

Université Paris-Diderot,

Laboratoire Matériaux et Phénomènes Quantiques

CNRS UMR 7162

10, rue A. Domon et L. Duquet,

75013 Paris, France

Abstract

This paper reports on photoionisation loading of Sr^+ ions in a linear Paul trap based on ultrafast pulses. We benefit from an autoionising resonance of strontium to photoionise the neutral atoms using two-photon absorption of femtosecond pulses at $\lambda = 431$ nm. We compare this technique to electron-bombardment and demonstrate clear advantages of using photoionisation. It actually allows the loading of a pure Sr^+ ions and the use of lower radio-frequency voltage amplitudes for the trapping. In these conditions, and in the absence of fine minimization of the micro-motion responsible of rf heating, up to 4×10^4 cooled Sr^+ ions could be trapped.

INTRODUCTION

Samples of laser-cooled ions confined in electromagnetic traps play a prominent role in several domains related to atomic physics: quantum information[1], metrology[2], quantum optics[3]. The traditional strategy used to load an ion trap with the desired species is to create the ions directly inside the trapping region by electron-bombardment (EB) of a neutral atomic beam. As already underlined by several groups[4, 5, 6] this technique, while very flexible because it applies to any atomic (and molecular) species, has several serious drawbacks. First, the electron beam can eventually charge some insulator present in the neighbouring of the trap affecting the trapping potential with a slowly varying non-controlled electric field. This situation result in an excess of radio-frequency (RF) heating that imposes to tune frequently compensation voltages[5, 6]. Second, the vacuum quality of the setup is deteriorated by the presence a hot filament near the trap. Third, the very small cross-section of the electron-impact ionisation requires huge atomic fluxes that negatively affect both the vacuum and the electrode surfaces. In recent years, several groups have developed photo-ionisation techniques that virtually eliminate these drawbacks. We can distinguish two different methods that have been introduced, depending on the excitation path from the neutral atom to the ionised state. In what we will call "two-step" photoionisation (TSPI) a first narrowband cw laser is tuned on an intermediate transition and the ionisation threshold is attained with a second photon (that can eventually have the same energy). This method needs a non-negligible population in the intermediate level (saturating stabilized laser). The other method is based on a two-photon transition that directly brings a neutral atom above the ionisation threshold (two-photon photo-ionisation, TPPI). TPPI requires high peak-power, normally associated to ultrafast pulses.

TSPI has the important advantage to be potentially isotope-selective through the isotopic-shifts associated to the intermediate transition. It has been applied to Mg [7, 8], Ca [4, 5, 8], Ba [9], Yb [10], and Sr [6]. On the other hand, TPPI could become an easier solution in case of far ultra-violet (UV) transitions (e.g. to create Cd^+ ions[11]). That is due to the large tunability-range of Ti:Sa femtosecond lasers and to the efficiency of frequency-doubling or tripling in the ultrafast regime.

This paper reports on photoionisation loading of Sr^+ in a linear Paul trap based on ultrafast pulses. We demonstrate that TPPI presents indeed several advantages with respect

to EB. In particular it allowed us to explore trapping regimes with low RF voltages, to obtain larger ion clouds and to improve the vacuum quality by lowering the power in the Sr oven. Such a lower atom flux is particularly interesting in the case of micro-fabricated traps because it improves the electrode lifetime and reduces heating associated to patch potentials[12]. Let us finally mention that, because of its two-photon character, TPPI has the potential advantage to define an extremely well-localized spatial region for the ion production associated with the waist of the ionising beam[13]. This facilitates the ion creation at the center of the trap with an initially low temperature.

EXPERIMENTAL SETUP

The level structure of Sr atom is similar to that of other atoms with two electrons on the external shell (the relevant levels are represented in Figure 1, upper part). The 1P_1 intermediate level has been used to implement TSPI technique in Sr[6], following the example of Ca[4] and Yb[10]. In the particular case of the Sr atom, the presence of a $(4d^2 + 5p^2)^1D_2$ autoionising state that lies above the ionisation threshold is very helpful to increase the photoionisation efficiency. The spectral characteristics of this state have been investigated earlier by two-color spectroscopy, in particular a cross section of 5600 Mb for the $^1P_1 \rightarrow (4d^2 + 5p^2)^1D_2$ transition has been measured[14]. Such an autoionising state can also be reached from the ground state by two-photon absorption: the $(5s^2)^0S_0 \rightarrow (4d^2 + 5p^2)^1D_2$ two-photon transition has a linewidth of 0.7 nm FWHM (due to the short lifetime of the final state) and can be resonantly driven by two photons centered at 431 nm [15]. The strategy of the TPPI loading experiment described in this paper is based on this transition (Figure 1).

The neutral Sr atom beam is produced in an oven placed below the trap and formed by a tungsten filament in which flows a maximum current of 1.15A, that corresponds to a heating power of 1.3W. The photoionising laser source is based on a homemade femtosecond Titanium:Sapphire oscillator that produces 50 fs pulses of 1.5 nJ energy at $\lambda = 862$ nm with a repetition rate of 100 MHz. The pulses are frequency doubled in a 0.5 μm thick Beta Barium Borate (BBO) nonlinear crystal. A pulse energy of more than 0.15 nJ is routinely obtained at $\lambda = 431$ nm. The beam is directed into the vacuum chamber and focused at the center of the ion trap by a lens of focal distance 200 mm. The measured photoionising

beam size at the waist is $20 \pm 2 \mu\text{m}$.

Sr^+ ions are trapped in a standard linear Paul trap (see Figure 2) [16]. Four parallel rod electrodes (diameter 6.35 mm, inner radius $r_0 = 3.2 \text{ mm}$) are used for the radio-frequency (rf) radial confinement. A 2.5 MHz rf potential with an amplitude V_{rf} in the range 50–400 V is applied to two of the diagonally opposed electrodes. The remaining two rods are normally grounded or used to resonantly excite the ion motion in the trap (see below). The Sr^+ ion radial movement is defined by a typical secular frequency $\nu_R = 400 \text{ kHz}$ for an applied voltage $V_{rf} = 500 \text{ V}$. Two annular "end caps" separated by 20 mm are used for the longitudinal confinement and brought to a static positive voltage ($V_{ec} = 500 \text{ V}$). The corresponding axial frequency is $\nu_A = 20 \text{ kHz}$. Trapped ions can be detected in a destructive way by an ejection sequence: an asymmetric potential is applied to the "end caps" that kicks out the cloud along the trap axis through an electrostatic lens towards an electron multiplier. This detection scheme allows us to perform ion-counting and to analyze the trapped species by performing mass-spectra. A mass spectrum is obtained by measuring the losses induced in successive realization of the cloud in the presence of a "tickle" sinusoidal excitation applied to two diagonally opposed rods. The result of this parametric excitation is a depletion of the ion signal when the tickle frequency matches an integer fraction of $2\nu_R$ (negative peak in the mass spectrum)[17].

Trapped Sr^+ ions are Doppler cooled using the $^2S_{1/2} \rightarrow ^2P_{1/2}$ optical transition at $\lambda = 421.7 \text{ nm}$ (natural linewidth, $\Gamma/2\pi = 20.2 \text{ MHz}$, Figure 2, bottom). This transition is driven using laser light generated by a commercial extended-cavity diode laser (Toptica DL100). Optical pumping into the metastable "clock" $^2D_{3/2}$ state is avoided by employing a commercial fiber-laser (Koheras Adjustik Y10) emitting light at $\lambda = 1091.8 \text{ nm}$. Let us mention that this setup allows us to perform both the measurement of the ion fluorescence, useful for optimizing the cooling phase, and the precise evaluation of the number of trapped ions by ejection and ion-counting detection. An example of a fluorescence image of a large cloud (2 mm length and $200 \mu\text{m}$ width, FWHM) containing $4 \times 10^4 \text{ Sr}^+$ ions is shown in Figure 3.

RESULTS

In a first experiment, we performed mass spectra of the trapped ions produced either by EB or by TPPI. Typical mass spectra corresponding to these techniques are shown on Figure 4. Basically two main peaks associated to Sr^+ are expected and observed, corresponding to a tickle frequency of $2\nu_R$ and ν_R (negative peaks in the mass spectrum). This frequency (400 kHz for the experimental conditions of Figure 4) depends on the trap potentials and the ion mass. Secondary satellite peaks of lower amplitude separated by $\nu_A \simeq 40$ kHz are visible at the second harmonic frequency $2\nu_R$ (800 kHz) that originate from the coupling between the axial and longitudinal movement in the large ion clouds[18]. The main difference between the two spectra is the value of the peak contrast compared to the background level. The contrast, of the saturated peak at $2\nu_R$ gives us information about the proportion of Sr^+ ions in the trap. TPPI produces a contrast of 100%, indicating that only Sr^+ ions are loaded, compared to 66% in the case of EB for which other ion species are also produced. This result demonstrates the advantage of selectivity offered by photoionisation for applications such as quantum memories based on large ion clouds[19]. As we demonstrated that only strontium ions are produced in the trap when using TPPI, we also confirmed that this ionisation process is actually based on two-photon absorption. Indeed, the ionisation rate as function of the ionising beam intensity shows the expected quadratic dependence (Figure 5).

In a second experiment, we compared the loading times and the loading levels in the two cases of EB and TPPI. The experiment is performed as follows: we turn-on the electron gun or the photoionising beam at time $\tau = 0$; after a certain time τ , we turn off the laser cooling and eject the loaded ions towards the ion counter. Laser cooling is stopped in the ejection phase to avoid spurious losses induced by damping. We show in Figure 6a the number of trapped ions, as function of τ , obtained using the two techniques. In both cases, saturation in the number of trapped ions is reached within a few tens of seconds. However, only TPPI allowed us to create clouds containing a large number of strontium ions: the EB technique saturates around 1000 trapped ions. In fact, with TPPI an optimum for the number of trapped Sr^+ ions is obtained for $V_{rf} = 125$ V, that is for a relatively shallow trap. In the case of EB, it is impossible to explore the same range for the parameter V_{rf} since too many spurious ionic species are produced and trapped in shallow potential wells. This

effect is clearly visible in the mass spectra of Figure 6b obtained with the two techniques for $V_{rf} = 350$ V (dotted curve). Therefore, in our case of a non-crystalized regime, the possibility to work at a low V_{rf} is crucial in order to obtain a large number of trapped ions. We interpret this result in terms of rf-heating process that increases with increasing V_{rf} [20]. In this situation, and in the presence of laser-cooling, the ion-density depends on the balance between cooling and heating. A maximum density is then be achievable by minimizing V_{rf} (i.e. the rf heating) within the stability region. But the total number of trapped ions also depends on the volume of the trap that is imposed by V_{rf} and V_{ec} . Numerical simulations carried by Simion software allowed us to evaluate the trap volume as a function of V_{rf} for $V_{ec} = 500$ V. An optimum (maximum) volume is obtained for $V_{rf} = 180$ V, slightly larger than the experimental value that maximizes the total number of trapped ions.

We also observed that, when working in the regime of relatively high V_{rf} =, that is optimal for EB ($V_{rf} = 500V$), 1100 ions can be trapped by EB to be compared to 1200 ions obtained by photoionisation. This small discrepancy might be explained by space charge effects due to electrons that can perturb the trap electrostatic potential or by the intrinsic possibility, and advantage, of photoionisation to produce ions at the centre of the trap.

CONCLUSION

We demonstrated two-photon photoionisation loading of Sr^+ in a linear Paul trap using ultrafast pulses centered at $\lambda = 431$ nm. We compared this technique to the electron bombardment loading and observed several advantages, already mentioned in previous experiments concerning other species or other photo-excitation paths. In particular this technique allowed us to explore trapping regimes with low RF voltages, to obtain larger ion clouds and to improve the vacuum quality by lowering the power in the Sr oven. In these conditions, and in the absence of fine minimization of the micro-motion responsible of rf heating, up to 4×10^4 cooled Sr^+ ions could be trapped.

ACKNOWLEDGMENTS

We thank P. Lepert for technical support and N. Sangouard for useful discussions. The authors would also like to thank M. Joffre for the lending of the femtosecond oscillator. This

work was supported by ANR "jeunes chercheuses et jeunes chercheurs" research contract JC05_61454.

-
- * Electronic address: Jean-Pierre.likforman@univ-paris-diderot.fr
- [1] H. Häffner, C. F. Roos, and R. Blatt, *Quantum computing with trapped ions* Phys. Rep. in press (2008) <http://dx.doi.org/10.1016/j.physrep.2008.09.003>; R. Blatt and D. Wineland, *Entangled states of trapped atomic ions* Nature, **453**, 1008 (2008)
- [2] P. Gill, *Optical frequency standards* Metrologia **42** S125-S137 (2005) <http://dx.doi.org/10.1088/0026-1394/42/3/S13>
- [3] G.R. Guthöhrlein, M. Keller, K. Hayasaka, W. Lange, H. Walther, *A single ion as a nanoscopic probe of an optical field* Nature **414**, 49 (2001) <http://dx.doi.org/10.1038/35102129>; J. Eschner, C. Raab, F. Schmidt-Kaler, R. Blatt, *Light interference from single atoms and their mirror images* Nature **413**, 495 (2001) <http://dx.doi.org/10.1038/35097017>; P. Herskind, A. Dantan, M.B. Langkilde-Lauesen, A. Mortensen, J.L. Sørensen, M. Drewsen, *Loading of large ion Coulomb crystals into a linear Paul trap incorporating an optical cavity* Appl. Phys. B **93** 373-379 (2008) <http://dx.doi.org/10.1007/s00340-008-3199-8>
- [4] S. Gulde, D. rotter, P. Barton, F. Schmidt-Kaler, R. Blatt, W. Hogervorst, *Simple and efficient photo-ionisation loading of ions for precision ion-trapping experiments* Appl. Phys. B **73**, 861-863 (2001) <http://dx.doi.org/10.1007/s003400100749>
- [5] D. M. Lucas, A. Ramos, J. P. Home, M. J. McDonnell, S. Nakayama, J.-P. Stacey, S. C. Webster, D. N. Stacey, and A. M. Steane, *Isotope-selective photoionisation for calcium ion trapping* Phys. Rev. A **69** 012711 (2004) <http://dx.doi.org/10.1103/PhysRevA.69.012711>
- [6] M. Brownnutt, V. Letchumanan, G. Wilpers, R. C. Thompson, P. Gill, A. G. Sinclair, *Controlled photoionisation loading of $^{88}\text{Sr}^+$ for precision ion-trap experiments* Appl. Phys. B **87** 411-415 (2007) <http://dx.doi.org/10.1007/s00340-007-2624-8>
- [7] D. N. Madsen, S. Balslev, M. Drewsen, N. Kjærgaard, Z. Videsen, and J. W. Thomsen, *Measurements on photo-ionisation of $3s3p\ ^1P_1$ magnesium atoms* J. Phys. B **33**, 4981-4988 (2000). <http://dx.doi.org/10.1088/0953-4075/33/22/302>
- [8] N. Kjærgaard, L. Hornekær, A. M. Thommesen, Z. Videsen, M. Drewsen *Isotope selective loading of an ion trap using resonance-enhanced two-photon ionisation* Appl. Phys. B **71**,

- 207?210 (2000) <http://dx.doi.org/10.1007/s003400000296>
- [9] A. V. Steele, L. R. Churchill, P. F. Griffin, and M. S. Chapman, *Photoionisation and photoelectric loading of barium ion traps* Phys. Rev. A **75** 053404 (2007) <http://dx.doi.org/10.1103/PhysRevA.75.053404>
- [10] K. Hosaka, S. A. Webster, P. J. Blythe, A. Stannard, D. Beaton, H. S. Margolis, S. N. Lea, and P. Gill, *An optical frequency standard based on the electric octupole transition in $^{171}\text{Yb}^+$* IEEE Trans. Instrum. Meas. **54**, 759-762 (2005) <http://dx.doi.org/10.1109/TIM.2004.843319>; Chr. Balzer, A. Braun, T. Hannemann, Chr. Paape, M. Ettl, W. Neuhauser, and Chr. Wunderlich, *Electrodynamically trapped Yb^+ ions for quantum information processing* Phys. Rev. A **73**, 041407R (2006). <http://dx.doi.org/10.1103/PhysRevA.73.041407>
- [11] L. Deslauriers, M. Acton, B. B. Blinov, K.-A. Brickman, P. C. Haljan, W. K. Hensinger, D. Hucul, S. Katnik, R. N. Kohn, Jr., P. J. Lee, M. J. Madsen, P. Maunz, S. Olmschenk, D. L. Moehring, D. Stick, J. Sterk, M. Yeo, K. C. Younge, and C. Monroe, *Efficient photoionisation loading of trapped ions with ultrafast pulses* Phys. Rev. A **74** 063421 (2006) <http://dx.doi.org/10.1103/PhysRevA.74.063421>.
- [12] Ralph G. DeVoe and Christian Kurtsiefer, *Experimental study of anomalous heating and trap instabilities in a microscopic ^{137}Ba ion trap* Phys. Rev. A **65** 063407 (2002) <http://dx.doi.org/10.1103/PhysRevA.65.063407>.
- [13] A. diaspro (Ed.), *Confocal and Two-Photon Microscopy: Foundations, Applications and Advances* Wiley-VCH, Weinheim, 2001; ISBN 0-471-40920-0.
- [14] W. Mende, K. Bartschat and M. Kock, *Near-threshold photoionisation from the $\text{Sr I } (5s5p)^1P_1^0$ state* J. Phys. B **28**, 2385?2393 (1995). <http://dx.doi.org/10.1088/0953-4075/28/12/008>
- [15] M. A. Baig, M. Yaseen, Raheel Ali, Ali Nadeem, S.A. Bhatti, *Near-threshold photoionisation spectra of strontium* Chem. Phys. Lett. **296**, 403?408 (1998). [http://dx.doi.org/10.1016/S0009-2614\(98\)01059-8](http://dx.doi.org/10.1016/S0009-2614(98)01059-8)
- [16] J. D. Prestage, G. J. Dick, and L. Maleki, *New ion trap for frequency standard applications* J. Appl. Phys. **66**, 1013 (1989). <http://dx.doi.org/10.1063/1.343486>
- [17] L. Guidoni, S. Guibal, T. Coudreau, B. Dubost and F. Grosshans, *Dynamique d'ions confinés dans un piège de Paul linéaire* J. Phys. IV France **135** 315-316, (2006). <http://dx.doi.org/10.1051/jp4:2006135104>
- [18] R. Dubessy, S. Removille, B. Dubost, S. Guibal, T. Coudreau, and L. Guidoni, *Nonlinear*

- couplings and cooling dynamics in a large Paul trap designed for quantum information* in CLEO/Europe and IQEC 2007 Conference Digest, (Optical Society of America, 2007), paper IC_17. http://www.opticsinfobase.org/abstract.cfm?URI=IQEC-2007-IC_17
- [19] T. Coudreau, F. Grosshans, S. Guibal and L. Guidoni, *Feasibility of a quantum memory for continuous variables based on trapped ions: from generic criteria to practical implementation*, J. Phys. B: At. Mol. Opt. Phys. **40**, 413 (2007) <http://dx.doi.org/10.1088/0953-4075/40/2/014>
- [20] R. Blümel, J. M. Chen, E. Peik, W. Quint, W. Schleich, Y. R. Shen, and H. Walther, *Phase transitions of stored laser-cooled ions* Nature **334**, 309 (1988). <http://dx.doi.org/10.1038/334309a0>




FIG. 1: Upper part: relevant energy levels of neutral Strontium involved in two-step photo-ionisation (TSPI, left) and in two-photon photo-ionisation (TPPI, right). Lower part: relevant energy levels of $Sr+$ ion involved in Doppler cooling.




FIG. 2: Schematic description of the linear Paul trap setup. Four parallel rod electrodes are used for the rf radial confinement. Two annular "end caps" separated by 20 mm are used for the longitudinal confinement. The trapped ions can be axially ejected and counted by an electron multiplier. The ion cloud fluorescence resulting from Doppler cooling is recorded on a ccd camera.




FIG. 3: Fluorescence image at $\lambda = 422$ nm of a trapped ion cloud (2 mm length and $200 \mu\text{m}$ width, FWHM) containing 3×10^4 Sr^+ ions




FIG. 4: comparison of mass spectra of the trapped ions produced either by electron-beam or by photoionisation. The two main peaks associated to Sr^+ are observed for a tickle frequency of $2\nu_R$ and ν_R (negative peaks in the mass spectrum). $V_{rf} = 500$ V.




FIG. 5: Photoionisation rate as function of the average power of the ionisation beam. A perfect quadratic dependence is observed as expected for a two-photon absorption ionisation process.




FIG. 6: a) Number of trapped ions as function of the loading time. In the case of photoionisation, the curves corresponding to several V_{rf} are traced. b) Mass spectra of the trapped ions produced either by electron-beam (dotted curve) or by photoionisation (solid curve) for $V_{rf} = 350$ V.