



**HAL**  
open science

## Distinction of some induced representations

Nadir Matringe

► **To cite this version:**

| Nadir Matringe. Distinction of some induced representations. 2009. hal-00340843v6

**HAL Id: hal-00340843**

**<https://hal.science/hal-00340843v6>**

Preprint submitted on 28 Feb 2009 (v6), last revised 21 Oct 2009 (v13)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Distinction of some induced representations

Nadir MATRINGE\*

February 28, 2009

## Abstract

Let  $K/F$  be a quadratic extension of  $p$ -adic fields,  $\sigma$  the nontrivial element of the Galois group of  $K$  over  $F$ , and  $\pi$  a smooth representation of  $GL(n, K)$ . Denoting by  $\pi^\vee$  the smooth contragredient of  $\pi$ , and by  $\pi^\sigma$  the representation  $\pi \circ \sigma$ , we show that the representation of  $GL(2n, K)$  obtained by normalized parabolic induction of the representation  $\pi^\vee \otimes \pi^\sigma$  is distinguished with respect to  $GL(2n, F)$ . This is a step towards the classification of distinguished generic representations of general linear groups over  $p$ -adic fields.

## Introduction

Let  $K/F$  be a quadratic extension of  $p$ -adic fields,  $\sigma$  the nontrivial element of the Galois group of  $K$  over  $F$ , and  $\pi$  a smooth representation of  $GL(n, K)$ . We denote by  $\sigma$  again the automorphism of  $M_{2n}(K)$  induced by  $\sigma$ .

A smooth representation  $\rho$  of  $GL(2n, K)$  is said to be distinguished if there is a nonzero  $GL(2n, F)$ -invariant linear form on its space. If  $\rho$  is irreducible, the space of such forms is of dimension at most 1.

Calling  $\pi^\vee$  the smooth contragredient of  $\pi$  and  $\pi^\sigma$  the representation  $\pi \circ \sigma$ , we denote by  $\pi^\sigma \times \pi^\vee$  the representation of  $GL(2n, K)$ , obtained by normalized induction of the representation  $\pi^\sigma \otimes \pi^\vee$  of the standard parabolic subgroup of type  $(n, n)$ . The aim of this work is to show that such a representation is distinguished.

The case  $n = 1$  is treated in [H] for unitary  $\pi^\sigma \times \pi^\vee$ , using a criterion characterizing distinction in terms of gamma factors. In [F3], Flicker defines a linear form on the space of  $\pi^\sigma \times \pi^\vee$  by a formal integral which would define the invariant linear form once the convergence is insured. Finally in [F-H], for  $n = 1$ , the convergence of this linear form is obtained for  $\pi^\sigma | \cdot |_K^s \times \pi^\vee | \cdot |_{\bar{K}}^{-s}$  and  $s$  of real part large enough, the conclusion follows from an analytic continuation argument.

We generalize this method here. The first part, which consists of showing that the representation  $\pi^\sigma | \cdot |_K^s \times \pi^\vee | \cdot |_{\bar{K}}^{-s}$  when  $Re(s)$  is large enough, is similar. The analytic continuation argument is a consequence of the theory of Igusa local zeta functions.

The first half of the first section is about notations and basic concepts used in the rest of the work. In the second half, we recall properties of the space of flat sections of an parabolically induced representation, and a theorem about rationality of Igusa zeta functions.

The second section is devoted to the proof of our main theorem 2.1, which asserts that the representation  $\pi^\sigma \times \pi^\vee$  is distinguished.

We end this introduction by recalling a conjecture about classification of distinguished generic representations:

**Conjecture.** *Let  $m$  be a positive integer, and  $\rho$  a generic representation of the group  $GL(m, K)$ , obtained by normalised parabolic induction of quasi-square-integrable representations  $\Delta_1, \dots, \Delta_t$ . It is distinguished if and only if there exists a reordering of the  $\Delta_i$ 's, and an integer  $r$  between 1 and  $t/2$ , such that we have  $\Delta_{i+1}^\sigma = \Delta_i^\vee$  for  $i = 1, 3, \dots, 2r - 1$ , and  $\Delta_i$  is distinguished for  $i > 2r$ .*

---

\*Nadir Matringe, IMJ, 2 place Jussieu, F-75251, Paris Cedex 05. Email: matringe@math.jussieu.fr.

We denote by  $\eta$  the nontrivial character of  $F^*$  trivial on the norms of  $K^*$ . According to Proposition 26 in [F1], Proposition 12 of [F2], Theorem 6 of [K], and Corollary 1.6 [A-K-T], our result reduces the proof of the conjecture to showing that representations of the form  $\Delta_1 \times \cdots \times \Delta_t$  with  $\Delta_{i+1}^\sigma = \Delta_i^\vee$  for  $i = 1, 3, \dots, 2r - 1$  for some  $r$  between 1 and  $t/2$ , and non isomorphic distinguished or  $\eta$ -distinguished  $\Delta_i$ 's for  $i > 2r$  are not distinguished whenever one of the  $\Delta_i$ 's is  $\eta$ -distinguished for  $i > 2r$ . According to [M3], the preceding conjecture implies the equality of the analytically defined Asai  $L$ -function and the Galois Asai  $L$ -function of a generic representation.

## 1 Notations and preliminary results

### 1.1 Notations

We denote by  $|\cdot|_K$  and  $|\cdot|_F$  the respective absolute values on  $K^*$ , by  $q_K$  and  $q_F$  the respective cardinalities of their residual field, and by  $R_K$  the valuation ring of  $K$ .

We call partition of a positive integer  $n$ , a family  $\bar{n} = (n_1, \dots, n_t)$  of positive integers (for a certain  $t$  in  $\mathbb{N} - \{0\}$ ), such that the sum  $n_1 + \cdots + n_t$  is equal to  $n$ . To such a partition, we associate a subgroup of  $GL(n, K)$  denoted by  $P_{\bar{n}}(K)$ , given by matrices of the form

$$\begin{pmatrix} g_1 & \star & \star & \star & \star \\ & g_2 & \star & \star & \star \\ & & \ddots & \star & \star \\ & & & g_{t-1} & \star \\ & & & & g_t \end{pmatrix},$$

with  $g_i$  in  $GL(n_i, K)$  for  $i$  between 1 and  $t$ . We call it the standard parabolic subgroup associated with the partition  $\bar{n}$ . We denote by  $N_{\bar{n}}(K)$  its unipotent radical subgroup, given by the matrices

$$\begin{pmatrix} I_{n_1} & \star & \star \\ & \ddots & \star \\ & & I_{n_t} \end{pmatrix}.$$

Finally we denote by  $P_n(K)$  the affine subgroup of  $GL(n, K)$  given by the matrices  $\begin{pmatrix} g & \star \\ & 1 \end{pmatrix}$ , with  $g$  in  $GL(n-1, K)$ .

Let  $X$  be a locally closed space of an  $l$ -group  $G$ , and  $H$  closed subgroup of  $G$ , with  $H.X \subset X$ . If  $V$  is a complex vector space, we denote by  $C^\infty(X, V)$  the space of smooth functions from  $X$  to  $V$ , and by  $C_c^\infty(X, V)$  the space of smooth functions with compact support from  $X$  to  $V$  (if one has  $V = \mathbb{C}$ , we simply denote it by  $C_c^\infty(X)$ ).

If  $\rho$  is a complex representation of  $H$  in  $V_\rho$ , we denote by  $C^\infty(H \backslash X, \rho, V_\rho)$  the space of functions  $f$  from  $X$  to  $V_\rho$ , fixed under the action by right translation of some compact open subgroup  $U_f$  of  $G$ , and which verify  $f(hx) = \rho(h)f(x)$  for  $h \in H$ , and  $x \in X$  (if  $\rho$  is a character, we denote this space by  $C^\infty(H \backslash X, \rho)$ ). We denote by  $C_c^\infty(H \backslash X, \rho, V_\rho)$  subspace of functions with support compact modulo  $H$  of  $C^\infty(H \backslash X, \rho, V_\rho)$ .

We denote by  $Ind_H^G(\rho)$  the representation by right translation of  $G$  in  $C^\infty(H \backslash G, \rho, V_\rho)$  and by  $ind_H^G(\rho)$  the representation by right translation of  $G$  in  $C_c^\infty(H \backslash G, \rho, V_\rho)$ . We denote by  $Ind'_H(\rho)$  the normalized induced representation  $Ind_H^G((\Delta_G/\Delta_H)^{1/2}\rho)$  and by  $ind'_H(\rho)$  the normalized induced representation  $ind_H^G((\Delta_G/\Delta_H)^{1/2}\rho)$ .

Let  $n$  be a positive integer, and  $\bar{n} = (n_1, \dots, n_t)$  be a partition of  $n$ , and suppose that we have a representation  $(\rho_i, V_i)$  of  $GL(n_i, K)$  for each  $i$  between 1 and  $t$ . Let  $\rho$  be the extension to  $P_{\bar{n}}(K)$  of the natural representation  $\rho_1 \otimes \cdots \otimes \rho_t$  of  $GL(n_1, K) \times \cdots \times GL(n_t, K)$ , by taking it trivial on  $N_{\bar{n}}(K)$ . We denote by  $\rho_1 \times \cdots \times \rho_t$  the representation  $Ind_{P_{\bar{n}}(K)}^{GL(nK)}(\rho)$ .

## 1.2 Flat sections

Let  $\pi$  be a representation of  $GL(n, K)$ , let  $(n_1, \dots, n_t)$  be a partition of  $n$ , and let  $\rho_1, \dots, \rho_t$  be smooth representations of  $GL(n_i, K)$ , such that  $\pi = \rho_1 \times \dots \times \rho_t$ . The map  $s = (s_1, \dots, s_t) \mapsto q_K^s = (q_K^{s_1}, \dots, q_K^{s_t})$  defines an isomorphism of varieties between  $\mathcal{D}_K^t = (\mathbb{C}/\frac{2i\pi}{\ln(q_K)}\mathbb{Z})^t$  and  $\mathbb{C}^{*t}$ . One can twist the representation  $\pi$  by an element  $s$  of  $\mathcal{D}_K^t$  into the representation  $\pi_s = \rho_1|_{q_K^{s_1}} \times \dots \times \rho_t|_{q_K^{s_t}}$ .

Denoting by  $(\rho, V_\rho)$  the tensor product of the  $\rho_i$ 's, the restrictions of functions of the space  $C_c^\infty(P_{\bar{n}}(K) \backslash GL(n, K), \Delta\rho_s, V_\rho)$  of  $\pi_s$  (where  $\Delta$  is  $(\Delta_{GL(n, K)}/\Delta_{P_{\bar{n}}(K)})^{1/2}$ ) to the maximal compact subgroup  $K_n = GL(n, R_K)$  of  $GL(n, K)$ , define the same space  $\mathcal{F}_\pi$ , which is called the space of flat sections of the series  $\pi_s$ . This restriction being injective, it identifies for each  $s$  in  $\mathcal{D}_K^t$ , the space of  $\pi_s$  with  $\mathcal{F}_\pi$ . Hence to every  $f$  in  $\mathcal{F}_\pi$  corresponds a unique function  $f_s$  in the space of  $\pi_s$ , and  $s \mapsto f_s$  can be seen as a section of the vector bundle over  $\mathcal{D}_K^t$  with fibers  $\pi_s$ , which is actually trivial, isomorphic to  $\mathcal{D}_K^t \times \mathcal{F}_\pi$ .

We still denote by  $\pi_s$  the representation of  $GL(n, K)$  on  $\mathcal{F}_\pi$ , defined by  $\pi_s(g)f = (\pi_s(g)f_s)|_{K_n}$  for  $f$  in  $\mathcal{F}_\pi$  and  $g$  in  $GL(n, K)$ . For fixed  $g$  in  $GL(n, K)$ , the map  $s \mapsto \pi_s(g)f$  from  $\mathcal{D}_K^t$  to  $\mathcal{F}_\pi$  is regular (i.e. belongs to the space  $\mathbb{C}[\mathcal{D}_K^t] \otimes \mathcal{F}_\pi$ ).

**Lemma 1.1.** (see [CS], Lemma 2.2) *Let the representation  $\pi$  be as above, and let  $f$  belong to the space  $\mathcal{F}_\pi$ . If we denote by  $N$  the unipotent radical of  $P_{\bar{n}}(K)$ , by  $N^-$  its image under transposition, and by  $w$  a Weyl element such that  $wNw^{-1}$  is equal to  $N^-$ .*

- *There exists a compact open subgroup  $U_f$  of  $GL(n, K)$ , such that for every  $s$  in  $\mathcal{D}_K^t$ , one has  $\pi_s(u)f_s = f_s$ .*
- *There exists a compact open subgroup  $N_f$  of  $N$ , such that for every  $s$  in  $\mathcal{D}_K^t$ , the function  $n \mapsto f_s(wn)$  for  $n$  in  $N$ , has its support in  $N_f$ .*

*Proof.* For the first point, we choose  $U_f$ , such that  $\pi_s(u)f = f$  for  $u$  in  $U_f$ .

For the second point, we take a countable exhaustive sequence of  $N$  of compact subgroups  $N_i$ . Denoting by  $\phi_{f_s}$  the function  $n \mapsto f_s(wn)$ , as  $wNw^{-1} = N^-$  has trivial intersection with  $P_{\bar{n}}(K)$ , we deduce that for every  $s$ , the function  $\phi_{f_s}$  has compact support in  $N$ . Hence denoting by  $\mathcal{D}_i$  the subset of  $\mathcal{D} = \mathcal{D}_K^t$ , consisting of  $s$  such that  $\phi_{f_s}$  has its support in  $N_i$ , we have  $\mathcal{D} = \cup_i \mathcal{D}_i$ . As the sets  $\mathcal{D}_i$  are closed, Baire's theorem then implies that one of the  $\mathcal{D}_i$ 's contains an open subset of  $\mathcal{D}$ . Now the condition “ $\phi_{f_s}$  has its support in  $N_i$ ”, is equivalent to the fact that for every  $n$  in  $N$ , outside  $N_i$ , the vector  $\phi_{f_s}(n) = (\pi_s(n)f)(w)$  is equal to 0. Calling  $\mathcal{D}_{i_0}$  one of the  $\mathcal{D}_i$ 's containing an open subset  $\mathcal{D}$ , then if  $n$  in  $N$  is outside of  $N_{i_0}$ , the regular function  $s \mapsto (\pi_s(n)f)(w)$  zero for  $s$  in  $\mathcal{D}_{i_0}$ , hence zero everywhere. From this we deduce that  $\mathcal{D}$  is equal to  $\mathcal{D}_{i_0}$ , and this concludes the proof.  $\square$

## 1.3 Igusa local zeta functions

Let  $m$  be a positive integer, and  $P$  be a non constant polynomial in  $F[X_1, \dots, X_m]$ . Then to such a polynomial  $P$ , one can associate the following functional on  $C_c^\infty(F^m)$ , depending on the complex variable  $s$ , formally defined for  $\phi$  in  $C_c^\infty(F^m)$  by the formula:

$$I_P(\phi, s) = \int_{F^m} |P(x)|_F^s \phi(x) dx,$$

for  $dx$  the Haar measure on  $F^m$ . It is clear that such an integral is absolutely convergent, hence well defined, for  $Re(s)$  positive. For fixed  $\phi$ , this defines a function of the complex parameter  $s$  in the variety  $\mathcal{D}_F = \mathbb{C}/\frac{2i\pi}{\ln(q_F)}\mathbb{Z} \simeq \mathbb{C}^*$ . Actually, we have the following fact, which is a consequence of a theorem of Igusa:

**Theorem 1.1.** ([I], theorem 8.2.1) *For every  $\phi$  in  $C_c^\infty(F^m)$ , the function  $s \mapsto I_P(\phi, s)$  is a rational function of  $\mathcal{D}_F$ .*

*Moreover, there is a polynomial  $Q$  in  $\mathbb{C}[X]$ , such that for every  $\phi$  in  $C_c^\infty(F^m)$ , the function  $s \mapsto Q(q_F^{-s})I_P(\phi, s)$  is regular (i.e. a polynomial in  $q_F^{-s}$ ) on  $\mathcal{D}_F$ .*

## 2 Distinction of representations $\pi^\sigma \times \pi^\vee$

We denote by  $G$  the group  $GL(2n, K)$ , by  $H$  its subgroup  $GL(2n, F)$ , by  $G'$  the group  $GL(n, K)$  and by  $M$  the group  $M_n(K)$ . We denote by  $P$  the group  $P_{(n,n)}(K)$ , and by  $N$  the group  $N_{(n,n)}(K)$ .

We denote by  $\bar{H}$  subgroup of  $G$  given by matrices of the form  $\begin{pmatrix} A & B \\ B^\sigma & A^\sigma \end{pmatrix}$ , and by  $\bar{T}$  the subgroup of  $\bar{H}$  of matrices  $\begin{pmatrix} A & 0 \\ 0 & A^\sigma \end{pmatrix}$ , with  $A$  in  $G'$ .

We call  $U$  be the matrix  $\begin{pmatrix} I_n & -\delta I_n \\ I_n & \delta I_n \end{pmatrix}$  of  $G$ , and  $W$  the matrix  $\begin{pmatrix} & -I_n \\ I_n & \end{pmatrix}$ . One has  $U^\sigma U^{-1} = W$  and the group  $H$  is equal to  $U^{-1}\bar{H}U$ .

**Lemma 2.1.** *The double class  $PUH$  is opened in  $G$ .*

*Proof.* Call  $S$  the space of matrices  $g$  in  $G$  verifying  $g^\sigma = g^{-1}$ , which is, from Proposition 3. of chapter 10 of [S], homeomorphic to the quotient space  $G/H$  by the map  $Q : g \mapsto g^\sigma g^{-1}$ . As the map  $Q$  sends  $U$  on  $W$ , the double class  $PUH$  corresponds to the open subset of matrices  $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$  in  $S$  such that  $\det(C) \neq 0$ , the conclusion follows.  $\square$

We prove the following integration formula.

**Lemma 2.2.** *There is a Haar measure  $d\dot{h}$  on the quotient space  $\bar{T}\backslash\bar{H}$ , and a Haar measure  $dB$  on  $M$ , such that for any measurable positive function  $\phi$  on the quotient space  $\bar{T}\backslash\bar{H}$ , then the integrals*

$$\int_{\bar{T}\backslash\bar{H}} \phi(\dot{h})d\dot{h}$$

and

$$\int_M \phi \left( \begin{pmatrix} I_n & B \\ B^\sigma & I_n \end{pmatrix} \right) \frac{dB}{|\det(I_n - BB^\sigma)|_K^n}$$

are equal.

*Proof.* It suffices to show this equality when  $\phi$  is positive, continuous with compact support in  $\bar{T}\backslash\bar{H}$ . We fix Haar measures  $dt$  on  $\bar{T}$  and  $dg$  on  $\bar{H}$ , such that  $d\dot{h}dt = dg$ . It is known that there exists some positive function  $\tilde{\phi}$  with compact support in  $\bar{H}$ , such that  $\phi = \tilde{\phi}^{\bar{T}}$ , which means that for any  $\dot{h}$  in  $\bar{H}$ , one has  $\phi(\dot{h}) = \int_{\bar{T}} \tilde{\phi}(tg)dt$ . One then has the relation

$$\int_{\bar{T}\backslash\bar{H}} \phi(\dot{h})d\dot{h} = \int_{\bar{H}} \tilde{\phi}(g)dg.$$

Now as  $\bar{H}$  is conjugate to  $H$ , there are Haar measures  $dA$  and  $dB$  on  $M$  such that  $dt$  is equal to  $d^*A = \frac{dA}{|\det(A)|_K^n}$ , and the Haar measure on  $\bar{H}$  is described by the relation

$$d \left( \begin{pmatrix} A & B \\ B^\sigma & A^\sigma \end{pmatrix} \right) = \frac{dAdB}{\left| \det \begin{pmatrix} A & B \\ B^\sigma & A^\sigma \end{pmatrix} \right|_F^{2n}} = \frac{dAdB}{\left| \det \begin{pmatrix} A & B \\ B^\sigma & A^\sigma \end{pmatrix} \right|_K^n}.$$

Hence we have

$$\begin{aligned} \int_{\bar{T}\backslash\bar{H}} \phi(\dot{h})d\dot{h} &= \int_{M \times M} \tilde{\phi} \left( \begin{pmatrix} A & B \\ B^\sigma & A^\sigma \end{pmatrix} \right) \frac{dAdB}{\left| \det \begin{pmatrix} A & B \\ B^\sigma & A^\sigma \end{pmatrix} \right|_K^n} \\ &= \int_{M \times M} \tilde{\phi} \left[ \begin{pmatrix} A & \\ & A^\sigma \end{pmatrix} \begin{pmatrix} I_n & A^{-1}B \\ (A^{-1}B)^\sigma & I_n \end{pmatrix} \right] \frac{dAdB}{|\det(A)|_K^{2n} |\det(I_n - A^{-1}B(A^{-1}B)^\sigma)|_K^n} \end{aligned}$$

as the complementary of  $G'$  is a negligible set of  $M$ .

This becomes after the change of variable  $B := A^{-1}B$  equal to

$$\int_{M \times M} \tilde{\phi} \left[ \begin{pmatrix} A & \\ & A^\sigma \end{pmatrix} \begin{pmatrix} I_n & B \\ B^\sigma & I_n \end{pmatrix} \right] \frac{dA}{|\det(A)|_K^n} \frac{dB}{|\det(I_n - BB^\sigma)|_K^n}$$

which is itself equal to

$$\int_{G' \times M} \tilde{\phi} \left[ \begin{pmatrix} A & \\ & A^\sigma \end{pmatrix} \begin{pmatrix} I_n & B \\ B^\sigma & I_n \end{pmatrix} \right] d^*A \frac{dB}{|\det(I_n - BB^\sigma)|_K^n}.$$

The conclusion follows from the fact that  $\tilde{\phi}^{\bar{T}}$  is equal to  $\phi$ . □

**Theorem 2.1.** *Let  $n$  be a positive integer, and let  $\pi$  be a smooth irreducible representation of  $G'$ . Then the representation  $\pi^\sigma \times \pi^\vee$  is a distinguished representation of  $G$ .*

*Proof.* We will first show that the representation  $\Pi_s = \pi^\sigma | \cdot |^s \times \pi^\vee | \cdot |^{-s}$  of  $G$  is distinguished for a complex number  $s$  of real part large enough. Then we show that this property still holds for  $s$  equal to 0.

Step 1.

The representation  $\pi^\sigma | \cdot |^s \otimes \pi^\vee | \cdot |^{-s}$  of  $G' \times G'$  is distinguished by  $\bar{T}$ . Call  $L$  a nonzero  $\bar{T}$ -invariant linear form on the space of  $\pi^\sigma | \cdot |^s \otimes \pi^\vee | \cdot |^{-s}$ .

We denote by  $\rho_s$  the representation  $P$ , which is the extension of  $\pi^\sigma | \cdot |^s \otimes \pi^\vee | \cdot |^{-s}$  to  $P$  by the trivial representation of  $N_{(n,n)}(K)$ . Let  $f_s$  belong to the space  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$  of  $\pi^\sigma | \cdot |^s \times \pi^\vee | \cdot |^{-s}$ , corresponding to  $f$  in the space of flat sections  $\mathcal{F}_\Pi$ , where  $\Pi = \Pi_0 = \pi^\sigma \times \pi^\vee$ . If  $A, B$  and  $g$  belong respectively to  $G', G'$  and  $G$ , one has the following relation:

$$f_s \left[ \begin{pmatrix} A & \star \\ 0 & B \end{pmatrix} g \right] = \frac{|\det(A)|_K^{n/2+s}}{|\det(B)|_K^{n/2+s}} \pi^\sigma(A) \otimes \pi^\vee(B) f(g).$$

In particular for  $f_s$  in  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$ , corresponding to  $f$ , the restriction to  $\bar{H}$  of the function  $L_{f_s} : g \mapsto L(f_s(g))$  belongs to the space  $C^\infty(\bar{T} \backslash \bar{H})$ , but its support modulo  $\bar{T}$  is generally not compact, we will show later that the space of functions obtained this way contains  $C_c^\infty(\bar{T} \backslash \bar{H})$  as a proper subspace. We must show that for  $s$  of real part large enough, the integral  $\int_{\bar{T} \backslash \bar{H}} |L_{f_s}(\dot{h})| d\dot{h}$  converges.

According to lemma 2.2, this integral is equal to

$$\int_M |L_{f_s}| \left( \begin{pmatrix} I_n & B \\ B^\sigma & I_n \end{pmatrix} \right) \frac{dB}{|\det(I_n - BB^\sigma)|_K^n}.$$

As before, we can suppose that  $B$  belongs to  $G'$ , hence the following decomposition holds

$$\begin{pmatrix} I_n & B \\ B^\sigma & I_n \end{pmatrix} = \begin{pmatrix} (I_n - BB^\sigma)B^{-\sigma} & I_n \\ B^\sigma & I_n \end{pmatrix} \begin{pmatrix} & -I_n \\ I_n & \end{pmatrix} \begin{pmatrix} I_n & B^{-\sigma} \\ & I_n \end{pmatrix}.$$

Denoting by  $\Phi_{f_s}$  the function

$$L(\phi_{f_s}) : B \mapsto L_{f_s} \left[ \begin{pmatrix} & -I_n \\ I_n & \end{pmatrix} \begin{pmatrix} I_n & B \\ & I_n \end{pmatrix} \right],$$

the following equalities follow:

$$\begin{aligned} \int_{\bar{T} \backslash \bar{H}} |L_{f_s}(\dot{h})| d\dot{h} &= \int_M \left( \frac{|\det(I_n - BB^\sigma)|_K}{|\det(B)|_K^2} \right)^{s+n/2} |\Phi_{f_s}(B^{-\sigma})| \frac{dB}{|\det(I_n - BB^\sigma)|_K^n} \\ &= \int_{G'} \left( \frac{|\det(I_n - BB^\sigma)|_K}{|\det(B)|_K^2} \right)^{s+n/2} |\Phi_{f_s}(B^{-\sigma})| \frac{|\det(B)|_K^n d^*B}{|\det(I_n - BB^\sigma)|_K^n} \\ &= \int_{G'} (|\det(I_n - C^{-\sigma}C^{-1})|_K |\det(C)|_K^2)^{s+n/2} |\Phi_{f_s}(C)| \frac{d^*C}{|\det(C)|_K^n |\det(I_n - C^{-\sigma}C^{-1})|_K^n} \\ &= \int_M |\det(CC^\sigma - I_n)|_K^{s-n/2} |\Phi_{f_s}(C)| dC \end{aligned}$$

Recalling that the map  $f_s$  belongs to  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$ , and as the map  $(P, N') \in P \times N \mapsto PWN' \in G$  is a homeomorphism, we deduce that

$$C \mapsto f_s \left[ \begin{pmatrix} & -I_n \\ I_n & \end{pmatrix} \begin{pmatrix} I_n & C \\ & I_n \end{pmatrix} \right]$$

has compact support in  $M$ , hence so does  $\Phi_{f_s}$ .

This eventually implies that  $\int_{\bar{T} \backslash \bar{H}} L(f_s(\dot{h})) d\dot{h}$  is convergent as soon as  $s$  has real part greater than  $n/2$ .

### Step 2.

Suppose that the complex number  $s$  has real part greater than  $n/2$ . We are going to show that the linear form  $\Lambda : f_s \mapsto \int_{\bar{T} \backslash \bar{H}} L_{f_s}(\dot{h}) d\dot{h}$  is nonzero. More precisely we are going to show that the space of functions  $L(f)$  on  $\bar{T} \backslash \bar{H}$  for  $f$  in  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$ , contain  $C_c^\infty(\bar{T} \backslash \bar{H})$ .

According to Lemma 2.1, the double class  $PUH$  is opened in  $G$ , hence the extension by zero outside  $PUH$  gives an injection of the space  $C_c^\infty(P \backslash PUH, \Delta_P^{-1/2} \rho_s)$  into the space  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$ . But the right translation by  $U$ , which is a vector space automorphism of  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$ , sends  $C_c^\infty(P \backslash PUH, \Delta_P^{-1/2} \rho_s)$  onto  $C_c^\infty(P \backslash P\bar{H}, \Delta_P^{-1/2} \rho_s)$ , hence  $C_c^\infty(P \backslash P\bar{H}, \Delta_P^{-1/2} \rho_s)$  is a subspace of  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$ .

Now restriction to  $\bar{H}$  defines an isomorphism between  $C_c^\infty(P \backslash P\bar{H}, \Delta_P^{-1/2} \rho_s)$  and  $C_c^\infty(\bar{T} \backslash \bar{H}, \rho_s)$  because  $\Delta_P$  has trivial restriction to the unimodular group  $\bar{T}$ . But then the map  $f \mapsto L(f)$  defines a morphism of  $\bar{H}$ -modules from  $C_c^\infty(\bar{T} \backslash \bar{H}, \rho_s)$  to  $C_c^\infty(\bar{T} \backslash \bar{H})$ , which is surjective because of the commutativity of the following diagram,

$$\begin{array}{ccc} C_c^\infty(\bar{H}) \otimes V_{\rho_s} & \xrightarrow{Id \otimes L} & C_c^\infty(\bar{H}) \\ \downarrow & & \downarrow \\ C_c^\infty(\bar{T} \backslash \bar{H}, \rho_s) & \longrightarrow & C_c^\infty(\bar{T} \backslash \bar{H}) \end{array},$$

where the vertical arrows defined in Lemma 2.9 of [M1] and the upper arrow are surjective.

We thus proved that space of restrictions to  $\bar{H}$  of functions of  $L(f)$ , for  $f$  in  $C_c^\infty(P \backslash G, \Delta_P^{-1/2} \rho_s)$ , contain  $C_c^\infty(\bar{T} \backslash \bar{H})$ , hence  $\Lambda$  is nonzero and the representation  $\pi^\sigma | \begin{smallmatrix} s & \\ & \pi^\vee \end{smallmatrix} | \begin{smallmatrix} s & \\ & \pi^\vee \end{smallmatrix}$  is distinguished for  $Re(s) \geq n/2$ .

### Step 3.

We are going to show that  $\Pi = \pi^\sigma \times \pi^\vee$  is distinguished. Let  $f$  belong to  $\mathcal{F}_\Pi$ , it is a consequence of Lemma 1.1 that there exists two compact open subgroups  $U_1 \subset U_2$  of  $(M_n(K), +)$ , such that for every  $s$  in  $\mathcal{D}_K$ , the function  $\Phi_{f_s}$  is invariant under addition by elements of  $U_1$ , and has support in  $U_2$ . Denoting by  $P$  the polynomial function on the finite dimension  $F$ -vector space  $M_n(K)$  given by  $C \mapsto \det(CC^\sigma - I_n)$ , we recall from the two first steps, using notations of subsection 1.3, that for  $Re(s) \geq n/2$ , the linear form  $f \mapsto I_P(\Phi_{f_s}, 2s - n)$  is nonzero and  $G_n(F)$ -invariant. For such  $s$ , one has  $I_P(\Phi_{f_s}, 2s - n) = \int_{U_2} |P(C)|_F^{2s-n} \Phi_{f_s}(C) dC = \sum_{C_i \in U_2/U_1} \Phi_{f_s}(C_i) \int_{C_i+U_1} |P(C)|_F^{2s-n} dC = \sum_{C_i \in U_2/U_1} \Phi_{f_s}(C_i) I_P(1_{C_i+U_1}, 2s - n)$ .

In particular, from subsection 1.2, we know that the functions  $s \mapsto \Phi_{f_s}(C_i)$  are polynomial in  $q_F^{-s}$ , and according to theorem 1.1, there is  $Q$  in  $\mathbb{C}[X]$ , such that for every  $f \in \mathcal{F}_\Pi$ , the function  $Q(q_F^{-s}) I_P(\Phi_{f_s}, 2s - n)$  is regular on  $\mathcal{D}_F$ . As the function  $s \mapsto I_P(\Phi_{f_s}, 2s - n)$  is meromorphic, one has for every  $h$  in  $GL(n, F)$  the equality  $I_P(\Phi_{\pi_s(h)f_s}, 2s - n) = I_P(\Phi_{f_s}, 2s - n)$  (because the equality holds for  $Re(s) \geq n/2$ ). Moreover one can choose an integer  $d$  such that the function  $(q_F^{-s} - 1)^d I_P(\Phi_{f_s}, 2s - n)$  is regular at zero for every  $f \in \mathcal{F}_\Pi$ , and doesn't vanish at  $s = 0$  for a good choice of  $f$ . Hence the linear form  $f_0 \mapsto \lim_{s \rightarrow 0} (q_F^{-s} - 1)^d I_P(\Phi_{f_s}, 2s - n)$  defines a nonzero  $G_n(F)$ -invariant linear form on the space of  $\Pi$ . We thus showed that the representation  $\Pi$  is distinguished.  $\square$

## References

- [A-K-T] U. K. Anandavardhanan, A.C. Kable and R.Tandon, *Distinguished representations and poles of twisted tensor L-functions*, Proc. Amer. Math. Soc., **132** (2004), No. 10, 2875-2883.
- [Ber] J. N. Bernstein, *P-invariant distributions on  $GL(N)$  and the classification of unitary representations of  $GL(N)$  (non-archimedean case)*, Lecture Notes in Math., vol 1041, Springer-Verlag, Berlin, (1983), 50-102.
- [CS] W. Casselman and J. Shalika, *The unramified principal series of  $p$ -adic groups. II. The Whittaker function*. Compositio Mathematica, 41 no. 2 (1980), p. 207-231
- [F1] Y. Flicker, *Twisted tensors and Euler products*, Bull. Soc. Math. France, **116** no.3, (1988), 295-313.
- [F2] Y. Flicker, *On distinguished representations*, J. Reine Angew. Math., **418** (1991), 139-172.
- [F3] Y. Flicker, *Distinguished representations and a Fourier summation formula*, Bull. Soc. Math. France, **120** (1992), 413-465.
- [F4] Y. Flicker, Appendix of *On zeroes of the twisted tensor L-function*, Math. Ann., **297**, (1993), 199-219.
- [F-H] Y. Flicker and J. Hakim, *Quaternionic distinguished representations*, Amer. J. Math., **116**, (1994), 683-736.
- [H] J. Hakim, *Distinguished  $p$ -adic Representations*, Duke Math. J., **62** (1991), 1-22.
- [I] Igusa, Jun-Ichi, *An Introduction to the Theory of Local Zeta Functions*, AMS/IP Studies in Advanced Mathematics, v. 14, 2000.
- [K] A. C. Kable, *Asai L-functions and Jacquet's conjecture*, Amer. J. Math., **126**, (2004), 789-820.
- [M1] N. Matringe, *Distinguished principal series representations for  $GL_n$  over a  $p$ -adic field*, Pacific J.Math., Vol. 239, No. 1, Jan 2009.
- [M2] N. Matringe, *Distinguished representations and exceptional poles of the Asai-L-function*, preprint, <http://hal.archives-ouvertes.fr/hal-00299528/fr/>, arXiv:0807.2748
- [M3] N. Matringe, *Conjectures about distinction and Asai L-functions of generic representations of general linear groups over local fields*, preprint, <http://arxiv.org/abs/0811.1410v2>
- [S] J.-P. Serre, *Corps locaux*, 2nd ed., Publications de l'Universit de Nancago 8, Hermann, Paris, 1968.
- [Z] A.V. Zelevinsky, *induced representations of reductive  $p$ -adic groups II*, Ann.Sc.E.N.S., 1980.