



**HAL**  
open science

## **14-3-3 regulates life span by both DAF-16-dependent and independent mechanisms in *Caenorhabditis elegans***

Caroline Araiz, Marie-Thérèse Château, Simon Galas

### ► To cite this version:

Caroline Araiz, Marie-Thérèse Château, Simon Galas. 14-3-3 regulates life span by both DAF-16-dependent and independent mechanisms in *Caenorhabditis elegans*. *Experimental Gerontology*, 2008, 43 (6), pp.505. 10.1016/j.exger.2008.03.001 . hal-00345888v1

**HAL Id: hal-00345888**

**<https://hal.science/hal-00345888v1>**

Submitted on 9 Jul 2010 (v1), last revised 12 Dec 2008 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Accepted Manuscript

14-3-3 regulates life span by both DAF-16-dependent and independent mechanisms in *Caenorhabditis elegans*

Caroline Araiz, Marie-Thérèse Château, Simon Galas


PII: S0531-5565(08)00083-1  
DOI: [10.1016/j.exger.2008.03.001](https://doi.org/10.1016/j.exger.2008.03.001)  
Reference: EXG 8455

To appear in: *Experimental Gerontology*

Received Date: 4 May 2007  
Revised Date: 13 February 2008  
Accepted Date: 4 March 2008

Please cite this article as: Araiz, C., Château, M-T., Galas, S., 14-3-3 regulates life span by both DAF-16-dependent and independent mechanisms in *Caenorhabditis elegans*, *Experimental Gerontology* (2008), doi: [10.1016/j.exger.2008.03.001](https://doi.org/10.1016/j.exger.2008.03.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**14-3-3 regulates life span by both DAF-16-dependent and independent mechanisms in *Caenorhabditis elegans*.**

Caroline Araiz<sup>1</sup>, Marie-Thérèse Château<sup>1,2</sup> and Simon Galas<sup>1,2</sup>

<sup>1</sup>CRBM- CNRS, 1919 route de Mende, 34293 Montpellier cedex 5, France;

<sup>2</sup>Université Montpellier1, Faculté de Pharmacie, 15 Avenue Charles Flahault - B.P. 14491, 34093 Montpellier cedex 5.

Keywords: [*C. elegans*, ageing, Insulin/IGF-1/DAF-2, FOXO/DAF-16, 14-3-3/*ftt1/ftt2*]

Running title: Ageing and stress regulation by 14-3-3 in *C. elegans*

---

Correspondence

Prof. Simon Galas, CRBM-CNRS, 1919 route de Mende, F34293 Montpellier cedex 5, France. Telephone +33 (0)4 67 61 34 32. Fax: +33 (0) 4 67 52 15 59. E-mail: [simon.galas@crbm.cnrs.fr](mailto:simon.galas@crbm.cnrs.fr)

**Abstract**

*Caenorhabditis elegans* life span, stress resistance and metabolism are regulated by the Insulin/IGF-1/DAF-2/DAF-16 pathway. DAF-16, a member of FOXO/Forkhead transcription factor family can be targeted by 14-3-3 proteins to promote stress resistance. We have identified a 14-3-3 *C. elegans* homolog which promotes life span by both DAF-2-dependent and -independent mechanisms and by an unexpected DAF-16 -independent mechanism. Our results demonstrate that *C. elegans* 14-3-3 proteins modulate stress responsive genes throughout adulthood. In conclusion, 14-3-3 can be considered as an acute stress-responsive regulator as well as a sustained modulator of the Insulin/IGF-1/DAF-2/DAF-16 regulatory pathway in promoting life expectancy of growing old worms.

ACCEPTED MANUSCRIPT

## Introduction

Cell regulatory pathways, including signal transduction, apoptosis, stress response and transformation can be targeted by one or several 14-3-3 gene isoform products in organisms ranging from yeast to human (Van Hemert et al., 2001). The 14-3-3 family members are discrete binding proteins with no intrinsic enzymatic activity. However, the 14-3-3 proteins can act as enzymatic regulators as well as localization anchors or adaptator molecules with scaffold activity that can, in turn, stimulate various protein-protein interactions (Van Hemert et al., 2001). The *Caenorhabditis elegans* (*C. elegans*) genome encodes two 14-3-3 genes: *ftt-1* and *ftt-2* (*ftt* for fourteen-three-three; Wang and Shakes, 1994). While a reported simultaneous expression of *ftt-1* and *ftt-2* during early embryogenesis may allow cooperative effects, the divergent expression patterns during pre-reproductive or adulthood stages argue for different respective biological functions (Wang and Shakes, 1997). Thus, while the germline enriched *ftt-1* gene product (Wang and Shakes, 1997) - also known as *par-5* - is involved in the asymmetric cortical localization of other *par* gene products that can, in turn, influence the general asymmetry determinism in early embryos (Morton et al., 2002), the *ftt-2* gene expression pattern is largely non overlapping with the former because of a reported somatic expression that also takes place at later developmental stages (Wang and Shakes, 1997). The expression pattern of a specific reporter transgene for *ftt-2* and the *ftt-1* further supports this view (Wang et al., 2006), altogether these reports make the *ftt-2* gene product a *bona fide* candidate for a transduction pathway regulator during post-embryonic or post-developmental stages of *C. elegans*.

Mammalian FOXO homologs are well known targets of 14-3-3 proteins (Van Hemert et al., 2001; Brunet et al. 1999) but it has been shown that active *C. elegans* DAF-16 transcription factor, a FOXO family transcription factor (Lin et al., 1997; Ogg et al., 1997; Kenyon 2005), can be inhibited by 14-3-3 -dependent and -independent pathways in cell culture (Cahill et al. 2001). The conserved DAF-2/IGF-1 signalling pathway (Kenyon 2005) is believed to regulate life span through inhibition of the *daf-16* gene product and it has been proposed that differential signalling may originate at the DAF-2/IGF-1 receptor to regulate *daf-16* gene product activity leading to the emergence of different phenotypic *daf-2* allele class mutants (Gems et al., 1998; Nanji et al., 2005). Moreover, the IGF-1 receptor itself, as well as downstream transducers such as 3-Phosphoinositide-dependent kinase-1 (PDK1) or IREs (insulin receptor substrates), are 14-3-3-reported targets (Van Hemert et al. 2001; Craparo et al., 1997; Sato et al., 2001). Recent reports demonstrated a positive worm life span modulation by 14-3-3 through a cooperative interaction mechanism involving both SIR-2.1 and FOXO/DAF-16: a mechanism fully suppressible by a *daf-16* loss-of-function mutation or RNAi inactivation (Wang et al., 2006; Berdichevsky et al. 2006; Li et al., 2007). This new regulatory pathway has been **linked** to a stress-dependent mechanism acting in parallel to the DAF-2/IGF-1 signalling pathway to activate FOXO/DAF-16 and promote life span (Berdichevsky and Guarente 2006).

This report demonstrates that *C. elegans* 14-3-3 gene products can promote life span by both FOXO/*daf-16* dependent and independent mechanisms. Furthermore we also reveal a specific *daf-2* class mutant allele requirement for 14-3-3 genes to promote life span, together with different effects on physiological traits including both thermal and oxidative stress resistance, development, dauer arrest and fat storage. Finally, our work extends *C. elegans* 14-3-3 involvements in biological pathways functioning with acute stress response and sustained life span regulation.

## Materials and Methods

## Strains and worm cultures

Worms were cultured and maintained as described previously (Brenner, 1974). We used the following strains: Wild type Bristol (N2), GR1307 *daf-16(mgDf50)*, CB13710 *daf-2(e1370)*, JT709 *pdk-1(sa709)*, JT9609 *pdk-1(sa680)*, GR1318 *pdk-1(mg142)*, CF1330 *muEx128* [pKL79 (*daf-16a::GFP*)+pRF4 *rol-6 (su1006)*], CL2070 *dvIs70*, CF1553 *mulS84* [pAD76(*sod-3::GFP*)], CL2166 *dvIs19* [pAF15(*gst-4::GFP::NLS*)] which were all provided by the Caenorhabditis Genetics Center (funded by the NIH National Center for Research Resources). DR1569 *daf-2(e1365)* was kindly provided by Donald L. Riddle, University of British Columbia, Canada. Unless specified, worms were cultured at 20°C or 15°C for JT709 *pdk-1(sa709)*.

## RNAi constructs

The *ftt-1* RNAi construct was obtained by subcloning a 709bp HindIII/BglIII fragment from the pCP5 vector into the pPD129.36 (L4440) vector (a generous gift of Johnathan Ewbank, CNRS, Marseille, France) while *ftt-2* RNAi construct was generated by subcloning a 531bp HindIII fragment from the pSE16 vector into the pPD129.36 (L4440) vector. Both pCP5 and pSE16 *ftt* cDNA clones were a generous gift from Diane Shakes (College of William and Mary, Virginia, USA). After transformation with either pPD129.36, *ftt-1* or *ftt-2* RNAi construct, HT115 (DE3) strain bacteria were grown in LB medium with both Ampicillin (50 mg/ml) and Tetracyclin (12,5 mg/ml) during eight hours, then seeded on NGM plates containing FUDR (98,5mg/ml), Ampicillin (50 mg/ml) and IPTG (1mM) and induced overnight.

## RNAi feeding and lifespan assays

Adult stage RNAi: L4 larvae fed with RNAi producing bacteria were allowed to grow on NGM plates at 20 °C until they died. To avoid offspring during experiments we used plates with FUDR (Mitchell et al., 1979). Dead worms were scored every day until they were all dead. Vitality was assessed by the animal response to one or several gentle nose touch. Worms that died by scrolling out of agar plates were scored as censored data. XLSTAT-life statistical software (Addinsoft, New York, NY, USA) was used to plot survival data by the Kaplan Meier method and differences between survival curves were calculated using the Log Rank test with 95% confidence.

Developmental stage RNAi and dauer assay : Five to ten L4/young adult hermaphrodites for each strain were permitted to egg-lay for two hours on plates seeded with transformed HT115 at 22,5°C. The progeny was then allowed to grow on these same conditions. Delay to reach L4 stage was checked comparing to N2 on control RNAi conditions. All the worms growing to L4/young adult stage were transferred onto NGM plates with only FUDR and E. Coli OP50 strain and scored for life span trials, whereas dauer larvae were scored until around 72 hours and transferred onto fresh RNAi plates and observed until they left this stage and scored for dauer exit. For statistical analysis of the life span data curves we considered only the trials that gave a high percentage of dauer in parallel.

## Real-time PCR

Synchronised N2 adult worms were transferred on RNAi plates at 20°C during one day and then collected by washing with M9 buffer and pelleted. Total RNA isolation was done using

TRIzol Reagent (Invitrogen) and RNA quantification was assessed by both gel electrophoresis and UV spectrophotometric analysis. cDNAs were synthesized with random hexamers by using Ready-To-Go™ You-Prime First-Strand Beads kit (GE Healthcare) according to the manufacturer's protocol. Real-time PCR reactions were performed using LightCycler Fast Start DNA Master<sup>plus</sup> SYBR Green I kit in capillaries. Triplicates for each sample were included for one single reaction.

The real-time PCR primers for TBA2 are: Forward primer: 5'-CCTCCTCCGAATGAATGAAA-3'; Reverse primer: 5'TCCGATACTGGAAACGGAAG-3'. Primers for *ftt-1* are: Forward primer: 5'-GTTTCAGAAGACTTGCTCAC-3'; Reverse primer: 5'-CTCGGTCACCTTCTTCATGGC-3'. Primers for *ftt-2* are: Forward primer: 5'-CAGTGACTCATCCACTTCAAGC-3'; Reverse primer: 5'-AACGCTCAGCCTGCTCGGCAAG-3'. All the PCR products were checked by sequencing. To avoid Light-Cycler cross amplification of RNA sequence included in RNAi-fed worms, both *ftt-1* and *ftt-2* specific forward primers were designed to hybridize the 5'-UTR region not included in the RNAi-targeted sequences. The corresponding reverse primers were designed to hybridize the boundary of the first and the second exons in order to prevent cross-amplification of unspliced RNAs. *tba-2* was used as the internal control and the RNA level of each gene of interest was normalized to the *tba-2* level for comparison. Real-time PCR reactions were performed three times using independent RNAi worms and RNA isolations and results were treated as described (Pfaffl, 2001). All of quantified PCR products in this study were previously sequenced for molecular identity verification.

### **Nile red fat staining and handling of worms for transgene observations**

We used Nile Red (9-diethylamino-5H-benzo[a]phenoxa-]phenoxazine-5-one) obtained from Molecular Probes (N-1142) dissolved with acetone and mixed with transformed HT115 DE3 *E. coli* strain, at a final concentration of 0,5µg/ml. Five L4/young adult hermaphrodites of each strain were allowed to egg-lay for two hours at 22,5°C under RNAi conditions and then removed. The progeny was left on these plates until it reached L4/young adult stage. Nile red treatment resulted in uptake and incorporation of the dye into lipid droplets in intestinal cells, the main site of worm fat storage.

L4/young adults from either CF1553, CL2070 or CL2166 strains were transferred onto NGM plates prepared for RNAi feeding induction with FUDR. After four and eighteen days of treatment, sample worms were collected and observed under epifluorescence microscopy.

### **Oxidative and heat shock stress resistance assays**

After one week of RNAi treatment at 20°C adult worms, were transferred onto 24-well plates (about 50 worms per well), with 10 mM of Menadione (Sigma Aldrich) and HT115 bacterias mixed with S complete medium to maintain the RNAi treatment during the assay. Complete S medium mixed with HT115 bacteria was used as a control. Experiments were carried out for 7 days at 20°C. We scored the number of alive and dead worms each day, using 1µM SYTOX Green Nucleic Acid Stain (Molecular Probes) to discriminate dead worms from alive ones. Survival histograms were analysed with XLSTAT software (Addinsoft, New York, NY, USA). For heat shock stress induction, adult worms fed on RNAi seeded plates at 20°C for one week were placed on a 37°C water bath for 2,5 hours then returned to 20°C and scored for dead worms.

### **Microscopy observation and pixel quantification**

Microscopy observations were performed on a Zeiss AxioImager.Z1 with a Photometrics CoolSnap HQ2 camera at either 40X or 100X magnification with Zeiss Plan Neofluar-1.3 or Zeiss Apochromat-1.4 objectives respectively. Nile red staining was assessed with a Zeiss FS00, BP530-585, FT600, LP615 red filter while GFP signal was assessed with a Zeiss GFP FS38HE, BP470/40, FT495, BPJ25/50 GFP filter. Pixel quantification was performed with Metamorph v7.1 software (Molecular Devices, USA) while data analysis by a student t-test was done using the XLSTAT software (Addinsoft, New York, NY, USA).

ACCEPTED MANUSCRIPT



## Experimental Results

### ***ftt-2* gene but not *ftt-1* gene modulates life span in *Daf-16* loss-of-function genetic background.**

We used RNA-mediated gene interference (RNAi) to reduce the expression of either *ftt-1* or *ftt-2* *C. elegans* genes in wild type worms (hereafter denoted as N2). We verified the the specificity of our RNAi constructs by quantifying *ftt-1* and *ftt-2* RNA level in N2 RNAi worms by real-time PCR (see materials and methods). As shown in Fig. 1, we reported both significant and specific gene knock down of both *ftt-1* or *ftt-2* mRNA expression in worms fed with the corresponding RNAi bacteria. Moreover, similar to previous reports (Morton et al., 2002; Wang et al., 2006; Berdichevsky et al., 2006) using *par-5* mutants or *ftt-1* RNAi, we observed almost 100% dead embryos in the progeny of adults with *ftt-1* RNAi.

In contrast, the *ftt-2* RNAi treatment didn't reveal any embryonic lethality. As previously reported (Berdichevsky et al., 2006), we observed that *ftt-1* and *ftt-2* RNAi treatment induced DAF-16::GFP nuclear localization (data not shown). Additionally, *ftt-2* RNAi during larval stage induced a low percentage of "bagging" phenotype (15% for wild type worms), as a result of internal hatching of the progeny and a small fraction (8% for wild type worms) of protruding vulvae phenotype in adults. No "bagging" phenotype was observed with *ftt-2* RNAi during adulthood. Therefore, the gene specificity of the RNAi treatment correlated with phenotypic traits.

We then assessed for 14-3-3-dependent effects on worm life span by subjecting N2 adults to either *ftt-1* or *ftt-2* RNAi. As shown in Fig. 2.A and Table 1, N2 worms showed a significant life span shortening while *ftt-1* RNAi yielded an intermediate life span shortening compared to *ftt-2* RNAi. This result gives an indication that N2 worms are more prone to *ftt-2* for life span modulation than to *ftt-1* when maintained in stress-free conditions since all of these experiments were conducted at 20°C.

The DAF-16 transcription factor is believed to be situated at the major crossroads of the various regulatory pathways that modulates life span and stress resistance in *C. elegans* (Lin et al., 1997; Ogg et al., 1997; Kenyon 2005; Dillin et al., 2002). Recent reports (Wang et al., 2006; Berdichevsky et al., 2006; Li et al., 2007) demonstrated new 14-3-3 regulations extending stress resistance and life span in a DAF-16 dependent manner. However, active DAF-16 inhibition by 14-3-3 dependent and independent pathways in cell culture has been reported (Cahill et al., 2001). To clarify the relationship between 14-3-3 and DAF-16-dependent life span modulation mechanisms, we assessed life span modulation in a *daf-16(mgDf50)* deletion mutant allele treated with *ftt* RNAi. As shown in Fig. 2.B and Table 1, we noticed a highly significant reduction in life span upon *ftt-2* RNAi. In contrast *ftt-1* RNAi did not induce any significant life span reduction. This unexpected result demonstrates, for the first time, that DAF-16-dependent and -independent mechanisms can be targeted by 14-3-3 to promote life span. Moreover, because life span modulation by *ftt-1* is fully suppressible by a *daf-16* loss-of-function mutation, we propose that *daf-16* is the only target of *ftt-1* for life span regulation in worms while *ftt-2* is able to modulate life span by an as yet unknown *daf-16*-independent mechanism together with a *daf-16*-dependent process.

Previous studies have already reported the RNAi potency to specifically target a given *ftt* gene product in worm (Morton et al., 2002; Berdichevsky et al., 2006; Wang et al., 2006; Li et al., 2007). Therefore, the data presented here strongly support that RNAi constructs can specifically target the corresponding *ftt-1* or *ftt-2* gene products without any cross-phenotypic consequences.

### **Worm life span modulation by 14-3-3 differs depending on *daf-2* mutant class allele background.**

Major biological outputs of the DAF-2/IGF-1 pathway are believed to elicit FOXO/DAF-16 activity modulation since strong loss-of-function mutations in the *daf-16* gene can suppress the induced phenotypes by point mutation in the DAF-2/IGF-1 receptor tyrosine kinase (Ogg et al., 1997; Lin et al., 2001). We therefore assessed 14-3-3 life span modulation effects in the *daf-2(e1365)* and *daf-2(e1370)* reduction-of-function mutant because they belong respectively to *daf-2* class 1 and class 2 phenotypic mutants (Gems et al., 1998). Amino-acid substitution (A580T) in the ligand-binding domain of *daf-2(e1365)* and in the tyrosin kinase receptor domain (P1465S) of *daf-2(e1370)* have been proposed to explain the phenotypic difference between both *daf-2* class mutants (Gems et al., 1998; Nanji et al., 2005). Different *daf-2* class mutants and corresponding phenotypic traits are believed to result from the mobilisation of different signalling intermediates that originate at the DAF-2/IGF-1 receptor and result in a subset of common and different downstream unknown complex processes, that can in turn, converge to regulate the *daf-16* gene product (Gems et al., 1998; Nanji et al., 2005). We expected these experiments to be highly informative with respect to 14-3-3 dependent regulations on worm life span by providing new insights into 14-3-3-dependent phenotypic traits expressing or not a *daf-2* class mutant requirement to be revealed. As shown in Fig. 2.C,D and Table 1, both *ftt-1* and *ftt-2* RNAi treatments induced a significant life span reduction when applied to both *daf-2* class mutants. However, as attested by the 50<sup>th</sup> percentile (see Table 1), the strongest life span reduction was observed upon *ftt-2* RNAi treatment on both *daf-2* class mutants. Moreover, relevant statistical difference between both *ftt-1* and *ftt-2* RNAi treatment was only evident on *daf-2(e1370)* and we were also surprised to note a return to a wild type mean life span of the *daf-2(e1370)* worms (compare mean life span of experiments code A and M on Table 1) upon *ftt-2* RNAi treatment only. Altogether, these observations indicate that *daf-2(e1370)* class 2 mutant is more prone to *ftt-2* -dependent life span modulation than *daf-2(e1365)* class 1 mutant.

It has been reported that DAF-16 activity regulation mainly operates by the 3-Phosphoinositide-dependent kinase-1 /AGE-1 signalling pathway in *daf-2(e1370)* class 2 rather than in *daf-2(e1365)* class 1 mutants (Gems et al., 1998; Nanji et al., 2005). Moreover, 3-Phosphoinositide-dependent kinase-1 (PDK1) has been reported to be negatively targeted in its activity by 14-3-3 (Sato et al., 2001). Our results are in agreement with the former observation if we consider that *ftt-2* may act by the same pathway in *C. elegans*. If 14-3-3 does so, we then expect to observe a pronounced life span shortening by *ftt-2* RNAi in a *daf-2(e1370)* class 2 mutant background because such effect may result in a slight rise in DAF-16 inhibition through a loss of 14-3-3-dependent negative regulation acting on PDK1. The results described above argue for such epistasis interaction. Moreover, if *pdk-1* also represents the main target for *ftt-2* regulation, one might expect to observe an unmodified life span of *pdk-1* knock out worms upon *ftt-2* RNAi while, on the other hand, the *ftt-1* RNAi should do so. We then submitted the *pdk-1(sa709)* loss-of-function allele to RNAi and, as expected (Fig. 2. E), the *ftt-2* RNAi failed to affect life span significantly (P=0.796; Table 1) while *ftt-1* RNAi remained with a statistically significant (P=0.0001; Table 1) life span modulation activity. This result argues for the fact that *ftt-2* dependent life span modulation operates mainly towards the *pdk-1* gene, a result fully correlated with the observed life span modulation induced by *ftt-2* RNAi in the *daf-2* class 2 mutant. However, *ftt-1* RNAi produced a negative modulation on life span during the same experiment (Fig. 2. E), while we previously reported an opposite effect in both *daf-2* class mutant (Fig. 2. C, D). It is possible that such *ftt-1* dependent negative modulation on life span, also independent from the *pdk-1* component of the DAF-2/DAF-16 pathway, was previously undetectable in both *daf-2* class-mutant where the strongest life span modulation is observed.

In an attempt to specify either *ftt-1* or *ftt-2* respective life span effects throughout PDK-1 signaling modulation, we next assessed a *pdk-1(mg142)* gain-of-function mutant for RNAi knock down experiment. As shown in Figure 2. F, the *ftt-2* RNAi but not the *ftt-1* RNAi

induced a significant life span modulation in comparizon with control. This result is relevant to a negative modulation of *pdk-1* activity by *ftt-2* that can be highlighted here by a life span shortening effect upon RNAi. The absence of a significant modulation effect on life span by *ftt-1* in the same genetic background also reinforces the view that *ftt-1* doesn't regulate life span towards *pdk-1* in the DAF-2/DAF-16 regulatory pathway (Fig. 2. F and table 1) and also indicate that *ftt-1* negative modulation on life span is also masked in the *pdk-1* gain-of-function background as observed below for both *daf-2* class mutant.

Thus, we show here for the first time a *daf-2* class mutant-dependent modulation of 14-3-3 on *C. elegans* life span regulation. These results indicate that 14-3-3 influences longevity not only when DAF-2 activity is high and DAF-16 low (as in N2), but also when DAF-2 activity is low and DAF-16 high (as in *daf-2* mutants). Therefore, while *ftt-1* dependent life span modulation is restricted to N2 and only one of both *daf-2* class mutants, the *tt-2* dependent life span modulations can affect N2 and both *daf-2* class mutants assessed, as well as a *daf-16* loss-of-function background. This observation argues for the predicted *ftt-2* gene function on life span modulation previously defined on the basis of its expression pattern (Morton et al., 2002; Wang et al., 2006) in comparison with the one of *ftt-1*. Moreover, on the basis of respective modulation of life span reported here with *ftt* RNAi on *daf-2* class mutants, we present here further evidence arguing that *ftt-2* is a more potent life span modulator than *ftt-1*. Altogether, our results indicate that *ftt-2* can modulate worm life span with different order of magnitude depending on the DAF-2/DAF-16 transduction level targeted that can be summarized for by a strong modulation effect when transduction levels targeted lie between *daf-2* and *pdk-1* then intermediate effect when transduction targeted levels are between *pdk-1* and *daf-16* and discrete modulation effect by a *daf-16* independent mechanism.

#### ***ftt-2* modulates intrinsic worm thermotolerance.**

*Daf-2* reduction-of-function mutants show an increased *intrinsic thermo tolerance (itt)* phenotype (Lithgow et al., 1995) and conversely, heat shock treatment can induce rapid nuclear localization of DAF-16 (Henderson and Johnson 2001; Henderson et al. 2006). Because this *itt* phenotype trait is believed to require DAF-16 activity (Lin et al., 2001) we then assessed for *ftt-1* and *ftt-2* involvement in this process. As shown in Fig. 3.A, no thermal resistance difference between N2 worms fed with either control RNAi or *ftt-2* RNAi was noted while we observed a slight and time limited (P=0.05) significant resistance difference when *ftt-1* RNAi. On the opposite, *daf-2(e1370)* or *daf-2(e1365)* worms subjected to same treatment after either *ftt-1* or *ftt-2* RNAi (Fig. 3. B and C) were less resistant. As attested by the level of significance observed (p=0.005), a protracted effect on *daf-2(e1370)* thermal stress resistance was observable upon *ftt-2* RNAi while a delayed but strongest effect was noted when *ftt-1* RNAi (Fig. B). A comparable modulation by both *ftt-1* and *ftt-2* RNAi on thermal stress resistance was observed for *daf-2(e1365)* genetic background (Fig. 3. C).

This observation may indicate, as we reported for life span, that intrinsic thermotolerance of both *daf-2* class mutant allele worms are prone to *ftt* -dependent regulation than wild type. More interestingly, the rise in thermal stress sensitivity when *daf-2* mutant worms were fed with *ftt-2* RNAi was not observable when *daf-16(mgDf50)* mutant or *pdk-1(sa680)* loss of function mutant (Fig. 3. E) where fed by the same (Fig. 3. D). Moreover, identical result was noted upon *ftt-1* RNAi (Fig. 3. D and E). This observation indicates that the *itt* trait phenotype is a part of the stress resistance mechanism allowing *ftt-2* and, to a lesser extend *ftt-1*, to specifically promote worm life span throughout DAF-16 dependent mechanism as described here and in Berdichevsky et al. (2006). The lack of *ftt-2* RNAi effect on N2 worms upon thermic stress may be due to a threshold sensibility difference in comparison with *daf-2* mutants, which are also regarded as constitutively stressed worms, thereby explaining, as described (Berdichevsky et al., 2006), the higher sensitivity of stress resistant mechanisms to

14-3-3-dependent regulation. However, these results are in agreement with the expected *itt* phenotype for the corresponding genetic backgrounds tested, because *daf-16(mgDf50)* appeared less heat-resistant than wild type or *daf-2(e1370)* mutant worms in control RNAi conditions.

### ***ftt-2* can modulate various stress responses.**

DAF-2/IGF-1 pathway regulates stress resistance in worm (Gems et al., 1998) and treating a *daf-2(e1370)* mutant with *daf-16* RNAi during the reproductive period can also modulate its life span (Dillin et al., 2002). DAF-16 gene products are believed to act as major cellular defense regulators when worms are facing stress. However, while some stress responsive genes, as for example *sod-3* (Henderson et al., 2006; Honda and Honda 1999; Murphy et al., 2003; Lee et al., 2003b; McElwee et al., 2003; Oh et al., 2006), are mainly FOXO/DAF-16 targets, other stress-inducible genes, such as heat-shock proteins (HSPs), can be synergistically regulated by FOXO/DAF-16 and HSF-1 transcription factors (Hsu et al., 2003; Morley and Morimoto 2004). Moreover, it has been demonstrated that DAF-16 is essential for maximal *hsp-16* expression and for life span extension conferred by *hsp-16* over expression (Walker and Lithgow 2003). Glutathione *S*-transferases (GSTs), oxidative stress responsive genes products, are believed to participate in metabolisation of cell compounds by post-translational glutathione tripeptide conjugation. Moreover, GSTs have already been reported to be up-regulated in *daf-2* mutant (McElwee et al., 2006) and by various physical or ROS-inducer chemical stressors (Leiers et al., 2003). We therefore decided to test for stress responsive gene modulation during adulthood of *ftt-2* RNAi-induced worms because it had already been reported (Dillin et al., 2002) that DAF-2/IGF-1 pathway could influence ageing and stress resistance during ageing. To this aim, we used a GFP reporter which had been used to assay for DAF-16 activity: The *sod-3::GFP* (Berdichevsky et al., 2006; Libina et al., 2003), as well as a HSPs reporter: *hsp-16::GFP* (Link et al., 2002) and the oxidative stress-inducible reporter transgene *gst-4::GFP* (Leiers et al., 2003; Link et al., 2002) whose constitutive or ROS-induced expression is not dependent on DAF-16 activity (Henderson et al., 2006). As shown in Fig. 4, while the expression level of these GFP reporters does not appear to change between adulthood day four to day eighteen (compare Fig. 4.A and B; Fig. 4.E and F; Fig. 4.I and J) in control worms, the expression pattern becomes greatly modified during the same adulthood interval when worms are fed with *ftt-2* RNAi (compare Fig. 4.C and D; Fig. 4.G and H; Fig. 4.K and L). However, we can clearly discriminate two groups of GFP reporter genes from which induction response to *ftt-2* RNAi treatment differs. One first group with a high expression induction level (*sod-3::GFP* and *gst-4::GFP*) and a second with a limited induction level (*hsp-16::GFP*). To increase discriminative analysis of reporter transgenes, we decided to perform pixel quantification analysis for both successive groups of growing old worms (see Material and Methods). As shown in Fig. 4.M, we detected a gradual rise in the expression level for each transgene during ageing and, as expected, to a lesser extent for *hsp-16::GFP* transgene. Moreover, the limited *hsp-16::GFP* rise in expression is also notable when we compare both adulthood periods tested. By contrast, we noted (particularly at day eighteen of adulthood) an important change for *sod-3::GFP* reporter expression level while *gst-4::GFP* showed an intermediate level of induction. Because these experiments were conducted in a stress-free condition, we conclude that stress-responsive GFP transgenes have been tested for their basal *ftt-2*-dependent regulation during normal aging. This observation sheds new light onto the regulatory links between the DAF-2/IGF-1 pathway - Believed to influence ageing and stress resistance in growing old worms by a *daf-16* dependent mechanism - and *sod-3::GFP* gene activation. However, we reported here that *hsp-16::GFP*, another stress inducible gene, behaves differently during the same recorded adulthood time, from *sod-3::GFP*. This observation may indicate that *hsp-16* gene expression regulatory mechanisms are less prone to *ftt-2* regulatory activity in an environment devoid of stress. Moreover, these results are in

complete agreement with our previous experiments showing a lack of involvement of 14-3-3/*ftt-2* for *itt* phenotypic trait in wild type worms. Altogether, our observations argue for a sustained involvement of *ftt-2* gene product for stress-responsive gene regulation during adulthood as reported for the DAF-2/IGF-1/DAF-16 pathway. Moreover the interesting observation of a pronounced *sod-3::GFP* transgene induction at day eighteen in comparison with day four upon treatment with *ftt-2* RNAi (compare Fig. 4.C and D) may also point out that regulatory mechanisms operating to maintain the physiological status of growing old worms are not the same during the whole post-developmental period. A recent report (Henderson et al., 2006) introduced that oxidative stress could induce nuclear translocation of DAF-16 without any detectable over-expression of *sod-3::GFP* reporter. However, our experiments clearly demonstrate a strict relationship between nuclear DAF-16 translocation and *sod-3::GFP* reporter induction in non-stress induction conditions although upon *ftt-2* RNAi.

### ***ftt-2* modulates worm oxidative stress resistance by a *daf-16*-independent mechanism.**

The DAF-2/DAF-16 regulatory pathway is believed to link worm oxidative stress resistance to life span (Finkel and Holbrook 2000; Hsu et al., 2003). We therefore investigated the effect of Menadione in different genetic backgrounds, believed to affect worm oxidative stress resistance. The Menadione (2-methyl-1,4-naphthoquinone or vitamin K3) is a lipid-soluble substance that has been extensively used as an oxidant stressor in yeast (Shackelford et al., 2000) as well as in *Arabidopsis thaliana* (Baxter et al., 2007). Menadione can generate reactive superoxide ions, which can be further oxidized to yield hydrogen peroxide and has been reported to deplete total glutathione concentration in yeast cells by generation of menadione S-glutathione conjugates (Zadzinski et al., 1998).

We examined the effect of Menadione by determining survival of adult worms subjected to 10mM Menadione after one week of either *ftt-1* or *ftt-2* RNAi treatment (see Material and Methods). As expected in control conditions, the *daf-16(mgDf50)* mutant was less resistant (Fig. 5. D) to Menadione treatment than N2 (Fig. 5. A), which also appeared less resistant to both *daf-2(e1370)* and *daf-2(e1365)* reduction-of-function mutants (Fig. 5. B and C). Moreover, we observed that *daf-16* mutant life span was significantly reduced ( $P=0.005$ ) upon both *ftt* RNAi in comparison with control RNAi (Fig. 5. D) while the same treatment did not induced a significant survival difference in either N2 or *daf-2(e1370)* genetic backgrounds (Fig. 5. A and B). However, we observed a time limited significant effect of *ftt-1* RNAi on *daf-2(e1365)* loss of function mutants survival (Fig. 5. C). Altogether, our results indicate that both *ftt-1* and *ftt-2* are involved in promoting Menadione oxidative stress resistance by acting through a *daf-16*-independent pathway. Although the mechanism through which *ftt-1* and *ftt-2* promotes *daf-16(mgDf50)* stress resistance is yet unknown, it is possible that different mechanism operates because we previously demonstrated that *ftt-2* is the only 14-3-3 ortholog able to promote life span in both *daf-16* loss-of-function and *daf-2* reduction-of-function genetic backgrounds. Altogether, these findings show that *ftt-2* can determinate stress resistance even in a non-stress resistant genetic background such as *daf-16(mgDf50)*.

### **Dauer formation can be enhanced by *ftt-2* inactivation.**

Several *daf-2* thermosensitive (*ts*) reduction-of-function mutations can result in constitutive dauer larval formation: a quiescent (*ie.* a non obligatory stress resistant arrested stage that differs from obligatory diapause stage affecting other worm species) larval stage with developmental arrest, non-feeding activity, stress resistance and long-lived characteristics (Klass M. and Hirsh D., 1976). Dauer induction fold seems to depend on *daf-2ts* class alleles that can be summarised as more severe for class 2 and weaker for class 1 mutant. Therefore, the semi-restrictive temperature (22,5°C) applied to *daf-2ts* mutants can modify the

population that constitutively enters the dauer stage (Gems et al., 1998). We believed that such semi-restrictive temperature would help us to reveal possible *ftt-1* or *ftt-2*-dependent dauer formation process by generating a facilitated constitutive dauer commitment sub-threshold. We therefore assessed relevant class 2 *daf-2(e1370)* and class 1 *daf-2(e1365)* allele for dauer formation upon *ftt-2* RNAi at 22.5°C.

While *ftt-1* RNAi produced no detectable effect on dauer formation in all genetic backgrounds tested (Fig. 6), we observed a highly significant difference (P=0.005) upon *ftt-2* RNAi on *daf-2* class 1 mutant and a less significant difference (P=0.05) on *daf-2* class 2 mutant (Fig. 6). This observation indicates that class 1 *daf-2ts* mutant alleles are more prone to *ftt-2* dependent regulation than class 2 for dauer commitment while we reported the opposite effect for life span assays. This observation indicates for the first time the existence of a *daf-2* class mutant allele-specific modulation for dauer formation by *ftt* genes. However, because dauer entry and exit are not equivalent physiological processes, as revealed by time course microarray analysis (Wang and Kim 2003), we next asked if dauer exit might behave with the same difference between both *daf-2* mutants alleles tested. We then checked dauer population and observed full dauer exit for both *daf-2* reduction-of-function allele populations 120 hours after hatching.

### ***ftt-2* influences worm life span by acting during development.**

While *daf-2* RNAi can influence aging when applied only during worm adulthood (Kenyon 2005; Dillin et al., 2002), *daf-2* deletion mutants are affected by non-conditional arrest at embryonic or first larval stage (Gems et al., 1998). In addition, genes encoding mitochondrial respiratory chain components have been reported to influence worm life span if reduction-of-function, mimicked by RNAi, was applied only before the reproductive stage (Dillin et al., 2002b; Lee et al., 2003a).

We observed a variation on the worm populations that did not arrest at dauer stage upon either *ftt-1* or *ftt-2* RNAi at semi-restrictive temperature. Thus, while 82% of *daf-2(e1370)* worms did not arrest at dauer stage on control RNAi (see Material and Methods), the corresponding rate decreased to 59% and reached up to 81% when worms were fed for *ftt-2* and *ftt-1* RNAi respectively.

Moreover, in the *daf-2(e1365)* genetic background, non-dauer arrested worms fraction reached 83% on control RNAi but only 39% on *ftt-2* RNAi while *ftt-1* RNAi induced a comparable rate to wild type as 80%. Wild type worms did not revealed any dauer formation on control or on both RNAi treatments at 22.5°C and 25°C too (data not shown).

We then assessed the worm fraction with uninterrupted larval stage progression for developmental rate when RNAi knock down. Both wild type and *daf-16(mgDf50)* worms fed by *ftt-2* or *ftt-1* RNAi protracted 24 hours to reach the fourth larval stage (96 hours after hatching) when compared to control worms. Moreover, both *daf-2(e1370)* and *daf-2(e1365)* worms protracted 48 hours to reach the fourth larval stage (120 hours after hatching) when *ftt-2* RNAi but only 24 hours (96 hours after hatching) when *ftt-1* RNAi. This observation indicates that *ftt-2*, and to a lesser extend *ftt-1*, reduction-of-function applied during worm development can induce a "slow living phenotype" by perhaps acting, more or less directly, through the modulation of a yet unknown transduction pathway able to induce others defects as reported here and in previous reports (Wang et al., 2006; Berdichevsky et al., 2006).

Because development was affected with the same order of magnitude on both *daf-2* class mutant at semi-restrictive temperature, we then addressed for a possible influence of this developmental modulation on subsequent post-developmental stages.

To this aim, worms that reached the fourth larval stage were shifted to *ftt-2* or *ftt-1* RNAi free condition at 22.5°C and assessed for life span (see Material and Methods). As shown in Fig. 7, we observed a highly significant (P=0.001) life span shortening for wild type worms induced with *ftt-2* RNAi than *ftt-1* RNAi in comparison with controls (Fig. 7.A and Table 2).

Moreover, both *daf-2(e1370)* and *daf-2(e1365)* worms showed a remarkable life span shortening on both RNAi (Fig. 7. C and D and Table 2) in comparison with non-induced worms and wild type worms after equivalent treatment (Table 2). This unexpected result highlights for the first time that *ftt-2*, and to a lesser extent *ftt-1*, RNAi knock down can shorten life span if administered only during development in both wild type and *daf-2* reduction-of-function mutants.

Moreover, the observed higher mortality rate for both *daf-2* class mutants indicates that they are more prone than wild type worms to *ftt* dependent regulations for development - A similar observation is also reported below for post-developmental life span experiments (Fig. 2; Table1).

However, because *daf-16(mgDf50)* worms behaved the same as both *daf-2* mutants (Fig.7. B), we also conclude that *ftt*-dependent control of development involves both DAF-2/DAF-16-dependent and -independent regulatory mechanisms.

The high mortality rate and mortality curve shape observed during our experiments may reflect possible non-specific sickness acting on life span by *ftt* RNAi knock down but other observations argue against such a simple conclusion. Thus, as shown in Table 2, it is possible to observe that *ftt* RNAi do not impair by the same worms life span with different genetic backgrounds. For example, while the *ftt-2* RNAi induced a 26% mean life span reduction in wild type, we showed a here 60% mean life span reduction for *daf-2(e1365)* and up to 67% reduction for *daf-2(e1370)* while *daf-16(mgDf50)* worms were less affected with a 44% mean life span reduction only.

If *ftt* RNAi knock downs produce a non-specific effect on life span, one might expect a proportional linear rise in life span reduction from short-living worms (as *daf-16(mgDf50)*) to long-living worms (*daf-2* class mutants). However, this was not the case because short-living worms were more affected than wild type worms. Moreover, the *daf-2* class 2 (*daf-2(e1370)*) life span was more affected upon *ftt-2* RNAi than *daf-2* class 1 (*daf-2(e1365)*) as already reported for post-developmental life span experiments below (Figure 2, table 1) were *daf-2(e1370)* and *daf-2(e1365)* life span were reduced by 36% and 32% respectively.

This observation provides new insight into both *ftt-2* and *ftt-1* -dependent regulations on life span that need to be specifically assessed during development or adulthood to avoid misinterpretation of results due to possible unrelated cumulative effects produced by *ftt* RNAi administered during pre- and post-reproductive periods. For example, while we reported both a small percentage of protruding vulvae phenotype (15%) and bagging phenotype in adult worms fed on OP50 bacteria after *ftt-2* RNAi induced during development, we noted that the same *ftt-2* RNAi treatment applied during adulthood induced only the life span reduction phenotype. Therefore, noticing that *daf-2* mutants life expectancy was more affected than the wild type clearly indicates that *ftt-2*, and to a lesser extent *ftt-1*, acts as a more potent modulator during development when DAF-2 signalling is low and DAF-16 activity high.

### ***C. elegans* fat storage can be modulated by *ftt-2*.**

During our dauer commitment analysis we noted a reproducible “dark appearance” shape variation between control and *ftt-2* RNAi induced worms. Such “dark appearance” shape phenotype can be explained by lipid droplet accumulation as already described for *daf-2* reduction-of-function mutation or dauer larvae (Kimura et al., 1997).

The increased fat content trait of *daf-2* mutants has been shown to be fully suppressible by *daf-16* mutants (Lin et al., 1997; Ogg et al., 1997) and we reported below (Fig.6) that *ftt-2* but not *ftt-1* RNAi can increase dauer larvae (a well characterized resistant stage with high lipid accumulation) commitment in two different *daf-2* class mutants together with a more significant effect for class 1 mutant allele. Moreover, it has been demonstrated that some *daf-2* mutations (class 2) can induce dauer-like trait expression in adults while others (class 1) do not and show a normal metabolism instead (Gems et al., 1998). However, recent microarray

analysis reported the identification of a transcriptional products group whose expression was highly similar in both dauers and *daf-2* mutant adults (McElwee et al., 2003; McElwee et al., 2006) while some other groups behaved differently. We therefore assessed if *ftt-2* or *ftt-1* could influence lipid metabolism in both *daf-2* class mutant alleles with the same order of difference as for dauer commitment. As shown in Fig. 8 (A-L), the apparent fat content of *daf-16(mgDf50)*: Fig. 8.D and F, and *daf-2(e1365)*: Fig. 8.G and I was reduced upon *ftt-2* RNAi while both wild type worms (Fig. 8.A and C) and *daf-2(e1370)* fat content (Fig. 8.J and L) appeared less affected. Nevertheless, the *ftt-1* RNAi did not induce apparent lightening in lipid staining excepted for *daf-2(e1370)* genetic background (Fig.8. J and K).

Therefore, relative quantification (see Material and Methods) of Nile Red lipid staining (Fig. 9) provided more insight with respect to the difference of fat storage that appeared more significantly impaired in *daf-2(e1365)* and *daf-16(mgDf50)* worms than in *daf-2(e1370)* and N2 worms when *ftt-2* RNAi while *ftt-1* RNAi produced a significant lipid staining reduction in *daf-2(e1370)* worms only.

This observation provides novel insights into the striking opposite effect of *ftt-2* RNAi on both dauer induction and its accompanying lipid droplet accumulation trait. More unexpectedly, we also found a significant fall of lipid droplet accumulation in a dauer defective genetic background mutant as *daf-16(mgDf50)*. On the basis of this experimental evidence, we propose that *ftt-2* can promote lipid accumulation by a *daf-16* -independent pathway. Therefore, *ftt-2* can increase dauer commitment in *daf-2* class 1 mutant allele, an observation indicating a modulation via alternative regulatory pathway that may (Gems et al., 1998; Nanji et al., 2005) or may not emerge from the DAF-2 receptor. Thus, while both dauers and *daf-2* adults contain large quantities of lipid droplets, it has been reported (Walker and Lithgow 2003) that triglyceride synthesis enzymes are not expressed in the same fashion during these two *C. elegans* life history stages leaving open possible differences in the regulatory pathways targeted by *ftt-2* besides of DAF-2/IGF-1 regulatory pathway. Moreover, a quantitative mass spectrometry analysis reported (Dong et al., 2007) a subset of significant abundant proteins in *daf-2(e1370)* mutant were able to either slow down or accelerate ageing rate when targeted by RNAi knock down then living open the possibility for compensatory mechanism with opposites effects on the rate of ageing but altogether with a positive global effect as observed in long living *daf-2* mutants. It is possible that some molecular targets of *ftt-2* may fall into both classes of protein thereby explaining opposite effects on dauer commitment and lipid accumulation reported in this study.

## Discussion

### The *ftt-2* can act as a life span determinant regulator in *C. elegans*.

Because of a reported lethal phenotype for both *ftt-1* and *ftt-2* loss-of-function mutants (Wang et al., 2006; Berdichevsky et al., 2006; Li et al., 2007), we used RNAi in an attempt to investigate the *C. elegans* 14-3-3 genes homolog involvement in life span regulation. The RNAi experiments described in our study allowed us to analyze developmental as well as post-developmental *ftt*-dependent modulation of various *C. elegans* life history traits.

We report here that *ftt-2* is a more potent life span modulator in all of the *C. elegans* genetic backgrounds that we assessed as well as in a closer worm species such as *C. briggsae* (Araiz and Galas; manuscript in preparation). As previously proposed in other reports (Cahill et al., 2001; Berdichevsky et al., 2006), the *C. elegans* *ftt* genes can promote cytoplasmic retention of phosphorylated DAF-16 but can also mediate activation of a 14-3-3/SIR-2.1/DAF-16 stress mediated-dependent pathway to elicit stress resistance and life span extension. Here we show that *ftt-2* can promote *C. elegans* life span in stress-free conditions. Moreover, we show for the first time the existence of two actions levels of *ftt-2* (as well as for *ftt-1*) for life span



modulation. Thus, while *ftt-2* gene knockout by RNAi can induce side morphological problems (protruding vulva or bagging phenotype, as also reported elsewhere (Wang et al., 2006; Berdichevsky et al., 2006)), together with a strong life span shortening, the same RNAi induction only during the post-developmental period induces a highly significant life span shortening without morphological side effects. We also demonstrate here the specificity of *ftt-1* and *ftt-2* RNAi induction by the observation of non-overlapping induced phenotypes.

In summary we show here that *ftt-2*, and to a lesser extent the *ftt-1*, can positively modulate the life expectancy during worm adulthood by acting at developmental stages only. This mechanism may reflect the 14-3-3 pleiotropic effects on worm signalling systems because both *daf-2* class mutant assessed in our RNAi knock down experiments appeared to be modulated to the same relative extent difference as the control. We also report in this work that *ftt-2* can promote worm life span throughout a negative modulation of the *pdk-1* while *ftt-1* does not. However, we show here that *daf-2* class 2 mutants are more prone to *ftt-2* modulation for life span than *daf-2* class 1 mutant alleles during adulthood. Previous reports (Gems et al., 1998; Nanji et al., 2005) have shown that *daf-2* class 2 mutant alleles are genetically characterised by a greater loss of PI3kinase signalling, which reflects multiple signal outputs from the DAF-2 receptor. It is possible to conclude that main effect on life span by *ftt-2* is mediated by this transduction pathway (as depicted in Fig. 9) because of a *pdk-1* requirement as revealed by our experiments. Altogether, the experiments described here demonstrate for the first time a *ftt-2* (as well as *ftt-1*) life span different modulation in both *daf-2* class 1 and class 2 mutants.

#### ***ftt-2* can modulate life span in a *daf-16* null genetic background**

In this report, we show that *daf-16* loss-of-function mutant retains a *ftt-2* but not *ftt-1*-dependent life span modulation (Fig. 9). This result indicates that additional mechanisms can regulate worm life span in a *daf-16*-independent but *ftt-2*-dependent manner. Moreover, we show for both *daf-2* class mutants assessed in this work, that *ftt-2* RNAi only during developmental life history period can also exert a deleterious effect on worm life span.

Previous reports introduced that SIR-2.1, a *C. elegans* NAD-dependent deacetylase homolog, can interact with FTT-1 and FTT-2 proteins to activate DAF-16 and extend the life span of stressed worms only (Wang et al., 2006; Berdichevsky et al., 2006; Li et al., 2007 and Fig. 8). Moreover, these reports showed that *sir-2.1* or *ftt-1* and *ftt-2* were not required for life span extension when reduced DAF-2/IGF-1 signalling although such studies only assessed one *daf-2* class mutant. We demonstrate here the existence of a new regulatory mechanism involving the *ftt-2* gene and acting via *daf-16* and the DAF-2/IGF-1 pathway together with another yet unknown *daf-16* -independent process for normal life span modulation, stress resistance and metabolism regulation (Fig. 8). Because previous studies did not assess for the FTT-2/Sir-2.1/DAF-16 stoichiometric regulatory complex and because it is always possible that cytoplasmic retention of DAF-16 upon *ftt-2* RNAi does not reflect the fate of the whole DAF-16 cellular pool, this new *ftt-2* target pathway may use part of the FTT-2 protein pool to regulate life span in a *daf-16* -independent manner.

#### **Dauer formation and lipid accumulation can be modulated by *ftt-2* knock down.**

It has been reported that dauer formation regulation of *daf-2* class 1 mutant occurs mainly via a parallel pathway from the *daf-2/lage-1/daf-16* pathway (Gems et al., 1998; Nanji et al., 2005). However we show here that dauer commitment of both *daf-2* class mutants are greatly enhanced when *ftt-2* is targeted by RNAi while *ftt-1* RNAi produces no detectable effect.

Because we show that *daf-2* class 2 allele mutants are more prone to positive life span regulation by *ftt-2* than class 1 mutant allele and because we show that dauer induction upon *ftt-2* RNAi can be greatly enhanced in *daf-2* class 1 mutant, we propose that an alternative

*daf-2/daf-16* pathway is more prone to *ftt-2* regulation for dauer formation but not for life span modulation. Moreover, because lipid droplet accumulation is a well reported accompanying metabolic phenotype of worm dauer induction as well as in dauer constitutive phenotypes, we demonstrate here that *ftt-2* must be unexpectedly involved in promoting lipid droplet accumulation together with a negative influence in dauer commitment while *ftt-1* is not involved in dauer commitment regulation but only for lipid accumulation in *daf-2* class 2 mutant.

***ftt-2* can modulate worm stress resistance responses during adulthood period.**

We reveal here that *ftt-2* promotes stress resistance by both *daf-2* dependent and *daf-16* - independent processes. We also show that this effect is correlated with the up-regulation of a subset of *daf-16* target genes that have been involved in stress resistance. However, because our experiments were done in stress-free conditions, we can conclude here for a positive modulation by *ftt-2* in promoting stress-responsive genes. Moreover, we also demonstrate that *ftt-2* gene can promote the up-regulation of DAF-16 target genes that have been involved in stress resistance. We also demonstrate by two independent observations (heat shock assay and HSP-16::GFP expression) that heat shock responsive gene expression is not modulated by *ftt-2* thereby providing new insights into the fine tuning regulations of DAF-16 molecular targets. Such fine tuned regulations may use the new emerging family of FOXO/DAF-16 co-factors that can account for simultaneous *daf-16* -dependent and -independent processes regulating life span programs in stress conditions in *C. elegans* (Berdichevsky and Guarente 2006).

In conclusion, this study demonstrates, for the first time, the ability of the *C. elegans* 14-3-3 genes to modulate the life expectancy during adulthood even in a stress-free environment.

The new modulation properties on the various physiological traits assessed in this work (Fig. 9) yet extend the involvement of 14-3-3-dependent regulations from short time events, such as the one counteracting the danger of sudden darkness in plants (Mackintosh 2004) or acute stress response in *C. elegans* to a more sustained event such as normal life span regulation.

## Acknowledgments

We thank Theresa Stiernagle and the *Caenorhabditis elegans* Genetics Center for providing strains. We are grateful to Donald L. Riddle for the *daf-2(e1365)* strain and Diane Shake for kindly providing us with the 14-3-3/*ftt* cDNAs clones. We are especially grateful to MRI ([www.mri.cnrs.fr](http://www.mri.cnrs.fr)) and its technical assistance. We thank Julia Pfitzner, Victoria Malchin, Nga Nguyen and Professor Ulrich Mok, from Lise-Meitner-Oberstufenzentrum, Berlin, Germany, for technical assistance. We thank Dr. May Morris and Dr. Jean Michel Bellanger and Dr. Andrew Burgess for critical reading of the manuscript. We thank Dr. Anne Blangy for helpful with qPCR experiments. This work was supported by the CNRS, C. Araiz. is a PhD fellow of AFM ([www.afm-france.org](http://www.afm-france.org)) and ARC ([www.arc.asso.fr/](http://www.arc.asso.fr/)).

## References

- Baxter CJ, Redestig H, Schauer N, Repsilber D, Patil KR, Nielsen J, Selbig J, Liu J, Fernie AR, Sweetlove LJ. (2007) The Metabolic Response of Heterotrophic Arabidopsis Cells to Oxidative Stress. *Plant Physiol.* 143, 312-325.
- Berdichevsky A, Viswanathan M, Horvitz H.R. and Guarente L. (2006) C. elegans SIR-2.2 interacts with 14-3-3 Proteins to activate DAF-16 and extend life span. *Cell*, 125, 1165-1177.
- Berdichevsky A, Guarente L. (2006) A stress response pathway involving sirtuins, forkheads and 14-3-3 proteins. *Cell Cycle*. 5, 2588-2591.
- Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics* 77, 71–94.
- Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, Anderson MJ, Arden KC, Blenis J, Greenberg ME. (1999) Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell*. 96, 857-868.
- Cahill CM, Tzivion G, Nasrin N, Ogg S, Dore J, Ruvkun G and Alexander-Bridges M. (2001) Phosphatidylinositol 3-Kinase signaling inhibits DAF-16 DNA Binding and function via 14-3-3-dependent and 14-3-3-independent pathways. *J. Biol. Chem.*, 276, 13402-13410.
- Craparo A, Freund R and Gustafson TA. (1997). 14-3-3 (epsilon) interacts with the insulin-like growth factor I receptor and insulin receptor substrate in a phospho-serine dependent manner. *J. Biol. Chem.*, 272, 11663-11669.
- Dillin A, Crawford DK, Kenyon C. (2002) Timing requirements for insulin/IGF-1 signaling in C. elegans. *Science*. 298, 830-834.
- Dillin A, Hsu AL, Arantes-Oliveira N, Lehrer-Graiwer J, Hsin H, Fraser AG, Kamath RS, Ahringer J, Kenyon C. (2002b) Rates of behavior and aging specified by mitochondrial function during development. *Science*. 298, 2398-2401.
- Finkel T, Holbrook NJ. (2000) Oxidants, oxidative stress and the biology of ageing. *Nature*. 408, 239-247.
- Gems D, Sutton AJ, Sundermeyer ML, Larson PL, Albert PS, King KV, Edgley M and Riddle DL (1998) Two pleiotropic classes of daf-2 mutation affect larval arrest, adult behavior, reproduction and longevity in *Caenorhabditis elegans*. *Genetics*, 150, 129-155.
- Henderson ST and Johnson TE (2001) daf-16 integrates developmental and environmental inputs to mediate aging in the nematode *Caenorhabditis elegans*. *Curr. Biol.*, 11, 1975-1980.
- Henderson ST, Bonafe M, Johnson TE. (2006) daf-16 protects the nematode *Caenorhabditis elegans* during food deprivation. *J Gerontol A Biol Sci Med Sci*. 61, 444-4460.
- Honda Y, Honda S. (1999) The daf-2 gene network for longevity regulates oxidative stress resistance and Mn-superoxide dismutase gene expression in *Caenorhabditis elegans*. *FASEB J*. 13, 1385-1393.
- Hsu AL, Murphy CT, Kenyon C. (2003) Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science*. 300, 1142-1145.
- Kenyon C. (2005) The plasticity of aging: insights from long-lived mutants. *Cell*. 120, 449-460.
- Kimura KD, Tissenbaum HA, Liu Y, Ruvkun G. (1997) daf-2, an insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science*, 277, 942-946.
- Klass M, Hirsh D. (1976) Non-ageing developmental variant of *Caenorhabditis elegans*. *Nature*. 260, 523-525.
- Lee SS, Lee RY, Fraser AG, Kamath RS, Ahringer J, Ruvkun G. (2003a) A systematic RNAi screen identifies a critical role for mitochondria in C. elegans longevity. *Nat Genet*. 33, 40-48.
- Lee SS, Kennedy S, Tolonen AC, Ruvkun G (2003b) DAF-16 target genes that control C. elegans life-span and metabolism. *Science*. 300, 644-647.
- Leiers B, Kampkotter A, Grevelding CG, Link CD, Johnson TE, Henkle-Duhrsen K (2003) stress-responsive glutathione S-transferase confers resistance to oxidative stress in *Caenorhabditis elegans*. *Free Radic Biol Med*. 34, 1405-1415.

- Libina N, Berman JR, Kenyon C (2003) Tissue-specific activities of *C. elegans* DAF-16 in the regulation of lifespan. *Cell*. 115, 489-502.
- Dong MQ, Venable JD, Au N, Xu T, Park SK, Cociorva D, Johnson JR, Dillin A, Yates JR 3rd. (2007) Quantitative mass spectrometry identifies insulin signaling targets in *C. elegans*. *Science*. 317, 660-663.
- Li J, Tewari M, Vidal M and Lee SS (2007) The 14-3-3 protein FTT-2 regulates DAF-16 in *Caenorhabditis elegans*. *Dev Biol*. 301, 82-91.
- Lin K, Dorman JB, Rodan A, Kenyon C (1997) *daf-16*: An HNF-3/forkhead family member that can function to double the life-span of *Caenorhabditis elegans*. *Science*. 278, 1319-1322.
- Lin K, Hsin H, Libina N and Kenyon C (2001) Regulation of the *Caenorhabditis elegans* longevity protein DAF-16 by insulin/IGF-1 and germline signaling., *Nat. Genet*. 28: 139-145.
- Link CD and Johnson CJ (2002) Reporter transgenes for study of oxidant stress in *Caenorhabditis elegans*. *Methods Enzymol*. 353, 497-505
- Lithgow GJ, White TM, Melov S, Johnson TE (1995) Thermotolerance and extended life-span conferred by single-gene mutations and induced by thermal stress. *Proc Natl Acad Sci USA*. 92, 7540-7544.
- Mackintosh C (2004) Dynamic interactions between 14-3-3 proteins and phosphoproteins regulate diverse cellular processes. *Biochem J*. 381, 329-342.
- McElwee J, Bubb K, Thomas JH (2003) Transcriptional outputs of the *Caenorhabditis elegans* forkhead protein DAF-16. *Aging Cell*. 2, 111-121.
- McElwee JJ, Schuster E, Blanc E, Thornton J, Gems D (2006) Diapause-associated metabolic traits reiterated in long-lived *daf-2* mutants in the nematode *Caenorhabditis elegans*. *Mech Ageing Dev*. 127, 458-472.
- Mitchell DH, Stiles JW, Santelli J, Sanadi DR (1979) Synchronous growth and aging of *Caenorhabditis elegans* in the presence of fluorodeoxyuridine. *J Gerontol*. 34, 28-36.
- Morley JF, Morimoto RI. (2004) Regulation of longevity in *Caenorhabditis elegans* by heat shock factor and molecular chaperones. *Mol Biol Cell*. 15, 657-664.
- Morton DG, Shakes DC, Nugent S, Dichoso D, Wang W, Golden A, Kemphues KJ. (2002) The *Caenorhabditis elegans* *par-5* gene encodes a 14-3-3 protein required for cellular asymmetry in the early embryo. *Dev Biol*. 241, 47-58.
- Murphy CT, McCarroll SA, Bargmann CI, Fraser A, Kamath RS, Ahringer J, Li H, Kenyon C. (2003) Genes that act downstream of DAF-16 to influence the lifespan of *Caenorhabditis elegans*. *Nature*. 424, 277-283.
- Nanji M, Hopper NA and Gems D. (2005) LET-60 modulates effects of insulin/IGF-1 signaling on development and aging in *Caenorhabditis elegans*. *Aging Cell*. 4, 235-245.
- Oh SW, Mukhopadhyay A, Dixit BL, Raha T, Green MR, Tissenbaum HA. (2006) Identification of direct DAF-16 targets controlling longevity, metabolism and diapause by chromatin immunoprecipitation. *Nat Genet*. 38, 251-257.
- Ogg S, Paradis S, Gottlieb S, Patterson GI, Lee L, Tissenbaum HA, Ruvkun G. (1997) The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature*. 389, 994-999.
- Pfaffl MW. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucl. Acid. Res.*, 29:e45.
- Sato S, Fujita N and Tsuruo T (2001) Regulation of kinase activity of 3-Phosphoinositide-dependent Protein Kinase-1 by binding to 14-3-3. *J. Biol. Chem.*, 277, 39360-39367.
- Shackelford RE, Kaufmann WK, Paules RS (2000) Oxidative stress and cell cycle checkpoint function. *Free Radic Biol Med*. 28, 1387-404.
- Van Hemert MJ, Steensma HY, Van Heusden GP (2001) 14-3-3 proteins: key regulators of cell division, signalling and apoptosis. *Bioessays*. 23, 936-946.
- Walker GA, Lithgow GJ. (2003) Lifespan extension in *C. elegans* by a molecular chaperone dependent upon insulin-like signals. *Aging Cell*. 2, 131-139.

- Wang W, Shakes DC. (1997) Expression patterns and transcript processing of *ftt-1* and *ftt-2*, two *C. elegans* 14-3-3 homologues. *J Mol Biol.* 268, 619-630.
- Wang W, Shakes DC. (1994) Isolation and sequence analysis of a *Caenorhabditis elegans* cDNA which encodes a 14-3-3 homologue. *Gene.* 147, 215-218.
- Wang J, Kim SK (2003) Global analysis of dauer gene expression in *Caenorhabditis elegans*. *Development.* 130, 1621-1634
- Wang Y, Wook Oh S, Deplancke B, Luo J, Walhout AJM and Tissenbaum H A (2006) *C. elegans* 14-3-3 proteins regulates life span and interact with SIR-2.1 and DAF-16/FOXO. *Mech Ageing Dev.* 127, 741-747.
- Zadzinski R, Fortuniak A, Bilinski T, Grey M and Bartosz G (1998) Menadione toxicity in *Saccharomyces cerevisiae* cells: activation by conjugation with glutathione. *Biochem Mol Biol Int.* 44, 747-759.

ACCEPTED MANUSCRIPT

## Table Legends

### Table 1 : Life span modulation by 14-3-3 depends on *C. elegans* genotype in stress-free conditions.

The tabulated data show the average results of four to eight independent RNAi trials. Worms of indicated genotypes were assayed for life span by feeding on the *E. coli* HT115 strain that express either *ftt-1*, *ftt-2* RNAi or control vector (see Material and Methods). XLSTAT-life statistical software (Addinsoft, New York, NY, USA) was used to plot survival data by the Kaplan Meier method and differences between survival curves were calculated using the LogRank test with 95% confidence.

(£) Represents the 50th percentile (the age when the survival fraction of animals reaches 0,50%.

(•) Experiment identification code.

(\$) Probability of being identical to other life span experiment indicated in parentheses.

(\*) Total death scored (number of censored values).

### Table 2 : *C. elegans* life span shortening by pre-reproductive *ftt-2* RNAi induction.

The tabulated data show the average results of four to five independent RNAi trials at 22.5°C. Worms of indicated genotypes were fed on *E. coli* HT115 strain bacteria expressing either *ftt-2*, *ftt-1* or control RNAi during pre-reproductive developmental period then transferred to RNAi-free conditions and assayed for life span (see Material and Methods).

XLSTAT-life statistical software (Addinsoft, New York, NY, USA) was used to plot survival data by the Kaplan Meier method and differences between survival curves were calculated using the LogRank test with 95% significance.

(@) Mean life span values (Standard Deviation).

(\$) Probability of being identical to other life span experiment indicated in parentheses.

(\*) Total death scored (number of censored values).

## Legends to Figures:

Figure 1. The *ftt-1* and *ftt-2* RNAi treatments induce specific mRNA knock down.

N2 were treated by RNAi feeding for either *ftt-1* or *ftt-2* gene specific knock down during adulthood (see materials and Methods). The y-axis indicates the relative RNA levels of either *ftt-1* or *ftt-2* normalized to the RNA level of tubulin (*tba-2*). One representative experiment with the standard error of the mean (S.E.M.) is shown. The *P*-value of <0.005 (\*\*) was determined by Student's *t*-test.

**Figure 2. Both *C. elegans* 14-3-3 can shorten adult life span but only *ftt-2* can do it in various genetic backgrounds.**

Adult worm life-span analysis were performed on RNAi plates as described in Material and Methods. (A-E) The squared lines represent the life span of animals of indicated genotype fed by bacteria carrying control RNAi vector. Circle lines and triangle lines represent life spans of indicated genotypes fed with bacteria carrying either the *ftt-2* or the *ftt-1* RNAi constructs.

(A) Both *ftt-2* and *ftt-1* RNAi shorten life span of wild type (N2) worms but with different order of magnitude.

(B) Only *ftt-2* RNAi shortens life span in a *daf-16* null genetic background.  
 (C-D) Only *ftt-2* RNAi can shorten life span in both *daf-2* class 1 and class 2 reduction-of-function mutants.  
 (E-F) The *ftt-2* life span modulation operates mainly throughout *pdk-1* gene.  
 See Table 1 for corresponding quantitative and statistical analysis. All experiments were performed at 20°C.

**Figure 3. *ftt-2* RNAi knock down shortens worm survival after heat shock induction in *daf-2* reduction-of-function background but not in both wild type (N2) or *daf-16* loss-of-function genetic background.**

(A-E) Young adults of the indicated genotype were fed by bacteria carrying either *ftt-1*, *ftt-2* or control RNAi construct, then subjected to heat shock (Material and Methods) and their viability scored at the indicated times. Results presented are mean values +/- S.E.M. of five independent experiments. Statistical analysis was done by a Student's *t*-test where \* *P* indicates *P* < 0.05 significance level. All experiments were performed at 20°C.

**Figure 4. *ftt-2* RNAi can promotes activation of stress responsive gene reporters during adulthood in non-stressing conditions.**

(A-L) Wild type (N2) young adult worms carrying either a *sod-3::GFP* (A-D), a *hsp-16::GFP* (E-H) or a *gst-4::GFP* (I-L) stress reporter transgene were fed by bacteria carrying either control RNAi vector (A-B; E-F; I-J) or *ftt-2* RNAi construct (C-D; G-H; K-L) and scored for transgene expression at fourth (A,C,E,G,I,K) and eighteenth days (B,D,F,H,J,L) of adulthood by fluorescence microscopy as described in Material and Methods. Note higher level of *sod-3::GFP* transgene expression at day eighteen of adulthood only when worms fed *ftt-2* RNAi construct (D) in comparison with same worms scored at fourth day of adulthood (C) or with worms fed on control RNAi at both days of adulthood (A-B). Both *sod-3::GFP* (C-D) and *gst-4::GFP* (K-L) transgene inductions during adulthood are greater than *hsp-16::GFP* (G-H) when compared to controls at both adulthood periods (*sod-3::GFP*: A-B; *hsp-16::GFP*: E-F; *gst-4::GFP*: I-J). All images were taken at 40X magnification using fluorescence microscopy with an exposure time of 1000ms for A-D; 200ms for E-H and 1000ms for I-L. All experiments were performed at 20°C. the positive transgene induction control tests are not shown. (M) *ftt-2* RNAi-dependent induction of stress responsive genes during adulthood. Wild type (N2) young adult worms carrying either *sod-3::GFP*, *hsp-16::GFP* or a *gst-4::GFP* stress reporter transgene were fed by bacteria carrying either control RNAi vector or *ftt-2* RNAi construct and scored for stress responsive transgene expression at fourth and eighteenth days of adulthood. Quantifications were done by fluorescence microscopy as described in Material and Methods on at least ten animals. Bars represent the standard error (SEM) and mean values are expressed after normalisation with respective control RNAi, which represent 100%. Statistical Student's *t*-test (\**P* value < 0.05; \*\* *P* value < 0.0005) comparison between day four and day eighteen values are: \*\* *P* = 0.0003 for *sod-3::GFP*; *P* = 0.09 for *hsp-16::GFP* and \* *P* = 0.01 for *gst-4::GFP*. All experiments were performed at 20°C.

**Figure 5. Both *ftt* RNAi knock down prevent oxidative stress resistance by a *daf-16* independent mechanism.**

(A-D) Young adults of the indicated genotype were fed by bacteria carrying either *ftt-1*, *ftt-2* or control RNAi construct, then subjected to oxidative treatment by 10mM Menadione (as described in Material and Methods) and their viability scored at the indicated time. Results presented are mean values +/- S.E.M. of at least five independent experiments. Statistical analysis was done by a Student *t*-test at \* *P* < 0.05 and \*\* *P* < 0.0005 significance level. The total number of worms scored for each genotype and RNAi condition are in parentheses: Wild type Control RNAi (N=218); Wild type-*ftt-1* RNAi (N=174); Wild type-*ftt-2* RNAi (N=238);



*daf-16(mgDf50)*-Control RNAi (N=169); *daf-16(mgDf50)-ftt-1* RNAi (N=587); *daf-16(mgDf50)-ftt-2* RNAi (N=180); *daf-2(e1365)*-Control RNAi (N=159); *daf-2(e1365)-ftt-1* RNAi (N=164); *daf-2(e1365)-ftt-2* RNAi (N=183); *daf-2(e1370)*-Control RNAi (N=219); *daf-2(e1370)-ftt-1* RNAi (N=277); *daf-2(e1370)-ftt-2* RNAi (N=193). All experiments were performed at 20°C.

**Figure 6. *ftt-2* RNAi knock down promotes dauer constitutive induction phenotype in both *daf-2* class 1 and 2 reduction-of-function mutants.**

Worms of indicated genotype were allowed to hatch on bacteria carryings either *ftt-1*, *ftt-2* or control RNAi construct at 22.5°C in order to create a dauer constitutive phenotype induction subthreshold. Dauer larval arrest was scored as described in Material and Methods. The total number of worms scored for each genotype and RNAi condition are in parentheses: Wild type Control RNAi (N=469); Wild type-*ftt-1* RNAi (N=130); Wild type-*ftt-2* RNAi (N=475); *daf-2(e1370)*-Control RNAi (N=1219); *daf-2(e1370)-ftt-1* RNAi (N=480); *daf-2(e1370)-ftt-2* RNAi (N=1052); *daf-2(e1365)*-Control RNAi (N=1138); *daf-2(e1365)-ftt-1* RNAi (N=621); *daf-2(e1365)-ftt-2* RNAi (N=1235). Results presented are mean values +/- S.E.M of at least five independent experiments. Statistical student's *t*-test was used to compare datas from control and *ftt* RNAi fed worms and give either \**P*<0.05 and \*\**P*<0.0005 threshold signification level.

**Figure 7. Reducing *ftt-2* by RNAi during development leads to apparent sickness at subsequent post-developmental stage.**

Worms of indicated genotype were allowed to hatch on bacteria carryings either *ftt-1*, *ftt-2* or control RNAi. Worm population that did not arrest as dauers was fed on bacteria carrying either control RNAi vector or *ftt-1* or *ftt-2* RNAi until they reached young adult stage then transferred on new plates and fed by classical OP50 *E. coli* strain then scored for life span. Shortening life span in all of genetic backgrounds was also characterised by sickness defects such bagging or protruding vulvae phenotype. See text for details and Table 2 for corresponding quantitative and statistical datas analysis. All of the experiments were conducted at 22.5°C.

**Figure 8. *ftt-2* can positively modulate lipid droplets reserves.**

(A-L) Worms of indicated genotype were allowed to hatch on bacteria carrying either control RNAi vector, *ftt-1* or *ftt-2* RNAi at 22.5°C together with Nile Red lipid staining until young adult stage. Pictures were taken at 40X magnification for 200ms by fluorescence microscopy as described in Material and Methods. (M) Lipids droplets quantification reveals a highly significant reduction of lipid content in *daf-16(mgDf50)* loss-of-function mutant upon *ftt-2* RNAi knock down only. Quantifications of Nile Red lipid staining were done by fluorescence microscopy as described in Material and Methods on at least ten animals. Bars represent the standard error (SEM) and mean values are expressed after normalisation with respective control RNAi, which represent 100%. Statistical Student's *t*-test comparison between *ftt* RNAi and respective control where \**P* value < 0.05 and \*\**P* value < 0.005, give the following results for *ftt-2*: \* *P* = 0.02 for Wild type; \* *P* = 0.03 for *daf-2(e1370)*; \* *P* = 0.009 for *daf-2(e1365)*; \**P* = 0.05 for *daf-16(mgDf50)* and for *ftt-1*: *P* = 0.26 for Wild type; \*\**P* = 0.0009 for *daf-2(e1370)*; *P* = 0.18 for *daf-2(e1365)*; *P* = 0.77 for *daf-16(mgDf50)*.

**Figure 9. A model depicting the involvement of 14-3-3 in DAF-16-dependent and -independent mechanisms regulating normal life span and acute stress response in *C. elegans*.**

We propose a two component model in which 14-3-3 can modulate (A) no stress and (B) stress dependent physiological responses.



(A) In non-stress conditions, 14-3-3 is permanently at work to modulate physiological traits in a DAF-2-dependent manner and we show that PDK1 is a target for 14-3-3 regulation. We demonstrate in this work that 14-3-3 can also modulate physiological traits by acting independently of DAF-16.







(B) In stress conditions, we showed that 14-3-3 promotes oxidative stress resistance in a DAF-16-independent manner. On the contrary, our work demonstrates that 14-3-3 can modulate thermotolerance by a DAF-2/DAF-16-dependent mechanisms introducing a new side regulatory stress responsive pathway element to the one already described by Berdichevsky et al., 2006.


ACCEPTED MANUSCRIPT


Strain	RNAi	50th percentile (£)	Mean life span days (+ /- SD)	Max life span	code (Ü)	Statistics (KM analysis) (\$)	N (censored) (*)
N2 (wt)	control	19	18,9 (0,1)	28	A	p<0,0001 (B) p<0,0001 (C)	1024 (67)
	<i>ftt-1</i>	17	16,4 (0,1)	29	B	p<0,0001 (A) p<0,0001 (C)	1590 (28)
	<i>ftt-2</i>	16	15,6 (0,1)	31	C	p<0,0001 (A) p<0,0001 (B)	1501 (49)
<i>daf-16</i> ( <i>mgDf50</i> )	control	14	13,9 (0,1)	24	D	p = 0,089 (E) p<0,0001 (F)	806 (24)
	<i>ftt-1</i>	14	14,0 (0,3)	29	E	p = 0,089 (D) p<0,0001 (F)	261 (0)
	<i>ftt-2</i>	13	12,8 (0,1)	20	F	p<0,0001 (D) p<0,0001 (E)	478 (36)
<i>daf-2</i> ( <i>e1365</i> )	control	34	31,3 (0,7)	42	G	p<0,0001 (H) p<0,0001 (J)	170 (17)
	<i>ftt-1</i>	26	25,4 (0,3)	38	H	p<0,0001 (G) p=0,623 (J)	380(18)
	<i>ftt-2</i>	23	23,4 (0,4)	38	J	p<0,0001 (G) p=0,623 (H)	432 (22)
<i>daf-2</i> ( <i>e1370</i> )	control	30	28,6 (0,4)	44	K	p<0,0001 (L) p<0,0001 (M)	277 (9)
	<i>ftt-1</i>	20	19,6 (0,2)	39	L	p<0,0001 (K) p<0,0001 (M)	953 (0)
	<i>ftt-2</i>	19	18,5 (0,1)	39	M	p<0,0001 (K) p<0,0001 (L)	1272 (4)
<i>pdk-1</i> ( <i>sa709</i> )	control	21	21,5 (0,5)	42	N	p<0,0001 (O) p=0,796 (P)	110 (58)
	<i>ftt-1</i>	24	24,7 (0,4)	45	O	p<0,0001 (N) p<0,0001 (P)	186 (74)
	<i>ftt-2</i>	21	21,5 (0,4)	43	P	p=0,796 (N) p<0,0001 (O)	184 (72)
<i>pdk-1</i> ( <i>mg142</i> )	control	20	19,1 (0,2)	26	Q	p=0,039 (R) p<0,0001 (S)	546 (136)
	<i>ftt-1</i>	20	19,5 (0,2)	28	R	p=0,039 (Q) p<0,0001 (S)	783 (75)
	<i>ftt-2</i>	17	17,6 (0,2)	31	S	p<0,0001 (Q) p<0,0001 (R)	736 (75)

Strain	RNAi during development	Mean life span days (+/- SD) (@)	Max life span	code	Statistics (KM analysis) (\$)	N (censored) (*)
N2 (wt)	control	19,0 (0,2)	28	A	p<0,0001 (B) p<0,0001 (C)	561 (11)
	<i>ftt-1</i>	16,4 (0,3)	25	B	p<0,0001 (A) p=0,007 (C)	241 (50)
	<i>ftt-2</i>	14,2 (0,3)	25	C	p<0,0001 (A) p=0,007 (B)	384 (72)
<i>daf-16</i> ( <i>mgDf50</i> )	control	13,0 (0,1)	19	D	p<0,0001 (E) p<0,0001 (F)	359 (17)
	<i>ftt-1</i>	8,7 (0,2)	16	E	p<0,0001 (D) p=0,153 (F)	208 (20)
	<i>ftt-2</i>	7,3 (0,3)	18	F	p<0,0001 (D) p=0,153 (E)	318 (9)
<i>daf-2</i> ( <i>e1365</i> )	control	28,6 (0,4)	40	J	p<0,0001 (K) p<0,0001 (L)	342 (6)
	<i>ftt-1</i>	12,5 (0,2)	35	K	p<0,0001 (J) p= 0,614 (L)	620 (95)
	<i>ftt-2</i>	11,4 (0,6)	33	L	p<0,0001 (J) p= 0,614 (K)	348 (0)
<i>daf-2</i> ( <i>e1370</i> )	control	26,1 (0,3)	39	G	p<0,0001 (H) p<0,0001 (I)	494 (13)
	<i>ftt-1</i>	12,0 (0,2)	24	H	p<0,0001 (G) p<0,0001 (I)	425 (68)
	<i>ftt-2</i>	8,4 (0,4)	32	I	p<0,0001 (G) p<0,0001 (H)	435 (0)






















(M)













M




## No Stress (A)



- Development
- Lipid accumulation
- Antioxidant and Chaperone genes (SOD-3, HSP-16)
- Life span
- Dauer

- Development
- Lipid accumulation
- Oxidative stress resistance (GST-4)
- Life span

## Stress (B)



- Oxidative stress resistance (Menadione)

- Thermotolerance
- Antioxidant genes
- Chaperone genes
- Longevity