



HAL
open science

On Hofmann's bilinear estimate

Pascal Auscher

► **To cite this version:**

| Pascal Auscher. On Hofmann's bilinear estimate. 2008. hal-00347845v2

HAL Id: hal-00347845

<https://hal.science/hal-00347845v2>

Preprint submitted on 19 Dec 2008 (v2), last revised 16 May 2009 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON HOFMANN'S BILINEAR ESTIMATE

PASCAL AUSCHER

Abstract. We generalize a result of S. Hofmann to systems, using the framework of a previous paper with Axelsson and McIntosh.

MSC classes: 35J25, 35J55, 47N20, 42B25

Keywords: elliptic systems; Dirichlet problem; quadratic estimates; Carleson measures

1. Introduction

S. Hofmann proved in [10] that weak solutions of

$$(1) \quad \operatorname{div}_{t,x} A(x) r_{t,x} U(t;x) = \sum_{i,j=0}^{X^n} \partial_i A_{i,j}(x) \partial_j U(t;x) = 0$$

on the upper half space $\mathbb{R}_+^{1+n} := \{(t;x) \in \mathbb{R} \times \mathbb{R}^n; t > 0, n \geq 1\}$, where the matrix $A = (A_{i,j}(x))_{i,j=0}^n \in L_1(\mathbb{R}^n; L(C^{1+n}))$ is assumed to be t -independent and within some L_1 neighborhood of a real symmetric strictly elliptic t -independent matrix, obey the following bilinear estimate

$$\iint_{\mathbb{R}_+^{1+n}} r_{t,x} U \overline{v} dt dx \leq C \|U\|_2 \|v\|_2$$

The trace of U at $t=0$ is assumed to be in the sense of non-tangential convergence a.e. and in $L_2(\mathbb{R}^n)$. See below for the definition of the square function $k_j k_j$ and the non-tangential maximal operator N . In addition, he proves that the solution operator $U_0 : U(t; \cdot) \rightarrow U(t; \cdot)$ defines a bounded C^* -semigroup on $L_2(\mathbb{R}^n)$ whose infinitesimal generator A has domain in $W^{1,2}(\mathbb{R}^n)$ with $\|k A f k_2 \leq \|k r f k_2$. This generalizes results of B. Dahlberg [7] corresponding to the case where $\operatorname{div} A r$ is the pullback of the Laplace operator on a special Lipschitz domain. Hofmann's results utilize the deep results of [1] and in particular Theorem 1.11 where the boundedness and invertibility of the layer potentials are obtained from a T(b) theorem, Rellich estimates in the case of real symmetric matrices and perturbation. This also generalizes somehow the case where $A_{0,i} = A_{i,0} = 0$ for $i = 1, \dots, n$ corresponding to the Kato square root problem.

The recent papers [3, 4] allow us to extend this further to systems, making clear in particular that specificities of real symmetric coefficients and of equations are not needed: it only depends on whether the Dirichlet problem is solvable. We use the solution operator constructed in [3] and this also makes transparent the para-product like character of this bilinear estimate. We also state a necessary and sufficient condition telling when the domain of the infinitesimal generator A is $W^{1,2}$.

2. Setting

We begin by giving the precise definition of well-posedness of the BVPs for system s . Throughout this note, we use the notation $X \approx Y$ and $X \lesssim Y$ for estimates to mean that there exists a constant $C > 0$, independent of the variables in the estimate, such that $X = C Y \leq C X$ and $X \leq C Y$, respectively.

We write $(t; x)$ for the standard coordinates for $\mathbb{R}^{1+n} = \mathbb{R} \times \mathbb{R}^n$, t standing for the vertical or normal coordinate. For vectors $v = (v_i)_{0 \leq i \leq n}^1 \in \mathbb{C}^{(1+n)m}$, we write $v_0 \in \mathbb{C}^m$ and $v_k \in \mathbb{C}^{nm}$ for the normal and tangential parts of v , i.e. $v_0 = (v_0)^1 \dots^m$ whereas $v_k = (v_i)_{1 \leq i \leq n}^1 \dots^m$.

For system s , gradient and divergence act as $(r_{t;x} U)_i = \partial_i U$ and $(\operatorname{div}_{t;x} F) = \sum_{i=0}^n \partial_i F_i$, with corresponding tangential versions $r_x U = (r_{t;x} U)_k$ and $(\operatorname{div}_x F) = \sum_{i=1}^n \partial_i F_i$. With $\operatorname{curl}_x F_k = 0$, we understand $\partial_j F_i = \partial_i F_j$, for all $i, j = 1; \dots; n; = 1; \dots; m$.

We consider divergence form second order elliptic systems

$$(2) \quad \sum_{i,j=0}^n \partial_i A_{i;j}^i(x) \partial_j U(t;x) = 0; \quad i,j=0 \dots n$$

on the half space $\mathbb{R}_+^{1+n} = \{f(t;x) \in \mathbb{R} \times \mathbb{R}^n; t > 0\}$, $n \geq 1$, where the matrix $A = (A_{i;j}^i(x))_{i,j=0; \dots; n}^{i=1; \dots; m} \in L_1(\mathbb{R}^n; L(\mathbb{C}^{(1+n)m}))$ is assumed to be t -independent with complex coefficients and strictly accretive on $N(\operatorname{curl}_x)$, in the sense that there exists $\delta > 0$ such that

$$(3) \quad \sum_{i,j=0}^n \operatorname{Re}(A_{i;j}^i(x) f_j(x) \overline{f_1(x)}) dx \geq \sum_{i=0}^n \int_{\mathbb{R}^n} f_1(x)^2 dx;$$

for all $f \in N(\operatorname{curl}_x) = \{fg \in L_2(\mathbb{R}^n; \mathbb{C}^{(1+n)m}); \operatorname{curl}_x(g_k) = 0\}$. By changing m to $2m$ we could assume that the coefficients are real-valued. But this does not simplify matters and we need the complex hermitian structure of our L_2 space anyway.

Definition 2.1. The Dirichlet problem (Dir-A) is said to be well-posed if for each $u \in L_2(\mathbb{R}^n; \mathbb{C}^m)$, there is a unique function

$$U_t(x) = U(t;x) \in C^1(\mathbb{R}_+; L_2(\mathbb{R}^n; \mathbb{C}^m))$$

such that $r_x U \in C^0(\mathbb{R}_+; L_2(\mathbb{R}^n; \mathbb{C}^{nm}))$, where U satisfies (2) for $t > 0$, $\lim_{t \downarrow 0} U_t = u$, $\lim_{t \downarrow 1} U_t = 0$, $\lim_{t \downarrow 1} r_{t;x} U_t = 0$ in L_2 norm, and $\int_{t_0}^{t_1} r_x U_s ds$ converges in L_2^1 when $t_0 \downarrow 0$ and $t_1 \downarrow 1$. More precisely, by U satisfying (2), we mean that $\int_t^1 (A r_{s;x} U_s)_k; r_x v ds = (A r_{t;x} U_t)_0; v$ for all $v \in C_0^1(\mathbb{R}^n; \mathbb{C}^m)$.

Restricting to real symmetric equations and their perturbations, this definition is not the one taken in [10]. However, a sufficient condition is provided in [3] to insure that the two methods give rise to the same solution. See also [1, Corollary 4.28]. It covers the matrices listed in Theorem 2.4 below. This definition is more akin to well-posedness for a Neumann problem (see Section 4).

Remark 2.2. In the case of block matrices, i.e. $A_{0;i}^i(x) = 0 = A_{i;0}^i(x)$, $1 \leq i \leq n; 1 \leq j \leq m$, the second order system (2) can be solved using semi-group theory: $V(t; \cdot) = e^{tL} u_0$ for $L = A_{00}^{-1} \operatorname{div}_x A_{kk} r_x$ acting as an unbounded operator on $L_2(\mathbb{R}^n; \mathbb{C}^{nm})$ (See below for the notation). This solution satisfies $V_t = V(t; \cdot) \in C^2(\mathbb{R}_+; L_2(\mathbb{R}^n; \mathbb{C}^m)) \setminus C^1(\mathbb{R}_+; D(L^{1/2}))$; $\lim_{t \downarrow 0} V_t = u_0$, $\lim_{t \downarrow 1} V_t = 0$ in L_2 norm,

and (2) holds in the strong sense in \mathbb{R}^n for all $t > 0$ (and in the sense of distributions in \mathbb{R}_+^{1+n}). Hence, the two notions of solvability are not a priori equivalent. That the solutions are the same follows indeed from the solution of the Kato square root problem for $L : D(L^{1/2}) = W^{1,2}(\mathbb{R}^n; \mathbb{C}^{nm})$ with $kL^{1/2}fk_2 = kr_xfk_2$. See [6] where this is explicitly proved when $A_{00} \in I$.

The following result is Corollary 3.4 of [3] (which, as we recall, furnishes a different proof of results obtained by combining [11] and [8] in the case of real symmetric matrices equations ($m = 1$)).

Theorem 2.3. Let $A \in L_1(\mathbb{R}^n; L(\mathbb{C}^{(1+n)m}))$ be a t -independent, complex matrix function which is strictly accretive on $N(\text{curl}_k)$ and assume that $(\text{Dir}-A)$ is well-posed. Then any function $U_t(x) \in C^1(\mathbb{R}_+; L_2(\mathbb{R}^n; \mathbb{C}^m))$ solving (2), with properties as in Definition 2.1, has estimates

$$\int_{\mathbb{R}^n} |u|^2 dx \leq \sup_{t > 0} \int_{\mathbb{R}^n} |U_t|^2 dx \leq \int_{\mathbb{R}^n} |\mathbb{N}(U)|^2 dx \leq k \int_{\mathbb{R}_+^{1+n}} |U|^2 dx;$$

where $u = U|_{\mathbb{R}^n}$. If furthermore A is real (not necessarily symmetric) and $m = 1$, then Moser's local boundedness estimate [14] gives the pointwise estimate $\mathbb{N}(U)(x) \leq N(U)(x)$, where the standard non-tangential maximal function is $N(U)(x) := \sup_{|y-x| < ct} \int_{\mathbb{R}^n} |U(t;y)| dy$ for fixed $0 < c < 1$.

We use the square function norm

$$kF_t k^2 := \int_0^t kF_s k^2 \frac{ds}{s} = \int_{\mathbb{R}_+^{1+n}} |F(t;x)|^2 \frac{dtdx}{t}$$

and the norm $k\mathbb{N}(F)k_2$, using the modified non-tangential maximal function

$$\mathbb{N}(F)(x) := \sup_{t > 0} t^{(1+n)/2} kF|_{Q(t;x)} k_{L_2(Q(t;x))};$$

where $Q(t;x) := [(1-c_0)t; (1+c_0)t] \times B(x; c_1 t)$, for some fixed constants $c_0 \in (0;1)$, $c_1 > 0$.

Next is Theorem 3.2 of [3], specialized to the Dirichlet problem.

Theorem 2.4. The set of matrices A for which $(\text{Dir}-A)$ is well-posed is an open subset of $L_1(\mathbb{R}^n; L(\mathbb{C}^{(1+n)m}))$. Furthermore, it contains

- (i) all Hermitian matrices $A(x) = A(x)^*$ (and in particular all real symmetric matrices),
- (ii) all block matrices where $A_{0i}^j(x) = 0 = A_{i0}^j(x)$, $1 \leq i \leq n; 1 \leq j \leq m$, and
- (iii) all constant matrices $A(x) = A$.

More importantly is the solution algorithm using an "infiniteesimal generator" T_A . Write $v \in C^{(1+n)m}$ as $v = [v_0; v_k]^T$, where $v_0 \in C^m$ and $v_k \in C^{nm}$, and introduce the auxiliary matrices

$$\bar{A} := \begin{pmatrix} A_{00} & A_{0k} \\ 0 & I \end{pmatrix}; \quad \underline{A} := \begin{pmatrix} 1 & 0 \\ A_{k0} & A_{kk} \end{pmatrix}; \quad \text{if } A = \begin{pmatrix} A_{00} & A_{0k} \\ A_{k0} & A_{kk} \end{pmatrix}$$

in the normal/tangential splitting of $C^{(1+n)m}$. The strict accretivity of A on $N(\text{curl}_k)$, as in (3), implies the pointwise strict accretivity of the diagonal block A_{00} . Hence

A_{00} is invertible, and consequently \overline{A} is invertible [This is not necessarily true for \underline{A} .] We define

$$T_A = \overline{A}^{-1} D \underline{A}$$

as an unbounded operator on $L_2(\mathbb{R}^n; \mathbb{C}^{(1+n)m})$ with D the first order self-adjoint operator given in the normal/tangential splitting by

$$D = \begin{pmatrix} 0 & \operatorname{div}_x \\ r_x & 0 \end{pmatrix} :$$

Proposition 2.5. Let $A \in L_1(\mathbb{R}^n; L(\mathbb{C}^{(1+n)m}))$ be a t -independent, complex matrix function which is strictly accretive on $N(\operatorname{curl}_k)$.

- (1) The operator T_A has quadratic estimates and a bounded holomorphic functional calculus on $L_2(\mathbb{R}^n; \mathbb{C}^{(1+n)m})$. In particular, for any holomorphic function ϕ on the left and right open half planes, with $z \mapsto \phi(z)$ and $z \mapsto \phi(z^{-1})$ qualitatively bounded, one has

$$\| \phi(T_A) f \|_2 \leq \| \phi \|_2 \| f \|_2 :$$

- (2) The Dirichlet problem $(\operatorname{Dir}-A)$ is well-posed if and only if the operator

$$S : \overline{R(\cdot + (T_A))} \rightarrow L_2(\mathbb{R}^n; \mathbb{C}^m); f \mapsto f_0$$

is invertible. Here, $\chi_+ = 1$ on the right open half plane and 0 on the left open half plane.

Item (1) is [3, Corollary 3.6] (and see [4] for an explicit direct proof) and item (2) can be found in [3, Section 4, proof of Theorem 2.2].

Lemma 2.6. Assume that $(\operatorname{Dir}-A)$ is well-posed. Let $u_0 \in L_2(\mathbb{R}^n; \mathbb{C}^m)$. Then the solution U of $(\operatorname{Dir}-A)$ in the sense of definition 2.1 is given by

$$U(t; \cdot) = (e^{tT_A} f)_0; \quad f = S^{-1} u_0 \in \overline{R(\cdot + (T_A))}$$

and furthermore

$$r_{t,x} U(t; \cdot) = \mathcal{G} e^{tT_A} f :$$

Proof. [3, Lemma 4.2] (See also [2, Lemma 2.55] with a slightly different formulation of the Dirichlet problem).

3. The bilinear estimate

We are now in position to state and prove the generalisation of Hormann's result.

Theorem 3.1. Assume that $(\operatorname{Dir}-A)$ is well-posed. Let $u_0 \in L_2(\mathbb{R}^n; \mathbb{C}^m)$ and U be the solution to $(\operatorname{Dir}-A)$ in the sense of definition 2.1. Then for all $v : \mathbb{R}_+^{1+n} \rightarrow \mathbb{C}^{(1+n)m}$ such that the right hand side is finite,

$$\int_{\mathbb{R}_+^{1+n}} \int_{\mathbb{R}_+^{1+n}} r_{t,x} U(t; \cdot) \overline{v} dt dx \leq C \| u_0 \|_2 (\| r_{t,x} v \|_2 + \| N v \|_2) :$$

The pointwise values of $v(t; x)$ in the non-tangential control $N v$ can be slightly improved to L^1 averages on balls having radii $\geq t$ for each fixed t . See the end of proof.

Proof. It follows from the previous result that there exists $f \in \overline{R(\cdot)(T_A)}$ such that $U(t; \cdot) = (e^{tT_A} f)_0$ and

$$r_{t,x} U(t; \cdot) = \mathcal{F} = T_A e^{tT_A} f; \quad F = e^{tT_A} f:$$

Integrating by parts with respect to t , we find

$$\int_{\mathbb{R}_+^{1+n}} r U \overline{v} \, dt dx = \int_{\mathbb{R}_+^{1+n}} t \partial_t F \overline{v} \, dt dx - \int_{\mathbb{R}_+^{1+n}} t \partial_t^2 F \overline{v} \, dt dx:$$

The boundary term vanishes because $t \partial_t F$ goes to 0 in L_2 when $t \rightarrow 0; 1$ (this uses $f \in \overline{R(\cdot)(T_A)}$) and $\sup_{t > 0} kv(t; \cdot)k < 1$ from $kN \cdot vk_2 < 1$.

For the first term, we use Cauchy-Schwarz inequality and that $kT_A F k \leq k_0 k_2$ from Theorem 2.3.

For the second term, we use the following identity: $T_A = \overline{A}^{-1} D B \overline{A}$ with $B = \overline{A} \overline{A}^{-1}$, which by [3, Proposition 3.2] is strictly accretive on $N(\text{curl}_k)$, and

$$\begin{aligned} t^2 \partial_t^2 F &= \overline{A}^{-1} (t D B)^2 e^{tD B} \overline{A} f \\ &= \overline{A}^{-1} (t D B) (I + (t D B)^2)^{-1} (t D B) \overline{A} f \\ &= \overline{A}^{-1} (t D B) (I + (t D B)^2)^{-1} \overline{A} (t T_A) (f) \end{aligned}$$

with

$$(z) = z(1 + z^2) e^{(\text{sgn} \text{Re } z)z}:$$

Thus,

$$\int_{\mathbb{R}_+^{1+n}} t \partial_t^2 F \overline{v} \, dt dx = \int_{\mathbb{R}_+^{1+n}} \overline{A}^{-1} (t T_A) (f) \overline{Q_t v_t} \, dt dx$$

with $Q_t = \overline{A}^{-1}$ and $v_t = (t D B) (I + (t D B)^2)^{-1}$ acting on $v_t = v(t; \cdot)$ for each fixed t . [The notation \overline{A} has nothing to do with complex conjugate and we apologize for any conflict this may cause.] It follows from the quadratic estimates proved in Proposition 2.5 that

$$k_j (t T_A) (f) k_j \leq k f k_2:$$

It remains to estimate $k Q_t v_t k_j$. To do that we follow the principal part approximation of [4] applied to Q_t instead of Δ_t there. That is, we write

$$(4) \quad Q_t v_t = Q_t \frac{I - P_t}{t(\cdot)^{1=2}} \Delta_t^{-1=2} v_t + (Q_t P_t - \Delta_t S_t P_t) v_t + \Delta_t S_t P_t v_t$$

where Δ_t is the Laplacian on \mathbb{R}^n , P_t is a nice scalar approximation to the identity acting componentwise on $L_2(\mathbb{R}^n; C^{(1+n)m})$ and Δ_t is the element of $L_{loc}^2(\mathbb{R}^n; L(C^{(1+n)m}))$ given by

$$\Delta_t(x) w := (Q_t w)(x)$$

for every $w \in C^{(1+n)m}$. We view w on the right-hand side of the above equation as the constant function valued in $C^{(1+n)m}$ defined on \mathbb{R}^n by $w(x) := w$. We identify $\Delta_t(x)$ with the (possibly unbounded) multiplication operator $\Delta_t : f(x) \mapsto \Delta_t(x) f(x)$. Finally, the dyadic averaging operator $S_t : L_2(\mathbb{R}^n; C^{(1+n)m}) \rightarrow L_2(\mathbb{R}^n; C^{(1+n)m})$ is given by

$$S_t u(x) := \frac{1}{|Q|} \int_Q u(y) \, dy$$

for every $x \in \mathbb{R}^n$ and $t > 0$, where Q is the unique dyadic cube in \mathbb{R}^n that contains x and has side length ℓ with $\ell = 2^{-j} t^{-1}$.

With this in hand, we apply the triple bar norm to (4).

Using the uniform L_2 boundedness of Q_t and that of $\frac{1}{t} P_t$, the first term in the RHS is bounded by $C \| \cdot \|_{L_2}^{1+2} \| \cdot \|_{L_2} \| \cdot \|_{L_2}$.

Following exactly the computation of Lemma 3.6 in [4], the second term in the RHS is bounded by $C \| \cdot \|_{L_2} \| \cdot \|_{L_2} \| \cdot \|_{L_2}$ using the uniform L_2 boundedness of P_t . This computation makes use of the off-diagonal estimates of Q_t , hence of Q_t , proved in [4, Proposition 3.11].

For the third term in the RHS, we observe that $Q_t(x)w = \overline{Q_t(A^{-1}w)}(x)$. Hence, the square function estimate on Q_t proved in [4, Theorem 1.1], the off-diagonal estimates of Q_t and the fact that $\overline{A^{-1}}$ is bounded imply that $Q_t(x) \int \frac{dtdx}{t}$ is a Carleson measure. Hence, from Carleson embedding theorem the third term contributes to $C \| \cdot \|_{L_2} \| \cdot \|_{L_2}$, which is controlled pointwise by the non-tangential maximal function in the statement with appropriate opening.

4. The domain of the Dirichlet semigroup

Assume $(D_{\text{Dir}}A)$ in the sense of definition 2.1 is well-posed. If we set

$$P_t u_0 = (e^{-tA} f)_0; \quad f = S^{-1} u_0 \in \overline{R(\cdot, T_A)}$$

for all $t > 0$, then Lemma 2.6 implies that $(P_t)_{t>0}$ is a bounded C_0 -semigroup on $L_2(\mathbb{R}^n; \mathbb{C}^m)$ [recall that well-posedness includes uniqueness and this allows to prove the semigroup property].

Furthermore, with our definition of the Dirichlet problem, the domain of the infinitesimal generator A of this semigroup is contained the Sobolev space $W^{1,2}(\mathbb{R}^n; \mathbb{C}^m)$ and $\| \cdot \|_{L_2} \| \cdot \|_{L_2} \leq \| \cdot \|_{L_2} \| \cdot \|_{L_2}$. Indeed, from Lemma 2.6 we have for all $t > 0$, $\partial_t e^{-tA} f = r_{t,x} U(t; \cdot)$. Also $e^{-tA} f \in \overline{R(\cdot, T_A)}$ and the invertibility of S tells that $r_{t,x} U(t; \cdot) = S^{-1}(\partial_t U(t; \cdot))$. Therefore

$$\| r_{t,x} U(t; \cdot) \|_{L_2} \leq \| \partial_t U(t; \cdot) \|_{L_2}$$

By definition of A , $\partial_t U(t; \cdot) = A U(t; \cdot)$, thus we have for all $t > 0$

$$\| r_{t,x} U(t; \cdot) \|_{L_2} \leq \| A U(t; \cdot) \|_{L_2}$$

The conclusion for the domain follows easily.

The question of whether this domain coincides with $W^{1,2}(\mathbb{R}^n; \mathbb{C}^m)$ is answered by the following theorem

Theorem 4.1. Assume that $(D_{\text{Dir}}A)$ and $(D_{\text{Dir}}A)$ are well-posed. Then the domain of the infinitesimal generator A of $(P_t)_{t>0}$ coincides with the Sobolev space $W^{1,2}(\mathbb{R}^n; \mathbb{C}^m)$ and $\| \cdot \|_{L_2} \| \cdot \|_{L_2} \leq \| \cdot \|_{L_2} \| \cdot \|_{L_2}$.

This theorem applies to the three situations listed in Theorem 2.4.

Proof. Combining [4, Lemma 4.2] (which says that $(D_{\text{Dir}}A)$ is equivalent to an auxiliary Neumann problem for A), [2, Proposition 2.52] (which says that this auxiliary Neumann problem is equivalent to a regularity problem for A : this is non-trivial) with the proof of Theorem 2.2 in [4] (giving the necessary and sufficient condition below for well-posedness of the regularity problem for A), we have that $(D_{\text{Dir}}A)$ is well-posed if and only if

$$R : \overline{R(\cdot, T_A)} \rightarrow L_2(\mathbb{R}^n; \mathbb{C}^{nm}); f \mapsto f_x$$

is invertible. This implies that for $f \in \overline{R(\cdot + (T_A))}$, we have that

$$\|kf\|_{k_2} \leq \|kf_0\|_{k_2};$$

Therefore, the conjunction of well-posedness for $(D_{\text{Dir}} - A)$ and $(D_{\text{Dir}} - A)$ gives

$$\|kf_0\|_{k_2} \leq \|kf\|_{k_2}; \quad f \in \overline{R(\cdot + (T_A))}:$$

From this, it is easy to conclude for the domain of A by an argument as before.

We have seen that invertibility of S reduces to that of R (up to taking adjoints). The only known way to prove it in such a generality (except for constant coefficients) is via a continuity method and the Rellich estimates showing that $\|kf\|_{k_2} \leq \|k(Af)\|_{k_2}$ for all $f \in \overline{R(\cdot + (T_A))}$: This method was first used in the context of Laplace equation on Lipschitz domains by Verchota [15]. This depends strongly of A . Various relations between Dirichlet, regularity and Neumann problems in the sense of non tangential approach for second order real symmetric equations are studied in [12, 13].

References

- [1] Alfonseca, M., Auscher, P., Axelsson, A., Hofmann, S., and Kim, S. Analyticity of layer potentials and L^2 solvability of boundary value problems for divergence form elliptic equations with complex L^1 coefficients. Preprint at arXiv:0705.0836v1 [math.AP].
- [2] Auscher, P., Axelsson, A., and Hofmann, S. Functional calculus of Dirac operators and complex perturbations of Neumann and Dirichlet problems. *Journal of Functional Analysis*, 255, 2 (2008), 374-448.
- [3] Auscher, P., Axelsson, A., and McIntosh, A. Solvability of elliptic systems with square integrable boundary data, preprint Preprint arXiv:0809.4968v1 [math.AP].
- [4] Auscher, P., Axelsson, A., and McIntosh, A. On a quadratic estimate related to the Kato conjecture and boundary value problems. Preprint arXiv:0810.3071v1 [math.CA].
- [5] Auscher, P., Hofmann, S., Lacey, M., McIntosh, A., and Tchamitchian, P. The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n . *Ann. of Math. (2)* 156, 2 (2002), 633-654.
- [6] Axelsson, A., Keith, S., and McIntosh, A. Quadratic estimates and functional calculi of perturbed Dirac operators. *Invent. Math.* 163, 3 (2006), 455-497.
- [7] Dahlberg, B. Poisson semigroups and singular integrals, *Proc. Amer. Math. Soc.* 97 (1986), 41-48.
- [8] Dahlberg, B., Jerison, D., and Kenig, C.E. Area integral estimates for elliptic differential operators with nonsmooth coefficients. *Ark. Mat.* 22, 1 (1984), 97-108.
- [9] Fabes, E., Jerison, D., and Kenig, C.E. Necessary and sufficient conditions for absolute continuity of elliptic-harmonic measure. *Ann. of Math. (2)* 119, 1 (1984), 121-141.
- [10] Hofmann, S. Dahlberg's bilinear estimate for solutions of divergence form complex elliptic equations. *Proc. Amer. Math. Soc.* 136 (2008), no. 12, 4223-4233.
- [11] Jerison, D.S., and Kenig, C.E. The Dirichlet problem in nonsmooth domains. *Ann. of Math. (2)* 113, 2 (1981), 367-382.
- [12] Kenig, C., and Pipher, J. The Neumann problem for elliptic equations with nonsmooth coefficients. *Invent. Math.* 113, 3 (1993), 447-509.
- [13] Kenig, C., and Pipher, J. The Neumann problem for elliptic equations with nonsmooth coefficients. Part II *Duke Math. J.* 81 1 (1995), 227-250.
- [14] Moser, J. On Hamack's theorem for elliptic differential equations. *Comm. Pure Appl. Math.* 14 (1961), 577-591.
- [15] Verchota, G. Layer potentials and regularity for the Dirichlet problem for Laplace's equation in Lipschitz domains. *J. Funct. Anal.* 59, 3 (1984), 572-611.

Pascal Auscher, Université de Paris-Sud, UMR du CNRS 8628, 91405 Orsay Cedex, France

E-mail address: pascal.auscher@math.u-psud.fr