



HAL
open science

On the homology of the space of singular knots

Hossein Abbaspour, David Chataur

► **To cite this version:**

Hossein Abbaspour, David Chataur. On the homology of the space of singular knots. 2008. hal-00348675v1

HAL Id: hal-00348675

<https://hal.science/hal-00348675v1>

Preprint submitted on 19 Dec 2008 (v1), last revised 10 Mar 2009 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE HOMOLOGY OF THE SPACE OF SINGULAR KNOTS

HOSSEIN ABBASPOUR AND DAVID CHATAUR

ABSTRACT. In this paper we introduce various associative products on the homology of the space of knots and singular knots in S^n . We prove that these products are related through a desingularization map. We also compute some of these products and prove the nontriviality of the desingularization morphism.

CONTENTS

1. Introduction	1
2. Long immersions and long knots	3
3. Stringy operations	4
4. Comparison with Chas-Sullivan product	6
5. Desingularization morphism	7
6. Some computations	10
References	15

1. INTRODUCTION

Various authors have tried to introduce a general framework for the structures introduced by Chas-Sullivan [CS1] and Cohen-Jones [CJ] on the homology and on the equivariant homology of the free loop space of a closed manifold. One approach, first used in [KS] and then formulated in [GS], is the notion of fiber-wise (homotopy) monoid. We use this formulation to introduce and calculate various algebra structures on the homology of knots, immersions, and singular knots with k double points.

In this paper the *knot space* $\text{Emb}(S^1, S^n)$ is the space of all embeddings of S^1 in the n -sphere S^n , and similarly $\text{Imm}(S^1, S^n)$ is the space of all *immersions* of S^1 in S^n . For a knot or an immersion $\gamma : S^1 \rightarrow S^n$, the point $\gamma(1)$ is called

Key words and phrases. knots, singular knots, long knots, free loops space, string operations.

The second author is partially supported by ANR grant 06-JCJC-0042 “Opérades, Bigèbres et Théories d’Homotopie”.

the *marked point* of γ . Here S^n is the unit sphere in \mathbb{R}^{n+1} and we make the identification $\mathbb{R}^n \simeq \{0\} \times \mathbb{R}^n \subset \mathbb{R}^{n+1}$. We think of $SO(n) \subset SO(n+1)$ as the stabilizer of $(1, 0, \dots, 0)$ and $SO(n-1) \subset SO(n+1)$ as the stabilizer of $(1, 0, \dots, 0)$ and $(0, 1, 0, \dots, 0)$. US^n is the unit tangent sphere bundle of S^n which can be identified with $SO(n+1)/SO(n-1)$. Let $LUS^n = C^\infty(S^1, US^n)$ be the free loop space of S^n .

The subspace $\text{Imm}'(S^1, S^n) \subset \text{Imm}(S^1, S^n)$ consists of those immersions which are not singular at the marked point and $\text{Imm}_k(S^1, S^n) \subset \text{Imm}'(S^1, S^n)$ is the space of immersions with exactly k double *points*.

We will be considering *uncompactified* or *long* version of the objects defined above which roughly corresponds to the one dimensional object obtained in \mathbb{R}^n using the *stereographic* map with the marked point as the center point.

For instance, the space of *long knots*¹ in \mathbb{R}^n is

$$\text{Emb}^l(\mathbb{R}, \mathbb{R}^n) = \{f : \mathbb{R} \times \{0\} \rightarrow \mathbb{R}^n, \text{supp}(f) \subset (-1, 1) \text{ and } f([-1, 1] \times \{0\}^{n-1}) \subset B^n\}$$

where $\text{supp}(f) = \{t \in \mathbb{R} \subset \mathbb{R}^n \mid f(t) \neq (t, 0, \dots, 0)\}$ and B^n is the unit ball. Similarly one defines $\text{Imm}^l(\mathbb{R}, \mathbb{R}^n)$ the space of long immersions and $\text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)$ the space of long immersion with k singular double points.

The purpose of this paper is to introduce various associative algebra structures on the homology of $\text{Emb}(S^1, S^n)$, $\text{Imm}'(S^1, S^n)$ and $\text{Imm}(S^1, S^n)$, and finally singular knot with double singular points $\coprod_k \text{Imm}'_k(S^1, S^n)$. In the case of singular knots, one naturally expects a compatibility with the Vassiliev spectral sequences. However this issue is not addressed in this paper. In order to introduce these products, for each space we consider a homotopically equivalent model as the total space of a fibration whose fibres form a continuous family of homotopy associative monoids and the base space is a closed oriented manifold. This should be compared with the loop space fibration $\Omega M \rightarrow LM \rightarrow M$ of a closed oriented manifold.

Theorem 3.1. $(H_{*+2n-1}(\text{Emb}(S^1, S^n)), \mu_{em}(-, -))$ is a graded commutative algebra.

Theorem 3.2. $(H_{*+2n-1}(\text{Imm}'(S^1, S^n)), \mu_{im}(-, -))$ is a graded commutative algebra.

The commutativity of the product follows from the computation of the homology by spectral sequences arguments.

Theorem 4.2. $H_{*+2n-1}(\text{Imm}(S^1, S^n))$ is a graded commutative algebra and the map induced by inclusion $i_* : H_{*+2n-1}(\text{Imm}'(S^1, S^n)) \rightarrow H_{*+2n-1}(\text{Imm}(S^1, S^n))$ is a map of algebras.

After restricting to knots with double singular points, one obtains:

¹There are some other models for the space of long knots, all homotopy equivalent to ours.

Theorem 4.3. *There is a collection of maps*

$$\mu_{im}^{k,l} : H_*(\text{Imm}'_k(S^1, S^n)) \times H_*(\text{Imm}'_l(S^1, S^n)) \rightarrow H_{*-2n+1}(\text{Imm}'_{k+l}(S^1, S^n))$$

such that

$$\mu_{im}^{k+l,m}(\mu_{im}^{k,l}(a, b), c) = \mu_{im}^{k,l+m}(a, \mu_{im}^{l,m}(b, c)).$$

In order to compare the algebra structure on the homologies of knot and singular knot spaces we introduce a desingularization map. Informally speaking, we resolve a singular knot at a double point in all possible ways, parameterized by unit vectors perpendicular to the tangent plane at the singularity. Of course this map has a certain degree and it is not a map of algebras. However it verifies a certain compatibility condition with respect to the number of singularities and the product. More precisely,

Theorem 5.1. *The desingularization morphisms $\sigma_k : H_*(\text{Imm}'_k(S^1, S^n)) \rightarrow H_{*-k(n-3)}(\text{Emb}(S^1, S^n))$, $k \geq 0$, are compatible with the products i.e. for $x \in H_*(\text{Imm}'_k(S^1, S^n))$ and $y \in H_*(\text{Imm}'_l(S^1, S^n))$*

$$\mu_{em}(\sigma_k(x), \sigma_l(y)) = \sigma_{k+l}(\mu_{im}^{k,l}(x, y))$$

$\text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$	Long knots in \mathbb{R}^n
$\text{Emb}(S^1, S^n)$	Embeddings in S^n (Knots)
$\text{Imm}^l(\mathbb{R}, \mathbb{R}^n)$	Long immersions in \mathbb{R}^n
$\text{Imm}'(S^1, S^n)$	Immersions in S^n without singularity at $t = 0$
$\text{Imm}'_k(\mathbb{R}, \mathbb{R}^n)$	Long immersions in \mathbb{R}^n with k double points
$\text{Imm}'_k(S^1, S^n)$	Immersions in S^n with k double points away from $t = 0$

TABLE 1. Notation

Acknowledgments: The first author would like to thank the Hausdorff Research Institute and the MPI in Bonn for its support while this paper was written up. The second author warmly thanks the Laboratoire Jean Leray at the University of Nantes for their invitation through the Matpyl program.

2. LONG IMMERSIONS AND LONG KNOTS

We start this section with a few remarks on the homotopy type of the space of knots $\text{Emb}(S^1, S^n)$ and immersions $\text{Imm}'(S^1, S^n)$, in terms of the long knots $\text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$ and long immersions $\text{Imm}^l(\mathbb{R}, \mathbb{R}^n)$. The main references are [B1, B2, BC].

Note that $SO(n-1)$ acts on $SO(n+1) \times \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$, by the natural action on the first factor. It acts on the second factor via the identification $\mathbb{R} \times \mathbb{R}^{n-1} =$

$\mathbb{R}^n \simeq \{0\} \times \mathbb{R}^n \subset \mathbb{R}^{n+1}$. One may form the quotient $SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$ which has the same homotopy type of $\text{Emb}(S^1, S^n)$ (see [B2]). A homotopy equivalence is given by the stereographic map with $(1, 0, \dots, 0) \in S^n \subset \mathbb{R}^{n+1} = \mathbb{R} \times \mathbb{R}^n$ as the center. Given a pair $(A, \gamma) \in SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$ one can compactify γ in \mathbb{R}^{n+1} to get a knot in $S^n \subset \mathbb{R}^{n+1}$ passing through $(1, 0, \dots, 0)$. Then, we let A act on γ to get an element of $\text{Emb}(S^1, S^n)$. This defines a homotopy equivalence map

$$(2.1) \quad \rho_{emb} : SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n) \rightarrow \text{Emb}(S^1, S^n)$$

which makes the following diagram to commute.

$$(2.2) \quad \begin{array}{ccc} SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n) & \xrightarrow{\rho_{emb}} & \text{Emb}(S^1, S^n) \\ \downarrow & & \downarrow ev \\ SO(n+1)/SO(n-1) & \longrightarrow & US^n \end{array}$$

Here $ev(\gamma) = (\gamma(0), \gamma'(0)/\|\gamma'(0)\|)$ for $\gamma \in \text{Emb}(S^1, S^n)$. Therefore, we have the isomorphism,

$$(2.3) \quad \rho_{em} : H_*(SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)) \rightarrow H_*(\text{Emb}(S^1, S^n)).$$

Similarly, one can consider the Borel construction $SO(n+1) \times_{SO(n-1)} \text{Imm}^l(\mathbb{R}, \mathbb{R}^n)$ and $SO(n+1) \times_{SO(n-1)} \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)$ the homotopy equivalences induced by the stereographic map which gives rise to the commutative diagram

$$(2.4) \quad \begin{array}{ccc} SO(n+1) \times_{SO(n-1)} \text{Imm}^l(\mathbb{R}, \mathbb{R}^n) & \xrightarrow{\rho_{im}} & \text{Imm}^l(S^1, S^n) \\ \downarrow & & \downarrow ev \\ SO(n+1)/SO(n-1) & \longrightarrow & US^n \end{array}$$

Similarly, one has the isomorphisms,

$$(2.5) \quad \rho_{im} : H_*(SO(n+1) \times_{SO(n-1)} \text{Imm}^l(\mathbb{R}, \mathbb{R}^n)) \rightarrow H_*(\text{Imm}^l(S^1, S^n))$$

$$(2.6) \quad \rho_{im}^k : H_*(SO(n+1) \times_{SO(n-1)} \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)) \rightarrow H_*(\text{Imm}_k^l(S^1, S^n))$$

3. STRINGY OPERATIONS

In this section we recall the general framework where some of the string topology operations can be defined. This was observed by several authors independently, including the authors of this article. However the main published reference is [GS], therefore we follow their approach. Let $F \rightarrow E \xrightarrow{\pi} M$ be a fiber bundle over a closed compact manifold M of dimension d . We suppose

that E is equipped with a fibrewise associative product, that is a fiber bundle map $m : E \times_B E \rightarrow E$ such that

$$(3.1) \quad m(x, m(y, z)) = m(m(x, y), z)$$

Since the associativity of the product is not achieved for most of the interesting cases, one must adapt this definition to the case where the product is homotopy associative and consider *fiberwise homotopy monoids*.

One can take one step further by acquiring an operadic approach in the monoidal category of spaces fibered over M . This is simply done by introducing the trivial bundles $\bar{\mathcal{C}}_n = \mathcal{C}_n \times M \rightarrow M$ where \mathcal{C}_n is the space of n little cubes (see [GS]). Then, one may consider a fiber bundle $E \rightarrow M$ which is an algebra over the bundle operad $\bar{\mathcal{C}} = \coprod_n \bar{\mathcal{C}}_n$ *i.e.* we have a collection of fibre bundle maps $E \times \bar{\mathcal{C}}_n \rightarrow E$ subject to the usual axioms for the algebras over an operad.

This formulation allows us to define a commutative and associative product of degree $-d$

$$\bullet : H_*(E) \times H_*(E) \rightarrow H_{*-d}(E)$$

and a bracket of degree $(1 - d)$

$$\{-, -\} : H_*(E) \times H_*(E) \rightarrow H_{*+1-d}(E)$$

which satisfies the Jacobi identity for a Lie bracket of degree $1 - d$. One could be more ambitious and consider algebras over the bundle operad of framed little cube operad in order to equip $H_*(E)$ with a BV-algebra structure. This feature is not discussed in this paper.

Applying this machinery to the fibration

$$\text{Emb}(S^1, S^n) \simeq SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n) \rightarrow SO(n+1)/SO(n-1),$$

one obtains immediately a commutative and associative product of degree $-2n+1$

$$(3.2) \quad \mu_{em}(-, -) = \rho_{em}(\rho_{em}^{-1}(-) \bullet \rho_{em}^{-1}(-)) : H_*(\text{Emb}(S^1, S^n)) \times H_*(\text{Emb}(S^1, S^n)) \rightarrow H_{*-2n+1}(\text{Emb}(S^1, S^n))$$

Theorem 3.1. (*Gruher-Salvatore, ...*) $(H_*(\text{Emb}(S^1, S^n)), \mu_{em})$ is a commutative and associative algebra.

This product is also introduced in Gruher and Salvatore's paper for the case $n = 3$ based on the R. Budney's work [B1]. Note that for all $n \geq 3$, the little cube operad acts on the space of long knots $\text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$, therefore this product is well-defined for all $n \geq 3$.

Similarly, one can consider the fibration

$$\mathrm{Imm}'(S^1, S^n) \simeq SO(n+1) \times_{SO(n-1)} \mathrm{Imm}'(\mathbb{R}, \mathbb{R}^n) \rightarrow SO(n+1)/SO(n-1).$$

which accommodates a fiberwise homotopy associative product but not the action of the full little cube bundle operad. For instance, we lose the homotopy commutativity of the product. In the case of embeddings, the product at the level of fibers is basically the connected sum of long knots.

(3.3)

$$\mu_{im}(-, -) = \rho_{im}(\rho_{im}^{-1}(-) \bullet \rho_{im}^{-1}(-)) : H_*(\mathrm{Imm}'(S^1, S^n)) \times H_*(\mathrm{Imm}'(S^1, S^n)) \rightarrow H_{*-2n+1}(\mathrm{Imm}'(S^1, S^n))$$

Theorem 3.2. $H_{*+2n-1}(\mathrm{Imm}'(S^1, S^n))$ equipped with μ_{im} is a graded associative algebra.

After restricting to immersions with k and l double points, it reads

(3.4)

$$\mu_{im}^{k,l} = \rho_{im}^{k+l}(\rho_{im}^{k-1}(-) \bullet \rho_{im}^{l-1}(-)) : H_*(\mathrm{Imm}'_k(S^1, S^n)) \times H_*(\mathrm{Imm}'_l(S^1, S^n)) \rightarrow H_{*-2n+1}(\mathrm{Imm}'_{k+l}(S^1, S^n))$$

Remark 3.3. We observe that all these theorems are corollaries of Gruher-Salvatore and Budney's work, we therefore don't propose a detailed proof.

4. COMPARISON WITH CHAS-SULLIVAN PRODUCT

So far we have only define the associative product on $H_*(\mathrm{Imm}'(S^1, S^n))$ and not on $H_*(\mathrm{Imm}(S^1, S^n))$. The algebra structure of the latter will be discussed in the last section where we prove Theorem 4.2 cited in the introduction. What is discussed in this section is the comparison of the algebra structure on $H_*(\mathrm{Imm}'(S^1, S^n))$ with that of $H_*(LUS^n)$ equipped with Chas-Sullivan loop product. The 1-jet map $\Psi : \gamma \mapsto (\gamma, \gamma'/\|\gamma'\|)$ is a map of the fiberwise homotopy monoids making the following diagrams to commute

$$(4.1) \quad \begin{array}{ccc} \mathrm{Imm}'(S^1, S^n) & \xrightarrow{\Psi} & LUS^n \\ \downarrow ev & & \downarrow ev_0 \\ US^n & \xrightarrow{id} & US^n \end{array}$$

and

$$(4.2) \quad \begin{array}{ccc} SO(n+1) \times_{SO(n-1)} \text{Imm}^l(\mathbb{R}, \mathbb{R}^n) & \xrightarrow{\Psi \circ \rho_{im}} & LUS^n \\ \downarrow & & \downarrow ev_0 \\ US^n & \xrightarrow{id} & US^n \end{array}$$

Recall that $ev(\gamma) = (\gamma(0), \gamma'(0)/\|\gamma'(0)\|)$ for $\gamma \in \text{Emb}(S^1, S^n)$ and $ev_0(\alpha) = \alpha(0)$ for $\alpha \in LUS^n$.

An immediate consequence of Proposition 11 in [GS] is

Theorem 4.1. *The map induced by 1-jet map $(H_*(\text{Imm}^l(S^1, S^n)), \mu_{im}) \rightarrow (H_*(LUS^n), \bullet)$ is a map of graded associative algebras. Here, \bullet is the Chas-Sullivan loop product.*

Using Hirsch-Smale theorem, one can state the theorem above differently.

Theorem (Hirsch-Smale) The 1-jet map $\text{Imm}(S^1, S^n) \rightarrow LUS^n$, given by $\gamma \mapsto (\gamma, \gamma'/\|\gamma'\|)$, is a homotopy equivalence.

One can transfer the associative commutative product from $H_*(LUS^n)$ to $H_*(\text{Imm}(S^1, S^n))$, and then Theorem 4.1 reads

Theorem 4.2. *$H_{*+2n-1}(\text{Imm}(S^1, S^n))$ is a graded associative algebra and the map induced by inclusion $H_{*+2n-1}(\text{Imm}^l(S^1, S^n)) \rightarrow H_{*+2n-1}(\text{Imm}(S^1, S^n))$ is a map of algebras.*

5. DESINGULARIZATION MORPHISM

We first introduce \mathfrak{Q}_k^n the *desingularization spaces* and the *desingularization morphism* $\sigma_k : \mathfrak{Q}_k^n \rightarrow \text{Emb}(\mathbb{R}, \mathbb{R}^n)$. One should think of \mathfrak{Q}_k^n as the space of k -singular knots decorated with k tangent vectors at the singular points, to be used to desingularize the knot (see [CCL]).

Let $G_{2,n} = SO(n)/SO(2) \times SO(n-2)$ be the Grassmanian manifold of oriented 2-plane in \mathbb{R}^n . One has the canonical fibration $Q_n \rightarrow G_{2,n}$ whose fibers are S^{n-3} . Here Q_n is the space of pairs (π, v) where π is a 2-plane and v is a vector perpendicular to π . More precisely,

$$Q_n = SO(n) \times_{(SO(2) \times SO(n-2))} S^{n-3}$$

where the action of $SO(2) \times SO(n-2)$ is trivial for the first factor and for the second factor is given by the natural action of $SO(n-2)$ on $S^{n-3} \subset \mathbb{R}^{n-2}$. Then $Q_n^{\times k}$ fibers over $G_{2,n}^{\times k}$ through the map $(\pi_i, v_i)_{1 \leq i \leq k} \mapsto (\pi_i)_{1 \leq i \leq k}$.

Then we define \mathfrak{Q}_n^k to be the total space of the pull back of the fibration $Q_n^{\times k} \rightarrow G_{2,n}^{\times k}$ via the natural map $r_k : \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n) \rightarrow G_{2,n}^{\times k}$, given by sending an immersion with k singular double points to the k 2-planes define at the singularities by the tangent vectors. We have the following commutative diagram of

fibrations,

$$(5.1) \quad \begin{array}{ccc} \mathfrak{Q}_n^k & \longrightarrow & Q_n^{\times k} \\ \downarrow & & \downarrow \\ \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n) & \xrightarrow{r_k} & G_{2,n}^{\times k} \end{array}$$

where the fibres are $(S^{n-3})^{\times k}$.

The *desingularization morphism* $\sigma : \coprod_{k \geq 1} \mathfrak{Q}_n^k \rightarrow \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$ is defined by resolving the singularities using the normal vectors given at those points. More explicitly, let $(\gamma, (v_i)_{1 \leq i \leq k}) \in \mathfrak{Q}_k$ where $\gamma : \mathbb{R} \rightarrow \mathbb{R}^n$ is a long immersion with double singular points $\gamma(t_1^1) = \gamma(t_2^1), \gamma(t_1^2) = \gamma(t_2^2), \dots, \gamma(t_1^k) = \gamma(t_2^k)$ and v_1, \dots, v_k are the vectors normal to the tangent plane at the singularities. Consider the bump functions,

$$(5.2) \quad \alpha_i(t) = \begin{cases} 0 & \text{if } t \notin [t_{a_i}^i - \epsilon, t_{a_i}^i + \epsilon] \\ (-1)^{a_i} \delta \exp\left(\frac{-1}{(t-t_{a_i}^i)^2 - \epsilon^2}\right) v_i & \text{if } t \in [t_{a_i}^i - \epsilon, t_{a_i}^i + \epsilon] \end{cases}$$

where each a_i is 1 or 2. Let the embedding $\sigma((\gamma, (v_i)_{i=1}^k)) : \mathbb{R} \rightarrow \mathbb{R}^n$ be

$$(5.3) \quad \sigma_k^n((\gamma, (v_i)_{i=1}^k))(t) = \gamma(t) + \sum_{i=1}^k \alpha_i(t).$$

To define σ_k^n one has to make a choice for a_i , but it turns out that these choices give rise to homologous cycles. Moreover, our construction obviously depends on good choices of ϵ and δ . One has to make sure that each t_1^i and t_2^i are at least ϵ away from each other and δ is small enough not to create any new singularity. To resolve this problem, one considers the space \mathfrak{Q}'_k^n consisting of triples $[(\gamma, (v_i)_{1 \leq i \leq k}), \epsilon, \delta]$ for which the construction works. One has the fibration $\mathfrak{Q}'_k^n \rightarrow \mathfrak{Q}_k^n$, given by projecting on the first factor $(\gamma, (v_i)_{1 \leq i \leq k})$, whose fibers are contractible. Therefore \mathfrak{Q}'_k^n and \mathfrak{Q}_k^n have the same homotopy type. Note that both \mathfrak{Q}'_k^n and \mathfrak{Q}_k^n thought of long immersions decorated with the some normal vectors at the singularities have a natural homotopy associative product.

Thus we have,

Lemma 5.1. (See [CCL]) *Desingularization morphism $\sigma_k^n : \mathfrak{Q}_k^n \rightarrow \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$ is homologically well-defined and by the abuse of notation we have*

$$\sigma_k^n : H_*(\mathfrak{Q}_k^n) \rightarrow H_*(\text{Emb}^l(\mathbb{R}, \mathbb{R}^n)).$$

which is a map of associative algebras with respect to the product induced on the homologies by the concatenation products on \mathfrak{Q}_k^n and $\text{Emb}^l(\mathbb{R}, \mathbb{R}^n)$.

Moreover, σ_k^n is clearly $SO(n-1)$ equivariant, therefore it induces a homologically well-defined map

$$\Sigma_k^n : H_*(SO(n+1) \times_{SO(n-1)} \mathfrak{Q}_k^n) \rightarrow H_*(SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)).$$

We have a map

$$(5.4) \quad \begin{array}{ccc} \coprod_k \mathfrak{Q}_k^n & \xrightarrow{\coprod_k \sigma_k^n} & \text{Emb}^l(\mathbb{R}, \mathbb{R}^n) \\ \downarrow & & \downarrow \\ \coprod_k SO(n+1) \times_{SO(n-1)} \mathfrak{Q}_k^n & \xrightarrow{\coprod_k \Sigma_k^n} & SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n) \\ \downarrow & & \downarrow \\ SO(n+1)/SO(n-1) & \xrightarrow{id} & SO(n+1)/SO(n-1) \end{array}$$

of fiberwise monoids, hence:

Lemma 5.2.

$\oplus_k \Sigma_k^n : (\oplus_k H_*(SO(n+1) \times_{SO(n-1)} \mathfrak{Q}_k^n), *) \rightarrow (H_*(SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)), \mu_{im})$
is map of associative algebras.

We need another map in order to complete the construction of the desingularization morphism from the homology of the space of immersions in S^n to the homology of knot space.

Proposition 5.3. (See [BG]) *Let $F \rightarrow E \rightarrow M$ be an oriented fibration whose fiber has the homotopy type of a closed manifold of dimension p . Then there is a natural map $f! : H_*(M) \rightarrow H_{*+p}(E)$. This is the dual of the map on cohomology given by integration over fibres.*

A description using spectral sequence is the following composite

$$E_{p,n}^2 = H_p(M, \mathcal{H}_n(F)) = H_p(M, \mathbb{F}) \hookrightarrow E_{p,n}^\infty \Rightarrow H_{p+n}(E)$$

Here $\mathcal{H}_q(F)$ stands for the local coefficient system which is trivial in this case and has the fundamental class of F as a generator (because the fibration is oriented). In this description one can see clearly the naturality property.

Applying this construction to the oriented fibration $\mathfrak{Q}_k^n \rightarrow \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)$ and $SO(n+1) \times_{SO(n-1)} \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n) \rightarrow SO(n+1) \times_{SO(n-1)} \mathfrak{Q}_k^n$

Proposition 5.4. *The fibration $\mathfrak{Q}_k^n \rightarrow \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)$ gives rises to maps*

$$\theta_n^k : H_*(\text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)) \rightarrow H_{*+k(n-3)}(\mathfrak{Q}_k^n)$$

and

$$\Theta_n^k : H_*(SO(n+1) \times_{SO(n-1)} \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)) \rightarrow H_{*+k(n-3)}(SO(n+1) \times_{SO(n-1)} \mathfrak{Q}_k^n)$$

It follows immediately from the spectral sequence description of the Gysin maps which can easily accomodate the multiplicative structure of the homologies involved.

Lemma 5.5. *The Gysin maps Θ_k^n 's are multiplicative in the sense that*

$$\Theta_k^n(x) * \Theta_l^n(y) = \Theta_{k+l}^n(\mu_{im}(x, y)).$$

Then the composition

$$(5.5) \quad \begin{aligned} \Sigma_k^n \circ \Theta_k^n : H_*(SO(n+1) \times_{SO(n-1)} \text{Imm}_k^l(\mathbb{R}, \mathbb{R}^n)) \rightarrow \\ H_{*-k(n-3)}(SO(n+1) \times_{SO(n-1)} \text{Emb}^l(\mathbb{R}, \mathbb{R}^n)) \end{aligned}$$

defines the desingularization morphism

$$(5.6) \quad \phi_k^n = \Sigma_k^n \circ \Theta_k^n : H_*(\text{Imm}_k(S^1, S^n)) \rightarrow H_{*-k(n-3)}(\text{Emb}(S^1, S^n))$$

Theorem 5.6. *The desingularization morphisms $\phi_k^n : H_*(\text{Imm}'_k(S^1, S^n)) \rightarrow H_{*-k(n-3)}(\text{Emb}(S^1, S^n))$, $k \geq 0$, are compatible with the products i.e. for $x \in \mathbb{H}_*(\text{Imm}'_k(S^1, S^n))$ and $y \in H_*(\text{Imm}'_l(S^1, S^n))$*

$$\mu_{em}(\phi_k^n(x), \phi_l^n(y)) = \phi_{k+l}^n(\mu_{im}^{k,l}(x, y))$$

Proof. For $x, y \in H_*(\text{Imm}'_k(S^1, S^n))$

$$(5.7) \quad \mu_{em}(\phi_k^n(x), \phi_l^n(y)) = \mu_{em}(\Sigma_k^n \circ \Theta_k^n(x), \Sigma_l^n \circ \Theta_l^n(y)) = \Sigma_{k+l}^n(\Theta_k^n(x) * \Theta_l^n(y))$$

by Lemma 5.2. The latter is $\Sigma_{k+l}^n \circ \Theta_{k+l}^n(\mu_{im}^{k,l}(x, y)) = \phi_{k+l}^n(\mu_{im}^{k,l}(x, y))$. □

6. SOME COMPUTATIONS

Let us consider the canonical inclusion $i : \text{Imm}'(S^1, S^n) \rightarrow \text{Imm}(S^1, S^n)$, the aim of this section is twofold

first we determine the morphism i_* induced in singular homology

secondly we prove that the desingularization morphism is non-trivial.

Immersion in \mathbb{R}^{n+1} . As a baby example let us first consider the case of long immersions $\text{Imm}^l(\mathbb{R}, \mathbb{R}^{n+1})$ and of singular knots $\text{Imm}(S^1, \mathbb{R}^{n+1})$ into an Euclidean space. Thanks to the Hirsch-Smale's theorem one knows that the maps, $\gamma \mapsto D\gamma$ and $\gamma \mapsto (\gamma, D\gamma)$ give homotopy equivalences between these spaces and the spaces ΩS^n and $LU\mathbb{R}^{n+1}$, where $D(\gamma) := \frac{\dot{\gamma}}{\|\dot{\gamma}\|}$.

$$(6.1) \quad \begin{array}{ccc} \text{Imm}^l(\mathbb{R}, \mathbb{R}^{n+1}) & \xrightarrow{D} & \Omega S^n \\ \downarrow & & \downarrow \\ \text{Imm}(S^1, \mathbb{R}^{n+1}) & \xrightarrow{(id, D)} & LU\mathbb{R}^{n+1} = L\mathbb{R}^{n+1} \times LS^n S^n \\ & \searrow D_0 & \downarrow ev_0 \\ & & S^n \end{array}$$

We identify $U\mathbb{R}^{n+1}$ together with $\mathbb{R}^{n+1} \times S^n$, and we consider the canonical projection $pr_2 : U\mathbb{R}^{n+1} \rightarrow S^n$. Thus the spaces $\text{Imm}^l(\mathbb{R}, \mathbb{R}^{n+1})$ and $\text{Imm}(S^1, \mathbb{R}^{n+1})$ are homotopy equivalent to ΩS^n and LS^n .

The concatenation of long immersions in $\text{Imm}^l(\mathbb{R}, \mathbb{R}^{n+1})$ obviously corresponds to the composition of based loops in ΩS^n , we recall that the singular homology of these spaces together with this multiplicative structure is isomorphic to the tensor algebra generated by an element u of degree $n - 1$. The evaluation map

$$D_0 : \text{Imm}(S^1, \mathbb{R}^{n+1}) \xrightarrow{D} LS^n \xrightarrow{ev_0} S^{n+1}$$

can be used to endow the singular homology of $\text{Imm}(S^1, \mathbb{R}^{n+1})$ with a product. We consider the diagram

$$(6.2) \quad \begin{array}{ccc} \text{Imm}(S^1, \mathbb{R}^{n+1}) \times \text{Imm}(S^1, \mathbb{R}^{n+1}) & \xleftarrow{j} & \text{Imm}(S^1, \mathbb{R}^{n+1}) \times_{S^n} \text{Imm}(S^1, \mathbb{R}^{n+1}) \\ & & \downarrow Tr \\ & & \text{Imm}_{(0, \dots, 0)}(\infty, \mathbb{R}^{n+1}) \end{array}$$

where $\text{Imm}(S^1, \mathbb{R}^{n+1}) \times_{S^n} \text{Imm}(S^1, \mathbb{R}^{n+1})$ is the space of immersions (γ_1, γ_2) such that $D_0(\gamma_1) = D_0(\gamma_2)$, the space $\text{Imm}_{(0, \dots, 0)}(\infty, \mathbb{R}^{n+1})$ is the subspace of $\text{Imm}(S^1, \mathbb{R}^{n+1}) \times_{S^n} \text{Imm}(S^1, \mathbb{R}^{n+1})$ such that $\gamma_1(0) = \gamma_2(0) = (0, \dots, 0)$. The map Tr is a translation map, it is defined by using the unique translation T_1 that sends $\gamma_1(0)$ to $(0, \dots, 0)$ and the unique translation T_2 that sends $\gamma_2(0)$ to $(0, \dots, 0)$, $Tr(\gamma_1, \gamma_2) = (T_1(\gamma_1), T_2(\gamma_2))$. We also have a composition map

$$comp : \text{Imm}_{(0, \dots, 0)}(\infty, \mathbb{R}^{n+1}) \rightarrow \text{Imm}(S^1, \mathbb{R}^{n+1}).$$

As j is a finite codimensional embedding we can define a Gysin morphism $j_!$ and a product :

$$- \bullet - := comp_* \circ Tr_* \circ j_! \circ - \times -.$$

This multiplicative structure is isomorphic to the Chas-Sullivan product on $H_{*+n}(LS^n)$, the isomorphism is induced by the map D .

The algebra $H_{*+n}(LS^n)$ has been computed by Cohen, Jones and Yan [CJY]. For a closed oriented manifold M they consider the Serre spectral sequence associated to the evaluation fibration

$$\Omega M \rightarrow LM \rightarrow M$$

enriched it with a multiplicative structure. Using the Pontryagin product on $H_*(\Omega M)$ and the intersection product on $\mathbb{H}_*(M) = H_{*+n}(M)$ the second page of this spectral sequence can be endowed with a product and

$$E_{p,q}^2 := \mathbb{H}_p(M, H_q(\Omega M)) \Rightarrow \mathbb{H}_{p+q}(LM) := H_{*+n}(LM).$$

For the spheres, they get

$$\mathbb{H}_*(\text{Imm}(S^1, \mathbb{R}^{n+1})) \cong \mathbb{H}_*(LS^n) \cong \begin{cases} \Lambda(a) \otimes \mathbb{Z}[u] & \text{for } n \text{ odd} \\ (\Lambda(b) \otimes \mathbb{Z}[a, v]) / (a^2, ab, 2av) & \text{for } n \text{ even} \end{cases}$$

with $a \in \mathbb{H}_{-n}(LS^n) \cong E_{-n,0}^\infty$, $b \in \mathbb{H}_{-1}(LS^n) \cong E_{-n,n-1}^\infty$, $u \in \mathbb{H}_{n-1}(LS^n) \cong E_{0,n-1}^\infty$ and $v \in \mathbb{H}_{2n-2}(LS^n) \cong E_{0,2n-2}^\infty$.

Immersions in S^n . The computation of $\mathbb{H}_*(\text{Imm}(S^1, S^n))$ has been completed by Le Borgne and the second author in [CL]. In fact Le Borgne in [L] considers the loop fibration

$$LF \rightarrow LE \rightarrow LB$$

associated to a smooth fiber bundle of oriented closed manifolds

$$F \rightarrow E \rightarrow B$$

then he enriches the Serre spectral sequence with a multiplicative structure coming from the Chas-Sullivan loop product on LB , LE and LF :

$$E_{p,q}^2 := \mathbb{H}_p(LB, \mathbb{H}_q(LF)) \rightarrow \mathbb{H}_{p+q}(LE).$$

He applies this new technique to the fibration

$$LS^{n-1} \rightarrow LUS^n \rightarrow LS^n$$

and proves that in the case n is odd that this spectral sequence collapses at the E^2 -term. In the even case he proves that this is the Serre spectral sequence of

$$\Omega US^n \rightarrow LUS^n \rightarrow US^n$$

that collapses at the E^2 -term. In each case there are serious extension issues which are all solved in [CL] by using Morse theoretical techniques. We have the following isomorphisms of algebras :

Even case. When n is even

$$\mathbb{H}_*(\text{Imm}(S^1, S^n)) \cong \mathbb{H}_*(US^n, H_*(\Omega US^n))$$

we also recall that

$$\mathbb{H}_*(US^n) \cong \Lambda(a_{-n+1}, b_{-2n+1}) / (2a_{-n+1}, a_{-n+1}b_{-2n+1})$$

and that

$$H_*(\Omega US^n) \cong \mathbb{Z}[u_{n-2}, v_{2n-2}]/(2u_{n-2}).$$

The homology of US^n can be computed using the Serre spectral sequence of the fibration

$$S^{n-1} \rightarrow US^n \rightarrow S^n$$

and $H_*(\Omega US^n)$ using the Serre spectral sequence of

$$\Omega S^{n-1} \rightarrow \Omega US^n \rightarrow \Omega S^n.$$

Odd case. When n is odd we have

$$\mathbb{H}_*(\text{Imm}(S^1, S^n)) \cong \mathbb{H}_*(LS^n) \otimes \mathbb{H}_*(LS^{n-1}).$$

In order to determine $i_* : H_*(\text{Imm}'(S^1, S^n)) \rightarrow H_*(\text{Imm}(S^1, S^n))$, we consider the two fibrations

$$\text{Imm}'_*(S^1, S^n) \rightarrow \text{Imm}'(S^1, S^n) \rightarrow S^n$$

$$\text{Imm}'_{*,v}(S^1, S^n) \rightarrow \text{Imm}'(S^1, S^n) \rightarrow US^n,$$

where Imm'_* denotes the space of based immersions and $\text{Imm}'_{*,v}$ is its subspace consisting of those with a given unit tangent vector at the base point.

We notice that the second fibration admits a section *geod* this map sends the unit tangent vector $v \in US^n$ to the unique great circle such that $\dot{\gamma}(0) = v$.

Theorem 6.1. *For n even number then*

$$\mathbb{H}_*(\text{Imm}'(S^1, S^n)) \cong \mathbb{H}_*(US^n, H_*(\Omega S^{n-1})) \cong \mathbb{H}_*(US^n) \otimes H_*(\Omega S^{n-1}).$$

The morphism i_ is induced by the canonical morphism*

$$\mathbb{H}_*(US^n, H_*(\Omega S^{n-1})) \rightarrow \mathbb{H}_*(US^n, H_*(\Omega US^n)) \cong \mathbb{H}_*(\text{Imm}(S^1, S^n))$$

which is the identity on $\mathbb{H}_(US^n)$ and the morphism*

$$j_* : H_*(\Omega S^{n-1}) \rightarrow H_*(\Omega US^n)$$

given by the inclusion of a fibre $j : S^{n-1} \rightarrow US^n$. For n odd number then

$$\mathbb{H}_*(\text{Imm}'(S^1, S^n)) \cong \mathbb{H}_*(S^n) \otimes \mathbb{H}_*(LS^{n-1}),$$

in this case the morphism i_ is the injection*

$$\mathbb{H}_*(S^n) \otimes \mathbb{H}_*(LS^{n-1}) \xrightarrow{c_* \otimes Id} \mathbb{H}_*(LS^n) \otimes \mathbb{H}_*(LS^{n-1}),$$

where $c : S^n \hookrightarrow LS^n$ is the inclusion of the constant loops.

Proof. (the even case). Let us consider the Serre spectral sequence associated to the fibration

$$\mathrm{Imm}'_{*,v}(S^1, S^n) \rightarrow \mathrm{Imm}'(S^1, S^n) \rightarrow US^n.$$

The E^2 term of this spectral sequence is given by

$$E_{p,q}^2 = \mathbb{H}_p(US^n, \mathbb{H}_q(\mathrm{Imm}'_{*,v}(S^1, S^n)))$$

which is isomorphic to

$$\mathbb{H}_p(US^n) \otimes H_q(\Omega S^{n-1}).$$

Let us recall that in the even case we have

$$\mathbb{H}_*(US^n) \cong \Lambda(a_{-n+1}, b_{-2n+1}) / (2a_{-n+1}, a_{-n+1}b_{-2n+1})$$

and

$$H_*(\Omega S^{n-1}) \cong \mathbb{Z}[u_{n-2}].$$

In order to compute the other terms, we use the fact that this spectral sequence is multiplicative. As it has a section one has that the differentials $d_k(a_{-n+1}) = d_k(b_{-2n+1}) = 0$ for $k \geq 2$, and we also have $d_k(u_{n-2}) = 0$ for degree reasons. We conclude that this spectral sequence collapses at the E^2 term. Also for degree reasons there is no extensions issues thus we have that

$$\mathbb{H}_*(\mathrm{Imm}'(S^1, S^n)) \cong \Lambda(a_{-n+1}, b_{-2n+1}) / (2a_{-n+1}, a_{-n+1}b_{-2n+1}) \otimes \mathbb{Z}[u_{n-2}].$$

(The odd case). Let us consider the Serre-spectral sequence associated to the fibration

$$\mathrm{Imm}'_*(S^1, S^n) \rightarrow \mathrm{Imm}'(S^1, S^n) \rightarrow S^n$$

as in the even case for degree reasons this spectral sequence collapses at the E_2 -term and there is no extensions issues. \square

6.1. Non-triviality of the desingularization morphism. Let us prove that the desingularization morphism

$$\phi_n^2 : H_0(\mathrm{Imm}_2(S^1, S^n)) \rightarrow H_{2n-6}(\mathrm{Emb}(S^1, S^n))$$

is non-trivial. We suppose that $n > 4$ and we follow R. Budney's construction, in [B3] he gives a generator of $\pi_{2n-6}(\mathrm{Emb}^l(\mathbb{R}, \mathbb{R}^n)) \cong \mathbb{Z}$ which is the first non-trivial homotopy group. Let us describe this generator geometrically, let us take a long immersion γ together with two regular double points : we have $\gamma(t_1) = \gamma(t_3)$ and $\gamma(t_2) = \gamma(t_4)$ with $t_1 < t_2 < t_3 < t_4$, such that the resolution is the trefoil knot. Associated to this long immersion we have a desingularization map

$$\mathrm{des} : S^{n-3} \times S^{n-3} \rightarrow \mathrm{Emb}^l(\mathbb{R}, \mathbb{R}^n).$$

If s_{2n-6} is the fundamental class of $S^{n-3} \times S^{n-3}$ then in homology $\mathrm{des}_*(s_{2n-6})$ is the generator of $H_{2n-6}(\mathrm{Emb}^l(\mathbb{R}, \mathbb{R}^n))$.

Let us take a closure $\gamma' \in \mathrm{Imm}_2(S^1, S^n)$ of γ then by an easy spectral sequence

argument one can show that

1) $H_{2n-6}(\text{Emb}(S^1, S^n)) \cong \mathbb{Z}$ by using the Serre's spectral sequence of the fibration $\text{Emb}(S^1, S^n) \rightarrow US^n$ and the fact that this spectral has a section we have $H_{2n-6}(\text{Emb}(S^1, S^n)) \cong E_2^{0,2n-6}$.

2) If k_{2n-6} is a generator of $H_{2n-6}(\text{Emb}(S^1, S^n))$ then we have $\phi_n^2(1_{\text{tref}}) = k_{2n-6}$ where 1_{tref} is the generator the 0-th homology group of the connected component containing γ' .

REFERENCES

- [BG] J. C. Becker, D. H. Gottlieb, A history of duality in algebraic topology, History of topology, 725-745, North-Holland, Amsterdam, 1999.
- [Br] J-L. Brylinski, *Loop spaces, characteristic classes and geometric quantization*, Progress in Mathematics, **107**, Birkhäuser Boston Inc., Boston, MA, 1993.
- [BC] R. Budney and F. Cohen, On the homology of the space of knots, preprint ArXiv math.GT/0504206.
- [B1] R. Budney, *Little cubes and long knots*. Topology 46 (2007), no. 1, 1–27.
- [B2] R. Budney, *Topology of spaces of knots in dimension 3*, preprint ArXiv math.GT/0506524.
- [B3] R. Budney, *A family of embedding spaces*, Geometry and Topology Monographs 13 (2007).
- [CCL] A. Cattaneo, P. Cotta-Ramusino, R. Longoni, *Configuration spaces and Vassiliev classes in any dimension*. Algebr. Geom. Topol. 2 (2002), 949–1000
- [CL] D. Chataur, J.-F. Le Borgne, *Homology of spaces of regular loops in the sphere*, preprint ArXiv.math.AT/08113319
- [CJY] R.L. Cohen, J.D.S. Jones, J. Yan; The Loop-homology algebra of Spheres and Projective Spaces, Categorical Decomposition Techniques in Algebraic Topology. Progress in Mathematics 215 - Birkhauser-Verlag- 2004.
- [CS1] M. Chas and D. Sullivan, *String topology*, preprint ArXiv math.GT/9911159.
- [CJ] R. Cohen, J. Jones, *A homotopy theoretic realization of string topology*, Math. Ann. **324** (2002), no. 4, 773–798.
- [CV] R. Cohen, S. Voronov, *Notes on String Topology*, in “String Topology and Cyclic Homology”, *Adv. Courses in Math: CRM, Barcelona*, Birkhauser (2006), 1-95.
- [GS] K. Gruher and P. Salvatore, Generalized string topology operations, preprint ArXiv math.AT/0602210.
- [KS] S. Kallel and S. Salvatore, Rational maps and string topology, preprint arXiv math.AT/0309137.
- [L] J.F. Le Borgne, The loop-product spectral sequence, Expositiones Mathematicae, February 2008, pages 25-40.
- [S] P. Salvatore, Knots, operads and double loop spaces, preprint ArXiv math.AT/0608490.
- [Sm] S. Smale, The classification of immersions of spheres in Euclidean spaces, *Ann. of Math.* (2) **69** 1959 327–344.

LABORATOIRE JEAN LERAY, UNIVERSITÉ DE NANTES, NANTES 44300, FRANCE.
E-mail address: `abbaspour@univ-nantes.fr`

LABORATOIRE PAUL PAINLEVÉ, UNIVERSITÉ DE LILLE 1, 59655 VILLENEUVE D'ASCQ
CÉDEX, FRANCE

E-mail address: `David.Chataur@math.univ-lille1.fr`