

Spin Configuration in the 1/3 Magnetization Plateau of Azurite determined by NMR

F. Aimo, S. Kramer, M. Klanjsek, M. Horvatic, C. Berthier, H. Kikuchi

▶ To cite this version:

F. Aimo, S. Kramer, M. Klanjsek, M. Horvatic, C. Berthier, et al.. Spin Configuration in the 1/3 Magnetization Plateau of Azurite determined by NMR. 2008. hal-00350473v1

HAL Id: hal-00350473 https://hal.science/hal-00350473v1

Preprint submitted on 6 Jan 2009 (v1), last revised 24 Feb 2009 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spin Configuration in the 1/3 Magnetization Plateau of Azurite determined by NMR

F. Aimo,¹ S. Krämer,¹ M. Klanjšek,^{1,*} M. Horvatić,^{1,†} C. Berthier,¹ and H. Kikuchi²

¹Grenoble High Magnetic Field Laboratory, CNRS, BP 166, 38042 Grenoble Cedex 9, France

²Department of Applied Physics, University of Fukui, Fukui 910-8507, Japan

(Dated: January 6, 2009)

High magnetic field 63,65 Cu NMR spectra were used to determine the local spin polarization in the 1/3 magnetization plateau of Azurite, Cu₃(CO₃)₂(OH)₂, which is a model system for the distorted diamond antiferromagnetic spin-1/2 chain. The spin part of the hyperfine field of the Cu₂ (dimer) sites is found to be field independent, negative and strongly anisotropic, corresponding to $\approx 10\%$ of fully polarized spin in a *d*-orbital. This is close to the expected configuration of the "quantum" plateau, where a singlet state is stabilized on the dimer. However, the observed non-zero spin polarization points to some triplet admixture, induced by strong asymmetry of the diamond bonds J_1 and J_3 .

PACS numbers: 75.10.Pq, 75.25.+z, 67.80.dk, 76.60.-k

The natural mineral Azurite, $Cu_3(CO_3)_2(OH)_2$, has been recently recognized [1] as a model system for a frustrated antiferromagnetic Heisenberg spin-1/2 chain of "distorted diamond" geometry defined in Fig. 1. Its most prominent feature is a large plateau in the magnetization curve at 1/3 of the saturation magnetization, which extends from 11 to 30 T when the applied magnetic field (H_0) is perpendicular to the chains. Such a "1/3 plateau" is usually associated to a classical collinear up-up-down (uud) type of spin arrangement, or rather to a quantum state which has this classical analogue. For example, a uud state is predicted for spins on a two-dimensional triangular lattice and observed in the Cs_2CuBr_4 compound [2]. The 1/3 plateau in Azurite is proposed to be of fundamentally different, "00u" type, where the dominant J_2 coupling ensures that the two "dimer" spins on the Cu2 sites (see Fig. 1) are in a singlet state, while the third "monomer" (Cu1) spin is completely polarized by the field. As this state is based on the presence of a singlet, it is of pure quantum nature without a classical analogue. Azurite is a good candidate to be the first system exhibiting such a 1/3 plateau state, but a direct experimental evidence is still missing. The point is that both types of plateaus are predicted for a diamond chain, the 00u type driven by dominant J_2 coupling and an analogue of the *uud* state in presence of dominant J_1 and J_3 [3, 4]. The Azurite is close to the phase boundary between them, and there is a controversy on the J values proposed from the magnetization, specific heat and neutron measurements [1, 5, 6, 7]. The two different plateau types are distinguished by very different local spin polarizations, which can in principle be directly accessed by performing nuclear magnetic resonance (NMR) on the on-site copper 63,65 Cu nuclei. In this letter we present such NMR data which show that in the 1/3 plateau the dimer spins are nearly in the singlet configuration and thus confirm the 00u type of plateau. However, we find a small non-zero spin polarization of these sites, estimated to approximately 10% of full polarization, which points to an important asymmetry of J_1 and J_3 couplings.

In general, the copper NMR spectrum of a single crystal consists of 6 NMR lines per each non-equivalent Cu site, corresponding to three transitions between energy levels of a spin I = 3/2 nucleus for each of the two ⁶³Cu and ⁶⁵Cu isotopes. In the crystallographic structure of Azurite, shown in Fig. 1, we recognize two different copper sites in two equivalent chains of different orientation with respect to the arbitrary direction of the applied magnetic field. We also note that the two Cu2 sites of each dimer are expected to be undistinguishable by NMR, which is ensured by the inversion symmetry



FIG. 1: (color online) Diamond chains formed by the exchange interactions, $J_{ij}\mathbf{S}_i \cdot \mathbf{S}_j$, between S = 1/2 spins of Cu^{2+} ions in the crystal structure of Azurite. There are two equivalent but differently oriented chains, related by the *ac*-plane of mirror symmetry. Chains contain "monomer" spins on the Cu1 sites coupled by J_1 and J_3 interactions to each spin of the dimer formed by the two Cu2 sites, mutually coupled by J_2 . There is an inversion symmetry on each Cu1 site and at the center of each dimer. For each Cu site 4 nearest neighboring oxygen atoms (connected by thin lines) define approximately the plane of the local symmetry of the wave functions and of the corresponding EFG tensor. Dotted vectors define the angle ϑ between the magnetic field and the Z principal axis of the EFG tensor. C and H atoms are not shown.

with respect to the center of each dimer. We thus expect that 63,65 Cu NMR spectrum has $6 \times 2 \times 2 = 24$ NMR lines. Spectra presented in Fig. 2 contain only 12 lines, meaning that we observe only one of the two Cu sites. A standard way for the identification of the observed site is to compare the symmetry of the local electric field gradient (EFG) tensor determined by NMR to what is expected from the local symmetry of the four nearest neighboring (NN) oxygen atoms (see Fig. 1). This rather technical procedure, explained in detail in the following paragraph, unambiguously determines that only Cu2, i.e. the dimer site is observed by NMR.

Each copper isotope (I = 3/2) generates a triplet of NMR lines whose average frequency reflects the Zeeman coupling to the total "effective" magnetic field $\mathbf{H}_{\mathrm{eff}}$, while the line splitting is induced by the "quadrupolar" coupling to the local EFG. The corresponding nuclear spin Hamiltonian, $\mathcal{H} = \hbar \gamma \mathbf{I} \cdot \mathbf{H}_{eff} + h\nu_{Q}[3I_{Z}^{2} - I(I+1) + \eta(I_{+}^{2} +$ $[I_{-}^{2})/2]/6$, is uniquely defined by 5 parameters: the EFG tensor described by the quadrupolar coupling $\nu_{\rm Q}$ and its asymmetry parameter η , and H_{eff} and its direction $(\vartheta_{\rm EFG}, \varphi_{\rm EFG})$ with respect to the principal axes (X,Y,Z) of the EFG tensor [8]. Knowing the gyromagnetic ratios for the two isotopes, ${}^{63}\gamma$ and ${}^{65}\gamma$, as well as the ratio of their quadrupolar couplings, ${}^{63}\nu_Q/{}^{65}\nu_Q = 1.0805$, these 5 parameters can be fit to provide the observed 6 NMR frequencies (for each chain). The $\nu_{\rm Q}$ and η parameters do not depend on the orientation of magnetic field and, in particular, they are common to spectra from two chains shown in Fig. 1. In this experiment, several complete spectra have been taken at different rotation angles α around the axis that was close to the crystal a-axis, as shown in Fig. 2. The smallest NMR line widths and thus the most precise fits are obtained when $\vartheta_{\rm EFG} \cong 90^{\circ}$, which for chain 1 corresponds to $\alpha \cong 45^{\circ}$. This particular orientation has therefore been used to determine $^{63}\nu_{\rm Q} = 36.5 \,\mathrm{MHz}$ and $\eta = 0.085$, and the obtained EFG values have been successfully used to produce all the other fits - this time by fitting only three parameters $(H_{\rm eff}, \vartheta_{\rm EFG}, \varphi_{\rm EFG})$ for each set of 6 NMR frequencies (see Fig. 2). The experimentally obtained $\vartheta_{\rm EFG}(\alpha)$ dependence could then be compared to a simple approximate estimate for this quantity, based on the crystallographic structure. We know that for an ideal tetragonal coordination a pure $d_{X^2-Y^2}$ orbital pointing towards 4 NN oxygens generates axially symmetric EFG with the strongest principal axis along the Z direction. Therefore, the best estimate for the Z axis is the normal to the plane approximately defined by 4 NN oxygens (see Fig. 1). The direction of \mathbf{H}_{eff} is approximated by the direction of the applied field, supposing that the rotation axis is precisely the *a*-axis of the crystal. These two directions define $\vartheta_{\text{Theory}}$ whose rotation (α -) dependence for both Cu sites and both chains are plotted in Fig. 3, together with the experimental $\vartheta_{\rm EFG}$ values. Neglecting small offset due to various approximations, one clearly



FIG. 2: (color online) Rotation dependence of copper NMR spectra of Azurite at T = 1.5 K. Magnetic field of 15.0 T is applied perpendicular to the rotation axis that was close to the crystal *a*-axis. (a) NMR spectrum taken at 85° and the corresponding fit (vertical lines), as explained in the text. (b) Angular dependence of the observed line positions (crosses, with lines to guide the eye) and the corresponding fits (open symbols: circles for the central transitions and squares and diamonds for the satellites). Color code is given in the figure: dark (light) grey lines for the chain 1 (2) and solid (dashed) lines for the $^{63(65)}$ Cu isotope. Vertical dotted line denotes $H_0 \parallel ac$ -plane orientation, where both chains are identical.

identifies that the observed NMR signal corresponds to dimer Cu2 sites.

The principal information obtained from the fits is the spin part of the hyperfine field, $\mathbf{H}_{\rm spin} = \mathbf{H}_{\rm eff} - (1 + \mathsf{K}_{\rm orb}) \mathbf{H}_0 = \mathsf{A} g \mu_{\rm B} \langle \mathbf{S} \rangle$, induced by the local spin polarization $\langle \mathbf{S} \rangle$ through the hyperfine coupling tensor A. The orbital (Van Vleck) shift tensor $\mathsf{K}_{\rm orb}$ is here a minor correction, because typical A values for a copper spin are as large as $A_Z \approx -20 \,\mathrm{T}/\mu_{\rm B}$, with large anisotropy $A_{\rm Z}/A_{\perp} \approx 10$. Knowing that the EFG and the hyperfine shift tensors are dominantly determined by the same wave function, we expect that the principal axes of both tensors are approximately the same, so that the $H_{\rm spin}$ vs. $\vartheta_{\rm EFG}$ dependence provides a complete information on the local spin polarization. In Fig. 3 we have plotted the rotational dependence of the experimental NMR line shift $(H_{\rm eff} - H_0)$ and its extrapolation by a sinu-



FIG. 3: (color online) Left scale (black lines and symbols): rotation dependence of the NMR shift obtained from the highest frequency ⁶⁵Cu NMR line (open circles), the fit to these data (solid line) and an estimate of the orbital shift (dashed line). These results provide an estimate of the hyperfine field $H_{\rm spin}^{\parallel} \approx -2.0 \,\mathrm{T}$ and its anisotropy $H_{\rm spin}^{\parallel}/H_{\rm spin}^{\perp} \approx 10$, which corresponds to 10% spin polarization in a *d*-orbital. Right scale (color / grey-scale lines and symbols): ϑ_{EFG} deduced by fitting the spectra shown in Fig. 2 (solid squares and diamonds), following closely the predictions for the dimer Cu2 site (solid lines) and not the Cu1 site (short-dashed lines), as explained in the text. Experimental $\vartheta_{EFG}(\alpha)$ dependence (diamonds) is also used to define the upper horizontal scale.

soidal fit, together with an estimate of the orbital shift $(K_{\text{orb}}^{Z} \approx 1.3 \%, K_{\text{orb}}^{\perp} \approx 0.3 \%)$. We thus estimate the spininduced hyperfine field to be $H_{\text{spin}}^{\parallel} \approx -2.0 \text{ T}$, with the anisotropy $H_{\text{spin}}^{\parallel}/H_{\text{spin}}^{\perp} \approx 10$. This corresponds to about 10 % spin polarization of a typical $\text{Cu}^{2+} d_{\text{X}^2-\text{Y}^2}$ orbital. The error in these values is estimated to be $\approx 20 \%$, dominantly from the extrapolation of the angular dependence to $H_{\text{spin}}^{\parallel}$ [9]. In particular, in Fig. 3 we clearly see that the maximum of the experimental line shift is shifted by 18° from the expected $\vartheta_{\text{EFG}} = 90^{\circ}$ value. This means that the principal axes of the EFG and the hyperfine tensors are not really parallel, pointing to a departure from the simplified picture of pure $d_{X^2-Y^2}$ orbital. Indeed, the electronic density observed by x-ray diffraction suggest significant admixture of other orbitals [10].

In a true magnetization plateau the magnetization should not vary with the magnetic field. In order to test this most prominent feature of a plateau in Azurite, we performed very high field measurements of the copper NMR spectra, in the field range 17–28 T and at 1.4 K, for H_0 applied close to the *c*-axis. For technical convenience these spectra have been taken at constant NMR frequency by sweeping the field, and the obtained line positions are plotted in Fig. 4. The field dependence is found to be linear and the fits show that the NMR line shift is indeed to a high precision field independent. For

a field variation from 19 to 26 T (i.e. 37 %) the change in the measured $|H_{\rm spin}|$ is found to be (1 ± 1) %, where the precision is limited by our estimate of the orbital shift tensor. In contrast to the total magnetization measurements, the orbital effects have been separated out in our analysis proving that the local spin polarization of dimers is field independent. This information is important to distinguish and/or constrain the possible effects of Dzvaloshinski-Moria (DM) interaction terms, which might induce some weak field dependence of the spin polarizations. The DM interaction on the dimer bond has been invoked to explain strong anisotropy of the width of the plateau [1]. However, the presence of an inversion center at the center of the dimer precludes such a term, and only DM interaction on J_1 and J_3 exchange paths are possible. Whether or not they can explain the observed anisotropy has to be elucidated.

Despite considerable efforts, we could not observe the NMR signal from the monomer Cu1 spin. We have tried to find it in a broad frequency range covering the $H_{\rm spin}$ values up to -20 T, at very low temperature and in the middle of the plateau to minimize the spin fluctuations by maximizing the gap over T ratio. However, while these conditions ensure that the longitudinal (T_1) relaxation is certainly reduced, this is not necessarily the case for the transverse (T_2) relaxation [11]. We know that the spin polarization at the Cu1 site has to be $\approx 80\%$, to provide the correct total polarization of the 1/3 plateau. This is ≈ 8 times more than at the dimer sites, and we can roughly estimate that the corresponding T_2 ratio is



FIG. 4: Magnetic field dependence of the 63,65 Cu NMR line positions in Azurite at 1.4 K, for the field orientation close to the *c*-axis. The spectra were taken at constant frequency by sweeping the field, and results plotted in the reduced frequency scale to eliminate the dominant field dependence $\langle \gamma \rangle H_0$, with $\langle \gamma \rangle = ({}^{63}\gamma + {}^{65}\gamma)/2 = 11.687 \text{ MHz/T}$. Lines are linear fits to the observed field dependence, and symbols fits to this linear interpolation at 19, 22.5 and 26 T, which confirm that the spin polarization of the dimer Cu2 site is magnetic field independent. Color/symbol code is the same as in Fig. 2.

of the order of $8^2 = 64$. Knowing that the dimer T_2 values are in the $10\text{--}100\,\mu\text{s}$ range (depending on the orientation), this factor is enough to reduce the Cu1 T_2 below the experimental dead time for the observation of an NMR signal. This provides a reasonable explanation why Cu1 spin could not be observed, but also an important information about the system: the longitudinal spin fluctuations (effective in T_2 relaxation) are not gapped.

Finally, we remark that here we have only considered the standard on-site hyperfine coupling to the copper spin, and not the transferred hyperfine coupling which could in principle couple the observed nuclear spin at the dimer site to the neighboring (strongly polarized) monomer spin. This latter mechanism typically relies on almost negligible admixture ($\sim 1\%$) of the on-site s-wave orbital in the extended Wannier wave function belonging to the neighboring electronic spin. While the on-site spin polarization induced in this way is negligible ($\sim 1\%$), very high hyperfine coupling of an s-wave orbital ($\sim 200 \,\mathrm{T}/\mu_{\rm B}$) can in principle provide significant transferred hyperfine field. However, this field is positive and isotropic, in obvious contradiction to what is observed in Azurite. Negative and strongly anisotropic $H_{\rm spin}$ necessarily implies significant on-site spin polarization of the Cu orbital.

To provide a simple discussion for the observed polarization of dimer spins, we note that the approximate unit cell wave functions proposed for the 1/3 plateau of a diamond chain [3] can be generalized to represent an arbitrary mixture of the three single-spin-flip states, $\Psi(\mp,\beta) = \cos\beta \left[|\downarrow\uparrow\uparrow\rangle\mp|\uparrow\downarrow\uparrow\rangle \right]/\sqrt{2} - \sin\beta |\uparrow\uparrow\downarrow\rangle$, which by construction has correct total and local spin polarization of $\mu_{\rm B}$ and $\mu_{\rm B} \times (\sin^2\beta, \ \sin^2\beta, \ \cos^2\beta - \sin^2\beta)$, respectively. In particular, there is equal spin polarization on the two dimer spins. In this notation the two reference plateau states [3] are $00u = \Psi(-, 0)$ and *uud* $= \Psi(+, \arccos(1/\sqrt{3})),$ where the sign difference corresponds to the different symmetry. Note that the "-" signs in the definition of Ψ ensure dominant lowering of the energy by the exchange terms $S_i^{\pm} S_i^{\mp}$. Using $\Psi(\mp, \beta)$ as a trial function to minimize the energy by optimizing β , one can easily see that the pure singlet $00u \ (\beta = 0)$ state is obtained only for the symmetric diamond couplings $J_1 = J_3$, while deviation from this case necessarily leads to some admixture of the triplet, meaning some non-zero polarization of the dimer spins ($\beta \neq 0$). Observed polarization on the dimer site means that this admixture is significant, $\sin\beta \approx \sqrt{0.1} \approx 0.3$ [12], and thus should correspond to an important asymmetry of couplings. A correct estimate of the corresponding Jcoupling values should rely on numerical solutions of the spin Hamiltonian, relating the observed spin polarization to the corresponding constraint on the J couplings, say the J_3/J_1 vs. J_2/J_1 dependence [13]. Combined with other constrains, as the width of the plateau from the magnetization data [1] and the energy of excitations from the neutron data [6], this should clearly define the couplings, or indicate whether the diamond chain model is too simple to describe Azurite. Here we recall a possible influence of inter chain couplings. For example, for each dimer spin one of its NN oxygens is at the same time the apical oxygen (in the pyramid having a basis of 4 NN oxygens) of a dimer spin in the NN chain along a-axis.

In conclusion, by copper NMR in the 1/3 magnetization plateau of Azurite we have determined the local spin polarization of the dimer spins to be $\approx 0.1 \,\mu_{\rm B}$. This provides the first direct evidence for a "quantum" type of a 1/3 plateau having *no* classical analogue, which consists of dimers in a singlet state and fully polarized monomers. The deviation from ideal zero polarization of the dimer implies important asymmetry of the diamond couplings, $J_1 \neq J_3$, and provides a strong constraint for the determination of their values.

We acknowledge fruitful discussions with B. Grenier, A. Honecker, O. Janson, F. Mila, H. Ohta, T. Sakai and T. Tonegawa. Part of this work has been supported by the European Commission through the Transnational Access Specific Support Action (Contract No. RITA-CT-2003-505474) and the EuroMagNET network (Contract No. RII3-CT-2004-506239).

- * Present address: Jožef Stefan Institute, Jamova cesta 39, 1000 Ljubljana, Slovenia
- [†] Email address: mladen.horvatic@grenoble.cnrs.fr
- [1] H. Kikuchi et al., Phys. Rev. Lett. 94, 227201 (2005).
- [2] T. Ono *et al.*, Phys. Rev. B **67**, 104431 (2003) and Refs. therein.
- [3] K. Okamoto, T. Tonegawa and M. Kaburagi, J. Phys.: Condens. Matter 15 5979 (2003).
- [4] B. Gu and G. Su, Phys. Rev. B 75, 174437 (2007).
- [5] B. Gu and G. Su, Phys. Rev. Lett. 97, 089701 (2006);
 H. Kikuchi *et al.*, Phys. Rev. Lett. 97, 089702 (2006).
- [6] K. C. Rule et al., Phys. Rev. Lett. 100, 117202 (2008).
- [7] H.-J. Mikeska and C. Luckmann, Phys. Rev. B 77, 054405 (2008).
- [8] A. Abragam, The Principles of Nuclear Magnetism, (Clarendon Press, Oxford, 1961).
- [9] A minor approximation is to neglect the fact that the strong anisotropy and values of the A tensor make $\mathbf{H}_{\mathrm{eff}}$ and \mathbf{H}_0 somewhat non-parallel.
- [10] E. L. Belokoneva, Yu. K. Gubina, and J. B. Forsyth, Phys. Chem. Miner. 28, 498 (2001).
- [11] Well known example is the relaxation in the superconducting state of high T_c superconductors, for a review see C. Berthier *et al.*, J. Phys. I France **6**, 2205 (1996).
- [12] This corresponds closely to the state $\Psi(-, \arcsin(1/3))$ which was identified at $J_1 : J_2 : J_3 = 1 : 2 : -0.5$ and proposed to be the third type of plateau state [4]. However, while two other states are clearly distinguished by their different parity upon the exchange of two dimer spins, this third state rather appears as 00u state modified by some admixture of the triplet (on dimer spins) without changing its symmetry.
- [13] One compatible set of values is $J_1: J_2: J_3 = 1: 1.2: 0.4$, T. Tonegawa, private communication.