



**HAL**  
open science

## Cell wall component and mycotoxin moieties involved in binding of fumonisin B1 and B2 by lactic acid bacteria.

Vincent Niderkorn, Diego Morgavi, Bettina Aboab, Marielle Lemaire, Hamid Boudra

### ► To cite this version:

Vincent Niderkorn, Diego Morgavi, Bettina Aboab, Marielle Lemaire, Hamid Boudra. Cell wall component and mycotoxin moieties involved in binding of fumonisin B1 and B2 by lactic acid bacteria.. Journal of Applied Microbiology, 2009, 106, pp.977-985. 10.1111/j.1365-2672.2008.04065.x . hal-00360818v2

**HAL Id: hal-00360818**

**<https://hal.science/hal-00360818v2>**

Submitted on 9 Mar 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Cell wall component and mycotoxin moieties involved in the binding of fumonisin B<sub>1</sub> and B<sub>2</sub> by lactic acid bacteria

V. Niderkorn<sup>1,2</sup>, D.P. Morgavi<sup>1</sup>, B. Aboab<sup>3</sup>, M. Lemaire<sup>3</sup> and H. Boudra<sup>1</sup>

<sup>1</sup>INRA, UR1213 Herbivores, Saint-Genès Champanelle, France

<sup>2</sup>Lallemand S.A.S, 19, rue des briquetiers, Blagnac, France

<sup>3</sup>CNRS, UMR 6504, SEESIB, University Blaise Pascal, Aubiere, France

Correspondence to Hamid Boudra, INRA, UR1213 Herbivores, Site de Theix, F-63122 Saint-Genès Champanelle, France. E-mail: [hboudra@clermont.inra.fr](mailto:hboudra@clermont.inra.fr)

## KEYWORDS

binding ; bioavailability ; detoxification ; fumonisins ; lactic acid bacteria ; peptidoglycan ; probiotics

## ABSTRACT

**Aims:** The ability of lactic acid bacteria (LAB) to bind fumonisins B<sub>1</sub> and B<sub>2</sub> (FB<sub>1</sub>, FB<sub>2</sub>) in fermented foods and feeds and in the gastrointestinal tract could contribute to decrease their bioavailability and toxic effects on farm animals and humans. The aim of this work was to identify the bacterial cell wall component(s) and the functional group(s) of FB involved in the LAB–FB interaction.


**Methods and Results:** The effect of physicochemical, enzymatic and genetic treatments of bacteria and the removal/inactivation of the functional groups of FB on toxin binding were evaluated. Treatments affecting the bacterial wall polysaccharides, lipids and proteins increased binding, while those degrading peptidoglycan (PG) partially decreased it. In addition, purified PG from Gram-positive bacteria bound FB in a manner analogue to that of intact LAB. For FB, tricarballic acid (TCA) chains play a significant role in binding as hydrolysed FB had less affinity for LAB.

**Conclusions:** Peptidoglycan and TCA are important components of LAB and FB, respectively, involved in the binding interaction.

**Significance and Impact of the Study:** Lactic acid bacteria binding efficiency seems related to the peptide moiety structure of the PG. This information can be used to select probiotics with increased FB binding efficiency.

## Introduction

Fumonisin, a structurally related mycotoxin group produced by *Fusarium verticillioides* and *Fusarium proliferatum*, are common contaminants of corn and corn-based products worldwide (Shephard *et al.* 1996). There are several identified fumonisins, but fumonisin B<sub>1</sub> (FB<sub>1</sub>) and B<sub>2</sub> (FB<sub>2</sub>) are the most important and constitute up to 70% of the fumonisins found in naturally contaminated foods and feeds. FB<sub>1</sub> is the diester of propane-1,2,3-tricarboxylic acid (tricarballic acid, TCA) and 2-amino-12,16-dimethyl-3,5,10,14,15-pentahydroxyeicosane, in which the C<sub>14</sub> and C<sub>15</sub> hydroxyl groups are esterified with the terminal carboxyl group of TCA. FB<sub>2</sub> is the C<sub>10</sub> deoxy analogue of FB<sub>1</sub>, in which the corresponding stereogenic units on the icosane backbone have the same configurations (Fig. 1).



**Figure 1** Absolute configuration of fumonisin B<sub>1</sub> (FB<sub>1</sub>) and B<sub>2</sub> (FB<sub>2</sub>)

Fumonisin B<sub>1</sub> and B<sub>2</sub> are phytotoxic to corn (Lamprecht *et al.* 1994), cytotoxic to various mammalian cell lines (Abbas *et al.* 1993) and FB<sub>1</sub> is a carcinogen in rat liver and kidney (IARC 2002). The occurrence of these analogues in home-grown corn has been associated with an increased risk of esophageal cancer in humans (Shephard *et al.* 2000). FB<sub>1</sub> is considered possible carcinogens to human and classified as class 2B (IARC 2002). These mycotoxins are the causal agent of two well described diseases in domestic animals: equine leukoencephalomalacia (Riley *et al.* 1997) and porcine pulmonary edema syndrome (Harrison *et al.* 1990). In addition, they have also been associated with nephrotoxic, hepatotoxic and immunosuppressing effects in various animal species (Morgavi and Riley 2007). The mechanism of action appears to involve mainly disruption of sphingolipid biosynthesis by the inhibition of the enzyme sphingosine *N*-acetyltransferase (ceramide synthase) (reviewed by Voss *et al.* 2007). FB are more toxic than their hydrolysed or *N*-acetylated derivatives (Gelderblom *et al.* 1993). The free amino group appears to play a specific role in the biological activity of fumonisins.

Binding of FB by lactic acid bacteria (LAB) from fermented foods and feeds, and by LAB present in the gastrointestinal tract (GIT) could contribute to decrease the toxin bioavailability. This property could also decrease the exposure of intestinal mucosa to FB. Gut tissues exposed to FB have a diminished immune response and an altered barrier function against colonization by pathogenic *Escherichia coli* (Bouhet *et al.* 2004). Viable and nonviable LAB are able to bind FB in a pH, genus, bacterial density and analogue (FB<sub>2</sub> > FB<sub>1</sub>) dependent manner *in vitro* (Niderkorn *et al.* 2006). FB binding is rapid and particularly effective in acidic conditions, forming a stable complex in the range of pH present in the GIT. This activity is probably present in a variety of fermented foods and feeds (Mokoena *et al.* 2005; Niderkorn *et al.* 2007) and might also operate in the stomach. Binding of other major mycotoxins: aflatoxin B<sub>1</sub> (Haskard *et al.* 2000), zearalenone (El-Nezami *et al.* 2002a) and certain trichothecenes (El-Nezami *et al.* 2002b) by some probiotic LAB has also been shown *in vitro*. In the absence of a simple detoxification method for foods and feeds contaminated by FB, the use of selected strains of LAB appears as a promising approach to reduce their toxicological effects. However, an understanding of the binding mechanism is required to allow the optimization and safe dietary application of this technology. The aim of this work was to identify the component of the bacterial cell wall and the chemical structure of FB involved in the mechanism of binding.

## **Materials and methods**

### **Bacteria and bacteria-derived materials**

Strains *Lactobacillus paraplantarum* CNRZ 1885 (CNRS, FRE2326 Strasbourg, France) and *Streptococcus thermophilus* RAR1 (LAB collection of the Research Unit for Food Process Engineering and Microbiology, INRA, Thivernal-Grignon, France) were used in most experiments. *Streptococcus thermophilus* CNRZ 1066 and its non-capsular, non-exopolysaccharide (EPS) producing mutant *Strep. thermophilus* JIM 8752 (delta epsE) were obtained from the Microbial Genetics Unit, INRA, Jouy-en-Josas, France. *Lactococcus lactis* subsp. *cremoris* MG1363 and mutants, in which the synthesis of certain cell wall components and adhesion properties are affected, were from the LAB and Opportunistic Pathogens Laboratory, INRA, Jouy-en-Josas, France. Bacterial strains were grown at optimal temperature (30 or 37°C) in De Man, Rogosa, Sharpe broth for lactobacilli and M17 broth (Oxoid Ltd., Basingstoke, UK), supplemented with 0.5% of glucose for lactococci or 10% of lactose for streptococci. Commercial purified peptidoglycans (PG) from Gram-positive bacteria *Micrococcus luteus* and *Bacillus subtilis* were purchased from Sigma, Steinheim, Germany.

### **Determination of the bacterial cell wall component involved in binding**

To identify the binding site, bacteria were subjected to different physicochemical and enzymatic treatments. Bacteria (*Lact. paraplantarum* CNRZ 1885 and *Strep. thermophilus* RAR1) were prepared in advance and stored at -18°C until use. Optimization tests showed that freezing did not negatively affect the binding ability of these strains (shown in results). For experiments, bacteria were thawed at room temperature, washed twice with 0.01 mol l<sup>-1</sup> phosphate-buffered saline (PBS), pH 7.4 and treated by one of the following methods: water (25 or 100°C, 15 min), hydrochloric acid (1 mol l<sup>-1</sup> HCl, 100°C, 15 min), sodium dodecyl sulphate (SDS, 2% w/v, 100°C, 15 min) or trichloroacetic acid (10% w/v, 100°C, 15 min). After treatment, suspensions were centrifuged (3000 g, 10 min, 5°C). For enzymatic treatments, washed

bacteria were resuspended in 1 ml lysozyme (Sigma; 45 000 U ml<sup>-1</sup> in phosphate buffer, pH 6), mutanolysin (Sigma; 5000 U ml<sup>-1</sup> in phosphate buffer, pH 6), pronase E (Sigma; 1 mg ml<sup>-1</sup> in 0.01 mol l<sup>-1</sup> PBS, pH 7.4), lipase (Sigma; 1 mg ml<sup>-1</sup> in 0.01 mol l<sup>-1</sup> PBS, pH 7.4) or trypsin (Sigma; 1 mg ml<sup>-1</sup> in Tris-HCl buffer, pH 8, 10 mmol l<sup>-1</sup> CaCl<sub>2</sub>). Suspensions were incubated at 37°C for 2 h with shaking (240 rev min<sup>-1</sup>) and centrifuged (12 000 g, 10 min, 5°C). All bacterial pellets from both the physicochemical and enzymatic treatments were washed three times with 4 ml of PBS and used for the binding assay. Non-treated controls were added at each experimental run. All experiments were performed in triplicate.

### **Determination of the functional group of fumonisins involved in binding**

To identify which functional group of FB can interact with bacteria, different chemical reactions were applied at different sites of FB derivatives. FB<sub>1</sub> and FB<sub>2</sub>, purchased from Sigma and Promec (Tygerberg, South Africa), respectively, were dissolved in an exact volume of acetonitrile–water in a 1 : 1 (v/v) ratio to achieve the desired concentration of stock solutions. Hydrolysed FB<sub>1</sub> (HFB<sub>1</sub>) and FB<sub>2</sub> (HFB<sub>2</sub>) were obtained according to Pagliuca *et al.* (2005). Total hydrolysis of pure FB<sub>1</sub> and FB<sub>2</sub> was checked by HPLC. The chromatograms showed absence of FB peak and presence of a single peak with retention times corresponding to the expected HFB product (Pagliuca *et al.* 2005). An optimized procedure was also used to determine the effect of the amine group in binding. Free amine of both FB was hidden by reaction with *ortho*-phthalaldehyde (OPA). This option was chosen because the fumonisins of the group A in which the free amine is naturally absent are not commercially available.

### ***In vitro* binding assay**

Treated and non-treated bacteria (10<sup>9</sup> or 10<sup>10</sup> CFU ml<sup>-1</sup> for certain experiments, see footnotes of tables) were tested as previously described (Niderkorn *et al.* 2007). Briefly, bacterial material was suspended in 1 ml of corn infusion adjusted to pH 4 with lactic acid and containing FB<sub>1</sub> and FB<sub>2</sub> (5 µg ml<sup>-1</sup> each) or their derivative compounds. The corn infusion was prepared by steeping dry whole-plant corn in water and filtering as described by Niderkorn *et al.* (2007). For each experiment, positive controls containing no bacterial material and a negative control containing no toxin were included. Assays and controls were incubated at 25°C for 1 h and centrifuged (3000 g, 10 min, 5°C). Supernatants and bacterial pellets were analysed for FB by reversed-phase HPLC to determine free and bound fractions respectively. Because of the instability of the FB-OPA derivative (Williams *et al.* 2004), assays with the free amine hidden were performed following an exact timing: At *t* = 0, a pure FB solution (800 µg ml<sup>-1</sup>) and reagent with (or without) OPA were mixed (1 : 1 v/v). At *t* = 2 min, 50 µl of this mixture was mixed to 950 µl of acidified corn infusion containing bacteria (10<sup>10</sup> CFU ml<sup>-1</sup>), then incubated for 9.25 min at 25°C. At *t* = 12 min, tubes were centrifuged (4500 g, 3 min, 4°C). At *t* = 20 min, supernatants containing free FB were derivatized with OPA. All samples were injected at *t* = 22 min. In these conditions, preliminary assays have shown that the complex FB-OPA remains sufficiently stable to carry out measurements. For this experiment, pellets were not analysed.

### **Fumonisin analysis**

Supernatants from all samples and pellets extracts were fourfold diluted in acetonitrile–water (1 : 1 v/v), then 40 µl were added to 60 µl 0.1 mol l<sup>-1</sup> borate buffer at pH 10 and 100 µl of OPA reagent were added. The preparation was mixed and allowed to react for 2 min before injection of 20 µl into the HPLC system. For FB extraction, 1 ml acetonitrile–water (1 : 1 v/v) was added to the bacterial pellets and this mixture was vigorously vortexed, placed in an ultrasonic bath for 6 min, then centrifuged (4500 g, 3 min, 5°C). Analysis of FB and their hydrolysed derivatives were done at room temperature by HPLC, using fluorimetric detection. The HPLC system consisted of a GOLD 126 solvent module (Beckman Coulter, Fullerton, CA, USA), an automatic sampler (Spectra-Physics, San Jose, CA, USA) equipped with a 100-µl loop and a fluorescence detector FL3000 (Spectra-System, San Jose, CA, USA). Separation of FB<sub>1</sub>, FB<sub>2</sub>, HFB<sub>1</sub> and HFB<sub>2</sub> was performed on a C<sub>18</sub> reversed-phase column (Prontosil, 150 × 4.6 mm, 3 µm, Bishoff Chromatography) with a gradient elution using acetonitrile (A) and water–methanol (1 : 1 v/v) acidified at pH 3.35 with pure acetic acid (B). The gradient was started at 10% of solvent A, which increased to 60% in 6 min, then maintained at 60% for 7 min, before returned to the initial condition in 1 min. The flow rate was 1 ml min<sup>-1</sup> and detection was set at 336 nm excitation and 440 nm emission.

The retention times of FB<sub>1</sub>, FB<sub>2</sub>, HFB<sub>1</sub> and HFB<sub>2</sub> were 9.9, 12.2, 10.2, 13.4 min respectively. The percentage of free (or bound) mycotoxin was calculated as 100× [Peak area of mycotoxin in the supernatant (or pellet extract)/Peak area of mycotoxin in the positive control].

## Statistical analysis

Data was subjected to the analysis of variance (ANOVA). A significant difference between means of controls and assays ( $P < 0.05$ ) was determined by Dunnett's test using the STATISTICAL ANALYSIS SYSTEM (SAS) software package, ver. 8 (SAS Institute Inc., Cary, NC, USA).

## Results

### Bacterial cell wall components affecting binding

None of the physicochemical treatments applied to bacteria decreased binding of FB<sub>1</sub> or FB<sub>2</sub>. On the contrary, freezing/thawing and thermal treatments of bacteria increased the bound fractions of FB<sub>1</sub> and FB<sub>2</sub> in both tested strains ( $P < 0.05$ ) (Table 1). Among the chemical treatments, trichloroacetic acid caused a large increase in bound FB proportion ( $P < 0.05$ ). HCl also produced the same effect although it was only significant on *Streptococcus* cells ( $P < 0.05$ ). For the enzymatic treatments, lysozyme and mutanolysin were the only treatments which caused a partial, but significant decrease of this activity ( $P < 0.05$ ) (Table 1). In contrast, lipase, trypsin and pronase E, an unspecific protease from *Streptomyces griseus*, had no effect on binding ( $P > 0.05$ ) (data not shown).

**Table 1** Effect of freezing, chemical and enzymatic treatments of bacteria on binding of fumonisin B<sub>1</sub> and B<sub>2</sub> by *Streptococcus thermophilus* RAR1 and *Lactobacillus paraplantarum* CNRZ 1885\*

Treatment	FB <sub>1</sub>						FB <sub>2</sub>					
	<i>Strep. thermophilus</i> RAR1			<i>Lact. paraplantarum</i> CNRZ 1885			<i>Strep. thermophilus</i> RAR1			<i>Lact. paraplantarum</i> CNRZ 1885		
	Free (%)‡	Bound (%)§	Recov. (%)	Free (%)‡	Bound (%)§	Recov. (%)	Free (%)‡	Bound (%)§	Recov. (%)	Free (%)‡	Bound (%)§	Recov. (%)
Freezing												
Fresh bacteria (control)	95 ± 3	3 ± 0	98	95 ± 5	2 ± 0	97	64 ± 6	41 ± 3	105	71 ± 0	27 ± 2	98
Thawed bacteria	93 ± 2	14 ± 1 ↑	107	93 ± 1	8 ± 0 ↑	101	41 ± 2 ↑	51 ± 1 ↑	92	60 ± 4 ↑	42 ± 0 ↑	102
Physicochemical treatments†												
Water, room temperature, 15 min (control)	78 ± 5	9 ± 0	87	79 ± 2	9 ± 0	88	30 ± 4	45 ± 1	75	36 ± 5	45 ± 1	81
HCl, 1 mol l <sup>-1</sup> , 100°C, 15 min	67 ± 1	24 ± 1 ↑	91	87 ± 3	7 ± 2	94	19 ± 2	65 ± 2 ↑	84	35 ± 7	51 ± 7	86
Trichloroacetic acid, 10% (w/v), 100°C, 15 min	54 ± 1 ↑	37 ± 1 ↑	91	69 ± 3 ↑	19 ± 4	88	9 ± 2 ↑	76 ± 2 ↑	85	20 ± 3	65 ± 8 ↑	85
Enzymatic treatments†												
Without treatment (control)	91 ± 4	14 ± 1	105	93 ± 3	7 ± 1	100	38 ± 1	53 ± 2	91	65 ± 5	43 ± 2	108
Lysozyme, 45 000 U ml <sup>-1</sup>	89 ± 3	9 ± 0 ↓	98	93 ± 4	7 ± 0	100	42 ± 11	41 ± 1 ↓	83	75 ± 3 ↓	34 ± 1 ↓	109
Mutanolysin, 5000 U ml <sup>-1</sup>	92 ± 4	10 ± 1 ↓	102	99 ± 5	5 ± 1 ↓	104	51 ± 7 ↓	42 ± 2 ↓	93	77 ± 7 ↓	27 ± 6 ↓	104

Data shown are means ± SD of triplicates.

↓ and ↑ within the same column, indicate that treatment decreased or increased binding ( $P < 0.05$ ) compared with corresponding control.

\*For all experiments, treated or not treated bacteria (10<sup>9</sup> CFU ml<sup>-1</sup>) were incubated in acidified corn infusion containing FB<sub>1</sub> and FB<sub>2</sub> (5 µg ml<sup>-1</sup> each) for 1 h at 25°C.

†Experiments were performed with thawed bacteria. Enzymatic treatments were done at 37°C for 2 h.

‡Free fraction of fumonisin remaining in supernatant (vs control without bacteria).

§Bound fraction of fumonisin remaining in bacterial pellet (vs bacterial pellet spiked with FB<sub>1</sub> and FB<sub>2</sub>, 5 µg each).


**Table 2** Effect of mutations affecting lipoteichoic acids and peptidoglycan biosynthesis in *Lactococcus lactis* subsp. *cremoris* on binding of fumonisin B<sub>1</sub> and B<sub>2</sub>\*

Genotype	Protein, function affected or phenotype	FB <sub>1</sub>			FB <sub>2</sub>		
		Free (%)	Bound (%)	Recov. (%)	Free (%)	Bound (%)	Recov. (%)
Derivatives from <i>L. lactis</i> subsp. <i>cremoris</i> MG1363							
MG1363	Wild type	92 ± 6	4 ± 1	96	25 ± 2	65 ± 3	90
<i>dltD</i>	LTA synthesis	88 ± 1 ↑	11 ± 1 ↑	99	30 ± 0	69 ± 4	99
<i>pbp 2A</i> <sup>-</sup>	PBP 2A (PG transpeptidase)	94 ± 1	4 ± 1	98	48 ± 8 ↓	47 ± 1 ↓	95
<i>pbp 2B</i> <sup>-</sup>	PBP 2B (PG transpeptidase)	95 ± 5	5 ± 0	100	52 ± 6 ↓	45 ± 3 ↓	97

Data shown are means ± SD of triplicates.  
 ↓ and ↑ within the same column, indicate that treatment decreased or increased binding ( $P < 0.05$ ) compared with the corresponding control.  
 PG, peptidoglycan; LTA, lipoteichoic acids; PBP, Penicillin binding protein.  
 \*Bacteria ( $10^{10}$  CFU ml<sup>-1</sup>) were incubated in corn infusion containing FB<sub>1</sub> and FB<sub>2</sub> (5 µg ml<sup>-1</sup> each) for 1 h at 25°C.

### Role of peptidoglycan

We observed decrease of FB<sub>2</sub> binding with mutants of *L. lactis* that had an altered PG structure because of perturbed transpeptidase functions (*pbp 2A*<sup>-</sup> and *2B*<sup>-</sup>) (Shohayeb and Chopra 1987) ( $P < 0.05$ ) (Table 2). However, mutants *acmA* and *ponA* from *L. lactis*, in which the immobilization property (phenomenon of adhesion, chain and biofilm formation) was modified (Mercier *et al.* 2002), had no effect on binding as compared with wild type *L. lactis* ( $P > 0.05$ ) (data not shown). To confirm the role of PG component, purified PG from Gram-positive bacteria *M. luteus* and *B. subtilis* at different concentrations (0, 0.1, 0.5, 1 and 1.5 mg ml<sup>-1</sup>) were tested in a similar way. Results showed that these polymers can bind FB in an analogue dependent manner (FB<sub>2</sub> > FB<sub>1</sub>) (Fig. 2). Significant bound fractions were observed even with the lowest concentration tested (0.1 mg PG ml<sup>-1</sup>). However, the binding efficiency varied between the two purified PG tested (*B. subtilis* > *M. luteus*).



**Figure 2** Fractions of fumonisin B<sub>1</sub> (□, ■) and B<sub>2</sub> (○, ●) bound to purified peptidoglycans from *Bacillus subtilis* (open symbols) and *Micrococcus luteus* (closed symbols). Data shown are the mean and standard deviations (error bars) of triplicates.

### Fumonisin structural component affecting binding

To identify the role of the main functional groups of FB in the formation of the mycotoxin–cell wall complex, the free amine and TCA arms were alternatively hidden or removed. When the free-amine group was hidden by derivatization with OPA, the proportions of FB<sub>1</sub> and FB<sub>2</sub> bound by *Lact. paraplantarum* CNRZ 1885 and *Strep. thermophilus* RAR1 were higher than those observed with unmodified toxins (Table 3). This effect was more pronounced for FB<sub>1</sub> ( $P < 0.05$ ) than FB<sub>2</sub> ( $P > 0.05$ ). Inversely, the binding rates of HFB<sub>1</sub> and HFB<sub>2</sub> for both strains was lower than those of FB<sub>1</sub> and FB<sub>2</sub> respectively ( $P < 0.05$ ). Whatever the applied treatment, binding appears to be greater for FB<sub>2</sub> comparing

with FB<sub>1</sub>. To explain the different binding behaviour between FB<sub>1</sub> and FB<sub>2</sub>, we investigated their three-dimensional structure by molecular modelling in conditions simulating those of the binding tests. For that, conformations were carried out in aqueous conditions applying ionized states of carboxyl and amine groups of FB in acidic conditions to generate the most stable conformer using MACROMODEL 8.0 (Shroedinger Inc, Portland, OR, USA). Results showed that a hydrogen bond in FB<sub>1</sub> structure is formed between the hydrogen of the hydroxyl group at C<sub>10</sub> and the oxygen of the carbonyl group of the TCA at C<sub>15</sub> (Fig. 3).

**Table 3** Effect of hydrolysis of fumonisin's tricarballylic acid chains and free amine group inactivation on binding of fumonisin B<sub>1</sub> and B<sub>2</sub> by *Lactobacillus paraplantarum* and *Streptococcus thermophilus*\*

Toxin	<i>Lact. paraplantarum</i> CNRZ 1885			<i>Strep. thermophilus</i> RAR1		
	Free (%)	Bound (%)	Recov. (%)	Free (%)	Bound (%)	Recov. (%)
FB <sub>1</sub>	73 ± 7	27 ± 2	100	59 ± 1	40 ± 1	99
HFB <sub>1</sub>	80 ± 6	14 ± 0 ↓	94	85 ± 10 ↓	22 ± 1 ↓	107
FB <sub>2</sub>	19 ± 1	74 ± 1	93	27 ± 1	71 ± 3	98
HFB <sub>2</sub>	43 ± 4 ↓	46 ± 11 ↓	89	48 ± 4 ↓	45 ± 3 ↓	93
FB <sub>1</sub> †	89 ± 1	ND‡		81 ± 1	ND	
FB <sub>1</sub> -OPA	68 ± 2 ↑	ND		66 ± 8 ↑	ND	
FB <sub>2</sub> †	56 ± 4	ND		31 ± 4	ND	
FB <sub>2</sub> -OPA	48 ± 0	ND		26 ± 0	ND	


Data shown are means ± SD of triplicates.

↓ and ↑ within the same column, indicate that treatment decreased or increased binding ( $P < 0.05$ ) compared with the corresponding control.

\*Bacterial density =  $10^{10}$  CFU ml<sup>-1</sup>.

†Treated in the same way as FB-OPA derivatives (see M&M for details).

‡Not determined.



**Figure 3** Molecular conformations of FB<sub>1</sub> and FB<sub>2</sub> in aqueous solution. Conformational analysis of molecules in water solution was performed using Monte-Carlo Multiple Method (Chang *et al.* 1989) with AMBER force field (Weiner *et al.* 1984; Cornell *et al.* 1995) and GB/SA solvation model (Still *et al.* 1990) of MACROMODEL 8.0 programme (Shroedinger Inc, Portland, OR, USA). To take account of pKas of fumonisins, conformations were carried out applying ionized states

of carboxyl and amine groups. In this case, four Na<sup>+</sup> and one Cl<sup>-</sup> were added in the solution to maintain neutral charge of the molecular system.

## Discussion

The ability of LAB to bind fumonisins might contribute to decrease the bioavailability and toxic effects of FB<sub>1</sub> and FB<sub>2</sub> in human and farm animals. The binding activity of LAB could be integrated in the criteria of selection of probiotics and starters used for the acidification of fermented corn meals and corn silage. However, the mechanism of binding is unknown and the LAB–FB interaction needs to be better understood to optimize the selection of strains. In this paper, we provided some insight into the interaction between FB and LAB that can explain different binding behaviour of FB<sub>1</sub> and FB<sub>2</sub>.

### Determination of the bacterial cell wall binding site

The cell wall of LAB has the typical Gram-positive structure made of a thick, multilayered PG sacculus in which proteins, teichoic acid (TA) and LTA and polysaccharides are associated (Delcour *et al.* 1999). We previously reported that all genera of LAB are capable to bind FB<sub>1</sub> and FB<sub>2</sub> (Niderkorn *et al.* 2007) suggesting that the binding site is a component largely conserved in the cell wall of these bacteria. This component is synthesized early in the bacterial growth cycle since binding was observed in the latency phase. Binding was observed throughout the growth cycle with a maximum at the end of the exponential phase (data not shown).

The increase in binding observed with heat- and acid-treated bacteria (Table 1) was also reported for other mycotoxins such as aflatoxin B<sub>1</sub> (El-Nezami *et al.* 1998) and zearalenone (El-Nezami *et al.* 2002a). It is known that these treatments degrade the surface of the cell wall. The trichloroacetic acid treatment to extract PG-associated cell wall polymers from Gram-positive bacteria is well established (Heckels and Virji 1988). Polysaccharides and TA are known targets of HCl and trichloroacetic acid treatments (Quiberoni *et al.* 2000). Our results suggest that binding takes place in the subsurface of the cell wall in sites exposed by the heat or acid treatments.

The results obtained with mutant strains are in agreement with the physicochemical and enzymatic treatments of bacteria. Taken together, results indicate that the binding site of FB are not surface polysaccharides, lipids or proteins, but may be rather the PG or compounds tightly associated to it, as it was suggested for the binding of aflatoxin B<sub>1</sub> (Lahtinen *et al.* 2004). It is worth mentioning that LTA are the main components responsible for the hydrophobicity of the cell wall and thus, for the adhesion properties of bacteria (Dahlback *et al.* 1981). However, these adhesion properties appear not to have any function in FB binding. Similarly, the mechanism of immobilization of bacteria, characterized by natural PG modifications consisting of little breaks in the PG structure (Ibrahim *et al.* 2004), seems not to be associated with FB binding.

Binding by commercially available purified PG from two strains of Gram-positive bacteria (Fig. 2) was consistent with our results obtained with intact LAB, thus, supporting the hypothesis that PG is likely the binding site of FB. The PG backbone is a conserved structure composed of linear glycan chains alternating *N*-acetyl glucosamine (GlcNAc) and *N*-acetyl muramic acid (MurNAc) in a β (1→4) linkage. These chains are crosslinked by means of short peptides. The specific amino acid sequence of peptide bridges and consequently, the molecular structure of PG vary with the bacterial species (Schleifer and Kandler 1972). As the bound fraction of FB varied between the PG tested, but also among genera of LAB (Niderkorn *et al.* 2007), it seems that the amino acid sequence play an important role in the efficiency of the mechanism. The PG structure vary mainly in the amino acid in position 3 (AA<sub>3</sub>) of the peptide bridge and in the cross-linking amino acids. *B. subtilis* and *M. luteus* differ in both the AA<sub>3</sub> and cross-linking amino acids (Schleifer and Kandler 1972). This difference could explain their dissimilar efficiency in binding FB. The higher binding efficiency of the *Streptococcus* genus compared with the *Lactobacillus* genus could be because of the amino acid sequence of the cross-bridge that is two to three molecules of L-Ala in the former and D-Asp in the latter (Schleifer and Kandler 1972; Bouhss *et al.* 2002).

### Relationship between fumonisins structure and binding

The higher binding rate of the FB-OPA derivative compared with unmodified FB suggest that the free-amine group possessing nucleophilic properties is not involved in FB<sub>1</sub> and FB<sub>2</sub> interaction with bacteria. In addition, in acidic conditions, the ionized state of this function could even decrease binding, in particular for FB<sub>1</sub>. Controls containing reagents other than OPA, *e.g.* mercaptoethanol, used in the



derivatization reaction were done to exclude possible interferences on binding. However, the exact function that OPA may have on the FB derivative and/or on the bacterial cell wall that could modify the interaction was unknown. The use of natural fumonisin derivatives such as *N*-acetylated FB<sub>1</sub> might be a better alternative to test the exact role of the free-amino group. Notwithstanding the reservations associated to the derivatization methodology, masking the free-amino group increased rather than decreased binding suggesting that this chemical function has not a positive effect on FB-bacteria interaction. Inversely, the lower binding rate of hydrolysed FB compared with the intact FB indicates that one or both TCA arms play a positive role in the mechanism.

In spite of their similar structure, FB<sub>2</sub> was in all experiments more bound than FB<sub>1</sub> (Tables 1 and 2 and Fig. 2). The same tendency between FB was recorded for OPA-treated or hydrolysed derivatives (FB<sub>2</sub>-OPA > FB<sub>1</sub>-OPA, HFB<sub>2</sub> > HFB<sub>1</sub>) (Table 3). These results are in agreement with those reported for other bacteria and other experimental conditions (Niderkorn *et al.* 2006, 2007). The only structural variation between FB<sub>1</sub> and FB<sub>2</sub> consists in an additional hydroxyl group in C<sub>10</sub> for FB<sub>1</sub> (Fig. 1). Thus, it is reasonable to postulate that this hydroxyl group plays directly or indirectly a negative role in binding. The spatial conformation induced by the hydrogen bond of FB<sub>1</sub> makes the molecule more coiled and apparently less favourable to binding by the bacterial cell wall. This conformation could disturb the interactions with the PG. These molecular conformations were conserved through pH variation as the addition of charges on functional groups of FB<sub>1</sub> and FB<sub>2</sub> did not affect the results of modelling. However, as HFB<sub>2</sub> was more bound than HFB<sub>1</sub>, it seems that the hydroxyl group in C<sub>10</sub> continues to be unfavourable to binding after TCA removal.

The objective of this work was to improve our understanding of the LAB–FB binding interaction. However, further work is needed before this methodology can be used to treat contaminated feeds. It is important to note that, differently from normal probiotic strains, LAB used to bind FB should have a low capacity to adhere to intestinal mucus and enterocytes to reduce the risk of toxin release in the GIT. The efficiency of bacterial strains to modulate intestinal toxin absorption and toxicity was already demonstrated *in vivo* for aflatoxin B1 in both rats (Gratz *et al.* 2006) and humans (El-Nezami *et al.* 2006).

## Conclusions

In this work, we demonstrated that PG of LAB and more generally PG of Gram-positive bacteria, are the most likely site of FB binding. This result helps to explain the widespread binding of fumonisins by LAB. Existing differences in binding capacity of different bacterial species can be rationally explained by the variation in PG structure. This observation should allow to select efficient strains in terms of FB binding, as fermentation starters and/or probiotic mixtures on the base of their PG-type. We also showed that at least one TCA arm of FB play an important role in their binding to bacteria. As it was reported that TCA arms also play a favourable role in the intestinal absorption of FB<sub>1</sub> (Dantzer *et al.* 1999; De Angelis *et al.* 2005), binding of FB<sub>1</sub> and FB<sub>2</sub> could decrease even more their absorption and their toxic effects on the intestinal mucosal cells. However, further quantitative *in vitro* and *in vivo* studies are needed to evaluate the real impact of LAB binding activity on the bioavailability of FB in higher organisms.

## References

- Abbas, H.K., Gelderblom, W.C., Cawood, M.E. and Shier, W.T. (1993) Biological activities of fumonisins, mycotoxins from *Fusarium moniliforme*, in jimsonweed (*Datura stramonium* L.) and mammalian cell cultures. *Toxicon* **31**, 345–353.
- Bouhet, S., Hourcade, E., Loiseau, N., Fikry, A., Martinez, S., Roselli, M., Galtier, P., Mengheri, E. *et al.* (2004) The mycotoxin fumonisin B1 alters the proliferation and the barrier function of porcine intestinal epithelial cells. *Toxicol Sci* **77**, 165–171.
- Bouhss, A., Josseume, N., Severin, A., Tabei, K., Hugonnet, J.E., Shlaes, D., Mengin-Lecreulx, D., van Heijenoort, J. *et al.* (2002) Synthesis of the L-alanyl-L-alanine cross-bridge of *Enterococcus faecalis* peptidoglycan. *J Biol Chem* **277**, 45935–45941.
- Chang, G., Guida, W.C. and Still, W.C. (1989) An internal-coordinate Monte Carlo method for searching conformational space. *J Am Chem Soc* **111**, 4379–4386.
- Cornell, W.D., Cieplak, P., Bayly, C.I., Gould, I.R., Merz, K.M., Ferguson, D.M., Spellmeyer, D.C., Fox, T. *et al.* (1995) A second generation force field for the simulation of proteins, nucleic acids, and organic molecules. *J Am Chem Soc* **117**, 5179–5197.
- Dahlback, B., Hermansson, M., Kjelleberg, S. and Norkrans, B. (1981) The hydrophobicity of bacteria – an important factor in their initial adhesion at the air–water interface. *Arch Microbiol* **128**, 267–270.

- Dantzer, W.R., Hopper, J., Mullin, K., Hendrich, S. and Murphy, P.A. (1999) Excretion of 14 C-fumonisin B, 14 C-hydrolyzed fumonisin B, and 14 C-fumonisin B-fructose in rats. *J Agric Food Chem* **47**, 4291–4296.
- De Angelis, I., Friggè, G., Raimondi, F., Stamatii, A., Zucco, F. and Caloni, F. (2005) Absorption of fumonisin B1 and aminopentol on an *in vitro* model of intestinal epithelium; the role of P-glycoprotein. *Toxicol* **45**, 285–291.
- Delcour, J., Ferain, T., Deghorain, M., Palumbo, E. and Hots, P. (1999) The biosynthesis and functionality of the cell-wall of lactic acid bacteria. *Antonie Van Leeuwenhoek* **76**, 159–184.
- El-Nezami, H., Kankaanpää, P., Salminen, S. and Ahokas, J. (1998) Physicochemical alterations enhance the ability of dairy strains of lactic acid bacteria to remove aflatoxin from contaminated media. *J Food Prot* **61**, 466–468.
- El-Nezami, H., Polychronaki, N., Salminen, S. and Mykkanen, H. (2002a) Binding rather than metabolism may explain the interaction of two food-grade *Lactobacillus* strains with zearalenone and its derivative  $\alpha$ -zearalenol. *Appl Environ Microbiol* **68**, 3545–3549.
- El-Nezami, H.S., Chrevatidis, A., Auriola, S., Salminen, S. and Mykkanen, H. (2002b) Removal of common *Fusarium* toxins *in vitro* by strains of *Lactobacillus* and *Propionibacterium*. *Food Addit Contam* **19**, 680–686.
- El-Nezami, H.S., Polychronaki, N.N., Ma, J., Zhu, H., Ling, W., Salminen, E.K., Juvonen, R.O., Salminen, S.J. *et al.* (2006) Probiotic supplementation reduces a biomarker for increased risk of liver cancer in young men from Southern China. *Am J Clin Nutr* **83**, 1199–1203.
- Gelderblom, W.C., Cawood, M.E., Snyman, S.D., Vlegaar, R. and Marasas, W.F. (1993) Structure-activity relationships of fumonisins in short-term carcinogenesis and cytotoxicity assays. *Food Chem Toxicol* **31**, 407–414.
- Gratz, S., Taubel, M., Juvonen, R.O., Viluksela, M., Turner, P.C., Mykkanen, H. and El-Nezami, H. (2006) *Lactobacillus rhamnosus* strain GG modulates intestinal absorption of aflatoxin B1 and its fecal excretion and toxicity in rats. *Appl Environ Microbiol* **72**, 7398–7400.
- Harrison, L.R., Colvin, B.M., Greene, J.T., Newman, L.E. and Cole, J.R. Jr (1990) Pulmonary edema and hydrothorax in swine produced by fumonisin B1, a toxic metabolite of *Fusarium moniliforme*. *J Vet Diagn Invest* **2**, 217–221.
- Haskard, C., Binnion, C. and Ahokas, J. (2000) Factors affecting the sequestration of aflatoxin by *Lactobacillus rhamnosus* strain GG. *Chem Biol Interact* **128**, 39–49.
- Heckels, J.E. and Virji, M. (1988) Separation and purification of surface components. In *Bacterial Cell Surface Techniques* ed. Hancock, I.C. and Poxton, I.R. pp. 67–135. New York: John Wiley and Sons.
- Ibrahim, M., Briandet, R., Mistou, M.Y., Chretien, A., Tremblay, J. and Kulakauskas, S. (2004) Immobilization of lactococci. *Lait* **84**, 103–114.
- International Agency for Research on Cancer. (2002) *IARC Monographs on the Evaluation of the Carcinogenic Risks to Humans: Some Traditional Herbal Medicines, some Mycotoxins, Naphthalene and Styrene*. Lyon, France: International Agency for Research on Cancer **82**, 301–366.
- Lahtinen, S.J., Haskard, C.A., Ouwehand, A.C., Salminen, S.J. and Ahokas, J.T. (2004) Binding of aflatoxin B1 to cell wall components of *Lactobacillus rhamnosus* strain GG. *Food Addit Contam* **21**, 158–164.
- Lamprecht, S.C., Marasas, W.F.O., Alberts, J.F., Cawood, M.E., Gelderblom, W.C.A., Shephard, G.S., Thiel, P.G. and Calitz, F.J. (1994) Phytotoxicity of fumonisins and TA-toxin to corn and tomato. *Phytopathology* **84**, 393–391.
- Mercier, C., Durrieu, C., Briandet, R., Domakova, E., Tremblay, J., Buist, G. and Kulakauskas, S. (2002) Positive role of peptidoglycan breaks in lactococcal biofilm formation. *Mol Microbiol* **46**, 235–243.
- Mokoena, M.P., Chelule, P.K. and Gqaleni, N. (2005) Reduction of fumonisin B1 and zearalenone by lactic acid bacteria in fermented maize meal. *J Food Prot* **68**, 2095–2099.
- Morgavi, D.P. and Riley, R.T. (2007) An historical overview of field disease outbreaks known or suspected to be caused by consumption of feeds contaminated with *Fusarium* toxins. *Anim Feed Sci Technol* **137**, 201–212.
- Niderkorn, V., Boudra, H. and Morgavi, D.P. (2006) Binding of *Fusarium* mycotoxins by fermentative bacteria *in vitro*. *J Appl Microbiol* **101**, 849–856.
- Niderkorn, V., Morgavi, D.P., Pujos, E., Tissandier, A. and Boudra, H. (2007) Screening of fermentative bacteria for their ability to bind and biotransform deoxynivalenol, zearalenone and fumonisins in an *in vitro* simulated corn silage model. *Food Addit Contam* **24**, 406–415.
- Pagliuca, G., Zironi, E., Ceccolini, A., Matera, R., Serrazanetti, G.P. and Piva, A. (2005) Simple method for the simultaneous isolation and determination of fumonisin B1 and its metabolite aminopentol-1 in swine liver by liquid chromatography-fluorescence detection. *J Chromatogr B* **819**, 97–103.
- Quiberoni, A., Stiefel, J.I. and Reinheimer, J.A. (2000) Characterization of phage receptors in *Streptococcus thermophilus* using purified cell walls obtained by a simple protocol. *J Appl Microbiol* **89**, 1059–1065.
- Riley, R.T., Showker, J.L., Owens, D.L. and Ross, P.F. (1997) Disruption of sphingolipid metabolism and induction of equine leukoencephalomalacia by *Fusarium proliferatum* culture material containing fumonisin B2 or B3. *Environ Toxicol Pharmacol* **3**, 221–228.
- Schleifer, K.H. and Kandler, O. (1972) Peptidoglycan types of bacterial cell walls and their taxonomic implications. *Bacteriol Rev* **36**, 407–477.
- Shephard, G.S., Thiel, P.G., Stockenstrom, S. and Sydenham, E.W. (1996) Worldwide survey of fumonisin contamination of corn and corn-based products. *J AOAC Int* **79**, 671–687.
- Shephard, G.S., Marasas, W.F.O., Leggott, N.L., Yazdanpanah, H., Rahimian, H. and Safavi, N. (2000) Natural occurrence of fumonisins in corn from Iran. *J Agric Food Chem* **48**, 1860–1864.
- Shohayeb, M. and Chopra, I. (1987) Mutations affecting penicillin-binding proteins 2 a, 2 b and 3 in *Bacillus subtilis* alter cell shape and peptidoglycan metabolism. *Microbiology* **133**, 1733–1742.
- Still, W.C., Tempczyk, A., Hawley, R.C. and Hendrickson, T. (1990) Semianalytical treatment of solvation for molecular mechanics and dynamics. *J Am Chem Soc* **112**, 6127–6129.

- Voss, K.A., Smith, G.W. and Haschek, W.M. (2007) Fumonisin: toxicokinetics, mechanism of action and toxicity. *Anim Feed Sci Technol* **137**, 299–325.
- Weiner, S.J., Kollman, P.A., Case, D.A., Singh, U.C., Ghio, C., Alagona, G., Profeta, S. and Weiner, P. (1984) A new force field for molecular mechanical simulation of nucleic acids and proteins. *J Am Chem Soc* **106**, 765–784.
- Williams, L.D., Meredith, F.I. and Riley, R.T. (2004) Fumonisin-ortho-phthalaldehyde derivative is stabilized at low temperature. *J Chromatogr B* **806**, 311–314.