

Limit distribution theory for maximum likelihood estimation of a log-concave density

Fadoua Balabdaoui, Kaspar Rufibach, Jon A. Wellner

▶ To cite this version:

Fadoua Balabdaoui, Kaspar Rufibach, Jon A. Wellner. Limit distribution theory for maximum likelihood estimation of a log-concave density. 2009. hal-00363228

HAL Id: hal-00363228 https://hal.science/hal-00363228

Preprint submitted on 20 Feb 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LIMIT DISTRIBUTION THEORY FOR MAXIMUM LIKELIHOOD ESTIMATION OF A LOG-CONCAVE DENSITY

By Fadoua Balabdaoui , Kaspar Rufibach $^{*}\,$, and Jon A. Wellner $^{\dagger}\,$

We find limiting distributions of the nonparametric maximum likelihood estimator (MLE) of a log-concave density, i.e. a density of the form $f_0 = \exp \varphi_0$ where φ_0 is a concave function on \mathbb{R} . Existence, form, characterizations and uniform rates of convergence of the MLE are given by Rufibach (2006) and Dümbgen and Rufibach (2007). The characterization of the log-concave MLE in terms of distribution functions is the same (up to sign) as the characterization of the least squares estimator of a convex density on $[0,\infty)$ as studied by Groeneboom, Jongbloed and Wellner (2001b). We use this connection to show that the limiting distributions of the MLE and its derivative are, under comparable smoothness assumptions, the same (up to sign) as in the convex density estimation problem. In particular, changing the smoothness assumptions of Groeneboom, Jongbloed and Wellner (2001b) slightly by allowing some higher derivatives to vanish at the point of interest, we find that the pointwise limiting distributions depend on the second and third derivatives at 0 of H_k , the "lower invelope" of an integrated Brownian motion process minus a drift term depending on the number of vanishing derivatives of $\varphi_0 = \log f_0$ at the point of interest. We also establish the limiting distribution of the resulting estimator of the mode $M(f_0)$ and establish a new local asymptotic minimax lower bound which shows the optimality of our mode estimator in terms of both rate of convergence and dependence of constants on population values.

1. Introduction.

1.1. Log-concave densities. A probability density f on the real line is called log-concave if it can be written as

$$f(x) = \exp \varphi(x)$$

1

^{*}Partially supported by the Swiss National Science Foundation

 $^{^{\}dagger}\text{Research}$ supported in part by NSF grant DMS-0503822 and NI-AID grant 2R01 AI291968-04

AMS 2000 subject classifications: Primary 62N01, 62G20; secondary 62G05

Keywords and phrases: asymptotic distribution, integral of Brownian motion, invelope process, log–concave density estimation, lower bounds, maximum likelihood, mode estimation, nonparametric estimation, qualitative assumptions, shape constraints, strongly unimodal, unimodal

2 FADOUA BALABDAOUI, KASPAR RUFIBACH, AND J. A. WELLNER

for some concave function $\varphi : \mathbb{R} \to [-\infty, \infty)$. We let \mathcal{LC} denote the class of all log-concave densities on \mathbb{R} . As shown by Ibragimov (1956), a density function f is log-concave if and only if its convolution with any unimodal density is again unimodal. Thus the class of log-concave densities is often referred to as the class of "strongly unimodal" densities. Furthermore, the class \mathcal{LC} of log-concave densities is exactly the class of Polya frequency functions of order 2, PFF_2 as noted by Pal, Woodroofe and Meyer (2007); see also Dharmadhikari and Joag-Dev (1988), page 150, and Marshall and Olkin (1979), page 492.

The log–concave shape constraint is appealing for many reasons:

(1) Many parametric models, for a certain range of their parameters, are in fact log-concave, e.g. normal, uniform, gamma (r, λ) for $r \ge 1$, beta(a, b) for $a \ge 1$ and $b \ge 1$, generalized Pareto, Gumbel, Fréchet, logistic or Laplace, to mention only some of these models. Therefore, assuming log-concavity offers a flexible non-parametric alternative to purely parametric models. Note that a log-concave density need not be symmetric.

(2) Every log-concave density is automatically unimodal. Furthermore, logconcavity of a density f immediately implies specific shape constraints for certain functions derived from f, see Barlow and Proschan (1975), Marshall and Olkin (2007), Marshall and Olkin (1979), Dharmadhikari and Joag-Dev (1988), An (1998), Bagnoli and Bergstrom (2005), and Section 7 in Dümbgen et al. (2007). Thus, having an estimator (and its limiting distribution) for f at hand provides almost automatically estimators (and limiting distributions) for those functions. Corollary 2.3 illustrates this for the hazard rate. (3) Although the nonparametric MLE of a unimodal density does not exist (see e.g. Birgé (1997), the nonparametric MLE of a log-concave density exists, is unique, and has desirable consistency and rates of convergence properties. Thus the class of log-concave (or strongly unimodal) densities may be a useful and valuable surrogate for the larger class \mathcal{U} of unimodal densities.

(4) Tests for multimodality and mixing can be based on a semiparametric model with densities of the form $f_{c,\varphi}(x) = \exp(\varphi(x) + cx^2)$ where φ is concave and c > 0 as shown by Walther (2002).

(5) Chang and Walther (2006) further show that the EM-algorithm can be extended to work for log-concave component densities.

(6) First attempts to estimate a log-concave density in \mathbb{R}^d were made by Cule et al. (2007).

(7) The log-concave density estimator can be used to improve accuracy in the estimation of the so-called "tail index" of a generalized Pareto distribution, see Müller and Rufibach (2006).

(8) It should be noted that no arbitrary choices such as bandwidth, kernel, or prior are involved in the estimation of a log-concave density; these are all obviated by this shape restriction.

(9) We expect good adaptivity properties of the MLE \hat{f}_n in the class \mathcal{LC} .

Here is some elaboration on this last point. It is fairly well-known that nonparametric maximum likelihood estimators of monotone functions have desirable adaptation properties with respect to the smoothness of the underlying true monotone function: see Birgé (1989) for a study of the Grenander estimator of a monotone density; see Low and Kang (2002) for the construction of an adaptive estimator (at a point) in the context of the white noise model; and see Cai and Low (2005) for an illuminating discussion of the issues. Cai and Low (2007) have initiated a study of adaptive estimation in the setting of convex function estimation, but much more remains to be done. Dümb
gen and Rufibach (2007) show that the MLE $\widehat{\varphi}_n$ of
 φ_0 adapts to the local smoothness of φ_0 in the following sense: if φ_0 is $\mathrm{H\ddot{o}lder}(\beta, L)$ on $[A, B] \subset$ $[x: f_0(x) > 0] \text{ where } 1 \le \beta \le 2, \text{ then } \|\widehat{\varphi}_n - \varphi_0\|_a^b = O_p((n^{-1}\log n)^{\beta/(2\beta+1)})$ for any A < a < b < B and $\|g\|_a^b := \sup_{a \le x \le b} |g(x)|$. This carries over similarly to yield local adaptivity properities of \widehat{f}_n . Current evidence suggests that the nonparametric MLE's of convex densities as studied in Groeneboom, Jongbloed and Wellner (2001b) and of log-concave densities as in Dümbgen and Rufibach (2007) are also adaptive to local smoothness in terms of their local limiting distributions. We intend to investigate this in more detail in future work.

For properties of (random variables with) log-concave densities we refer to Dharmadhikari and Joag-Dev (1988), Marshall and Olkin (1979), and Rufibach (2006). Log-concavity of a density f implies certain shape constraints for functions derived from f, such as the distribution function, the tail or hazard function. See An (1998) for comparisons with the related notion of a log-convex density.

1.2. Log-concave density estimation. Now let $X_{(1)} < X_{(2)} < \cdots < X_{(n)}$ be the order statistics of n independent random variables X_1, \ldots, X_n distributed according to a log-concave probability density $f_0 = \exp \varphi_0$ on \mathbb{R} . The distribution function corresponding to f_0 is denoted by F_0 .

The maximum likelihood estimator (MLE) of a log-concave density was introduced in Rufibach (2006) and Dümbgen and Rufibach (2007). Algorithmic aspects were treated in Rufibach (2007) and in a more general framework in Dümbgen, Hüsler and Rufibach (2007), while consistency with respect to the Hellinger metric was established by Pal, Woodroofe and Meyer (2007), and rates of convergence of \hat{f}_n and \hat{F}_n were established by Dümbgen and

Rufibach (2007). Since the derivation of the MLE of a log–concave density is extensively treated in these references, we only briefly recall its definition and the properties relevant for this paper.

If \mathcal{C} denotes the class of all concave functions $\varphi : \mathbb{R} \to [-\infty, \infty)$, the estimator $\widehat{\varphi}_n$ of φ_0 is the maximizer of the "adjusted" criterion function

$$L(\varphi) = \int_{\mathbb{R}} \varphi(x) d\mathbb{F}_n(x) - \int_{\mathbb{R}} \exp \varphi(x) dx$$

over \mathcal{C} , where \mathbb{F}_n is the empirical distribution function of the observations. The log-concave density estimator is then $\widehat{f}_n := \exp \widehat{\varphi}_n$, which exists and is unique.

1.3. Some properties of $\widehat{\varphi}_n$. For any continuous piecewise linear function $h_n : [X_{(1)}, X_{(n)}] \to \mathbb{R}$ such that the knots of h_n coincide with (some of) the order statistics $X_{(1)}, \ldots, X_{(n)}$, introduce the set of knots $\widehat{\mathcal{S}}_n(h_n)$ of h_n as

$$\widehat{\mathcal{S}}_n(h_n) := \{ t \in (X_{(1)}, X_{(n)}) : h'_n(t-) > h'_n(t+) \} \cup \{ X_{(1)}, X_{(n)} \}$$

Dümbgen and Rufibach (2007) found that $\widehat{\varphi}_n$ is piecewise linear, that $\widehat{\varphi}_n = -\infty$ on $\mathbb{R} \setminus [X_{(1)}, X_{(n)}]$ and that the knots of $\widehat{\varphi}_n$ only occur at (some of the) ordered observations $X_{(1)} < \cdots < X_{(n)}$. The latter property is entirely different from the estimation of a k-monotone density for k > 1 (see below), where the knots fall strictly between observations with probability equal to 1. According to Theorem 2.4 in Dümbgen and Rufibach (2007), the estimator $\widehat{\varphi}_n$ has the following characterization. For $x \geq X_{(1)}$ (recall that $\widehat{\varphi}_n := 0$ outside $[X_{(1)}, X_{(n)}]$) define the processes

$$\begin{split} \widehat{F}_n(x) &:= \int_{X_{(1)}}^x \exp(\widehat{\varphi}_n(t)) dt, \quad \widehat{H}_n(x) := \int_{X_{(1)}}^x \widehat{F}_n(t) dt, \\ \mathbb{H}_n(x) &:= \int_{X_{(1)}}^x \mathbb{F}_n(t) dt = \int_{-\infty}^x \mathbb{F}_n(t) dt. \end{split}$$

Then the concave function $\widehat{\varphi}_n$ is the MLE of the log–density φ_0 if and only if

(1.1)
$$\widehat{H}_n(x) \quad \begin{cases} \leq \mathbb{H}_n(x) & \text{for all } x \geq X_{(1)} \\ = \mathbb{H}_n(x) & \text{if } x \in \widehat{\mathcal{S}}_n(\widehat{\varphi}_n). \end{cases}$$

1.4. Other shape constraints. Maximum likelihood estimation of a monotone density f_0 on $[0, \infty)$ was first studied by Grenander (1956) who found that a function \hat{f}_n is the estimator of f_0 if and only if it is equal to the

left derivative of the least concave majorant of \mathbb{F}_n . Prakasa Rao (1969) established the asymptotic distribution theory at a point $x_0 > 0$ such that $f'_0(x_0) < 0$ and f'_0 is continuous in a neighborhood of x_0 :

$$n^{1/3}(\hat{f}_n(x_0) - f_0(x_0)) \rightarrow_d |f'_0(x_0)f_0(x_0)/2|^{1/3}\mathbb{Z}$$

where \mathbb{Z} is the slope at zero of the (least) concave majorant of the process $W(t) - t^2$, $t \in \mathbb{R}$ for two-sided Brownian motion W starting at 0.

Under the assumption that the true density f_0 is convex and non-increasing on $[0, \infty)$, Groeneboom, Jongbloed and Wellner (2001b) defined and characterized the MLE \hat{f}_n as well as the least squares estimator of f_0 . Here, at any point $x_0 > 0$ such that $f''_0(x_0) > 0$ and f''_0 is continuous in a neighborhood of x_0 ,

$$n^{2/5}(\hat{f}_n(x_0) - f_0(x_0)) \to_d (24^{-1}f_0^2(x_0)f_0''(x_0))^{1/5} \mathbb{H}''(0),$$

where \mathbb{H} is a random cubic spline such that \mathbb{H}'' is convex and \mathbb{H} stays above integrated two-sided Brownian motion $+t^4$ and touches the Gaussian process exactly at those points where \mathbb{H}'' changes its slope, see Groeneboom, Jongbloed and Wellner (2001a).

The classes of monotone and convex decreasing densities are particular cases of the class of k-monotone densities. A density function p on $[0, \infty)$ is 1-monotone if it is non-increasing; it is 2-monotone if it is non-increasing and convex; and it is k-monotone for $k \geq 3$ if and only if $(-1)^j p^{(j)}$ is nonnegative, non-increasing, and convex for $j = 0, \ldots, k - 2$. Balabdaoui and Wellner (2007) were able to adapt the approach of Groeneboom, Jongbloed and Wellner (2001b) to this general class of densities. However, their result depends on the validity of a conjecture about an upper bound for the error in a particular Hermite interpolation via odd-degree splines.

We find that log-concave estimation shares many similarities with the aforementioned shape-constrained estimation problems. In particular the limiting distribution of the MLE, our nonparametric estimator, involves a stochastic process whose second derivative is concave, and which stays below an integrated Brownian motion minus t^{k+2} . The even integer k determines the number of vanishing derivatives of the true concave function φ_0 at the estimation point x_0 .

1.5. Organization of the paper. In Section 2, we establish the limiting distributions of the MLE estimators, $\hat{\varphi}_n$ and \hat{f}_n , at a fixed point $x_0 \in \mathbb{R}$ under some specified working assumptions. The characterization of either $\hat{\varphi}_n$ or \hat{f}_n given in (1.1) coincides, except for the direction of the inequality, with

that of the least squares estimator of a convex decreasing density, studied by Groeneboom et al. (2001b); see their Lemma 2.2, p. 1657. This enables us to adopt the general scheme of the proof in their paper.

Log–concave densities f and their logarithm φ can easily have vanishing second and higher derivatives at fixed points as can be seen from the density function

$$f_0(x) = \sqrt{2} \frac{\Gamma(3/4)}{\pi} \exp(-x^4), \ x \in \mathbb{R}.$$

In this case $\varphi_0^{(j)}(x_0) = 0, j = 1, 2, 3$ for $x_0 = 0$, and $\varphi_0^{(4)}(x_0) \neq 0$. The following "tilted" version of f_0 shows that vanishing second derivatives of φ_0 can also occur at points other than the mode of f:

$$\tilde{f}_0(x) = \exp(a + bx)f_0(x) = \tilde{a}\exp(bx - x^4)$$

where $\tilde{a} = \tilde{a}(b) := 1/\int_{\mathbb{R}} \exp(bx - x^4) dx$; in this case $\tilde{\varphi}_0 := \log \tilde{f}_0$ satisfies $\tilde{\varphi}_0''(0) = 0$, but the mode $\tilde{m}_0 := M(\tilde{f}_0) = (b/4)^{1/3} > 0$ when b > 0, and $\tilde{\varphi}_0''(\tilde{m}_0) = -12(b/4)^{2/3} < 0$. Thus the formulation of our asymptotic results allows higher derivatives of the concave function φ_0 to vanish at the estimation point. This is somewhat more general than the assumptions of Groeneboom, Jongbloed and Wellner (2001b) (where a natural assumption is that the second derivative is positive at the point of interest, but similar vanishing of second derivatives and existence of a non-zero higher order derivative can also easily occur), but it is analogous to the results of Wright (1981) and Leurgans (1982) for nonparametric estimation of a monotone regression function. Similar results for the Grenander estimator of a monotone tone density are stated by Anevski and Hössjer (2002). We find that the respective limiting distributions of the MLE and its first derivative depend on a stochastic process, H_k , equal almost surely to the "lower invelope" (or just "invelope") on \mathbb{R} of the integrated Brownian motion minus t^{k+2} where k is the order of the first non-zero derivative of φ_0 at the point of interest.

In Section 3, the estimation point x_0 is taken to be equal to the mode, m_0 , defined to be the smallest point in the modal interval of the log-concave density f_0 . A natural estimator of m_0 , which we denote by \widehat{M}_n , can be taken to be the smallest number maximizing the MLE $\widehat{\varphi}_n$, or equivalently the smallest number maximizing the MLE \widehat{f}_n . In this section, we establish our second main result: the asymptotic distribution of \widehat{M}_n . Under the asymptotic that the second derivative $f''_0(m_0) < 0$, we show that this distribution depends on the random variable defined to be the argmax or mode of $H_2^{(2)}$ on \mathbb{R} . When the second, third, and higher derivatives of order k - 1 or lower vanish at

 m_0 but $f_0^{(k)}(m_0) < 0$, then the limit distribution depends on the mode of $H_k^{(2)}$.

Proofs are deferred to Section 4.

To illustrate all the quantities for which we provide limiting distributions, in Figure 1 we give plots of \hat{f}_n , $\hat{\varphi}_n$, \hat{F}_n , and $\hat{\lambda}_n = \hat{f}_n/(1-\hat{F}_n)$, based on two samples of sizes n = 20 and n = 200 drawn from a Gamma(2, 1) density $f_0(x) = xe^{-x}\mathbf{1}_{[0,\infty)}(x)$. All these plots were generated using the R-package logcondens, see Rufibach and Dümbgen (2007).

2. Limiting distribution theory. To state the main result, we make the following assumptions.

2.1. Assumptions. Fix $x_0 \in \mathbb{R}$. We suppose that the true density $f_0 = \exp \varphi_0$ satisfies the following assumptions:

- (A1) The density function $f_0 \in \mathcal{LC}$.
- (A2) $f_0(x_0) > 0.$
- (A3) The function φ_0 is at least twice continuously differentiable in a neighborhood of x_0 .
- (A4) If $\varphi_0''(x_0) \neq 0$, then k = 2. Otherwise, suppose that k is the smallest integer such that $\varphi_0^{(j)}(x_0) = 0, j = 2, \dots, k-1, \varphi_0^{(k)}(x_0) \neq 0$, and $\varphi_0^{(k)}$ is continuous in a neighborhood of x_0 .

Note that concavity of φ_0 and A3 and A4 imply that k is necessarily even and that $\varphi_0^{(k)}(x_0) < 0$. Indeed, suppose that k > 2. Using Taylor expansion of φ_0'' up to degree k - 2, there exists a small h > 0 for which we can write

$$\varphi_0''(x) = \frac{\varphi_0^{(k)}(x_0)}{(k-2)!} (x-x_0)^{k-2} + o((x-x_0)^{k-2}), \qquad x \in [x_0-h, x_0+h].$$

Since $\varphi_0''(x) \leq 0$ for all $x \in [x_0 - h, x_0 + h]$, it follows that k - 2 is even; i.e. k even and $\varphi_0^{(k)}(x_0) < 0$.

2.2. Notation. Let W denote two-sided Brownian motion, starting at 0. For $t \in \mathbb{R}$, define:

(2.2)
$$Y_k(t) = \begin{cases} \int_0^t W(s)ds - t^{k+2} & \text{if } t \ge 0\\ \int_t^0 W(s)ds - t^{k+2} & \text{if } t < 0. \end{cases}$$

For the uniform norm of a bounded function f we write $||f||_{\infty} = \sup_{x \in \mathbb{R}} |f(x)|$. The derivative of $\widehat{\varphi}_n$ at $x \in \mathbb{R}$ is as usual denoted by $\widehat{\varphi}'_n(x)$. However, if $x \in \widehat{\mathcal{S}}_n(\widehat{\varphi}_n)$, then we define $\widehat{\varphi}'_n(x)$ as the left-derivative.



FIG 1. Examples for log-concave density, log-density, CDF, and hazard rate estimation for n = 20,200 (-- true functions, - estimators). The dotted vertical lines indicate the set $\hat{S}_n(\hat{\varphi}_n)$. The \cdot -- vertical lines are placed at the mode of the estimated density.

THEOREM 2.1. Suppose that A1 - A4 hold. Then

$$\begin{pmatrix} n^{k/(2k+1)} \left(\widehat{f}_n(x_0) - f_0(x_0) \right) \\ n^{(k-1)/(2k+1)} \left(\widehat{f}'_n(x_0) - f'_0(x_0) \right) \end{pmatrix} \to_d \begin{pmatrix} c_k(x_0,\varphi_0) \ H_k^{(2)}(0) \\ d_k(x_0,\varphi_0) \ H_k^{(3)}(0) \end{pmatrix}$$

and

$$\begin{pmatrix} n^{k/(2k+1)} \left(\widehat{\varphi}_n(x_0) - \varphi_0(x_0) \right) \\ n^{(k-1)/(2k+1)} \left(\widehat{\varphi}'_n(x_0) - \varphi'_0(x_0) \right) \end{pmatrix} \to_d \begin{pmatrix} C_k(x_0, \varphi_0) \ H_k^{(2)}(0) \\ D_k(x_0, \varphi_0) \ H_k^{(3)}(0) \end{pmatrix}.$$

where H_k is the "lower invelope" of the process Y_k ; that is,

$$\begin{split} H_k(t) &\leq Y_k(t) \quad \text{for all } t \in \mathbb{R}; \\ H_k^{(2)} \text{ is concave}; \\ H_k(t) &= Y_k(t) \text{ if the slope of } H_k^{(2)} \text{ decreases strictly at } t. \end{split}$$
The constants c_k , d_k , C_k , and D_k are given by

(2.3)
$$c_k(x_0,\varphi_0) = \left(\frac{f_0(x_0)^{k+1}|\varphi_0^{(k)}(x_0)|}{(k+2)!}\right)^{1/(2k+1)}$$

(2.4)
$$d_k(x_0,\varphi_0) = \left(\frac{f_0(x_0)^{k+2}|\varphi_0^{(k)}(x_0)|^3}{[(k+2)!]^3}\right)^{1/(2k+1)}$$

(2.5)
$$C_k(x_0,\varphi_0) = \left(\frac{|\varphi_0^{(k)}(x_0)|}{f_0(x_0)^k(k+2)!}\right)^{1/(2k+1)}$$

(2.6)
$$D_k(x_0,\varphi_0) = \left(\frac{|\varphi_0^{(k)}(x_0)|^3}{f_0(x_0)^{k-1}[(k+2)!]^3}\right)^{1/(2k+1)}$$

COROLLARY 2.2. Suppose that A1 - A4 hold with k = 2. Then

$$\left(\begin{array}{c} n^{2/5} \left(\widehat{f}_n(x_0) - f_0(x_0)\right) \\ n^{1/5} \left(\widehat{f}'_n(x_0) - f'_0(x_0)\right) \end{array}\right) \to_d \left(\begin{array}{c} c_2(x_0,\varphi_0) \ H_2^{(2)}(0) \\ d_2(x_0,\varphi_0) \ H_2^{(3)}(0) \end{array}\right)$$

and

$$\left(\begin{array}{c} n^{2/5} \left(\widehat{\varphi}_n(x_0) - \varphi_0(x_0)\right) \\ n^{1/5} \left(\widehat{\varphi}'_n(x_0) - \varphi'_0(x_0)\right) \end{array}\right) \to_d \left(\begin{array}{c} C_2(x_0,\varphi_0) \ H_2^{(2)}(0) \\ D_2(x_0,\varphi_0) \ H_2^{(3)}(0) \end{array}\right).$$

where H_2 is the (concave) invelope of the process Y_2 ; that is,

$$H_2(t) \leq Y_2(t) \quad for \ all \ t \in \mathbb{R};$$

10 FADOUA BALABDAOUI, KASPAR RUFIBACH, AND J. A. WELLNER

 $H_2^{(2)}$ is concave; $H_2(t) = Y_2(t)$ if the slope of $H_2^{(2)}$ decreases strictly at t. The constants c_2 , d_2 , C_2 and D_2 are given by (2.3) - (2.6) with k = 2.

Note that the constants $C_2(x_0, \varphi_0)$ and $D_2(x_0, \varphi_0)$, up to inversion of $f_0(x_0)$, exhibit a structure very similar to that of the constants given by Groeneboom, Jongbloed and Wellner (2001b) in the problem of estimating a convex density g_0 on $[0, \infty)$. We recall here that in the latter problem, those constants are found to be equal to (we use our notation to make the comparison easy)

$$c_2(x_0,g_0) = \left(\frac{g_0(x_0)^2 g_0^{(2)}(x_0)}{4!}\right)^{1/5}, \quad d_2(x_0,g_0) = \left(\frac{g_0(x_0)(g_0^{(2)}(x_0))^3}{(4!)^3}\right)^{1/5}.$$

It is clear that φ_0 in the log–concave problem plays exactly the same role as f_0 in the problem of estimating a convex density. However, in the first case estimation is based on observations which are distributed according to $\exp \varphi_0$, whereas in the latter the data come from f_0 itself. A good insight into the difference between the expressions of the asymptotic constants can be gained from the proof of Theorem 4.6 in Section 4. There, we show that the leading coefficient of the drift of the limiting process Y_k depends on $\varphi_0^{(k)}(x_0)f_0(x_0) = f_0^{(k)}(x_0) - (\varphi_0'(x_0))^k f_0(x_0)$, where the second term is "filtered out" in the Taylor expansion of the estimation error in the neighborhood of x_0 . Hence, $|\varphi_0^{(k)}(x_0)| \cdot f_0(x_0)$ can be viewed as the dominating term replacing $|g_0^{(k)}(x_0)|$ in the convex estimation problem. For k=2 the constants $c_2(x_0, \varphi_0)$ and $d_2(x_0, \varphi_0)$ given in (2.3) and (2.4) with k = 2 match closely with $c_2(x_0, g_0)$ and $d_2(x_0, g_0)$ obtained by Groeneboom, Jongbloed and Wellner (2001b) in the convex estimation problem, with $f_0(x_0)$ in the numerator, whereas $f_0(x_0)$ shows up in the denominator in the asymptotic constants $C_2(x_0, \varphi_0)$ and $D_2(x_0, \varphi_0)$. This results from applying the deltamethod to $\widehat{f}_n(x_0) = \exp(\widehat{\varphi}_n(x_0))$ and $\widehat{f}'_n(x_0) = \widehat{\varphi}'_n(x_0)\widehat{f}_n(x_0)$ which yields $C_2(x_0,\varphi_0)$ and $D_2(x_0,\varphi_0)$

Finally, and in order to compare also the random parts of the limits in the convex and log-concave estimation problems, we would like to note that for our lower invelope process H_k , $-H_k$ has the same distribution as the "upper invelope" of $-Y_k$, which was called just the "invelope" in the case k = 2 by Groeneboom et al. (2001b): The process $-Y_k$ has a drift equal to plus t^{k+2} which specializes to t^4 in the convex density problem with k = 2. This "upper invelope" stays above $-Y_k$ and admits a convex second derivative. Since -W has the same distribution as W, it follows that the

upper and lower invelopes \mathbb{H}_k and H_k (associated with estimation of convex and concave functions respectively) satisfy $\mathbb{H}_k \stackrel{d}{=} -H_k$. Since the derivatives at zero $\mathbb{H}_k^{(2)}(0)$ and $\mathbb{H}_k^{(3)}(0)$ of \mathbb{H}_k are distributed symmetrically about zero, the same is true of the derivatives at zero $H_k^{(2)}(0)$ and $H_k^{(3)}(0)$ of H_k .

As shown by Barlow and Proschan (1975), Lemma 5.8, page 77, (see also Marshall and Olkin (1979), p. 493; Marshall and Olkin (2007), p. 102; An (1998); and Bagnoli and Bergstrom (2005)), if f_0 is log-concave, then the hazard function

$$\lambda_0(x) = \frac{f_0(x)}{1 - F_0(x)} \mathbf{1}_{\{x < F_0^{-1}(1)\}}$$

is monotone non-decreasing. Defining the estimator of λ_0 based on f_n as

$$\hat{\lambda}_n(x) = \frac{\hat{f}_n(x)}{1 - \hat{F}_n(x)} \mathbf{1}_{\{x < X_{(n)}\}},$$

application of the delta-method yields the following corollary.

COROLLARY 2.3. Suppose that A1 - A4 hold. Then

$$\begin{pmatrix} n^{k/(2k+1)} \left(\widehat{\lambda}_n(x_0) - \lambda_0(x_0) \right) \\ n^{(k-1)/(2k+1)} \left(\widehat{\lambda}'_n(x_0) - \lambda'_0(x_0) \right) \end{pmatrix} \to_d \begin{pmatrix} g_k(x_0,\varphi_0) \ H_k^{(2)}(0) \\ h_k(x_0,\varphi_0) \ H_k^{(3)}(0) \end{pmatrix}$$

where the constants g_k and h_k are given by

$$g_k(x_0, \varphi_0) = c_k(x_0, \varphi_0) / (1 - F_0(x_0))$$

$$h_k(x_0, \varphi_0) = d_k(x_0, \varphi_0) / (1 - F_0(x_0))$$

For a more thorough discussion of the implications for the hazard rate if f_0 is log-concave see Dümbgen and Rufibach (2007), at the end of Section 3 and Dümbgen et al. (2007), Section 7.

3. Inference about the mode of f_0 . Estimation of the mode of a unimodal density has been considered by many authors; see e.g. Parzen (1962), Chernoff (1964), Grenander (1965), Dalenius (1965), Venter (1967), Wegman (1970a), Wegman (1970b), Wegman (1971), Eddy (1980), Eddy (1982), Hall (1982), Müller (1989), Romano (1988), Vieu (1996), and more recently Meyer (2001), and Herrmann and Ziegler (2004).

12 FADOUA BALABDAOUI, KASPAR RUFIBACH, AND J. A. WELLNER

Empirical studies of the performance of various estimators are given by Dalenius (1965), Ekblom (1972), Meyer (2001), and Meyer and Woodroofe (2004). Many of the methods considered for estimating the mode of a unimodal smooth density use kernel estimation, but others are based on the principle of substitution with another choice of estimator of the population density: for example, the estimators of Venter (1967), are related to nearest - neighbor estimators of the density f_0 . All the estimators of the mode in the class of unimodal densities known to us involve some more more or less ad-hoc choice essentially because the maximum likelihood estimator of a unimodal density is not well-defined as is nicely explained by Birgé (1997). (Note that Wegman (1970b), Wegman (1971) discussed the nonparametric MLE of a unimodal density subject to a constraint on the height of the mode; without some constraint of this type, the MLE does not exist.)

For virtually all of the estimators of which we are aware, some choice of a smoothing parameter or bandwidth or constraint is required. Empirical choice of smoothing parameters has been studied by Müller (1989) who studied local methods of choosing the smoothing parameter, Grund and Hall (1995) who studied bootstrap methods, and Ziegler (2004) who studied plugin methods. Klemelä (2005) gave a construction of adaptive estimators based on Lepski's method (Lepskiĭ (1992)). For nonparametric Bayes estimators of unimodal densities and hence of the mode, see Brunner and Lo (1989), Ho (2006a), and Ho (2006b); for these estimators, choice of a prior is equivalent to a choice of smoothing parameters.

In contrast, estimation in the (large!) subclass of log-concave (or strongly unimodal) densities is much simpler, avoiding bandwidth or smoothing parameter choices completely. Since the maximum likelihood estimator exists, we can simply estimate the mode by the mode (or smallest point in a modal interval) of the MLE \hat{f}_n . Using the notation introduced by Eddy (1982) (and also used by Romano (1988)), we let $\widehat{M}_n := M(\widehat{f}_n)$ where M denotes the mode functional (or "smallest argmax" functional) given by

$$M(g) := \min\{t : g(t) = \max_{u \in \mathbb{R}} g(u)\}.$$

Because of the adaptive properties of the MLE's \widehat{f}_n of f_0 and $\widehat{\varphi}_n$ of φ_0 discussed in Section 1, we expect \widehat{M}_n to adapt to different local smoothness (or peakedness) hypotheses on f_0 (much as the Grenander estimator is locally adaptive in the case of estimating a monotone density, see e.g. Birgé (1989), page 1535). Here we study \widehat{M}_n as an estimator of the mode $M(f_0) := m_0$ under just the condition that f_0 has a continuous second derivative f''_0 in a neighborhood of m_0 with $f''_0(m_0) < 0$. We begin in the next subsection with

a new asymptotic minimax lower bound for estimation of m_0 under this hypothesis. The following subsection gives our new limiting distribution result for the MLE \widehat{M}_n of the mode m_0 .

3.1. New lower bounds for estimating the mode.

Has'minskii (1979) established a lower bound for estimation of the mode m_0 of a unimodal density $f \in \mathcal{U}$ assuming that f satisfies $f''(m_0) < 0$. He showed that the best local asymptotic minimax rate of convergence for any estimator of m_0 is $n^{-1/5}$. Has'minskii based his proof on a sequence of parametric submodels of the form

$$f_n(x,\theta) = f(x) + \theta n^{-2/5} g(n^{1/5}(x-m_0))$$

where, for $a := -f''(m_0)$,

$$g(x) := g_a(x) = \begin{cases} x, & \text{if } |x| \le 1/a, \\ 0, & \text{if } |x| \ge K > 1/a, \end{cases}$$

and $g := g_a$ satisfies g(-x) = -g(x) and |g''(x)| < a/2 for all $x \in \mathbb{R}$. However, Has'minskii (1979) did not study the dependence of the local minimax bound on $a = -f''(m_0)$ and $f(m_0)$, leaving his bound in terms of $c_0^2 := f(m_0) / \int g_a^2(x) dx$ involving the still unspecified function $g = g_a$.

Here we consider different parametric submodels and derive the dependence of the constant in local asymptotic minimax lower bound for estimation of the mode m_0 in the family \mathcal{LC} of log-concave (or strongly unimodal) densities.

We want to derive asymptotic lower bounds for the local minimax risks for estimating the mode M(f). The L_1 - minimax risk for estimating a functional ν of f_0 based on a sample X_1, \ldots, X_n of size n from f_0 which is known to be in a subset $\mathcal{LC}_{n,\tau}$ of \mathcal{LC} is defined by

(3.7)
$$MMR_1(n, T_n, \mathcal{LC}_{n,\tau}) := \inf_{T_n} \sup_{f \in \mathcal{LC}_{n,\tau}} E_f |T_n - \nu(f)|$$

where the infimum ranges over all possible measurable functions $T_n = t_n(X_1, \ldots, X_n)$ mapping \mathbb{R}^n to \mathbb{R} . The shrinking classes $\mathcal{LC}_{n,\tau}$ used here are Hellinger balls centered at f_0 :

$$\mathcal{LC}_{n,\tau} = \left\{ f \in \mathcal{LC} : H^2(f, f_0) = \frac{1}{2} \int_{-\infty}^{\infty} \left(\sqrt{f(z)} - \sqrt{f_0(z)} \right)^2 dz \le \tau/n \right\}.$$

Consider estimation of

(3.8)
$$\nu(f) := M(f) = \inf\{t \in \mathbb{R} : t = \sup_{u \in \mathbb{R}} f(u)\}.$$

Let $f_0 \in \mathcal{LC}$ and $m_0 = M(f_0)$ be fixed such that f_0 is twice continuously differentiable at m_0 and $f''_0(m_0) < 0$. Consider the family $\{\varphi_{\epsilon}\}_{\epsilon>0}$ and resulting family $\{f_{\epsilon}\}_{\epsilon>0}$ defined as follows:

$$\varphi_{\epsilon}(x) = \begin{cases} \varphi_0(x), & x < m_0 - \epsilon c_{\epsilon} \\ \varphi_0(x), & x > m_0 + \epsilon, \\ \varphi_0(m_0 + \epsilon) \\ + \varphi'_0(m_0 + \epsilon)(x - m_0 - \epsilon), & x \in [m_0 - \epsilon, m_0 + \epsilon] \\ \varphi_0(m_0 - \epsilon c_{\epsilon}) \\ + \varphi'_0(m_0 - \epsilon c_{\epsilon})(x - m_0 + \epsilon c_{\epsilon}), & x \in [m_0 - \epsilon c_{\epsilon}, m_0 - \epsilon) \end{cases}$$

where c_{ϵ} is chosen so that φ_{ϵ} is continuous at $m_0 - \epsilon$. Note that if $\varphi_0(x) = \gamma - \gamma_0 (x - m_0)^2$, then $c_{\epsilon} = 3$ for all ϵ , and $c_{\epsilon} \to 3$ as $\epsilon \downarrow 0$ since $f_0''(m_0) < 0$. Now define

$$h_{\epsilon}(x) := \exp(\varphi_{\epsilon}(x)), \text{ and } f_{\epsilon}(x) := \frac{h_{\epsilon}(x)}{\int h_{\epsilon}(y)dy}.$$

Then f_{ϵ} is log-concave for each $\epsilon > 0$ with mode $m_0 - \epsilon$ by construction, so with $\nu(f_{\epsilon}) := M(f_{\epsilon}) :=$ the mode of f_{ϵ} we have

$$\nu(f_{\epsilon}) - \nu(f_{0}) = M(f_{\epsilon}) - M(f_{0}) = m_{0} - \epsilon - m_{0} = -\epsilon.$$

Furthermore, the following lemma holds.

LEMMA 3.1. Under the above assumptions

$$H^{2}(f_{\epsilon}, f_{0}) = \frac{2f_{0}''(m_{0})^{2}}{5f_{0}(m_{0})}\epsilon^{5} + o(\epsilon^{5}) := \rho\epsilon^{5} + o(\epsilon^{5}).$$

Proof. Proceeding as in Jongbloed (1995),

$$H^{2}(f_{\epsilon}, f_{0}) = \frac{1}{2} \int_{-\infty}^{\infty} [\sqrt{f_{\epsilon}(x)} - \sqrt{f_{0}(x)}]^{2} dx$$

$$= \frac{1}{2} \int_{m_{0} - \epsilon c_{\epsilon}}^{m_{0} + \epsilon} [\sqrt{f_{\epsilon}(x)} - \sqrt{f_{0}(x)}]^{2} dx$$

$$= \frac{2}{5} f_{0}(m_{0}) \varphi_{0}''(m_{0})^{2} \epsilon^{5} + o(\epsilon^{5}) = \frac{2}{5} \frac{f_{0}''(m_{0})^{2}}{f_{0}(m_{0})} \epsilon^{5} + o(\epsilon^{5})$$

as $\epsilon \downarrow 0$. Calculations similar to those of Jongbloed (1995) (see also Jongbloed (2000)) and Groeneboom et al. (2001b)) complete the proof of the lemma.

Taking $\epsilon = cn^{-1/5}$ and defining $f_n := f_{cn^{-1/5}}$ yields

$$\nu(f_n) - \nu(f_0) = M(f_n) - M(f_0) = -cn^{-1/5}$$

and

$$nH^2(f_n, f_0) = \frac{2}{5} \frac{f_0''(m_0)^2}{f_0(m_0)} c^5 + o(1) := \rho c^5 + o(1).$$

Plugging these into the lower bound Lemma 4.1 of Groeneboom (1996) with $\ell(x) := |x|$ yields

$$\liminf_{n} \inf_{T_n} n^{1/5} \max \left\{ E_{n,P_n} | T_n - M(f_n) |, E_{n,P} | T_n - M(f_0) | \right\}$$
$$\geq \frac{1}{4} c \exp(-2\rho c^5) = \frac{e^{-1/5}}{4 \cdot 10^{1/5}} \rho^{-1/5} = (.15512) \left(\frac{f_0(m_0)}{f_0''(m_0)^2} \right)^{1/5}$$

by choosing $c = (10\rho)^{-1/5}$. This yields the following proposition.

PROPOSITION 3.2. (Minimax risk lower bound). Suppose that $\nu(f) = M(f)$ as defined in (3.8), and that $\mathcal{LC}_{n,\tau}$ is as defined above where f_0'' is continuous in a neighborhood of $m_0 = M(f_0)$ with $f_0''(m_0) < 0$. Then

$$\sup_{\tau>0} \limsup_{n\to\infty} n^{1/5} \inf_{T_n} \sup_{f\in\mathcal{LC}_{n,\tau}} E_f |T_n - M(f)| \\ \geq \left(\frac{5/2}{4^5 \cdot e \cdot 10}\right)^{1/5} \left(\frac{f_0(m_0)}{f_0''(m_0)^2}\right)^{1/5} \doteq (.15512) \left(\frac{f_0(m_0)}{f_0''(m_0)^2}\right)^{1/5}$$

REMARK 3.3. Note that the constant $b(f_0, m_0) := (f_0(m_0)/f_0''(m_0)^2)^{1/5}$ appearing on the right side of this lower bound is scale equivariant in exactly the right way: if $f_c(x) := f_0(m_0 + (x - m_0)/c)/c$ for c > 0, then $b(f_c, m_0) = cb(f_0, m_0)$ for all c > 0. The constant $b(f_0, m_0)$ will appear in the limit distribution appearing in the next subsection.

REMARK 3.4. If \mathcal{LC} is replaced by the class \mathcal{U} of unimodal densities on \mathbb{R} and $\mathcal{LC}_{n,\tau}$ is replaced by $\mathcal{U}_{n,\tau}$ defined analogously where f_0 satisfies $f''_0(m_0) < 0$ and f''_0 continuous in a neighborhood of m_0 , then a minimax lower bound of the same form as Proposition 3.2 holds with exactly the same dependence on $b(f_0, m_0) = (f_0(m_0)/f''_0(m_0)^2)^{1/5}$, but with the absolute constant .15512... replaced by .19784.... This can be seen by taking the perturbations $\{f_{\epsilon}\}_{\epsilon>0}$ defined by

$$f_{\epsilon}(x) = \begin{cases} f_0(x), & x \le x_0 - \epsilon, \\ f_0(x), & x > x_0 + \epsilon, \\ f_0(x_0) + b_{\epsilon}(x - x_0 + \epsilon), & x_0 - \epsilon \le x \le x_0 + \epsilon \end{cases}$$

where b_{ϵ} is chosen so that $f_{\epsilon}(x_0 + \epsilon) > f_0(x_0 + \epsilon)$ and $\int_{x_0-\epsilon}^{x_0+\epsilon} f_{\epsilon}(x)dx = \int_{x_0-\epsilon}^{x_0+\epsilon} f_0(x)dx$.

REMARK 3.5. If φ_0 is continuously k-times differentiable in a neighborhood of the mode m_0 , $\varphi_0^{(j)}(m_0) = 0$ for $j = 2, \ldots, k - 1$, and $\varphi_0^{(k)}(m_0) \neq 0$ (Assumption A4), then it can be shown that the minimax rate of convergence is $n^{1/(2k+1)}$ and that the minimax lower bound is proportional to

$$\left(\frac{1}{f_0(m_0)\varphi_0^{(k)}(m_0)^2}\right)^{1/(2k+1)} = \left(\frac{f_0(m_0)}{f_0^{(k)}(m_0)^2}\right)^{1/(2k+1)}$$

where the proportionality constant depends on the largest root of the polynomial $x^k - (k/(k-1))x^{k-1} - (2k-1)/(k-1)$ (which equals 3 when k = 2).

3.2. Limiting distribution for the MLE \widehat{M}_n in \mathcal{LC} .

Now let \widehat{f}_n be the MLE of f in the class \mathcal{LC} of log-concave densities, and let $\widehat{M}_n = M(\widehat{f}_n), m_0 = M(f_0)$. Here is our result concerning the limiting distribution of \widehat{M}_n under the same assumptions on f_0 as in the previous section on lower bounds.

THEOREM 3.6. Suppose that f_0'' is continuous in a neighborhood of $m_0 = M(f_0)$ and that $f_0''(m_0) < 0$. Then

$$n^{1/5}(\widehat{M}_n - m_0) \to_d \left(\frac{(4!)^2 f_0(m_0)}{f_0''(m_0)^2}\right)^{1/5} M(H_2^{(2)}).$$

Note that the limiting distribution depends on a multiple of the same constant $b(f_0, m_0)$ which appears in the asymptotic minimax lower bound of Proposition 3.2, times a universal term $M(H_2^{(2)})$, the mode of the "estimator" $H_2^{(2)}(t)$ of the canonical concave function $-12t^2$ in the limit Gaussian problem: estimate the mode of $f_0(t) = -12t^2$ based on observation of $Y(t) = \int_0^t X(s) ds$ when

$$dX(t) = f_0(t)dt + dW(t).$$

We expect that this distribution, namely the distribution of

$$M(H_2^{(2)}) = \operatorname{argmax}_{t \in \mathbb{R}} H_2^{(2)}(t)$$

will occur in several other problems involving nonparametric estimation of the mode or antimode of convex or concave functions under similar second

derivative hypotheses: for example it seems clear that it will occur as the limiting distribution of the nonparametric estimator of the antimode of a convex bathtub shaped hazard (in the setting of Jankowski and Wellner (2007)); as the limiting distribution of the nonparametric estimator of the antimode of a convex regression function in the setting of Groeneboom, Jongbloed and Wellner (2001b); and as the limiting distribution of the nonparametric estimator of the nonparametric estimator of the mode of a concave regression function.

When $\varphi_0^{(j)}(m_0) = 0$ for j = 2, ..., k - 1, $\varphi_0^{(k)}(m_0) \neq 0$, and $\varphi_0^{(k)}$ is continuous in a neighborhood of m_0 , then an analogous result (with a completely similar proof) holds:

$$n^{1/(2k+1)}(\widehat{M}_n - m_0) \to_d \left(\frac{(k+2)!^2}{f_0(m_0)|\varphi_0^{(k)}(m_0)|^2}\right)^{1/(2k+1)} M(H_k^{(2)}).$$

In particular, when k = 4, the rate of convergence is $n^{1/9}$ and the limit distribution becomes that of

$$\left(\frac{6!^2 f_0(m_0)}{f_0^{(4)}(m_0)^2}\right)^{1/9} M(H_4^{(2)}).$$

Apparently estimation of m_0 becomes considerably more difficult when the second and possibly higher order derivatives of φ_0 vanish.

On the other hand, if φ_0 (or equivalently f_0) is cusp-shaped at m_0 , then the rate of convergence of \widehat{M}_n is $n^{1/3}$ and the local asymptotic minimax rate of convergence is also $n^{1/3}$; we will pursue these issues elsewhere.

4. Proofs for Sections 2 and 3.

Throughout this section we fix k and let

$$\begin{aligned} r_n &:= n^{(k+2)/(2k+1)}, \qquad s_n &:= n^{-1/(2k+1)}, \\ x_n(t) &:= x_{n,k}(t) &:= x_0 + s_n t &:= x_0 + n^{-1/(2k+1)}t, \\ \mathcal{I} &:= \mathcal{I}(x_0, n, k, t) &:= \begin{cases} [x_0, x_n(t)], & t \ge 0, \\ [x_n(t), x_0], & t < 0. \end{cases} \end{aligned}$$

4.1. Preparation: Technical Lemmas and Tightness Results.

First, some notation.

18 FADOUA BALABDAOUI, KASPAR RUFIBACH, AND J. A. WELLNER

 $Local \ processes:$ The local processes \mathbb{Y}_n^{loc} and \widehat{H}_n^{loc} are defined for $t\in\mathbb{R}$ by

$$\mathbb{Y}_{n}^{loc}(t) := r_{n} \int_{x_{0}}^{x_{n}(t)} \left(\mathbb{F}_{n}(v) - \mathbb{F}_{n}(x_{0}) - \int_{x_{0}}^{v} \left(\sum_{j=0}^{k-1} \frac{f_{0}^{(j)}(x_{0})}{j!} (u - x_{0})^{j} \right) du \right) dv$$

and

$$\begin{aligned} \widehat{H}_n^{loc}(t) &:= r_n \int_{x_0}^{x_n(t)} \int_{x_0}^v \left(\widehat{f}_n(u) - \sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j \right) du dv \\ &+ \widehat{A}_n t + \widehat{B}_n \end{aligned}$$

where

(4.9)
$$\hat{A}_n = r_n s_n \left(\widehat{F}_n(x_0) - \mathbb{F}_n(x_0) \right)$$

(4.10)
$$B_n = r_n (H_n(x_0) - \mathbb{H}_n(x_0)).$$

We also define the "modified" local processes

(4.11)
$$\mathbb{Y}_{n}^{locmod}(t) := \frac{r_{n}}{f_{0}(x_{0})} \int_{x_{0}}^{x_{n}(t)} \left(\mathbb{F}_{n}(v) - \mathbb{F}_{n}(x_{0}) - \int_{x_{0}}^{v} \left(\sum_{j=0}^{k-1} \frac{f_{0}^{(j)}(x_{0})}{j!} (u - x_{0})^{j} \right) du \right) dv$$
$$- r_{n} \int_{x_{0}}^{x_{n}(t)} \int_{x_{0}}^{v} \widehat{\Psi}_{k,n,2}(u) du dv,$$

and

$$\mathbb{H}_{n}^{locmod}(t) := r_{n} \int_{x_{0}}^{x_{n}(t)} \int_{x_{0}}^{v} \left(\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0}) - (u - x_{0})\varphi_{0}'(x_{0})\right) du dv$$
(4.12)
$$+ \frac{\hat{A}_{n}t + \hat{B}_{n}}{f_{0}(x_{0})}$$

where $\widehat{\Psi}_{k,n,2}$ is defined below in (4.34).

LEMMA 4.1. Let \mathcal{F} be a collection of functions defined on $[x_0 - \delta, x_0 + \delta]$, with $\delta > 0$ small and let s > 0. Suppose that for a fixed $x \in [x_0 - \delta, x_0 + \delta]$ and R > 0 such that $[x, x + R] \subseteq [x_0 - \delta, x_0 + \delta]$, the collection

$$\mathcal{F}_{x,R} = \{ f_{x,y} := f \mathbb{1}_{[x,y]}, \quad f \in \mathcal{F}, \ x \le y \le x + R \}$$

admits an envelope $F_{x,R}$ such that

$$EF_{x,R}^2(X_1) \leq KR^{2d-1}, \qquad R \leq R_0,$$

for some $d \ge 1/2$ and K > 0 depending only on x_0 and δ Moreover, suppose that

(4.13)
$$\sup_{Q} \int_{0}^{1} \sqrt{\log N(\eta \| F_{x,R} \|_{Q,2}, \mathcal{F}_{x,R}, L_2(Q))} d\eta < \infty.$$

Then, for each $\epsilon > 0$ there exist random variables M_n of order $O_p(1)$ (not depending on x or y and $R_0 > 0$ such that

$$\left|\int f_{x,y}d(\mathbb{F}_n - F_0)\right| \leq \epsilon |y - x|^{s+d} + n^{-\frac{s+d}{2s+1}}M_n \quad for \ |y - x| \leq R_0.$$

Proof. See Kim and Pollard (1990) and Balabdaoui and Wellner (2007), Lemmas 4.4 and 6.1. The special case s = 1 = d is Lemma 4.1 of Kim and Pollard (1990).

LEMMA 4.2. If A3 and A4 hold, then

(4.14)
$$f_0^{(j)}(x_0) = [\varphi_0'(x_0)]^j f_0(x_0)$$
 for $j = 1, \dots, k-1$,

and for j = k,

$$f_0^{(k)}(x_0) = (\varphi_0^{(k)}(x_0) + [\varphi_0'(x_0)]^k) f_0(x_0).$$

Proof. The expressions for $f_0^{(j)}(x_0)$ follow immediately from a recursive argument using the identity $f_0 = \exp \varphi_0$ and the assumption $\varphi_0^{(j)}(x_0) = 0$ for $j = 2, \ldots, k-1$ if k > 2.

Now let $\tau_n^+ := \inf\{t \in \widehat{\mathcal{S}}(\widehat{\varphi}_n) : t > x_0\}$, and $\tau_n^- := \sup\{t \in \widehat{\mathcal{S}}(\widehat{\varphi}_n) : t < x_0\}$.

THEOREM 4.3. If A1 - A4 hold, then

Theorem 4.3 should be compared to Theorem 3.3 of Dümbgen and Rufibach (2007). When their Theorem 3.3 is specialized to the case $\beta = 2$ so that that $\varphi_0''(x) \leq C < 0$ for all $x \in T := [A, B]$, then it yields the following:

If m_n denotes the number of elements in $\mathcal{S}_n(\widehat{\varphi}_n) \cap T$, then for any successive knot points t_{i-1} and t_i in $\mathcal{S}_n(\widehat{\varphi}_n) \cap T$,

(4.16)
$$\sup_{i=2,\dots,m_n} (t_i - t_{i-1}) = O_p(\rho_n^{1/5})$$

where $\rho_n = \log(n)/n$. This is "weaker" in the sense that only the supremum over all knots on a compact interval was considered; but Theorem 3.3 of Dümbgen and Rufibach (2007) is more general in the sense that it provides the correct rate over a whole range of Hölder classes, not only for twice differentiable φ_0 . Bounds generalizing (4.16) to $1 \leq \beta \leq 2$ are then used to get upper bounds on the uniform rate of convergence for $\widehat{F}_n - \mathbb{F}_n$ on T. However, for the estimation problem treated in this paper the localized version of the gap problem given in Theorem 4.3 provides the results of interest here concerning local limiting distribution theory.

Proof of Theorem 4.3. From the first characterization of the estimator \hat{f}_n in Dümbgen and Rufibach (2007), for every function Δ such that $\hat{\varphi}_n + t\Delta$ is concave for a t > 0 small enough, we know that

(4.17)
$$\int_{\mathbb{R}} \Delta(x) d\mathbb{F}_n(x) \leq \int_{\mathbb{R}} \Delta(x) d\widehat{F}_n(x).$$

This is equivalent to

(4.18)
$$\int_{\mathbb{R}} \Delta(x) d(\mathbb{F}_n(x) - F_0(x)) \leq \int_{\mathbb{R}} \Delta(x) (\widehat{f}_n(x) - f_0(x)) dx.$$

Using specific indicator functions for Δ , one can furthermore show that

(4.19)
$$\widehat{F}_n(\tau) \in [\mathbb{F}_n(\tau) - 1/n, \mathbb{F}_n(\tau)]$$

for every $\tau \in \widehat{S}_n(\widehat{\varphi}_n)$, see Rufibach (2006) and Corollary 2.5 of Dümbgen and Rufibach (2007).

Now, the idea is to choose a particular permissible perturbation function Δ that satisfies the following two conditions:

- 1. Δ is "local", i.e. compactly supported on $[\tau_n^-, \tau_n^+]$.
- 2. Δ should "filter" out the unknown error $\hat{f}_n f_0$.

The second requirement means that Δ should be chosen so that

(4.20)
$$\int_{\tau_n^-}^{\tau_n^+} \Delta(x) dx = 0 \qquad \int_{\tau_n^-}^{\tau_n^+} \Delta(x) (x - \overline{\tau}) dx = 0,$$

where $\overline{\tau} := (\tau_n^- + \tau_n^+)/2$ is the mid-point of $[\tau_n^-, \tau_n^+]$. If this is guaranteed, then the right side of (4.18) in the end will only depend on the distance $\tau_n^+ - \tau_n^-$ and $f_0(x_0)$.

Define Δ_0 by

$$\Delta_0(x) = (x - \tau_n^-) \mathbf{1}_{[\tau_n^-, \overline{\tau}]}(x) + (\tau_n^+ - x) \mathbf{1}_{[\overline{\tau}, \tau_n^+]}(x).$$

Since $\widehat{\varphi}_n + t\Delta_0$ is concave for small t > 0, Δ_0 is permissible. It is also compactly supported. However, since Δ_0 is nonnegative, there is no hope that it fulfills the second of the requirements above. We therefore introduce a modified perturbation function

$$\Delta_1(x) = \Delta_0(x) - \frac{1}{4}(\tau_n^+ - \tau_n^-) \mathbf{1}_{[\tau_n^-, \tau_n^+]}(x), \ x \in \mathbb{R}.$$

Clearly, existence of a t > 0 such that $\hat{\varphi}_n + t\Delta_1$ is concave is no longer guaranteed. However, using (4.19),

$$\int \Delta_1(x) d(\mathbb{F}_n - F_0)(x)$$

$$= \int \Delta_1(x) d(\mathbb{F}_n - \widehat{F}_n)(x) + \int \Delta_1(x) d(\widehat{F}_n - F_0)(x)$$

$$(4.21) \leq \frac{\tau_n^+ - \tau_n^-}{4} \left| \int_{\tau_n^-}^{\tau_n^+} d(\mathbb{F}_n - \widehat{F}_n)(x) \right| + \int \Delta_1(x) d(\widehat{F}_n - F_0)(x)$$

(4.22)
$$\leq \frac{\tau_n^+ - \tau_n^-}{2n} + \int \Delta_1(x) (\widehat{f}_n - f_0)(x) dx.$$

To get the inequality in (4.21), we used (4.17) with $\Delta = \Delta_0$ and (4.19). The next step is to get bounds for the integrals in the crucial inequality (4.22). Define

$$R_{1n} := \int \Delta_1(x)(\widehat{f}_n - f_0)(x)dx \quad \text{and} \\ R_{2n} := \int \Delta_1(x)d(\mathbb{F}_n - F_0)(x).$$

Rearranging the inequality in (4.22) and use these definitions yields

$$-R_{1n} \le \frac{\tau_n^+ - \tau_n^-}{2n} - R_{2n}.$$

Consistency of $\widehat{\varphi}_n$ together with $\varphi_0^{(k)}(x_0) < 0$ implies $\tau_n^+ - \tau_n^- = o_p(1)$. Thus it follows from Lemma 4.4 that

$$M_k \left(-\varphi_0^{(k)}(x_0) \right) (\tau_n^+ - \tau_n^-)^{k+2} (1 + o_p(1))$$

$$\leq o_p(1)n^{-1} + O_p(r_n^{-1}) = O_p(r_n^{-1}).$$

This yields the claimed rate, $O_p(n^{-1/(2k+1)})$, for the distance between τ_n^+ and τ_n^- .

LEMMA 4.4. Suppose A1-A4 hold. Then

$$R_{2n} = O_p(r_n^{-1})$$

and

$$R_{1n} = M_k f_0(x_0) \varphi_0^{(k)}(x_0) (\tau_n^+ - \tau_n^-)^{k+2} + o_p((\tau_n^+ - \tau_n^-)^{k+2})$$

where $M_k > 0$ depends only on k and $\varphi_0^{(k)}(x_0) < 0$.

Proof. Define the function $p_n(t) = \hat{\varphi}_n(t) - \varphi_0(t)$ for any $t \in [\tau_n^-, \tau_n^+]$. Then, using Taylor expansion of $h \mapsto \exp(h)$ up to order k, we can find $\theta_{t,n} \in [\tau_n^-, \tau_n^+]$ such that

$$R_{1n} = \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) f_0(t) \Big(\sum_{j=1}^{k-1} \frac{p_n(t)^j}{j!} + \frac{1}{k!} \exp(\theta_{t,n}) p_n(t)^k \Big) dt := \sum_{j=1}^k \frac{S_{nj}}{j!}$$

where

$$S_{nj} := \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) f_0(t) p_n(t)^j dt \quad \text{for } 1 \le j \le k-1 \text{ and}$$

$$S_{nk} := \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) f_0(t) \exp(\theta_{t,n}) p_n(t)^k dt.$$

If we expand $f_0(t)$ around the mid-point $\overline{\tau}$ of $[\tau_n^-, \tau_n^+]$ we get for $1 \le j \le k-1$ and a $\eta_{n,t,j} \in [\tau_n^-, \tau_n^+]$,

$$S_{nj} = \sum_{l=0}^{k-1} \frac{f_0^{(l)}(\bar{\tau})}{l!} \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t)(t-\bar{\tau})^l p_n(t)^j dt + \int_{\tau_n^-}^{\tau_n^+} \frac{f_0^{(k)}(\eta_{n,t,j})}{k!} \Delta_1(t)(t-\bar{\tau})^k p_n(t)^j dt$$

and for j = k,

$$S_{nk} = \sum_{l=0}^{k-1} \frac{f_0^{(l)}(\overline{\tau})}{l!} \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) \exp(\theta_{t,n}) (t-\overline{\tau})^l p_n(t)^k dt + \int_{\tau_n^-}^{\tau_n^+} \frac{f_0^{(k)}(\eta_{n,t,k})}{k!} \Delta_1(t) \exp(\theta_{t,n}) (t-\overline{\tau})^k p_n(t)^k dt.$$

It turns out that the dominating term in R_{1n} is the first term in the Taylor expansion of S_{n1} . All the other terms are of smaller order since both p_n and $(t-\bar{\tau})^l, l>0$ are $o_p(1)$ uniformly in $t \in [\tau_n^-, \tau_n^+]$. We denote this dominating term by Q_{n1} . Since $\hat{\varphi}_n$ is linear on $[\tau_n^-, \tau_n^+]$ write $\hat{\varphi}_n(t) = \hat{\varphi}_n(\bar{\tau}) + (t-\bar{\tau})\hat{\varphi}'_n(\bar{\tau})$. By Taylor expansion of p_n around $\bar{\tau}$ we get

$$\begin{aligned} \frac{Q_{1n}}{f_0(\bar{\tau})} &= \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) p_n(t) dt \\ &= p_n(\bar{\tau}) \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) dt + p'_n(\bar{\tau}) \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) (t - \bar{\tau}) dt \\ &- \sum_{j=2}^k \frac{\varphi_0^{(j)}(\bar{\tau})}{j!} \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) (t - \bar{\tau})^j dt - \int_{\tau_n^-}^{\tau_n^+} \epsilon_n(t) \Delta_1(t) (t - \bar{\tau})^k dt \end{aligned}$$

where the first two terms are zero since (4.20) holds when $\Delta = \Delta_1$ and $\|\epsilon_n\|_{\infty} \to_p 0$ as $\tau_n^+ - \tau_n^- \to_p 0$. Using the fact that

$$\int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) (t - \bar{\tau})^j dt$$

$$(4.23) \qquad = \begin{cases} 0, & \text{for } j = 0 \text{ and } j \text{ odd,} \\ (\tau_n^+ - \tau_n^-)^{j+2} \left(\frac{-j}{2^{(j+2)}(j+1)(j+2)}\right), & \text{for } j \text{ even,} \end{cases}$$

we conclude that

$$Q_{1n} = \frac{k}{2^{(k+2)}k!(k+1)(k+2)} f_0(\bar{\tau})\varphi_0^{(k)}(\bar{\tau}) \left((\tau_n^+ - \tau_n^-)^{k+2} + o_p(1) \right)$$

and the claimed form of R_{1n} in the lemma follows.

For R_{2n} , we proceed along the lines of the proof of Lemma 4.1 in Groeneboom, Jongbloed and Wellner (2001b). This means, we have to line up with the assumption of Theorem 2.14.1 in van der Vaart and Wellner (1996). Therefore, define a generalized version of R_{2n} :

$$R_{2n}^{x,y} = \int_x^y \Delta_1(z) d(\mathbb{F}_n - F_0)(z)$$

24 FADOUA BALABDAOUI, KASPAR RUFIBACH, AND J. A. WELLNER

for $-\infty < x \leq y$. With this function we have for some R > 0 specified later,

$$\sup_{\substack{y:0 \le y-x \le R}} |R_{2n}^{x,y}|$$

= $2 \sup_{\substack{y:0 \le y-x \le R}} \left| \int_{x}^{(x+y)/2} \left(z - x - \frac{1}{4}(y-x) \right) d(\mathbb{F}_n - F_0)(z) \right|$
= $2 \sup_{\substack{y:0 \le y-x \le R}} \left| \int h_{x,y}(z) d(\mathbb{F}_n - F_0)(z) \right|$

where

$$h_{x,y}(z) = \left(z - x - \frac{1}{4}(y - x)\right) \mathbf{1}_{[x,(x+y)/2]}(z) = h(z)\mathbf{1}_{[x,(x+y)/2]}(z).$$

Then the collection of functions

$$\mathcal{F}_{x,R} = \{h1_{[x,(x+y)/2]} : x \le y \le x+R\}$$

is a Vapnik-Chervonenkis subgraph class with envelope function

$$F_{x,R}(z) = ((z-x) + R/4) \mathbf{1}_{[x,x+R]}(z).$$

Finally, Theorem 2.6.7 in van der Vaart and Wellner (1996) yields the entropy condition (4.13).

A log–concave density is always unimodal and the value at the mode is finite, and hence $K := ||f_0||_{\infty}$ is finite. Therefore

$$\begin{split} & EF_{x,R}^2(X_1) \\ & = \int_x^{x+R} (z-x)^2 f_0(z) dz + \frac{R}{2} \int_x^{x+R} (z-x) f_0(z) dz + \frac{R^2}{16} \int_x^{x+R} f_0(z) dz \\ & \leq \left(\frac{K}{3} (z-x)^3 + \frac{RK}{4} (z-x)^2 + \frac{R^2K}{16} z \right) \Big|_{z=x}^{x+R} \\ & = \left. \frac{31}{48} K R^3 . \end{split}$$

It follows from Lemma 4.1 with d = 2 and s = k that $R_{2n} = O_p(r_n^{-1})$. \Box

4.2. Proofs for Section 2.

LEMMA 4.5. For any M > 0, we have

(4.24)
$$\sup_{|t| \le M} |\widehat{\varphi}'_n(x_0 + s_n t) - \varphi'_0(x_0)| = O_p(s_n^{k-1}),$$

(4.25)
$$\sup_{|t| \le M} \left| \widehat{\varphi}_n(x_0 + s_n t) - \varphi_0(x_0) - s_n t \varphi_0'(x_0) \right| = O_p(s_n^k).$$

Furthermore, if we define for any $u \in \mathbb{R}$

$$\hat{e}_n(u) = \hat{f}_n(u) - \sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j - f_0(x_0) \frac{[\varphi_0'(x_0)]^k}{k!} (u - x_0)^k,$$

then

$$\sup_{|t| \le M} \left| \hat{e}_n(x_0 + s_n t) - f_0(x_0) \left(\widehat{\varphi}_n(x_0 + s_n t) - \varphi_0(x_0) - s_n t \varphi_0'(x_0) \right) \right|$$
(4.26)
$$= o_p(s_n^k).$$

Proof. The proof of (4.24) and (4.25) is identical to that of Lemma 4.4 in Groeneboom, Jongbloed and Wellner (2001b) since the characterization of \hat{f}_n given in (1.1) is (up to the direction of the inequality) equivalent to that of the least squares estimator of a convex density.

Now, we prove (4.26). Using Taylor expansion of $h \mapsto \exp(h)$ up to order k around zero, we can write

$$\widehat{f}_{n}(u) - f_{0}(x_{0}) = f_{0}(x_{0}) [\exp(\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0})) - 1]$$

$$(4.27) = f_{0}(x_{0}) \sum_{j=1}^{k} \frac{1}{j!} \left(\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0})\right)^{j} + f_{0}(x_{0})\widehat{\Psi}_{k,n,1}(u)$$

where

$$\widehat{\Psi}_{k,n,1}(u) = \sum_{j=k+1}^{\infty} \frac{1}{j!} (\widehat{\varphi}_n(u) - \varphi_0(x_0))^j.$$

But for any $j \ge 1$,

$$\begin{aligned} (\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0}))^{j} \\ &= [\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0}) - (u - x_{0})\varphi_{0}'(x_{0}) + (u - x_{0})\varphi_{0}'(x_{0})]^{j} \\ &= \sum_{r=1}^{j} {j \choose r} [\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0}) - (u - x_{0})\varphi'(x_{0})]^{r} [\varphi_{0}'(x_{0})]^{j-r} (u - x_{0})^{j-r} \\ (4.28) &+ [\varphi_{0}'(x_{0})]^{j} (u - x_{0})^{j}. \end{aligned}$$

26 FADOUA BALABDAOUI, KASPAR RUFIBACH, AND J. A. WELLNER

Hence, using (4.25) and (A3), we get on the set $\{u \ : \ |u-x_0| \le Mn^{-1/(2k+1)}\}$

$$(\widehat{\varphi}_n(u) - \varphi_0(x_0))^j = o_p(n^{-k/(2k+1)})$$

for all $j \ge k+1$.

In particular, this implies that

(4.29)
$$\widehat{\Psi}_{k,n,1}(u) = o_p(n^{-k/(2k+1)})$$

uniformly in $u \in [x_0 - tn^{-1/(2k+1)}, x_0 + tn^{-1/(2k+1)}]$ where $|t| \le M$, and

$$\widehat{f}_n(u) - f_0(x_0) - f_0(x_0) \left(\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0) \varphi'_0(x_0) \right) - f_0(x_0) \sum_{j=1}^k \frac{\varphi_0^{(j)}(x_0)}{j!} (u - x_0)^j = o_p(n^{-k/(2k+1)}).$$

Using Lemma 4.2, the latter can be rewritten as

$$\widehat{f}_{n}(u) - f_{0}(x_{0}) - f_{0}(x_{0}) \left(\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0}) - (u - x_{0})\varphi_{0}'(x_{0})\right) - \sum_{j=1}^{k-1} \frac{f_{0}^{(j)}(x_{0})}{j!} (u - x_{0})^{j} - f_{0}(x_{0}) \frac{\varphi_{0}^{(k)}(x_{0})}{k!} (u - x_{0})^{k} = o_{p}(n^{-k/(2k+1)}),$$

or equivalently

$$\left| \hat{e}_n(x_0 + tn^{-1/(2k+1)}) - f_0(x_0) \left(\widehat{\varphi}_n(x_0 + tn^{-1/(2k+1)}) - \varphi_0(x_0) - n^{-1/(2k+1)} t \varphi_0'(x_0) \right) \right| = o_p(n^{-k/(2k+1)})$$

uniformly in $|t| \leq M$.

-		-
L		

THEOREM 4.6. Let K > 0.

(i) If $\{Y_k(t), t \in \mathbb{R}\}$ is the canonical process defined in (2.2), then the localized process $\gamma_1 \mathbb{Y}_n^{locmod}(\gamma_2 \cdot)$ converges weakly in C[-K, K] to Y_k where

(4.30)
$$\gamma_1 = \left(\frac{f_0(x_0)^{k-1}|\varphi_0^{(k)}(x_0)|^3}{[(k+2)!]^3}\right)^{1/(2k+1)}$$

(4.31)
$$\gamma_2 = \left(\frac{f_0(x_0)|\varphi_0^{(k)}(x_0)|^2}{[(k+2)!]^2}\right)^{1/(2k+1)}$$

Equivalently, \mathbb{Y}_n^{locmod} converges weakly in C[-K,K] to the "driving process" $Y_{a,k,\sigma}$ where

(4.32)
$$Y_{k,a,\sigma}(t) := a \int_0^t W(s) ds - \sigma t^{k+2}$$

- and where $a = 1/\sqrt{f_0(x_0)}$, $\sigma = |\varphi_0^{(k)}(x_0)|/(k+2)!$. (ii) The localized processes satisfy $\mathbb{Y}_n^{locmod}(t) \widehat{H}_n^{locmod}(t) \ge 0$ for all $t \in \mathbb{R}$, with equality for all t such that $x_n(t) = x_0 + tn^{-1/(2k+1)} \in \widehat{S}_n(\widehat{\varphi}_n)$.
- (iii) Both \ddot{A}_n and \ddot{B}_n defined above in (4.9) and (4.10) are tight.
- (iv) The vector of processes

$$\left(\widehat{H}_{n}^{locmod}, (\widehat{H}_{n}^{locmod})^{(1)}, (\widehat{H}_{n}^{locmod})^{(2)}, \mathbb{Y}_{n}^{locmod}, (\widehat{H}_{n}^{locmod})^{(3)}, (\mathbb{Y}_{n}^{locmod})^{(1)}\right)$$

converges weakly in $(C[-K,K])^4 \times (D[-K,K])^2$ endowed with the product topology induced by the uniform topology on the spaces C[-K, K]and the Skorohod topology on the spaces D[-K, K] to the process

$$\left(H_{k,a,\sigma}, H_{k,a,\sigma}^{(1)}, H_{k,a,\sigma}^{(2)}, Y_{k,a,\sigma}, H_{k,a,\sigma}^{(3)}, Y_{k,a,\sigma}^{(1)}\right)$$

where $H_{k,a,\sigma}$ is the unique process on \mathbb{R} satisfying

(4.33)
$$\begin{cases} H_{k,a,\sigma}(t) \leq Y_{k,a,\sigma}(t) \text{ for all } t \in \mathbb{R}, \\ \int (H_{k,a,\sigma}(t) - Y_{k,a,\sigma}(t)) dH_{k,a,\sigma}^{(3)}(t) = 0, \\ H_{k,a,\sigma}^{(2)} \text{ is concave.} \end{cases}$$

Proof. (i) The first step will be to modify the local processes, i.e. going from the "density" to the "log-density" level, in order to be able to exploit concavity of φ_0 and $\hat{\varphi}_n$ and to connect the local process to the limiting

distribution obtained by Groeneboom, Jongbloed and Wellner (2001b) for estimating a convex density.

First, by Lemma 4.2, (4.27) and (A3), we can write

$$\begin{split} f_0(x_0)^{-1} & \left(\widehat{f}_n(u) - \sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j \right) \\ &= f_0(x_0)^{-1} \left(\widehat{f}_n(u) - f_0(x_0) - f_0(x_0) \sum_{j=1}^{k-1} \frac{[\varphi_0'(x_0)]^j}{j!} (u - x_0)^j \right) \\ &= \widehat{\Psi}_{k,n,1}(u) + \sum_{j=1}^k \frac{1}{j!} \left[\widehat{\varphi}_n(u) - \varphi_0(x_0) \right]^j - \sum_{j=1}^{k-1} \frac{[\varphi_0'(x_0)]^j}{j!} (u - x_0)^j \\ &= \widehat{\Psi}_{k,n,1}(u) + \left(\widehat{\varphi}_n(u) - \varphi_0(x_0) - \varphi_0'(x_0) (u - x_0) \right) + \\ &+ \sum_{j=2}^k \frac{1}{j!} \left[\widehat{\varphi}_n(u) - \varphi_0(x_0) \right]^j - \sum_{j=2}^{k-1} \frac{[\varphi_0'(x_0)]^j}{j!} (u - x_0)^j \\ &=: \left(\widehat{\varphi}_n(u) - \varphi_0(x_0) - \varphi_0'(x_0) (u - x_0) \right) + \widehat{\Psi}_{k,n,2}(u) \end{split}$$

introducing the new remainder term

(4.34)
$$\widehat{\Psi}_{k,n,2}(u) = \widehat{\Psi}_{k,n,1}(u) + \sum_{j=2}^{k} \frac{1}{j!} \left[\widehat{\varphi}_{n}(u) - \varphi_{0}(x_{0}) \right]^{j} - \sum_{j=2}^{k-1} \frac{[\varphi_{0}'(x_{0})]^{j}}{j!} (u - x_{0})^{j}.$$

Using (4.28) and (4.29) yields

$$\begin{split} \int_{\mathcal{I}} \int_{x_0}^{v} \widehat{\Psi}_{k,n,2}(u) du dv \\ &= t^2 n^{-2/(2k+1)} \sup_{u \in [x_0,v], v \in \mathcal{I}} |\widehat{\Psi}_{k,n,1}(u)| + \sum_{j=2}^{k} \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^{v} [\widehat{\varphi}_n(u) - \varphi_0(x_0)]^j du dv \\ &- \sum_{j=2}^{k-1} \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^{v} [\varphi_0'(x_0)]^j (u - x_0)^j du dv \end{split}$$

$$\begin{split} &= o_p(r_n^{-1}) \\ &+ \sum_{j=2}^k \frac{1}{j!} \sum_{l=1}^j {j \choose l} \int_{\mathcal{I}} \int_{x_0}^v [\widehat{\varphi}_n(u) - \varphi_0(x_0) \\ &- (u - x_0) \varphi'_0(x_0)]^l (u - x_0)^{j-l} [\varphi'_0(x_0)]^{j-l} du dv \\ &+ \sum_{j=2}^k \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^v [\varphi'_0(x_0)]^j (u - x_0)^j du dv \\ &- \sum_{j=2}^{k-1} \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^v [\varphi'_0(x_0)]^j (u - x_0)^j du dv \\ &= o_p(r_n^{-1}) \\ &+ \sum_{j=2}^k \frac{1}{j!} \sum_{l=1}^j {j \choose l} \int_{\mathcal{I}} \int_{x_0}^v [\widehat{\varphi}_n(u) - \varphi_0(x_0) \\ &- (u - x_0) \varphi'_0(x_0)]^l (u - x_0)^{j-l} [\varphi'_0(x_0)]^{j-l} du dv \\ &+ \frac{1}{k!} \int_{\mathcal{I}} \int_{x_0}^v (u - x_0)^k [\varphi'_0(x_0)]^k du dv. \end{split}$$

But by Lemma 4.5 one can easily show that for $j = 2, \ldots, k$ and $l = 1, \ldots, j$

$$r_n \int_{\mathcal{I}} \int_{x_0}^{v} [\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi_0'(x_0)]^l (u - x_0)^{j-l} [\varphi_0'(x_0)]^{j-l} du dv$$

= $O_p(n^{-[k(l-1)+(j-l)]/(2k+1)}) = o_p(1)$

uniformly in $|t| \leq M$. Similarly,

$$r_n \int_{\mathcal{I}} \int_{x_0}^{v} (u - x_0)^k [\varphi_0'(x_0)]^k du dv = \frac{[\varphi_0'(x_0)]^k}{(k+1)(k+2)} t^{k+2}.$$

Hence, it follows that

$$r_n \int_{\mathcal{I}} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) du dv = \frac{[\varphi_0'(x_0)]^k}{(k+2)!} t^{k+2} + o_p(1)$$

as $n \to \infty$ and uniformly in $|t| \le M$.

We turn now to the modified local processes, \mathbb{Y}_n^{locmod} and \widehat{H}_n^{locmod} defined in (4.11) and (4.12). It is not difficult to show that

(4.35)
$$\mathbb{Y}_n^{locmod}(t) = \frac{\mathbb{Y}_n^{loc}(t)}{f_0(x_0)} - r_n \int_{\mathcal{I}} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) du dv$$

and

(4.36)
$$\widehat{H}_n^{locmod}(t) = \frac{\widehat{H}_n^{loc}(t)}{f_0(x_0)} - r_n \int_{\mathcal{I}} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) du dv.$$

Note that the process \widehat{H}_n^{locmod} is in fact similar to \widehat{H}_n^{loc} , except that it is defined in terms of the log–density φ_0 instead of the density f_0 . This can be more easily seen from its original expression given in (4.12). The second expression of \widehat{H}_n^{locmod} given above is only useful for showing that it stays below \mathbb{Y}_n^{locmod} while touching it at points t such that $x_n(t) = x_0 + tn^{-1/(2k+1)} \in \widehat{S}_n(\widehat{\varphi}_n)$. The biggest advantage of considering this modified version is to be able to use concavity of φ_0 the same way Groeneboom, Jongbloed and Wellner (2001b) used convexity of the true estimated density g_0 . Their process \widetilde{H}_n^{loc} resembles \widehat{H}_n^{locmod} to a large extent (see p. 1688), and by combining arguments similar to theirs with Lemma 4.2 and the results obtained above, it follows that

$$\begin{aligned} & \exists_{n}^{locmod}(t) \\ \Rightarrow \quad [f_{0}(x_{0})]^{-1/2} \quad \int_{0}^{t} W(s)ds + \frac{f_{0}^{(k)}(x_{0})}{(k+2)!f_{0}(x_{0})}t^{k+2} - \frac{[\varphi_{0}'(x_{0})]^{k}}{(k+2)!} \ t^{k+2} \\ & = \quad [f_{0}(x_{0})]^{-1/2} \int_{0}^{t} W(s)ds + \frac{\varphi_{0}^{(k)}(x_{0})}{(k+2)!} \ t^{k+2} \\ & = \quad Y_{k,a,\sigma}(t) \qquad \text{in} \quad C[-K,K] \end{aligned}$$

where $a := [f_0(x_0)]^{-1/2}$, $\sigma := |\varphi_0^{(k)}(x_0)|/(k+2)!$, as in (4.32). Now let γ_1 and γ_2 be chosen so that

$$\gamma_1 Y_{k,a,\sigma}(\gamma_2 t) \stackrel{d}{=} Y_k(t)$$

as processes where Y_k is the integrated Gaussian process defined in (2.2). Using the scaling property of Brownian motion; that is, $\alpha^{-1/2}W(\alpha t) =_d W(t)$ for any $\alpha > 0$, we get

$$\gamma_1 \gamma_2^{3/2} = a^{-1}$$
 and $\gamma_1 \gamma_2^{k+2} = \sigma^{-1}$.

This yields γ_1 and γ_2 as given in (4.30) and (4.31), and hence

$$\begin{pmatrix} n^{k/(2k+1)} \left(\widehat{\varphi}_n(x_0) - \varphi_0(x_0) \right) \\ n^{(k-1)/(2k+1)} \left(\widehat{\varphi}'_n(x_0) - \varphi'_0(x_0) \right) \\ \rightarrow_d \quad f_0(x_0)^{-1} \begin{pmatrix} c_k(x_0,\varphi_0) \ H_k^{(2)}(0) \\ d_k(x_0,\varphi_0) \ H_k^{(3)}(0) \end{pmatrix}$$

We get the explicit expression of the asymptotic constants $c_k(x_0, \varphi_0)$ and $d_k(x_0, \varphi_0)$ using the following relations:

(4.37)
$$f_0(x_0)^{-1}c_k(x_0,\varphi_0) = (\gamma_1\gamma_2^2)^{-1}$$
 and

(4.38)
$$f_0(x_0)^{-1}d_k(x_0,\varphi_0) = (\gamma_1\gamma_2^3)^{-1}.$$

This is completely analogous to the derivations on p. 1689 in Groeneboom, Jongbloed and Wellner (2001b), precisely

(4.39)
$$(\gamma_1 \widehat{H}_n^{locmod}(\gamma_2 t))^{(2)}(0) = \gamma_1 \gamma_2^2 (\widehat{H}_n^{locmod})^{(2)}(0)$$
$$= n^{k/(2k+1)} f_0(x_0) c_k(x_0, \varphi_0)^{-1} \left(\widehat{\varphi}_n(x_0) - \varphi_0(x_0)\right)$$

and

$$(\gamma_1 \widehat{H}_n^{locmod}(\gamma_2 t))^{(3)}(0) = \gamma_1 \gamma_2^3 (\widehat{H}_n^{locmod})^{(3)}(0)$$

(4.40)
$$= n^{(k-1)/(2k+1)} f_0(x_0) d_k(x_0, \varphi_0)^{-1} \bigg(\widehat{\varphi}_n'(x_0) - \varphi_0'(x_0) \bigg).$$

From (4.37) and (4.38) we get $c_k(x_0, \varphi_0)$ and $d_k(x_0, \varphi_0)$ as given in (2.3) and (2.4), and $C_k(x_0, \varphi_0)$ and $D_k(x_0, \varphi_0)$ as in (2.5) and (2.6).

(ii) Note that we can write

$$\mathbb{Y}_n^{loc}(t) - \widehat{H}_n^{loc}(t) = r_n \left(\mathbb{H}_n(x_n(t)) - \widehat{H}_n(x_n(t)) \right) \ge 0$$

by making use of (1.1) and the specific choice of \hat{A}_n and \hat{B}_n . But since we connect \hat{H}_n^{locmod} and \mathbb{Y}_n^{locmod} to the "invelope" the latter property needs primarily to hold for the modified processes. This can easily be established by considering (4.35) and (4.36), and hence it follows that

$$\mathbb{Y}_n^{locmod}(t) - \widehat{H}_n^{locmod}(t) \ge 0$$

for all $t \in \mathbb{R}$, with equality if $x_n(t) = x_0 + tn^{-1/(2k+1)} \in \widehat{\mathcal{S}}_n(\widehat{\varphi}_n)$.

(iii) To show that \hat{A}_n and \hat{B}_n are tight. By Theorem 4.3, we know that there exists M > 0 and $\tau \in \widehat{\mathcal{S}}(\widehat{\varphi}_n)$ such that $0 \le x_0 - \tau \le M n^{-1/(2k+1)}$ with

large probability. Now using (4.19) we can write

$$\begin{aligned} |\hat{A}_{n}| &\leq r_{n}s_{n} \left| \left(\widehat{F}_{n}(x_{0}) - \widehat{F}_{n}(\tau) \right) - \left(\mathbb{F}_{n}(x_{0}) - \mathbb{F}_{n}(\tau) \right) \right| + r_{n}/n \\ &\leq r_{n}s_{n} \left| \int_{\tau}^{x_{0}} \left(\widehat{f}_{n}(u) - \sum_{j=0}^{k-1} \frac{f_{0}^{(j)}(x_{0})}{j!} (u - x_{0})^{j} \right) du \right| \\ &+ r_{n}s_{n} \left| \int_{\tau}^{x_{0}} \left(\sum_{j=0}^{k-1} \frac{f_{0}^{(j)}(x_{0})}{j!} (u - x_{0})^{j} - f_{0}(u) du \right| \right. \\ &+ r_{n}s_{n} \left| \int_{\tau}^{x_{0}} d(\mathbb{F}_{n} - F_{0}) \right| + n^{-k/(2k+1)} \\ &:= \hat{A}_{n1} + \hat{A}_{n2} + \hat{A}_{n3} + n^{-k/(2k+1)}. \end{aligned}$$

Now,

$$\begin{aligned} |\hat{A}_{n1}| \\ &\leq r_n s_n \left| \int_{\tau}^{x_0} \hat{e}_n(u) du - f_0(x_0) \left(\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0) \varphi'_0(x_0) \right) du \right| \\ &+ r_n s_n f_0(x_0) \left| \int_{\tau}^{x_0} \frac{[\varphi'_0(x_0)]^k}{k!} (u - x_0)^k \right) du \right| + \\ &+ r_n s_n f_0(x_0) \left| \int_{\tau}^{x_0} \left(\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0) \varphi'_0(x_0) \right) du \right| \\ &\leq o_p(1) + O_p(r_n s_n(\tau - x_0)^{k+1}) + O_p(r_n s_n(\tau - x_0) n^{-k/(2k+1)}) \\ &= O_p(1), \end{aligned}$$

where we used (4.26) and (4.25) to bound the first and last terms. To bound \hat{A}_{n2} we use Taylor approximation of $f_0(u)$ around x_0 to get

$$\hat{A}_{n2} \leq r_n \left| \int_{\tau}^{x_0} \frac{f_0^{(k)}(x_0)}{k!} (u - x_0)^k du \right| + r_n \left| \int_{\tau}^{x_0} (u - x_0)^k \epsilon_n(u) du \right|$$

= $O_p(1),$

where ϵ_n is a function such that $\|\epsilon_n\| \to_p 0$ as $x_0 - \tau \to_p 0$. To bound \hat{A}_{n3} , similar derivations as the ones used for bounding R_{2n} (see the proof of Lemma 4.4) can be employed where the perturbation function Δ_1 needs to be replaced by $\Delta_2(x) = 1_{[\tau, x_0]}(x)$. At "one higher integration level", similar computations can be used to

At "one higher integration level", similar computations can be used to show tightness of \hat{B}_n .

(iv) The proof of this last part of the theorem is basically identical to that of Theorem 6.2 for the LSE in Groeneboom, Jongbloed and Wellner

(2001b), and arguments similar to those of Groeneboom, Jongbloed and Wellner (2001a) or, alternatively, tightness plus uniqueness arguments along the lines of Groeneboom, Maathuis, and Wellner (2007). \Box

Proof of Theorem 2.1. The claimed joint convergence involving $\widehat{\varphi}_n$ and $\widehat{\varphi}'_n$ follows from part (iv) of Theorem 4.6 and the relations (4.39) and (4.40). The joint limiting distribution of $\widehat{f}_n(x_0) - f_0(x_0)$ and $\widehat{f}'_n(x_0) - f'_0(x_0)$ follows immediately by applying the delta-method.

4.3. Proofs for Section 3.

Proof of Theorem 3.6. We first use the simple fact that \widehat{M}_n is the only point $x \in \mathbb{R}$ which satisfies

(4.41)
$$\widehat{\varphi}'_n(t) \begin{cases} >0, & \text{if } t < x \\ \le 0, & \text{if } t \ge x \end{cases}$$

This follows immediately from concavity of $\widehat{\varphi}_n$ and the definition of \widehat{M}_n . Note that $\widehat{\varphi}_n$ may have a flat region or "modal interval"; in this case, there exists an entire interval of points where the maximum is attained, and \widehat{M}_n is the left endpoint of this interval.

A tightness property of the process $H_2^{(3)}$, which follows from Lemma 2.7 of Groeneboom, Jongbloed and Wellner (2001b), is also needed to establish the limiting distribution of \widehat{M}_n : for any $\epsilon > 0$ and $t \in \mathbb{R}$, there exists $C = C(\epsilon)$ such that

$$P\left(\left|H_2^{(3)}(t) + 24t\right| > C\right) \le \epsilon.$$

In other words, one can view $H_2^{(3)}(t)$ as an "estimator" of the odd function -24t. Since C is independent of t, it follows that for a fixed ϵ , $H_2^{(3)}(t) < 0$ (resp. $H_2^{(3)}(t) > 0$) for t > 0 (resp. -t < 0) big enough, with probability greater than $1 - \epsilon$.

The sign of $H_2^{(3)}$ and uniqueness of \widehat{M}_n turn out to be crucial in determining the limiting distribution of the latter. From Theorem 4.6 and the two derivative relations (4.39) and (4.40) it follows that

$$\begin{pmatrix} n^{k/(2k+1)} \left(\hat{\varphi}_n(x_0 + tn^{-1/(2k+1)}) - \varphi_0(x_0) - tn^{-1/(2k+1)} \varphi'_0(x_0) \right) \\ n^{(k-1)/(2k+1)} \left(\hat{\varphi}'_n(x_0 + tn^{-1/(2k+1)}) - \varphi'_0(x_0) \right) \end{pmatrix}$$

$$(4.42) \qquad \Rightarrow \begin{pmatrix} H_{k,a,\sigma}^{(2)}(t) \\ H_{k,a,\sigma}^{(3)}(t) \end{pmatrix} \quad \text{in } C[-K,K] \times D[-K,K]$$

for each K > 0 with the product topology induced by the uniform topology on C[-K, K] and the Skorohod topology on D[-K, K]. Here $H_{k,a,\sigma}$ is the

unique process on \mathbb{R} satisfying (4.33). A similar result holds for the MLE of the log-concave density f_0 . When x_0 is replaced by the population mode $m_0 = M(f_0)$ and k = 2 the second weak convergence implies that

$$n^{1/5}\left(\hat{\varphi}'_n(m_0 - Tn^{-1/5}) - \varphi'_0(m_0)\right) \to_d H^{(3)}_{2,a,\sigma}(-T),$$

and

$$n^{1/5}\left(\hat{\varphi}'_n(m_0+Tn^{-1/5})-\varphi'_0(m_0)\right) \to_d H^{(3)}_{2,a,\sigma}(T).$$

For T > 0 large enough, this in turn implies that for $\epsilon > 0$, we can find $N \in \mathbb{N} \setminus \{0\}$ such that for all n > N we have that

$$P\left(\hat{\varphi}'_n(m_0 - Tn^{-1/5}) > 0 \text{ and } \hat{\varphi}'_n(m_0 + Tn^{-1/5}) < 0\right) > 1 - \epsilon$$

using the property of \widehat{M}_n in (4.41), it follows that

$$P\left(\widehat{M}_n \in \left[m_0 - Tn^{-1/5}, \ m_0 + Tn^{-1/5}\right]\right) > 1 - \epsilon$$

for all n > N.

We first conclude that $\widehat{M}_n - m_0 = O_p(n^{-1/5})$. Then we note that

$$n^{1/5}(\widehat{M}_n - m_0) = M(\mathbb{Z}_n)$$

where

$$\mathbb{Z}_{n}(t) = n^{2/5} (\hat{\varphi}_{n}(m_{0} + tn^{-1/5}) - \varphi_{0}(m_{0}))$$

$$\Rightarrow \mathbb{Z}(t) := H_{2,a,\sigma}^{(2)}(t) \quad \text{in } C([-K,K])$$

for each K > 0 by (4.42) with k = 2. Thus by the argmax continuous mapping theorem (see e.g. van der Vaart and Wellner (1996), page 286) it follows that

$$M(\mathbb{Z}_n) \to_d M(\mathbb{Z}) = M(H_{2,a,\sigma}^{(2)})$$

where $\mathbb{Z} = H_{2,a,\sigma}^{(2)}$, $a = 1/\sqrt{f_0(m_0)}$, and $\sigma = |\varphi_0^{(2)}(m_0)|/4!$. Note that $H_{2,a,\sigma}$ is related to the "driving process" $Y_{2,a,\sigma}$ with a = 1

Note that $H_{2,a,\sigma}$ is related to the "driving process" $Y_{2,a,\sigma}$ with $a = 1/\sqrt{f_0(m_0)}$, $\sigma = |\varphi_0^{(2)}(m_0)|/4!$ as in (4.32) with k = 2. Now $\gamma_1 Y_{2,a,\sigma}(\gamma_2 t) \stackrel{d}{=} Y_2(t)$ as processes where $Y_2 := Y_{2,1,1}$. Thus it also holds that

$$\gamma_1 H_{2,a,\sigma}(\gamma_2 t) \stackrel{d}{=} H_2(t) \text{ and } \gamma_1 \gamma_2^2 H_{2,a,\sigma}^{(2)}(\gamma_2 t) \stackrel{d}{=} H_2^{(2)}(t),$$

or, equivalently, $H_{2,a,\sigma}^{(2)}(v) \stackrel{d}{=} H_2^{(2)}(v/\gamma_2)/(\gamma_1\gamma_2^2)$. Since $M(dg(c \cdot)) = c^{-1}M(g)$ for c, d > 0, it follows that

$$M(H_{2,a,\sigma}^{(2)}) \stackrel{d}{=} M\left(\frac{1}{\gamma_1\gamma_2^2}H_2^{(2)}(\cdot/\gamma_2)\right) \stackrel{d}{=} \gamma_2 M(H_2^{(2)})$$

where

$$\gamma_2 = \left(f_0(m_0) \frac{|\varphi_0^{(2)}(m_0)|^2}{(4!)^2}\right)^{-1/5} = \left(\frac{(4!)^2 f_0(m_0)}{f_0^{''}(m_0)^2}\right)^{1/5}$$

by direct computation using $f'_0(m_0) = 0 = \varphi'_0(m_0)$ and Lemma 4.2.

Acknowledgments. This research was initiated while Kaspar Rufibach was visiting the Institute for Mathematical Stochastics at the University of Göttingen, Germany and while Fadoua Balabdaoui was visiting the Institute of Mathematical Statistics and Actuarial Science at the University of Bern, Switzerland. We would like to thank both institutions for their hospitality.

REFERENCES

- AN, M. Y. (1998). Logconcavity versus logconvexity: a complete characterization. J. Economic Theory 80 350 – 369.
- ANEVSKI, D. and HÖSSJER, O. (2002). Monotone regression and density function estimation at a point of discontinuity. J. Nonparametr. Stat. 14 279–294.
- BAGNOLI, M. and BERGSTROM, T. (2005). Log-concave probability and its applications. *Econom. Theory* 26 445–469.
- BALABDAOUI, F. and WELLNER, J. A. (2007). Estimation of a k-monotone density: limiting distribution theory and the spline connection. Ann. Statist. **35** to appear.
- BARLOW, R. E. and PROSCHAN, F. (1975). Statistical theory of reliability and life testing. Holt, Rinehart and Winston, Inc., New York. Probability models, International Series in Decision Processes, Series in Quantitative Methods for Decision Making.
- BIRGÉ, L. (1989). The Grenander estimator: a nonasymptotic approach. Ann. Statist. 17 1532–1549.
- BIRGÉ, L. (1997). Estimation of unimodal densities without smoothness assumptions. Ann. Statist. 25 970–981.
- BRUNNER, L. J. and LO, A. Y. (1989). Bayes methods for a symmetric unimodal density and its mode. Ann. Statist. 17 1550–1566.
- CAI, T. and LOW, M. (2007). Estimating convex functions. Tech. rep., University of Pennsylvania. Talk at JSM, Salt Lake City, July 31, 2007; announced with Mark Low as speaker; presented by T. Cai.
- CAI, T. T. and Low, M. G. (2005). On adaptive estimation of linear functionals. Ann. Statist. **33** 2311–2343.
- CHANG, G. and WALTHER, G. (2006). Clustering with mixtures of log-concave distributions. Tech. rep., Stanford University. Available at http://www-stat.stanford.edu/ walther/flexclust.pdf.
- CHERNOFF, H. (1964). Estimation of the mode. Ann. Inst. Statist. Math. 16 31-41.
- CULE, M., GRAMACY, R. and SAMWORTH, R. (2007). LogConcDEAD: Maximum likelihood estimation of a log-concave density. R package version 1.0-5.
- DALENIUS, T. (1965). The mode—a neglected statistical parameter. J. Roy. Statist. Soc. Ser. A 128 110–117.
- DHARMADHIKARI, S. and JOAG-DEV, K. (1988). Unimodality, convexity, and applications. Probability and Mathematical Statistics, Academic Press Inc., Boston, MA.

- DÜMBGEN, L., HÜSLER, A. and RUFIBACH, K. (2007). Active set and EM algorithms for log-concave densities based on complete and censored data. Tech. rep., University of Bern. Available at arXiv:0707.4643.
- DÜMBGEN, L. and RUFIBACH, K. (2007). Maximum likelihood estimation of a log-concave density and its density and its distribution function. Tech. rep., University of Bern. Available at arXiv:0709.0334.
- EDDY, W. F. (1980). Optimum kernel estimators of the mode. Ann. Statist. 8 870–882.
- EDDY, W. F. (1982). The asymptotic distributions of kernel estimators of the mode. Z. Wahrsch. Verw. Gebiete 59 279–290.
- EKBLOM, H. (1972). A Monte Carlo investigation of mode estimators in small samples. Applied Statistics **21** 177–184.
- GRENANDER, U. (1956). On the theory of mortality measurement. II. Skand. Aktuarietidskr. **39** 125–153 (1957).
- GRENANDER, U. (1965). Some direct estimates of the mode. Ann. Math. Statist **36** 131–138.
- GROENEBOOM, P. (1996). Lectures on inverse problems. In Lectures on probability theory and statistics (Saint-Flour, 1994), vol. 1648 of Lecture Notes in Math. Springer, Berlin, 67–164.
- GROENEBOOM, P., JONGBLOED, G. and WELLNER, J. A. (2001a). A canonical process for estimation of convex functions: the "invelope" of integrated Brownian motion $+t^4$. Ann. Statist. **29** 1620–1652.
- GROENEBOOM, P., JONGBLOED, G. and WELLNER, J. A. (2001b). Estimation of a convex function: characterizations and asymptotic theory. *Ann. Statist.* **29** 1653–1698.
- GRUND, B. and HALL, P. (1995). On the minimisation of L^p error in mode estimation. Ann. Statist. **23** 2264–2284.
- HALL, P. (1982). Asymptotic theory of Grenander's mode estimator. Z. Wahrsch. Verw. Gebiete 60 315–334.
- HAS'MINSKIĬ, R. Z. (1979). Lower bound for the risks of nonparametric estimates of the mode. In *Contributions to statistics*. Reidel, Dordrecht, 91–97.
- HERRMANN, E. and ZIEGLER, K. (2004). Rates on consistency for nonparametric estimation of the mode in absence of smoothness assumptions. *Statist. Probab. Lett.* 68 359–368.
- Ho, M.-W. (2006a). Bayes estimation of a symmetric unimodal density via S-paths. J. Comput. Graph. Statist. 15 848–860.
- Ho, M.-W. (2006b). Bayes estimation of a unimodal density via S-paths. Technical report, National University of Singapore. Available at http://arxiv.org/math/0609056.
- IBRAGIMOV, I. A. (1956). On the composition of unimodal distributions. Teor. Veroyatnost. i Primenen. 1 283–288.
- JANKOWSKI, H. and WELLNER, J. A. (2007). Nonparametric estimation of a convex bathtub-shaped hazard function. Technical report, University of Washington.
- JONGBLOED, G. (1995). *Three Statistical Inverse Problems*. Ph.D. thesis, Delft University of Technology.
- JONGBLOED, G. (2000). Minimax lower bounds and moduli of continuity. Statist. Probab. Lett. 50 279–284.
- KIM, J. and POLLARD, D. (1990). Cube root asymptotics. Ann. Statist. 18 191–219.
- KLEMELÄ, J. (2005). Adaptive estimation of the mode of a multivariate density. J. Nonparametr. Stat. 17 83–105.
- LEPSKIĬ, O. V. (1992). On problems of adaptive estimation in white Gaussian noise. In *Topics in nonparametric estimation*, vol. 12 of *Adv. Soviet Math.* Amer. Math. Soc., Providence, RI, 87–106.

- LEURGANS, S. (1982). Asymptotic distributions of slope-of-greatest-convex-minorant estimators. Ann. Statist. 10 287–296.
- Low, M. G. and KANG, Y.-G. (2002). Estimating monotone functions. Statist. Probab. Lett. 56 361–367.
- MARSHALL, A. W. and OLKIN, I. (1979). Inequalities: theory of majorization and its applications, vol. 143 of Mathematics in Science and Engineering. Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York.
- MARSHALL, A. W. and OLKIN, I. (2007). *Life distributions*. Springer Series in Statistics, Springer, New York. Structure of nonparametric, semiparametric, and parametric families.
- MEYER, M. C. (2001). An alternative unimodal density estimator with a consistent estimate of the mode. *Statist. Sinica* **11** 1159–1174.
- MEYER, M. C. and WOODROOFE, M. (2004). Consistent maximum likelihood estimation of a unimodal density using shape restrictions. *Canad. J. Statist.* **32** 85–100.
- Müller, H.-G. (1989). Adaptive nonparametric peak estimation. Ann. Statist. 17 1053–1069.
- Müller, S. and RUFIBACH, K. (2006). Smoothed tail index estimation. Tech. rep., Stanford University. Available at arXiv:math/0612140.
- PAL, J. K., WOODROOFE, M. B. and MEYER, M. C. (2007). Estimating a polya frequency function. In Complex Datasets and Inverse Problems: Tomography, Networks, and Beyond, vol. 54 of IMS Lecture Notes-Monograph Series. IMS, 239–249.
- PARZEN, E. (1962). On estimation of a probability density function and mode. Ann. Math. Statist. 33 1065–1076.
- PRAKASA RAO, B. L. S. (1969). Estimation of a unimodal density. Sankhyā Ser. A **31** 23–36.
- ROMANO, J. P. (1988). On weak convergence and optimality of kernel density estimates of the mode. Ann. Statist. 16 629–647.
- RUFIBACH, K. (2006). Log-concave density estimation and bump hunting for I.I.D. observations. Ph.D. thesis, Universities of Bern and Göttingen.
- RUFIBACH, K. (2007). Computing maximum likelihood estimators of a log-concave density function. J. Statist. Comp. Sim. 77 561–574.
- RUFIBACH, K. and DÜMBGEN, L. (2007). logcondens: Estimate a Log-Concave Probability Density from iid Observations. R package version 1.3.1.
- URL http://www.stanford.edu/ kasparr , http://www.stat.unibe.ch/ duembgen VAN DER VAART, A. W. and WELLNER, J. A. (1996). Weak Convergence and Empirical Processes with Applications in Statistics. Springer, New York.
- VENTER, J. H. (1967). On estimation of the mode. Ann. Math. Statist 38 1446–1455.
- VIEU, P. (1996). A note on density mode estimation. Statist. Probab. Lett. 26 297-307.
- WALTHER, G. (2002). Detecting the presence of mixing with multiscale maximum likelihood. J. Amer. Statist. Assoc. 97 508–513.
- WEGMAN, E. J. (1970a). Maximum likelihood estimation of a unimodal density function. Ann. Math. Statist. 41 457–471.
- WEGMAN, E. J. (1970b). Maximum likelihood estimation of a unimodal density. II. Ann. Math. Statist. 41 2169–2174.
- WEGMAN, E. J. (1971). A note on the estimation of the mode. Ann. Math. Statist. 42 1909–1915.
- WRIGHT, F. T. (1981). The asymptotic behavior of monotone regression estimates. Ann. Statist. 9 443–448.
- ZIEGLER, K. (2004). Adaptive kernel estimation of the mode in a nonparametric random design regression model. Probab. Math. Statist. 24 213–235.

38 FADOUA BALABDAOUI, KASPAR RUFIBACH, AND J. A. WELLNER

CENTRE DE RECHERCHE EN MATHEMATIQUES DE LA DECISIONINSTITUTE OF SOCIAL ANDUNIVERSITE PARIS-DAUPHINEPREVENTIVE MEDICINE, BIPARIS, FRANCEUNIVERSITY OF ZURICHE-MAIL: fadoua@ceremade.dauphine.frCH-8001 ZURICH, SWITZER

INSTITUTE OF SOCIAL AND PREVENTIVE MEDICINE, BIOSTATISTICS UNIT UNIVERSITY OF ZURICH CH-8001 ZURICH, SWITZERLAND E-MAIL: kaspar.rufibach@ifspm.uzh.ch

Department of Statistics, Box 354322 University of Washington Seattle, WA 98195-4322 E-mail: jaw@stat.washington.edu