

Role of environmental factors for the vertical distribution (0–1000 m) of marine bacterial communities in the NW Mediterranean Sea

J. F. Ghiglione, C. Palacios, J. C. Marty, G. Mével, C. Labrune, P. Conan, Mireille Pujo-Pay, N. Garcia, M. Goutx

▶ To cite this version:

J. F. Ghiglione, C. Palacios, J. C. Marty, G. Mével, C. Labrune, et al.. Role of environmental factors for the vertical distribution (0–1000 m) of marine bacterial communities in the NW Mediterranean Sea. Biogeosciences Discussions, 2008, 5 (3), pp.2131-2164. hal-00365586v1

HAL Id: hal-00365586 https://hal.science/hal-00365586v1

Submitted on 18 Jun 2008 (v1), last revised 29 Jul 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2131

Biogeosciences Discuss., 5, 2131–2164, 2008 www.biogeosciences-discuss.net/5/2131/2008/ © Author(s) 2008. This work is distributed under the Creative Commons Attribution 3.0 License.

Biogeosciences Discussions is the access reviewed discussion forum of Biogeosciences

Role of environmental factors for the vertical distribution (0–1000 m) of marine bacterial communities in the NW Mediterranean Sea

J. F. Ghiglione^{1,2}, C. Palacios³, J. C. Marty⁴, G. Mével^{5,6,7}, C. Labrune^{1,2}, P. Conan^{1,2}, M. Pujo-Pay^{1,2}, N. Garcia⁸, and M. Goutx⁸

¹CNRS, UMR7621, Laboratoire d'Océanographie Biologique de Banyuls, Avenue Fontaulé, BP44, 66650 Banyuls-sur-Mer, France
²UPMC Univ Paris 06, UMR 7621, Laboratoire ARAGO, Avenue Fontaulé, BP44, 66650 Banyuls-sur-Mer, France
³UMR 5244 CNRS UPVD EPHE, Biologie et Ecologie Tropicale et Méditerranéenne, Université de Perpignan Via Domitia, 52 Avenue Paul Alduy, 66860 Perpignan Cedex, France
⁴Observatoire Océanologique de Villefranche-sur-mer; Université Pierre et Marie Curie-Paris6; Institut National des Sciences de l'Univers (INSU), CNRS (UMR 7093), BP 28, 06234 Villefranche sur mer, France
⁵CNRS, UMR 7144, Equipe de Chimie Marine, Station Biologique de Roscoff, 29682 Roscoff, France



Biogeosciences

Discussions

5, 2131–2164, 2008

Environmental influence on bacterial community structure





⁶UPMC, Univ Paris 06, UMR 7144, Station Biologique de Roscoff, 29682 Roscoff, France
⁷UBO, Institut Universitaire Européen de la Mer, 29280 Plouzané, France
⁸Laboratoire de Microbiologie, Géochimie et Ecologie Marines; Université de la Méditerranée; Centre d'Océanologie de Marseille, CNRS (UMR 6117), Campus de Luminy, case 901, 13288
Marseille Cedex 09, France

Received: 26 March 2008 - Accepted: 21 April 2008 - Published: 26 May 2008

Correspondence to: J. F. Ghiglione (ghiglione@obs-banyuls.fr)

BGD

5, 2131–2164, 2008

Environmental influence on bacterial community structure





Abstract

Bacterioplankton play a central role in energy and matter fluxes at the sea, yet the factors that constrain their variation in the marine systems are still poorly understood. Here we show the explanatory power of multivariate statistical analysis of bacterial community structures coupled with fine measurements of numerous environmental parameters. We gathered and analysed data from a one month sampling period from the surface to 1000 m depth at the JGOFS-DYFAMED station (NW Mediterranean Sea). This station is characterized by very poor horizontal advection currents what makes it an ideal model to test hypothesis on the causes of vertical stratification of bacterial communities. Capillary electrophoresis single strand conformation polymorphism (CE-SSCP) fingerprinting profiles analyzed using multivariate statistical methods demonstrated a vertical zonation of bacterial assemblages in three layers, above, in or just below the chlorophyll maximum and deeper, that remained stable during the entire sampling period. Through the use of direct gradient multivariate ordination analyses

- ¹⁵ we demonstrate that a complex array of biogeochemical parameters is the driving force behinds bacterial community structure shifts in the water column. Physico-chemical parameters such as phosphate, nitrate, salinity and to a lesser extend temperature, oxygen, dissolve organic carbon and photosynthetically active radiation acted in synergy to explain bacterial assemblages changes with depth. Analysis of lipid biomarkers of the
- organic matter sources and fates suggested that bacterial community structure at the surface layers was in part explained by lipids from chloroplast origin. Further detailed analysis of pigment-based phytoplankton diversity gave evidence of a compartmentalized influence of several phytoplankton groups on bacterial community structure in the first 150 m depth. This study is probably the first example of an analysis employing
- ²⁵ a complex environmental dataset in combination with microbial community profiles to unravel the mechanisms underneath bacterial assemblages in marine systems.

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





1 Introduction

Microorganisms represent the most abundant, biogeochemically important biological component in the oceans. At concentrations ranging between 10⁴ and 10⁶ cells ml⁻¹, an estimated 20 to 50% of marine primary productivity is channeled through bacteri-oplankton (Azam et al., 1983; Cho and Azam, 1988, 1990). Bacterial species composition is an important variable controlling the rates and patterns of organic matter remineralization (Martinez et al., 1996; Pinhassi et al., 1997). Thus, knowledge of the composition of bacterial communities, and how that composition varies over space and time, is likely to be of major importance for understanding the role of bacteria in marine biogeochemistry (Riemann et al., 1999). The application of bacterial community changes with depth as demonstrated in a wide variety of oligotrophic seawaters.

including the Pacific and Atlantic Oceans (Lee and Fuhrman, 1991), the Mediterranean Sea (Acinas et al., 1997; Moeseneder et al., 2001; Ghiglione et al., 2005, 2007), the

- ¹⁵ Arabian Sea (Riemann et al., 1999) and Antarctic zones (Murray et al., 1998). The presence of the same pattern in such widely distant ocean basins suggests that vertical stratification is a very important factor determining the bacterial species inhabiting the ocean (Acinas et al., 1997). Previous studies formulated different hypotheses to explain the vertical distribution of bacterial communities pointing to by depth-related
- changes in environmental conditions or interactions with other planktonic components. Thus, the existing uncertainty underlines the need for a more robust investigation of this question. More generally, the factors that constrain microbial community composition and its variation in marine systems are still poorly understood in marine systems (Polz et al., 2006). We hypothesized that the combination of high-throughput diver-
- sity composition assessments in combination with in-depth environmental parameters measurements and appropriate multivariate statistical analyses should shed light on the actual factors responsible for bacterioplankton communities shifts with depth.

The NW Mediterranean Sea is characterized by oligotrophic conditions and strong

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





isothermic stratification of the water column during the summer. The great depth (2300 m), and weakness of horizontal advection currents (Andersen and Prieur, 2000), makes the French JGOFS-DYFAMED sampling area situated at the central zone of the Ligurian Sea (NW Mediterranean) a great model zone to follow the mineralization
 fluxes from the prokaryotic compartment along the water column. In this work, we examined the vertical distribution (0–1000 m) of bacterial community structure at this station during a one month cruise, by using capillary electrophoresis-single strand conformation polymorphism (CE-SSCP). This technique, originally developed for mutation

- detection, has been widely used for genetic profiling of microbial communities within
 a variety of ecosystems (Lee et al., 1996; Sunnucks et al., 2000; Godon et al., 2001; Ghiglione et al., 2005) and presents similar or higher detection limits than DGGE or T-RFLP fingerprinting techniques (Hong et al., 2007; Smalla et al., 2007). CE-SSCP high reproducibility permits comparisons of microbial community fingerprinting profiles from a large set of samples (Ghiglione et al., 2005). As a first objective we sought to de-
- termine the consistency of the vertical distribution of bacterial communities during the late summer–autumn transition period by using multivariate non-parametric statistical methods. Secondly, we took quantitatively unparallel measurements of environmental parameters including physico-chemical variables, lipid class biotracers and phytoplankton pigments, to investigate their influence as potential driving forces for shifts
 in bacterial community structure along the water column using multivariate statistical
- direct gradient ordination methods. In a companion paper, the vertical and temporal dynamics of bacterial abundance and activity during this cruise are reported in Mével et al. (2008).

BGD 5, 2131-2164, 2008 **Environmental** influence on bacterial community structure J. F. Ghiglione et al. **Title Page** Introduction Abstract Conclusions References Tables **Figures**

Close



Full Screen / Esc

Printer-friendly Version

Interactive Discussion

14

Back

2 Materials and methods

2.1 Study site and general sampling methods

Sampling took part during a one month cruise conducted from mid-September to mid-October 2004 at the French-JGOFS time-series station DYFAMED (43°25′2 N, 07°51′8 E; 2350 m max depth) in the NW Mediterranean Sea on the RV "Thalassa". We collected samples once a week on 19 September (Julian Day – JD 263; CTD 46) (sample A), 26 September (JD 270; CTD 101) (sample B), 6 October (JD 280; CTD 167) (sample C) and 12 October (JD 286, CTD 215) (sample D) using a rosette system equipped with 24 Niskin bottles. For each sampling day, samples were collected at depths between the surface and 1000 m (i.e. 5, 20, 40, 60, 80, 150, 200, 400, 500, 700, 1000 m).

2.2 Physico-chemical parameters

15

We used standard sensors of the CTD – SeaBird SBE 911 plus installed in the rosette system to measure pressure, temperature, conductivity, oxygen, salinity and photosynthetically active radiation (PAR).

For nutrients (nitrate, nitrite, phosphate, silicate) seawater was collected into polyethylene flasks and analyses were performed on board on a Technicon AutoAnalyzer (AAII) according to Tréguer and LeCorre (1975). Ammonia was measured on board according to Holmes et al. (1999) on a Turner Design TD-700 fluorimeter.

 For dissolved organic carbon (DOC), seawater was filtered through two precombusted (24 h, 450°C) glass fiber filters (Whatman GF/F, 25 mm) and collected into precombusted glass tubes closed with a screw cap and a Teflon liner. Samples were acidified with orthophosphoric acid (H₃PO₄) and DOC was analyzed immediately on board on a Shimadzu TOC-V analyser using the high temperature catalytic
 oxidation (HTCO) technique (Sugimura and Suzuki, 1988; Cauwet, 1994). Analytical accuracy of measurements was less than 2 μM. Deep Sargasso Sea reference

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





water $(47 \text{ mol I}^{-1} \text{ C}, \pm 0.5 \text{ SE}, \text{http://www.rsmas.miami.edu/groups/biogeochem/CRM.}$ html) was injected every 10 samples to insure stable operating conditions.

Colored dissolved organic matter (CDOM) substances were extracted from 500 ml sea water samples by adsorption on C18 microcolumns. CDOM was eluated from the microcolumn by methanol and quantified by light-absorbance at 412 nm.

2.3 Lipid biomarkers

5

25

Four liters of seawater were filtered onto pre-combusted glass fiber filters (Whatman GF/F, 47 mm). Dissolved lipids were extracted from 21 of the filtrate by liquid-liquid extraction, using dichloromethane (two extractions at sea water natural pH, followed by two extractions at pH 2). Each lipid extract was separated into classes of com-10 pounds according to polarity, using chromarods SIII (0.9 mm diameter, 150 mm length, $75\,\mu m$ silica thick) and guantified using a thin layer chromatography-flame ionization detection (TLC-FID) latroscan TH10 apparatus model MK-IV (latron, Japan; hydrogen flow, 160 ml min⁻¹; air flow, 2000 ml min⁻¹). The elution scheme previously described (Striby et al., 1999) separated twelve classes of acyl-lipids and free lipids, including 15 reserves (wax esters (WE) and triacylglycerols (TG)), lipids from chloroplasts (LC) (pigments and glycolipids), cellular membrane lipids (phospholipids (PL) which include phosphoglycerides, phosphatidylethanolamines and phosphatidylcholines), and their hydrolysis metabolites (Met) (alcohols, free fatty acids, monoglycerides and 1,2- and 1,3-diglycerides). 20

2.4 Phytoplankton pigments

Two liters and a half of seawater were filtered on glass fiber filters (Whatman GF/F, 25 mm), which were immediately frozen in liquid nitrogen until analysis. Pigments analysis employed a Agilent Technologies HPLC system with a RP-C8 (Agilent) column 150 mm long, 3 mm internal diameter, $3.5 \,\mu$ m granulometry (Zorbax Eclipse) and a Diode array detector (1100 model). All steps of extraction and analysis were per-

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





formed under dim light conditions. The use of an internal standard (Vitamin E Acetate) allowed for the correction of possible losses during extraction as well as for the presence of water in the filter. The analytical method was based on a separation gradient between a methanol: ammonium acetate (70:30 v:v) mixture and 100% methanol as in Vidussi et al. (1996) with a few improvements in sensitivity.

Identification of pigments was performed using retention-time data and by comparison of on-line collected absorption spectra with those of a library established from standards and reference cultures. Response factors for eight external pigment standards (peridinin, 19'-butanoyloxyfucoxanthin, fucoxanthin, 19'-hexanoyloxyfucoxanthin, allox-

anthin, zeaxanthin, chlorophyll-*b*, chlorophyll-*a*) were determined by spectrophotometry (Perkin Elmer) followed by HPLC analysis. These response factors were then derived to compute the specific extinction coefficients of these pigments for the HPLC system. The specific extinction coefficients from others pigments were derived from the literature (Jeffrey et al., 1997). The concentration of key chlorophyll and carotenoid
 pigments was used to determine phytoplankton community composition.

A review on taxonomic pigments can be found in Jeffrey and Vesk (1997). Divinyl chlorophyll-*a* and Divinyl chlorophyll-*b* are typical markers of picoplanktonic prochlorophytes (Goericke and Repeta, 1992; Claustre et al., 1994), and zeaxanthin is contained in both prochlorophytes and synechoccocus. 19' hexanoyloxyfucoxanthin is a marker of prymnesiophytes (Wright and Jeffrey, 1987) and 19' butanoyloxyfucoxanthin is mainly associated with pelagophytes (Andersen et al., 1993) but also with prymnesiophytes (Jeffrey et al., 1997). In the microphytoplankton range fucoxanthin is associated with diatoms (Barlow et al., 1997) but is also present in prymnesiophytes and pelagophytes. Peridinin is a specific marker of autotrophic dinoflagellates (Jeffrey and Vesk, 1997).

²⁵ The CHEMTAX matrix factorization program developed by Mackey et al. (1998) was applied to the phytoplankton data set to determine the contribution of principal taxonomic phytoplankton groups to the total chlorophyll-*a* (Marty et al., 2008¹).

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





¹Marty J. C., Garcia, N., and Raimbault, P.: Phytoplankton dynamics and primary production under late summer conditions (DYNAPROC II cruise Sep/Oct 2004, NW Mediterranean Sea),

Two liters of seawater were pre-filtered through 3μ m-pore-size filters (47 mm, Nucleopore) to remove most eukaryotic organisms and to prevent clogging of the final filter. Bacterial cells were concentrated onto 0.22μ m-pore-size filters (47 mm, PC Nucleo-⁵ pore) and stored into 2 ml Eppendorf tubes at -20° C until extraction. The procedure used for DNA extraction was modified from Furhman et al. (1988). This is a relatively gentle method that employs enzymatic and detergent-based lysis to avoid excessive shearing of DNA, producing DNA of suitable quality for PCR amplification, and reducing the risk of chimera formation during PCR. Previously frozen filters were cut with sterilized scissors into small strips and vortexed briefly in 840 μ l of alkaline lysis buffer (50 mM Tris hydrochloride pH=8.3, 40 mM EDTA, 0.75 M sucrose). Cells were lysed by an initial incubation for 45 min at 37°C after adding 50 μ l of freshly prepared lysosyme solution (20 mg ml⁻¹), and a second incubation at 50°C for 10 min after adding 100 μ l of 10% sodium dodecyl sulfate and 10 μ l of a proteinase K solution (20 mg ml⁻¹).

- ¹⁵ The lysate was extracted with an equal volume of phenol:chloroform:isoamyl alcohol (25:24:1) and an equal volume of chloroform:isoamyl alcohol (24:1). Nucleic acids were precipitated at –20°C overnight with 1 volume of isopropanol, and resuspended in TE (10 mM Tris, 1 mM EDTA). Molecular weight distribution and purity of the DNA were assessed by 1% agarose gel electrophoresis and quantified either by visual comparison
- with molecular weight markers in ethidium bromide stained agarose gels (rough estimate) or by optical density measurements using GeneQuant II (Pharmacia Biotech.). PCR amplification was performed by using primers w49 (5'-ACG GTC CAG ACT CCT ACG GG-3') and w34 (5'-TTA CCG CGG CTG CTG GCA C-3'), which amplify the variable V3 region of the 16S rDNA (Escherichia coli positions 329–533) (Brosius et al., 1221). The second state of the second state.
- ²⁵ 1981). The primer w34 was fluorescently labelled at the 5'-end with phosphoramidite (TET, Applied Biosystems). Both primers were synthesized commercially (Eurogentec). Each $50 \,\mu$ I reaction mixture contained $50 \,\mu$ M of each primer, 1X Pfu reaction

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure

J. F. Ghiglione et al.





Deep-Sea Res. I, in revision, 2008.

buffer, 20 mM dNTPs, 1.0 U of Pfu DNA polymerase (Promega) and 0.1 μg of template DNA. PCR amplification consisted of an initial denaturation step of 94°C for 2 min, followed by 25 cycles of denaturation at 94°C for 30 s, annealing at 61°C for 30 s, and extension at 72°C for 30 s, and a final elongation step at 72°C for 10 min using a Robocycler (Stratagene). Correct size (ca. 200 bp length) of PCR products were determined by 2% agarose gel electrophoresis with a DNA size standard (Low DNA Mass Ladder, GIBCO BRL).

- 2.6 Capillary Electrophoresis Single Strand Conformation Polymorphism (CE-SSCP) procedure
- Labelled PCR amplicons were purified with a QIAquick PCR purification kit (Qiagen) according to manufacturers instructions followed by quantification by visual inspection as above. Samples were diluted in sterile TE (10 mM Tris, 1 mM EDTA) in order to obtain 10 ng μl⁻¹ of PCR product. From the resulting dilution, 1 μl of PCR product was mixed with 18 μl of formamide (Applera) and 1 μl of an internal size standard
 GeneScan-400 Rox (Applied Biosystems). The mixture was then denatured for 5 min at 94°C and immediately cooled on ice for at least 5 min. The procedure used for CE-SSCP analysis was a modification of that described by Delbès et al. (1998). Samples were electrokinetically injected (5 s, 12 kV) in a capillary tube (47 cm×50 μm) filled with a mixture composed of 5.6% GeneScan polymer (Applied Biosystems) and 10%
- ²⁰ autoclaved glycerol in sterile TBE buffer (90 mM Tris-borate, 2 mM EDTA [pH 8.0]). Electrophoresis was carried out at 15 kV and 30°C for 30 min per sample and phosphoramidite (TET)-labelled amplicons were detected by a laser with a virtual filter C (detection wavelengths 532, 537, and 584 nm). Data were collected with ABI Prism 310 collection software (Applied Biosystems). In order to normalize mobilities from different
- ²⁵ runs, all electropherograms were calibrated by fixing the positions of peaks produced by the size standard and by using a second-order least square curve (i.e. linear regression) to provide the best interlane comparison (Genescan analysis software, Applied Biosystems). Electropherograms were then transferred to the software SAFUM (Zemb



et al., 2007), which renders a profile of fluorescence intensity as a function of retention time per sample taking therefore into account the presence and intensity of each individual signal. This software corrects for sample intensity differences rendering normalized profiles for all samples.

5 2.7 Ecological statistical analysis

Comparative analysis of CE-SSCP fingerprints was carried out with the PRIMER 5 software (PRIMER-E, Ltd., UK). Ordination of Bray-Curtis similarities among normalized sample profiles was performed by non-metric multidimensional scaling (MDS). We used this ordination technique to determine the relationships among sample profiles as representative of the bacterial community structure of each sample site. This method attempts to preserve the ranked order of the similarity of any two communities as an inverse function of the distance between plotted sample points (Kruskal, 1964). The degree to which the plot matches the similarity matrix can be judged by examining stress values (Kruskal's stress). MDS was performed using 30 random starting configurations of sample points to avoid local minima. Additionally, hierarchical agglomerative clustering of Bray-Curtis similarities was performed using the group average method

- of PRIMER software. To test the null hypothesis that there was no difference between bacterial communities of different depth layers we conducted an analysis of similarities with the subroutine ANOSIM of PRIMER. ANOSIM is a nonparametric technique de-
- ²⁰ signed to allow statistical comparisons for multivariate data sets in a manner similar to univariate techniques (ANOVA) (Clarke and Warwick, 2001). ANOSIM first calculates the *R* statistic that displays the degree of separation between groups. Complete separation is indicated by R=1, whereas R=0 suggests no separation. Having determined *R*, ANOSIM randomly assigns samples to different groups to generate a null distribu-
- tion of *R* (Monte Carlo test) to test whether within-group samples are more closely related to each other than would be expected by chance.

To investigate relationships between bacterioplankton community structure and physico-chemical parameters, lipids biomarkers and phytoplankton pigments, we used



a direct gradient approach, canonical correspondence analysis (CCA) using the software package CANOCO, version 4.5 for Windows (ter Braak and Šmilauer, 2002). CCA generates an ordination plot that shows the main pattern of variation in community composition as accounted for by measured environmental variables. In other

- ⁵ words, the resulting ordination axes are linear combinations of the environmental variables that best explain microbial diversity composition data (ter Braak, 1986). We first imported OTUs abundance data from spreadsheets using WCanoImp program within the CANOCO package. We then used Canoco program to perform CCA with species scaling on intersample distances so that samples and environmental variables formed
- a biplot. To statistically evaluate the significance of the first canonical axis and of all canonical axes together we used Monte Carlo permutation full model test with 199 unrestricted permutations. Finally, to represent biplots we used the program CANODRAW within CANOCO package. Additionally, Spearman's rank pairwise correlations between the environmental variables mentioned above helped to determine their significance for further ecological analysis.

3 Results

3.1 Bacterial community structure analysis

Bacterial community structure was characterized through CE-SSCP fingerprinting profiling once per week within 0–1000 m depth during September–October 2004. An example of the community fingerprints obtained at different depths for julian day 263 is shown in Fig. 1. On a mean of 49 peaks (SD=0.82, *n*=4) found at each date, 19.4% (±5.23) were found at all depth whereas 14.9% (±0.82) appeared only once or twice in the water column. For comparative analysis of community fingerprints we used multivariate MDS and hierarchical agglomerative clustering. Both analyses show a re²⁵ current in-depth pattern along the sampling period with distinct bacterial communities at three different layers: 0–40 m, 60–150 m and 200–1000 m (Fig. 2). MDS ordination

5, 2131-2164, 2008

Environmental influence on bacterial community structure





plot stress value was low (0.14) which indicates that the ordinations unlikely to mislead interpretation (Clarke and Warwick, 2001). We then used ANOSIM to determine the significant differences among bacterial communities from the three depth layers. This analysis showed the highest difference between surface and deeper community structures (R=0.754, p=0.001). Bacterial communities from the intermediate layer were closer to the upper layer than to the deeper layer but still significantly different from both (R=0.32 p=0.001 and R=0.22, p=0.004, respectively).

5

3.2 Bacterial community structure and physico-chemical variables ecological analysis

We first performed CCA using all physico-chemical parameters as constrained variables of the bacterial community diversity from CE-SSCP fingerprints of 0-1000 m depth samples. CCA revealed values higher than 20 of the variance inflation factor for nitrate and silicate, which indicated that these variables were almost perfectly correlated with other variables in the dataset. A strong Spearman's rank pairwise correlation between nitrate and silicate (R=0.91, p<0.01) allowed us to use nitrate as a proxy of silicate to perform CCA together with the rest of physico-chemical parameters. A summary of the results of this CCA is shown in Table 1. The total variance of fitted 0-1000 m samples diversity data when using physico-chemical variables to explain their bacterial community structure was 0.237, as indicated by the sum of all canonical eigenvalues (ter Braak and Smilauer, 2002). The cumulative percentage variance of species-environment relation indicates that the first and second canonical 20 axes account for 38.4 and 17.9% of this variance respectively (Table 1). Consequent axes accounted for less than 10% of the variance each, and are not further considered here. The eigenvalues of the ordination analyses, which also measure their relative

importance are small (<0.3) implying that CCA does not strongly separates species niches and therefore the biplot rule must be used to interpret the fitted relative abundances of samples with respect to physico-chemical variables in the CCA ordination diagram (Fig. 3). Monte Carlo test for first and all canonical axes were highly significant (p<0.01) indicating that the parameters selected are good explanatory variables of

BGD 5, 2131-2164, 2008 **Environmental** influence on bacterial community structure J. F. Ghiglione et al. **Title Page** Introduction Abstract Conclusions References **Tables Figures** Back Close Full Screen / Esc **Printer-friendly Version** Interactive Discussion

the community structure. First canonical axis is highly negatively correlated with depth, salinity, nitrate and PO_4 (ca. -0.5) and positively correlated with oxygen and DOC (ca. 0.4) and to a lesser extend with temperature and PAR (ca. 0.2). This first synthetic gradient clearly separates samples in the three depth layers observed when using di-

- ⁵ versity data alone indicating that their community composition varies with variation in concentrations of these parameters. Because depth was not collinear with the other variables correlated with first axis, we include this variable in the analysis as a proxy of pressure effect on bacterial community structure. Surface water samples further spread out along the second canonical axis (Fig. 3). We infer that 60 m depth samples have a proving the second canonical axis (Fig. 3).
- ¹⁰ have similar concentrations of nitrite, ammonia and CDOM variables elements than 40 m samples and together form a group independent from that of the 5–20 m samples (with the exception of C-20m sample). The significant contribution of these variables in defining surface bacterial community structures was confirmed by additional CCAs using only shallower samples (data not shown).
- 15 3.3 Bacterial community structure and lipid biomarkers ecological analysis

CCA total variance of the fitted CE-SSCP diversity from 0–1000 m depth samples when using dissolve lipid biomarkers to explain this diversity is 0.093 with the first two axes accounting for 72.5% of this variance (Table 1). Eigenvalues for these axes were 0.050 and 0.018, respectively. The Monte Carlo test for first and all canonical axes rendered
 a significant p-value of 0.02. The lipid biomarkers explain therefore the difference in bacterial community structures among samples according to the CCA biplot in Fig. 4. The first synthetic gradient was strongly positively correlated with lipids from chloroplast origin (LC, ca. 0.8) followed by triglycerides (TG, ca. 0.4). Samples from the upper 0–20 m layer, and especially samples from time B-5 m and 20 m, time C-5 m and 20 m

and time D-20 m were separated from the rest according to this first axis. The rest of samples formed a relatively tighter group with no clear difference between layers at this first axis indicating the homogeneity of LC at these samples (Fig. 4). Some samples from this last deeper group further spread out along the second axis, which is slightly Printer-friendly Version

Interactive Discussion



positively correlated with TG (ca. 0.3), and to a lesser extent with the remaining lipid biomarkers analyzed in this study. However, samples appear mixed with respect to their depth along this axis.

3.4 Distribution of phytoplankton pigments

25

- ⁵ Because phytoplankton pigments were almost undetectable below 150 m depth, all analyses with phytoplankton diversity data included 0–150 m depth samples only. Total chlorophyll-*a* (TChl-*a*) maxima increased during the one month cruise period of this study from 0.45–0.44 µg l⁻¹ at 50 m depth for times A and B, respectively, to 0.60 µg l⁻¹ at 40 m depth for time C and 0.80 µg l⁻¹ at 50 m depth for time D. The study period was
 ¹⁰ characterized by the predominance of small species (nano- and pico-phytoplankton) over microphytoplankton. The principal classes observed were prochlorophytes, cyanobacteria (*Synechococcus*) together with prymnesiophytes, pelagophytes and prasinophytes (with concentrations of these classes varying from 15 to 19% of TChl-*a*). Diatoms and dinoflagellates were lesser constituents of the phytoplankton (from 15 to 5% of TChl-*a*). We observed very important variations in the vertical distribution
- ¹⁵ 3 to 5% of TChl-*a*). We observed very important variations in the vertical distribution of specific biomasses. As an example, euglenophytes were minor constituents of the TChl-*a* biomass (<2%) and present at very low concentrations during the whole sampling period except for their high concentration (almost 10% of TChl-*a*) in the surface layer 0–20 m at date A (Marty et al., 2008¹).
- 20 3.5 Bacterial community structure and phytoplankton pigments ecological analysis

CCA when using pigment analysis-based phytoplankton diversity to explain CE-SSCP bacterial community structure gave a total variance of fitted 0–150 m samples diversity data of 0.200 (Table 1). Axes 1 and 2 accounted for 56.2% of this variance being the other axes negligible. Eigenvalues for the two first axes were 0.064 and 0.048 respectively. Monte Carlo permutation tests of first and all canonical axes were highly significant (p-values<0.01). Canonical axis 1 is highly positively correlated with *Syne*-





chococcus and euglenophytes (ca. ~0.5) and, to a lesser extend, positively correlated with dinoflagelates and negatively to pelagophytes and diatoms (ca. \pm 0.3) (Fig. 5). This first axis clearly distinguished between samples from the two shallower depth layers described above based on higher concentrations of *Synechococcus*, euglenophytes and

- dinoflagelates in the 0–20 m depth samples vs. the higher levels of pelagophytes and diatoms in the deeper 60–150 m layer. The second axis was correlated with higher abundances of prasinophytes and prochlorophytes further distinguishing the bacterial communities from samples at 40–60 m with the exception of B-60 m and C-60 m samples. Primnesiophytes, chlorophytes and cryptophytes variation had slight influence on the distribution of applicated by their apple areas.
- the distribution of samples as indicated by their small arrow length (Fig. 5).

4 Discussion

4.1 CE-SSCP incomes to bacterial community profiling studies

Culture-independent, sequence-based community fingerprinting techniques such as CE-SSCP (Lee et al., 1996), terminal restriction fragment length polymorphism (T RFLP) (Marsh, 1999), or automated ribosomal intergenic spacer analysis (A-RISA) (Borneman and Triplett, 1997) allow comparison of a theoretically infinite number of samples. Application of these techniques to environmental samples yields considerable amount of information on their microbial diversity composition that can be used to assess how environmental factors contribute to changes in microbial community structure. However, the introduction of experimental biases during analysis and the interpretation of the complex data generated still represent major challenges for the ap-

- plication of these methods in microbial ecology. Biases during nucleic acid extraction, PCR amplification (Wintzingerode et al., 1997) and 16S rDNA copy number variability (Farrelly et al., 1995) may lead to a distorted view of bacterial community structure. In this study, several tests were performed to investigate the efficiency and reproducibility
- of the DNA extraction and PCR–CE–SSCP protocols. Cell lysis was monitored several

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





times by microscopy, which assured that there was no cells recalcitrant to our lysis procedure. The quality of the DNA was carefully assessed before proceeding to 16S rDNA gene amplification. Only single PCR products were used for subsequent analyses. The use of an internal standard co-migrating with each sample PCR product

- ⁵ allowed us to normalize peak mobilities for comparative sample analysis. This procedure results in a high reproducibility of both, peak pattern and peak intensities, as previously observed (Ghiglione et al., 2005). Finally, because all samples in this study are subject to similar biases, comparative analysis of microbial community structure and relative abundances are still possible (Hughes et al., 2001). On the other hand,
- the use of adequate numerical and statistical tools overcomes the difficulty of analyzing complex datasets. Until now most studies have used indirect gradient analysis such as principal component analysis or cluster analysis to infer the influence of environmental variables on microbial diversity data, leaving the use of direct gradient analysis approaches such as redundancy analysis or canonical correspondence analysis (CCA)
- neglected (Ramette, 2007). Here we compared both indirect and direct gradient multivariate approaches to explore relationships between bacterial community structure as obtained by CE-SSCP with respect to concomitant measurements of physico-chemical parameters, lipid class biotracers and pigment analysis-based phytoplankton diversity.

4.2 Indirect gradient multivariate analysis to study bacterial community composition

- The DYFAMED offshore station is an ideal sampling site for 1-D studies because it is far enough from the Ligurian current to be protected from lateral transport (Andersen and Prieur, 2000). During the late summer-autumn oligotrophic conditions, biotic parameters showed a temporarily stable organization along the 0–1000 m water column. Bacterial community structure appeared differentiated in three layers, shallow
- (5-40 m), intermediate (60-150 m) and mesopelagic (200-1000 m), when using multivariate (Fig. 2) and cluster analyses. The robustness of this strong stratification with depth pattern was confirmed by a greater similarity within than between samples from each of these layers. This study adds to previous findings of a strong stratification pat-

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





tern with depth of bacterial communities in oligotrophic conditions (Lee and Fuhrman, 1991; Acinas et al., 1997; Murray et al., 1998; Riemann et al., 1999; Moeseneder et al., 2001; Ghiglione et al., 2007). Vertical and temporal dynamics of bacterial abundance and activity during this cruise are depicted in Mével et al. (2008).

5 4.3 Linking bacterial community profiling to the environment by direct gradient multivariate analysis

To unravel the environmental parameters eventually responsible for the diversity pattern observed, we chose direct gradient canonical correspondence analysis (CCA). Long ago developed by community ecologists to elucidate the relationships between biological assemblages of species and their environment, it is only recently that CCA has been applied to microbial assemblages from marine (Cordova-Kreylos et al., 2006; Klaus et al., 2007; Sapp et al., 2007), lake (Yannarell and Triplett, 2005) and soil (Salles et al., 2004) systems. Problems with this method may arise when explanatory variables are intercorrelated, making indistinguishable their separate effects. To avoid this problem, we performed separate analysis of measured environmental variables by organizing them in three ranks: physico-chemical variables, lipid biomarkers and phytoplankton pigments. We also observed Spearman's rank pairwise correlations between variables within each rank and avoided co-linear variables when performing CCA.

4.4 CCA with physico-chemical parameters

- ²⁰ Our CCA analyses, using physico-chemical data as explanatory variables, showed a temporarily stable vertical stratification of the bacterial community. At the DYFAMED station, under the oligotrophic conditions observed, the diversity data alone is statistically significantly related to changes in concentrations of salinity, nitrate and phosphate variables and to a lesser extend oxygen, DOC, PAR and temperature (Fig. 3). Physical
- ²⁵ and chemical parameters are usually studied in combination with bacterial activity or bacterial growth to demonstrate their influence on the prokaryotic component of the ma-

BGD			
5, 2131–2164, 2008			
Environmental influence on bacterial community structure J. F. Ghiglione et al.			
Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
14	►I.		
•	4 Þ		
Back	Close		
Full Screen / Esc			
Printer-friendly Version			
Interactive Discussion			

rine food web (Caron, 1994; Downing et al., 1999), sometimes with specific changes along the water column (Riemann et al., 1999). At the DYFAMED station, bacterial activity was limited by phosphorus above the deep-chlorophyll maximum (DCM) level (0–40 m) and by carbon at intermediate depth (60–150 m) for the same seasonal sampling time as in our study (Van Wambeke et al., 2002). To our knowledge, however, no studies have demonstrated the direct influence of environmental parameters on the bacterial community structure of natural environmental gradients. The originality of the present study is to demonstrate that several physico-chemical parameters are acting in synergy with nearly equal contribution to the bacterial vertical stratification. A complex interaction of physico-chemical parameters is therefore necessary to predict bacterial community structure.

4.5 CCA with lipid class biotracers

Since lipid classes are useful markers to determine sources and fate of organic matter in marine systems (Parrish, 1988) they have widely been used as biological tracers (Parrish et al., 1995; Derieux et al., 1998; Goutx et al., 2000), but only once in conjunc-15 tion with bacterial community structure changes (Ghiglione et al., 2007). In the present study CCA analysis showed that dissolved lipids from chloroplast origin (LC), which indicate phytoplankton source of DOM (Goutx et al., 1990), were the main biotracer associated with bacterial community structure, due surely to their high concentration at particular samples of 5-20 m depths (Fig. 4). Inputs of LC from microalgae to the 20 dissolved carbon pool might occur through cell lysis, grazing by herbivorous and/or viral lysis. We found a significant correlation between dissolved LC concentrations and both Synechoccocus cells number and Synechoccocus to total picophytoplankton ratio (R^2 =0.55 and R^2 =0.61, p<0.01, n=36, respectively). This result, together with pico-phytoplancton (Synechococcus and Prochlorococcus) dominance in the sampling 25

site (see Results) let us infer that *Synechoccocus* cells are the major source of LC pools in our samples. The other four classes of dissolved lipid biomarkers contributed to a lesser extent to bacterial community structure differences among samples along





the water column perhaps because their biological signature is more difficult to track. Phospholipids (PL) are membrane compounds in biological cells (Gérin and Goutx, 1993), wax esters (WE), are storage compounds in zooplankton and bacteria (Sargent et al., 1977; Ishige et al., 2003), trigycerides (TG) are reserve compounds in zooplankton, microalgae, bacteria and flagellates (Lee et al., 1971; Parrish and Wangersky, 5 1987; Alvarez and Steinbüchel, 2002; Méjanelle et al., 2005) and metabolites (Met) are products of acyl-lipid hydrolysis, free fatty acids, free alcohols, diglycerides and monoglycerides (Goutx et al., 2003). Their marginal effect upon the sample scores in the ordination diagram is probably because, taken together, these biomarkers indicated the presence of freshly biosynthesized organic matter not exclusively from phytodetritus.

CCA with pigment analysis-based phytoplankton diversity 4.6

10

It is well known that marine phytoplankton biomass and species distributions vary along vertical profiles, an obvious example being the deep chlorophyll maximum found in many areas of the world's oceans. This distribution may affect prokaryotic assemblages as well, since phytoplankton distribution appears to be one of the most important biotic variables influencing planktonic bacterial growth (Azam et al., 1983; Cole et al., 1988). Some studies have proposed a possible relation between bacterial community structure and phytoplankton, but most of them were limited to the study of chlorophyll-a pigments. Phytoplankton pigment signatures obtained by high-pressure liquid chro-20 matography are widely used as a proxy of phytoplankton diversity in marine systems (Bustillos-Guzman et al., 1995; Mackey et al., 1998; Marty et al., 2002). We used this signature to determine how phytoplankton taxonomic composition explained bacterial community CE-SSCP profiles. We found that the stratification of bacterial commu-

nity structures in two shallow layers, 0-40 m and 60-150 m, coincided with the higher 25 presence of Synechococcus, euglenophytes and dinoflagelates in the first 40 m samples, while diatoms and pelagophytes were preferentially present at samples bellow 60 m. Furthermore, separation of samples situated at the chlorophyll-a maximum cor-



responds with a higher abundance of prochlorophytes and prasinophytes at this depth. The general preferential abundance of some of these groups along the water column coincides with our findings, particularly for *Synechococcus* (Veldhuis et al., 2005), diatoms (Latasa et al., 1992), and pelagophytes (Claustre et al., 1994). It is also the case for prochlorophytes which are preferentially located at the base of the chlorophyll maximum (Li and Wood, 1988; Marty et al., 2002) and for prasinophytes which have been described as forming localized maxima at depth in the Mediterranean Sea (Marie et al., 2006). Euglenophytes and dinoflagelates presence at the surface layers is particular to this study. On the other hand, because of their uniform distribution in the first

¹⁰ 150 m depth, it is not surprising that prymnesiophytes, cryptophytes and chlorophytes had low influence on bacterial community changes with depth. Our results are the first study to demonstrate a compartmentalized influence of each phytoplankton group on bacterial community structure stratification in the first 150 m of the water column.

5 Concluding remarks

- ¹⁵ Our study is perhaps the first example of such a complex biogeochemical dataset, which includes physico-chemical parameters, lipid class biotracers and pigment analysis-based phytoplankton diversity, analyzed together with microbial community profiles to unravel the mechanisms responsible for bacterial community structuring with depth in the marine environment. Direct gradient multivariate canonical analyses using
- these parameters as explanatory environmental variables of the molecular fingerprinting profiles along the water column revealed that the observed vertical zonation of bacterial communities in three layers, above, in or just below the chlorophyll maximum and deeper, is due to synergetic driving forces pertaining to the three biogeochemical levels analyzed in this study. Similarly, the interaction of several of these parameters
- ²⁵ further distinguished samples situated above and in the chlorophyll maximum (up to 150 m). The role of the quality of the organic matter (DOC and CDOM) is significant in the upper layers (Fig. 3) and we have been able to attribute it to a phytoplankton

BGD 5, 2131-2164, 2008 **Environmental** influence on bacterial community structure J. F. Ghiglione et al. **Title Page** Introduction Abstract Conclusions References **Tables Figures** 14 Back Close Full Screen / Esc

Printer-friendly Version

Interactive Discussion



origin (Figs. 4 and 5). The influence of phytoplankton was not limited to exudates released at the surface levels since different phytoplankton phylogenetic groups explain the bacterial community structure zonation in the first 150 m depth (Fig. 5). This finding is consistent with a notion of specific association between bacteria and phytoplankton.

⁵ This study set the basis to further investigations of the plausible mechanisms responsible of bacterioplankton distribution in marine systems. For instance, by enlarging the array of ecological factors explored such as virus or protozoan predation or competition between bacteria for carbon and nutrient sources.

Acknowledgements. We thank the captains and the crews of the RV "Thalassa" for their help
 during sampling. We thank the people involved in the physico-chemical parameters sampling
 and measurements, and especially R. Demesmay, A. Momzikoff and L. Prieur. We specifically acknowledge the contribution of our friend Valérie Andersen who was chief scientist during the Dynaproc2 cruise and passed away before she could participate in the valorisation of her oceanographic campaign. This work was supported by PROOF Grant provided by the
 INSU-CNRS and was part of the PECHE project "Production and exportation of carbon: control by heterotrophic organisms at small time scales"). We are grateful to John Dolan and France Van Wambeke for helpful comments on the manuscript.

References

20

Acinas, S. G., Rodriguez-Valera, F., and Pedros-Alio, C.: Spatial and temporal variation in marine bacterioplankton diversity as shown by RFLP fingerprinting of PCR amplified 16S

rDNA, FEMS Microbiol. Ecol., 24, 27–40, 1997.

Alvarez, H. M. and Steinbüchel, A.: Triacylglycerols in prokaryotic microorganisms, Appl.Microbiol. Biotechnol., 60, 367–376, 2002.

Andersen, R. A., Saunders, G. W., Paskind, M. P., and Sexton, J. P.: Ultrastructure and 18S

rRNA gene sequence for Pelagomonas calceolate gen. et sp. nov. and the description of a new algal class, the Pelagophyceae classis nov., J. Phycol., 29, 701–715, 1993.

Andersen, V. and Prieur, L.: One-month study in the open NW Mediterranean Sea (DYNAPROC experiment, May 1995): overview of the hydrobiogeochemical structures and effects of wind events, Deep-Sea Res. I, 47, 397–422, 2000.

BGD 5, 2131-2164, 2008 **Environmental** influence on bacterial community structure J. F. Ghiglione et al. **Title Page** Introduction Abstract Conclusions References **Tables Figures** Back Close Full Screen / Esc **Printer-friendly Version** Interactive Discussion

- Azam, F., Fenchel, T., Field, J. G., Gray, J. S., Meyerreil, L. A., and Thingstad, F.: The ecological role of water column microbes in the sea, Mar. Ecol. Prog. Ser., 10, 257–263, 1983.
- Barlow, R. G., Mantoura, R. F. C., Cummings, D. G., and Fileman, T. W.: Pigment chemotaxonomic distributions of phytoplankton during summer in the western Mediterranean, Deep-Soa Ros, Part II, 44, 832–850, 1997
- ⁵ Sea Res. Part II, 44, 833–850, 1997.

15

- Borneman, J. and Triplett, E. W.: Molecular microbial diversity in soil from Eastern Amazonia: evidence for unusual microorganisms and microbial population shifts associated with deforestation, Appl. Environ. Microbiol., 63, 2647–2653, 1997.
- Brosius, J., Dull, T., Sleeter, D. D., and Noller, H. F.: Gene organization and primary structure of a ribosomal RNA operon from Escherichia coli, J. Mol. Biol., 148, 107–127, 1981.
- Bustillos-Guzman, J., Claustre, H., and Marty, J. C.: Specific phytoplankton signatures and their relationship to hydrographic conditions in the coastal northwestern Mediterranean Sea, Mar. Ecology Prog. Ser., 124, 247–258, 1995.

Caron, D. A.: Inorganic nutrients, bacteria, and the microbial loop, Microb. Ecol., 28, 295–298, 1994.

- Cauwet, G.: HTCO method for dissolved organic carbon analysis in seawater: influence of catalyst on blank estimation, Mar. Chem., 47, 55–64, 1994.
 - Cho, B. C. and Azam, F.: Major role of bacteria in biogeochemical fluxes in the oceans interior, Nature, 332, 441–443, 1988.
- ²⁰ Cho, B. C. and Azam, F.: Biogeochemical significance of bacterial biomass in the ocean's euphotic zone, Mar. Ecol. Prog. Ser., 63, 253–259, 1990.
 - Clarke, K. R. and Warwick, R. W.: Change in marine communities: an approach to statistical analysis and interpretation, 2nd edition, Primer-E: Plymouth, 2001.

Claustre, H., Kerhervé, P., Marty, J. C., Prieur, L., Videau, C., and Hecq, J. H.: Phytoplankton

- dynamics associated with a geostrophic front: ecological and biogeochemical implications,
 J. Mar. Res., 52, 711–742, 1994.
 - Cole, J. J., Findlay, S., and Pace, M. L.: Bacterial production in fresh and saltwater ecosystems: a cross-system overview, Mar. Ecol. Prog. Ser., 43, 1–10, 1988.
 - Cordova-Kreylos, A. L., Cao, Y., Green, P. G., Hwang, H. M., Kuivila, K. M., Lamontagne, M.
- ³⁰ G., et al.: Diversity, composition, and geographical distribution of microbial communities in California salt marsh sediments, Appl. Environ. Microbiol., 72, 3357–3366, 2006.
 - Delbès, C., Godon, J. J., and Moletta, R.: 16S rDNA sequence diversity of a culture-accessible part of an anaerobic digestor bacterial community, Anaerobe, 4, 267–269, 1998.

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure



- Derieux, S., Fillaux, J., and Saliot, A.: Lipid class and fatty acid distributions in particulate and dissolved fractions in the North Adriatic Sea, Org. Geochem., 29, 1609–1621, 1998.
- Downing, J. A., Osenberg, C. W., and Sarnelle, O.: Meta-analysis of marine nutrient-enrichment experiments: variation in the magnitude of nutrient limitation, Ecology, 80, 1157–1167, 1999.
- 5 Farrelly, V., Rainey, F. A., and Stackebrandt, E.: Effect of genome size and rrn gene copy number on PCR amplification of 16S ribosomal-RNA genes from a mixture of bacterial species, Appl. Environ. Microbiol., 61, 2798–2801, 1995.
 - Fuhrman, J. A., Comeau, D. E., Hagstrom, A., and Chan, A. M.: Extraction from natural planktonic microorganisms of DNA suitable for molecular biological studies, Appl. Environ. Micro-

Ghiglione, J. F., Larcher, M., and Lebaron, P.: Spatial and temporal scales of variation in bacterioplankton community structure in the NW Mediterranean Sea, Aquat. Microb. Ecol., 40, 229-240, 2005.

15

20

Ghiglione, J. F., Mével, G., Pujo-Pay, M., Mousseau, L., Lebaron, P., and Goutx, M.: Diel and seasonal variations in abundance, activity, and community Structure of particle-attached and free-living bacteria in NW Mediterranean Sea, Microb. Ecol., 54, 217–231, 2007.

Godon, J. J., Duthoit, F., Delbes, C., Millet, L., and Montel, M. C.: Use of molecular fingerprint for the study of complex microbial ecosystem. Application to AOC Salers cheese, Lait, 81,

257-262, 2001.

Goericke, R. and Repeta, D.: The pigments of Prochlorococcus marinus: the presence of divinyl-chlorophylls a and b in a marine prokaryote, Limnol. Oceanogr., 37, 425–433, 1992.

Goutx, M., Gérin, C., and Bertrand, J. C.: An application of latroscan thin-layer chromatogra-

- phy with flame ionization detection lipids classes of microorganisms as biomarkers in the 25 marine environment, Org. Geochem., 16, 1231-1237, 1990.
 - Goutx, M., Guigue, C., and Striby, L.: Triacylglycerol biodegradation experiment in marine environmental conditions : definition of a new lipolysis index, Org. Geochem., 34, 1465-1473, 2003.
- 30 Goutx, M., Momzikoff, A., Striby, L., Andersen, V., Marty, J. C., and Vescovali, I.: High frequency fluxes of labile compounds in the Central Ligurian Sea, Northwestern Mediterranean, Deep Sea Res. I. 47, 533-556, 2000.

Holmes, R. M., Aminot, A., Kérouel, R., Hooker, B. A., and Petersen, B. J.: A simple and precise

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure

Title Page				
Abstract	Introduction			
Conclusions	References			
Tables	Figures			
	P1			
•	•			
Back	Close			
Full Screen / Esc				
Printer-friendly Version				
Interactive Discussion				



biol., 54, 1426–1429, 1988. 10

Gérin, C. and Goutx, M.: Separation and quantitation of phospholipids in marine bacteria by latroscan TLC/FID analysis, J. Plan. Chromatogr., 6, 307-312, 1993.

method for measuring ammonium in marine and freshwater ecosystems, Can. J. Fish. Aquat. Sci., 56, 1801–1808, 1999.

Hong, H., Pruden, A., and Reardon, K. F.: Comparison of CE-SSCP and DGGE for monitoring a complex microbial community remediating mine drainage, J. Microbiol. Meth., doi:

5 10.1016/j.mimet.2006.11.016, 2007.

10

Hughes, J. B., Hellmann, J. J., Ricketts, T. H., and Bohannan, B. J. M.: Counting the uncountable: Statistical approaches to estimating microbial diversity, Appl. Environ. Microbiol., 67, 4399–4406, 2001.

Ishige, T., Tani, A., Sakai, Y., and Kato, N.: Wax ester production by bacteria, Curr. Opin. Microbiol., 6, 244–250, 2003.

Jeffrey, S. W., Mantoura, R. F. C., and Bjornland, T.: Data for the identification of 47 key phytoplankton pigments, in: Phytoplankton pigments in oceanography, edited by: Jeffrey, S. W., Mantoura, R. F. C., and Wright, S. W., UNESCO, Paris, 449–559, 1997.

Jeffrey, S. W. and Vesk, M.: Introduction to marine phytoplankton and their pigment signatures,

- in: Phytoplankton pigments in oceanography, edited by: Jeffrey, S. W., Mantoura, R. F. C., and Wright, S. W., UNESCO, Paris, 37–84, 1997.
 - Klaus, J. S., Janse, I., Heikoop, J. M., Sanford, R. A., and Fouke, B. W.: Coral microbial communities, zooxanthellae and mucus along gradients of seawater depth and coastal pollution, Environ. Microbiol. Rev., 9, 1291–1305, 2007.
- 20 Kruskal, J. B.: Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis, Psychometrika, 29, 1–27, 1964.
 - Latasa, M., Estrada, M., and Delgado, M.: Plankton-pigment relationships in the Northwestern Mediterranean during stratification, Mar. Ecol. Prog. Ser., 88, 61–73, 1992.

Lee, D. H., Zo, Y. G., and Kim, S. J.: Nonradioactive method to study genetic profiles of natu-

- ral bacterial communities by PCR-single-strand-conformation polymorphism, Appl. Environ.
 Microbiol., 62, 3112–3120, 1996.
 - Lee, R. H., Nevenzel, J. C., and Paffenhofer, G. A.: Importance of wax esters and other lipids in the marine food chain: phytoplankton and copepods, Mar. Biol., 9, 99–108, 1971.

Lee, S. H. and Fuhrman, J. A.: Spatial and temporal variation of natural bacterioplankton as-

- semblages studied by total genomic DNA cross-hybridization, Limnol. Oceanogr., 36, 1277– 1287, 1991.
 - Li, W. K. W. and Wood, M.: Vertical distribution of North Atlantic ultraplankton: analysis by flow cytometry and epifluorescence microscopy, Deep-Sea Res. I, 35, 1615–1638, 1988.

5, 2131–2164, 2008

Environmental influence on bacterial community structure





Mackey, D. J., Higgins, H. W., Mackey, M. D., and Holdsworth, D.: Algal class abundances in the western equatorial Pacific: estimation from HPLC measurements of chloroplast pigments using CHEMTAX, Deep-Sea Res. I, 45, 1141–1168, 1998.

Marie, D., Zhu, F., Balaguie, V., Ras, J., and Vaulot, D.: Eukaryotic picoplankton communities

- of the Mediterranean Sea in summer assessed by molecular approaches (DGGE, TTGE, QPCR), FEMS Microb. Ecol., 55, 403–415, 2006.
 - Marsh, T. L.: Terminal restriction fragment length polymorphism (T-RFLP): an emerging method for characterizing diversity among homologous populations of amplification products, Curr. Opin. Microbiol., 2, 323–327, 1999.
- Martinez, J., Smith, D. C., Steward, G. F., and Azam, F.: Variability in ectohydrolytic enzyme activities of pelagic marine bacteria and its significance for substrate processing in the sea, Aquat. Microb. Ecol., 10, 223–230, 1996.

Marty, J. C., Chiaverini, J., Pizay, M. D., and Avril, B.: Seasonal and interannual dynamics of nutrients and phytoplankton pigments in the western Mediterranean Sea at the DYFAMED time-series station (1991–1999), Deep Sea Res. II, 49, 1965–1985, 2002.

- Méjanelle, L., Laureillard, J., and Rassoulzadegan, F.: Polar lipids biomarkers of free living bacteria from oligotrophic marine bacteria, Biogeochem., 72, 365–383, 2005.
 - Mével, G., Vernet, M., and Ghiglione, J. F.: Short-term temporal variations of heterotrophic bacterial abundance and production in the open NW Mediterranean Sea, Biogeosciences
- ²⁰ Discuss., 5, 1899–1932, 2008,

15

http://www.biogeosciences-discuss.net/5/1899/2008/.

- Moeseneder, M., Winter, C., and Herndl, G. J.: Horizontal and vertical complexity of attached and free-living bacteria in the eastern Mediterranean Sea, determined by 16S rDNA and 16S rRNA fingerprints, Limnol. Oceanogr., 46, 95–107, 2001.
- ²⁵ Murray, A. E., Preston, C. M., Massana, R., Taylor, L. T., Blakis, A., Wu, K., and DeLong, E. F.: Seasonal and spatial variability of bacterial and archaeal assemblages in coastal waters near Anvers Island, Antartica, Appl. Environ. Microbiol., 64, 2585–2595, 1998.

Parrish, C. C.: Dissolved and particulate marine lipid classes: a review, Mar. Chem., 23, 17–40, 1988.

Parrish, C. C. and Wangersky, P. J.: Particulate and dissolved ipid classes in cultures of Phaeodactylum tricornutum grown in cage culture turbidostats with a range of nitrogen supply rates, Mar. Ecol. Progr. Ser., 35, 119–128, 1987.

Parrish, C. C., McKenzie, C. H., MacDonald, B. A., and Hatfield, E. A.: Seasonal studies of

BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure





seston lipids in relation to microplankton species composition and scallop growth in South Broad Cove, Newfoundland, Mar. Ecol. Prog. Ser., 129, 151–164, 1995.

- Pinhassi, J., Zweifel, U. L., and Hagstrom, A.: Dominant marine bacterioplankton species found among colony-forming bacteria, Appl. Environ. Microbiol., 63, 3359–3366, 1997.
- ⁵ Polz, M. F., Hunt, D. E., Preheim, S. P., and Weinreich, D. M.: Patterns and mechanisms of genetic and phenotypic differentiation in marine microbes, Philosoph, Transact. Royal Soc. B-Biol. Sci., 361, 2009–2021, 2006.

Ramette, A.: Multivariate analyses in microbial ecology, FEMS Microbiol. Ecol., 62, 142–160, 2007.

Riemann, L., Steward, G. F., Fandino, L. B., Campbell, L., Landry, M. R., and Azam, F.: Bacterial community composition during two consecutive NE Monsoon periods in the Arabian Sea studied by denaturing gradient gel electrophoresis (DGGE) of rRNA genes, Deep-Sea Res. II, 46, 1791–1811, 1999.

Salles, J. F., van Veen, J. A., and van Elsas, J. D.: Multivariate analyses of Burkholderia species

¹⁵ in soil: effect of crop and land use history, Appl. Environ. Microbiol, 70, 4012–4020, 2004. Sapp, M., Wichels, A., Wiltshire, K. H., and Gerdts, G.: Bacterial community dynamics during winter-spring transition in the North Sea, FEMS Microbiol. Ecol., 59, 622–637, 2007.

Sargent, J. R., Gatten, R. R., and MacIntosh, R.: Wax esters in the marine environment: their occurence, formation, transformation and ultimate fate, Mar. Chem., 5, 573–584, 1977.

²⁰ Smalla, K., Oros-Sichler, M., Milling, A., Heuer, H., Baumgarte, S., Becker, R., et al.: Bacterial diversity of soils assessed by DGGE, T-RFLP and SSCP fingerprints of PCR-amplified 16S rRNA gene fragments: Do the different methods provide similar results?, J. Microbiol. Meth., 69, 470–479, 2007.

Striby, L., Lafont, R., and Goutx, M.: Improvement in the latroscan thin-layer chromatographic-

- flame ionisation detection analysis of marine lipids. Separation and quantitation of monoacylglycerols and diacylglycerols in standards and natural samples, J. Chromatogr., 849, 371– 380, 1999.
 - Sugimura, Y. and Suzuki, Y.: A high-temperature catalytic oxidation method for the determination of non-volatile dissolved organic carbon in seawater by direct injection of a liquid sample,
- ³⁰ Mar. Chem., 24, 105–131, 1988.
 - Sunnucks, P., Wilson, A. C. C., Beheregaray, L. B., Zenger, K., French, J., and Taylor, A. C.: SSCP is not so difficult: the application and utility of single-stranded conformation polymorphism in evolutionary biology and molecular ecology, Mol. Ecol., 9, 1699–1710, 2000.

BGD

5, 2131–2164, 2008

Environmental influence on bacterial community structure

Title Page				
Abstract	Introduction			
Conclusions	References			
Tables	Figures			
	►I			
•	•			
Back	Close			
Full Screen / Esc				
Printer-friendly Version				
Interactive Discussion				



- Ter Braak, C. J. F.: Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis, Ecology, 67, 1167–1179, 1986.
- Ter Braak, C. J. F. and Šmilauer, P.: CANOCO Reference Manual and CanoDraw for Windows User's Guide: Software for Canonical Community Ordination (version 4.5), Microcomputer Power, Ithaca NY, USA, 2002.

5

10

- Tréguer, P. and LeCorre, P.: Manuel d'analyse des sels nutritifs dans l'eau de mer, Utilisation de l'AutoAnalyser II Technicon. 2nd ed., 1975.
- Van Wambeke, F., Christaki, U., Giannakourou, A., Moutin, T., and Souvemerzoglou, K.: Longitudinal and vertical trends of bacterial limitation by phosphorus and carbon in the Mediterranean Sea, Microb. Ecol., 43, 119–133, 2002.
- Veldhuis, M. J. W., Timmermans, K. R., Croot, P., and Van der Wagt, B.: Picophytoplankton: a comparative study of their biochemical composition and photosynthetic properties, J. Sea Res., 53, 7–24, 2005.

Vidussi, F., Claustre, H., BustillosGuzman, J., Cailliau, C., and Marty, J. C.: Determination

- of chlorophylls and carotenoids of marine phytoplankton: separation of chlorophyll a from divinyl-chlorophyll a and zeaxanthin from lutein, J. Plankton Res., 18, 2377–2382, 1996.
 - Wintzingerode, F., Göbel, U. B., and Stackebrandt, E.: Determination of microbial diversity in environmental samples: pitfalls of PCR-based rRNA analysis, FEMS Microbiol. Ecol. Rev., 21, 213–229, 1997.
- Wright, S. W. and Jeffrey, S. W.: Fucoxanthin pigment markers of marine phytoplankton analysed by HPLC and HPTLC, Mar. Ecol. Progr. Ser., 38, 259–266, 1987.
 - Yannarell, A. C. and Triplett, E. W.: Geographic and environmental sources of variation in lake bacterial community composition, Appl. Environ. Microbiol., 71, 227–239, 2005.
- Zemb, O., Haegeman, B., Delgenes, J. P., Lebaron, P., and Godon, J. J.: Safum: statistical analysis of SSCP fingerprints using PCA projections, dendrograms and diversity estimators,
- Mol. Ecol. Notes, 7(5), 767–770, doi:10.1111/j.1471-8286.2007.01882.x., 2007.

BGD

5, 2131–2164, 2008

Environmental influence on bacterial community structure





Table 1. Summary of results from canonical correspondence analyses of the bacterioplankton community structure data when constrained by physico-chemical, lipid biomarkers or phyto-plankton environmental variables, respectively.

Environmental variables	Physico-chemical		Lipid biomarkers		Phytoplankton	
	(0–1000 m)		(0–1000 m)		(0–150 m)	
Total inertia	0.518		0.415		0.449	
Sum of all canonical eigenvalues	0.237		0.093		0.200	
	Axis 1	Axis 2	Axis 1	Axis 2	Axis 1	Axis 2
Eigenvalues Species-environment correlations Cumulative percentage variance of – species data – species-environment relation	0.091 0.926 17.6 38.4	0.042 0.876 25.8 56.3	0.050 0.761 9.6 53.6	0.018 0.714 13.0 72.5	0.064 0.959 14.3 32.0	0.048 0.850 25.1 56.2

BGD

5, 2131–2164, 2008

Environmental influence on bacterial community structure

Title Page				
Abstract	Introduction			
Conclusions	References			
Tables	Figures			
	►I			
•	•			
Back	Close			
Full Screen / Esc				
Printer-friendly Version				
Interactive Discussion				







Printer-friendly Version

Full Screen / Esc











Fig. 3. Canonical correspondence analysis of bacterioplankton community structure from samples from 0 to 1000 m depth using physico-chemical parameters. Arrows point in the direction of increasing values of each variable. The length of the arrows indicates the degree of correlation with the represented axes. The position of samples relative to arrows is interpreted by projecting the points on the arrow and indicates the extent to which a sample bacterial community composition is influenced by the environmental parameter represented by that arrow. Samples from different depth groups (0–20 m, 60–150 m and 200–1000 m) are indicated by different symbols.





BGD

5, 2131-2164, 2008

Environmental influence on bacterial community structure

J. F. Ghiglione et al.





Fig. 4. Canonical correspondence analysis of bacterioplankton community structure from samples from 0 to 1000 m depth using lipid biomarker parameters. For interpretation see Fig. 3. LC: lipids from chloroplastic origin, TG: triglycerides, Met: metabolites, PL: phospholipids, WE: wax esters.



BGD 5, 2131-2164, 2008 **Environmental** influence on bacterial community structure J. F. Ghiglione et al. **Title Page** Introduction Abstract Conclusions References Tables **Figures** 14 Back Close Full Screen / Esc **Printer-friendly Version** Interactive Discussion



Fig. 5. Canonical correspondence analysis of bacterioplankton community structure from samples from 0 to 150 m depth using phytoplankton pigments variability. For interpretation see Fig. 3.