



**HAL**  
open science

# The Gevrey Hypoellipticity for linear and non-linear Fokker-Planck equations

Hua Chen, Wei-Xi Li, Chao-Jiang Xu

► **To cite this version:**

Hua Chen, Wei-Xi Li, Chao-Jiang Xu. The Gevrey Hypoellipticity for linear and non-linear Fokker-Planck equations. *Journal of Differential Equations*, 2009, 246, pp.320-339. hal-00369846

**HAL Id: hal-00369846**

**<https://hal.science/hal-00369846>**

Submitted on 22 Mar 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# THE GEVREY HYPOELLIPTICITY FOR LINEAR AND NON-LINEAR FOKKER-PLANCK EQUATIONS

Hua Chen<sup>1</sup> & Wei-Xi Li<sup>1</sup> & Chao-Jiang Xu<sup>1,2</sup>

<sup>1</sup>School of Mathematics and Statistics, Wuhan University,  
Wuhan 430072, China  
email: chenhua@whu.edu.cn  
wei-xi.li@whu.edu.cn

<sup>2</sup>Université de Rouen, UMR 6085-CNRS, Mathématiques  
Avenue de l'Université, BR.12,76801 Saint Etienne du Rouvray, France  
email: Chao-Jiang.Xu@univ-rouen.fr

ABSTRACT. In this paper, we study the Gevrey regularity of weak solution for a class of linear and quasilinear Fokker-Planck equations.

## 1. INTRODUCTION

Recently, a lot of progress has been made on the study for the spatially homogeneous Boltzmann equation without angular cutoff, cf. [2, 3, 8, 22] and references therein, which shows that the singularity of collision cross-section yields some gain of regularity in the Sobolev space frame on weak solutions for Cauchy problem. That means, this gives the  $C^\infty$  regularity of weak solution for the spatially homogeneous Boltzmann operator without angular cutoff. The local solutions having the Gevrey regularity have been constructed in [21] for initial data having the same Gevrey regularity, and a general Gevrey regularity results have given in [17] for spatially homogeneous and linear Boltzmann equation of Cauchy problem for any initial data. In the other word, there is the smoothness effect similar to heat equation.

However, there is no general theory for the spatially inhomogeneous problems. It is now a kinetic equation in which the diffusion part is nonlinear operator of velocity variable. In [1], by using the uncertainty principle and microlocal analysis, they obtain a  $C^\infty$  regularity results for linear spatially inhomogeneous Boltzmann equation without angular cutoff.

In this paper, we will study the Gevrey regularity of weak solution for the the following Fokker-Planck operator in  $\mathbb{R}^{2n+1}$

$$(1.1) \quad \mathcal{L} = \partial_t + v \cdot \partial_x - a(t, x, v) \Delta_v,$$

where  $\Delta_v$  is Laplace operator of velocity variables  $v$ .

The motivation of study for this class of operator is , as in [15], attempt to study inhomogeneous Boltzmann equations without angular cutoff and non linear Vlasov-Fokker-Planck equation (see [11, 12]).

Before stating the result, we recall the definition of Gevrey class function. Let  $U$  be an open subset of  $\mathbb{R}^N$  and  $f$  be a real function defined in  $U$ . We say  $f \in G^s(U)$  ( $s \geq 1$ ) if  $f \in C^\infty(U)$  and for any

---

Research partially supported by NSFC.

compact subset  $K$  of  $U$ , there exists a constant  $C = C_K$ , depending only on  $K$ , such that for all multi-indices  $\alpha \in \mathbb{N}^N$  and for all  $x \in K$

$$(1.2) \quad |\partial^\alpha f(x)| \leq C_K^{|\alpha|+1} (|\alpha|!)^s.$$

Denote by  $\bar{U}$  the closure of  $U$  in  $\mathbb{R}^N$ . We say  $f \in G^s(\bar{U})$  if  $f \in G^s(W)$  for some open neighborhood  $W$  of  $\bar{U}$ . The estimate (1.2) for  $x \in K$  is valid if and only if the following one is valid ( cf.Chen hua-Rodino[5] or Rodino[18]):

$$\|\partial^\alpha f\|_{L^2(K)} \leq C_K^{|\alpha|+1} (|\alpha|)^{s|\alpha|}.$$

In this paper, we use the above estimate in  $L^2$ .

We say an operator  $P$  is  $G^s$  hypoelliptic in  $U$  if  $u \in \mathcal{D}'$ ,  $Pu \in G^s(U)$  implies  $u \in G^s(U)$ . Likewise, we say an operator  $P$  is  $C^\infty$  hypoelliptic in  $U$  if  $u \in \mathcal{D}'$ ,  $Pu \in C^\infty(U)$  implies  $u \in C^\infty(U)$ .

The operator  $\mathcal{L}$  satisfies the Hörmander' condition. By virtue of the results of Hörmander [5], we know that  $\mathcal{L}$  is  $C^\infty$  hypoelliptic. In the aspect of Gevrey class, Derridj-Zuily [7] proved that  $\mathcal{L}$  is  $G^s$ -hypoelliptic for  $s > 6$  in a general form of Hörmander's operators.

In this paper, we improve firstly the results of [7] for Fokker-Planck operators as the following theorems.

**Theorem 1.1.** *For any  $s \geq 3$ , if the coefficient  $a$  is in  $G^s(\mathbb{R}^{2n+1})$  and  $a(t, x, v) \geq c_0 > 0$ , then the operator  $\mathcal{L}$  given in (1.1) is  $G^s$  hypoelliptic in  $\mathbb{R}^{2n+1}$ ,*

Of course, Theorem 1.1 is also true for the following general operators,

$$\tilde{\mathcal{L}} = \partial_t + A(v) \cdot \partial_x - \sum_{j,k=1}^n a_{jk}(t, x, v) \partial_{v_j}^2 \partial_{v_k}$$

in an open domain  $U$  of  $\mathbb{R}^{2n+1}$ , where  $A$  is a non singular  $n \times n$  constant matrix,  $(a_{jk}(t, x, v))$  is positive defined on  $U$  and belongs to  $G^s(U)$ .

**Remark** Our results is a local and interior regularity results, that means if there exists a weak solution in  $\mathcal{D}'$ , then this solution is in Gevrey class in interior of domain. So that if the weak solution is a solution of the Cauchy problem, we don't need the regularity of initial data.

Secondly, we consider the quasi-linear equation

$$(1.3) \quad \partial_t u + v \cdot \nabla_x u - a \Delta_v u = F(t, x, v, u, \nabla_v u)$$

where  $F$  is nonlinear function of real variable  $(t, x, v, w, p)$ .

**Theorem 1.2.** *Let  $u$  be a weak solution of equation (1.3) such that  $u, \nabla_v u \in L_{loc}^\infty(\mathbb{R}^{2n+1})$ , then*

$$u \in G^s(\mathbb{R}^{2n+1})$$

for any  $s \geq 3$ , provided the coefficients  $a$  is in  $G^s(\mathbb{R}^{2n+1})$ ,  $a(t, x, v) \geq c_0 > 0$  and nonlinear function  $F(t, x, v, w, p)$  is in  $G^s(\mathbb{R}^{2n+2+n})$ .

**Remark :** If the nonlinear term  $F$  is independent of  $p$  or in the form of

$$\nabla_v(F(t, x, v, u)),$$

we can suppose that the weak solution  $u \in L_{loc}^\infty(\mathbb{R}^{2n+1})$ .

The plan of this paper is as follows : In section 2, we obtain a sharp subelliptic estimate for the Fokker-Planck operator  $\mathcal{L}$  via direct computation, and then prove the Gevrey hypoellipticity of  $\mathcal{L}$ . In section 3, we prove the Gevrey regularity for the weak solutions of the quasi-linear Fokker-Planck equation.

## 2. SUBELLIPTIC ESTIMATE

We recall firstly some notations,  $\|\cdot\|_\kappa, \kappa \in \mathbb{R}$ , is the classical Sobolev norm in  $H^\kappa(\mathbb{R}^{2n+1})$ , and  $(h, k)$  is the inner product of  $h, k \in L^2(\mathbb{R}^{2n+1})$ . Moreover if  $f, g \in C_0^\infty(\mathbb{R}^{2n+1})$ , from Hölder inequality and Young inequality, for any  $\varepsilon > 0$ ,

$$(2.1) \quad |(f, g)| \leq \|h\|_\kappa \|g\|_{-\kappa} \leq \frac{\varepsilon \|h\|_\kappa^2}{2} + \frac{\|g\|_{-\kappa}^2}{2\varepsilon}.$$

We have also the interpolation inequality for Sobolev space, for any  $\varepsilon > 0$  and any  $r_1 < r_2 < r_3$ ,

$$(2.2) \quad \|h\|_{r_2} \leq \varepsilon \|h\|_{r_3} + \varepsilon^{-(r_2-r_1)/(r_3-r_2)} \|h\|_{r_1}.$$

Let  $\Omega$  be an open subset of  $\mathbb{R}^{2n+1}$ . We denote by  $S^m = S^m(\Omega), m \in \mathbb{R}$ , the symbol space of classical pseudo-differential operator and  $P = P(t, x, v, D_t, D_x, D_v) \in \text{Op}(S^m)$  a pseudo-differential operator of symbol  $p(t, x, v; \tau, \xi, \eta) \in S^m$ . If  $P \in \text{Op}(S^m)$ , then  $P$  is a continuous operator from  $H_c^\kappa(\Omega)$  to  $H_{loc}^{\kappa-m}(\Omega)$ . Here  $H_c^\kappa(\Omega)$  is the subspace of  $H^\kappa(\mathbb{R}^{2n+1})$  consisting of the distributions having their compact support in  $\Omega$ , and  $H_{loc}^{\kappa-m}(\Omega)$  consists of the distributions  $h$  such that  $\phi h \in H^{\kappa-m}(\mathbb{R}^{2n+1})$  for any  $\phi \in C_0^\infty(\Omega)$ . The more properties can be found in the Treves' book [20]. Remark that if  $P_1 \in \text{Op}(S^{m_1}), P_2 \in \text{Op}(S^{m_2})$ , then  $[P_1, P_2] \in \text{Op}(S^{m_1+m_2-1})$ .

Now we show a sharp subelliptic estimate for the operator  $\mathcal{L}$ , our proof bases on the work of Bouchut [4] and Morimoto-Xu [15].

**Proposition 2.1.** *Let  $K$  be a compact subset of  $\mathbb{R}^{2n+1}$ . Then for any  $r \geq 0$ , there exists a constant  $C_{K,r}$ , depending only on  $K$  and  $r$ , such that for any  $f \in C_0^\infty(K)$ ,*

$$(2.3) \quad \|f\|_r \leq C_{K,r} \{ \|\mathcal{L}f\|_{r-2/3} + \|f\|_0 \}.$$

To simplify the notation, in this section we will denote by  $C_K$  the different suitable constants depending only on  $K$ . We have firstly the following three lemmas, which establish the gain of regularity in the velocity variable  $v$ , in the space variable  $x$  and in the time variable  $t$ , respectively.

**Lemma 2.2.** *There exists a constant  $C_K$  such that for any  $f \in C_0^\infty(K)$ ,*

$$\|\nabla_v f\|_0 \leq C_K \{ |(\mathcal{L}f, f)| + \|f\|_0 \}.$$

Moreover, for any  $r \geq 0$ , for any  $\varepsilon > 0$ ,

$$\|\nabla_v f\|_r \leq \varepsilon \|\mathcal{L}f\|_r + C_{K,\varepsilon} \|f\|_r.$$

We get a gain of regularity of order 1 for  $v$  variable. This is obtained directly by the positivity of coefficient  $a$  and compact support of  $f$ . For the space variable  $x$ , we have also the following subelliptic estimate.

**Lemma 2.3.** *There exists a constant  $C_K$  such that for any  $f \in C_0^\infty(K)$ ,*

$$\|D_x^{2/3} f\|_0 \leq C_K (\|\mathcal{L}f\|_0 + \|f\|_0),$$

where  $D_x^{2/3} = (-\Delta_x)^{1/3}$ .

This is a result of [4], and it is deduced by following two estimates

$$\|D_x^{2/3} f\|_0 \leq C_K \|\Delta_v f\|_0^{1/3} \|\partial_t f + v \cdot \partial_x f\|_0^{2/3},$$

and

$$\|\Delta_v f\|_0 \leq C_K (\|\mathcal{L}f\|_0 + \|f\|_0).$$

For the time variable  $t$ , we have also a gain of regularity of order 2/3.

**Lemma 2.4.** *There exists a constant  $C_K$  such that for any  $f \in C_0^\infty(K)$ ,*

$$\|\partial_t f\|_{-1/3} \leq C_K (\|\mathcal{L}f\|_0 + \|f\|_0).$$

In fact, we have

$$\|\partial_t f\|_{-1/3} = \|\Lambda^{-1/3} \partial_t f\|_0 \leq \|\Lambda^{-1/3} (\partial_t + v \cdot \partial_x) f\|_0 + \|\Lambda^{-1/3} v \cdot \partial_x f\|_0,$$

where  $\Lambda = (1 + |D_t|^2 + |D_x|^2 + |D_v|^2)^{1/2}$ . From Lemma 2.3, we have

$$\|\Lambda^{-1/3} v \cdot \partial_x f\|_0 \leq C_K \|D_x^{2/3} f\|_0 \leq C_K (\|\mathcal{L}f\|_0 + \|f\|_0).$$

The estimation for the term  $\|\Lambda^{-1/3} (\partial_t + v \cdot \partial_x) f\|_0$  can be obtained by direct calculus as in [15].

**Proof of Proposition 2.1.** The Lemma 2.2, Lemma 2.3 and Lemma 2.4 deduce immediately

$$(2.4) \quad \|f\|_{2/3} \leq C_K \{ \|\mathcal{L}f\|_0 + \|f\|_0 \}.$$

Moreover, choose a function  $\psi \in C_0^\infty(\mathbb{R}^{2n+1})$  such that  $\psi|_K \equiv 1$ ,  $\text{Supp } \psi$  is a neighborhood of  $K$ . Then for any  $f \in C_0^\infty(K)$  and any  $r \geq 0$ ,

$$\|f\|_r = \|\psi f\|_r \leq C_K \{ \|\psi \Lambda^{r-2/3} f\|_{2/3} + \|[\Lambda^{r-2/3}, \psi] f\|_{2/3} \}.$$

By virtue of (2.4) and the interpolation inequality (2.2), we have

$$\begin{aligned} \|f\|_r &\leq C_K \{ \|\mathcal{L} \psi \Lambda^{r-2/3} f\|_0 + \|f\|_{r-2/3} \} \\ &\leq C_{\varepsilon, K} \{ \|\mathcal{L} \psi \Lambda^{r-2/3} f\|_0 + \|f\|_0 \} + \varepsilon \|f\|_r. \end{aligned}$$

Taking  $\varepsilon$  small enough, we get

$$\|f\|_r \leq C_K \{ \|\mathcal{L}f\|_{r-2/3} + \|f\|_0 + \|[\mathcal{L}, \psi \Lambda^{r-2/3}] f\|_0 \}.$$

Direct verification gives

$$\begin{aligned} [\mathcal{L}, \psi \Lambda^{r-2/3}] &= [\partial_t + v \cdot \partial_x, \psi \Lambda^{r-2/3}] - \sum_{j=1}^n \{ \|[a, \psi \Lambda^{r-2/3}] \partial_{v_j}^2 \\ &\quad + a[\partial_{v_j}, [\partial_{v_j}, \psi \Lambda^{r-2/3}]] + 2a[\partial_{v_j}, \psi \Lambda^{r-2/3}] \partial_{v_j} \}, \end{aligned}$$

This along with Lemma 2.2 yields

$$\begin{aligned} \|[\mathcal{L}, \psi \Lambda^{r-2/3}] f\|_0 &\leq C_K \{ \|f\|_{r-2/3} + \sum_{j=1}^n \|\partial_{v_j} f\|_{r-2/3} \} \\ &\leq C_K \{ \|\mathcal{L}f\|_{r-2/3} + \|f\|_{r-2/3} \}. \end{aligned}$$

These three estimates gives immediately

$$\|f\|_r \leq C_K \{ \|\mathcal{L}f\|_{r-2/3} + \|f\|_0 + \|f\|_{r-2/3} \}.$$

Applying interpolation inequality (2.2) again and taking  $\varepsilon$  small enough, we prove Proposition 2.1.

We consider now the commutators of the operators  $\mathcal{L}$  with derivation and cut-off function.

**Proposition 2.5.** *Let  $K$  be a compact subset of  $\mathbb{R}^{2n+1}$ . Then for any  $r \geq 0$ , there exist constants  $C_{K,r}, C_{K,r,\varphi}$  such that for any  $f \in C_0^\infty(K)$ ,*

$$\|[\mathcal{L}, D]f\|_r \leq C_{K,r} \{ \|\mathcal{L}f\|_{r+1-2/3} + \|f\|_0 \},$$

and

$$\|[\mathcal{L}, \varphi]f\|_r \leq C_{K,r,\varphi} \{ \|\mathcal{L}f\|_{r-1/3} + \|f\|_0 \},$$

where  $\varphi \in C_b^\infty(\mathbb{R}^{2n+1})$  and we denote by  $D$  the differential operator  $\partial_t, \partial_x$  or  $\partial_v$ .

**Proof.** By using the positivity of coefficient  $a$ , we have

$$\|\Delta_v f\|_r \leq C_K \{ \|\mathcal{L}f\|_r + \|f\|_{r+1} \}.$$

And  $[\mathcal{L}, D] = [\partial_t + v \cdot \partial_x, D] - [a, D]\Delta_v$  deduce

$$\|[\mathcal{L}, D]f\|_r \leq C_K \{ \|f\|_{r+1} + \|\Delta_v f\|_r \}.$$

The above two inequalities along with the subelliptic estimate (2.3) yield the first desired inequality in Proposition 2.5.

To treat  $\|[\mathcal{L}, \varphi]f\|_r$ , the subelliptic estimate (2.3) give

$$\|\nabla_v f\|_r \leq C_K (\|\mathcal{L}f\|_{r-1/3} + \|f\|_0).$$

Now simple verification gives

$$\begin{aligned} \|[\mathcal{L}, \varphi]f\|_r &\leq C_K \left\{ \|f\|_r + \sum_{j=1}^n \|\partial_{v_j} f\|_r \right\} \\ &\leq C_{K,r} \{ \|\mathcal{L}f\|_{r-1/3} + \|f\|_0 \}. \end{aligned}$$

This completes the proof of Proposition 2.5.

We prove now the Gevrey hypoellipticity of  $\mathcal{L}$ . Our starting point is the following result due to M.Durand [9]:

**Proposition 2.6.** *Let  $P$  be a linear differential operator with smooth coefficients in  $\mathbb{R}_y^m$  and  $\varrho, \varsigma$  two fixed positive numbers. If for any  $r \geq 0$ , any compact  $K \subseteq \mathbb{R}^m$  and any  $\varphi \in C^\infty(\mathbb{R}^m)$ , there exist constants  $C_{K,r}$  and  $C_{K,r}(\varphi)$  such that for all  $f \in C_0^\infty(K)$ , the following conditions are fulfilled:*

$$\begin{aligned} (H_1) \quad & \|f\|_r \leq C_{K,r} (\|Pf\|_{r-\varrho} + \|f\|_0), \\ (H_2) \quad & \|[P, D_j]f\|_r \leq C_{K,r} (\|Pf\|_{r+1-\varsigma} + \|f\|_0), \\ (H_3) \quad & \|[P, \varphi]f\|_r \leq C_{K,r}(\varphi) (\|Pf\|_{r-\varsigma} + \|f\|_0), \end{aligned}$$

where

$$D_j = \frac{1}{i} \frac{\partial}{\partial y_j}, j = 1, 2, \dots, m.$$

Then for  $s \geq \max(1/\varsigma, 2/\varrho)$ ,  $P$  is  $G^s(\mathbb{R}^m)$  hypoelliptic, provided the coefficients of  $P$  are in the class of  $G^s(\mathbb{R}^m)$ .

Proposition 2.1 shows that the operator  $\mathcal{L}$  satisfies the conditions  $(H_1)$  with  $\varrho = 2/3$ , Proposition 2.5 assures the conditions  $(H_2)$  and  $(H_3)$  with  $\varsigma = 1/3$ . Then  $\mathcal{L}$  is  $G^s(\mathbb{R}^{2n+1})$  hypoelliptic,  $s \geq 3$ , and we have proved Theorem 1.1.

### 3. GEVREY REGULARITY OF NONLINEAR EQUATIONS

Let  $u \in L_{loc}^\infty(\mathbb{R}^{2n+1})$  be a weak solution of (1.3). Firstly, we will prove  $u \in C^\infty(\mathbb{R}^{2n+1})$ . And we need the following nonlinear composition results (see for example [23]).

**Lemma 3.1.** *Let  $F(t, x, v, w, p) \in C^\infty(\mathbb{R}^{2n+2+n})$  and  $r \geq 0$ . If  $u, \nabla_v u \in L_{loc}^\infty(\mathbb{R}^{2n+1}) \cap H_{loc}^r(\mathbb{R}^{2n+1})$ , then  $F(\cdot, u(\cdot), \nabla_v u(\cdot)) \in H_{loc}^r(\mathbb{R}^{2n+1})$ , and*

$$(3.1) \quad \|\phi_1 F(\cdot, u(\cdot), \nabla_v u(\cdot))\|_r \leq \bar{C} \{ \|\phi_2 u\|_r + \|\phi_2 \nabla_v u\|_r \},$$

where  $\phi_1, \phi_2 \in C_0^\infty(\mathbb{R}^{2n+1})$  and  $\phi_2 = 1$  on the support of  $\phi_1$ , and  $\bar{C}$  is a constant depending only on  $r, \phi_1, \phi_2$ .

**Remark.** If the nonlinear term  $F$  is independent of  $p$  or in the form of

$$\nabla_v(F(t, x, v, u)),$$

Then that  $u \in L_{loc}^\infty(\mathbb{R}^{2n+1}) \cap H_{loc}^r(\mathbb{R}^{2n+1})$  yields  $F(\cdot, u(\cdot), \nabla_v u(\cdot)) \in H_{loc}^r(\mathbb{R}^{2n+1})$ .

**Lemma 3.2.** *Let  $u, \nabla_v u \in H_{loc}^r(\mathbb{R}^{2n+1}), r \geq 0$ . Then we have*

$$(3.2) \quad \|\varphi_1 \nabla_v u\|_r \leq C \|\varphi_2 u\|_r,$$

where  $\varphi_1, \varphi_2 \in C_0^\infty(\mathbb{R}^{2n+1})$  and  $\varphi_2 = 1$  on the support of  $\varphi_1$ , and  $C$  is a constant depending only on  $r, \varphi_1, \varphi_2$ .

In fact, we have

$$\|\varphi_1 \nabla_v u\|_r \leq \|[\nabla_v, \varphi_1]u\|_r + \|\nabla_v \varphi_1 u\|_r.$$

Clearly, the first term on the right is bounded by  $C \|\varphi_2 u\|_r$ . For the second term, combining the second inequality in Lemma 2.2 and (3.1), we get the desired estimate (3.2) at once. This completes the proof of Lemma 3.2.

Now we are ready to prove

**Proposition 3.3.** *Let  $u$  be a weak solution of (1.3) such that  $u, \nabla_v u \in L_{loc}^\infty(\mathbb{R}^{2n+1})$ . Then  $u$  is in  $C^\infty(\mathbb{R}^{2n+1})$ .*

In fact, from the subelliptic estimate (2.3) and the fact  $\mathcal{L}u(\cdot) = F(\cdot, u(\cdot), \nabla_v u(\cdot))$ , it then follows

$$(3.3) \quad \|\psi_1 u\|_{r+2/3} \leq \tilde{C} \{ \|\psi_2 F(\cdot, u(\cdot), \nabla_v u(\cdot))\|_r + \|\psi_2 u\|_0 \},$$

where  $\psi_1, \psi_2 \in C_0^\infty(\mathbb{R}^{2n+1})$  and  $\psi_2 = 1$  on the support of  $\psi_1$ . Combining (3.1), (3.2) and (3.3), we have  $u \in H_{loc}^\infty(\mathbb{R}^{2n+1})$  by standard iteration. This completes the proof of Proposition 3.3.

Now starting from the smooth solution, we prove the Gevrey regularity. It suffices to show the regularity in the open unit ball

$$\Omega = \{(t, x, v) \in \mathbb{R}^{2n+1} : t^2 + |x|^2 + |v|^2 < 1\}.$$

Set

$$\Omega_\rho = \{(t, x, v) \in \Omega : (t^2 + |x|^2 + |v|^2)^{1/2} < 1 - \rho\}, \quad 0 < \rho < 1.$$

Let  $U$  be an open subset of  $\mathbb{R}^{2n+1}$ . Denote by  $H^r(U)$  the space consisting of the functions which are defined in  $U$  and can be extended to  $H^r(\mathbb{R}^{2n+1})$ . Define

$$\|u\|_{H^r(U)} = \inf \{ \|\tilde{u}\|_{H^s(\mathbb{R}^{2n+1})} : \tilde{u} \in H^s(\mathbb{R}^{2n+1}), \tilde{u}|_U = u \}.$$

We denote  $\|u\|_{r,U} = \|u\|_{H^r(U)}$ , and

$$\|D^j u\|_r = \sum_{|\beta|=j} \|D^\beta u\|_r.$$

In order to treat the nonlinear term  $F$  on the right hand of (1.3), we need the following two lemmas. The first one (see [23] for example) concerns weak solution in some algebra, and the second is an analogue of Lemma 1 in [10]. In the sequel  $C_j > 1$  will be used to denote suitable constants depending only on  $n$  or the function  $F$ .

**Lemma 3.4.** *Let  $r > (2n+1)/2$  and  $u_1, u_2 \in H^r(\mathbb{R}^{2n+1})$ , Then  $u_1 u_2 \in H^r(\mathbb{R}^{2n+1})$ , moreover*

$$(3.4) \quad \|u_1 u_2\|_r \leq \tilde{C} \|u_1\|_r \|u_2\|_r,$$

where  $\tilde{C}$  is a constant depending only on  $n, r$ .

**Lemma 3.5.** *Let  $M_j$  be a sequence of positive numbers and for some  $B_0 > 0$ , the  $M_j$  satisfy the monotonicity conditions*

$$(3.5) \quad \frac{j!}{i!(j-i)!} M_i M_{j-i} \leq B_0 M_j, \quad (i = 1, 2, \dots, j; j = 1, 2, \dots).$$

Suppose  $F(t, x, v, u, p)$  satisfy

$$(3.6) \quad \left\| \left( D_{t,x,v}^j D_u^l D_p^m F \right) (\cdot, u(\cdot), \nabla_v u(\cdot)) \right\|_{r+n+1, \Omega} \leq C_1^{j+l+m} M_{j-2} M_{m+l-2}, \quad j, m+l \geq 2,$$

where  $r$  is a real number satisfying  $r+n+1 > (2n+1)/2$ . Then there exist two constants  $C_2, C_3$  such that for any  $H_0, H_1$  satisfying  $H_0, H_1 \geq 1$  and  $H_1 \geq C_2 H_0$ , if  $u(t, x, v)$  satisfy the following conditions

$$(3.7) \quad \|D^j u\|_{r+n+1, \Omega_{\bar{\rho}}} \leq H_0, \quad 0 \leq j \leq 1,$$

$$(3.8) \quad \|D^j u\|_{r+n+1, \Omega_{\bar{\rho}}} \leq H_0 H_1^{j-2} M_{j-2}, \quad 2 \leq j \leq N,$$

$$(3.9) \quad \|D_v D^j u\|_{r+n+1, \Omega_{\bar{\rho}}} \leq H_0 H_1^{j-2} M_{j-2}, \quad 2 \leq j \leq N.$$

Then for all  $\alpha$  with  $|\alpha| = N$ ,

$$(3.10) \quad \left\| \psi_N D^\alpha [F(\cdot, u(\cdot), \nabla_v u(\cdot))] \right\|_{r+n+1} \leq C_3 H_0 H_1^{N-2} M_{N-2},$$

where  $\psi_N \in C_0^\infty(\Omega_{\bar{\rho}})$  is an arbitrary function.

**Proof.** Denote  $p = (p_1, p_2, \dots, p_n) = \nabla_v u$  and  $k = (k_1, k_2, \dots, k_n)$ . From Faa di Bruno' formula,  $\psi_N D^\alpha [F(\cdot, u(\cdot), \nabla_v u(\cdot))]$  is the linear combination of terms of the form

$$(3.11) \quad \frac{\psi_N \partial^{|\tilde{\alpha}|+l+|k|} F}{\partial_{t,x,v}^{\tilde{\alpha}} \partial u^l \partial p_1^{k_1} \dots \partial p_n^{k_n}} \prod_{j=1}^l D^{\gamma_j} u \cdot \prod_{i=1}^n \prod_{j_i=1}^{k_i} D^{\beta_{j_i}} (\partial_{v_i} u),$$

where  $|\tilde{\alpha}| + l + |k| \leq |\alpha|$  and

$$\sum_{j=1}^l \gamma_j + \sum_{i=1}^n \sum_{j_i}^{k_i} \beta_{j_i} = \alpha - \tilde{\alpha},$$

and if  $\gamma_i$  or  $\beta_{j_i}$  equals to 0, we just mean  $D^{\gamma_i} u$  or  $D^{\beta_{j_i}} u$  doesn't appear in (3.11). Choose a function  $\tilde{\psi} \in C_0^\infty(\Omega_{\bar{\rho}})$  such that  $\tilde{\psi} = 1$  on Supp  $\psi_N$ . Note that  $n+1+r > (2n+1)/2$ , and hence applying Lemma 3.4, we have

$$(3.12) \quad \begin{aligned} & \left\| \frac{\psi_N \partial^{|\tilde{\alpha}|+l+|k|} F}{\partial_{t,x,v}^{\tilde{\alpha}} \partial u^l \partial p_1^{k_1} \dots \partial p_n^{k_n}} \prod_{j=1}^l D^{\gamma_j} u \cdot \prod_{i=1}^n \prod_{j_i=1}^{k_i} D^{\beta_{j_i}} (\partial_{v_i} u) \right\|_{r+n+1} \\ &= \left\| \frac{\psi_N \partial^{|\tilde{\alpha}|+l+|k|} F}{\partial_{t,x,v}^{\tilde{\alpha}} \partial u^l \partial p_1^{k_1} \dots \partial p_n^{k_n}} \prod_{j=1}^l \tilde{\psi} D^{\gamma_j} u \cdot \prod_{i=1}^n \prod_{j_i=1}^{k_i} \tilde{\psi} \partial_{v_i} D^{\beta_{j_i}} u \right\|_{r+n+1} \\ &\leq \tilde{C} \left\| \psi_N (\partial^{|\tilde{\alpha}|+l+|k|} F) \right\|_{r+n+1} \cdot \prod_{j=1}^l \left\| \tilde{\psi} D^{\gamma_j} u \right\|_{r+n+1} \times \prod_{i=1}^n \prod_{j_i=1}^{k_i} \left\| \tilde{\psi} \partial_{v_i} D^{\beta_{j_i}} u \right\|_{r+n+1} \\ &\leq C_0 \left\| (\partial^{|\tilde{\alpha}|+l+|k|} F) \right\|_{r+n+1, \Omega} \cdot \prod_{j=1}^l \|D^{\gamma_j} u\|_{r+n+1, \Omega_{\bar{\rho}}} \times \prod_{i=1}^n \prod_{j_i=1}^{k_i} \|\partial_{v_i} D^{\beta_{j_i}} u\|_{r+n+1, \Omega_{\bar{\rho}}}. \end{aligned}$$

In virtue of (3.7)-(3.9) and (3.12), the situation is entirely similar to [10]. The only difference is that we replace the Hölder norm  $|u|_j$  by  $\|D^j u\|_{r+n+1, \Omega_{\bar{\rho}}}$  and  $\|D_v D^j u\|_{r+n+1, \Omega_{\bar{\rho}}}$ . Then the same argument as the proof of Lemma 1 in [10] yields (3.10). This completes the proof of Lemma 3.5.



**Proposition 3.6.** *Let  $s \geq 3$ . Suppose  $u \in C^\infty(\bar{\Omega})$  is a solution of (1.3), and  $a(t, x, v) \in G^s(\mathbb{R}^{2n+1})$ ,  $F(t, x, v, w, p) \in G^s(\mathbb{R}^{2n+2+n})$  and  $a \geq c_0 > 0$ . Then there exists a constant  $A$  such that for any  $r \in [0, 1]$  and any  $N \in \mathbb{N}$ ,  $N \geq 3$ ,*

$$(E)_{r,N} \quad \begin{aligned} & \|D^\alpha u\|_{r+n+1, \Omega_\rho} + \|D_v D^\alpha u\|_{r-1/3+n+1, \Omega_\rho} \\ & \leq \frac{A^{|\alpha|-1}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{sr}, \quad \forall |\alpha| = N, \quad \forall 0 < \rho < 1. \end{aligned}$$

From  $(E)_{r,N}$ , we have immediately

**Proposition 3.7.** *Under the same assumption as Proposition 3.6, we have  $u \in G^s(\Omega)$ .*

In fact, for any compact subset  $K$  of  $\Omega$ , we have  $K \subset \Omega_{\rho_0}$  for some  $\rho_0$ ,  $0 < \rho_0 < 1$ . For any  $\alpha$ ,  $|\alpha| \geq 3$ , letting  $r = 0$  in  $(E)_{r,N}$ , we have

$$\|D^\alpha u\|_{L^2(K)} \leq \|D^\alpha u\|_{n+1, \Omega_{\rho_0}} \leq \frac{A^{|\alpha|-1}}{\rho_0^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s \leq \left(\frac{A}{\rho_0^s}\right)^{|\alpha|+1} (|\alpha|!)^s.$$

This completes the proof of Proposition 3.7.

**Proof of Proposition 3.6.** We use induction on  $N$ . Assuming  $(E)_{r, N-1}$  holds for any  $r$  with  $0 \leq r \leq 1$ , and we will show  $(E)_{r, N}$  still holds for any  $r \in [0, 1]$ . For any  $\alpha$ ,  $|\alpha| = N$ , and for any  $\rho$ ,  $0 < \rho < 1$ , choose a function  $\varphi_{\rho, N} \in C_0^\infty(\Omega_{\frac{N-1}{N}\rho})$  such that  $\varphi_{\rho, N} = 1$  in  $\Omega_\rho$ . It is easy to see

$$\sup |D^\gamma \varphi_{\rho, N}| \leq C_\gamma (\rho/N)^{-|\gamma|} \leq C_\gamma (N/\rho)^{|\gamma|}, \quad \forall \gamma.$$

And we will proceed to prove the truth of  $(E)_{r, N}$  by the following lemmas.

**Lemma 3.8.** *For  $r = 0$ , we have*

$$\|D^\alpha u\|_{n+1, \Omega_\rho} + \|D_v D^\alpha u\|_{-1/3+n+1, \Omega_\rho} \leq \frac{C_7 A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s, \quad \forall 0 < \rho < 1.$$

**Proof .** Write  $|\alpha| = |\beta| + 1$ , then  $|\beta| = N - 1$ . Denote  $\frac{N-1}{N}\rho$  by  $\tilde{\rho}$ . In the sequel we will use the following fact frequently

$$\frac{1}{\rho^{sk}} \leq \frac{1}{\tilde{\rho}^{sk}} = \frac{1}{\rho^{sk}} \times \left(\frac{N}{N-1}\right)^{sk} \leq \frac{C_4}{\rho^{sk}}, \quad k = 1, 2, \dots, N-3.$$

Note that  $\varphi_{\rho, N} = 1$  in  $\Omega_\rho$  and hence

$$\begin{aligned} \|D^\alpha u\|_{n+1, \Omega_\rho} & \leq \|\varphi_{\rho, N} D^\alpha u\|_{n+1} \leq \|\varphi_{\rho, N} D^\beta u\|_{1+n+1} + \|(D\varphi_{\rho, N}) D^\beta u\|_{n+1} \\ & \leq C_5 \{ \|D^\beta u\|_{1+n+1, \Omega_{\tilde{\rho}}} + (N/\rho) \|D^\beta u\|_{n+1, \Omega_{\tilde{\rho}}} \}. \end{aligned}$$

Since  $(E)_{r, N-1}$  holds by assumption for any  $r$  with  $0 \leq r \leq 1$ , we have immediately

$$\begin{aligned} & \|D^\beta u\|_{1+n+1, \Omega_{\tilde{\rho}}} + (N/\rho) \|D^\beta u\|_{n+1, \Omega_{\tilde{\rho}}} \\ & \leq \frac{A^{|\beta|-1}}{\tilde{\rho}^{s(|\beta|-3)}} ( (|\beta|-3)! )^s (N/\tilde{\rho})^s + (N/\rho) \frac{A^{|\beta|-1}}{\tilde{\rho}^{s(|\beta|-3)}} ( (|\beta|-3)! )^s \\ & \leq \frac{2A^{|\alpha|-2}}{\tilde{\rho}^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/(N-3))^s \\ & \leq \frac{C_6 A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s. \end{aligned}$$

Thus

$$(3.13) \quad \|D^\alpha u\|_{n+1, \Omega_\rho} \leq \frac{C_5 C_6 A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s.$$

The same arguments as above shows that

$$\|D_v D^\alpha u\|_{-1/3+n+1, \Omega_\rho} \leq \frac{C_5 C_6 A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s.$$

This along with (3.13) yields the conclusion.

**Lemma 3.9.** *For any  $0 \leq r \leq 1/3$ , we have*

$$\|D^\alpha u\|_{r+n+1, \Omega_\rho} + \|D_v D^\alpha u\|_{r-1/3+n+1, \Omega_\rho} \leq \frac{C_{35} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{rs}, \quad \forall 0 < \rho < 1.$$

**Proof .** We firstly prove the conclusion is true for  $r = 1/3$ , i.e., to show

$$\|D^\alpha u\|_{1/3+n+1, \Omega_\rho} + \|D_v D^\alpha u\|_{n+1, \Omega_\rho} \leq \frac{C_{35} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}, \quad \forall 0 < \rho < 1.$$

And we will proceed in the following four steps.

**Step 1.** Claim

$$(3.14) \quad \|[\mathcal{L}, \varphi_{\rho, N} D^\alpha] u\|_{-1/3+n+1} \leq \frac{C_{19} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}.$$

In fact, write  $\mathcal{L} = X_0 - a\Delta_v$  with  $X_0 = \partial_t + v \cdot \partial_x$ . Then direct verification deduces

$$\begin{aligned} \|[\mathcal{L}, \varphi_{\rho, N} D^\alpha] u\|_{-1/3+n+1} &\leq \| [X_0, \varphi_{\rho, N} D^\alpha] u \|_{-1/3+n+1} + \| a[\Delta_v, \varphi_{\rho, N} D^\alpha] u \|_{-1/3+n+1} \\ &\quad + \| \varphi_{\rho, N} [a, D^\alpha] \Delta_v u \|_{-1/3+n+1} \\ &=: (I) + (II) + (III). \end{aligned}$$

Denote  $[X_0, D^\alpha]$  by  $D^{\alpha_0}$ . Then  $|\alpha_0| \leq |\alpha|$  and

$$\begin{aligned} (I) &\leq \| [X_0, \varphi_{\rho, N} D^\alpha] u \|_{n+1} + \| \varphi_{\rho, N} D^{\alpha_0} u \|_{n+1} \\ &\leq C_8 \{ (N/\rho) \| D^\alpha u \|_{n+1, \Omega_{\tilde{\rho}}} + \| D^{\alpha_0} u \|_{n+1, \Omega_{\tilde{\rho}}} \}. \end{aligned}$$

Note that  $s \geq 3$ . Using Lemma 3.8, we have

$$(3.15) \quad (I) \leq C_8 (N/\rho + 1) \frac{C_7 A^{|\alpha|-2}}{\tilde{\rho}^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s \leq \frac{C_9 A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}.$$

Next we will estimate (II). It is easy to see that

$$(3.16) \quad \begin{aligned} \|[\Delta_v, \varphi_{\rho, N} D^\alpha] u\|_{-1/3+n+1} &\leq 2 \| [D_v, \varphi_{\rho, N}] D_v D^\alpha u \|_{-1/3+n+1} \\ &\quad + \| [D_v, [D_v, \varphi_{\rho, N}]] D^\alpha u \|_{-1/3+n+1}. \end{aligned}$$

We firstly treat the first term on the right hand. Using Lemma 3.8 again, we have

$$(3.17) \quad \begin{aligned} \| [D_v, \varphi_{\rho, N}] D_v D^\alpha u \|_{-1/3+n+1} &\leq (N/\rho) \| D_v D^\alpha u \|_{-1/3+n+1, \Omega_{\tilde{\rho}}} \\ &\leq (N/\rho) \frac{C_7 A^{|\alpha|-2}}{\tilde{\rho}^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s \\ &\leq \frac{C_{10} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}. \end{aligned}$$

Next we treat  $\|[D_v, [D_v, \varphi_{\rho, N}] D^\alpha u]\|_{-1/3+n+1}$ , and we compute

$$\begin{aligned}
& \|[D_v, [D_v, \varphi_{\rho, N}] D^\alpha u]\|_{-1/3+n+1} \\
& \leq \|(D^2 \varphi_{\rho, N}) D^\beta u\|_{2/3+n+1} + \|(D^3 \varphi_{\rho, N}) D^\beta u\|_{-1/3+n+1} \\
& \leq C_{11} \{ (N/\rho)^2 \|D^\beta u\|_{2/3+n+1, \Omega_{\tilde{\rho}}} + (N/\rho)^3 \|D^\beta u\|_{n+1, \Omega_{\tilde{\rho}}} \} \\
& \leq C_{11} \{ (N/\rho)^2 \frac{A^{|\beta|-1}}{\tilde{\rho}^{s(|\beta|-3)}} ((|\beta|-3)!)^s (N/\tilde{\rho})^{2s/3} \\
& \quad + (N/\rho)^3 \frac{A^{|\beta|-1}}{\tilde{\rho}^{s(|\beta|-3)}} ((|\beta|-3)!)^s \} \\
& \leq C_{11} \{ (N/\rho)^2 (N/\tilde{\rho})^{-s/3} \frac{A^{|\alpha|-2}}{\tilde{\rho}^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s \\
& \quad + (N/\rho)^3 (N/\tilde{\rho})^{-s} \frac{A^{|\alpha|-2}}{\tilde{\rho}^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s \} \\
& \leq \frac{C_{12} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}.
\end{aligned}$$

This along with (3.16) and (3.17) shows at once

$$(3.18) \quad (II) \leq \frac{C_{13} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}.$$

It remains to treat (III), and using Leibniz' formula,

$$\begin{aligned}
(III) & \leq \sum_{0 < |\gamma| \leq |\alpha|} \binom{\alpha}{\gamma} \|\varphi_{\rho, N} (D^\gamma a) \Delta_v D^{\alpha-\gamma} u\|_{-1/3+n+1} \\
& \leq \sum_{0 < |\gamma| \leq |\alpha|} \binom{\alpha}{\gamma} \|D^\gamma a\|_{n+1, \Omega} \cdot \|\varphi_{\rho, N} \Delta_v D^{\alpha-\gamma} u\|_{-1/3+n+1}.
\end{aligned}$$

Since  $a \in G^s(\mathbb{R}^{2n+1})$ , then

$$\|D^\gamma a\|_{n+1, \Omega} \leq C_{14}^{|\gamma|-2} ((|\gamma|-3)!)^s, \quad |\gamma| \geq 3,$$

and

$$\|D^\gamma a\|_{n+1, \Omega} \leq C_{14}, \quad |\gamma| = 1, 2.$$

Moreover, note  $|\alpha| - |\gamma| + 1 \leq N$ , and hence applying Lemma 3.8, we have for any  $\gamma$ ,  $|\gamma| \leq |\alpha| - 2$ ,

$$\begin{aligned}
\|\varphi_{\rho, N} \Delta_v D^{\alpha-\gamma} u\|_{-1/3+n+1} & \leq \|D_v D^{\alpha-\gamma+1} u\|_{-1/3+n+1, \Omega_{\tilde{\rho}}} \\
& \leq \frac{C_7 A^{|\alpha|-|\gamma|+1-2}}{\tilde{\rho}^{s(|\alpha|-|\gamma|-2)}} ((|\alpha|-|\gamma|-2)!)^s \\
& \leq \frac{C_{15} A^{|\alpha|-|\gamma|+1-2}}{\rho^{s(|\alpha|-|\gamma|-2)}} ((|\alpha|-|\gamma|-2)!)^s.
\end{aligned}$$

Consequently, we compute

$$\begin{aligned}
& \sum_{2 \leq |\gamma| \leq |\alpha| - 2} \binom{\alpha}{\gamma} \|D^\gamma a\|_{n+1, \Omega} \cdot \|\varphi_{\rho, N} \Delta_v D^{\alpha - \gamma} u\|_{-1/3 + n + 1} \\
& \leq \sum_{2 \leq |\gamma| \leq |\alpha| - 2} \binom{\alpha}{\gamma} C_{14}^{|\gamma| - 2} ((|\gamma| - 2)!)^s \frac{C_{15} A^{|\alpha| - |\gamma| + 1 - 2}}{\rho^{s(|\alpha| - |\gamma| - 2)}} ((|\alpha| - |\gamma| - 2)!)^s \\
& \leq \frac{C_{15} A^{|\alpha| - 2}}{\rho^{s(|\alpha| - 3)}} \sum_{2 \leq |\gamma| \leq |\alpha| - 2} \left(\frac{C_{14}}{A}\right)^{|\gamma| - 1} |\alpha|! ((|\gamma| - 2)!)^{s-1} ((|\alpha| - |\gamma| - 2)!)^{s-1} \\
& \leq \frac{C_{15} A^{|\alpha| - 2}}{\rho^{s(|\alpha| - 3)}} ((|\alpha| - 3)!)^s \sum_{2 \leq |\gamma| \leq |\alpha| - 2} \left(\frac{C_{14}}{A}\right)^{|\gamma| - 1} |\alpha| \frac{(|\alpha| - 1)(|\alpha| - 2)}{(|\alpha| - 3)^{s-1}} \\
& \leq \frac{C_{16} A^{|\alpha| - 2}}{\rho^{s(|\alpha| - 3)}} ((|\alpha| - 3)!)^s (N/\rho)^{s/3} \sum_{2 \leq |\gamma| \leq |\alpha| - 2} \left(\frac{C_{14}}{A}\right)^{|\gamma| - 1}.
\end{aligned}$$

Taking  $A$  large enough such that  $\sum_{2 \leq |\gamma| \leq |\alpha| - 2} \left(\frac{C_{14}}{A}\right)^{|\gamma| - 1} \leq 1$ , then we get

$$\sum_{2 \leq |\gamma| \leq |\alpha| - 2} \binom{\alpha}{\gamma} \|D^\gamma a\|_{n+1, \Omega} \cdot \|\varphi_{\rho, N} \Delta_v D^{\alpha - \gamma} u\|_{-1/3 + n + 1} \leq \frac{C_{16} A^{|\alpha| - 2}}{\rho^{s(|\alpha| - 3)}} ((|\alpha| - 3)!)^s (N/\rho)^{s/3}.$$

For  $|\gamma| = 1$ ,  $|\alpha| - 1$  or  $|\alpha|$ , we can compute directly

$$\binom{\alpha}{\gamma} \|D^\gamma a\|_{n+1, \Omega} \cdot \|\varphi_{\rho, N} \Delta_v D^{\alpha - \gamma} u\|_{-1/3 + n + 1} \leq \frac{C_{17} A^{|\alpha| - 2}}{\rho^{s(|\alpha| - 3)}} ((|\alpha| - 3)!)^s (N/\rho)^{s/3}.$$

Combination of the above two inequalities give that

$$(III) \leq \frac{C_{18} A^{|\alpha| - 2}}{\rho^{s(|\alpha| - 3)}} ((|\alpha| - 3)!)^s (N/\rho)^{s/3}.$$

This along with (3.15) and (3.18) yields the conclusion (3.14).

**Step 2.** Claim

$$(3.19) \quad \|\varphi_{\rho, N} D^\alpha [F(\cdot, u(\cdot), \nabla_v u(\cdot))]\|_{-1/3 + n + 1} \leq \frac{C_{21} A^{|\alpha| - 2}}{\rho^{s(|\alpha| - 3)}} ((|\alpha| - 3)!)^s (N/\rho)^{s/3}.$$

Firstly, we will prove  $F$  and  $u$  satisfy the conditions (3.7)-(3.9) for some  $M_j$ . By Lemma 3.8, we have

$$(3.20) \quad \|D^j u\|_{-1/3 + n + 1, \Omega_{\tilde{\rho}}} \leq \|D^j u\|_{n+1, \Omega_{\tilde{\rho}}} \leq \frac{C_7 A^{j-2}}{\tilde{\rho}^{s(j-3)}} ((j-3)!)^s, \quad 3 \leq j \leq N,$$

$$(3.21) \quad \|D_v D^j u\|_{-1/3 + n + 1, \Omega_{\tilde{\rho}}} \leq \frac{C_7 A^{j-2}}{\tilde{\rho}^{s(j-3)}} ((j-3)!)^s, \quad 3 \leq j \leq N,$$

and

$$(3.22) \quad \|D^j u\|_{-1/3 + n + 1, \Omega_{\tilde{\rho}}} \leq C_7, \quad 0 \leq j \leq 2.$$

Since  $F \in G^s(\mathbb{R}^{2n+1} \times \mathbb{R})$ , then

$$(3.23) \quad \|(D_{t,x,v}^k \partial_u^l D_p^m F)(\cdot, u(\cdot), \nabla_v u(\cdot))\|_{-1/3 + n + 1, \Omega} \leq C_{20}^{k+l} ((k-3)!)^s ((l-3)!)^s, \quad k, m+l \geq 3.$$

Define  $M_j, H_0, H_1$  by setting

$$H_0 = C_7, \quad H_1 = A, \quad M_0 = C_7, \quad M_j = \frac{((j-1)!)^s}{\tilde{\rho}^{s(j-1)}}, \quad j \geq 1.$$

We can choose  $A$  large enough such that  $H_1 = A \geq C_2 H_0$ . Then (3.20)-(3.23) can be rewritten

$$(3.24) \quad \|D^j u\|_{-1/3+n+1, \Omega_{\tilde{\rho}}} \leq H_0, \quad 0 \leq j \leq 1,$$

$$(3.25) \quad \|D^j u\|_{-1/3+n+1, \Omega_{\tilde{\rho}}} \leq H_0 H_1^{j-2} M_{j-2}, \quad 2 \leq j \leq |\alpha| = N,$$

$$(3.26) \quad \|D_v D^j u\|_{-1/3+n+1, \Omega_{\tilde{\rho}}} \leq H_0 H_1^{j-2} M_{j-2}, \quad 2 \leq j \leq |\alpha| = N,$$

$$(3.27) \quad \|(D_{t,x,v}^k \partial_u^l D_p^m F)\|_{-1/3+n+1, \Omega} \leq C_{20}^{k+l} M_{k-2} M_{m+l-2}, \quad k, m+l \geq 2.$$

For each  $j$ , note that  $s \geq 3$  and hence

$$(3.28) \quad \begin{aligned} \frac{j!}{i!(j-i)!} M_i M_{j-i} &= \frac{j!}{i!(j-i)!} ((i-1)!)^{s-1} ((j-i-1)!)^{s-1} \tilde{\rho}^{-s(i-1)} \tilde{\rho}^{-s(j-i-1)} \\ &\leq (j!)((j-2)!)^{s-1} \tilde{\rho}^{-s(j-1)} \\ &\leq \frac{j}{(j-1)^{s-1}} (j-1)! ((j-1)!)^{s-1} \tilde{\rho}^{-s(j-1)} \\ &\leq M_j. \end{aligned}$$

Thus  $M_j$  satisfy the monotonicity condition (3.5). In virtue of (3.24)-(3.28), using Lemma 3.5, we have

$$\begin{aligned} \|\varphi_{\rho, N} D^\alpha [F(\cdot, u(\cdot))]\|_{-1/3+n+1} &\leq C_3 H_0 H_1^{|\alpha|-2} M_{|\alpha|-2} \\ &\leq \frac{C_3 C_7 A^{|\alpha|-2}}{\tilde{\rho}^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s \\ &\leq \frac{C_{21} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}. \end{aligned}$$

This completes the proof of conclusion (3.19).

**Step 3.** Claim

$$(3.29) \quad \|\mathcal{L} \varphi_{\rho, N} D^\alpha u\|_{-1/3+n+1} \leq \frac{C_{23} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ((|\alpha|-3)!)^s (N/\rho)^{s/3}.$$

In fact,

$$\begin{aligned} \|\mathcal{L} \varphi_{\rho, N} D^\alpha u\|_{-1/3+n+1} &\leq C_{22} \{ \|\mathcal{L}, \varphi_{\rho, N} D^\alpha u\|_{-1/3+n+1} + \|\varphi_{\rho, N} D^\alpha \mathcal{L} u\|_{-1/3+n+1} \} \\ &= C_{22} \{ \|\mathcal{L}, \varphi_{\rho, N} D^\alpha u\|_{-1/3+n+1} \\ &\quad + \|\varphi_{\rho, N} D^\alpha [F(\cdot, u(\cdot), \nabla_v u(\cdot))]\|_{-1/3+n+1} \}. \end{aligned}$$

This along with (3.14), (3.19) in step 1 and step 2 yields immediately the conclusion (3.29).

**Step 4.** Claim

$$(3.30) \quad \|\varphi_{\rho,N}D^\alpha u\|_{1/3+n+1} + \|\varphi_{\rho,N}D_v D^\alpha u\|_{1/3-1/3+n+1} \leq \frac{C_{31}A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{s/3}.$$

In fact, applying the subelliptic estimate (2.3), we obtain

$$\|\varphi_{\rho,N}D^\alpha u\|_{1/3+n+1} \leq C_{24} \{ \|\mathcal{L}\varphi_{\rho,N}D^\alpha u\|_{-1/3+n+1} + \|\varphi_{\rho,N}D^\alpha u\|_{n+1} \}.$$

Combining Lemma 3.8 and (3.29) in Step 3, we have

$$(3.31) \quad \|\varphi_{\rho,N}D^\alpha u\|_{1/3+n+1} \leq \frac{C_{25}A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{s/3}.$$

Now it remains to treat  $\|\varphi_{\rho,N}D_v D^\alpha u\|_{1/3-1/3+n+1}$ , and

$$\|\varphi_{\rho,N}D_v D^\alpha u\|_{1/3-1/3+n+1} \leq \|D_v \varphi_{\rho,N}D^\alpha u\|_{n+1} + \|[D_v, \varphi_{\rho,N}]D^\alpha u\|_{n+1}.$$

Firstly, we treat the first term on the right. By direct calculation, it follows that

$$\begin{aligned} & \|D_v \varphi_{\rho,N}D^\alpha u\|_{n+1}^2 \\ &= \operatorname{Re}(\mathcal{L}\varphi_{\rho,N}D^\alpha u, a^{-1}\Lambda^{2n+2}\varphi_{\rho,N}D^\alpha u) - \operatorname{Re}(X_0\varphi_{\rho,N}D^\alpha u, a^{-1}\Lambda^{2n+2}\varphi_{\varepsilon,k\varepsilon}D^\alpha u) \\ &= \operatorname{Re}(\mathcal{L}\varphi_{\rho,N}D^\alpha u, a^{-1}\Lambda^{2n+2}\varphi_{\rho,N}D^\alpha u) - \frac{1}{2}(\varphi_{\rho,N}D^\alpha u, [a^{-1}\Lambda^{2n+2}, X_0]\varphi_{\rho,N}D^\alpha u) \\ &\quad - \frac{1}{2}(\varphi_{\rho,N}D^\alpha u, [\Lambda^{2n+2}, a^{-1}]X_0\varphi_{\rho,N}D^\alpha u) \\ &\leq C_{26} \{ \|\mathcal{L}\varphi_{\rho,N}D^\alpha u\|_{-1/3+n+1}^2 + \|\varphi_{\rho,N}D^\alpha u\|_{1/3+n+1}^2 \}. \end{aligned}$$

This along with (3.29) and (3.31) shows at once

$$\|D_v \varphi_{\rho,N}D^\alpha u\|_{r-1/3+n+1} \leq \frac{C_{27}A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{s/3}.$$

Moreover Lemma 3.8 yields

$$\begin{aligned} \|[D_v, \varphi_{\rho,N}]D^\alpha u\|_{n+1} &\leq C_{28}(N/\rho)\|D^\alpha u\|_{n+1,\Omega_{\tilde{\rho}}} \\ &\leq \frac{C_{28}C_7A^{|\alpha|-2}}{\tilde{\rho}^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{s/3} \\ &\leq \frac{C_{29}A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{s/3}. \end{aligned}$$

From the above two inequalities, we have

$$\|\varphi_{\rho,N}D_v D^\alpha u\|_{1/3+n+1} \leq \frac{C_{30}A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{s/3}.$$

This completes the proof of Step 4.

It's clear for any  $\rho$ ,  $0 < \rho < 1$ ,

$$\|D^\alpha u\|_{1/3+n+1,\Omega_\rho} + \|D_v D^\alpha u\|_{1/3-1/3+n+1,\Omega_\rho} \leq \|\varphi_{\rho,N}D^\alpha u\|_{1/3+n+1} + \|\varphi_{\rho,N}D_v D^\alpha u\|_{1/3-1/3+n+1}.$$

Thus from Step 4, it follows that the conclusion in Lemma 3.9 is true for  $r = 1/3$ .

Moreover for any  $0 < r < 1/3$ , using the interpolation inequality (2.2), we have

$$\begin{aligned} \|D^\alpha u\|_{r+n+1,\Omega_\rho} &\leq \|\varphi_{\rho,N}D^\alpha u\|_{r+n+1} \\ &\leq \varepsilon\|\varphi_{\rho,N}D^\alpha u\|_{1/3+n+1} + \varepsilon^{-r/(1/3-r)}\|\varphi_{\rho,N}D^\alpha u\|_{n+1} \\ &\leq \varepsilon\frac{C_{31}A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{s/3} + \varepsilon^{-r/(1/3-r)}\frac{C_{32}A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s, \end{aligned}$$

Taking  $\varepsilon = (N/\rho)^{s(r-1/3)}$ , then

$$\|D^\alpha u\|_{r+n+1, \Omega_\rho} \leq \frac{C_{33} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{rs}.$$

Similarly,

$$\|D_v D^\alpha u\|_{r-1/3+n+1, \Omega_\rho} \leq \frac{C_{34} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{rs}.$$

This completes the proof of Lemma 3.9.

Inductively, we have the following

**Lemma 3.10.** *For any  $r$  with  $1/3 \leq r \leq 2/3$ ,*

$$(3.32) \quad \|D^\alpha u\|_{r+n+1, \Omega_\rho} + \|D_v D^\alpha u\|_{r-1/3+n+1, \Omega_\rho} \leq \frac{C_{38} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^{sr}, \quad \forall 0 < \rho < 1.$$

Moreover, the above inequality still holds for any  $r$  with  $2/3 \leq r \leq 1$ .

**Proof.** Repeating the proof of Lemma 3.9, we have the truth of (3.32) for  $1/3 \leq r \leq 2/3$ . The case  $2/3 \leq r \leq 1$  is a little different. The conclusion in Step 1 in the above proof still holds for  $r = 1$ , and corresponding to Step 2, we have to make some modification to prove

$$\|\varphi_{\rho, N} D^\alpha [F(\cdot, u(\cdot), \nabla_v u(\cdot))]\|_{1/3+n+1} \leq \frac{C_{36} A^{|\alpha|-2}}{\rho^{s(|\alpha|-3)}} ( (|\alpha|-3)! )^s (N/\rho)^s.$$

From the truth of (3.32) for  $1/3 \leq r \leq 2/3$ , it follows

$$\|D^j u\|_{1/3+n+1, \Omega_{\tilde{\rho}}} \leq \frac{C_{37} A^{j-2}}{\tilde{\rho}^{s(j-3)}} ( (j-3)! )^s (j/\tilde{\rho})^{s/3}, \quad 3 \leq j \leq N,$$

$$\|D_v D^j u\|_{1/3+n+1, \Omega_{\tilde{\rho}}} \leq \|D_v D^j u\|_{2/3-1/3+n+1, \Omega_{\tilde{\rho}}} \leq \frac{C_{37} A^{j-2}}{\tilde{\rho}^{s(j-3)}} ( (j-3)! )^s (j/\tilde{\rho})^{2s/3}, \quad 3 \leq j \leq N,$$

and

$$\|D^j u\|_{1/3+n+1, \Omega_{\tilde{\rho}}} \leq C_{37}, \quad 0 \leq j \leq 2,$$

Hence we need define a new sequence  $\bar{M}_j$  by setting

$$\bar{M}_0 = C_{37}, \quad \bar{M}_j = \frac{((j-1)!)^s}{\tilde{\rho}^{s(j-1)}} ((j+2)/\tilde{\rho})^{2s/3}, \quad j \geq 1.$$

For each  $j$ , note that  $s \geq 3$  and hence direct computation deduces that for  $0 < i < j$ ,

$$\begin{aligned} \frac{j!}{i!(j-i)!} \bar{M}_i \bar{M}_{j-i} &= \frac{j!}{i!(j-i)} ((i-1)!)^{s-1} ((j-i-1)!)^{s-1} \\ &\quad \times (i+2)^{2s/3} (j-i+2)^{2s/3} \tilde{\rho}^{-s(j-2)} \tilde{\rho}^{-4s/3} \\ &\leq 4(j!) ((j-2)!)^{s-1} (j+2)^{2s/3-1} (j+1)^{2s/3-1} \tilde{\rho}^{-s(j-1)} \tilde{\rho}^{-2s/3} \tilde{\rho}^{-2s/3} \\ &\leq \frac{4j(j+1)^{2s/3-1}}{(j-1)^{s-1}} (j-1)! ((j-1)!)^{s-1} \tilde{\rho}^{-s(j-1)} ((j+2)/\tilde{\rho})^{2s/3} \\ &\leq C_{39} \bar{M}_j. \end{aligned}$$

In the last inequality we used the fact  $s-1 \geq 2s/3$ . Thus  $\bar{M}_j$  satisfy the monotonicity condition (3.5). Now the left is entirely similar to the proof of Lemma 3.9. And thus (3.32) holds for  $r = 1$  and hence for  $2/3 \leq r \leq 1$  by interpolation inequality (2.2). This completes the proof of Lemma 3.10.

Recall  $C_7$ ,  $C_{35}$  and  $C_{35}$  are the constants appearing in Lemma 3.8, Lemma 3.9 and Lemma 3.10. Now taking  $A$  large enough such that  $A \geq \max\{C_7, C_{35}, C_{38}\}$ , and then by the above three Lemmas we have the truth of  $(E)_{r,N}$  for any  $r \in [0, 1]$ . This complete the proof of Proposition 3.6.

## REFERENCES

- [1] R. Alexandre, S. Ukai, Y. Morimoto, C.-J. Xu, T. Yang, Uncertainty principle and regularity for Boltzmann equation,
- [2] R. Alexandre, L. Desvillettes, C. Villani, B. Wennberg, Entropy dissipation and long-range interactions, *Arch. Rational Mech. Anal.* **152** (2000) 327-355.
- [3] R. Alexandre, M. Safadi, Littlewood Paley decomposition and regularity issues in Boltzmann equation homogeneous equations. I. Non cutoff and Maxwell cases, *M3AM* (2005) 8-15.
- [4] F. Bouchut, Hypoelliptic regularity in kinetic equations. *J. Math. Pure Appl.* **81** (2002), 1135-1159.
- [5] Chen Hua, L.Rodino, General theory of PDE and Gevrey class. *General theory of partial differential equations and microlocal analysis*(Trieste 1995), Pitman Res. Notes in Math. Ser., **349**, Longman, Harlow, 6-81, (1996).
- [6] Chen Hua, L.Rodino, Paradifferential calculus in Gevrey class . *J. Math. Kyoto Univ.* **41**, (2001), 1-31.
- [7] M.Derridj, C.Zuily, Sur la régularité Gevrey des opérateurs de Hörmander. *J.Math.Pures et Appl.* **52** (1973), 309-336.
- [8] L. Desvillettes, B. Wennberg, Smoothness of the solution of the spatially homogeneous Boltzmann equation without cutoff. *Comm. Partial Differential Equations* **29** (2004), no. 1-2, 133–155.
- [9] M.Durand, Régularité Gevrey d'une classe d'opérateurs hypo-elliptiques. *J.Math.Pures et Appl.* **57** (1978), 323-360.
- [10] A.Friedman, On the Regularity of the solutions of Non-linear Elliptic and Parabolic Systems of Partial Differential Equations. *J. Math. Mech.* **7** (1958), 43-59.
- [11] B. Helffer, F. Nier, Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians. *Lecture Notes in Mathematics*, **1862** Springer-Verlag, Berlin, 2005.
- [12] F. Hérau, F. Nier, Isotropic hypoellipticity and trend to equilibrium for the Fokker-Planck equation with a high-degree potential. *Arch. Ration. Mech. Anal.* **171** (2004), no. 2, 151–218.
- [13] L.Hörmander, Hypoelliptic second order differential equations. *Acta Math.* **119** (1967), 147-171.
- [14] J.Kohn, Lectures on degenerate elliptic problems. Pseudodifferential operators with applications, C.I.M.E., Bresanone 1977, 89-151(1978).
- [15] Y.Morimoto, C.-J. Xu, Hypoellipticity for a class of kinetic equations. to appear at J. Math. Kyoto U.
- [16] Y. Morimoto and C.-J. Xu, Logarithmic Sobolev inequality and semi-linear Dirichlet problems for infinitely degenerate elliptic operators, *Astérisque* **284** (2003), 245–264.
- [17] Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, Regularity of solutions to the spatially homogeneous Boltzmann equation without Angular cutoff, preprint.
- [18] L.Rodino, *Linear partial differential operators in Gevrey class*. World Scientific, Singapore, 1993.
- [19] L.P.Rothschild, E.M.Stein, Hypoelliptic differential operators and nilpotent groups. *Acta.Math.* **137** (1977), 248-315.
- [20] F.Treves, *Introduction to Pseudodifferential and Fourier Integral Operators*. Plenum, New York, 1980.
- [21] S. Ukai, Local solutions in Gevrey classes to the nonlinear Boltzmann equation without cutoff, *Japan J. Appl. Math.***1**(1984), no. 1, 141–156.
- [22] C. Villani, On a new class of weak solutions to the spatially homogeneous Boltzmann and Landau equations, *Arch. Rational Mech. Anal.*, **143**, 273–307, (1998).
- [23] C.-J. Xu, Nonlinear microlocal analysis. *General theory of partial differential equations and microlocal analysis*(Trieste 1995), Pitman Res. Notes in Math. Ser., **349**, Longman, Harlow, 155-182, (1996).