



HAL
open science

The integral homology of PSL_2 of imaginary quadratic integers of class number two

Alexander Rahm, Mathias Fuchs

► **To cite this version:**

Alexander Rahm, Mathias Fuchs. The integral homology of PSL_2 of imaginary quadratic integers of class number two. 2009. hal-00370722v1

HAL Id: hal-00370722

<https://hal.science/hal-00370722v1>

Preprint submitted on 26 Mar 2009 (v1), last revised 13 Sep 2010 (v12)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE INTEGRAL HOMOLOGY OF PSL_2 OF IMAGINARY QUADRATIC INTEGERS OF CLASS NUMBER TWO

ALEXANDER RAHM AND MATHIAS FUCHS

ABSTRACT. We calculate the integral homology of the non-Euclidean Bianchi groups $\mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-m}]})$ for $m = 5, 6, 10, 13$, making essential use of an equivariant cellular decomposition of a retract of hyperbolic three-space due to Flöge.

CONTENTS

1. Introduction	1
2. Flöge's complex, contractibility and the spectral sequence	3
3. Computations of the integral homology of $\mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-m}]})$	11
3.1. $m = 13$	11
3.2. $m = 5$	19
3.3. $m = 10$	22
3.4. $m = 6$	27
4. Appendix: The equivariant retraction	30
References	33

1. INTRODUCTION

The object of study of this paper are the PSL_2 -groups Γ of the ring of integers \mathcal{O} of an imaginary quadratic number field $\mathbb{Q}[\sqrt{-m}]$, where m is a square-free positive integer. For m congruent 1 or 2 modulo 4, we have $\mathcal{O}_{\mathbb{Q}[\sqrt{-m}]} = \mathbb{Z}[\sqrt{-m}]$.

The arithmetic groups under study have often been called Bianchi groups, because their study began with the works of Bianchi [4] back in 1892. These groups have since been extensively studied, see for instance the monographs [6, 7, 13].

A space that lends itself naturally to the study of these 2-by-2-matrix groups is the associated symmetric space, hyperbolic three-space \mathcal{H} . We use the upper-half-space model of \mathcal{H} and identify its boundary with $\mathbb{C} \cup \infty \cong \mathbb{C}P^1$. Bianchi exposed a fundamental domain for the action of Γ on the space \mathcal{H} .

Date: March 26, 2009.

2000 Mathematics Subject Classification. 11F75, Cohomology of arithmetic groups. 22E40, Discrete subgroups of Lie groups. 57S30, Discontinuous groups of transformations.

The first named author is supported by a research grant of the Ministère de l'Enseignement supérieur et de la Recherche, and partially supported by DFH-UFA grant CT-26-07-I and by a DAAD (German Academic Exchange Service) grant.

The second named author was supported by a GK Gruppen und Geometrie post-doctoral fellowship of the DFG.

However, the virtual cohomological dimension of arithmetic groups which are lattices in $\mathrm{SL}_2(\mathbb{C})$ is two, so it is desirable to define a proper action on a contractible cellular two-dimensional space. Moreover, this space should be cofinite.

In principle, this has been achieved by Mendoza [14] and Flöge [8] using the reduction theory for binary Hermitian forms of Minkowski, Bianchi, Humbert, Harder and others.

These two approaches have in common that they consider two-dimensional Γ -equivariant retracts which are cocompact and are endowed with a natural CW-structure such that the action of Γ is cellular and the quotient is a finite CW-complex.

In the cases of non-trivial ideal class group, there is a difference between the approaches of Mendoza and Flöge. The elements of the class group of the number field are in bijection with the Γ -orbits of the cusps, where the cusps are the elements of the number field $\mathbb{Q}[\sqrt{-d}]$ and ∞ thought of as elements of the canonical boundary $\mathbb{C}P^1$. The cusps which represent a non-trivial element of the class group are commonly called *singular* points. While Mendoza retracts away from all cusps, Flöge retracts away only from the non-singular ones. Rather than the space \mathcal{H} itself, he considers the space $\widehat{\mathcal{H}}$ obtained from \mathcal{H} by adjoining the Γ -orbits of the singular points. This has the effect that it is possible to retract away only from the non-singular cusps, but onto the singular ones in the appropriate way. It turns out that this gives a suitable contractible 2-dimensional Γ -complex also in the case of non-trivial class group.

Using Mendoza's complex, Schwermer and Vogtmann [16] showed how to calculate the integral group homology in the cases of trivial class group $m = 1, 2, 3, 7, 11$. Vogtmann [20] computed the rational homology as the homology of the quotient in many cases of non-trivial class group. The integral cohomology in the cases $m = 2, 3, 5, 6, 7, 10, 11$ has been determined by Berkove [3], based on Flöge's presentation of the groups with generators and relations.

It is the purpose of the present paper to show how to calculate the integral homology of Bianchi groups with non-trivial class group using Flöge's complex. In the cases considered, Bianchi has computed the fundamental polyhedron. With an implementation in Pari/GP [2], due to the first named author, of Swan's algorithm [19] we re-obtain this polyhedron. In the cases $m = 5, 6, 10$, Flöge has computed the cell stabilizers and cell identifications; and with our Pari/GP program, we redo Flöge's computations and do the same computation in the case $m = 13$. We use the equivariant Euler characteristic to check our computations. With GAP [10] as well as by hand, we find some conjugations between the occurring stabilizer matrices. Then we follow the lines of Schwermer and Vogtmann [16], encountering a spectral sequence which degenerates on the E^3 -page and not already on the E^2 -page as it does in the cases of trivial class group. This is because we have adjoined to our complex the singular points, which have infinite stabilizers. So we have some additional use of homological algebra to obtain the homology of the Bianchi group. Our results are as follows:

For $m = 5$:

$$\mathrm{H}_q(\Gamma; \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^2 \oplus \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^2 & q = 1 \\ \mathbb{Z} \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2 & q = 2 \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^q & q \geq 3; \end{cases}$$

for $m = 10$:

$$\mathrm{H}_q(\Gamma, \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^3 \oplus (\mathbb{Z}/2)^2, & q = 1, \\ \mathbb{Z}^2 \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2, & q = 2, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^q, & q \geq 3; \end{cases}$$

for $m = 6$:

$$\mathrm{H}_q(\Gamma; \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^2 \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2, & q = 1, \\ \mathbb{Z} \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^2, & q = 2, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+2}, & q = 6k + 3, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+1}, & q = 6k + 4, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+4}, & q = 6k + 5, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+1}, & q = 6k + 6, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k}, & q = 6k + 1, q \geq 7, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+3}, & q = 6k + 2, q \geq 8; \end{cases}$$

and for $m = 13$:

$$\mathrm{H}_q(\Gamma, \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^3 \oplus (\mathbb{Z}/2)^2, & q = 1, \\ \mathbb{Z}^2 \oplus \mathbb{Z}/4 \oplus (\mathbb{Z}/3)^2 \oplus \mathbb{Z}/2, & q = 2, \\ (\mathbb{Z}/2)^q \oplus (\mathbb{Z}/3)^2, & q = 4k + 3, \quad k \geq 0, \\ (\mathbb{Z}/2)^q, & q = 4k + 4, \quad k \geq 0, \\ (\mathbb{Z}/2)^q, & q = 4k + 1, \quad k \geq 1, \\ (\mathbb{Z}/2)^q \oplus (\mathbb{Z}/3)^2, & q = 4k + 2, \quad k \geq 1. \end{cases}$$

Thus for $q \geq 2$, the torsion for $m = 5$ is the same as for $m = 10$, analogously to the cohomology results of Berkove [3]. The free part of these homology groups is in accordance with the rational homology results of Vogtmann [20]. We give the full details for our homology computation in the cases $m = 10, 13$. We then give slightly less details in the cases $m = 5, 6$.

The authors would like to thank Philippe Elbaz-Vincent for his inspiration and advice, and the first named author would like to thank Bill Allombert for his help with programming in Pari/GP.

2. FLÖGE'S COMPLEX, CONTRACTIBILITY AND THE SPECTRAL SEQUENCE

Denote the hyperbolic three-space by $\mathcal{H} = \mathbb{C} \times \mathbb{R}_+^*$. We will not use its smooth structure, only its structure as a homogeneous $\mathrm{SL}_2(\mathbb{C})$ -space. The action is given by the formula

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (z, r) := \left(\frac{(\bar{d} - \bar{c}z)(az - b) - r^2 \bar{c}a}{|cz - d|^2 + r^2 |c|^2}, \frac{r}{|cz - d|^2 + r^2 |c|^2} \right);$$

where $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C})$. As usual, we extend the action of $\mathrm{SL}_2(\mathbb{C})$ to the boundary $\mathbb{C}P^1$ which we identify with $\{r = 0\} \cup \infty = \mathbb{C} \cup \infty$. The action passes continuously to the boundary, where it reduces to the usual action

by Möbius transformations $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot z = \frac{az-b}{-cz+d}$. As $-1 \in \mathrm{SL}_2(\mathbb{C})$ acts trivially, the action passes to $\mathrm{PSL}_2(\mathbb{C})$. Let now $m \in \mathbb{N}$ be square-free, and let $R = \mathcal{O}_{\mathbb{Q}[\sqrt{-m}]}$, $\Gamma = \mathrm{PSL}_2(R)$. When the class number of $K = \mathbb{Q}[\sqrt{-m}]$ is one, then classical reduction theory provides a natural equivariant deformation retract of \mathcal{H} which is a CW-complex. This complex is defined as follows. One first considers the union of all hemispheres

$$S_{\mu,\lambda} := \left\{ (z, r) : \left| z - \frac{\lambda}{\mu} \right|^2 + r^2 = \frac{1}{|\mu|^2} \right\} \subset \mathcal{H},$$

for any two μ, λ with $R\mu + \lambda R = R$. Then one considers the “space above the hemispheres”

$$B := \{ (z, r) : |cz - d|^2 + r^2 |c|^2 \geq 1 \\ \text{for all } c, d \in R, c \neq 0 \text{ such that } Rc + Rd = R \}$$

and its boundary ∂B inside \mathcal{H} . For nontrivial class group, the following definition comes to work.

Definition 1. A point $s \in \mathbb{C}P^1 - \{\infty\}$ is called a singular point if for all $c, d \in R$, $c \neq 0$, $Rc + Rd = R$ we have $|cs - d| \geq 1$.

The singular points modulo the action of Γ on $\mathbb{C}P^1$ are in bijection with the nontrivial elements of the class group [18].

In [8], Flöge extends the hyperbolic three-space \mathcal{H} to a larger space $\widehat{\mathcal{H}}$, defined as follows.

Definition 2. As a set, $\widehat{\mathcal{H}} \subset \mathbb{C} \times \mathbb{R}^{\geq 0}$ is the closure under the Γ -action of the union $\widehat{B} := B \cup \{\text{singular points}\}$. The topology is generated by the topology of \mathcal{H} together with the following neighborhoods of the translates s of singular points:

$$\widehat{U}_\epsilon(s) := \{s\} \cup \begin{pmatrix} s & 0 \\ -1 & s^{-1} \end{pmatrix} \cdot \{(z, r) \in \mathcal{H} : r > \epsilon^{-1}\}.$$

Remark 3. The matrix $\begin{pmatrix} s & 0 \\ -1 & s^{-1} \end{pmatrix}$ maps the point at infinity into s , thus giving the point s the topology of ∞ . The neighborhood $\widehat{U}_\epsilon(s)$ is sometimes called a “horoball” because in the upper-half space model it is a Euclidean ball, but with the hyperbolic metric it has “infinite radius”.

The space $\widehat{\mathcal{H}}$ is endowed with the natural Γ -action. Now the essential aspect of Flöge’s construction is the following consequence of Flöge’s theorem [9, 6.6], which we append as theorem 23.

Corollary 4. There is a retraction ρ from $\widehat{\mathcal{H}}$ onto the set $X \subset \widehat{\mathcal{H}}$ of all Γ -translates of $\partial \widehat{B}$, i. e. there is a continuous map $\rho : \widehat{\mathcal{H}} \rightarrow X$ such that $\rho(p) = p$ for all $p \in X$. The set X admits a natural structure as a cellular complex X^\bullet , such that Γ acts cellularly on X^\bullet .

Remark 5. (1) We show with the below lemma that ρ is a homotopy equivalence, without giving a continuous path of maps $\widehat{\mathcal{H}} \rightarrow \widehat{\mathcal{H}}$ connecting ρ to the identity on $\widehat{\mathcal{H}}$.

- (2) The map ρ is Γ -equivariant because its fibers are geodesics. But we do not make use of this fact, as we do not need to show that the homotopy type of $\Gamma \backslash \widehat{\mathcal{H}}$ is the same as that of $\Gamma \backslash X$. This would be useful in the case of trivial class group, i. e. the case of a proper action, to compute the rational homology $H_*(\Gamma; \mathbb{Q}) \cong H_*(\Gamma \backslash \mathcal{H}; \mathbb{Q})$.
- (3) We will provide X^\bullet with a cellular structure which is fine enough to make the cell stabilizers fix the cells pointwise.

Lemma 6. *Let Y be a CW-complex which admits an inclusion i into a contractible topological space A , such that i is a homeomorphism between Y with its cellular topology and the image $i(Y)$ with the subset topology of A . Let $p : A \rightarrow Y$ be a continuous map with $p \circ i = \mathrm{id}_Y$. Then p is a homotopy equivalence.*

Proof. For all $n \in \mathbb{N}$, the induced maps on the homotopy groups $(\mathrm{id}_Y)_* = (p \circ i)_* : \pi_n(Y) \rightarrow \pi_n(Y)$ factor through $\pi_n(A) = 0$, hence are the zero map; and $\pi_n(Y) = 0$. Denote by c the constant map from A to the one-point space. Then $c \circ i$ is a morphism of CW-complexes, and the zero maps it induces on the homotopy groups are isomorphisms. Thus by Whitehead's Theorem [12], $c \circ i$ is a homotopy equivalence. As A is contractible, the composition $(c \circ i) \circ p = c$ is a homotopy equivalence, so the same holds already for p . \square

Together with the lemma below, we obtain a crucial fact underlying this paper.

Corollary 7. *X^\bullet is contractible.*

The following is an observation on Flöge's construction.

Lemma 8. *The space $\widehat{\mathcal{H}}$ is contractible.*

Proof. One can identify the boundary of $\mathcal{H} \cong \{(z, r) \in \mathbb{C} \times \mathbb{R} \mid r > 0\}$ with $\mathbb{C}P^1 \cong \mathbb{C} \cup \infty \cong \{r = 0\} \cup \infty$. By viewing the singular points as part of the boundary, one arrives at an upper half-space model of $\widehat{\mathcal{H}}$.

Now consider $\mathcal{H}_1 := \{(z, r) \in \widehat{\mathcal{H}} : r \geq 1\}$ with the subspace topology of $\widehat{\mathcal{H}}$. The idea of the proof is to consider a vertical retraction onto \mathcal{H}_1 , and to show by an explicit argument that preimages of open sets are open.

Flöge [9, Korollar 5.8] suggests to use the map

$$\phi : \widehat{\mathcal{H}} \times [0, 1] \rightarrow \widehat{\mathcal{H}}, ((z, r), t) \mapsto \begin{cases} (z, r) & \text{for all } t \in [0, 1], & \text{if } r \geq 1 \\ (z, r + t(1 - r)), & & \text{if } r < 1. \end{cases}$$

Let us now check that this is a continuous family of continuous maps. Consider the collection of open balls with respect to the Euclidean metric on $\mathbb{C} \times \mathbb{R}_+$ as soon as they are either contained in $\mathbb{C} \times \mathbb{R}_+^*$, or touch the boundary $\mathbb{C} \times \{0\}$ in a cusp in $\widehat{\mathcal{H}} - \mathcal{H}$. This is a basis for the topology of $\widehat{\mathcal{H}}$. Consider one such open ball \mathcal{B} , and its preimage under some ϕ_t , $t \in [0, 1)$. This either lies entirely in \mathcal{H} , and is open, or it has boundary points. In the latter case, consider the inverse of ϕ_t on $\widehat{\mathcal{H}} - \mathcal{H}_1$, given by

$$\phi_t^{-1} = \left(z, \frac{r-t}{1-t} \right),$$

if this is in $\widehat{\mathcal{H}}$. Suppose there is a cusp s with $s \in \widehat{\mathcal{H}} - \mathcal{H}$ and $\phi_t(s, 0) \in \mathcal{B}$. As \mathcal{B} is open, we find $\beta > 0$ and $\delta > 0$ such that $(s, t + \beta)$ and $(s + \delta, t)$ are in \mathcal{B} . Since

$$\begin{cases} \phi_t\left(s, \frac{\beta}{1-t}\right) = (s, t + \beta) \in \mathcal{B} \\ \phi_t(s + \delta, 0) = (s + \delta, t) \in \mathcal{B}, \end{cases}$$

we know that $(s, \frac{\beta}{1-t})$ and $(s + \delta, 0)$ are in the preimage of \mathcal{B} under ϕ_t . We deduce that the whole horosphere of Euclidean diameter $\min\{\beta, \delta\}$ touching at the cusp s is included in the preimage of \mathcal{B} . Thus each point of the preimage has a neighborhood entirely contained in the preimage, which therefore also is open. The continuity at $t = 1$ as well as the continuity in the variable t follow from very similar arguments.

The space \mathcal{H}_1 is homeomorphic to $\mathbb{C} \times \mathbb{R}_+$, thus contractible. \square

2.0.1. The spectral sequence.

Corollary 7 gives us a contractible complex X^\bullet on which Γ acts cellularly. As a consequence, the integral homology $H_*(\Gamma; \mathbb{Z})$ can be computed as the hyperhomology $\mathbb{H}_*(\Gamma; C_\bullet(X))$ of Γ with coefficients in the cellular chain complex associated to X . This hyperhomology is computable because there is a spectral sequence as in [5, VII] which is also the one used in [16]. It is the spectral sequence associated to the double complex $\Theta_\bullet^\Gamma \otimes_{\mathbb{Z}\Gamma} C_\bullet(X)$ computing the hyperhomology, where we denote by Θ_\bullet^Γ the bar resolution of the group Γ . This spectral sequence can be rewritten (see [16, 1.1]) to yield

$$E_{p,q}^1 = \bigoplus_{\sigma \in \Gamma \backslash X^p} H_q(\Gamma_\sigma; \mathbb{Z}) \implies H_{p+q}(\Gamma; \mathbb{Z}),$$

where Γ_σ denotes the stabilizer of (the chosen representative for) the p -cell σ . We have stated the above E^1 -term with trivial \mathbb{Z} -coefficients in $H_q(\Gamma_\sigma; \mathbb{Z})$, because we use a fundamental domain which is strict enough to give X a cell structure on which Γ acts without inversion of cells.

We shall also make extensive use of the description of the d^1 -differential given in [16].

The technical difference to the cases of trivial class group, treated by [16], is that the stabilizers of the singular points are free abelian groups of rank two. In particular, the Γ -action on our complex X^\bullet is not a *proper action* in the sense that all stabilizers would be finite. As a consequence, the considered spectral sequence does not degenerate on the E^2 -level as it does in Schwermer and Vogtmann's cases.

So we compute a nontrivial differential d^2 , making some additional use of homological algebra, in particular the below lemma and its corollary.

Remark 9. It would be possible to shift the technical difficulty away from homological algebra, using a topological modification of our complex. In our cases of class number two, there is one singular point in the fundamental domain, representing the nontrivial element of the class group. Its stabilizer is free abelian of rank two, and contributes the homology of a torus to the zeroth column of the E^2 -term of our spectral sequence: $H_1(\mathbb{Z}^2; \mathbb{Z}) \cong \mathbb{Z}^2$, $H_2(\mathbb{Z}^2; \mathbb{Z}) \cong \mathbb{Z}$ and $H_q(\mathbb{Z}^2; \mathbb{Z}) = 0$ for $q > 2$. One could modify our complex in order to make the Γ -action on it proper, by replacing each singular point by an \mathbb{R}^2 with the former stabilizer \mathbb{Z}^2 now acting properly. Then the

nontriviality of our differential is equivalent to the existence of a nontrivial homology relation induced by adjoining the torus $\mathbb{R}^2/\mathbb{Z}^2$ to the fundamental domain.

The following lemma will be useful for computing our d^2 -differential. In order to state it, let Γ_σ be a finite subgroup of Γ , let M be a $\mathbb{Z}\Gamma_\sigma$ -module, and let $\ell : \Gamma/\Gamma_\sigma \rightarrow \Gamma$ be a set-theoretical section of the quotient map $\pi : \Gamma \rightarrow \Gamma/\Gamma_\sigma$. Furthermore, denote the standard bar resolution of a discrete group Γ by Θ_\bullet^Γ .

Lemma 10. *The section ℓ defines a map of $\mathbb{Z}\Gamma_\sigma$ -complexes*

$$\hat{\varepsilon}_\ell : \Theta_\bullet^\Gamma \longrightarrow \Theta_\bullet^{\Gamma_\sigma}$$

of degree zero which is a retraction of the resolution Θ_\bullet^Γ of the group Γ to the resolution $\Theta_\bullet^{\Gamma_\sigma}$ of Γ_σ . The map $\hat{\varepsilon}_\ell$ is induced on $\Theta_0^\Gamma = \mathbb{Z}\Gamma$ by

$$\begin{aligned} \Gamma &\xrightarrow{\varepsilon_\ell} \mathbb{Z}\Gamma_\sigma, \\ \gamma &\mapsto (\ell(\pi(\gamma)))^{-1}\gamma \end{aligned}$$

and is continued as a tensor product $\hat{\varepsilon}_\ell = \varepsilon_\ell \otimes \dots \otimes \varepsilon_\ell = \varepsilon_\ell^{\otimes(n+1)}$ on Θ_n^Γ .

Remark 11. (1) Attention: ε_ℓ is a $\mathbb{Z}\Gamma_\sigma$ -linear map because Γ_σ acts from the right.

(2) Note that the resulting isomorphism in homology

$$\mathrm{H}_*(\Theta_\bullet^\Gamma \otimes_{\mathbb{Z}\Gamma_\sigma} M) \longrightarrow \mathrm{H}_*(\Theta_\bullet^{\Gamma_\sigma} \otimes_{\mathbb{Z}\Gamma_\sigma} M)$$

is independent of the choice of ℓ , and consistent with the canonical isomorphisms of both sides with $\mathrm{H}_*(\Gamma_\sigma; M)$.

(3) Note that in the above lemma, it is not necessary to require $\ell(\pi(1)) = 1$.

This would imply that ε_ℓ is the identity on $\Theta_\bullet^{\Gamma_\sigma}$. However, we will choose $\ell(\pi(1)) = 1$ for simplicity.

(4) In explicit terms, the map ε_ℓ is described as follows.

$$\varepsilon_\ell : \mathbb{Z}\Gamma \rightarrow \mathbb{Z}\Gamma_\sigma,$$

$$\sum_{\gamma \in \Gamma} a_\gamma \gamma = \sum_{\gamma_\sigma \in \Gamma_\sigma} \sum_{\rho \in \Gamma/\Gamma_\sigma} a_{\gamma_\sigma \ell(\rho)} \gamma_\sigma \ell(\rho) \mapsto \sum_{\gamma_\sigma \in \Gamma_\sigma} \left(\sum_{\rho \in \Gamma/\Gamma_\sigma} a_{\gamma_\sigma \ell(\rho)} \right) \gamma_\sigma,$$

where the a_γ are coefficients from \mathbb{Z} . The map ε_ℓ restricts to the identity on $\mathbb{Z}\Gamma_\sigma$ and gives an isomorphism of \mathbb{Z} -modules

$$\mathbb{Z}[\ell(\rho)\Gamma_\sigma] \rightarrow \mathbb{Z}\Gamma_\sigma$$

for every Γ_σ -orbit $\ell(\rho)\Gamma_\sigma$.

Proof (of the lemma). In fact, the statement holds for any chain map $\hat{\varepsilon}$ in the place of $\hat{\varepsilon}_\ell$ that satisfies the following conditions. They are easily checked to hold for the maps $\hat{\varepsilon}_\ell$.

(1) $\hat{\varepsilon}$ is $\mathbb{Z}\Gamma_\sigma$ -linear.

(2) The augmentation $\Theta_0^\Gamma \rightarrow \mathbb{Z}$ is the composition of $\hat{\varepsilon}$ with the augmentation $\Theta_0^{\Gamma_\sigma} \rightarrow \mathbb{Z}$.

Then the statement follows from the comparison theorem [21, 2.2.6] of fundamental homological algebra. In fact, the properties imply that $\hat{\varepsilon}$ is a chain map of resolutions lifting the identity on \mathbb{Z} . An inverse is given by the canonical inclusion $\Theta_{\bullet}^{\Gamma_\sigma} \rightarrow \Theta_{\bullet}^{\Gamma}$, and since the composition is unique up to chain homotopy equivalence, it must be homotopic to the identity. \square

The group Γ_σ acts diagonally on $\Theta_1^\Gamma \cong \mathbb{Z}\Gamma \otimes_{\mathbb{Z}} \mathbb{Z}\Gamma$, and trivially on \mathbb{Z} , so we can consider $\Theta_1^\Gamma \otimes_{\mathbb{Z}\Gamma_\sigma} \mathbb{Z}$.

Corollary 12. *Let a cycle $(\sum_i (a_i \otimes_{\mathbb{Z}} b_i) \otimes_{\mathbb{Z}\Gamma_\sigma} 1) \in \Theta_1^\Gamma \otimes_{\mathbb{Z}\Gamma_\sigma} \mathbb{Z}$ be given, where $a_i, b_i \in \Gamma$. The ensuing element in $H_1(\Gamma_\sigma, \mathbb{Z})$ is then given by*

$$\sum_i \varepsilon(a_i) \overline{\varepsilon_\ell(a_i)^{-1} \varepsilon_\ell(b_i)},$$

where ε is the augmentation from $\mathbb{Z}\Gamma$ to \mathbb{Z} .

This expression makes sense because $\varepsilon_\ell(a_i)$ is invertible in $\mathbb{Z}\Gamma_\sigma$. Note that the cycle condition on $\sum_i (a_i \otimes_{\mathbb{Z}} b_i) \otimes_{\mathbb{Z}\Gamma_\sigma} 1$ says that

$$\sum_i (b_i - a_i) \otimes_{\mathbb{Z}\Gamma_\sigma} 1 = 0.$$

which means that $\sum_i a_i$ is equivalent to $\sum_i b_i$ modulo $\mathbb{Z}\Gamma_\sigma$.

Proof. Using the lemma 10, we just need to apply the map

$$(\varepsilon_\ell \otimes_{\mathbb{Z}} \varepsilon_\ell) \otimes_{\mathbb{Z}\Gamma_\sigma} 1 : (\mathbb{Z}\Gamma \otimes_{\mathbb{Z}} \mathbb{Z}\Gamma) \otimes_{\mathbb{Z}\Gamma_\sigma} \mathbb{Z} \rightarrow (\mathbb{Z}\Gamma_\sigma \otimes_{\mathbb{Z}} \mathbb{Z}\Gamma_\sigma) \otimes_{\mathbb{Z}\Gamma_\sigma} \mathbb{Z}$$

to get

$$\begin{aligned} \sum (\varepsilon_\ell \otimes_{\mathbb{Z}} \varepsilon_\ell \otimes_{\mathbb{Z}\Gamma_\sigma} 1)(a_i \otimes_{\mathbb{Z}} b_i \otimes_{\mathbb{Z}\Gamma_\sigma} 1) &= \sum (\varepsilon_\ell(a_i) \otimes_{\mathbb{Z}} \varepsilon_\ell(b_i)) \otimes_{\mathbb{Z}\Gamma_\sigma} 1 \\ &= \sum (1 \otimes_{\mathbb{Z}} \varepsilon_\ell(a_i)^{-1} \varepsilon_\ell(b_i)) \otimes_{\mathbb{Z}\Gamma_\sigma} \varepsilon(a_i). \end{aligned}$$

In bar notation, this is $\sum [\varepsilon_\ell(a_i)^{-1} \varepsilon_\ell(b_i)] \otimes_{\mathbb{Z}\Gamma_\sigma} \varepsilon(a_i)$, and is mapped to

$$\sum_i \varepsilon(a_i) \overline{\varepsilon_\ell(a_i)^{-1} \varepsilon_\ell(b_i)} \in H_1(\Gamma_\sigma, \mathbb{Z}),$$

and to

$$\sum_i \varepsilon(a_i) \varepsilon_\ell(a_i)^{-1} \varepsilon_\ell(b_i) \pmod{[\Gamma_\sigma, \Gamma_\sigma]}$$

by the isomorphism into the abelianization of Γ_σ described in [5, page 36]. \square

2.0.2. The mass formula for the Euler characteristic.

We will use the Euler characteristic to check the geometry of the quotient $\Gamma \backslash X$. Recall the following definitions and proposition, which we include for the reader's convenience.

Definition 13 (Euler characteristic). *Suppose Γ' is a torsion-free group. Then we define its Euler characteristic as*

$$\chi(\Gamma') = \sum_i (-1)^i \dim H_i(\Gamma'; \mathbb{Q}).$$

Suppose further that Γ' is a torsion-free subgroup of finite index in a group Γ . Then we define the Euler characteristic of Γ as

$$\chi(\Gamma) = \frac{\chi(\Gamma')}{[\Gamma : \Gamma']}.$$

This is well-defined because of [5, IX.6.3].

Definition 14 (Equivariant Euler characteristic). *Suppose X is a Γ -complex such that*

- (1) every isotropy group Γ_σ is of finite homological type;
- (2) X has only finitely many cells mod Γ .

Then we define the Γ -equivariant Euler characteristic of X as

$$\chi_\Gamma(X) := \sum_{\sigma} (-1)^{\dim \sigma} \chi(\Gamma_\sigma),$$

where σ runs over the orbit representatives of cells of X .

Proposition 15 ([5, IX.7.3 e']). *Suppose X is a Γ -complex such that $\chi_\Gamma(X)$ is defined. If Γ is virtually torsion-free, then Γ is of finite homological type and $\chi(\Gamma) = \chi_\Gamma(X)$.*

Let now Γ be $\mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-m}]})$. Then the above proposition applies to X taken to be Flöge's (or still, Mendoza's) Γ -equivariant deformation retract of \mathcal{H} . Using $\chi(\Gamma_\sigma) = \frac{1}{\mathrm{card}(\Gamma_\sigma)}$ for Γ_σ finite, the fact that the singular points have stabilizer \mathbb{Z}^2 , and the torsion-free Euler characteristic

$$\chi(\mathbb{Z}^2) = \sum_i (-1)^i \mathrm{rank}_{\mathbb{Z}}(\mathrm{H}_i \mathbb{Z}^2) = 1 - 2 + 1 = 0,$$

we get the formula

$$\chi(\Gamma) = \sum_{\sigma} (-1)^{\dim \sigma} \frac{1}{\mathrm{card}(\Gamma_\sigma)},$$

where σ runs over the orbit representatives of cells of X with finite stabilizers.

Proposition 16. *The Euler characteristic $\chi(\Gamma)$ vanishes.*

Remark 17. This, together with the formula

$$0 = \chi(\Gamma) = \chi_\Gamma(X) = \sum_{\sigma} (-1)^{\dim \sigma} \frac{1}{\mathrm{card}(\Gamma_\sigma)},$$

allows to check the joint data of the geometry of the fundamental domain, cell stabilizers and cell identifications.

Proof of the latter proposition. Denote by ζ_K the Dedekind zeta function associated to the number field $K := \mathbb{Q}[\sqrt{-m}]$.

Brown [5, below (IX.8.7)] deduces the following formula from Harder's result [11, page 453]:

$$\chi(\mathrm{SL}_n(\mathcal{O}_K)) = \prod_{j=2}^n \zeta_K(1-j),$$

so especially

$$\chi(\mathrm{SL}_2(\mathcal{O}_K)) = \zeta_K(-1).$$

As any cell σ in the interior of hyperbolic space has a stabilizer $SL_2(\mathcal{O}_K)_\sigma$ of twice the cardinality of Γ_σ , we have

$$\chi(\Gamma) = \frac{1}{2}\chi(SL_2(\mathcal{O}_K)) = \frac{1}{2}\zeta_K(-1).$$

Using the functional equation of ζ_K [15] and the fact that K has no real places because it is imaginary quadratic, we get

$$\zeta_K(-1) = 0.$$

□

Remark 18. One can prove the above proposition without using the Dedekind zeta function. This alternative proof applies to *any* cofinite arithmetically defined subgroup Γ of $PSL(2, \mathbb{C})$. It is the main theorem of Harder's article on the Gauss-Bonnet theorem [11] that the Euler characteristic is the covolume of Γ with respect to the Euler-Poincaré form μ on \mathcal{H} , i. e. $\chi(\Gamma) = \int_Y d\mu$, where Y is a fundamental domain for the action of Γ on \mathcal{H} . This extends the classical Gauss-Bonnet theorem from the theory of the Euler-Poincaré form, see [17, paragraph 3] (here the theorem is hidden as the existence assertion of the Euler-Poincaré measure) to non-cocompact but cofinite discrete subgroups. The measure μ is a fundamental datum associated to the symmetric space, without reference to any discrete group. In [17, paragraph 3,2a] it is shown that $\mu = 0$ on any odd-dimensional space. Since $\dim \mathcal{H} = 3$, we have $\chi(\Gamma) = 0$.


3. COMPUTATIONS OF THE INTEGRAL HOMOLOGY OF $\mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-m}]})$

Throughout this section, we mean trivial \mathbb{Z} -coefficients wherever we leave out the coefficients. We will always label the singular point in the fundamental domain by s ; and we use the notation

$$\otimes_{\sigma} := \otimes_{\mathbb{Z}[\Gamma_{\sigma}]}.$$

Let $\omega := \sqrt{-m}$. We have $\Gamma = \mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-m}]}) = \mathrm{PSL}_2(\mathbb{Z}[\omega])$ in the cases $m = 5, 6, 10, 13$ considered.

3.1. $m = 13$. The fundamental domain for Γ looks as follows.



The coordinates in Hyperbolic space of the vertices can be found in [4]. We make the following definitions.

$$\begin{aligned} A &:= \pm \begin{pmatrix} 9 & 7\omega \\ \omega & -10 \end{pmatrix}, & B &:= \pm \begin{pmatrix} -2-\omega & 2-\omega \\ 4 & 2+1\omega \end{pmatrix}, & C &:= \pm \begin{pmatrix} -1-\omega & 8-\omega \\ 3 & 1+2\omega \end{pmatrix}, \\ D &:= \pm \begin{pmatrix} 5 & 2\omega \\ \omega & -5 \end{pmatrix}, & E &:= \pm \begin{pmatrix} -\omega & 6 \\ 2 & \omega \end{pmatrix}, & J &:= \pm \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}, \\ S &:= \pm \begin{pmatrix} & -1 \\ 1 & 1 \end{pmatrix}, & K &:= \pm \begin{pmatrix} 11+4\omega & -17+7\omega \\ -8+\omega & -10-3\omega \end{pmatrix}, & M &:= \pm \begin{pmatrix} 4-2\omega & 12+\omega \\ 4+\omega & -4+2\omega \end{pmatrix}, \\ U &:= \pm \begin{pmatrix} 1 & \omega \\ & 1 \end{pmatrix}, & V &:= \pm \begin{pmatrix} -\omega & 6-\omega \\ 2 & 2+\omega \end{pmatrix}, & W &:= \pm \begin{pmatrix} 14-\omega & 13+6\omega \\ 2\omega & -12+\omega \end{pmatrix}, \\ P &:= V^{-1}D, & T &:= P^{-1}S^2, & R &:= TU^{-1}S^2U. \end{aligned}$$

We observe the relations $T = CKCA(CKC)^{-1}$, $V^{-1} = CAC^{-1}M$ and $S^2 = BS^{-1}BS$. The matrix U acts as a vertical translation by $-\omega$ on this fundamental domain. There are seventeen orbits of vertices, which have the

following stabilizers.

$$\begin{aligned}
 \Gamma_o &= \langle J|J^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_e &= \langle A, U^{-1}JU|A^3 = (U^{-1}JU)^2 = (AU^{-1}JU)^2 = 1 \rangle \cong S_3, \\
 \Gamma_f &= \langle D, E|D^2 = E^2 = (DE)^2 = 1 \rangle \cong D_2, \\
 \Gamma_g &= \langle J, T|J^2 = T^3 = (JT)^2 = 1 \rangle \cong S_3, \\
 \Gamma_h &= \langle E, AU^{-1}JU|E^2 = (AU^{-1}JU)^2 = (EAU^{-1}JU)^2 = 1 \rangle \cong D_2, \\
 \Gamma_j &= \langle S|S^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_s &= \langle V, W|VW = WV \rangle \cong \mathbb{Z}^2, \\
 \Gamma_t &= \langle R, U^{-1}SU|R^2 = (U^{-1}SU)^3 = (RU^{-1}SU)^2 = 1 \rangle \cong S_3, \\
 \Gamma_u &= \langle B|B^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_v &= \langle D|D^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_w &= \langle B, S|B^2 = S^3 = (BS)^2 = 1 \rangle \cong S_3, \\
 \Gamma_x &= \Gamma_z = \langle CAC^{-1}|(CAC^{-1})^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_a &= \langle S^{-1}BS|(S^{-1}BS)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_b &= \Gamma_c = \langle M|M^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_y &= \langle T|T^3 = 1 \rangle \cong \mathbb{Z}/3.
 \end{aligned}$$

There are twenty-eight orbits of edges. The edge stabilizers of isomorphy type $\mathbb{Z}/2$ are given on the chosen edge orbit representatives as

$$\begin{aligned}
 \Gamma_{(f,v)} &= \langle D|D^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(h,u')} &= \langle EAU^{-1}JU|(EAU^{-1}JU)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(t,b'')} &= \langle R|R^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(w,a)} &= \langle S^{-1}BS|(S^{-1}BS)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(b,c)} &= \langle M|M^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(a'',c'')} &= \langle C^{-1}S^{-1}BSC|(C^{-1}S^{-1}BSC)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(v',t)} &= \langle RU^{-1}SU|(RU^{-1}SU)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(w,u)} &= \langle B|B^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(h,e)} &= \langle AU^{-1}JU|(AU^{-1}JU)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(g,f)} &= \langle DE|(DE)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(f,h)} &= \langle E|E^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(o,g)} &= \langle J|J^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(o',e)} &= \langle U^{-1}JU|(U^{-1}JU)^2 = 1 \rangle \cong \mathbb{Z}/2.
 \end{aligned}$$

and the edge stabilizers of isomorphy type $\mathbb{Z}/3$ are given on the chosen edge orbit representatives as

$$\begin{aligned}
 \Gamma_{(e,x')} &= \langle A|A^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(x,z)} &= \langle CAC^{-1}|(CAC^{-1})^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(g,y)} &= \langle T|T^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(j,w)} &= \langle S|S^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(t,j')} &= \langle U^{-1}SU|(U^{-1}SU)^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(y',z')} &= \langle KCA(KC)^{-1}|(KCA(KC)^{-1})^3 = 1 \rangle \cong \mathbb{Z}/3.
 \end{aligned}$$

We use the identifications $C \cdot x' = x$, $U \cdot j' = j$, $C \cdot y' = y$ and $K \cdot z = z'$. We find nine edge orbits with the trivial stabilizer, thirteen edge orbit representatives with stabilizer type $\mathbb{Z}/2$, and six with stabilizer type $\mathbb{Z}/3$. The singular vertex has stabilizer type \mathbb{Z}^2 , and there are six vertex orbit representatives with stabilizer type $\mathbb{Z}/2$, two with D_2 , four with S_3 and four with $\mathbb{Z}/3$. Furthermore, there are twelve orbits of faces with trivial stabilizers. The above data gives the Γ -equivariant Euler characteristic of X :

$$\chi_\Gamma(X) = 6 \cdot \frac{1}{2} + 4 \cdot \frac{1}{3} + 2 \cdot \frac{1}{4} + 4 \cdot \frac{1}{6} - 9 - 13 \cdot \frac{1}{2} - 6 \cdot \frac{1}{3} + 12 = 0,$$

in accordance with remark 17.

3.1.1. E^1 -page for $m = 13$.

We obtain for the row $q = 0$ in the columns $p = 0, 1, 2$:

$$\mathbb{Z}^{17} \xleftarrow{d_{1,0}^1} \mathbb{Z}^{28} \xleftarrow{d_{2,0}^1} \mathbb{Z}^{12},$$

for q congruent 1 modulo 4.

For $q = 1$, there is an additional module $H_1(\Gamma_s) \cong \mathbb{Z}^2$ on the target side, which can not be hit because the edge stabilizers are only torsion.

Remark 19. So, the 3-torsion in $H_1(\Gamma)$ has already been killed by the d^1 differential. This shows that there is no injection

$$H_1(\mathrm{PSL}_2(\mathbb{Z})) \rightarrow H_1(\Gamma).$$

We verify this fact by considering the generator of the 3-torsion in $H_1(\mathrm{PSL}_2(\mathbb{Z}))$, which is induced by the matrix S . In the group Γ for $m = 13$, the matrix S of order three is subject to the relation $S^2 = BS^{-1}BS$ where B is a matrix of order two defined above. The right hand side of this equation can be simplified to the unit element when we pass it to $\Gamma^{ab} \cong H_1(\Gamma)$. So, the above non-injectivity is based on the fact that S does not survive abelianizing Γ while it survives abelianizing $\mathrm{PSL}_2(\mathbb{Z})$.

3.1.3. *The rows with q even.*

There is a zero map arriving at $\bigoplus_{\sigma \in \Gamma \backslash X^0} H_q(\Gamma_\sigma) = (\mathbb{Z}/2)^q$ for q bigger than 2, and respectively at

$$\bigoplus_{\sigma \in \Gamma \backslash X^0} H_2(\Gamma_\sigma) = \mathbb{Z} \oplus (\mathbb{Z}/2)^2.$$

3.1.4. *The E^2 -page.* In the rows with $q \geq 2$, $E_{p,q}^2$ is concentrated in the columns $p = 0$ and $p = 1$ given as follows:

$$\begin{array}{l|ll} q = 4k + 1, & q \geq 5 & (\mathbb{Z}/2)^q & (\mathbb{Z}/3)^2 \\ q \text{ even,} & q \geq 4 & (\mathbb{Z}/2)^q & 0 \\ q = 4k + 3, & q \geq 3 & (\mathbb{Z}/3)^2 \oplus (\mathbb{Z}/2)^q & 0 \\ \dots & & \dots & \dots \\ q = 2 & & \mathbb{Z} \oplus (\mathbb{Z}/2)^2 & 0 \end{array}$$

In the rows $q = 0$ and $q = 1$, $E_{p,q}^2$ is concentrated in the columns $p = 0, 1, 2$:

$$\begin{array}{cccc} q = 1 & \mathbb{Z}^2 \oplus (\mathbb{Z}/2)^2 & (\mathbb{Z}/3)^2 \oplus \mathbb{Z}/2 & 0 \\ & \swarrow d^2 & & \\ q = 0 & \mathbb{Z} & \mathbb{Z}^2 & \mathbb{Z}^2 \end{array}$$

3.1.5. *Computation of the differential d^2 .*

The only nontrivial d^2 -arrow is determined on the E^0 -level by the following maps connecting $E_{2,0}^0$ with $E_{0,1}^0$:

$$\begin{array}{ccc} \bigoplus_{\sigma \in \Gamma \backslash X^0} \Theta_1 \otimes_\sigma \mathbb{Z} & \xleftarrow{1 \otimes \delta} & \bigoplus_{\sigma \in \Gamma \backslash X^1} \Theta_1 \otimes_\sigma \mathbb{Z} \\ & & \downarrow d_\Theta \otimes 1 \\ & & \bigoplus_{\sigma \in \Gamma \backslash X^1} \Theta_0 \otimes_\sigma \mathbb{Z} \xleftarrow{1 \otimes \delta} \bigoplus_{\sigma \in \Gamma \backslash X^2} \Theta_0 \otimes_\sigma \mathbb{Z} \end{array}$$

where d_Θ is the differential of the bar resolution Θ_\bullet for Γ , and δ is the differential of Flöge's cellular complex. The generators of the abelian group $E_{2,0}^2 \cong \mathbb{Z}^2$ are represented by the face (c, s, c', z) and the union of two faces $(b, x, b', v', y', a', u, a, y, v) =: F$, whose quotients by Γ are homeomorphic to 2-spheres. We observe the edge identifications $CAC^{-1} \cdot (c, z) = (c', z)$,

$$V^{-1} \cdot (s, c) = (s, c'), \quad CAC^{-1} \cdot (b, x) = (b', x), \quad V^{-1} \cdot (b, v) = (b', v'), \\ P \cdot (y, v) = (y', v'), \quad S^2 \cdot (a, y) = (a', y'), \quad \text{and } B \cdot (a, u) = (a', u).$$

The above d^2 -arrow is induced by

$$\delta((c, s, c', z)) = (CAC^{-1} - 1) \cdot (c, z) + (V^{-1} - 1) \cdot (s, c)$$

and

$$\delta((b, x, b', v', y', a', u, a, y, v)) = (CAC^{-1} - 1) \cdot (x, b) + (V^{-1} - 1) \cdot (b, v) \\ + (P - 1) \cdot (v, y) + (S^2 - 1) \cdot (y, a) + (B - 1) \cdot (a, u).$$

The lift $1 \otimes_F 1$ in $E_{2,0}^0$ of the generator of $E_{2,0}^2$ represented by $F = (b, x, b', v', y', a', u, a, y, v)$ is mapped as follows:

$$\begin{array}{ccc} (1, CAC^{-1}) \otimes_b 1 - (1, CAC^{-1}) \otimes_x 1 & & (1, CAC^{-1}) \otimes_{(x,b)} 1 \\ + (1, V^{-1}) \otimes_v 1 - (1, V^{-1}) \otimes_b 1 & & + (1, V^{-1}) \otimes_{(b,v)} 1 \\ + (1, P) \otimes_y 1 - (1, P) \otimes_v 1 & \longleftarrow & + (1, P) \otimes_{(v,y)} 1 \\ + (1, S^2) \otimes_a 1 - (1, S^2) \otimes_y 1 & & + (1, S^2) \otimes_{(y,a)} 1 \\ + (1, B) \otimes_u 1 - (1, B) \otimes_a 1 & & + (1, B) \otimes_{(a,u)} 1 \\ & & \downarrow d_{\Theta} \otimes 1 \\ & & (CAC^{-1} - 1) \otimes_{(x,b)} 1 \\ & & + (V^{-1} - 1) \otimes_{(b,v)} 1 \\ & & + (P - 1) \otimes_{(v,y)} 1 \longleftarrow 1 \otimes_F 1 \\ & & + (S^2 - 1) \otimes_{(y,a)} 1 \\ & & + (B - 1) \otimes_{(a,u)} 1 \end{array}$$

The passage to E^1 . We attribute the symbols t_σ to the part of this sum lying in $\Theta_1 \otimes_\sigma \mathbb{Z}$:

$$\begin{aligned} t_x &:= -(1, CAC^{-1}) \otimes_x 1, \\ t_b &:= (1, CAC^{-1}) \otimes_b 1 - (1, V^{-1}) \otimes_b 1, \\ t_v &:= (1, V^{-1}) \otimes_v 1 - (1, P) \otimes_v 1, \\ t_y &:= (1, P) \otimes_y 1 - (1, S^2) \otimes_y 1, \\ t_a &:= (1, S^2) \otimes_a 1 - (1, B) \otimes_a 1, \\ t_u &:= (1, B) \otimes_u. \end{aligned}$$

With the formula in our corollary 12, we find the classes \bar{t}_σ in $H_1(\Theta_* \otimes_\sigma \mathbb{Z})$ as follows: As $V^{-1}M = CAC^{-1}$ and $\Gamma_b = \langle M \mid M^2 = 1 \rangle$,

$$t_b = [CAC^{-1}] \otimes_b 1 - [V^{-1}] \otimes_b 1 = [V^{-1}M] \otimes_b 1 - [V^{-1}] \otimes_b 1$$

gives the cycle

$$\overline{VV^{-1}M} - \overline{VV^{-1}} = \overline{M} \in \langle \overline{M} \mid 2\overline{M} = 0 \rangle \cong H_1(\Gamma_b, \mathbb{Z}).$$

As $V^{-1} = PD$ and $\Gamma_v = \langle D \mid D^2 = 1 \rangle$,

$$t_v = [V^{-1}] \otimes_v 1 - [P] \otimes_v 1 = [PD] \otimes_v 1 - [P] \otimes_v 1$$

gives the cycle

$$\overline{P^{-1}PD} - \overline{P^{-1}P} = \overline{D} \in \langle \overline{D} \mid 2\overline{D} = 0 \rangle \cong H_1(\Gamma_v, \mathbb{Z}).$$

As $S^2 = BS^{-1}BS$ and $\Gamma_a = \langle S^{-1}BS \mid (S^{-1}BS)^2 = 1 \rangle$,

$$t_a = [S^2] \otimes_a 1 - [B] \otimes_a 1 = [BS^{-1}BS] \otimes_a 1 - [B] \otimes_a 1$$

gives the cycle

$$\overline{B^{-1}BS^{-1}BS} - \overline{B^{-1}B} = \overline{S^{-1}BS} \in \langle \overline{S^{-1}BS} \mid 2\overline{S^{-1}BS} = 0 \rangle \cong H_1(\Gamma_a, \mathbb{Z}).$$

Finally, $t_u = [B] \otimes_u 1$ gives the cycle

$$\overline{B} \in \langle \overline{B} \mid 2\overline{B} = 0 \rangle \cong H_1(\Gamma_u, \mathbb{Z}).$$

The term $E_{0,1}^2$ having no 3-torsion, the 3-torsion part $\overline{t}_x + \overline{t}_y$ of the above sum makes no contribution to the image of d^2 .

The 2-torsion part, $\overline{t}_b + \overline{t}_a + \overline{t}_v + \overline{t}_u$, equals the image $d_{1,1}^1(\overline{t}_{(b,c)} + \overline{t}_{(c'',a'')} + \overline{t}_{(v,f)} + \overline{t}_{(f,h)} + \overline{t}_{(h,u'')})$, where \overline{t}_σ stands for the generator of $H_1(\Gamma_\sigma, \mathbb{Z}) \cong \mathbb{Z}/2$. Hence it makes no contribution neither, and we obtain $d^2(F) = 0$.

The lift $1 \otimes_{(c,s,c',z)} 1$ of the generator (c, s, c', z) is mapped as follows:

$$\begin{array}{ccc} \begin{array}{l} (1, CAC^{-1}) \otimes_z 1 \\ -(1, CAC^{-1}) \otimes_c 1 \\ +(1, V^{-1}) \otimes_c 1 \\ -(1, V^{-1}) \otimes_s 1 \end{array} & \xleftarrow{1 \otimes \delta} & \begin{array}{l} (1, CAC^{-1}) \otimes_{(c,z)} 1 \\ +(1, V^{-1}) \otimes_{(s,c)} 1 \end{array} \\ & & \downarrow d_\Theta \otimes 1 \\ & & \begin{array}{l} (CAC^{-1} - 1) \otimes_{(c,z)} 1 \\ +(V^{-1} - 1) \otimes_{(s,c)} 1 \end{array} \xleftarrow{1 \otimes \delta} 1 \otimes_{(c,s,c',z)} 1 \end{array}$$

The passage to E^1 . We attribute the symbols t_σ to the part of this sum lying in $\Theta_1 \otimes_\sigma \mathbb{Z}$:

$$\begin{aligned} t_z &:= (1, CAC^{-1}) \otimes_z 1, \\ t_c &:= (1, V^{-1}) \otimes_c 1 - (1, CAC^{-1}) \otimes_c 1, \\ t_s &:= -(1, V^{-1}) \otimes_s 1. \end{aligned}$$

With the formula in our corollary 12, we find the classes \overline{t}_σ in $H_1(\Theta_* \otimes_\sigma \mathbb{Z})$ as follows:

$$t_z = [CAC^{-1}] \otimes_z 1$$

gives the cycle

$$\overline{CAC^{-1}} \in \langle \overline{CAC^{-1}} \mid 3\overline{CAC^{-1}} = 0 \rangle \cong H_1(\Gamma_z, \mathbb{Z}).$$

As $V^{-1}M = CAC^{-1}$ and $\Gamma_c = \langle M \mid M^2 = 1 \rangle$,

$$t_c = [V^{-1}] \otimes_c 1 - [CAC^{-1}] \otimes_c 1 = [V^{-1}] \otimes_c 1 - [V^{-1}M] \otimes_c 1$$

gives the cycle

$$\overline{VV^{-1}} - \overline{VV^{-1}M} = -\overline{M} \in \langle \overline{M} \mid 2\overline{M} = 0 \rangle \cong H_1(\Gamma_c, \mathbb{Z}).$$

Finally,

$$t_s = -[V^{-1}] \otimes_s 1$$

gives the cycle

$$\overline{V} \in \langle \overline{V}, \overline{W} \rangle \cong H_1(\Gamma_s, \mathbb{Z}) \cong \mathbb{Z}^2.$$

The term $E_{0,1}^2$ having no 3-torsion, the 3-torsion part \overline{t}_z of the above sum makes no contribution to the image of d^2 .

However the 2-torsion part, $\overline{t}_c = \overline{M}$, passes to the E^2 -page because no chain of edges can have the single point c as its boundary. Furthermore, \overline{V} is one of the generators of the free part of $E_{0,1}^2 \cong \mathbb{Z}^2 \oplus (\mathbb{Z}/2)^2$, so we obtain $d^2((c, s, c', z)) = \overline{M} + \overline{V}$, which is of infinite order and has the following property: There is no element $\eta \in E_{0,1}^2$ with $k\eta = \overline{M} + \overline{V}$ for an integer $k > 1$. As we have seen that $d^2(F) = 0$, we obtain the quotient

$$E_{0,1}^3 \cong \mathbb{Z} \oplus (\mathbb{Z}/2)^2.$$

Hence we obtain for integral homology the following short exact sequences:

$$\begin{cases} 0 \rightarrow (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow (\mathbb{Z}/3)^2 \rightarrow 0, & q = 4k + 2, \\ 0 \rightarrow (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0, & q = 4k + 1, \\ 0 \rightarrow (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0, & q = 4k + 4, \\ 0 \rightarrow (\mathbb{Z}/3)^2 \oplus (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0, & q = 4k + 3, \\ 0 \rightarrow \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \rightarrow H_2(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z} \oplus (\mathbb{Z}/3)^2 \oplus \mathbb{Z}/2 \rightarrow 0, \\ 0 \rightarrow \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \rightarrow H_1(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}^2 \rightarrow 0. \end{cases}$$

3.1.6. *The spectral sequence for $m = 13$ with $\mathbb{Z}/2$ -coefficients.*

From [16, lemma 4.2(3)] we compute by elementary means that $H_q(\mathcal{S}_3; \mathbb{Z}/2) \cong \mathbb{Z}/2$, $H_q(\mathbb{Z}/2; \mathbb{Z}/2) \cong \mathbb{Z}/2$ and $H_q(\mathcal{D}_2; \mathbb{Z}/2) \cong (\mathbb{Z}/2)^{q+1}$ for all $q \geq 0$. Also by elementary means, we get

$$H_q(\mathbb{Z}^2, \mathbb{Z}/2) \cong \begin{cases} 0, & q \geq 3, \\ \mathbb{Z}/2, & q = 2, \\ (\mathbb{Z}/2)^2, & q = 1. \end{cases}$$

3.1.7. *E^1 -page with $\mathbb{Z}/2$ -coefficients.*

We can apply the functor $- \otimes \mathbb{Z}/2$ to the row $q = 0$ and obtain in the columns $p = 0, 1, 2$:

$$(\mathbb{Z}/2)^{17} \xleftarrow{d_{1,0}^1} (\mathbb{Z}/2)^{28} \xleftarrow{d_{2,0}^1} (\mathbb{Z}/2)^{12}.$$

The rest of this row are zeroes. The matrix $d_{1,0}^1$ has rank 16 and the matrix $d_{2,0}^1$ has rank 10. In the rows with $q > 0$, the differential d^1 is given by a single arrow $d_{1,q}^1$ from $E_{1,q}^1 = (H_q(\mathbb{Z}/2; \mathbb{Z}/2))^{13} \oplus (H_q(\mathbb{Z}/3; \mathbb{Z}/2))^6 \cong (\mathbb{Z}/2)^{13}$ to

$$E_{0,q}^1 = H_q(\mathbb{Z}^2; \mathbb{Z}/2) \oplus (H_q(\mathbb{Z}/2; \mathbb{Z}/2))^6 \oplus (H_q(\mathcal{D}_2; \mathbb{Z}/2))^2 \oplus (H_q(\mathcal{S}_3; \mathbb{Z}/2))^4,$$

and the rest of these rows are zeroes.

For $q = 1$, we have $d_{1,1}^1$ of rank 12 arriving at $E_{0,1}^1 \cong (\mathbb{Z}/2)^{16}$.

For $q \geq 3$, we have $d_{1,q}^1$ of rank 13 arriving at $E_{0,q}^1 \cong (\mathbb{Z}/2)^{12+2q}$.

For $q = 2$, we have $d_{1,2}^1$ of rank 13 arriving at $E_{0,2}^1 \cong (\mathbb{Z}/2)^{17}$. The only difficulty in seeing this is to compute the maps $H_q(\mathbb{Z}/2; \mathbb{Z}/2) \rightarrow H_q(\mathcal{D}_2; \mathbb{Z}/2)$ induced by the inclusions $f : \mathbb{Z}/2 \rightarrow \mathcal{D}_2$. For this task we take the resolutions of $\mathbb{Z}/2 \cong \langle t \mid t^2 = 1 \rangle$ and $\mathcal{D}_2 \cong \langle D, E \mid D^2 = E^2 = (DE)^2 = 1 \rangle$ proposed by [16] and compute the chain map induced by extending f to a ring homomorphism $f : \mathbb{Z}[\mathbb{Z}/2] \rightarrow \mathbb{Z}[\mathcal{D}_2]$. We can then apply the functor $- \otimes_{\mathbb{Z}[G]} \mathbb{Z}/2$ to this chain map (where $\mathbb{Z}/2$ is the trivial $\mathbb{Z}[G]$ -module for $G = \mathbb{Z}/2, \mathcal{D}_2$).

3.1.8. *The E^2 -page for $m = 13$ with $\mathbb{Z}/2$ -coefficients.* We obtain in the rows with $q \geq 2$ the E^2 -term concentrated in the column $p = 0$,

$$\begin{array}{c|c} q \geq 3 & (\mathbb{Z}/2)^{2q-1} \\ q = 2 & (\mathbb{Z}/2)^4, \end{array}$$

and in the rows $q = 0, q = 1$ it is concentrated in the columns $p = 0, 1, 2$:

$$\begin{array}{ccccccc}
 q = 1 & & (\mathbb{Z}/2)^4 & & \mathbb{Z}/2 & & 0 \\
 & & & \swarrow & & & \\
 & & & & d_{2,0}^2 & & \\
 & & & & & & \\
 q = 0 & & \mathbb{Z}/2 & & (\mathbb{Z}/2)^2 & & (\mathbb{Z}/2)^2.
 \end{array}$$

Computation of the differential $d_{2,0}^2$. The basis $\{(c, s, c', z), F\}$ of $E_{2,0}^2$ with \mathbb{Z} -coefficients induces a basis of $E_{2,0}^2$ with $\mathbb{Z}/2$ -coefficients. The Universal Coefficient Theorem gives us an isomorphism

$$0 \rightarrow H_1(\Gamma_\sigma; \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Z}/2 \xrightarrow{\cong} H_1(\Gamma_\sigma; \mathbb{Z}/2) \rightarrow 0 = \mathrm{Tor}_1^{\mathbb{Z}}(H_0(\Gamma_\sigma; \mathbb{Z}), \mathbb{Z}/2)$$

which we will use to transfer the elements $\bar{t}_\sigma \in H_1(\Gamma_\sigma; \mathbb{Z})$ computed in subsection 3.1.5 to $H_1(\Gamma_\sigma; \mathbb{Z}/2)$.

For $d_{2,0}^2((c, s, c', z))$ the computation is as follows.

As \bar{t}_c generates $H_1(\Gamma_c; \mathbb{Z}) \cong \mathbb{Z}/2$, it is transferred to the generator of

$$H_1(\Gamma_c; \mathbb{Z}/2) \cong \mathbb{Z}/2.$$

Since \bar{t}_s can be completed with a second element to a \mathbb{Z} -basis of $H_1(\Gamma_s; \mathbb{Z}) \cong \mathbb{Z}^2$, it is transferred to a nontrivial element of

$$H_1(\Gamma_s; \mathbb{Z}/2) \cong (\mathbb{Z}/2)^2.$$

The element \bar{t}_z vanishes because

$$H_1(\Gamma_z; \mathbb{Z}) \otimes \mathbb{Z}/2 \cong \mathbb{Z}/3 \otimes \mathbb{Z}/2 = 0.$$

The sum $\bar{t}_c + \bar{t}_s$ is quotiented to a nontrivial element on the E^2 -page because $H_1(\Gamma_s; \mathbb{Z}/2)$ is not hit by the d^1 -differential. So $d_{2,0}^2(\langle (c, s, c', z) \rangle) \cong \mathbb{Z}/2$.

For $d_{2,0}^2(F)$ the computation is as follows.

The 3-torsion vanishing when tensoring with $\mathbb{Z}/2$, the 3-torsion part $\bar{t}_x + \bar{t}_y$ of the sum makes no contribution to the image of d^2 .

The 2-torsion part, $\bar{t}_b + \bar{t}_a + \bar{t}_v + \bar{t}_u$, equals the image

$$d_{1,1}^1(\overline{t_{(b,c)}} + \overline{t_{(c'',a'')}} + \overline{t_{(v,f)}} + \overline{t_{(f,h)}} + \overline{t_{(h,u')}}),$$

where \bar{t}_σ , $\sigma \in \{b, a, v, u, (b, c), (c'', a''), (v, f), (f, h), (h, u')\}$ now stands for the generator of $H_1(\Gamma_\sigma; \mathbb{Z}/2) \cong \mathbb{Z}/2$. Hence it makes no contribution neither, and we obtain $d^2(F) = 0$. Thus $d_{2,0}^2$ has rank 1.


So the $E^3 = E^\infty$ -page gives us immediately

$$H_q(\Gamma, \mathbb{Z}/2) \cong \begin{cases} (\mathbb{Z}/2)^{2q-1}, & q \geq 3, \\ (\mathbb{Z}/2)^6, & q = 2, \\ (\mathbb{Z}/2)^5, & q = 1 \end{cases}$$

and we conclude

$$H_q(\Gamma, \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^3 \oplus (\mathbb{Z}/2)^2, & q = 1, \\ \mathbb{Z}^2 \oplus \mathbb{Z}/4 \oplus (\mathbb{Z}/3)^2 \oplus \mathbb{Z}/2, & q = 2, \\ (\mathbb{Z}/2)^q \oplus (\mathbb{Z}/3)^2, & q = 4k + 3, \quad k \geq 0, \\ (\mathbb{Z}/2)^q, & q = 4k + 4, \quad k \geq 0, \\ (\mathbb{Z}/2)^q, & q = 4k + 1, \quad k \geq 1, \\ (\mathbb{Z}/2)^q \oplus (\mathbb{Z}/3)^2, & q = 4k + 2, \quad k \geq 1. \end{cases}$$

3.2. $m = 5$. The fundamental domain for $\Gamma := \mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-5}]})$ looks as follows.



We will make use of the following matrices:

$$\begin{aligned} A &:= \pm \begin{pmatrix} & -1 \\ 1 & \end{pmatrix}, & B &:= \pm \begin{pmatrix} -\omega & 2 \\ 2 & \omega \end{pmatrix}, & M &:= \pm \begin{pmatrix} -\omega & 4 \\ 1 & \omega \end{pmatrix}, & S &:= \pm \begin{pmatrix} & -1 \\ 1 & 1 \end{pmatrix}, \\ U &:= \pm \begin{pmatrix} 1 & \omega \\ & 1 \end{pmatrix}, & V &:= \pm \begin{pmatrix} -\omega & 2-\omega \\ 2 & 2+\omega \end{pmatrix}, & W &:= \pm \begin{pmatrix} 6-\omega & 5+2\omega \\ 2\omega & \omega-4 \end{pmatrix}, \end{aligned}$$

which are subject to the relations $UMU^{-1} = A$, $UWS(UW)^{-1} = S$, $WABW^{-1} = MB$ and $S = ABV$. There are five orbits of vertices, with stabilizers

$$\begin{aligned} \Gamma_b &= \langle A, B | A^2 = B^2 = 1 \rangle \cong \mathcal{D}_2, \\ \Gamma_u &= \langle B, M | B^2 = M^2 = 1 \rangle \cong \mathcal{D}_2, \\ \Gamma_a &= \langle AB | AB^2 = 1 \rangle \cong \mathbb{Z}/2, \\ \Gamma_v &= \langle S | S^3 = 1 \rangle \cong \mathbb{Z}/3, \\ \Gamma_s &= \langle V, W | VW = WV \rangle \cong \mathbb{Z}^2 \end{aligned}$$

and identifications $UW \cdot a = a_1$, $V^{-1} \cdot a = a_2$, $S^2 \cdot a = a_2$, $U \cdot u = u_1$ and $UW \cdot v = v_1$. There are seven orbits of edges, with stabilizers

$$\begin{aligned} \Gamma_{(b,a)} &= \langle AB | AB^2 = 1 \rangle \cong \mathbb{Z}/2, \\ \Gamma_{(v,v_1)} &= \langle S | S^3 = 1 \rangle \cong \mathbb{Z}/3, \\ \Gamma_{(a_3,u)} &= \langle MB | MB^2 = 1 \rangle \cong \mathbb{Z}/2, \\ \Gamma_{(u,b)} &= \langle B | B^2 = 1 \rangle \cong \mathbb{Z}/2, \\ \Gamma_{(u_1,b)} &= \langle A | A^2 = 1 \rangle \cong \mathbb{Z}/2; \end{aligned}$$

(a, v) and (a, s) having the trivial stabilizer. There are three orbits of faces, with trivial stabilizers.

The above data gives the Γ -equivariant Euler characteristic of X :

$$\chi_{\Gamma}(X) = \frac{1}{4} + \frac{1}{4} + \frac{1}{2} + \frac{1}{3} - \frac{1}{2} - \frac{1}{3} - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} - 2 + 3 = 0,$$

in accordance with remark 17.

3.2.1. *The zeroth row of the E^1 -page.* This row identifies with the cellular chain complex of the quotient complex $\Gamma \backslash X$.

We obtain for the row $q = 0$ in the columns $p = 0, 1, 2$:

$$\mathbb{Z}^5 \xleftarrow{d_{1,0}^1} \mathbb{Z}^7 \xleftarrow{d_{2,0}^1} \mathbb{Z}^3$$

where 1 is the only elementary divisor of the differential matrices, with multiplicity four for $d_{1,0}^1$, and multiplicity two for $d_{2,0}^1$. The homology of $\Gamma \backslash X$ is generated in degree 1 by the loop represented by the edge (v, v_1) , and in degree 2 by the quotient of the face (a_2, s, a, v) , which is homeomorphic to a 2-sphere.

3.2.2. *Odd rows of the E^1 -page.* We start by investigating the morphism

$$\mathbb{Z}^2 \oplus \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^5 \xleftarrow{d_{1,1}^1} \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^4$$

and the morphism

$$\mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{q+4} \xleftarrow{d_{1,q}^1} \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^4$$

for $q \geq 3$. On the 3-torsion, $d_{1,q}^1$ is zero.

On the 2-torsion, $d_{1,q}^1$ it is given by the matrix

$$(d_{1,q}^1)_{(2)} = \begin{array}{c|cccc} & (b, a) & (a_3, u) & (u, b) & (u_1, b) \\ \hline a & 1 & -1 & 0 & 0 \\ b & -1 & 0 & 0 & 1 \\ \dots & \dots & \dots & \dots & \dots \\ b & -1 & 0 & 1 & 0 \\ u & 0 & 1 & -1 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ u & 0 & 1 & 0 & -1, \end{array}$$

where we fill in $\frac{q-1}{2}$ zero rows into each dotted line, except that in the columns with a “-1” both above and below the dots, we write “-1” into all entries of this column which are between the two “-1”’s.

Thus $d_{1,1}^1$ has rank 3 and $d_{1,q}^1$ has rank 4 for $q \geq 3$.

3.2.3. *Even rows of the E^1 -term.* There is a zero map arriving at $E_{0,2}^1 \cong \mathbb{Z} \oplus (\mathbb{Z}/2)^2$.

For $q \geq 4$, there is a zero map arriving at $E_{0,q}^1 \cong (\mathbb{Z}/2)^q$.

The rest of the E^1 -page are zeroes.

3.2.4. *The E^2 -page.*

The following periodic part appears in the columns $p = 0$ and $p = 1$:

$$\begin{array}{l|ll} q \geq 4 \text{ even} & (\mathbb{Z}/2)^q & 0 \\ q \geq 3 \text{ odd} & (\mathbb{Z}/2)^q \oplus \mathbb{Z}/3 & \mathbb{Z}/3; \end{array}$$

the lowest three rows are given as follows in the columns $p = 0, 1, 2$:

$$\begin{array}{rcccl} q = 2 & \mathbb{Z} \oplus (\mathbb{Z}/2)^2 & & 0 & 0 \\ q = 1 & \mathbb{Z}^2 \oplus (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3 & & \mathbb{Z}/2 \oplus \mathbb{Z}/3 & 0 \\ q = 0 & \mathbb{Z} & & \mathbb{Z} & \mathbb{Z} \end{array} \quad \begin{array}{l} \\ \\ \swarrow d^2 \end{array}$$

and the rest of the E^2 -page are zeroes.

Let us compute the only nontrivial d^2 -arrow. The generator of $E_{2,0}^2$ comes from the 2-cell (a_2, s, a, v) . Among its vertices, we have the identifications $S^2 \cdot a = a_2$ and $V^{-1} \cdot a = a_2$, where the matrices V of infinite order stabilizes the singular point s , and the matrix S of order three stabilizes the point v . The lift $1 \otimes_{(a_2, s, a, v)} 1$ of the generator of $E_{2,0}^2$ is mapped as follows in the

E^0 -page:

$$\begin{array}{ccc}
 (V^{-1}, 1) \otimes_s 1 - (V^{-1}, 1) \otimes_a 1 & \xleftarrow{1 \otimes \delta} & (V^{-1}, 1) \otimes_{(a,s)} 1 \\
 + (1, S^2) \otimes_v 1 - (1, S^2) \otimes_a 1 & & + (1, S^2) \otimes_{(a,v)} 1 \\
 & & \downarrow d_{\Theta} \otimes 1 \\
 & & 1 \otimes_{(a,s)} 1 - V^{-1} \otimes_{(a,s)} 1 \\
 & & + S^2 \otimes_{(a,v)} 1 - 1 \otimes_{(a,v)} 1 \xleftarrow{\quad} 1 \otimes_{(a_2, s, a, v)} 1
 \end{array}$$

It passes to

$$(\overline{V}, 2\overline{S}, \overline{AB}) \in \langle \overline{V}, \overline{W} \rangle \oplus \langle \overline{S} \mid 3\overline{S} = 0 \rangle \oplus (\mathbb{Z}/2)^2 \cong E_{0,1}^2,$$

which is of infinite order and has the following property: There is no element $\eta \in E_{0,1}^2$ with $k\eta = (\overline{V}, 2\overline{S}, \overline{AB})$ for an integer $k > 1$. So,

$$E_{0,1}^3 \cong \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3.$$

Thus the E^∞ -page gives us the following short exact sequences:

$$\begin{cases}
 0 \rightarrow (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}/3 \rightarrow 0 & q \geq 4 \text{ even,} \\
 0 \rightarrow \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0 & q \geq 3 \text{ odd,} \\
 0 \rightarrow \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \rightarrow H_2(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}/3 \oplus \mathbb{Z}/2 \rightarrow 0, \\
 0 \rightarrow \mathbb{Z} \oplus \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^2 \rightarrow H_1(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z} \rightarrow 0.
 \end{cases}$$

The result

$$H_q(\Gamma; \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^2 \oplus \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^2 & q = 1 \\ \mathbb{Z} \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2 & q = 2 \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^q & q \geq 3 \end{cases}$$

is obtained after resolving the ambiguity of the extension $H_2(\Gamma, \mathbb{Z})$ by a reflection like the one on [16, page 587], for which we have to recompute the spectral sequence with $\mathbb{Z}/2$ -coefficients, giving us

$$H_q(\Gamma; \mathbb{Z}/2) \cong \begin{cases} (\mathbb{Z}/2)^4 & q = 1 \\ (\mathbb{Z}/2)^5 & q = 2 \\ (\mathbb{Z}/2)^{2q-1} & q \geq 3. \end{cases}$$

Remark 20. There are the following checks of the computations we made. Let us look at the low term short exact sequence

$$0 \rightarrow E_{0,1}^\infty \rightarrow \Gamma^{\mathrm{ab}} \rightarrow E_{1,0}^\infty \rightarrow 0$$

of the spectral sequence. We have $E_{1,0}^\infty = H_1(\Gamma \backslash X) = (\pi_1(\Gamma \backslash X))^{ab}$, and the projection is the abelianization of the map $\Gamma \rightarrow \pi_1(\Gamma \backslash X)$ given as follows. Choose a fixed base point $x \in X$. For every $\gamma \in \Gamma$, choose a continuous path in X from x to γx . This gives a well-defined loop in $\Gamma \backslash X$ since X is contractible.


As Flöge shows, an inspection of the complex X and the associated stabilizer group yields, together with [1, theorem 4.5], a presentation of Γ by means of generators and relations. (It should be even easier to deduce only the abelianization of Γ , however we didn't figure out the details for this.) So suppose Γ^{ab} is known. The group $E_{0,1}^\infty = E_{0,1}^3$ is then the kernel of the projection $\Gamma^{\mathrm{ab}} \rightarrow \pi_1(\Gamma \backslash X)$.

For $m = 5$, this check looks as follows. The abelianization is the abelian group $\Gamma^{\mathrm{ab}} \cong \langle \overline{A}, \overline{B}, \overline{S}, \overline{U} : 2\overline{A} = 0, 2\overline{B} = 0, 3\overline{S} = 0 \rangle$. Since the fundamental group of the quotient is free, only the parabolic elements U and V can define

nontrivial loops in the quotient. The element U generates a nontrivial loop, while V generates a trivial loop.

So it follows that $E_{0,1}^\infty \cong \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3$, generated by $\bar{V}, \bar{A}, \bar{B}$ and \bar{S} . This is consistent with the computation above, involving the detailed analysis of the d^2 -differential.

3.3. $m = 10$. The fundamental domain for $\Gamma := \mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-10}]})$ looks as follows.



We will use the following definitions:

$$\begin{aligned}
 A &:= \pm \begin{pmatrix} & -1 \\ 1 & \end{pmatrix}, & B &:= \pm \begin{pmatrix} -\omega & 3 \\ 3 & \omega \end{pmatrix}, & C &:= \pm \begin{pmatrix} -1-\omega & 4-\omega \\ 2 & 1+\omega \end{pmatrix}, \\
 D &:= \pm \begin{pmatrix} \omega-1 & -4 \\ 3 & 1+\omega \end{pmatrix}, & L &:= \pm \begin{pmatrix} \omega & 3 \\ 3 & -\omega \end{pmatrix}, & R &:= \pm \begin{pmatrix} 5+\omega & 2\omega-3 \\ \omega-3 & -4-\omega \end{pmatrix}, \\
 S &:= \pm \begin{pmatrix} & -1 \\ 1 & 1 \end{pmatrix}, & U &:= \pm \begin{pmatrix} 1 & \omega \\ & 1 \end{pmatrix}, & V &:= \pm \begin{pmatrix} 1-\omega & 5 \\ 2 & 1+\omega \end{pmatrix}, \\
 W &:= \pm \begin{pmatrix} 11 & 5\omega \\ 2\omega & -9 \end{pmatrix}, & Y &:= \pm \begin{pmatrix} \omega-2 & -5 \\ 3 & 2+\omega \end{pmatrix}.
 \end{aligned}$$

The matrix U acts as a vertical translation by $-\omega$ on this fundamental domain. There are nine orbits of vertices, labelled $a, b, r, u, v, w, x, y, s$. We have the following identifications: $UWa = a_1$, $Wa = a_2$, $Va = a_3$; $S^{-1}v = v_1$, $U^{-1}Dv = v_2$; $Dw = w_1$, $U^{-1}Dw = w_2$; $Db = b_1$, $Cb = b_2$; $Dr = r_1$; $UWx = x_1$. The stabilizers of the vertex orbit representatives are

$$\begin{aligned}
 \Gamma_a &= \Gamma_b = \langle R \mid R^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_w &= \langle S \mid S^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_y &= \langle A, L \mid A^2 = L^2 = (AL)^2 = 1 \rangle \cong \mathcal{D}_2, \\
 \Gamma_u &= \langle A, B \mid A^2 = B^2 = (AB)^2 = 1 \rangle \cong \mathcal{D}_2, \\
 \Gamma_r &= \langle C \mid C^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_v &= \langle AB \mid (AB)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_x &= \langle B \mid B^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_s &= \langle V, W \mid VW = WV \rangle \cong \mathbb{Z}^2.
 \end{aligned}$$

There are fifteen orbits of edges, labelled $(b, v), (r, w), (b, r), (v, w), (a_2, w_2), (y, r_1), (x, a), (u, y), (a, b), (u, v), (a, s), (w, b_1), (r, v_2), (y, x_1), (x, u)$.

Amongst their stabilizers only

$$\begin{aligned}
 \Gamma_{(a_2, w_2)} &= \Gamma_{a_2} = W^{-1}\Gamma_a W = \langle W^{-1}RW \mid (W^{-1}RW)^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(a, b)} &= \Gamma_a = \Gamma_b = \langle R \mid R^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(w, b_1)} &= \Gamma_{b_1} = \Gamma_w = \langle S \mid S^3 = 1 \rangle \cong \mathbb{Z}/3, \\
 \Gamma_{(y, r_1)} &= \Gamma_{r_1} = D\Gamma_r D^{-1} = \langle AL = DCD^{-1} \mid (DCD^{-1})^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(u, v)} &= \Gamma_v = \langle AB \mid (AB)^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(r, v_2)} &= \Gamma_{v_2} = \Gamma_r = \langle C \mid C^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(y, x_1)} &= \Gamma_{x_1} = UW\Gamma_x(UW)^{-1} = \langle L \mid L^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(x, u)} &= \Gamma_x = \langle B \mid B^2 = 1 \rangle \cong \mathbb{Z}/2, \\
 \Gamma_{(u, y)} &= \langle A \mid A^2 = 1 \rangle \cong \mathbb{Z}/2
 \end{aligned}$$

are nontrivial. Furthermore, there are seven orbits of faces, with trivial stabilizers.

With the above information on the isomorphy types of the cell stabilizers, we get the Γ -equivariant Euler characteristic of X :

$$\chi_\Gamma(X) = 3 \cdot \frac{1}{3} + 2 \cdot \frac{1}{4} + 3 \cdot \frac{1}{2} - 3 \cdot \frac{1}{3} - 6 \cdot \frac{1}{2} - 6 + 7 = 0,$$

in accordance with remark 17.

3.3.1. *The row $q = 0$ in the E^1 -page for $m = 10$.*

We obtain for the row $q = 0$ in the columns $p = 0, 1, 2$:

$$\mathbb{Z}^9 \xleftarrow{d_{1,0}^1} \mathbb{Z}^{15} \xleftarrow{d_{2,0}^1} \mathbb{Z}^7,$$

where 1 is the only elementary divisor of the differential matrices, with multiplicity eight for $d_{1,0}^1$, and multiplicity five for $d_{2,0}^1$. The rest of this row are zeroes.

3.3.2. *Odd rows of the E^1 -term.*

For odd q , the morphism

$$\bigoplus_{\sigma \in \Gamma \backslash X^0} \mathrm{H}_q(\Gamma_\sigma) \xleftarrow{d_{1,q}^1} \bigoplus_{\sigma \in \Gamma \backslash X^1} \mathrm{H}_q(\Gamma_\sigma)$$

is for $q \geq 3$ of the form

$$(\mathbb{Z}/3)^3 \oplus (\mathbb{Z}/2)^{q+6} \leftarrow (\mathbb{Z}/3)^3 \oplus (\mathbb{Z}/2)^6.$$

For $q = 1$, we have to add $\mathrm{H}_1(\Gamma_s) \cong \mathbb{Z}^2$ on the target side of the morphism $d_{1,q}^1$, but the incoming torsion must reach it trivially.

On the 3-primary part, $d_{1,q}^1$ is given by the matrix

$$(d_{1,q}^1)_{(3)} = \begin{array}{c|ccc} & (a, b) & (Db, w) & (Wa, U^{-1}Dw) \\ \hline a & -1 & 0 & -1 \\ w & 0 & 1 & 1 \\ b & 1 & -1 & 0. \end{array}$$

This matrix has rank 2, so its image is isomorphic to $(\mathbb{Z}/3)^2$ and its kernel is of type $\mathbb{Z}/3$.

On the 2-primary part, $d_{1,q}^1$ is for odd q given by the matrix

$$(d_{1,q}^1)_{(2)} = \begin{array}{c|cccccc} & (y, r_1) & (u, v) & (r, v_2) & (y, x_1) & (x, u) & (u, y) \\ \hline u & 0 & -1 & 0 & 0 & 0 & -1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ u & 0 & -1 & 0 & 0 & 1 & 0 \\ y & -1 & 0 & 0 & 0 & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ y & -1 & 0 & 0 & -1 & 0 & 0 \\ x & 0 & 0 & 0 & 1 & -1 & 0 \\ r & 1 & 0 & -1 & 0 & 0 & 0 \\ v & 0 & 1 & 1 & 0 & 0 & 0, \end{array}$$

where we fill in $\frac{q-1}{2}$ zero rows into each of the two dotted lines, except that in the columns with a “-1” both above and below the dots, we write “-1” into all entries of this column which are between the two “-1”’s. This is due to the diagonal map

$$(\mathbb{Z}/2)^{\frac{q+3}{2}} \cong H_q(\mathcal{D}_2) \leftarrow H_q(\mathbb{Z}/2) \cong \mathbb{Z}/2, \quad \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \leftarrow 1,$$

which is induced by the inclusion $\mathcal{D}_2 \leftarrow \mathbb{Z}/2$ hitting the product of the two fixed generators of \mathcal{D}_2 . The above matrix $(d_{1,q}^1)_{(2)}$ has rank 5 for $q = 1$, and full rank 6 for $q \geq 3$.

3.3.3. The rows with q even.

There is a zero map arriving for q greater than 2 at $\bigoplus_{\sigma \in \Gamma \setminus X^0} H_q(\Gamma_\sigma) \cong (\mathbb{Z}/2)^q$, and for $q = 2$ at $\bigoplus_{\sigma \in \Gamma \setminus X^0} H_2(\Gamma_\sigma) \cong \mathbb{Z} \oplus (\mathbb{Z}/2)^2$.

3.3.4. The E^2 -page for $m = 10$:

The following periodic part appears in the columns $p = 0$ and $p = 1$:

$$\begin{array}{l|ll} q \geq 4 \text{ even} & (\mathbb{Z}/2)^q & 0 \\ q \geq 3 \text{ odd} & (\mathbb{Z}/2)^q \oplus \mathbb{Z}/3 & \mathbb{Z}/3; \end{array}$$

the lowest three rows are as follows in the columns $p = 0, 1, 2$:

$$\begin{array}{rcccl} q = 2 & \mathbb{Z} \oplus (\mathbb{Z}/2)^2 & 0 & 0 & \\ & & & & \\ q = 1 & \mathbb{Z}^2 \oplus (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3 & \mathbb{Z}/2 \oplus \mathbb{Z}/3 & 0 & \\ & & \swarrow d^2 & & \\ q = 0 & \mathbb{Z} & \mathbb{Z}^2 & \mathbb{Z}^2 & \end{array}$$

and the rest of the E^2 -page are zeroes.

3.3.5. Computation of the differential d^2 .

The generators of the abelian group $E_{2,0}^2 \cong \mathbb{Z}^2$ are represented by the 2-cell (a, s, a_3, x) and the union of two 2-cells (v_1, b_2, r, b, v, w) , whose quotients by Γ are homeomorphic to 2-spheres. On the vertices of (a, s, a_3, x) , we have the identifications $B \cdot a = a_3$ and $V \cdot a = a_3$, where the matrix B fixes x and the matrix V fixes s . For (v_1, b_2, r, b, v, w) , we have the identifications of vertices $Cb = b_2$, $Cr = r$, $S^2v = v_1$ and $S^2w = w$; and we pay particular

attention to the matrix $CR = S^2AB$ identifying the edge $(b, v) \cong (b_2, v_1)$. Thus the only nontrivial d^2 -arrow is induced by

$$\delta((a, s, a_3, x)) = (a, s) + V \cdot (s, a) + B \cdot (a, x) + (x, a)$$

and

$$\delta((v_1, b_2, r, b, v, w)) = (b, r) - C \cdot (b, r) + CR \cdot (b, v) + S^2 \cdot (v, w) - (v, w) - (b, v).$$

The lift $1 \otimes_{(v_1, b_2, r, b, v, w)} 1$ of the generator obtained from (v_1, b_2, r, b, v, w) is mapped as follows:

$$\begin{array}{ccc} (C, 1) \otimes_r 1 - (C, 1) \otimes_b 1 & & (C, 1) \otimes_{(b, r)} 1 \\ + (1, CR) \otimes_v 1 - (1, CR) \otimes_b 1 & \xleftarrow{1 \otimes \delta} & + (1, CR) \otimes_{(b, v)} 1 \\ + (1, S^2) \otimes_w 1 - (1, S^2) \otimes_v 1 & & + (1, S^2) \otimes_{(v, w)} 1 \\ & & \downarrow d_{\Theta} \otimes 1 \\ & & 1 \otimes_{(b, r)} 1 - C \otimes_{(b, r)} 1 \\ & & + CR \otimes_{(b, v)} 1 - 1 \otimes_{(b, v)} 1 \xleftarrow{1 \otimes_{(v_1, b_2, r, b, v, w)} 1} \\ & & + S^2 \otimes_{(v, w)} 1 - 1 \otimes_{(v, w)} 1 \end{array}$$

We attribute the symbols t_σ to the part of this sum lying in $\Theta_1 \otimes_\sigma \mathbb{Z}$.

$$\begin{aligned} t_r &:= (C, 1) \otimes_r 1, \\ t_b &:= -(1, CR) \otimes_b 1 - (C, 1) \otimes_b 1, \\ t_w &:= (1, S^2) \otimes_w 1, \\ t_v &:= (1, CR) \otimes_v 1 - (1, S^2) \otimes_v 1. \end{aligned}$$

With the formula in our corollary 12, we find the classes \overline{t}_σ in $H_1(\Theta_* \otimes_\sigma \mathbb{Z})$ in the following way.

$$\begin{aligned} \overline{t}_r &= \overline{C} \in \langle \overline{C} | 2\overline{C} = 0 \rangle = \Gamma_r^{ab} \cong H_1(\Gamma_r, \mathbb{Z}), \\ \overline{t}_b &= 2\overline{R} \in \langle \overline{R} | 3\overline{R} = 0 \rangle = \Gamma_b^{ab} \cong H_1(\Gamma_b, \mathbb{Z}), \\ \overline{t}_w &= 2\overline{S} \in \langle \overline{S} | 3\overline{S} = 0 \rangle = \Gamma_w^{ab} \cong H_1(\Gamma_w, \mathbb{Z}), \\ \overline{t}_v &= \overline{AB} \in \langle \overline{AB} | 2\overline{AB} = 0 \rangle = \Gamma_v^{ab} \cong H_1(\Gamma_v, \mathbb{Z}). \end{aligned}$$

As $E_{-1,1}^1 = 0$, the whole of $E_{0,1}^1$ is in the kernel of d^1 . So for the passage to E^2 , we just have to look at the image of d^1 .

The chain $\overline{t}_v + \overline{t}_r$ equals the image $d_{1,1}^1(-r, v_2)$, so it passes to the zero element in the quotient $E_{0,1}^2$. Meanwhile, the cycle $\overline{t}_b + \overline{t}_w$ is a representative of the generator of the 3-torsion in $E_{0,1}^2$, as we see from the matrix for the 3-primary part of $d_{1,1}^1$. Thus, $d_{2,0}^2(\langle (v_1, b_2, r, b, v, w) \rangle) \cong \mathbb{Z}/3$.

The lift $1 \otimes_{(a, s, a_3, x)} 1$ of the generator obtained from (a, s, a_3, x) is mapped

$$\begin{array}{ccc} (V, 1) \otimes_s 1 - (V, 1) \otimes_a 1 & & (V, 1) \otimes_{(a, s)} 1 \\ + (1, B) \otimes_x 1 - (1, B) \otimes_a 1 & \xleftarrow{1 \otimes \delta} & + (1, B) \otimes_{(a, x)} 1 \\ & & \downarrow d_{\Theta} \otimes 1 \\ & & 1 \otimes_{(a, s)} 1 - V \otimes_{(a, s)} 1 \\ & & + B \otimes_{(a, x)} 1 - 1 \otimes_{(a, x)} 1 \xleftarrow{1 \otimes_{(a, s, a_3, x)} 1} \end{array}$$

We attribute the symbols t_σ to the part of this sum lying in $\Theta_1 \otimes_\sigma \mathbb{Z}$,

$$\begin{aligned} t_s &:= (V, 1) \otimes_s 1, \\ t_x &:= (1, B) \otimes_x 1, \\ t_a &:= -(V, 1) \otimes_a 1 - (1, B) \otimes_a 1. \end{aligned}$$

We find the class $\overline{t}_s = -\overline{V} \in \langle \overline{V}, \overline{W} \rangle = \Gamma_s^{ab} \cong H_1(\Gamma_s, \mathbb{Z}) \cong \mathbb{Z}^2$, which is a generator of the free part of $E_{0,1}^1$. It can not be the image of a torsion

element from $E_{1,1}^1 = (\mathbb{Z}/3)^3 \oplus (\mathbb{Z}/2)^2$. Therefore, it is preserved when passing from $E_{0,1}^1$ to $E_{0,1}^2$. The cycles $\overline{t_x}$ and $\overline{t_a}$ are torsion, so the fact that $\overline{t_s}$ is a generator of the free part determines that the image $d_{2,0}^2(\langle (a, s, a_3, x) \rangle)$ is of infinite order and has the following property: There is no element $\eta \in E_{0,1}^2 \cong \mathbb{Z}^2 \oplus \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^2$ with $k\eta = d_{2,0}^2(\langle (a, s, a_3, x) \rangle)$ for an integer $k > 1$. Together with the isomorphism $d_{2,0}^2(\langle (v_1, b_2, r, b, v, w) \rangle) \cong \mathbb{Z}/3$, we obtain

$$E_{0,1}^3 \cong \mathbb{Z} \oplus (\mathbb{Z}/2)^2.$$

Thus the E^∞ -page gives the following short exact sequences:

$$\begin{cases} 0 \rightarrow (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}/3 \rightarrow 0, & \text{for } q \geq 4 \text{ even,} \\ 0 \rightarrow \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^q \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0, & \text{for } q \geq 3 \text{ odd,} \\ 0 \rightarrow \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \rightarrow H_2(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z} \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2 \rightarrow 0, \\ 0 \rightarrow \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \rightarrow H_1(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}^2 \rightarrow 0. \end{cases}$$

From here, we easily see the results,

$$H_q(\Gamma, \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^3 \oplus (\mathbb{Z}/2)^2, & q = 1, \\ \mathbb{Z}^2 \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2, & q = 2, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^q, & q \geq 3; \end{cases}$$

except for the ambiguity in the 3-torsion and the 2-torsion of the short exact sequence for $H_2(\Gamma; \mathbb{Z})$. To resolve it, we will compute homology with $\mathbb{Z}/2$ -coefficients,

$$H_q(\Gamma, \mathbb{Z}/2) \cong \begin{cases} (\mathbb{Z}/2)^{2q-1}, & q \geq 3 \\ (\mathbb{Z}/2)^6, & q = 2, \\ (\mathbb{Z}/2)^5, & q = 1, \end{cases}$$

and then use the Universal Coefficient Theorem in the form

$$H_2(\Gamma, \mathbb{Z}/2) \cong H_2(\Gamma; \mathbb{Z}) \otimes (\mathbb{Z}/2) \oplus \mathrm{Tor}_1^{\mathbb{Z}}(H_1(\Gamma; \mathbb{Z}), \mathbb{Z}/2).$$

and we do the same for $\mathbb{Z}/3$ -coefficients.


Remark 21. For $m = 10$, the check introduced in remark 20 takes the following form. The abelianization is the group

$$\Gamma^{\mathrm{ab}} \cong \langle \overline{A}, \overline{B}, \overline{D}, \overline{U}, \overline{W} : 2\overline{A} = 2\overline{B} = 0 \rangle.$$

The parabolic elements are U and W . The element U generates nontrivial loops in the quotient, while W and D generate trivial loops.

So it follows that $E_{0,1}^\infty = \mathbb{Z} \oplus (\mathbb{Z}/2)^2$, generated by $\overline{W}, \overline{A}$ and \overline{B} . This is consistent with the computation above.

3.4. $m = 6$. The fundamental domain for $\Gamma := \mathrm{PSL}_2(\mathcal{O}_{\mathbb{Q}[\sqrt{-6}]})$ looks as follows.



The matrix $U := \pm \begin{pmatrix} 1 & \omega \\ & 1 \end{pmatrix}$ performs a vertical translation by $-\omega$ of the fundamental domain for Γ . The following matrices occur in the cell stabilizers.

$$\begin{aligned} A &:= \pm \begin{pmatrix} & -1 \\ 1 & \end{pmatrix}, & B &:= \pm \begin{pmatrix} -1-\omega & 2-\omega \\ 2 & 1+\omega \end{pmatrix}, & R &:= \pm \begin{pmatrix} -\omega & 5-\omega \\ 1 & 1+\omega \end{pmatrix}, \\ S &:= \pm \begin{pmatrix} & -1 \\ 1 & 1 \end{pmatrix}, & V &:= \pm \begin{pmatrix} 1-\omega & 3 \\ 3 & 1+\omega \end{pmatrix}, & W &:= \pm \begin{pmatrix} 7 & 3\omega \\ 2\omega & -5 \end{pmatrix}. \end{aligned}$$

There are five orbits of vertices, labelled b, a, u, v, s , with stabilizers

$$\begin{aligned} \Gamma_u &= \langle B, S \mid B^2 = S^3 = (BS)^3 = 1 \rangle \cong \mathcal{A}_4, \\ \Gamma_v &= \langle B, R \mid B^2 = R^3 = (BR)^3 = 1 \rangle \cong \mathcal{A}_4, \\ \Gamma_{v_1} &= \langle UBU^{-1}, S \mid (UBU^{-1})^2 = S^3 = (UBU^{-1}S)^3 = 1 \rangle \cong \mathcal{A}_4, \\ \Gamma_a &= \langle SB \mid (SB)^3 = 1 \rangle \cong \mathbb{Z}/3, \\ \Gamma_{a_2} &= \langle RB \mid (RB)^3 = 1 \rangle \cong \mathbb{Z}/3, \\ \Gamma_b &= \langle A \mid A^2 = 1 \rangle \cong \mathbb{Z}/2, \\ \Gamma_s &= \langle V, W \mid VW = WV \rangle \cong \mathbb{Z}^2, \end{aligned}$$

and identifications $UW \cdot a = a_1$, $W \cdot a = a_2$, $V \cdot a = a_3$, $A \cdot a = a_3$, $UW \cdot b = b_1$ and $U \cdot v = v_1$. There are seven orbits of edges, labelled (b, a) , (a, s) , (a, u) , (u, v) , (a_2, v) , (b, b_1) and (u, v_1) , amongst whose stabilizers only

$$\begin{aligned} \Gamma_{(a_2, v)} &= \langle RB \mid (RB)^3 = 1 \rangle = \Gamma_{a_2} \cong \mathbb{Z}/3, \\ \Gamma_{(u, v_1)} &= \langle S \mid S^3 = 1 \rangle \cong \mathbb{Z}/3, \\ \Gamma_{(a, u)} &= \langle SB \mid (SB)^3 = 1 \rangle = \Gamma_a \cong \mathbb{Z}/3, \\ \Gamma_{(u, v)} &= \langle B \mid B^2 = 1 \rangle \cong \mathbb{Z}/2, \\ \Gamma_{(b, b_1)} &= \langle A \mid A^2 = 1 \rangle = \Gamma_b = \Gamma_{b_1} \cong \mathbb{Z}/2 \end{aligned}$$

are nontrivial; and three orbits of faces with trivial stabilizers. The above data gives the Γ -equivariant Euler characteristic of X :

$$\chi_\Gamma(X) = \frac{1}{12} + \frac{1}{12} + \frac{1}{3} + \frac{1}{2} - 1 - 1 - \frac{1}{3} - \frac{1}{3} - \frac{1}{3} - \frac{1}{2} - \frac{1}{2} + 3 = 0,$$

in accordance with remark 17.

3.4.1. Zeroth row of the E^1 -term.

We obtain in the columns $p = 0, 1, 2$:

$$\mathbb{Z}^5 \xleftarrow{d_{1,0}^1} \mathbb{Z}^7 \xleftarrow{d_{2,0}^1} \mathbb{Z}^3$$

where 1 is the only occurring elementary divisor of the differential matrices, with multiplicity four for $d_{1,0}^1$, and multiplicity two for $d_{2,0}^1$. The homology of this sequence is generated by the cycle (b, b_1) in degree one and by the face (a, s, a_3, b) in degree two.

3.4.2. Even rows of the E^1 -term.

The even rows are the zero map to $E_{0,2}^1 \cong \mathbb{Z} \oplus (\mathbb{Z}/2)^2$, and to $E_{0,q}^1 \cong (\mathrm{H}_q(\mathcal{A}_4))^2$ for the degree $q \geq 4$.

3.4.3. Odd rows of the E^1 -term.

The map $d_{1,q}^1$ is on the 2-primary part induced by the inclusion of $\Gamma_{(u,v)} \cong \mathbb{Z}/2$ into Γ_v and Γ_u which are of isomorphism type \mathcal{A}_4 . By [16, lemma 4.5(2)], every inclusion of $\mathbb{Z}/2$ into \mathcal{A}_4 induces injections on homology in degrees greater than 1, and is zero on H_1 . So the morphism

$$\mathbb{Z}^2 \oplus \mathbb{Z}/2 \oplus (\mathbb{Z}/3)^3 \xleftarrow{d_{1,1}^1} (\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^3$$

has $\mathbb{Z}/2$ -rank 0 on the 2-primary part, and

$$\mathbb{Z}/3 \oplus \mathbb{Z}/2 \oplus (\mathrm{H}_q(\mathcal{A}_4))^2 \xleftarrow{d_{1,q}^1} (\mathbb{Z}/2)^2 \oplus (\mathbb{Z}/3)^3$$


in the odd rows of degree $q \geq 3$ has $\mathbb{Z}/2$ -rank 1 on the 2-primary part.

On the 3-primary part, $d_{1,q}^1$ is for all odd q given by the following rank 2 matrix.

$$(d_{1,q}^1)_{(3)} = \begin{array}{c|ccc} & (a,u) & (a_2,v) & (u,v_1) \\ \hline a & -1 & -1 & 0 \\ u & 1 & 0 & -1 \\ v & 0 & 1 & 1. \end{array}$$

In order to determine it, we make use of the following facts.

First, by [16, lemma 4.5], each of the occurring group inclusions induces an injection in homology. So we have to determine the relative positions of the images coming from the edges in each direct summand over the points. In order to find out if cancelling occurs between terms with positive and negative signs, let us look at the following diagram. The symbol ΔW denotes the isomorphism given by conjugation with W , δ denotes an inner automorphism, ι denotes any canonical inclusion, and the arrows emanating from $\mathbb{Z}/3$ are labeled with the image of the canonical generator.



Applying homology H_q for odd q and taking into account that the fact that inner automorphisms act trivially on homology, we get a similar slightly smaller commutative diagram. One can then unambiguously identify all occurring groups $\mathrm{H}_q(\mathbb{Z}/3) \cong \mathbb{Z}/3$ and its images in $\mathrm{H}_q(\mathcal{A}_4)$ with the “abstract” $\mathrm{H}_q(\mathbb{Z}/3) \cong \mathbb{Z}/3$ in the middle. This gives a basis for the 3-primary parts of the source and a subspace of the image. In this basis, the 3-primary map

is given by the following matrix, followed by an injection which does not influence the homology.

3.4.4. The E^2 -page.

The following periodic part appears in the columns $p = 0$ and $p = 1$:

$$\begin{array}{l|ll} q = 6k + 2, q \geq 8 & (\mathbb{Z}/2)^{2k+2} & 0 \\ q = 6k + 1, q \geq 7 & (\mathbb{Z}/2)^{2k} \oplus \mathbb{Z}/3 & \mathbb{Z}/3 \oplus \mathbb{Z}/2 \\ q = 6k + 6 & (\mathbb{Z}/2)^{2k} & 0 \\ q = 6k + 5 & (\mathbb{Z}/2)^{2k+4} \oplus \mathbb{Z}/3 & \mathbb{Z}/3 \oplus \mathbb{Z}/2 \\ q = 6k + 4 & (\mathbb{Z}/2)^{2k} & 0 \\ q = 6k + 3 & (\mathbb{Z}/2)^{2k+2} \oplus \mathbb{Z}/3 & \mathbb{Z}/3 \oplus \mathbb{Z}/2; \end{array}$$

the lowest three rows are in the columns $p = 0, 1, 2$:

$$\begin{array}{cccc} q = 2 & \mathbb{Z} \oplus (\mathbb{Z}/2)^2 & 0 & 0 \\ q = 1 & \mathbb{Z}^2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/3 & (\mathbb{Z}/2)^2 \oplus \mathbb{Z}/3 & 0 \\ q = 0 & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \end{array}$$

and the rest of the E^2 -page are zeroes.

3.4.5. The $E^3 = E^\infty$ -term. For the calculation of the d^2 -differential, we have

$$\begin{aligned} \delta(a, s, a_3, b) &= (a_3, s) + (s, a) + (a, b) + (b, a_3) \\ &= (V \cdot a, s) + (s, a) + (a, b) + (b, A \cdot a) \\ &= V(a, s) - (a, s) - (b, a) + A(b, a), \\ 1 \otimes \delta(1 \otimes_{(a,s,a_3,b)} 1) &= 1 \otimes_{V(a,s)} 1 - 1 \otimes_{(a,s)} 1 - 1 \otimes_{(b,a)} 1 + 1 \otimes_{A(b,a)} 1 \\ &= (V - 1) \otimes_{(a,s)} 1 + (A - 1) \otimes_{(b,a)} 1 \\ &= (d_\Theta \otimes 1) \left((1, V) \otimes_{(a,s)} 1 + (1, A) \otimes_{(b,a)} 1 \right) \\ &= (d_\Theta \otimes 1) \left([V] \otimes_{(a,s)} 1 + [A] \otimes_{(b,a)} 1 \right). \end{aligned}$$

We then get

$$1 \otimes \delta \left([V] \otimes_{(a,s)} 1 + [A] \otimes_{(b,a)} 1 \right) = [V] \otimes_s 1 - [V] \otimes_a 1 + [A] \otimes_a 1 - [A] \otimes_b 1.$$

As $[V] \otimes_s 1$ and $[W] \otimes_s 1$ represent the generators of the torsion-free part of $E_{0,1}^2 \cong \mathbb{Z}^2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/3$, we see that the above computed element of $E_{0,1}^0$ represents an element $\nu \in E_{0,1}^2$ of infinite order with the following property: There is no element $\eta \in E_{0,1}^2$ with $k\eta = \nu$ for an integer $k > 1$. So, $E_{0,1}^3 \cong \mathbb{Z} \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2$ and $E_{2,0}^3 = 0$.

3.4.6. The short exact sequences.

We thus obtain for integer homology the following short exact sequences:

$$\left\{ \begin{array}{ll} 0 \rightarrow (\mathbb{Z}/2)^{2k+2} \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}/3 \oplus \mathbb{Z}/2 \rightarrow 0, & q = 6k + 2, q \geq 8 \\ 0 \rightarrow (\mathbb{Z}/2)^{2k} \oplus \mathbb{Z}/3 \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0, & q = 6k + 1, q \geq 7 \\ 0 \rightarrow (\mathbb{Z}/2)^{2k} \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}/3 \oplus \mathbb{Z}/2 \rightarrow 0, & q = 6k + 6, \\ 0 \rightarrow (\mathbb{Z}/2)^{2k+4} \oplus \mathbb{Z}/3 \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0, & q = 6k + 5, \\ 0 \rightarrow (\mathbb{Z}/2)^{2k} \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}/3 \oplus \mathbb{Z}/2 \rightarrow 0, & q = 6k + 4, \\ 0 \rightarrow (\mathbb{Z}/2)^{2k+2} \oplus \mathbb{Z}/3 \rightarrow H_q(\Gamma; \mathbb{Z}) \rightarrow 0, & q = 6k + 3, \\ 0 \rightarrow \mathbb{Z} \oplus (\mathbb{Z}/2)^2 \rightarrow H_2(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^2 \rightarrow 0, & \\ 0 \rightarrow \mathbb{Z} \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2 \rightarrow H_1(\Gamma; \mathbb{Z}) \rightarrow \mathbb{Z} \rightarrow 0. & \end{array} \right.$$

Summarizing, we obtain:

$$H_q(\Gamma; \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^2 \oplus \mathbb{Z}/3 \oplus \mathbb{Z}/2, & q = 1, \\ \mathbb{Z} \oplus \mathbb{Z}/4 \oplus \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^2, & q = 2, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+2}, & q = 6k + 3, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+1}, & q = 6k + 4, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+4}, & q = 6k + 5, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+1}, & q = 6k + 6, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k}, & q = 6k + 1, q \geq 7, \\ \mathbb{Z}/3 \oplus (\mathbb{Z}/2)^{2k+3}, & q = 6k + 2, q \geq 8. \end{cases}$$

We easily see these results except for the ambiguity in the 3-torsion of the short exact sequence for $H_2(\Gamma; \mathbb{Z})$ and in the 2-torsion for all even degrees. To resolve it, we compute homology with $\mathbb{Z}/2$ -coefficients,

$$H_q(\Gamma; \mathbb{Z}/2) \cong \begin{cases} (\mathbb{Z}/2)^3, & q = 1, \\ (\mathbb{Z}/2)^5, & q = 2, \\ (\mathbb{Z}/2)^{4k+5}, & q = 6k + 3, \\ (\mathbb{Z}/2)^{4k+3}, & q = 6k + 4, \\ (\mathbb{Z}/2)^{4k+5}, & q = 6k + 5, \\ (\mathbb{Z}/2)^{4k+5}, & q = 6k + 6, \\ (\mathbb{Z}/2)^{4k-2}, & q = 6k + 1, q \geq 7 \\ (\mathbb{Z}/2)^{4k+3}, & q = 6k + 2, q \geq 8. \end{cases}$$

and then use the Universal Coefficient Theorem in the form

$$H_{q+1}(\Gamma; \mathbb{Z}/2) \cong H_{q+1}(\Gamma; \mathbb{Z}) \otimes (\mathbb{Z}/2) \oplus \mathrm{Tor}_1^{\mathbb{Z}}(H_q(\Gamma; \mathbb{Z}), \mathbb{Z}/2).$$

We do the same for $\mathbb{Z}/3$ -coefficients.

Remark 22. For $m = 6$, the check introduced in remark 20 takes the following form. The abelianization is the abelian group $\Gamma^{\mathrm{ab}} \cong \langle \overline{A}, \overline{R}, \overline{U}, \overline{W} : 2\overline{A} = 0, 3\overline{R} = 0 \rangle$. The parabolic elements are U and W . The element U generates nontrivial loops in the quotient, while W generates trivial loops. So it follows that $E_{0,1}^\infty \cong \mathbb{Z} \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/3$, generated by $\overline{W}, \overline{A}$ and \overline{R} . This is consistent with the computation above.

4. APPENDIX: THE EQUIVARIANT RETRACTION

In this section, we give Flöge's proof of the existence of a retraction ρ from $\widehat{\mathcal{H}}$ to the cell complex X^\bullet . We do not show the fact that ρ is Γ -equivariant, which can be observed since the fibers of ρ are geodesics.

Theorem 23 ([9, theorem 6.6]). *X is a retract of $\widehat{\mathcal{H}}$, i. e. there is a continuous map $\rho : \widehat{\mathcal{H}} \rightarrow X$ such that $\rho(p) = p$ for all $p \in X$.*

The map ρ is first defined as the orthogonal projection π from \widehat{B} to $\partial\widehat{B}$, and is then continued to the whole of $\widehat{\mathcal{H}}$ by Γ .

Bianchi [4] has shown that a nearly strict fundamental domain for the action of Γ on \mathcal{H} can be chosen in the form of a Euclidean vertical column D inside B . Define

$$\widehat{D} := \{(z, r) \in \widehat{B} \mid 0 \leq \mathrm{Re}(z) \leq 1, \quad 0 \leq \mathrm{Im}(z) \leq \sqrt{m}\},$$

and denote by S the set of singular points in \widehat{D} . Finally, $D := \widehat{D} - S$.

Remark 24 ([9], D is Γ -normal). For every $p \in \mathcal{H}$, there exists a neighborhood U of p in \mathcal{H} such that there are at most finitely many $g \in \Gamma$ with $gD \cap U \neq \emptyset$.

We will use the following lemmas to prove the theorem 23.

Lemma 25 ([9, lemma 6.5]). *For any subset $A \subset D$ which is closed in \mathcal{H} and any $p \in \mathcal{H}$, there exists an open neighborhood U_p of p such that we have for all $g \in \Gamma$:*

$gA \cap U_p \neq \emptyset$ if and only if $p \in gA$.

Proof. By the above remark, there is a neighborhood U of p in \mathcal{H} for which $\{g \in \Gamma \mid gD \cap U \neq \emptyset\}$ is finite. So especially its subset

$$\Gamma_o := \{g \in \Gamma \mid gA \cap U \neq \emptyset \text{ and } p \notin gA\}$$

is finite. Therefore, A being closed, $\bigcup_{g \in \Gamma_o} gA$ is closed in \mathcal{H} . Thus

$U_p := U - (\bigcup_{g \in \Gamma_o} gA)$ is open in \mathcal{H} and satisfies to the requested condition. \square

Lemma 26 ([9, lemma 6.3]). *There is an $\varepsilon_0 > 0$ such that for all singular points $s, s' \in S$, for all $\varepsilon \leq \varepsilon_0$ and $g \in \Gamma$ we have the following statement:*


$$g\widehat{U}_\varepsilon(s) \cap \widehat{U}_\varepsilon(s') \neq \emptyset \text{ implies } gs = s'.$$

For class number two, as we obtain a fundamental domain for the action of Γ on $\widehat{\mathcal{H}}$ (stricter than \widehat{D}) containing just one singular point, this lemma states only that Γ acts discontinuously on $\widehat{\mathcal{H}}$ (with respect to its topology which is finer than the subset topology of \mathbb{R}^3); and we skip Flöge's proof which is useful for class number three or greater.

Lemma 27 ([9, lemma 6.4]). *There exists an $\varepsilon_1 > 0$ with the following property:*

If $\varepsilon \leq \varepsilon_1$ and $(z, r) \in \widehat{D}$ with $r < \varepsilon$, then there is an $s' \in S$ such that $(z, r) \in \widehat{U}_{2\varepsilon}(s')$.

Flöge draws the following sketch of the situation in a vertical half-plane, which we reproduce here with his kind permission:



He gives only some hints on the proof, which we want to make slightly more explicit here.

Sketch of proof. We consider the Euclidean geometry of the upper-half space model for $\widehat{\mathcal{H}}$ and write coordinates in $\mathbb{C} \times \mathbb{R}^{\geq 0}$. Denote by ε_1 the “height of the lowest non-singular vertex”, more precisely the minimum of the values $r > 0$ occurring as the real coordinate of the non-singular vertices $(z, r) \in \mathcal{H}$

of the fundamental domain $\rho(\widehat{D})$ for Γ . Then $\{(z, r) \in \widehat{D} \mid r < \varepsilon_1\}$ consists of one connected component for each singular point $s' \in S$. We will denote by $\widehat{D}_{s'}$ the connected component containing s' . Now fix $s' \in S$. There are finitely many hemispheres limiting \widehat{D} from below and touching s' . We will consider the situation in a vertical half-plane containing s' . The most critical vertical half-planes for our assertion contain the intersection arc of two such hemispheres, because the other vertical half-planes contain circle segments of $\partial\widehat{D}$ of greater radius. The intersection of two non-identical Euclidean 2-spheres which have more than one point in common, is a circle with center on the line segment connecting the two 2-sphere centers. Thus the intersection of the two hemispheres mentioned above is a semicircle with center in the plane $r = 0$. Denote by ζ the radius of this semicircle. Then $\varepsilon_1 \leq \zeta$, because an edge of our fundamental domain, connecting s' with a non-singular vertex, lies on this semicircle. Now it is easy to see that $\widehat{D}_{s'}$ is a subset of the truncated cone obtained as the convex envelope of s' and the horizontal disk with radius ζ and center (s', ζ) . We conclude that for all $\varepsilon < \varepsilon_1$, $\varepsilon > 0$, the set $\{(z, r) \in \widehat{D}_{s'} \mid r < \varepsilon\}$ is a subset of the horoball $\widehat{U}_{2\varepsilon}(s')$. So we have seen that ε_1 has the property claimed in the lemma. \square

Proof of theorem 23. For any $(z, r) \in \widehat{D}$ there is a unique $r_z \geq 0$ such that $(z, r_z) \in \widehat{D} \cap \partial\widehat{B} =: \widehat{G}$, in fact $r_z = \min \{r' : (z, r') \in \widehat{D}\}$. We can thus define the map $\pi : \widehat{D} \rightarrow \widehat{G}$ by $\pi(z, r) := (z, r_z)$. The map π is continuous with respect to the subset topology of \mathbb{R}^3 , and by [9, corollary 5.10] also with respect to the topology of $\widehat{\mathcal{H}}$. Furthermore, we have $\pi(p) = p$ for all $p \in \widehat{G}$.

We now extend π to a map $\rho : \widehat{\mathcal{H}} \rightarrow X$ as follows.

Because of $\left\{ \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix} : b \in R \right\} \cdot \widehat{D} = \widehat{G}$, we find for any $p \in \widehat{\mathcal{H}}$ a $\gamma \in \Gamma$ such that $\gamma(p) \in \widehat{D}$. We set $\rho(p) := \gamma^{-1} \circ \pi \circ \gamma(p)$. In order to show that this makes sense, we have to show that $p \in \gamma^{-1}\widehat{D} \cap \xi^{-1}\widehat{D}$ implies

$\gamma^{-1} \circ \pi \circ \gamma(p) = \xi^{-1} \circ \pi \circ \xi(p)$, where $\gamma, \xi \in \Gamma$. We have $\xi(p) \in \xi\gamma^{-1}\widehat{D} \cap \widehat{D}$, then $\gamma\xi^{-1}(\xi(p)) = \gamma(p) \in \widehat{D} \cap \gamma\xi^{-1}\widehat{D}$, and either $\xi(p), \gamma(p)$ are both from \widehat{G} , or both from $\widehat{D} \cap B^\circ$. In the first case, it immediately follows that $\gamma^{-1} \circ \pi \circ \gamma(p) = \xi^{-1} \circ \pi \circ \xi(p) = p$, and $\xi^{-1} \circ \xi(p) = p$. In the second case, we have by [9, lemma 3.4] that if $\gamma\xi^{-1} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, the entry c must vanish. So $\gamma\xi^{-1}$ is the product $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \begin{pmatrix} 1 & db \\ 0 & 1 \end{pmatrix}$. Both of the latter two matrices commute with π since any such element ζ satisfies $\zeta(\partial\widehat{B}) = \partial\widehat{B}$, and ζ maps vertical half-lines to vertical half-lines.

So we have $(\gamma\xi^{-1} \circ \pi \circ \xi\gamma^{-1})p' = \pi p'$ for all $p' \in \widehat{D}$ with $\xi\gamma^{-1}p' \in \widehat{D}$, and then it follows that

$$\xi^{-1} \circ \pi \circ \xi(p) = \gamma \circ \gamma(\xi^{-1} \circ \pi \circ \xi)\gamma^{-1}\gamma(p) = \gamma^{-1} \circ \pi \circ \gamma(p) = \gamma^{-1} \circ \pi \circ \gamma(p).$$

Thus, ρ is well-defined. Furthermore, $\pi(p) = p$ for all $p \in \widehat{G}$ implies $\rho(p) = p$ for all $p \in X$. It remains to show that ρ is continuous at any $p \in \widehat{\mathcal{H}}$.

1st case. In the case $p \in \mathcal{H}$, by lemma 25, p has an open neighborhood U_p such that: for any $\gamma \in \Gamma$, we have $\gamma U_p \cap D \neq \emptyset \iff \gamma(p) \in D$. Furthermore, the set $\{\gamma \in \Gamma : \gamma(p) \in D\}$ is finite [9, remark 3.6], say $\gamma_1, \dots, \gamma_n$.

Let now V be an open neighborhood of $\rho(p)$. Because of the continuity of

all γ_i, γ_i^{-1} and the continuity of $\pi : \widehat{D} \rightarrow \widehat{G}$, there exist neighborhoods U_i of p such that $\gamma_i^{-1} \circ \pi \circ \gamma_i(U_i) \subset V$. Note that for all γ_i we have $\gamma_i^{-1} \circ \pi \circ \gamma_i(p) = \rho(p)$.

Setting $U := U_p \cap (\bigcap_{i=1}^n U_i)$, we have $\rho(U) \subset V$, i. e. ρ is continuous at the point p .

2nd case. In the case $p \in \widehat{\mathcal{H}} \cap \mathbb{C}$, let ϵ_0, ϵ_1 and ϵ_s for $s \in S$ be positive real numbers as in lemma 26, lemma 27 and [9, lemma 5.9]; and let $\epsilon > 0$ be less than the minimum of $\frac{\epsilon_0}{2}, \epsilon_1, \epsilon_s$ for $s \in S$. Because of $\left\{ \begin{pmatrix} 1 & b \\ a & 1 \end{pmatrix} : b \in R \right\} \cdot \widehat{D} = \widehat{G}$, there exist $s \in S, \xi = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ such that $\xi s = p$ and by [9, remark 5.5(a)], we have $\xi \widehat{U}_\epsilon(s) = \widehat{U}_{\frac{\epsilon}{|cs-d|^2}}(p)$. Let us now show that $\rho(\widehat{U}_{\frac{\epsilon}{|cs-d|^2}}(p)) \subset \widehat{U}_{2\epsilon}(p)$.

Let $p' \in \widehat{U}_{\frac{\epsilon}{|cs-d|^2}}(p)$, and let $\gamma \in \Gamma$ with $\gamma p' \in \widehat{D}$. Then $\rho(p') = \gamma^{-1} \circ \pi \circ \gamma(p')$.

By [9, remark 5.5(b)], applied to s and $\gamma\xi$ it follows that

$\gamma p' = \gamma\xi(\xi^{-1}p') \in \widehat{U}_\epsilon(\gamma\xi s) = \widehat{U}_\epsilon(\gamma p)$, and by [9, remark 5.6] all conditions of lemma 27 are satisfied. So there is an $s' \in S$ such that $\gamma p' \in \widehat{U}_{2\epsilon}(s')$.

This means that $\gamma\xi(\widehat{U}_{2\epsilon}(s)) \cap \widehat{U}_{2\epsilon}(s') \neq \emptyset$, and by lemma 26 it follows that $s' = \gamma\xi s = \gamma p$.

Let us now consider $\gamma p'$ again. Since $\gamma p' \in \widehat{U}_\epsilon(\gamma p) = \widehat{U}_\epsilon(s') = U_\epsilon(s')$ and $\pi(U_\epsilon(s')) \subset U_\epsilon(s')$; and by [9, lemma 5.9] we have $U_\epsilon(s') \cap \widehat{B} \subset \widehat{U}_{2\epsilon}(s')$. So $\pi \circ \gamma p' \in \widehat{U}_{2\epsilon}(s')$. By [9, remark 5.5(b)] it finally follows that

$$\rho(p') = \gamma^{-1} \circ \pi \circ \gamma p' \in \gamma^{-1} \widehat{U}_{2\epsilon}(s') \subset \widehat{U}_{2\epsilon}(\gamma^{-1} s') = \widehat{U}_{2\epsilon}(p),$$

and we are done. \square

REFERENCES

- [1] Herbert Abels, *Generators and relations for groups of homeomorphisms*, Transformation groups (Proc. Conf., Univ. Newcastle upon Tyne, Newcastle upon Tyne, 1976), Cambridge Univ. Press, Cambridge, 1977, pp. 3–20. London Math. Soc. Lecture Note Series, No. **26**. MR **0474339** (**57** #13985), Zbl 0344.57014
- [2] Bill Allombert, Karim Belabas, Henri Cohen, Xavier Roblot, Ilya Zakharevitch, and the PARI group, *PARI/GP*, specialized computer algebra system. Documentation and free download on <http://pari.math.u-bordeaux.fr/>, 2008.
- [3] Ethan Berkove, *The integral cohomology of the Bianchi groups*, Trans. Amer. Math. Soc. **358** (2006), no. 3, 1033–1049 (electronic). MR **2187644** (**2006h**:20073), Zbl pre02237880
- [4] Luigi Bianchi, *Sui gruppi di sostituzioni lineari con coefficienti appartenenti a corpi quadratici immaginari*, Math. Ann. **40** (1892), no. 3, 332–412 (Italian). MR **1510727**, JFM 24.0188.02
- [5] Kenneth S. Brown, *Cohomology of groups*, Graduate Texts in Mathematics, vol. **87**, Springer-Verlag, New York, 1982. MR **672956** (**83k**:20002), Zbl 0584.20036
- [6] Jürgen Elstrodt, Fritz Grunewald, and Jens Mennicke, *Groups acting on hyperbolic space*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998. Harmonic analysis and number theory. MR **1483315** (**98g**:11058), Zbl 0888.11001
- [7] Benjamin Fine, *Algebraic theory of the Bianchi groups*, Monographs and Textbooks in Pure and Applied Mathematics, vol. **129**, Marcel Dekker Inc., New York, 1989. MR **1010229** (**90h**:20002), Zbl 0760.20014
- [8] Dieter Flöge, *Zur Struktur der PSL_2 über einigen imaginär-quadratischen Zahlringen*, Math. Z. **183** (1983), no. 2, 255–279 (German). MR **704107** (**85c**:11043), Zbl 0498.20036

- [9] ———, *PhD thesis: Zur Struktur der PSL_2 über einigen imaginär-quadratischen Zahlringen*, Johann-Wolfgang-Goethe-Universität, Fachbereich Mathematik, Frankfurt am Main, 1980 (German). Zbl 0482.20032
- [10] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4.12*, <http://www.gap-system.org>, 2008.
- [11] Günter Harder, *A Gauss-Bonnet formula for discrete arithmetically defined groups*, Ann. Sci. École Norm. Sup. (4) **4** (1971), 409–455. MR **0309145** (46 #8255), Zbl 0232.20088
- [12] Allen Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. MR **1867354** (2002k:55001), Zbl 1044.55001
- [13] Colin Maclachlan and Alan W. Reid, *The arithmetic of hyperbolic 3-manifolds*, Graduate Texts in Mathematics, vol. **219**, Springer-Verlag, New York, 2003. MR **1937957** (2004i:57021), Zbl 1025.57001
- [14] Eduardo R. Mendoza, *Cohomology of PGL_2 over imaginary quadratic integers*, Bonner Mathematische Schriften [Bonn Mathematical Publications], **128**, Dissertation, Rheinische Friedrich-Wilhelms-Universität, Mathematisches Institut, Bonn, 1979. MR **611515** (82g:22012), Zbl 0464.12005
- [15] Władysław Narkiewicz, *Elementary and analytic theory of algebraic numbers. 3rd ed.*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2004. MR **2078267** (2005c:11131), Zbl 0717.11045
- [16] Joachim Schwermer and Karen Vogtmann, *The integral homology of SL_2 and PSL_2 of Euclidean imaginary quadratic integers*, Comment. Math. Helv. **58** (1983), no. 4, 573–598. MR **728453** (86d:11046), Zbl 0545.20031
- [17] Jean-Pierre Serre, *Cohomologie des groupes discrets*, Prospects in mathematics (Proc. Sympos., Princeton Univ., Princeton, N.J., 1970), Princeton Univ. Press, Princeton, N.J., 1971, pp. 77–169. Ann. of Math. Studies, No. **70** (French). MR **0385006** (52 #5876), Zbl 0235.22020
- [18] ———, *A course in arithmetic*, Springer-Verlag, New York, 1973. Translated from the French; Graduate Texts in Mathematics, No. **7**. MR **0344216** (49 #8956), Zbl 0256.12001
- [19] Richard G. Swan, *Generators and relations for certain special linear groups*, Advances in Math. **6** (1971), 1–77 (1971). MR **0284516** (44 #1741), Zbl 0221.20060
- [20] Karen Vogtmann, *Rational homology of Bianchi groups*, Math. Ann. **272** (1985), no. 3, 399–419. MR **799670** (87a:22025), Zbl 0545.20031
- [21] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. **38**, Cambridge University Press, Cambridge, 1994. MR **1269324** (95f:18001), Zbl 0797.18001

I3M, UNIVERSITÉ MONTPELLIER 2 AND MATH. INSTITUT, UNIVERSITÄT GÖTTINGEN
E-mail address: alexander.rahm@math.univ-montp2.fr
URL: www.math.univ-montp2.fr/SPIP/_RAHM-Alexander_

DEPARTMENT OF BIOINFORMATICS, CENTER OF INFORMATICS, STATISTICS AND EPIDEMIOLOGY UMG, UNIVERSITY OF GÖTTINGEN
E-mail address: mfu@bioinf.med.uni-goettingen.de
URL: www.uni-math.gwdg.de/fuchs