



HAL
open science

Khovanov homology for signed divides

Olivier Couture

► **To cite this version:**

Olivier Couture. Khovanov homology for signed divides. Algebraic and Geometric Topology, 2009, 9 (4), pp.1987-2026. 10.2140/agt.2009.9.1987 . hal-00371949

HAL Id: hal-00371949

<https://hal.science/hal-00371949v1>

Submitted on 31 Mar 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Khovanov homology for signed divides.

Olivier Couture

Institut de Mathématiques de Bourgogne
Université de Bourgogne
UFR Sciences et Techniques
9 avenue Alain Savary - B.P 47870
21078 DIJON Cedex - France
olivier.couture@u-bourgogne.fr

Dedicated to Bernard Perron

Abstract

The purpose of this paper is to interpret polynomial invariants of strongly invertible links in terms of Khovanov homology theory. To a divide, that is a proper generic immersion of a finite number of copies of the unit interval and circles in a 2-disc, one can associate a strongly invertible link in the 3-sphere. This can be generalized to signed divides : divides with + or - sign assignment to each crossing point. Conversely, to any link L that is strongly invertible for an involution j , one can associate a signed divide. Two strongly invertible links that are isotopic through an isotopy respecting the involution are called strongly equivalent. Such isotopies give rise to moves on divides. In a previous paper of the author [2], one can find an exhaustive list of moves that preserves strong equivalence, together with a polynomial invariant for these moves, giving therefore an invariant for strong equivalence of the associated strongly invertible links. We prove in this paper that this polynomial can be seen as the graded Euler characteristic of a graded complex of vector spaces. Homology of such complexes is invariant for the moves on divides and so is invariant through strong equivalence of strongly invertible links.

57M27

Keywords: Strongly invertible links; Divides; Morse signed divides; Khovanov homology.

Introduction.

A *divide* Γ is the image of a proper generic immersion of a finite number of intervals and circles into the unit 2-disc \mathbf{D}^2 of \mathbb{R}^2 . With a divide Γ , N.A'Campo [1] associates a link $\mathcal{L}(\Gamma)$ in the unit 3-sphere of the tangent space $T\mathbf{D}^2$:

$$\mathcal{L}(\Gamma) = \{(p, v) \in T\mathbf{D}^2 : p \in \Gamma, v \in T_p\Gamma, \|p\|^2 + \|v\|^2 = 1\}.$$

This link has natural orientation and is strongly invertible with respect to the involution $j(p, v) = (p, -v)$.

In Couture–Perron [3] is given a generalization of divides. Let (x, y) be coordinates in \mathbf{D}^2 such that the restriction to Γ of the projection $\pi_1 : (x, y) \mapsto x$ is a Morse function. A *Morse signed divide* (MS–divide) relative to π_1 stands for such a divide with $+$ or $-$ sign assignment to each double point of Γ . Furthermore, if there exists $a \in]0, 1[$ such that all maxima (resp. minima) of $\pi_{1|\Gamma}$ project on a (resp. $-a$) and all double points in $] - a; a[$, the MS–divide is called *ordered (OMS–divide)*.

We also define a link associated with a MS–divide (see [3]), which is strongly invertible with respect to the involution $j(p, v) = (p, -v)$. If all signs are positive, this link is no more than the link of the divide without signs. The interest of OMS–divides rather than MS–divides is to obtain an immediate braid presentation of the link from the divide.

Strongly invertible links are closely related with OMS–divides. Let L be an oriented link in \mathbb{S}^3 and j be an involution of (\mathbb{S}^3, L) with non empty fixed point set, which preserves the orientation of \mathbb{S}^3 and reverses the orientation of L . Then (L, j) is called a strongly invertible link. As we said above, the link of a divide is strongly invertible for the implicit strong inversion $j(p, v) = (p, -v)$. Two strongly invertible links (L, j) and (L', j') are *strongly equivalent* if there exists an isotopy φ_t , $t \in [0, 1]$ of \mathbb{S}^3 sending L to L' such that $\varphi_1 \circ j = j' \circ \varphi_1$.

Isotopies through MS–divides give rise to strong equivalence of associated links. Also, one can find in Couture [2] an (exhaustive) list of elementary moves on MS–divides that preserve strong equivalence classes of the associated links. As a particular case, given a MS–divide, we can always construct another one, using these moves, which is an OMS–divide. Besides, we can transpose the moves on MS–divides directly to moves on OMS–divides. Two OMS–divides obtained one from the other by those moves on OMS–divides are called \mathcal{M} –equivalent (see the list of moves in Section 1.3). Then as an essential result of [2], we have:

- Theorem** (cf. [2]).
1. *Every strongly invertible link is strongly equivalent to the link of an OMS–divide.*
 2. *The links of two OMS–divides are strongly equivalent if and only if the OMS–divides are \mathcal{M} –equivalent.*

As the Jones polynomial is invariant under Reidemeister moves on links diagrams, there exists a Laurent polynomial for an OMS–divide with integral coefficients (see Couture [2]), which is invariant under \mathcal{M} –equivalence and so invariant under strong equivalence of strongly invertible links. Modulo 2, this polynomial coincide with Jones polynomial of the link. The purpose of this paper is to interpret the polynomial of an OMS–divide as the graded Euler characteristics of a graded complex of \mathbb{Z}_2 –vector spaces (theorem 3.16). Besides, if we call Khovanov homology of an OMS–divide the homology of this complex, then we have a stronger result:

Theorem. 3.17 *Khovanov homology of OMS–divides is invariant under \mathcal{M} –equivalence.*

Corollary. 3.18 *Khovanov homology is an invariant for strong equivalence of strongly invertible links.*

Eventually, Khovanov homology of OMS–divides is a refinement of the polynomial invariant of OMS–divides.

1 Divides and OMS–divides.

1.1 Divides and links of divides.

A *divide* of the unit 2–disc \mathbf{D}^2 of $\mathbb{R}^2(\simeq \mathbb{C})$ is the image Γ of a proper generic immersion γ :

$$(1.1) \quad \gamma : (J, \partial J) \rightarrow (\mathbf{D}^2, \partial \mathbf{D}^2), \quad J = \left(\bigsqcup_{j=1}^r I_j \right) \sqcup \left(\bigsqcup_{j=1}^s S_j \right).$$

where I_j and S_j are respectively copies of $[0, 1]$ and $\mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$, generic meaning that the only singularities of γ are ordinary double points and $\Gamma = \gamma(J)$ intersects $\partial \mathbf{D}^2$ transversally. Every $\gamma(I_j)$ (resp. $\gamma(S_j)$) is called *interval* (resp. *circular*) branch.

Let $S(\mathbf{D}^2) = \{(p, v) \in T\mathbf{D}^2 : |p|^2 + |v|^2 = 1\}$ be the unit sphere of the tangent space $T\mathbf{D}^2 \simeq \mathbf{D}^2 \times \mathbb{C}$. With a divide, A'Campo [1] associates a link $\mathcal{L}(\Gamma)$ in $S(\mathbf{D}^2)$:

$$(1.2) \quad \mathcal{L}(\Gamma) = \{(p, v) \in S(\mathbf{D}^2) : p \in \Gamma, v \in T_p \Gamma\}.$$

This link has a natural orientation induced by the two possible orientations of the branches of Γ and is strongly invertible for the involution $j(p, v) = (p, -v)$ of $S(\mathbf{D}^2)$ with axis $\text{Fix}(j) = \partial \mathbf{D}^2 \times \{0\}$ (see Section 1.3 below). Each interval branch of Γ leads to a strongly invertible component of $\mathcal{L}(\Gamma)$ and each circular branch of Γ to two components of $\mathcal{L}(\Gamma)$ interchanged by j .

1.2 OMS–Divides.

Let Γ be a divide. Suppose there exists $(a, b) \in]0, 1[\times]0, 1[$, $a^2 + b^2 < 1$ such that:

1. $\Gamma \subset \{x + iy \in \mathbf{D}^2 : -b < y < b\}$ and the restriction $\rho|_{\Gamma}$ is a Morse function;
2. all double points of Γ are contained in $] -a, a[\times] -b, b[$;
3. all maxima (resp. minima) of $\rho|_{\Gamma}$, called *vertical tangent points*, project on a (resp. $-a$).

Then Γ is called an *ordered Morse divide*. Double points and vertical tangent points will be called *singular points* of Γ . Now let ϵ be a function that associates $+$ or $-$ sign with each double point of Γ . Then (Γ, ϵ) is called an *ordered Morse signed divide* (OMS–divide) (relative to the projection $\rho(x + iy) = x$) (see [3]).

Let's associate an oriented j –strongly invertible link $\mathcal{L}(\Gamma, \epsilon)$ with an OMS–divide (Γ, ϵ) . This link coincides with $\mathcal{L}(\Gamma)$ except in solid tori $TD_p \cap S(\mathbf{D}^2) \simeq D_p \times \mathbb{S}^1$ over small discs D_p around negative double points p of (Γ, ϵ) where we change the two j –symmetric crossings from over to under. More precisely, in such a solid torus $TD_p \cap S(\mathbf{D}^2)$, the link is defined according to Figure 1.1. If $\epsilon = +$ for all double points then $\mathcal{L}(\Gamma, \epsilon) = \mathcal{L}(\Gamma)$.

Besides, from a divide Γ one can construct an OMS–divide (Γ', ϵ) by a succession of moves and isotopies, such that $\mathcal{L}(\Gamma', \epsilon)$ and $\mathcal{L}(\Gamma)$ are isotopic (see [3]) by an isotopy that respects the involution j (see Section 1.3 below for a more precise definition).




Figure 1.1: The link $\mathcal{L}(\Gamma, \epsilon)$ over a negative double point.

Remark 1.1. For simplicity, we will only consider an OMS-divide (Γ, ϵ) in $[-a, a] \times [-b, b]$, omitting its trivial part outside this rectangle. After rescaling, we also suppose that $a = b = 1$. Since we will often consider diagrams of local parts of OMS-divides (Γ, ϵ) , we distinguish end points of Γ , i.e. points of Γ in $\{-1, 1\} \times [-1, 1]$ without vertical tangent by a big point (see Figure 1.2).




Figure 1.2: A representative OMS-divide for the strongly invertible knot 5_2 .

Moreover, we will simply denote Γ instead of (Γ, ϵ) if no ambiguity occurs in the context.

1.3 M-equivalence for OMS-divides.

Two OMS-divides Γ and Γ' are \mathcal{M} -equivalent if we obtain one from the other by isotopy through OMS-divides and a finite sequence of the moves described on Figure 1.3 or symmetric situations with respect to horizontal and vertical directions (see [2]): Let j be an orientation




Figure 1.3: Moves of \mathcal{M} -equivalence.

preserving involution of \mathbb{S}^3 with non empty fix point set (i.e. $\text{Fix}(j)$ is trivial knot according to the solution of Smith conjecture [4]). An oriented link $L \subset \mathbb{S}^3$ is j -strongly invertible if j sends L to itself with opposite orientation. The couple (L, j) is called a strongly invertible link. With the link of an OMS-divide, we implicitly associate natural orientation and involution $j(p, v) = (p, -v)$ as in Section 1.1: such a link is strongly invertible.

Two strongly invertible links (L, j) and (L', j') are called *strongly equivalent*¹ if there exists

¹A same link L may have two strong inversions j and j' such that (L, j) and (L, j') are not strongly equivalent.

and isotopy φ_t , $t \in [0, 1]$ of \mathbb{S}^3 sending L to L' such that $\varphi_1 \circ j = j' \circ \varphi_1$. One can easily prove that \mathcal{M} -equivalent OMS-divides give rise to strongly equivalent strongly invertible links. Conversely, let's recall the following crucial theorem relating OMS-divides with strongly invertible links.

Theorem 1.2. [2]

1. Every strongly invertible link is strongly equivalent to the link of an OMS-divide.
2. The links of two OMS-divides are strongly equivalent if and only if the OMS-divides are \mathcal{M} -equivalent.

2 The polynomial of an OMS-divide.

Let's denote by Θ_0 and Θ_1 the local splittings of an OMS-divide (Γ, ϵ) in a neighborhood of a double point or vertical tangent point described in Figure 2.1 (Θ_0 "smooths" the OMS-divide whereas Θ_1 introduces horizontal cusps).




Figure 2.1: Local splittings.

Definition 2.1.

1. We extend the notion of OMS-divide: a cuspidal divide $\Gamma := (\Gamma, \epsilon)$ is a signed diagram like an OMS-divide except that it has a finite number of horizontal cusps (as in the result of type Θ_1 splittings). For instance, a partially (or totally) transformed OMS-divide through Θ_0 and Θ_1 is a cuspidal divide.
2. Let (Γ, ϵ) be an OMS-divide (or more generally of a cuspidal divide) with double and vertical tangent points numbered by p_1, \dots, p_n . Let $[k]$ be the word $k_1 k_2 \dots k_n$ $k_i \in \{0, 1\}$. A state $(S, \Theta_{[k]})$ of (Γ, ϵ) is the combination of:
 - a succession of local splittings $\Theta_{[k]} = (\Theta_{k_1}, \dots, \Theta_{k_n})$ at p_1, \dots, p_n .
 - the cuspidal divide $S = \Theta_{[k]}(\Gamma, \epsilon)$ without double points nor vertical tangent points obtained by transforming Γ through $\Theta_{[k]}$.
For simplification, we will often identify the cuspidal divide S with the state $(S, \Theta_{[k]})$. We denote by $St(\Gamma, \epsilon)$ the set of all states of (Γ, ϵ) .

One can define a j -strongly invertible link $\mathcal{L}(\Gamma, \epsilon)$ associated with a cuspidal divide (Γ, ϵ) exactly in the same way we have done for OMS-divide. However, such a link is generally unoriented precisely because of the introduction of cusps. Each local splitting at a double point of (Γ, ϵ) corresponds to simultaneously smoothing two symmetric crossing points of the corresponding representative closed braid diagram of $\mathcal{L}(\Gamma, \epsilon)$ (see [3]) whereas each local splitting at a vertical tangent point corresponds to smoothing a crossing point through the axis of the inversion j (see Figure 2.2).




Figure 2.2: The links $\mathcal{L}(\Gamma, \epsilon)$ and $\mathcal{L}(\Theta(\Gamma, \epsilon))$, $\Theta = \Theta_{1110010}$.

Let $\Gamma := (\Gamma, \epsilon)$ be an OMS-divide. Let $n = n_+ + n_- + n_0$ be the number of singular points of Γ where n_+ , n_- are respectively the numbers of positive and negative double points, and n_0 the number of vertical tangent points. Let's call:

$$(2.1) \quad w(\Gamma) = 2n_+ - 2n_- + n_0$$

the *writhe* of Γ (i.e. the writhe of the representative closed braid diagram of $\mathcal{L}(\Gamma, \epsilon)$ (see [3]) with $2n_+ + 2n_- + n_0$ crossings obtained from (Γ, ϵ)).

For a state $S \in St(\Gamma, \epsilon)$, let $cl(S)$ be the number of closed connected components and $op(S)$ be the number of open connected components (i.e. with two end points). Let $r_+(S)$, $r_-(S)$ and $r_0(S)$ be the numbers of Θ_1 local splittings (Figure 2.1) for positive double points, negative double points and vertical tangent points respectively to obtain S from (Γ, ϵ) . Let's set:

$$(2.2) \quad i(S) = r_+(S) - r_-(S) + r_0(S) \quad k(S) = w(\Gamma) + 2i(S) - r_0(S).$$

Definition 2.2. (cf. [2]) *The polynomial of an OMS-divide Γ (and more generally of a cuspidal divide) is the Laurent polynomial (of the variable \sqrt{t}) defined by:*

$$(2.3) \quad W_\Gamma(t) = \sum_{S \in St(\Gamma, \epsilon)} (-1)^{i(S)} (\sqrt{t})^{k(S)} \left(\frac{1}{t} + t\right)^{cl(S)} \left(\frac{1}{\sqrt{t}} + \sqrt{t}\right)^{op(S)-1}$$

Proposition 2.3. (cf. [2]) *The polynomial of an OMS-divide is invariant under \mathcal{M} -equivalence of OMS-divides and so is an invariant for strong equivalence of strongly invertible links.*

Definition 2.4. *A state S with + or - assignment to each connected component is called an enhanced state, and is denoted by \tilde{S} . The set of enhanced states of (Γ, ϵ) is denoted by $\tilde{St}(\Gamma, \epsilon)$, and S is called the underlying state of \tilde{S} .*

Let \tilde{S} be an enhanced state with underling state S . The numbers $i(\tilde{S}) := i(S)$ and $k(\tilde{S}) := k(S)$ in (2.2) do not depend of the signs of the components. The subset of enhanced states \tilde{S} of (Γ, ϵ) such that $i(\tilde{S}) = i$ is denoted by $\tilde{St}_i(\Gamma, \epsilon)$.

Let's denote by $\delta_{cl}(\tilde{S})$ (resp. $\delta_{op}(\tilde{S})$) the difference between the number of positive and negative closed (resp. open) components of \tilde{S} . Then we define the *degree* $j(\tilde{S})$ of the enhanced state \tilde{S} , which depends of the signs of the components of S by:

$$(2.4) \quad j(\tilde{S}) = k(\tilde{S}) + 2\delta_{cl}(\tilde{S}) + \delta_{op}(\tilde{S}).$$

We can now reformulate the polynomial of an OMS–divide Γ :

$$(2.5) \quad W_\Gamma(t) = \frac{\sqrt{t}}{1+t} \sum_{\tilde{S} \in \tilde{St}(\Gamma)} (-1)^{i(\tilde{S})} (\sqrt{t})^{j(\tilde{S})}$$

Remark 2.5. $j(\tilde{S})$ always has the same parity as half the number of end points of Γ . We also have the inequalities:

$$(2.6) \quad -n_- \leq i(\tilde{S}) \leq n_+ + n_0 \quad 2n_+ - 4n_- + n_0 \leq k(\tilde{S}) \leq 4n_+ - 2n_- + 2n_0.$$

3 Categorification.

3.1 Complex associated with an OMS–divide.

In this section, we define a graded complex of \mathbb{Z}_2 -vector spaces² associated with a divide. We follow here Viro’s approach of Khovanov homology for links [5], based on the Kauffman state model for the Jones polynomial: the polynomial of a divide also have been defined in [2] by state model.

Let $\Gamma := (\Gamma, \epsilon)$ be an OMS–divide (or a cuspidal divide). For $i \in \mathbb{Z}$, let $[[\Gamma]]_i = \mathbb{Z}_2\{\tilde{St}_i(\Gamma)\}$ be the finite dimensional \mathbb{Z}_2 -vector space generated by enhanced states \tilde{S} of Γ such that $i(\tilde{S}) = i$ (if $i < -n_-$ or $i > n_+ + n_0$, $[[\Gamma]]_i = \{0\}$). Degree $j(\tilde{S})$ defines a grading on $[[\Gamma]]_i$ and we denote:

$$(3.1) \quad [[\Gamma]] = ([[\Gamma]])_{i \in \mathbb{Z}} \quad [[\Gamma]]_i = \bigoplus_{j \in \mathbb{Z}} [[\Gamma]]_{i,j}$$

where $[[\Gamma]]_{i,j} = \mathbb{Z}_2\{\tilde{S} \in St_i(\Gamma) : j(\tilde{S}) = j\}$.

Now we define a differential on $[[\Gamma]]$ to obtain a (finite) complex of graded \mathbb{Z}_2 -vector spaces.

Definition 3.1. Let $\tilde{S}_1, \tilde{S}_2 \in \tilde{St}(\Gamma, \epsilon)$. We say that \tilde{S}_2 is adjacent to \tilde{S}_1 ($\tilde{S}_1 \rightsquigarrow \tilde{S}_2$) if:

1. S_1 and S_2 coincide outside a neighborhood D_p of a singular point p of (Γ, ϵ) ;
2. One can pass from \tilde{S}_1 to \tilde{S}_2 by one of the following transformations T in D_p :
 - $T = \Theta_1 \circ \Theta_0^{-1}$ if p is a positive double point or a vertical tangent point of (Γ, ϵ) ;
 - $T = \Theta_0 \circ \Theta_1^{-1}$ if p is a negative double point;
3. Signs rules described in [Figure 3.1](#), [3.2](#), [3.3](#) are fulfilled, signs of other components being unchanged.

(In these figures, black color is used for open components and gray for closed components, a dotted line means that the points are related in the state outside D_p . Lack of dotted line means that the points are not related outside D_p).

If \tilde{S}_2 is adjacent to \tilde{S}_1 then:

$$(3.2) \quad j(\tilde{S}_1) = j(\tilde{S}_2), \quad i(S_1) = i(S_2) - 1.$$




Figure 3.1: Case of a splitting at a positive double point p ($\epsilon(p) = +$).




Figure 3.2: Case of a splitting at a negative double point p ($\epsilon(p) = -$).




Figure 3.3: Case of a splitting at a vertical tangent point p .

The differential $d = (d_i)_{i \in \mathbb{Z}}$ on $[[\Gamma]]$, $d_i : [[\Gamma]]_i \rightarrow [[\Gamma]]_{i+1}$ is now defined in the following way: the matrix of d_i has coefficients defined by the incidence numbers $(\tilde{S}_1 : \tilde{S}_2)$, $\tilde{S}_1 \in \tilde{St}_i(\Gamma)$, $\tilde{S}_2 \in \tilde{St}_{i+1}(\Gamma)$:

$$(3.3) \quad (\tilde{S}_1 : \tilde{S}_2) = \begin{cases} 1 & \text{if } \tilde{S}_1 \rightsquigarrow \tilde{S}_2 \\ 0 & \text{else.} \end{cases}$$

From (3.2), d respects the degree j , i.e. $d_i = \bigoplus_j d_{i,j} : \bigoplus_j [[\Gamma]]_{i,j} \rightarrow \bigoplus_j [[\Gamma]]_{i+1,j}$.

²Here we choose \mathbb{Z}_2 -vector spaces for simplification to avoid signs. We can easily generalize taking for instance \mathbb{Z} -modules or \mathbb{Q} -vector spaces

Remark 3.2. We have dual roles for T_1^+ and T_1^- , T_2^+ and T_3^- , T_3^+ and T_2^- , T_4^+ and T_5^- , T_5^+ and T_4^- in [Figure 3.1](#) and [Figure 3.2](#). To go further about duality property, we could have introduced “negative tangent points” to interpret dual arrows of T_6 and T_7 in [Figure 3.3](#). However, we didn’t choose this option, since such “negative tangent points” can be replaced by:



Also, we can see that T_i^+ and T_i^- , $i \in \{1, 2, 3, 4\}$ give rise to analogous situations.

Proposition 3.3. $([[\Gamma]], d) = ([[\Gamma]], d_i)_{i \in \mathbb{Z}} = (\bigoplus_{j \in \mathbb{Z}} [[\Gamma]]_{i,j}, \bigoplus_{j \in \mathbb{Z}} d_{i,j})_{i \in \mathbb{Z}}$ is a finite complex of graded \mathbb{Z}_2 -vector spaces (each $[[\Gamma]]_i$ is finitely graded by degree j).

Proof. It suffices to verify all such diagrams:

$$(3.4) \quad \begin{array}{ccc} \tilde{S}_1 & \rightsquigarrow & \tilde{S}_2 \\ \downarrow \zeta & & \downarrow \zeta \\ \tilde{S}'_2 & \rightsquigarrow & \tilde{S}_3 \end{array}$$

corresponding to splitting two singular points are commutative. Since we have \mathbb{Z}_2 -vector spaces, commutative diagrams induce the relations $d_{i+1}d_i = 0$. Notice that from the previous remark, we can strongly reduce the number of cases to check (see also the proof of proposition [3.5](#)). \square

3.2 Alternative point of view.

Here we present an alternative (more algebraic) way to see the complex $([[\Gamma]], d)$, in terms of Frobenius algebra: probably can we link complexes of OMS-divides with 1+1-TQFT (or more precisely with some 1+1-TQFT with symmetry property).

Let $\mathcal{A} := \mathbb{Z}_2\{v_-, v_+\}$ be the graded \mathbb{Z}_2 -vector space generated by two elements v_- and v_+ such that $\deg(v_-) = -1$ and $\deg(v_+) = 1$. We define a commutative product $\mu_1 : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$, a unit $\eta_1 : \mathbb{Z}_2 \rightarrow \mathcal{A}$ and a non degenerate symmetric bilinear pairing $\beta_1 : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathbb{Z}_2$ by:

$$(3.5) \quad \begin{aligned} \mu_1(v_+ \otimes v_+) &= v_+, & \mu_1(v_+ \otimes v_-) &= \mu_1(v_- \otimes v_+) = v_-, & \mu_1(v_- \otimes v_-) &= 0 \\ \eta_1(0) &= 0, & \eta_1(1) &= v_+ \\ \beta_1(v_+ \otimes v_+) &= 0, & \beta_1(v_+ \otimes v_-) &= \beta_1(v_- \otimes v_+) = 1, & \beta_1(v_- \otimes v_-) &= 0. \end{aligned}$$

The form β_1 induces a duality isomorphism $\mathcal{A} \xrightarrow{\cong} \mathcal{A}^*$ and \mathcal{A} is a commutative Frobenius algebra with adjoint co-product $\delta_1 : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ and co-unit $\varepsilon_1 : \mathcal{A} \rightarrow \mathbb{Z}_2$:

$$(3.6) \quad \begin{aligned} \delta_1(v_+) &= v_+ \otimes v_- + v_- \otimes v_+ & \varepsilon_1(v_+) &= 0 \\ \delta_1(v_-) &= v_- \otimes v_- & \varepsilon_1(v_-) &= 1. \end{aligned}$$

Let $\phi_1 : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$ be the flip morphism: $\phi_1(a \otimes a') = a' \otimes a$, and $A : \mathcal{A} \rightarrow \mathcal{A}$ the identity morphism. The morphisms $\mu_1, \delta_1, \eta_1, \varepsilon_1$ are homogeneous with respective degrees $-1, -1, 1, 1$

and verify the relations of associativity, commutativity, co-associativity, co-commutativity:

$$(3.7) \quad \begin{cases} \mu_1 \circ \phi_1 = \mu_1 \\ \mu_1 \circ (\mu_1 \otimes A) = \mu_1 \circ (A \otimes \mu_1) \\ \mu_1 \circ (\eta_1 \otimes A) = A \end{cases} \quad \begin{cases} \phi_1 \circ \delta_1 = \delta_1 \\ (\delta_1 \otimes A) \circ \delta_1 = (A \otimes \delta_1) \circ \delta_1 \\ (\varepsilon_1 \otimes A) \circ \delta_1 = (A \otimes \varepsilon_1) \circ \delta_1 = A \end{cases}$$

and

$$\delta_1 \circ \mu_1 = (\mu_1 \otimes A) \circ (A \otimes \delta_1).$$

The vector space $\mathcal{A} \otimes \mathcal{A}$ has an induced structure of graded commutative Frobenius algebra with product, co-product, unit and co-unit :

$$(3.8) \quad \begin{aligned} \mu_1^\otimes &= (\mu_1 \otimes \mu_1) \circ (A \otimes \phi \otimes A) & \eta_1^\otimes &= \eta_1 \otimes \eta_1 \\ \delta_1^\otimes &= (A \otimes \phi \otimes A) \circ (\delta_1 \otimes \delta_1) & \varepsilon_1^\otimes &= \varepsilon_1 \otimes \varepsilon_1. \end{aligned}$$

Let $\mathcal{B} := \mathbb{Z}_2\{w_-, w_+\}$ be the graded \mathbb{Z}_2 -vector space generated by two elements w_- and w_+ , $\deg(w_-) = -2$, $\deg(w_+) = 2$. Let's consider respectively the injection and surjection $\iota : \mathcal{B} \rightarrow \mathcal{A} \otimes \mathcal{A}$ and $\pi : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{B}$ defined by:

$$(3.9) \quad \begin{aligned} \iota(w_+) &= v_+ \otimes v_+ & \pi(v_+ \otimes v_+) &= w_+ \\ \iota(w_-) &= v_- \otimes v_- & \pi(v_+ \otimes v_-) &= \pi(v_- \otimes v_+) = 0 \\ & & \pi(v_- \otimes v_-) &= w_- \end{aligned}$$

Then \mathcal{B} canonically inherits from $\mathcal{A} \otimes \mathcal{A}$ of a structure of graded commutative Frobenius algebra with product, co-product, unit and co-unit $\mu_2, \delta_2, \eta_2, \varepsilon_2$ with respective degrees $-2, -2, 2, 2$ satisfying :

$$(3.10) \quad \mu_2 = \pi \circ \mu_1^\otimes \circ (\iota \otimes \iota) \quad \delta_2 = (\pi \otimes \pi) \circ \delta_1^\otimes \circ \iota \quad \eta_2 = \pi \circ \eta_1^\otimes \quad \varepsilon_2 = \varepsilon_1^\otimes \circ \iota.$$

The morphisms ι and π are adjoint with degree -2 . We denote by $\phi_2 : \mathcal{B} \otimes \mathcal{B} \rightarrow \mathcal{B} \otimes \mathcal{B}$ the flip morphism $\phi_2(b \otimes b') = b' \otimes b$, and $B : \mathcal{B} \rightarrow \mathcal{B}$ the identity morphism.

For each (non enhanced) state S of Γ , let's number all open components with $p \in \{1, \dots, op(S)\}$ and all closed components with $q \in \{1, \dots, cl(S)\}$. Then for an enhanced state \tilde{S} with underlying state S , we define the tensor product:

$$(3.11) \quad t(\tilde{S}) = \bigotimes_{p=1}^{op(S)} v_{\pm_{op}(p)} \otimes \bigotimes_{q=1}^{cl(S)} w_{\pm_{cl}(q)} \in \mathcal{A}^{\otimes op(S)} \otimes \mathcal{B}^{\otimes cl(S)}$$

where $\pm_{op}(p)$ and $\pm_{cl}(q)$ are the $+$ or $-$ signs of the p -th open and the q -th closed components of \tilde{S} respectively. The degree of $t(\tilde{S})$ does not correspond to the degree $j(\tilde{S})$ of \tilde{S} :

$$(3.12) \quad \deg(t(\tilde{S})) = \delta_{op}(\tilde{S}) + 2\delta_{cl}(\tilde{S}) = j(\tilde{S}) - k(\tilde{S}) = j(\tilde{S}) - k(S).$$

So we introduce the following definition:

Definition 3.4. *Translation of the degree of a graded vector space.*

Let $\mathcal{V} = \bigoplus_{j \in \mathbb{Z}} \mathcal{V}_j$ be a graded \mathbb{Z}_2 -vector space. The translated graded \mathbb{Z}_2 -vector space $\mathcal{V}\{\ell\}$ is defined by: $\mathcal{V}\{\ell\}_j = \mathcal{V}_{j-\ell}$.

Now we translate the degree of $t(\tilde{S})$ by $k(S)$ and we define:

$$(3.13) \quad \mathcal{C}(\Gamma) = (\mathcal{C}_i(\Gamma))_{i \in \mathbb{Z}} \quad \text{where} \quad \mathcal{C}_i(\Gamma) = \bigoplus_{\substack{S \in St(\Gamma) \\ i(S)=i}} (\mathcal{A}^{\otimes op(S)} \otimes \mathcal{B}^{\otimes cl(S)}) \{k(S)\}.$$

Proposition 3.5. *The map $t : \widetilde{St}(\Gamma) \rightarrow \mathcal{C}(\Gamma)$ defined by 3.11 extends to an isomorphism of complexes: $t : \llbracket \Gamma \rrbracket \rightarrow \mathcal{C}(\Gamma)$.*

Proof. The incidence relations T_i^\pm , $1 \leq i \leq 5$ of Figure 3.1, 3.2 induce morphisms of \mathbb{Z}_2 -vector spaces denoted by T_i , which have degree -2 :

$$(3.14) \quad \begin{aligned} T_1 &= \delta_1 \circ \mu_1 & T_3 &= \mu_1 \circ (A \otimes \mu_1) \circ (A \otimes \iota) & T_5 &= \mu_2 \\ T_2 &= (A \otimes \pi) \circ (A \otimes \delta_1) \circ \delta_1 & T_4 &= \delta_2. \end{aligned}$$

and the incidence relations T_6, T_7 of Figure 3.3 induce morphisms denoted by the same letters T_6, T_7 , which have degree -1 :

$$(3.15) \quad T_6 = \pi \circ \delta_1 \quad T_7 = \mu_1.$$

More precisely, we have:

$$(3.16) \quad \begin{aligned} T_1 : \mathcal{A} \otimes \mathcal{A} &\rightarrow (\mathcal{A} \otimes \mathcal{A}) & T_3 : \mathcal{A} \otimes \mathcal{B} &\rightarrow \mathcal{A} & T_5 : \mathcal{B} \otimes \mathcal{B} &\rightarrow \mathcal{B} \\ \left. \begin{array}{l} v_+ \otimes v_+ \mapsto v_+ \otimes v_- + v_- \otimes v_+ \\ v_+ \otimes v_- \\ v_- \otimes v_+ \end{array} \right\} &\mapsto v_- \otimes v_- & \left. \begin{array}{l} v_+ \otimes w_+ \mapsto v_+ \\ v_- \otimes w_+ \end{array} \right\} &\mapsto v_- & \left. \begin{array}{l} w_+ \otimes w_+ \mapsto w_+ \\ w_+ \otimes w_- \\ w_- \otimes w_+ \end{array} \right\} &\mapsto w_- \\ v_- \otimes v_- &\mapsto 0 & \left. \begin{array}{l} v_+ \otimes w_- \\ v_- \otimes w_- \end{array} \right\} &\mapsto 0 & w_- \otimes w_- &\mapsto 0 \\ \\ T_2 : \mathcal{A} &\rightarrow (\mathcal{A} \otimes \mathcal{B}) & T_4 : \mathcal{B} &\rightarrow (\mathcal{B} \otimes \mathcal{B}) \\ v_+ &\mapsto v_+ \otimes w_- & w_+ &\mapsto w_+ \otimes w_- + w_- \otimes w_+ \\ v_- &\mapsto v_- \otimes w_- & w_- &\mapsto w_- \otimes w_- \\ \\ T_6 : \mathcal{A} &\rightarrow \mathcal{B} & T_7 : \mathcal{A} \otimes \mathcal{A} &\rightarrow \mathcal{A} \\ v_+ &\mapsto 0 & \left. \begin{array}{l} v_+ \otimes v_+ \mapsto v_+ \\ v_+ \otimes v_- \\ v_- \otimes v_+ \end{array} \right\} &\mapsto v_- \\ v_- &\mapsto w_- & v_- \otimes v_- &\mapsto 0 \end{aligned}$$

Using these morphisms, we transfer the differential on $\llbracket \Gamma \rrbracket$ to a differential on $\mathcal{C}(\Gamma)$. Notice according to remark 3.2 that T_1 is self-adjoint and that T_2 and T_3 (resp. T_4 and T_5) are adjoint. Moreover, T_2 and T_4 are injective whereas T_3 and T_5 are surjective. Also T_7 is surjective. The relation $d \circ d = 0$ (induced by commutative diagrams (3.4) in the proof of proposition 3.3) is recovered using the following relations:

– symmetry properties:

$$\begin{aligned} T_1 \circ \phi_1 &= \phi_1 \circ T_1 = T_1 & T_4 &= \phi_2 \circ T_4 \\ (A \otimes T_2) \circ \phi_1 &= (\phi_1 \otimes B) \circ (A \otimes T_2) & T_5 &= T_5 \circ \phi_2 \\ \phi_1 \circ (A \otimes T_3) &= (A \otimes T_3) \circ (\phi_1 \otimes B) & T_7 &= T_7 \circ \phi_1 \end{aligned}$$

– commutativity properties corresponding to the splitting of two double points:

$$\begin{aligned} (T_1 \otimes A) \circ (A \otimes T_1) &= (A \otimes T_1) \circ (T_1 \otimes A) & (T_4 \otimes B) \circ T_4 &= (B \otimes T_4) \circ T_4 \\ (A \otimes T_2) \circ T_1 &= (T_1 \otimes B) \circ (A \otimes T_2) & T_5 \circ (T_5 \otimes B) &= T_5 \circ (B \otimes T_5) \\ T_1 \circ (A \otimes T_3) &= (A \otimes T_3) \circ (T_1 \otimes B) & T_4 \circ T_5 &= (B \otimes T_5) \circ (T_4 \otimes B) \end{aligned}$$

$$\begin{aligned}
T_1 \circ T_1 &= (A \otimes T_3) \circ (\phi_1 \otimes B) \circ (A \otimes T_2) = 0 \\
(T_2 \otimes B) \circ T_2 &= (A \otimes \phi_2) \circ (T_2 \otimes B) \circ T_2 = (A \otimes T_4) \circ T_2 \\
T_3 \circ (T_3 \otimes B) &= T_3 \circ (T_3 \otimes B) \circ (A \otimes \phi_2) = T_3 \circ (A \otimes T_5) \\
T_2 \circ T_3 &= (T_3 \otimes B) \circ (A \otimes \phi_2) \circ (T_2 \otimes B) \\
&= (T_3 \otimes B) \circ (A \otimes T_4) = (A \otimes T_5) \circ (T_2 \otimes B)
\end{aligned}$$

– commutativity properties corresponding to the splitting of a double point and a vertical tangent point:

$$\begin{aligned}
T_4 \circ T_6 &= (T_6 \otimes B) \circ T_2 & T_7 \circ (A \otimes T_3) &= T_3 \circ (T_7 \otimes B) \\
T_6 \circ T_3 &= T_5 \circ (T_6 \otimes B) & T_2 \circ T_7 &= (T_7 \otimes B) \circ (A \otimes T_2) \\
T_2 \circ T_7 &= (A \otimes T_6) \circ T_1 & T_7 \circ T_1 &= 0 = T_3 \circ (A \otimes T_6)
\end{aligned}$$

– commutativity properties corresponding to the splitting of two vertical tangent points:

$$T_1 \circ (A \otimes T_7) = (A \otimes T_7) \circ (T_1 \otimes A). \quad \square$$

Remark 3.6. *The units and co-units $\eta_1, \eta_2, \varepsilon_1, \varepsilon_2$ of \mathcal{A} and \mathcal{B} correspond respectively to the creation of a positive open component, the creation of a positive closed component, the destruction of a negative open component and the destruction of a negative closed component. Besides, $A \otimes \varepsilon_2$ and $B \otimes \varepsilon_2$ are left inverses of T_2 and T_4 whereas $A \otimes \eta_2$, $B \otimes \eta_2$ and $A \otimes \eta_1$ are right inverses of T_3 , T_5 and T_7 . In [Section 4](#), we will often refer to these morphisms together with the following ones:*

$$\begin{array}{ccc}
\bar{\eta}_1 : \mathbb{Z}_2 \rightarrow \mathcal{A} & \bar{\eta}_2 : \mathbb{Z}_2 \rightarrow \mathcal{B} & \bar{\varepsilon}_1 : \mathcal{A} \rightarrow \mathbb{Z}_2 & \bar{\varepsilon}_2 : \mathcal{B} \rightarrow \mathbb{Z}_2 \\
1 \mapsto v_- & 1 \mapsto w_- & v_+ \mapsto 1 & w_+ \mapsto 1 \\
& & v_- \mapsto 0 & w_- \mapsto 0
\end{array}$$

which correspond respectively to the creation of a negative open component, the creation of a negative closed component, the destruction of a positive open component and the destruction of a positive closed component, and to the following composed morphisms:

$$\begin{array}{ccc}
\tau = \eta_1 \varepsilon_1 : \mathcal{A} \rightarrow \mathcal{A} & \sigma = \bar{\eta}_1 \varepsilon_2 : \mathcal{B} \rightarrow \mathcal{A} \\
v_+ \mapsto 0 & w_+ \mapsto 0 \\
v_- \mapsto v_+ & w_- \mapsto v_-
\end{array}$$

3.3 Review of basic facts about complexes.

Let $\mathcal{C} := (\mathcal{C}, d) = (\mathcal{C}_i, d_i)_{i \in \mathbb{Z}}$ be a complex of \mathbb{Z}_2 -vector spaces. We denote by $\mathcal{H}(\mathcal{C})$ its homology:

$$(3.17) \quad \mathcal{H}(\mathcal{C}) = (H_i)_{i \in \mathbb{Z}} \quad H_i = \text{Ker } d_i / \text{Im } d_{i-1}.$$

A complex is *acyclic* if its homology is null.

Definition 3.7. *Shift of the grading of a complex.*

Let $(\mathcal{C}, d) = (\mathcal{C}_i, d_i)_{i \in \mathbb{Z}}$ be a complex of \mathbb{Z}_2 -vector spaces. We define the complex:

$$(\mathcal{C}, d)[k] = (\mathcal{C}[k], d[k]) \quad \text{by } \mathcal{C}[k]_i = \mathcal{C}_{i-k} \quad \text{and} \quad d[k]_i = d_{i-k}.$$

(If $(C, d) = (\bigoplus_{j \in \mathbb{Z}} C_{i,j}, \bigoplus_{j \in \mathbb{Z}} d_{i,j})_{i \in \mathbb{Z}}$ is a complex of graded \mathbb{Z}_2 -vector spaces, then we can translate twice the grading of the complex and the degree of the vector spaces :

$$(C, d)[k]\{\ell\} = (C, d)\{\ell\}[k] \quad \text{is defined by} \quad C[k]\{\ell\}_{i,j} = C_{i-k,j-\ell} \quad d[k]\{\ell\}_{i,j} = d_{i-k,j-\ell}.$$

A morphism of complexes of \mathbb{Z}_2 -vector spaces $f : (C^0, d^0) \rightarrow (C^1, d^1)$ is a sequence $f = (f_i)_{i \in \mathbb{Z}}$ of linear maps $f_i : C_i^0 \rightarrow C_i^1$ such that³: $f d^0 = d^1 f$ (i.e. $\forall i, d_i^1 f_i = f_{i+1} d_i^0$).

Definition 3.8. For a morphism of complexes $f : (C^0, d^0) \rightarrow (C^1, d^1)$, the cone of f is the complex denoted by $\text{Cone}(f) = (C_i, D_i)_{i \in \mathbb{Z}}$ and defined by :

$$(3.18) \quad C_i = C_i^0 \oplus C_{i-1}^1 = C_i^0 \oplus (C^1[1])_i \quad D_i = \begin{pmatrix} d_i^0 & 0 \\ f_i & d_{i-1}^1 \end{pmatrix}.$$

(Notice that (C^0, d^0) and $(C^1, d^1)[1]$ are sub-complexes of (C, D)).

A morphism of complexes $f : (C^0, d^0) \rightarrow (C^1, d^1)$ induces an isomorphism in homology if and only if $\text{Cone}(f)$ is acyclic. This is the case if f is a *homotopy equivalence* of complexes, i.e. there exist a morphism of complexes $g : (C^1, d^1) \rightarrow (C^0, d^0)$ and sequences $h^0 = (h_i^0)_{i \in \mathbb{Z}}, h^1 = (h_i^1)_{i \in \mathbb{Z}}$ of linear maps (homotopies) $h_i^0 : C_{i+1}^0 \rightarrow C_i^0$ and $h_i^1 : C_{i+1}^1 \rightarrow C_i^1$ such that:

$$(3.19) \quad gf = \text{id} + h^0 d^0 + d^0 h^0 \quad \text{and} \quad fg = \text{id} + h^1 d^1 + d^1 h^1.$$

(i.e. $\forall i \quad g_i f_i = \text{id} + h_i^0 d_i^0 + d_{i-1}^0 h_{i-1}^0$ and $f_i g_i = \text{id} + h_i^1 d_i^1 + d_{i-1}^1 h_{i-1}^1$)

Remark 3.9. As a particular case, if $h^0 = 0$, the complex (C^0, d^0) is called a *strong deformation retract* of (C^1, d^1) , with inclusion map f , retraction g and homotopy map h^1 . Besides, up to changing h^1 to a new homotopy h , we can always suppose that $hh = 0$, $hf = 0$ and $gh = 0$. We will assume these properties are always satisfied in the definition of *strong deformation retraction*.

Proposition 3.10. Let (\bar{C}^1, \bar{d}^1) be a strong deformation retract of (C^1, d^1) with retraction r , inclusion j and homotopy map h such that $hh = 0$, $rh = 0$, $hj = 0$. Let $f : (C^0, d^0) \rightarrow (C^1, d^1)$ be a morphism of complexes. Then $\text{Cone}(rf)$ is a strong deformation retract of $\text{Cone}(f)$ with retraction $R = \begin{pmatrix} \text{id} & 0 \\ 0 & r \end{pmatrix}$, inclusion $J = \begin{pmatrix} \text{id} & 0 \\ hf & j \end{pmatrix}$ and homotopy $H = \begin{pmatrix} 0 & 0 \\ 0 & h \end{pmatrix}$ such that $HH = 0$, $RH = 0$ and $HJ = 0$.

Proof. Immediate. □

A *double complex* (C, d, ∂) is a sequence of complexes $(C^k, d^k)_{k \in \mathbb{Z}}$ of \mathbb{Z}_2 -vector spaces and morphisms of complexes $(\partial^k)_{k \in \mathbb{Z}}$:

$$\dots \xrightarrow{\partial^{k-1}} (C^k, d^k) \xrightarrow{\partial^k} (C^{k+1}, d^{k+1}) \xrightarrow{\partial^{k+1}} (C^{k+2}, d^{k+2}) \xrightarrow{\partial^{k+2}} \dots$$

³Since we are working with \mathbb{Z}_2 field, commutativity and anti-commutativity coincide so that we have equivalently $d^1 f + f d^0 = 0$.

such that for all $k \in \mathbb{Z}$, $\partial^{k+1}\partial^k = 0$. A *morphism of two double complexes* is a sequence of morphisms of complexes $f = (f^k)_{k \in \mathbb{Z}}$:

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\bar{\partial}^{k-1}} & (\bar{\mathcal{C}}^k, \bar{d}^k) & \xrightarrow{\bar{\partial}^k} & (\bar{\mathcal{C}}^{k+1}, \bar{d}^{k+1}) & \xrightarrow{\bar{\partial}^{k+1}} & (\bar{\mathcal{C}}^{k+2}, \bar{d}^{k+2}) \xrightarrow{\bar{\partial}^{k+2}} \cdots \\ & & \downarrow f^k & & \downarrow f^{k+1} & & \downarrow f^{k+2} \\ \cdots & \xrightarrow{\partial^{k-1}} & (\mathcal{C}^k, d^k) & \xrightarrow{\partial^k} & (\mathcal{C}^{k+1}, d^{k+1}) & \xrightarrow{\partial^{k+1}} & (\mathcal{C}^{k+2}, d^{k+2}) \xrightarrow{\partial^{k+2}} \cdots \end{array}$$

such that for all $k \in \mathbb{Z}$, $f^{k+1}\bar{\partial}^k = \partial^k f^k$. We also have notions of homotopy equivalence and strong deformation retraction for double complex. A morphism of double complexes $f : (\bar{\mathcal{C}}, \bar{d}, \bar{\partial}) \rightarrow (\mathcal{C}, d, \partial)$ is a *homotopy equivalence (of double complexes)* if there exists a morphism of double complexes $g : (\mathcal{C}, d, \partial) \rightarrow (\bar{\mathcal{C}}, \bar{d}, \bar{\partial})$ and homotopy maps $\bar{h} = (\bar{h}^k)_{k \in \mathbb{Z}}$, $h = (h^k)_{k \in \mathbb{Z}}$ (sequences of morphisms of complexes) $\bar{h}^k : (\bar{\mathcal{C}}^{k+1}, \bar{d}^{k+1}) \rightarrow (\bar{\mathcal{C}}^k, \bar{d}^k)$ and $\bar{h}^k : (\mathcal{C}^{k+1}, d^{k+1}) \rightarrow (\mathcal{C}^k, d^k)$ such that for all k :

$$g^k f^k = \text{id} + \bar{h}^k \bar{\partial}^k + \bar{\partial}^{k-1} \bar{h}^{k-1} \quad \text{and} \quad f^k g^k = \text{id} + h^k \partial^k + \partial^{k-1} h^{k-1}.$$

If $\bar{h} = 0$, $(\bar{\mathcal{C}}, \bar{d}, \bar{\partial})$ is called a *strong deformation retract* of the double complex $(\mathcal{C}, d, \partial)$ with inclusion f and retraction g . Again, up to changing the homotopy h , we assume that it satisfies: $hh = 0$, $hf = 0$ and $gh = 0$.

Now we extend the definition of cone to a finite sequence of morphisms of complexes. A double complex $(\mathcal{C}, d, \partial)$ is ∂ -finite if (\mathcal{C}^k, d^k) is trivial except for a finite number of values of k .

Definition 3.11. Let $(\mathcal{C}^0, d^0) \xrightarrow{\partial^0} \cdots \xrightarrow{\partial^{n-1}} (\mathcal{C}^n, d^n)$ be a ∂ -finite double complex. Let's denote:

$$\begin{aligned} \tilde{\partial}_i^0 : \mathcal{C}_i^0 &\rightarrow \mathcal{C}_i^1 \oplus \mathcal{C}_{i-1}^2 \oplus \cdots \oplus \mathcal{C}_{i-n+1}^n \\ u &\mapsto (\partial_i^0(u), 0, \dots, 0) \end{aligned}$$

Then the cone of $(\partial^0, \dots, \partial^{n-1})$ is the complex defined by:

$$(3.20) \quad \text{Cone}(\partial^0, \dots, \partial^{n-1}) = \text{Cone}(\tilde{\partial}^0, \text{Cone}(\partial^1, \dots, \partial^{n-1})).$$

Suppose that $f = (f^k)_{0 \leq k \leq n}$ is a morphism from a $\bar{\partial}$ -finite double complexes $(\bar{\mathcal{C}}, \bar{d}, \bar{\partial})$ to a ∂ -finite $(\mathcal{C}, d, \partial)$:

$$(\bar{\mathcal{C}}^0, \bar{d}^0) \xrightarrow{\bar{\partial}^0} \cdots \xrightarrow{\bar{\partial}^{n-1}} (\bar{\mathcal{C}}^n, \bar{d}^n) \quad \text{and} \quad (\mathcal{C}^0, d^0) \xrightarrow{\partial^0} \cdots \xrightarrow{\partial^{n-1}} (\mathcal{C}^n, d^n).$$

Let's set $F_i = f_i^0 \oplus f_{i-1}^1 \oplus \cdots \oplus f_{i-n}^n$. Then f induces a morphism of complexes:

$$C(f) = (F_i)_{i \in \mathbb{Z}} : \text{Cone}(\bar{\partial}^0, \dots, \bar{\partial}^{n-1}) \longrightarrow \text{Cone}(\partial^0, \dots, \partial^{n-1}).$$

If f^k are isomorphisms, $C(f)$ is also an isomorphism.

Proposition 3.12. If $(\bar{\mathcal{C}}, \bar{d}, \bar{\partial})$ is a $\bar{\partial}$ -finite double complex, $(\mathcal{C}, d, \partial)$ a ∂ -finite double complex and $f : (\bar{\mathcal{C}}, \bar{d}, \bar{\partial}) \rightarrow (\mathcal{C}, d, \partial)$ a homotopy equivalence with inverse g , then:

$$C(f) : \text{Cone}(\bar{\partial}^0, \dots, \bar{\partial}^{n-1}) \longrightarrow \text{Cone}(\partial^0, \dots, \partial^{n-1})$$

is a homotopy equivalence of complexes with inverse $C(g)$. So $C(f)$ induces an isomorphism in homology.

Furthermore if $(\bar{\mathcal{C}}, \bar{d}, \bar{\partial})$ is a strong deformation retract of $(\mathcal{C}, d, \partial)$ with inclusion map f and retraction g then $\text{Cone}(\bar{\partial}^0, \dots, \bar{\partial}^{n-1})$ is a strong deformation retract of $\text{Cone}(\partial^0, \dots, \partial^{n-1})$ with inclusion map $C(f)$ and retraction $C(g)$, and so $C(f)$ induces an isomorphism in homology.

Proof. Let $\bar{h} = (\bar{h}^k)_{1 \leq k < n}$ and $h = (h^k)_{0 \leq k < n}$ be homotopies associated with f and g . Then we have :

$$\begin{aligned} \bar{h}_{i+1}^{k-1} \bar{d}_i^k &= \bar{d}_i^{k-1} \bar{h}_i^{k-1} & \bar{h}_i^k \bar{\partial}_i^k + \bar{\partial}_i^{k-1} \bar{h}_i^{k-1} + \text{id} &= g_i^k f_i^k \\ h_{i+1}^{k-1} d_i^k &= d_i^{k-1} h_i^{k-1} & h_i^k \partial_i^k + \partial_i^{k-1} h_i^{k-1} + \text{id} &= f_i^k g_i^k \end{aligned}$$

Let $H = (H_i)_{i \in \mathbb{Z}}$ be the sequence of linear maps defined by:

$$\begin{aligned} H_i : \mathcal{C}_{i+1}^0 \oplus \mathcal{C}_i^1 \oplus \cdots \oplus \mathcal{C}_{i-n+1}^n &\longrightarrow \mathcal{C}_i^0 \oplus \mathcal{C}_{i-1}^1 \oplus \cdots \oplus \mathcal{C}_{i-n}^n \\ (x_0, x_1, \dots, x_n) &\longmapsto (h_i^0(x_1), h_{i-1}^1(x_2), \dots, h_{i-n+1}^{n-1}(x_n), 0) \end{aligned}$$

and $\bar{H} = (\bar{H}_i)_{i \in \mathbb{Z}}$ defined analogously on $(\bar{\mathcal{C}}, \bar{d}, \bar{\partial})$. Then if \bar{D} and D are the differentials of $\text{Cone}(\bar{\partial}^0, \dots, \bar{\partial}^{n-1})$ and $\text{Cone}(\partial^0, \dots, \partial^{n-1})$, we have:

$$FG = \text{id} + HD + DH \quad \text{and} \quad GF = \text{id} + \bar{H}\bar{D} + \bar{D}\bar{H}. \quad \square$$

3.4 Fundamental splitting lemmas.

Let (Γ, ϵ) be an OMS-divide or a cuspidal divide. Let p be a double point or a vertical tangent point of Γ . Let Γ^0 and Γ^1 be the cuspidal divides obtained from Γ by applying Θ_0 and Θ_1 at p respectively. Then each enhanced state of Γ can be identified with either an enhanced state of Γ^0 or Γ^1 i.e.:

$$(3.21) \quad \widetilde{St}(\Gamma)^{1-1} \simeq \widetilde{St}(\Gamma^0) \sqcup \widetilde{St}(\Gamma^1).$$

Consequently, $\llbracket \Gamma^0 \rrbracket$ and $\llbracket \Gamma^1 \rrbracket$ can be seen as sub-complexes of $\llbracket \Gamma \rrbracket$ up to translations of the grading i and the degree j . More precisely, we have :

Lemma 3.13. *Let d be the differential of $\llbracket \Gamma \rrbracket$.*

1. *If p is a positive double point then d induces the differentials d^0 and d^1 of $\llbracket \Gamma^0 \rrbracket \{2\}$ and $\llbracket \Gamma^1 \rrbracket \{4\}$ and a morphism $\llbracket \Gamma^0 \rrbracket \{2\} \xrightarrow{d^\bullet} \llbracket \Gamma^1 \rrbracket \{4\}$ such that $\llbracket \Gamma \rrbracket = \text{Cone}(d^\bullet)$.*
2. *If p is a negative double point then d induces the differentials d^0 and d^1 of $\llbracket \Gamma^0 \rrbracket \{-2\}$ and $\llbracket \Gamma^1 \rrbracket \{-4\}$ and a morphism $\llbracket \Gamma^1 \rrbracket \{-4\} \xrightarrow{d^\bullet} \llbracket \Gamma^0 \rrbracket \{-2\}$ such that $\llbracket \Gamma \rrbracket = \text{Cone}(d^\bullet)[-1]$.*
3. *If p is vertical tangent point, then d induces the differentials d^0 and d^1 of $\llbracket \Gamma^0 \rrbracket \{1\}$ and $\llbracket \Gamma^1 \rrbracket \{2\}$ and a morphism $\llbracket \Gamma^0 \rrbracket \{1\} \xrightarrow{d^\bullet} \llbracket \Gamma^1 \rrbracket \{2\}$ such that $\llbracket \Gamma \rrbracket = \text{Cone}(d^\bullet)$.*

Proof. Suppose that p is a positive double point of Γ . Then Γ^0 and Γ^1 have one positive double point less than Γ so that the writhes of Γ , Γ^0 and Γ^1 are related by:

$$w(\Gamma) = w(\Gamma^0) + 2 = w(\Gamma^1) + 2$$

Let \tilde{S} be an enhanced state of $\widetilde{St}_i(\Gamma)$ with degree $j = j(\tilde{S})$. If \tilde{S} is obtained from Γ using Θ_0 (resp. Θ_1) at p , then \tilde{S} can be seen as an enhanced state of $\widetilde{St}_i(\Gamma^0)$ with degree $j - 2$ (resp. of $\widetilde{St}_{i-1}(\Gamma^1)$ with degree $j - 4$). Besides, if $\tilde{S}_0 \in \widetilde{St}_i(\Gamma)$ and $\tilde{S}_1 \in \widetilde{St}_{i+1}(\Gamma)$ are adjacent enhanced states of degree j then it involves three cases:

- either $\tilde{S}_0 \in \tilde{S}t_i(\Gamma^0)$ and $\tilde{S}_1 \in \tilde{S}t_{i+1}(\Gamma^0)$ are adjacent enhanced states of Γ^0 with degrees $j-2$, so the differential d^0 of $[[\Gamma^0]]\{2\}$ coincide with the restriction of d to $[[\Gamma^0]]\{2\}$;
- either $\tilde{S}_0 \in \tilde{S}t_{i-1}(\Gamma^1)$ and $\tilde{S}_1 \in \tilde{S}t_i(\Gamma^1)$ are adjacent enhanced states of Γ^1 with degrees $j-4$, so the differential d^1 of $[[\Gamma^1]]\{4\}[1]$ coincide with the restriction of d to $[[\Gamma^1]]\{4\}[1]$;
- or $\tilde{S}_0 \in \tilde{S}t_i(\Gamma^0)$ with degree $j-2$ and $\tilde{S}_1 \in \tilde{S}t_i(\Gamma^1)$ with degree $j-4$, then d induces a map $d^\bullet : [[\Gamma^0]]\{2\} \rightarrow [[\Gamma^1]]\{4\}$ which is a morphism of complexes since from the proof of proposition 3.3 $dd = 0$ implies $d^\bullet d^0 = d^1 d^\bullet$.

Hence $[[\Gamma]] = \text{Cone}(d^\bullet)$. Similar arguments hold for the two other cases. \square

More generally, consider $k = k_+ + k_- + k_0$ double vertical tangent points p_1, \dots, p_k such that the k_+ first ones are positive double points, the next k_- ones negative double points and the last k_0 ones vertical tangent points. For each words $[a] = a_1 a_2 \dots a_{k_+}$, $a_i \in \{0, 1\}$, $[b] = b_1 b_2 \dots b_{k_-}$, $b_i \in \{0, 1\}$, $[c] = c_1 c_2 \dots c_{k_0}$, $c_i \in \{0, 1\}$, let $[a][b][c]$ be the word obtained by concatenation of $[a]$, $[b]$, $[c]$ and denote by $(\Gamma^{[a][b][c]}, \epsilon^{[a][b][c]})$ the cuspidal divide obtained from (Γ, ϵ) by performing:

$$\begin{cases} \Theta_{a_i} & \text{splitting at } p_i & \text{for } 1 \leq i \leq k_+ \\ \Theta_{b_i} & \text{splitting at } p_i & \text{for } k_+ < i \leq k_+ + k_- \\ \Theta_{c_i} & \text{splitting at } p_i & \text{for } k_+ + k_- < i \leq k = k_+ + k_- + k_0. \end{cases}$$

Let $1_{[a]}$, $1_{[b]}$ and $1_{[c]}$ be the numbers of occurrences of 1 in $[a]$, $[b]$ and $[c]$ and $\text{gr}([a][b][c]) = 1_{[a]} - 1_{[b]} + 1_{[c]}$. By restriction, the differential d of $[[\Gamma]]$ coincide with the differential $d^{[a][b][c]}$ of $[[\Gamma^{[a][b][c]}]]$. By iterating lemma 3.13, using same arguments, just following the incidence relations, we have:

Lemma 3.14. *For each ℓ , $-k_- \leq \ell \leq k_+ + k_0$, we can identify the complex:*

$$(\mathcal{C}^\ell, D^\ell) = \bigoplus_{\text{gr}([a][b][c])=\ell} [[\Gamma^{[a][b][c]}]]\{2(1_{[a]} - 1_{[b]} + k_+ - k_-) + 1_{[c]} + k_0\}$$

with a sub-complex of $[[\Gamma]]$, with differential $D^\ell = \bigoplus_{\text{gr}([a][b][c])=\ell} d^{[a][b][c]}$. The differential d induces a structure of double complex :

$$(\mathcal{C}^{-k_-}, D^{-k_-}) \xrightarrow{\Delta^{-k_-}} (\mathcal{C}^{-k_-+1}, D^{-k_-+1}) \xrightarrow{\Delta^{-k_-+1}} \dots \xrightarrow{\Delta^{k_++k_0-1}} (\mathcal{C}^{k_++k_0}, D^{k_++k_0})$$

such that $[[\Gamma]] = \text{Cone}(\Delta^{-k_-}, \dots, \Delta^{k_++k_0-1})[-k_-]$.

In the sequel, such a double complex will be called a *splitting diagram* of $[[\Gamma]]$.

3.5 Khovanov homology of OMS-divides.

Definition 3.15. *We call Khovanov homology $\mathcal{H}(\Gamma)$ of an OMS-divide (or a cuspidal divide) $\Gamma = (\Gamma, \epsilon)$ the homology of the complex $[[\Gamma]] = ([[\Gamma]]_i)_{i \in \mathbb{Z}}$:*

$$(3.22) \quad \mathcal{H}(\Gamma) = (\mathcal{H}_i(\Gamma))_{i \in \mathbb{Z}} \quad \mathcal{H}_i(\Gamma) = \bigoplus_{j \in \mathbb{Z}} \mathcal{H}_{i,j}(\Gamma) \quad \mathcal{H}_{i,j}(\Gamma) = \text{Ker } d_{i,j} / \text{Im } d_{i-1,j}.$$

Proposition 3.16. *If $\Gamma = (\Gamma, \epsilon)$ is an OMS-divide, then the polynomial W_Γ and the graded Euler characteristics of $\mathcal{H}(\Gamma)$ are related by:*

$$(3.23) \quad W_\Gamma(t^2) = \frac{t}{1+t^2} \chi_{gr}(\mathcal{H}(\Gamma)) = \frac{t}{1+t^2} \sum_{i \in \mathbb{Z}} (-1)^i \dim_{gr} \mathcal{H}_i(\Gamma)$$

where the graded dimension is: $\dim_{gr} \mathcal{H}_i(\Gamma) = \sum_{j \in \mathbb{Z}} t^j \dim_{\mathbb{Z}_2} \mathcal{H}_{i,j}(\Gamma)$.

Proof. Immediate from formula (2.5). □

We can now formulate our main theorem:

Theorem 3.17. *Khovanov homology of OMS-divides is invariant under \mathcal{M} -equivalence.*

Combined with theorem 1.2, we obtain:

Corollary 3.18. *Khovanov homology of OMS-divides is an invariant for strong equivalence of strongly invertible links.*

Section 4 is devoted to the proof of this theorem. Notice that from proposition 3.16, this theorem 3.17 is a refinement of proposition 2.3.

3.6 Examples.

(1) Figure 3.4 shows a divide for the link 3_1 , and its splitting diagram.

The associated complex and homology entries are:

$i \setminus j$	1	3	5	7	9
0	\mathbb{Z}_2	\mathbb{Z}_2			
1			\mathbb{Z}_2	\mathbb{Z}_2	
2				\mathbb{Z}_2	\mathbb{Z}_2

$$(\mathcal{A} \otimes \mathcal{A})\{3\} \xrightarrow{d^0} (\mathcal{A} \otimes \mathcal{A})\{5\} \oplus \mathcal{A}\{4\} \xrightarrow{d^1} (\mathcal{A} \otimes \mathcal{B})\{6\}$$

(2) Figure 3.5 shows a divide for the link 4_1 and its splitting diagram. The associated complex is:

$$(\mathcal{A}^{\otimes 3})\{-2\} \xrightarrow{d^{-1}} (\mathcal{A}^{\otimes 3}) \oplus (\mathcal{A}^{\otimes 2} \oplus \mathcal{A}^{\otimes 2})\{-1\} \xrightarrow{d^0} (\mathcal{A}^{\otimes 2} \oplus \mathcal{A}^{\otimes 2})\{1\} \oplus \mathcal{A} \xrightarrow{d^1} (\mathcal{A} \otimes \mathcal{B})\{2\}$$

and homology entries:

$i \setminus j$	-5	-3	-1	1	3	5
-1	\mathbb{Z}_2	\mathbb{Z}_2				
0		\mathbb{Z}_2	$(\mathbb{Z}_2)^2$	\mathbb{Z}_2		
1				\mathbb{Z}_2	\mathbb{Z}_2	
2					\mathbb{Z}_2	\mathbb{Z}_2




Figure 3.4:




Figure 3.5:

4 Invariance under \mathcal{M} -equivalence.

4.1 Invariance under type I moves.

Let Γ and $\tilde{\Gamma}$ be OMS-divides which differ only by a type I move (see Figure 4.1).



Figure 4.1: Type I move.

Proposition 4.1. *The complexes $[[\Gamma]]$ and $[[\tilde{\Gamma}]]$ have the same homology.*

Let's denote by Γ^{st} (resp. $\tilde{\Gamma}^{st}$), $s, t \in \{0, 1\}$ the cuspidal divides obtained by performing Θ_s, Θ_t splittings respectively at the $+$ double point and the vertical tangent point of Γ (resp. of $\tilde{\Gamma}$) in Figure 4.1, without changing any other singular point of these divides. From lemma 3.14, we have splitting diagrams given in Figure 4.2.




Figure 4.2: Splitting diagram for type I move.

Notice that $[[\Gamma^{00}]] = [[\tilde{\Gamma}^{00}]]$. In other words we have:

Lemma 4.2. *Let's denote $\Delta^0 = \begin{pmatrix} d^{0\bullet} \\ d^{\bullet 0} \end{pmatrix}$, $\Delta^1 = \begin{pmatrix} d^{\bullet 1} & d^{1\bullet} \end{pmatrix}$, $\tilde{\Delta}^0 = \begin{pmatrix} \tilde{d}^{0\bullet} \\ \tilde{d}^{\bullet 0} \end{pmatrix}$ and $\tilde{\Delta}^1 = \begin{pmatrix} \tilde{d}^{\bullet 1} & \tilde{d}^{1\bullet} \end{pmatrix}$. Then $[[\Gamma]] = \text{Cone}(\Delta^0, \Delta^1)$ and $[[\tilde{\Gamma}]] = \text{Cone}(\tilde{\Delta}^0, \tilde{\Delta}^1)$:*

$$(4.1) \quad \begin{array}{c} [[\Gamma^{00}]]\{3\} \xrightarrow{\Delta^0} [[\Gamma^{01}]]\{4\} \oplus [[\Gamma^{10}]]\{5\} \xrightarrow{\Delta^1} [[\Gamma^{11}]]\{6\} \\ \parallel \\ [[\tilde{\Gamma}^{00}]]\{3\} \xrightarrow{\tilde{\Delta}^0} [[\tilde{\Gamma}^{01}]]\{4\} \oplus [[\tilde{\Gamma}^{10}]]\{5\} \xrightarrow{\tilde{\Delta}^1} [[\tilde{\Gamma}^{11}]]\{6\} \end{array}$$

Proof. The morphism $d^{\bullet 1}$ corresponds to T_2 or T_4 and the morphism $d^{1\bullet}$ to T_3 or T_5 in (3.14). The lemma is a direct consequence of remark 3.6 (see also Figure 3.1 and Figure 3.2). \square

Proof of proposition 4.4. Consider the diagram:

$$\begin{array}{ccc} \llbracket \Gamma^{00} \rrbracket = \llbracket \Gamma_0 \rrbracket & & J = \begin{pmatrix} \text{id} \\ \eta_2 d^{\bullet 0} \end{pmatrix} \quad R = (\text{id} \quad d^{0\bullet} \varepsilon_2) \\ R \uparrow \downarrow J & & \\ \llbracket \Gamma^{01} \rrbracket \{-2\} \xrightarrow[H^0]{\Delta^0} \llbracket \Gamma^{00} \rrbracket \oplus \llbracket \Gamma^{11} \rrbracket \xrightarrow[H^1]{\Delta^1} \llbracket \Gamma^{10} \rrbracket \{2\} & & H^0 = (0 \quad \varepsilon_2) \quad H^1 = \begin{pmatrix} 0 \\ \eta_2 \end{pmatrix} \end{array}$$

From the previous two lemmas, $\llbracket \Gamma_0 \rrbracket$ is a strong deformation retract of $\llbracket \Gamma \rrbracket = \text{Cone}(\Delta^0, \Delta^1)[-1]$ with retraction R , inclusion J , and homotopy (H^0, H^1) :

$$R\Delta^0 = 0, \quad \Delta^1 J = 0, \quad H^0 \Delta^0 = \text{id}, \quad R J = \text{id}, \quad J R = \text{id} + \Delta^0 H^0 + H^1 \Delta^1, \quad \Delta^1 H^1 = \text{id}.$$

Hence from proposition 3.12 they have the same homology. \square

4.3 Invariance under type III move.

In this section, we only consider the case of move III_+ . The case of III_- can be checked in a similar way: we have dual situations as is said in remark 3.2 and in the proof of proposition 3.5.

Let Γ_1 and Γ_2 be OMS-divides which differ only by a type III_+ move (see Figure 4.7).



Figure 4.7: Type III_+ move.

Proposition 4.7. *The complexes $\llbracket \Gamma_1 \rrbracket$ and $\llbracket \Gamma_2 \rrbracket$ have the same homology.*

Let Γ_1^{stu} , $s, t, u \in \{0, 1\}$ be the cuspidal divides obtained by performing Θ_s , Θ_t and Θ_u splittings at the double points shown on the figure of Γ_1 (see Figure 4.7). From lemma 3.14, we have the following splitting diagram of $\llbracket \Gamma_1 \rrbracket$ (see Figure 4.8):

$$\begin{array}{ccccccc} & & \llbracket \Gamma_1^{100} \rrbracket \{8\} = \llbracket \text{diagram} \rrbracket \{8\} & \xrightarrow{d^{1\bullet 0}} & \llbracket \Gamma_1^{110} \rrbracket \{10\} = \llbracket \text{diagram} \rrbracket \{10\} & & \\ & \nearrow^{d^{\bullet 00}} & \oplus & \nearrow^{d^{10\bullet}} & \oplus & \searrow^{d^{11\bullet}} & \\ \llbracket \Gamma_1^{000} \rrbracket \{6\} = \llbracket \text{diagram} \rrbracket \{6\} & \xrightarrow{d^{0\bullet 0}} & \llbracket \Gamma_1^{010} \rrbracket \{8\} = \llbracket \text{diagram} \rrbracket \{8\} & \xrightarrow{d^{\bullet 10}} & \llbracket \Gamma_1^{101} \rrbracket \{10\} = \llbracket \text{diagram} \rrbracket \{10\} & \xrightarrow{d^{1\bullet 1}} & \llbracket \Gamma_1^{111} \rrbracket \{12\} = \llbracket \text{diagram} \rrbracket \{12\} \\ & \searrow^{d^{00\bullet}} & \oplus & \searrow^{d^{01\bullet}} & \oplus & \nearrow^{d^{\bullet 11}} & \\ & & \llbracket \Gamma_1^{001} \rrbracket \{8\} = \llbracket \text{diagram} \rrbracket \{8\} & \xrightarrow{d^{0\bullet 1}} & \llbracket \Gamma_1^{011} \rrbracket \{10\} = \llbracket \text{diagram} \rrbracket \{10\} & & \end{array}$$

Figure 4.8: Splitting diagram for type III_+ move.

Let's denote $\mathcal{C}^0 = \llbracket \Gamma_1^{000} \rrbracket \{6\}$, $\mathcal{C}^3 = \llbracket \Gamma_1^{111} \rrbracket \{12\}$, $\mathcal{C}^1 = \hat{\mathcal{C}}^1 \oplus \check{\mathcal{C}}^1$ and $\mathcal{C}^2 = \hat{\mathcal{C}}^2 \oplus \check{\mathcal{C}}^2$

where:

$$\begin{array}{ll} \hat{\mathcal{C}}^1 = \llbracket \Gamma_1^{100} \rrbracket \{8\} \oplus \llbracket \Gamma_1^{010} \rrbracket \{8\} & \check{\mathcal{C}}^1 = \llbracket \Gamma_1^{001} \rrbracket \{8\} \\ \hat{\mathcal{C}}^2 = \llbracket \Gamma_1^{110} \rrbracket \{10\} \oplus \llbracket \Gamma_1^{101} \rrbracket \{10\} & \check{\mathcal{C}}^2 = \llbracket \Gamma_1^{011} \rrbracket \{10\}. \end{array}$$

Lemma 4.8. $\llbracket \Gamma_1 \rrbracket = \text{Cone}(\Delta^0, \Delta^1, \Delta^2)$ where: $\mathcal{C}^0 \xrightarrow{\Delta^0} \mathcal{C}^1 \xrightarrow{\Delta^1} \mathcal{C}^2 \xrightarrow{\Delta^2} \mathcal{C}^3$ and:

$$\Delta^0 = \begin{pmatrix} d^{\bullet 00} \\ d^{0 \bullet 0} \\ d^{00 \bullet} \end{pmatrix} \quad \Delta^1 = \begin{pmatrix} d^{1 \bullet 0} & d^{\bullet 10} & 0 \\ d^{10 \bullet} & 0 & d^{\bullet 01} \\ 0 & d^{01 \bullet} & d^{0 \bullet 1} \end{pmatrix} \quad \Delta^2 = (d^{11 \bullet} \quad d^{1 \bullet 1} \quad d^{\bullet 11}).$$

Now we can use the same arguments as in Section 4.2. The morphism $d^{0 \bullet 1}$ (corresponding to T_2 and T_4) is injective and $d^{\bullet 11}$ (corresponding to T_3 and T_5) is surjective. Let $\varepsilon_2 : \check{\mathcal{C}}^2 \rightarrow \check{\mathcal{C}}^1$ be the “destruction” morphism of complexes and $\eta_2 : \mathcal{C}^3 \rightarrow \check{\mathcal{C}}^2$ be the “creation” morphism of complexes (see Figure 4.9).

Figure 4.9:

Lemma 4.9. The sequence: $0 \rightarrow \mathcal{C}^3 \xrightarrow{\eta_2} \check{\mathcal{C}}^2 \xrightarrow{\varepsilon_2} \check{\mathcal{C}}^1 \rightarrow 0$ is exact and $d^{\bullet 11}$, $d^{0 \bullet 1}$ are respectively sections of η_2 and ε_2 :

$$(4.6) \quad d^{\bullet 11} \eta_2 = \text{id}, \quad \varepsilon_2 d^{0 \bullet 1} = \text{id}, \quad d^{0 \bullet 1} \varepsilon_2 + \eta_2 d^{\bullet 11} = \text{id} + \eta_2 d^{\bullet 11} d^{0 \bullet 1} \varepsilon_2.$$

Proof. We have a similar situation as in lemma 4.6: $d^{0 \bullet 1}$ corresponds to T_2 or T_4 and $d^{\bullet 11}$ corresponds to T_3 or T_5 in (3.14). The lemma is a direct consequence of remark 3.6 (see also Figure 3.1). \square

Lemma 4.10. Let $\delta = d^{\bullet 01} \varepsilon_2 d^{01 \bullet} : \llbracket \Gamma_1^{010} \rrbracket \{8\} \rightarrow \llbracket \Gamma_1^{101} \rrbracket \{10\}$.

The following sequence is a double-complex:

$$(\mathcal{C}^0, d^0) \xrightarrow{\hat{\Delta}^0} (\hat{\mathcal{C}}^1, \hat{d}^1) \xrightarrow{\hat{\Delta}^1} (\hat{\mathcal{C}}^2, \hat{d}^2) \quad \text{where} \quad \hat{\Delta}^0 = \begin{pmatrix} d^{\bullet 00} \\ d^{0 \bullet 0} \end{pmatrix} \quad \hat{\Delta}^1 = \begin{pmatrix} d^{1 \bullet 0} & d^{\bullet 10} \\ d^{10 \bullet} & \delta \end{pmatrix}.$$

Proof. Since $\Delta^1 \Delta^0 = 0$, we obtain from lemma 4.9:

$$\hat{\Delta}^1 \hat{\Delta}^0 = \begin{pmatrix} d^{1 \bullet 0} d^{\bullet 00} + d^{\bullet 10} d^{0 \bullet 0} \\ d^{10 \bullet} d^{\bullet 00} + \delta d^{0 \bullet 0} \end{pmatrix} = \begin{pmatrix} 0 \\ d^{10 \bullet} d^{\bullet 00} + d^{\bullet 01} \varepsilon_2 d^{01 \bullet} d^{0 \bullet 0} \end{pmatrix}$$

and: $d^{10 \bullet} d^{\bullet 00} + d^{\bullet 01} \varepsilon_2 d^{01 \bullet} d^{0 \bullet 0} = d^{\bullet 01} d^{00 \bullet} + d^{\bullet 01} \varepsilon_2 d^{0 \bullet 1} d^{00 \bullet} = 0.$ \square

Lemma 4.11. Let $\hat{\mathcal{C}} = \text{Cone}(\hat{\Delta}^0, \hat{\Delta}^1)$. Then the complex $\hat{\mathcal{C}}$ is a strong deformation retract of $\llbracket \Gamma_1 \rrbracket$ and so they have the same homology.

Proof. Consider the diagram:

$$\begin{array}{ccccc} \mathcal{C}^0 & \xrightarrow{\hat{\Delta}^0} & \hat{\mathcal{C}}^1 & \xrightarrow{\hat{\Delta}^1} & \hat{\mathcal{C}}^2 \\ \downarrow \text{id} & & R^1 \uparrow \downarrow J^1 & & R^2 \uparrow \downarrow J^2 \\ \mathcal{C}^0 & \xrightarrow{\Delta^0} & \hat{\mathcal{C}}^1 \oplus \check{\mathcal{C}}^1 & \xrightarrow[\cong]{\Delta^1} & \hat{\mathcal{C}}^2 \oplus \check{\mathcal{C}}^2 & \xrightarrow[\cong]{\Delta^2} & \mathcal{C}^3 \\ & & & \cong \uparrow H^1 & & \cong \uparrow H^2 & \end{array}$$

$$\begin{aligned}
J^1 &= \begin{pmatrix} \text{id} & 0 \\ 0 & \text{id} \\ 0 & \varepsilon_2 d^{01\bullet} \end{pmatrix} & R^1 &= \begin{pmatrix} \text{id} & 0 & 0 \\ 0 & \text{id} & 0 \end{pmatrix} & H^1 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \varepsilon_2 \end{pmatrix} \\
J^2 &= \begin{pmatrix} \text{id} & 0 \\ 0 & \text{id} \\ \eta_2 d^{11\bullet} & \eta_2 d^{1\bullet 1} \end{pmatrix} & R^2 &= \begin{pmatrix} \text{id} & 0 & 0 \\ 0 & \text{id} & d^{\bullet 01} \varepsilon_2 \end{pmatrix} & H^2 &= \begin{pmatrix} 0 \\ 0 \\ \eta_2 \end{pmatrix}
\end{aligned}$$

From lemma 4.9, we easily verify the relations:

$$\begin{aligned}
J^1 \widehat{\Delta}^0 &= \Delta^0, & J^2 \widehat{\Delta}^1 &= \Delta^1 J^1, & \Delta^2 J^2 &= 0, & R^1 \Delta^0 &= \widehat{\Delta}^0, & R^2 \Delta^1 &= \widehat{\Delta}^1 R^1, & R^1 J^1 &= \text{id} \\
R^2 J^2 &= \text{id}, & J^1 R^1 &= \text{id} + H^1 \Delta^1, & J^2 R^2 &= \text{id} + H^2 \Delta^2 + \Delta^1 H^1, & 0 &= \text{id} + \Delta^2 H^2.
\end{aligned}$$

So downward arrows define an inclusion map, upward ones a retraction and H_1, H_2 homotopy maps of double complexes. We can apply proposition 3.12: $\widehat{\mathcal{C}}$ is a deformation retract of $\llbracket \Gamma_1 \rrbracket$ so they have the same homology. \square

Proof of proposition 4.7. Up to isotopy, the drawn part of Γ_1^{000} on Figure 4.8 is symmetric with respect to horizontal direction. Also the drawn part of Γ_1^{100} is symmetric to the one of Γ_1^{010} as well as the drawn part of Γ_1^{110} is symmetric to the one of Γ_1^{101} . The morphisms $d^{\bullet 00}$ and $d^{0\bullet 0}$ (resp. $d^{10\bullet}$ and $d^{\bullet 10}$) clearly play symmetric roles. Besides, the morphisms $\delta : \llbracket \Gamma_1^{010} \rrbracket \{8\} \rightarrow \llbracket \Gamma_1^{101} \rrbracket \{10\}$ and $d^{1\bullet 0} : \llbracket \Gamma_1^{100} \rrbracket \{8\} \rightarrow \llbracket \Gamma_1^{110} \rrbracket \{10\}$ also play symmetric roles. Since the drawn parts of Γ_1 and Γ_2 in Figure 4.7 also are symmetric with respect to horizontal direction, we deduce that twice the complexes $\llbracket \Gamma_1 \rrbracket$ and $\llbracket \Gamma_2 \rrbracket$ have the same homology as the complex $\widehat{\mathcal{C}}$. \square

4.4 Invariance under type IV moves.

Let Γ_1 and Γ_2 be OMS-divides which differ only by a type IV move (see Figure 4.10).



Figure 4.10: Type IV move.

Proposition 4.12. *The complexes $\llbracket \Gamma_1 \rrbracket$ and $\llbracket \Gamma_2 \rrbracket$ have the same homology.*

Before proving this proposition, we first introduce the following intermediate result.

Lemma 4.13. *Let Γ and $\widetilde{\Gamma}$ be two cuspidal divides which differ only in the following way:*



Figure 4.11:

(or symmetric situation with respect to horizontal direction). Then $\llbracket \widetilde{\Gamma} \rrbracket \{6\} [1]$ is a strong deformation retract of $\llbracket \Gamma \rrbracket$.




Figure 4.12: Splitting diagram of $[[\Gamma]]$.

Proof. With the same arguments as in the previous sections, using lemma 3.14 we have a splitting diagram of $[[\Gamma]]$ (Figure 4.12):

Let's denote:

$$\begin{aligned} \mathcal{C}^0 &= [[\Gamma^{000}]]\{4\} & \check{\mathcal{C}}^1 &= [[\Gamma^{001}]]\{5\} & \check{\mathcal{C}}^2 &= [[\Gamma^{011}]]\{6\} & \mathcal{C}^3 &= [[\Gamma^{111}]]\{8\} \\ \hat{\mathcal{C}}^1 &= [[\Gamma^{100}]]\{6\} \oplus [[\Gamma_1^{010}]]\{5\} & \hat{\mathcal{C}}^2 &= [[\Gamma^{110}]]\{7\} \oplus [[\Gamma_1^{101}]]\{7\}. \end{aligned}$$

Then $[[\Gamma]] = \text{Cone}(\Delta^0, \Delta^1, \Delta^2)$ where: $\mathcal{C}^0 \xrightarrow{\Delta^0} \hat{\mathcal{C}}^1 \oplus \check{\mathcal{C}}^1 \xrightarrow{\Delta^1} \hat{\mathcal{C}}^2 \oplus \check{\mathcal{C}}^2 \xrightarrow{\Delta^2} \mathcal{C}^3$

$$\Delta^0 = \begin{pmatrix} d^{\bullet 00} \\ d^{0\bullet 0} \\ d^{00\bullet} \end{pmatrix} = \begin{pmatrix} \hat{\Delta}^0 \\ d^{00\bullet} \end{pmatrix} \quad \Delta^1 = \begin{pmatrix} d^{1\bullet 0} & d^{\bullet 10} & 0 \\ d^{10\bullet} & 0 & d^{\bullet 01} \\ 0 & d^{01\bullet} & d^{0\bullet 1} \end{pmatrix} = \begin{pmatrix} \hat{\Delta}^1 & U \\ L & d^{\bullet 01} \end{pmatrix}$$

and

$$\Delta^2 = (d^{11\bullet} \quad d^{1\bullet 1} \quad d^{\bullet 11}) = \begin{pmatrix} \hat{\Delta}^2 & d^{\bullet 11} \end{pmatrix}.$$

Consider the creation / destruction morphisms of complexes (see Figure 4.13):

$$\begin{array}{llll} [[\Gamma^{100}]]\{6\} \xrightarrow{\tau} [[\Gamma^{000}]]\{4\} & [[\Gamma^{110}]]\{7\} \xrightarrow{\eta_2} [[\Gamma^{010}]]\{5\} & [[\Gamma^{101}]]\{8\} \xrightarrow{\eta_2} [[\Gamma^{011}]]\{5\} \\ [[\Gamma^{010}]]\{5\} \xrightarrow{\sigma} [[\Gamma^{000}]]\{4\} & [[\Gamma^{101}]]\{7\} \xrightarrow{\eta_1} [[\Gamma^{100}]]\{5\} & [[\tilde{\Gamma}]]\{6\} \xrightarrow{\eta_1} [[\Gamma^{001}]]\{5\} \\ & [[\Gamma^{101}]]\{7\} \xrightarrow{\eta_2} [[\Gamma^{010}]]\{5\} & [[\Gamma^{001}]]\{5\} \xrightarrow{\bar{\varepsilon}_1} [[\tilde{\Gamma}]]\{6\} \end{array}$$




Figure 4.13:

Let's define $\sigma' = (\text{id} + \tau d^{\bullet 00})\sigma : [[\Gamma^{010}]]\{5\} \rightarrow [[\Gamma^{000}]]\{4\}$ and:

$$\hat{H}^1 = \begin{pmatrix} 0 & \eta_1 \\ \eta_2 & \eta_2 \end{pmatrix} \quad \hat{H}^0 = (\tau \quad \sigma').$$

Then we have a short exact sequence $0 \rightarrow \hat{\mathcal{C}}^2 \xrightarrow{\hat{H}^1} \hat{\mathcal{C}}^1 \xrightarrow{\hat{H}^0} \mathcal{C}^0 \rightarrow 0$ such that $\hat{\Delta}^1$ and $\hat{\Delta}^0$ are sections of \hat{H}^1 and \hat{H}^0 :

$$(4.7) \quad \hat{\Delta}^1 \hat{H}^1 = \text{id}, \quad \hat{H}^0 \hat{\Delta}^0 = \text{id}, \quad \hat{\Delta}^0 \hat{H}^0 + \hat{H}^1 \hat{\Delta}^1 = \text{id} + \hat{H}^1 \hat{\Delta}^1 \hat{\Delta}^0 \hat{H}^0.$$

Moreover, for the compositions $\llbracket \tilde{\Gamma} \rrbracket \{6\} \xrightarrow{\eta_1} \check{\mathcal{C}}^1 \xrightarrow{d^{0\bullet 1}} \check{\mathcal{C}}^2$ and $\check{\mathcal{C}}^1 \xrightarrow{U} \hat{\mathcal{C}}^2 \xrightarrow{\hat{H}^1} \hat{\mathcal{C}}^1 \xrightarrow{L} \check{\mathcal{C}}^2$ we have:

$$(4.8) \quad d^{0\bullet 1} \eta_1 = 0 \quad L \hat{H}^1 U = d^{0\bullet 1} \eta_2 d^{\bullet 0 1} = 0.$$

Let's define homotopies $H^0 : \hat{\mathcal{C}}^1 \oplus \check{\mathcal{C}}^1 \rightarrow \mathcal{C}^0$, $H^1 : \hat{\mathcal{C}}^2 \oplus \check{\mathcal{C}}^2 \rightarrow \hat{\mathcal{C}}^1 \oplus \check{\mathcal{C}}^1$ and $H^2 : \mathcal{C}^3 \rightarrow \hat{\mathcal{C}}^2 \oplus \check{\mathcal{C}}^2$ by:

$$H^0 = \begin{pmatrix} \hat{H}^0 & 0 \\ 0 & 0 \end{pmatrix} \quad H^1 = \begin{pmatrix} \hat{H}^1 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \quad H^2 = \begin{pmatrix} 0 \\ 0 \\ \eta_2 \end{pmatrix}$$

and consider the injection J and retraction R :

$$\begin{array}{c} \llbracket \tilde{\Gamma} \rrbracket \{6\} \\ \begin{array}{c} \uparrow R \\ \downarrow J \end{array} \\ \mathcal{C}^0 \xrightleftharpoons[H^0]{\Delta^0} \hat{\mathcal{C}}^1 \oplus \check{\mathcal{C}}^1 \xrightleftharpoons[H^1]{\Delta^1} \hat{\mathcal{C}}^2 \oplus \check{\mathcal{C}}^2 \xrightleftharpoons[H^2]{\Delta^2} \mathcal{C}^3 \end{array}$$

$$R = \begin{pmatrix} 0 & 0 & \bar{\varepsilon}_1 \end{pmatrix} (\text{id} + \Delta^0 H^0) \\ = \begin{pmatrix} \bar{\varepsilon}_1 d^{00\bullet} \tau & \bar{\varepsilon}_1 d^{00\bullet} \sigma' & \bar{\varepsilon}_1 \end{pmatrix} \quad J = (\text{id} + H^1 \Delta^1) \begin{pmatrix} 0 \\ 0 \\ \eta_1 \end{pmatrix} = \begin{pmatrix} \eta_1 d^{\bullet 0 1} \eta_1 \\ \eta_2 d^{\bullet 0 1} \eta_1 \\ \eta_1 \end{pmatrix}$$

(see Figure 4.14 for retraction R). From (4.7) and (4.8), we easily verify the relations:




Figure 4.14: The retraction R .

$$\begin{aligned} R \Delta^0 &= 0, & \Delta^1 J &= 0, & H^0 J &= 0, & R H^1 &= 0, & H^0 H^1 &= 0, & H^1 H^2 &= 0 \\ R J &= \text{id}, & J R &= \text{id} + \Delta^0 H^0 + H^1 \Delta^1, & \text{id} + \Delta^1 H^1 + H^2 \Delta^2 &= 0, & \text{id} + \Delta^2 H^2 &= 0. \end{aligned}$$

Then $\llbracket \tilde{\Gamma} \rrbracket \{6\}[1]$ is a strong deformation retract of $\llbracket \Gamma \rrbracket = \text{Cone}(\Delta^0, \Delta^1, \Delta^2)$ from proposition 3.12. \square

Proof of proposition 4.12. Let Γ_1^s (resp. Γ_2^s), $s \in \{0, 1\}$ be the cuspidal divides obtained by performing Θ_s splittings at the left hand + double point of Γ_1 (resp. Γ_2) in Figure 4.10. Let's also denote, according to lemma 4.13, the cuspidal divides $\tilde{\Gamma}_1^1$ (resp. $\tilde{\Gamma}_2^1$) obtained by "retracting" Γ_1^1 (resp. Γ_2^1) (see Figure 4.15). Notice that $\llbracket \Gamma_1^0 \rrbracket = \llbracket \Gamma_2^0 \rrbracket$ and $\llbracket \tilde{\Gamma}_1^1 \rrbracket = \llbracket \tilde{\Gamma}_2^1 \rrbracket$.




Figure 4.15:

From lemma 3.14, the differential d_1 on $[[\Gamma_1]]$ (resp. d_2 on $[[\Gamma_2]]$) gives us the following cone:

$$[[\Gamma_1]] = \text{Cone} \left([[\Gamma_1^0]]\{2\} \xrightarrow{d_1^\bullet} [[\Gamma_1^1]]\{4\} \right) \quad \left(\text{resp.} \quad [[\Gamma_2]] = \text{Cone} \left([[\Gamma_2^0]]\{2\} \xrightarrow{d_2^\bullet} [[\Gamma_2^1]]\{4\} \right) \right).$$

From lemma 4.13, there exist strong deformation retractions :

$$[[\Gamma_1^1]] \xrightleftharpoons[J_1]{R_1} [[\tilde{\Gamma}_1^1]]\{6\}[1] \quad [[\Gamma_2^1]] \xrightleftharpoons[J_2]{R_2} [[\tilde{\Gamma}_2^1]]\{6\}[1].$$

So from proposition 3.10, $\text{Cone}(R_1 d_1^\bullet)$ (resp. $\text{Cone}(R_2 d_2^\bullet)$) is a strong deformation retract of $[[\Gamma_1]]$ (resp. $[[\Gamma_2]]$). Hence it suffices to show that $R_1 d_1^\bullet = R_2 d_2^\bullet$. Let's consider the splitting diagram of $[[\Gamma_1^0]] = [[\Gamma_2^0]]$ as $\text{Cone}(D^0, D^1, D^2)$ (see Figure 4.16).




Figure 4.16:

Now $R_1 d_1^\bullet$ and $R_2 d_2^\bullet$ corresponds to the diagram of Figure 4.17.




Figure 4.17:

By combining Figure 4.14 with Figure 3.1 we easily verify that $R_1 d_1^\bullet = R_2 d_2^\bullet$. □

4.5 Invariance under type V moves.

Let Γ_0 and Γ_+ (resp. Γ_-) be OMS-divides which differ only by a type V_+ (resp. type V_-) move (see Figure 4.18).



Figure 4.18: Type V moves.

$$[[\Gamma_+^0]]\{1\} = \left[\begin{array}{c} \text{---} \bullet \\ \text{---} \bullet \\ \text{---} \bullet \end{array} \right] \{1\} \xrightarrow{d_+^\bullet} [\Gamma_+^1]\{2\} = \left[\text{---} \bullet \right] \{2\}$$

Figure 4.19: Splitting diagram for type V_+ move.

Proposition 4.14. *The complexes $[[\Gamma_0]]$, $[[\Gamma_+]]$ and $[[\Gamma_-]]$ have the same homology.*

Proof. From lemma 3.14, we can see $[[\Gamma_+]]$ as the cone of the surjective morphism:

Let's consider the creation / destruction morphisms $\eta_1 : [[\Gamma_+^1]]\{2\} \rightarrow [[\Gamma_+^0]]\{1\}$ (right inverse of d_+^\bullet), $\bar{\eta}_1 : [[\Gamma]] \rightarrow [[\Gamma_+^0]]\{1\}$ and $\varepsilon_1 : [[\Gamma_+^0]]\{1\} \rightarrow [[\Gamma]]$ defined in Figure 4.20. Let $j =$




Figure 4.20:

$(\text{id} + \eta_1 d_+^\bullet) \bar{\eta}_1$ and $r = \varepsilon_1$. Then from the following diagram:

$$\begin{array}{ccc} [[\Gamma_0]] & & d_+^\bullet j = 0, \quad rj = \text{id}, \\ j \downarrow \uparrow r & & \\ [[\Gamma_+^0]]\{1\} & \xrightarrow{d_+^\bullet} & [[\Gamma_+^1]]\{2\} \end{array} \quad \begin{array}{l} jr = \text{id} + \eta_1 d_+^\bullet, \quad d_+^\bullet \eta_1 = \text{id} \\ \eta_1 \end{array}$$

we deduce using proposition 3.12 that $[[\Gamma_0]]$ is a strong deformation retract of $[[\Gamma_+]] = \text{Cone}(d_+^\bullet)$. So they have the same homology.

On the other hand, from lemma 3.14, $[[\Gamma_-]] = \text{Cone}(\Delta^0, \Delta^1)[-1]$:

$$[[\Gamma_-^{10}]]\{-3\} \xrightarrow{\Delta^0} [[\Gamma_-^{00}]]\{-1\} \oplus [[\Gamma_-^{11}]]\{-2\} \xrightarrow{\Delta^1} [[\Gamma_-^{01}]]\{3\} \quad \Delta^0 = \begin{pmatrix} d_-^{0\bullet} \\ d_-^{1\bullet} \end{pmatrix} \quad \Delta^1 = (d_-^{0\bullet} \ d_-^{1\bullet})$$





Figure 4.21: Splitting diagram for type V_- move.

Let's consider the morphisms $[[\Gamma_-^{00}]]\{-1\} \xrightarrow{\tau} [[\Gamma_-^{10}]]\{-3\}$, $[[\Gamma_-^{11}]]\{-2\} \xrightarrow{\sigma} [[\Gamma_-^{10}]]\{-3\}$, $[[\Gamma_-^{01}]] \xrightarrow{\eta_1} [[\Gamma_-^{00}]]\{-1\}$ and $[[\Gamma_-^{01}]] \xrightarrow{\eta_2} [[\Gamma_-^{11}]]\{-2\}$ defined in Figure 4.22.



Then from the diagram:

$$\begin{array}{c}
 \llbracket \Gamma_0 \rrbracket \\
 J \updownarrow R \\
 \llbracket \Gamma_-^{10} \rrbracket \{-3\} \xrightleftharpoons[H^0]{\Delta^0} \llbracket \Gamma_-^{00} \rrbracket \{-1\} \oplus \llbracket \Gamma_-^{11} \rrbracket \{-2\} \xrightleftharpoons[H^1]{\Delta^1} \llbracket \Gamma_-^{01} \rrbracket
 \end{array}$$

where $J = \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}$, $R = (0 \ \bar{\varepsilon}_2)$, $H^0 = (\tau \ (\text{id} + \tau d^{\bullet 0}) \sigma)$ and $H^1 = \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}$ we deduce that $\llbracket \Gamma_0 \rrbracket$ is a strong deformation retract of $\llbracket \Gamma_- \rrbracket = \text{Cone}(\Delta^0, \Delta^1)[-1]$. They have the same homology. \square

4.6 Invariance under type VI moves.

Let Γ and Γ_+ (resp. Γ_-) be OMS-divides which differ only by a type VI_+ (resp. type VI_-) move (see Figure 4.23).



Figure 4.23: Type VI moves.

Proposition 4.15. *The complexes $\llbracket \Gamma_0 \rrbracket$, $\llbracket \Gamma_+ \rrbracket$ and $\llbracket \Gamma_- \rrbracket$ have the same homology.*

We will break down the proof in two steps: the result is an immediate consequence of the following two lemmas.

Lemma 4.16. *Let Γ and $\tilde{\Gamma}$ be cuspidal divides defined by:*

$$\Gamma = \text{---} \text{---} \text{---} \text{---} \text{---} \quad \tilde{\Gamma} = \text{---} \text{---} \text{---} \text{---} \text{---}$$

Figure 4.24:

Then $\llbracket \tilde{\Gamma} \rrbracket$ is a strong deformation retract of $\llbracket \Gamma \rrbracket$.

Proof. Let's apply lemma 3.14 to $\llbracket \Gamma \rrbracket$. We have a splitting diagram :




Figure 4.25: Splitting diagram of $[[\Gamma]]$.

Let's denote :

$$\begin{aligned} \mathcal{C}^0 &= [[\Gamma^{100}]]\{-2\} & \mathcal{C}^1 &= [[\Gamma^{110}]]\{-1\} \oplus \left([[\Gamma^{101}]]\{-1\} \oplus [[\Gamma^{000}]] \right) \\ \mathcal{C}^2 &= [[\Gamma^{111}]] \oplus \left([[\Gamma^{010}]]\{1\} \oplus [[\Gamma^{001}]]\{1\} \right) & \mathcal{C}^3 &= [[\Gamma^{011}]]\{2\} \\ \Delta^0 &= \begin{pmatrix} d^{1\bullet 0} \\ d^{10\bullet} \\ d^{\bullet 00} \end{pmatrix} & \Delta^1 &= \begin{pmatrix} d^{11\bullet} & d^{1\bullet 1} & 0 \\ d^{\bullet 10} & 0 & d^{0\bullet 0} \\ 0 & d^{\bullet 01} & d^{00\bullet} \end{pmatrix} & \Delta^2 &= (d^{\bullet 11} \quad d^{01\bullet} \quad d^{0\bullet 1}). \end{aligned}$$

Then $[[\Gamma]] = \text{Cone}(\Delta^0, \Delta^1, \Delta^2)[-1]$ where $\mathcal{C}^0 \xrightarrow{\Delta^0} \mathcal{C}^1 \xrightarrow{\Delta^1} \mathcal{C}^2 \xrightarrow{\Delta^2} \mathcal{C}^3$. Let's consider the creation / destruction morphisms:

$$\begin{aligned} [[\Gamma^{101}]]\{-1\} &\xrightarrow{\sigma} [[\Gamma^{100}]]\{-2\} & [[\Gamma^{111}]] &\xrightarrow{\sigma} [[\Gamma^{110}]]\{-1\} & [[\Gamma^{011}]]\{2\} &\xrightarrow{\eta_2} [[\Gamma^{111}]] \\ [[\Gamma^{000}]] &\xrightarrow{\tau} [[\Gamma^{100}]]\{-2\} & [[\Gamma^{010}]]\{1\} &\xrightarrow{\eta_1} [[\Gamma^{000}]] & [[\tilde{\Gamma}]] &\xrightarrow{\eta_1} [[\Gamma^{110}]]\{-1\} \\ & & [[\Gamma^{001}]]\{1\} &\xrightarrow{\eta_2} [[\Gamma^{101}]]\{-1\} & [[\Gamma^{110}]]\{-1\} &\xrightarrow{\bar{\varepsilon}_1} [[\tilde{\Gamma}]] \end{aligned}$$

defined by:

$$\begin{aligned} \begin{array}{c} \begin{array}{c} a \\ b \end{array} \begin{array}{c} c \\ \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\sigma} \begin{cases} 0 & \text{if } c = + \\ \begin{array}{c} a \\ b \end{array} \begin{array}{c} \bar{c} \\ \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} & \text{if } c = - \end{cases} & \begin{array}{c} a \\ c \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\sigma} \begin{cases} 0 & \text{if } c = + \\ \begin{array}{c} a \\ \bar{c} \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} & \text{if } c = - \end{cases} & \begin{array}{c} a \\ \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\eta_2} \begin{array}{c} a \\ \dagger \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \\ \begin{array}{c} \begin{array}{c} c \\ a \end{array} \\ b \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\tau} \begin{cases} 0 & \text{if } c = + \\ \begin{array}{c} a \\ b \end{array} \begin{array}{c} \dagger \\ \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} & \text{if } c = - \end{cases} & \begin{array}{c} a \\ b \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\eta_1} \begin{array}{c} \dagger \\ a \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} & \begin{array}{c} a \\ \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\eta_1} \begin{array}{c} a \\ \dagger \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \\ \begin{array}{c} a \\ b \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\eta_2} \begin{array}{c} a \\ b \end{array} \begin{array}{c} \dagger \\ \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} & \begin{array}{c} a \\ c \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} \xrightarrow{\bar{\varepsilon}_1} \begin{cases} 0 & \text{if } c = + \\ \begin{array}{c} a \\ \end{array} \begin{array}{c} \diagup \\ \diagdown \end{array} & \text{if } c = - \end{cases} \end{aligned}$$

$a, b, c \in \{-, +\}$

Figure 4.26:

We define homotopies $H^0 : \mathcal{C}^1 \rightarrow \mathcal{C}^0$, $H^1 : \mathcal{C}^2 \rightarrow \mathcal{C}^1$ and $H^2 : \mathcal{C}^3 \rightarrow \mathcal{C}^2$:

$$H^0 = \begin{pmatrix} 0 & (1 + \tau d^{\bullet 00})\sigma & \tau \end{pmatrix} \quad H^1 = \begin{pmatrix} \sigma & 0 & 0 \\ \eta_2 d^{00\bullet} \eta_1 d^{\bullet 10} \sigma & \eta_2 d^{00\bullet} \eta_1 & \eta_2 \\ \eta_1 d^{\bullet 10} \sigma & \eta_1 & 0 \end{pmatrix} \quad H^2 = \begin{pmatrix} \eta_2 \\ 0 \\ 0 \end{pmatrix}$$

together with retraction and inclusion maps $R : \mathcal{C}^1 \rightarrow [[\tilde{\Gamma}]]$ and $J : [[\tilde{\Gamma}]] \rightarrow \mathcal{C}^1$:

$$\begin{aligned} R &= (\bar{\varepsilon}_1 \quad 0 \quad 0) (\text{id} + \Delta^0 H^0) \\ &= (\bar{\varepsilon}_1 \quad \bar{\varepsilon}_1 d^{1\bullet 0} \tau d^{\bullet 00} \sigma \quad \bar{\varepsilon}_1 d^{1\bullet 0} \tau) \quad J = (\text{id} + H^1 \Delta^1) \begin{pmatrix} \eta_1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} \eta_1 \\ \eta_2 d^{00\bullet} \eta_1 d^{\bullet 10} \eta_1 \\ \eta_1 d^{\bullet 10} \eta_1 \end{pmatrix} \end{aligned}$$

such that we have the diagram:

$$\begin{array}{ccccccc}
& & & & \llbracket \tilde{\Gamma} \rrbracket & & \\
& & & & \uparrow J & \downarrow R & \\
\mathcal{C}^0 & \xrightleftharpoons[H^0]{\Delta^0} & \mathcal{C}^1 & \xrightleftharpoons[H^1]{\Delta^1} & \mathcal{C}^2 & \xrightleftharpoons[H^2]{\Delta^2} & \mathcal{C}^3
\end{array}$$

We easily verify that :

$$\begin{aligned}
R\Delta^0 &= 0, & \Delta^1 J &= 0, & H^0 J &= 0, & RH^1 &= 0, & H^0 H^1 &= 0, & H^1 H^2 &= 0 \\
RJ &= \text{id}, & JR &= \text{id} + \Delta^0 H^0 + H^1 \Delta^1, & \text{id} + \Delta^1 H^1 + H^2 \Delta^2 &= 0, & \text{id} + \Delta^2 H^2 &= 0.
\end{aligned}$$

Then from proposition 3.12, $\llbracket \tilde{\Gamma} \rrbracket$ is a strong deformation retract of $\llbracket \Gamma \rrbracket$. \square

Lemma 4.17. *We have strong deformation retractions r_+ (resp. r_-) with injection j_+ (resp. j_-):*

$$\llbracket \Gamma_0 \rrbracket = \llbracket \text{---} \rrbracket \xrightleftharpoons[r_+]{j_+} \llbracket \tilde{\Gamma}_+ \rrbracket = \llbracket \text{---} \text{---} \text{---} \rrbracket \quad \left(\text{resp. } \llbracket \Gamma_0 \rrbracket = \llbracket \text{---} \rrbracket \xrightleftharpoons[r_-]{j_-} \llbracket \tilde{\Gamma}_- \rrbracket = \llbracket \text{---} \text{---} \text{---} \rrbracket \right)$$

Figure 4.27:

Proof. From lemma 3.14, $\llbracket \tilde{\Gamma}_+ \rrbracket = \text{Cone}(d_+)$ and $\llbracket \tilde{\Gamma}_- \rrbracket = \text{Cone}(d_-)[-1]$ where:

$$\llbracket \text{---} \text{---} \text{---} \rrbracket \{2\} \xrightarrow{d_+} \llbracket \text{---} \text{---} \text{---} \rrbracket \{4\} \quad \llbracket \text{---} \text{---} \text{---} \rrbracket \{-4\} \xrightarrow{d_-} \llbracket \text{---} \text{---} \text{---} \rrbracket \{-2\}$$

Figure 4.28:

For $\llbracket \Gamma_+ \rrbracket$, using proposition 3.12, we deduce the strong deformation retraction from the following diagram of Figure 4.29 since $d_+ \eta_2 = \text{id}$, $r_+ \eta_2 = 0$, $d_+ j_+ = 0$, $r_+ j_+ = \text{id}$ and $j_+ r_+ = \text{id} + \eta_2 d_+$.

$$\begin{array}{ccc}
\llbracket \text{---} \text{---} \text{---} \rrbracket \{2\} \xrightarrow{d_+} \llbracket \text{---} \text{---} \text{---} \rrbracket \{4\} & \llbracket \text{---} \text{---} \text{---} \rrbracket \xrightarrow{\eta_2} \llbracket \text{---} \text{---} \text{---} \rrbracket^+ & \\
\uparrow \eta_2 & \llbracket \text{---} \text{---} \text{---} \rrbracket \xrightarrow{\bar{\eta}_2} \llbracket \text{---} \text{---} \text{---} \rrbracket^- & \\
\llbracket \text{---} \text{---} \text{---} \rrbracket & \llbracket \text{---} \text{---} \text{---} \rrbracket \xrightarrow{\varepsilon_2} \begin{cases} 0 & \text{if } c = + \\ \llbracket \text{---} \text{---} \text{---} \rrbracket & \text{if } c = - \end{cases} & \\
r_+ \uparrow j_+ & & \\
r_+ = \varepsilon_2 & j_+ = (\text{id} + \eta_2 d_+) \bar{\eta}_2 &
\end{array}$$

Figure 4.29:

Similarly for $\llbracket \Gamma_- \rrbracket$ from the diagram of Figure 4.30:

since $\varepsilon_2 d_- = \text{id}$, $r_- d_- = 0$, $\varepsilon_2 j_- = 0$, $r_- j_- = \text{id}$ and $j_- r_- = \text{id} + d_- \varepsilon_2$. \square




Figure 4.30:

References

- [1] N. A'Campo - "Real deformations and complex topology of plane singularity" - Ann. Fac. Sc. Toulouse 8 (1999).
- [2] O. Couture - "Strongly invertible links and divides" Topology 47 (2008) 316-350.
- [3] O. Couture, B. Perron - "Representative braids for links associated to plane immersed curves" - J.Knot th. and Ram. Vol. 9, No. 1 (2000) 1-30.
- [4] J.W. Morgan, H. Bass - "The Smith conjecture" - Academic Press (1984).
- [5] O. Viro - "Remarks on the definition of Khovanov homology" - arXiv:math. GT/0202199 (2002).