HAL
open science

Non-relativistic conformal symmetries and Newton-Cartan structures

Christian Duval, Péter A. Horvathy

To cite this version:

Christian Duval, Péter A. Horvathy. Non-relativistic conformal symmetries and Newton-Cartan structures. Journal of Physics A: Mathematical and Theoretical, 2009, 42, pp.465206. hal-00373011v5

HAL Id: hal-00373011

https://hal.science/hal-00373011v5

Submitted on 26 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non-relativistic conformal symmetries and Newton-Cartan structures

C. DUVAL[‡]
Centre de Physique Théorique, CNRS, Luminy, Case 907
F-13288 Marseille Cedex 9 (France)[§]

P. A. HORVÁTHY[¶]
Laboratoire de Mathématiques et de Physique Théorique
Université de Tours, Parc de Grandmont
F-37200 TOURS (France)

26 September 2009

Abstract

This article provides us with a unifying classification of the conformal infinitesimal symmetries of non-relativistic Newton-Cartan spacetime. The Lie algebras of non-relativistic conformal transformations are introduced via the Galilei structure. They form a family of infinite-dimensional Lie algebras labeled by a rational "dynamical exponent", z. The Schrödinger-Virasoro algebra of Henkel et al. corresponds to $z = 2$. Viewed as projective Newton-Cartan symmetries, they yield, for timelike geodesics, the usual Schrödinger Lie algebra, for which $z = 2$. For lightlike geodesics, they yield, in turn, the Conformal Galilean Algebra (CGA) and Lukierski, Stichel and Zakrzewski [alias "$\mathfrak{alt}$" of Henkel], with $z = 1$. Physical systems realizing these symmetries include, e.g., classical systems of massive, and massless non-relativistic particles, and also hydrodynamics, as well as Galilean electromagnetism.

`arXiv:0904.0531v5 [hep-th]`. MSC2000: 37J15, 37K05, 70G65, 53B15.

Keywords: Schrödinger algebra, Conformal Galilei algebra, Newton-Cartan Theory.

[‡]mailto: duval-at-cpt.univ-mrs.fr

[§]UMR 6207 du CNRS associée aux Universités d'Aix-Marseille I and II and Université du Sud Toulon-Var; Laboratoire affilié à la FRUMAM-FR2291.

[¶]mailto: horvathy-at-lmpt.univ-tours.fr

Contents

1 Introduction **3**

2 Newton-Cartan structures **6**

2.1 Galilei structures and Newton-Cartan connections 6

2.2 NC-gauge transformations, and NC-Milne structures 7

2.2.1 Gauge transformations 7

2.2.2 NC-Milne structure 8

3 Conformal Galilei transformations, Schrödinger-Virasoro Lie algebra **8**

3.1 The Lie algebra, $\mathfrak{cgal}$, of conformal Galilei transformations 8

3.2 Conformal Galilei transformations, $\mathfrak{cgal}_z$, with dynamical exponent z 9

4 Conformal Newton-Cartan transformations **11**

4.1 The Schrödinger Lie algebra 12

4.1.1 The expanded Schrödinger Lie algebra, $\widetilde{\mathfrak{sch}}$, of projective Galilei conformal transformations 12

4.1.2 The Schrödinger Lie algebras, $\mathfrak{sch}_z$, with dynamical exponent z 14

4.2 Transformation law of NC-connections under conformal Galilei rescalings 15

4.3 Conformal NC transformations: lightlike geodesics 16

4.3.1 The conformal Newton-Cartan Lie algebra, $\mathfrak{cnc}$, of null-projective conformal Galilei transformations 17

4.3.2 The Lie algebra, $\mathfrak{cmil}$, of null-projective conformal transformations of NC-Milne spacetime 19

4.3.3 Conformal NC-Milne Lie algebras, $\mathfrak{cmil}_z$, with dynamical exponent z; the CGA Lie algebra 21

4.3.4 The finite-dimensional conformal Galilei Lie algebras, $\mathfrak{alt}_{2/N}(d)$ 22

5 Conformal Galilean symmetries of physical systems **24**

5.1 Galilean massive particles 26

5.2 Galilean symmetry in hydrodynamics 27

5.3 Galilean massless particles 33

5.4 Galilean Electromagnetism 36

6 Conclusion **38**

1 Introduction

Non-relativistic conformal symmetries, which are attracting much present interest [1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12], are of two types.

Firstly, it has been recognized almost forty years ago [14, 15] that the free Schrödinger equation of a massive particle has, beyond the obvious Galilean symmetry, two more "conformal" symmetries. They are generated by the "Schrödinger" spacetime vector fields, called *dilation*

$$D = 2t\frac{\partial}{\partial t} + x^A\frac{\partial}{\partial x^A} \tag{1.1}$$

and *expansion* (or inversion)

$$K = t^2\frac{\partial}{\partial t} + tx^A\frac{\partial}{\partial x^A} \tag{1.2}$$

where the dummy index A runs from 1 to d, the dimension of space.

Schrödinger dilations and expansions span, with time translations, $H = \partial/\partial t$, a Lie algebra isomorphic to $\mathfrak{so}(2,1)$. Adding dilations and expansions to the Galilei group yields a two-parameter extension of the latter, dubbed as the (centerless) *Schrödinger group*, $\mathrm{Sch}(d)$.[1]

Using the word "conformal" has been contested [1], hinting at its insufficiently clear relation to some conformal structure. This criticism is only half-justified, however. The Schrödinger symmetry has in fact been related to the Newton-Cartan structure of non-relativistic spacetime [21, 22, 23, 24], but this relation has remained rather confidential.

A different point of view was put forward in Ref. [26], where it has been shown that non-relativistic theories can be studied in a "Kaluza-Klein type" framework, whereas the "non-relativistic conformal" transformations appear as those, genuine, conformal transformations of a relativistic spacetime in one higher dimension, which commute with translations in the "vertical" direction. The latter provides us, furthermore, with the central extension required by the mass [15, 27, 28].

Secondly, after the pioneering work of Henkel [29], in [1, 2, 3, 7, 9, 10, 11, 13], attention has been directed to another, less-known and more subtle aspect. It has been shown, in fact, that a specific group contraction, applied to the relativistic conformal group $\mathrm{O}(d+1,2)$, provides, for *vanishing mass*, $m = 0$, a *second type of*

[1] The physical realizations of the Schrödinger group, in spatial dimension $d \geq 3$, admit one more parameter, associated with the *mass*. Adding it yields the *extended Schrödinger group*, which is the "non-relativistic conformal" extension of the one-parameter central extension of the Galilei group, called the *Bargmann group*. (See, e.g., [33, 21, 16, 25] for a geometrical account on the Bargmann group.) In the plane, $d = 2$, the Galilei group also has, apart from the previous one, a second, "exotic", central extension widely studied during the last decade [17, 18, 19, 20].

conformal extension of the Galilei group. Since group contraction does not change the number of generators, the new extension, called the *Conformal Galilean Group* [1] has the same dimension as its relativistic counterpart. Its Lie algebra, the *Conformal Galilei Algebra* is spelled as the CGA in the above-mentioned reference. The CGA is spanned by the vector fields[2]

$$X = \Big(\frac{1}{2}\kappa t^2 + \lambda t + \varepsilon\Big)\frac{\partial}{\partial t} + \Big(\omega^A_B\, x^B + \lambda x^A + \kappa t x^A - \frac{1}{2}\alpha^A t^2 + \beta^A t + \gamma^A\Big)\frac{\partial}{\partial x^A} \quad (1.3)$$

with $\boldsymbol{\omega} \in \mathfrak{so}(d)$, $\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma} \in \mathbb{R}^d$, and $\lambda, \kappa, \varepsilon \in \mathbb{R}$.

The new dilations and expansions, associated with λ and κ close, with time translations parametrized by ε, into an $\mathfrak{so}(2,1)$ Lie subalgebra [1, 2, 3, 9], acting *differently* from that of the Schrödinger case: unlike the "Schrödinger" one, (1.1), the CGA dilation in (1.3) dilates space and time at the same rate. Note also the factor $\frac{1}{2}$ in the time component of the new expansions. The vector $\boldsymbol{\alpha}$ generates, in turn, "accelerations" [1]. See also [29] for another approach, and [2] where the CGA was called $\mathfrak{alt}(d)$.

The Lie algebra (1.3) can be further generalized [2, 3, 9], in terms of infinitesimal "time redefinition" and time-dependent translations,

$$X = \xi(t)\frac{\partial}{\partial t} + \big(\xi'(t)x^A + \eta^A(t)\big)\frac{\partial}{\partial x^A} \quad (1.4)$$

where $\xi(t)$, and $\boldsymbol{\eta}(t)$ are arbitrary functions of time, t. The new expansions and accelerations are plainly recovered choosing $\xi(t) = \frac{1}{2}\kappa\, t^2$ and $\boldsymbol{\eta}(t) = -\frac{1}{2}\boldsymbol{\alpha}\, t^2$, respectively. Promoting the infinitesimal rotations, $X = \omega^A_B(t)x^B\partial_A$, to be also time-dependent yields an infinite-dimensional conformal extension of the CGA.

The purpose of the present paper, a sequel and natural extension of earlier work devoted to Galilean isometries [25], is to trace-back all these "conformal" symmetries to the *structure of non-relativistic spacetime.*

Our clue is to define *non-relativistic conformal transformations* in the framework of Newton-Cartan spacetime [30, 31, 32, 34], ideally suited to deal with those symmetries in a purely geometric way. In contradistinction to the (pseudo-)Riemannian framework, the degeneracy of the Galilei "metric" allows, as we shall see, for infinite-dimensional Lie algebras of conformal Galilei infinitesimal transformations, with a wealth of finite-dimensional Lie subalgebras, including the Schrödinger Lie algebra and the above-mentioned CGA.

Both the Schrödinger and Conformal Galilean transformations turn out to be special cases, related to our choice of the relative strength of space and time dilations, characterized by a dynamical exponent [2, 3].

[2]The central extensions of the CGA have been discussed in Refs. [1, 3].

Our paper is organized as follows.

After reviewing, in Section 2, the *Newton-Cartan structures* of $(d+1)$-dimensional non-relativistic spacetime, we introduce, in Section 3, the notion of *conformal Galilei transformation*. The latter is only concerned with the (singular) "metric", γ, and the "clock", represented by a closed one-form θ. Infinitesimal conformal Galilei transformations form an infinite-dimensional Virasoro-like Lie algebra, denoted by $\mathfrak{cgal}(d)$ in the case of ordinary Galilei spacetime. A geometric definition of the dynamical exponent, z, allows us to define the conformal Galilei Lie algebras, $\mathfrak{cgal}_z(M,\gamma,\theta)$ of an arbitrary Galilei structure, with prescribed z.

Now, Newton-Cartan structures also involve a connection, Γ, which is *not* entirely determined by the previous structures. Preserving the geodesic equations adds, in the generic case, extra conditions, which are explicitly derived in Section 4.

Those help us to reduce the infinite-dimensional conformal Galilei Lie algebra to that of the Schrödinger Lie algebra, $\mathfrak{sch}(d)$, for *timelike* geodesics of the flat NC-structure (with dynamical exponent $z=2$). This is reviewed in Section 4.1.

For *lightlike* geodesics, we get, in turn, a novel, infinite-dimensional, conformal extension, $\mathfrak{cnc}(d)$, of the (centerless) Galilei Lie algebra, which is worked out in Section 4.3. This conformal Newton-Cartan Lie algebra admits, indeed, infinite-dimensional Lie subalgebras defined by an arbitrary (rational) dynamical exponent, z. Also, the maximal Lie algebra of conformal automorphisms of a *Milne structure*, i.e., a NC-structure with a preferred geodesic and irrotational observer field, shows up as a finite-dimensional Lie algebra, denoted by $\mathfrak{cmil}(d)$ in the case of flat spacetime. The CGA (1.3) finally appears as a Lie subalgebra of $\mathfrak{cmil}(d)$ defined by the dynamical exponent $z=1$. The Lie algebras $\mathfrak{alt}_{2/N}(d)$ first defined in [29] appear plainly as the Lie algebras of polynomial vector fields of degree $N=1,2,3,\ldots$ in $\mathfrak{cgal}_z(d)$ with $z=2/N$. A geometric definition for the latter Lie algebras is still missing, though.

The general theory is illustrated, in Section 5, on various examples. Schrödinger symmetry is shown to be present for a Galilean massive particle and in hydrodynamics. The massless non-relativistic particle of Souriau exhibits, as a symmetry, an infinite-dimensional conformal extension of the centerless Galilei Lie algebra. At last, the Le Bellac-Lévy-Leblond theory of (magnetic-like) Galilean electromagnetism carries, apart of the Schrödinger symmetry, also the CGA.

2 Newton-Cartan structures

2.1 Galilei structures and Newton-Cartan connections

Let us recall that a Newton-Cartan (NC) spacetime structure, $(M, \gamma, \theta, \Gamma)$, consists of a smooth, connected, $(d+1)$-dimensional manifold M, a twice-contravariant symmetric tensor field $\gamma = \gamma^{ab}\,\partial_a \otimes \partial_b$ (where $a, b = 0, 1, \ldots, d$) of signature $(0, +, \ldots, +)$ whose kernel is spanned by the one-form $\theta = \theta_a dx^a$. Also Γ is a Galilei connection, i.e., a symmetric linear connection compatible with γ and θ [30, 31, 32, 34, 35, 36].

Now, in contradistinction to the relativistic framework, such a connection is *not* uniquely determined by the Galilei spacetime structure (M, γ, θ). Therefore, in order to reduce the ambiguity, one usually introduces NC-connection as Galilei connections subject to the nontrivial symmetry of the curvature: $R_{a\ c}^{\ b\ d} = R_{c\ a}^{\ d\ b}$ (where $R_{a\ c}^{\ b\ d} \equiv \gamma^{bk}\, R_{akc}^{\ \ \ d}$); the latter may be thought of as part of the covariant Newtonian gravitational field equations [31, 32, 33, 34].

Under mild geometric conditions, the quotient $T = M/\ker(\theta)$ is a well-behaved one-dimensional manifold, interpreted as the time axis endowed with the closed one-form θ, interpreted as the Galilei *clock*. The tensor field γ then defines a Riemannian metric on each of the (spacelike) fibers of the projection $M \to T$.

The standard example of a NC-structure is given by $M \subset \mathbb{R} \times \mathbb{R}^d$ together with $\gamma = \delta^{AB}\partial_A \otimes \partial_B$ (where $A, B = 1, \ldots, d$), and $\theta = dx^0$; the nonzero components of the connection, $\Gamma^A_{00} = \partial_A V$, host the Newtonian scalar potential, V. The above coordinate system $(x^0, \ldots, x^d)$ will be called Galilean.

The *flat* NC-structure corresponds to the subcase where $M = \mathbb{R} \times \mathbb{R}^d$, and

$$\gamma^{ab} = \delta^a_A \delta^b_B\, \delta^{AB}, \qquad \theta_a = \delta^0_a, \qquad \Gamma^c_{ab} = 0 \tag{2.1}$$

for all $a, b, c = 0, \ldots, d$. Such a coordinate system will be called (NC-)inertial.

Since we will be dealing with "conformal" Galilean spacetime transformations that preserve the directions of the Galilei structure, we must bear in mind that the transformation law of the NC-connection, Γ, will have to be specified independently of that of the Galilei "metric" (γ, θ), which is clearly due to the fact that there are extra degrees of freedom associated with NC-connections. Let us, hence, describe the precise geometric content of NC-connections.

It has been shown [32] that NC-connections can be decomposed according to[3]

$$\Gamma^c_{ab} = {}^U\Gamma^c_{ab} + \theta_{(a} F_{b)k} \gamma^{kc} \tag{2.2}$$

where [43]

$${}^U\Gamma^c_{ab} = \gamma^{ck}\Big(\partial_{(a}\, {}^U\gamma_{b)k} - \frac{1}{2}\partial_k\, {}^U\gamma_{ab}\Big) + \partial_{(a}\theta_{b)}\, U^c \tag{2.3}$$

[3]Round brackets denote symmetrization, and square ones will denote skew-symmetrization.

is the unique NC-connection for which the unit spacetime vector field U (i.e., such that $\theta_a U^a = 1$) is geodesic and curlfree, F being an otherwise arbitrary closed two-form. Here ${}^U\gamma$ is the symmetric, twice-covariant, tensor field uniquely determined by ${}^U\gamma_{ak}\gamma^{kb} = \delta_a^b - U^b\theta_a$ and ${}^U\gamma_{ak}U^k = 0$. From a mechanical standpoint, the above two-form, F, of M encodes Coriolis-like accelerations relatively to the observer U.

For example, if $M \subset \mathbb{R} \times \mathbb{R}^3$, the constant, future-pointing, vector field $U = \partial_0$ will represent the four-velocity of an observer. Now, F being closed, one has, locally, $F = dA$ for some one-form A, e.g., $A = -V(t, \mathbf{x})dt + \omega(t)_{BC}\, x^B dx^C$, where $V(t, \mathbf{x})$ is the Newtonian (plus centrifugal) potential, and $\boldsymbol{\omega}(t) \in \mathfrak{so}(3)$ the time-dependent angular velocity of the observer relatively to the Galilei frame associated with the coordinates $t = x^0$, and $\mathbf{x} = (x^1, x^2, x^3)$. Anticipating the equations of free fall, we check that the equations of NC-geodesics (4.1) — with the choice of time, t, as an affine parameter — yield, with the help of (2.2), the familiar equations

$$\ddot{t} = 0, \qquad \ddot{\mathbf{x}} = -\boldsymbol{\nabla} V + \dot{\boldsymbol{\omega}} \times \mathbf{x} + 2\boldsymbol{\omega} \times \dot{\mathbf{x}} \tag{2.4}$$

governing the motion of a massive particle in a rotating Galilei coordinate system.[4]

2.2 NC-gauge transformations, and NC-Milne structures

2.2.1 Gauge transformations

We have seen that, in view of (2.2), we can usefully parametrize NC-connections, Γ, by the previously introduced pairs (U, F) which are, themselves, not entirely fixed by the NC-connection. (This arbitrariness in the expression of the NC-connection can be traced-back to the degeneracy of the Galilei structure ; this does not occur in the pseudo-Riemannian case where the Levi-Civita connection is uniquely determined by the metric.)

Let us mention [32, 34] that for a given, *fixed*, Galilei structure (γ, θ), the pair (U', F') defines the same NC-connection, Γ, as (U, F) does iff both are *gauge*-related by a so-called *Milne boosts* [24, 25]

$$U' = U + \gamma(\Psi), \qquad F' = F + d\Phi \tag{2.5}$$

where $\Psi = \Psi_a dx^a$ is an arbitrary one-form of M, which may be interpreted as a boost,[5] and $\Phi = \Phi_a dx^a$ is such that

$$\Phi_a = \Psi_a - \Big(\Psi_b U^b + \frac{1}{2}\gamma^{bc}\Psi_b\Psi_c\Big)\theta_a. \tag{2.6}$$

[4]The non-trivial components of the NC-connection (2.2) read, in this case, $\Gamma^A_{00} = \partial_A V - \dot{\omega}^A_B x^B$, and $\Gamma^A_{B0} = -\omega^A_B$, for all $A, B = 1, 2, 3$.

[5]Two observers U, and U' are related by a boost, i.e., an acceleration $A = U' - U$ which is necessarily spacelike, $\theta(A) = 0$, hence of the form $A = \gamma(\Psi)$, as specified in (2.5).

The infinitesimal versions of the preceding gauge transformations read, accordingly,

$$\delta U = \gamma(\psi), \qquad \delta F = d\phi \tag{2.7}$$

where ψ is an arbitrary one-form of M (an infinitesimal boost), and

$$\phi = {}^{U}\gamma(\delta U). \tag{2.8}$$

One readily checks that, indeed, $\delta\Gamma = 0$.

2.2.2 NC-Milne structure

In fact, given a NC-connection, Γ, and an arbitrary observer, U, one uniquely determines the (closed) "Coriolis" two-form, F, via the fundamental relation [32, 34]

$$F_{ab} = -2\,{}^{U}\gamma_{c[a}\nabla_{b]}U^c \tag{2.9}$$

where ∇ stands for the covariant derivative associated with the NC-connection, Γ. This implies that the *geodesic acceleration*, $\dot{U}^a = U^b\nabla_b U^a$, of the observer U reads

$$\dot{U}^a = -F^a_b U^b \tag{2.10}$$

while its *curl* is of the form

$$2\nabla^{[a}U^{b]} = F^{ab} \tag{2.11}$$

where coordinate indices have lifted using γ, e.g., $F^a_b = \gamma^{ac}F_{cb}$. An inertial and non-rotating observer, U, will therefore be characterized by $F = 0$. Whenever such an observer exists, it will be called an *ether*, in the spirit of [24].

We call *NC-Milne structure* a NC-structure admitting an observer U such that

$$F_{ab} = 0 \tag{2.12}$$

for all $a, b = 0, \ldots, d$. We will denote this special NC-structure by $(M, \gamma, \theta, {}^{U}\Gamma)$; see (2.2).

3 Conformal Galilei transformations, Schrödinger-Virasoro Lie algebra

3.1 The Lie algebra, $\mathfrak{cgal}$, of conformal Galilei transformations

In close relationship to the Lorentzian framework, we call *conformal Galilei* transformation of (M, γ, θ) any diffeomorphism of M that preserves the direction of γ.

Owing to the fundamental constraint $\gamma^{ab}\theta_b = 0$, it follows that conformal Galilei transformation automatically preserve the direction of the time one-form θ.

In terms of infinitesimal transformations, a *conformal Galilei* vector field of (M, γ, θ) is a vector field, X, of M that Lie-transports the direction of γ; we will thus define $X \in \mathfrak{cgal}(M, \gamma, \theta)$ iff

$$L_X\gamma = f\gamma \qquad \text{hence} \qquad L_X\theta = g\,\theta \tag{3.1}$$

for some smooth functions f, g of M, depending on X. Then, $\mathfrak{cgal}(M, \gamma, \theta)$ becomes a Lie algebra whose bracket is the Lie bracket of vector fields.

The one-form θ being parallel-transported by the NC-connection, one has necessarily $d\theta = 0$; this yields $dg \wedge \theta = 0$, implying that g is (the pull-back of) a smooth function on T, i.e., that $g(t)$ depends arbitrarily on time $t = x^0$, which locally parametrizes the time axis. We thus have $dg = g'(t)\theta$.

Let us work out the expression of the generators of the *conformal Galilei Lie algebra*, $\mathfrak{cgal}(d) = \mathfrak{cgal}(\mathbb{R} \times \mathbb{R}^d, \gamma, \theta)$, of the flat NC-structure (2.1). Those are the vector fields, $X = X^0\partial_0 + X^A\partial_A$, solutions of (3.1), namely such that[6]

$$\partial_A X_B + \partial_B X_A = -f\,\delta_{AB} \tag{3.2}$$

$$\partial_A X^0 = 0 \tag{3.3}$$

$$\partial_0 X^0 = g \tag{3.4}$$

for all $A, B = 1, \ldots, d$. (We have put $X_A = \delta_{AB}X^B$.)

We readily find that $X \in \mathfrak{cgal}(d)$ iff[7]

$$X = \xi(t)\frac{\partial}{\partial t} + \Big(\omega^A_B(t)x^B + \eta^A(t) + \kappa^A(t)x_Bx^B - 2x^A\kappa_B(t)x^B + \chi(t)x^A\Big)\frac{\partial}{\partial x^A} \tag{3.5}$$

where $\boldsymbol{\omega}(t) \in \mathfrak{so}(d)$, $\boldsymbol{\eta}(t)$, $\boldsymbol{\kappa}(t)$, $\chi(t)$, and $\xi(t)$ are arbitrary functions of time, t; those are clearly interpreted as time-dependent infinitesimal rotations, space translations, expansions (or inversions), space dilations, and time reparametrizations.

We note, *en passant*, that the $\mathfrak{cgal}(d)$-generators (3.5) project as vector fields of the time axis; therefore, there exists a canonical Lie algebra homomorphism: $\mathfrak{cgal}(d) \to \mathrm{Vect}(\mathbb{R})$ given by $X \mapsto \xi(t)\partial_t$, onto the Lie algebra of vector fields of $T \cong \mathbb{R}$, i.e., the (centerless) *Virasoro Lie algebra*.

3.2 Conformal Galilei transformations, $\mathfrak{cgal}_z$, with dynamical exponent z

One can, at this stage, try and seek non-relativistic avatars of general relativistic infinitesimal conformal transformations. Given a Lorentzian (ore, more generally,

[6] Let us recall the general expressions of the Lie derivatives of γ and θ along the vector field $X = X^a\partial_a$ of M, namely $L_X\gamma^{ab} = X^c\partial_c\gamma^{ab} - 2\partial_c X^{(a}\gamma^{b)c}$, and $L_X\theta_a = \partial_a(\theta_b X^b)$.

[7] We will assume $d > 1$.

a pseudo-Riemannian) manifold (M, g), the latter Lie algebra is generated by the vector fields, X, of M such that

$$L_X(\mathrm{g}^{-1} \otimes \mathrm{g}) = 0 \tag{3.6}$$

where g^{-1} denotes the inverse of the metric $\mathrm{g} : TM \to T^*M$.

It has been shown [35] that one can expand a Lorentz metric in terms of the small parameter $1/c^2$, where c stands for the speed of light, as $\mathrm{g} = c^2\theta \otimes \theta - {}^U\gamma + \mathcal{O}(c^{-2})$, and $\mathrm{g}^{-1} = -\gamma + c^{-2}U \otimes U + \mathcal{O}(c^{-4})$, with the same notation as before. Then, a non-relativistic limit of Equation (3.6) would be $L_X \lim_{c\to\infty}(c^{-2}\,\mathrm{g}^{-1} \otimes \mathrm{g}) = 0$, viz.,

$$L_X(\gamma \otimes \theta \otimes \theta) = 0. \tag{3.7}$$

This is merely one of the possibilities at hand in our formalism. In fact, having at our disposal a Galilei structure on M, we will introduce, instead of (3.7), a more flexible condition. Indeed, owing to the degeneracy of the Galilei "metric" (γ, θ), we will deal with the following condition, namely,

$$L_X(\gamma^{\otimes m} \otimes \theta^{\otimes n}) = 0 \tag{3.8}$$

for some $m = 1, 2, 3, \ldots$, and $n = 0, 1, 2, \ldots$, to be further imposed on the vector fields $X \in \mathfrak{cgal}(M, \gamma, \theta)$. This is equivalent to Equation (3.1) together with the extra condition

$$f + q\,g = 0 \qquad \text{where} \qquad q = \frac{n}{m}. \tag{3.9}$$

Indeed, $L_X(\gamma^{\otimes m} \otimes \theta^{\otimes n}) = 0$ implies $L_X\gamma = f\gamma$ and $L_X\theta = g\,\theta$ for some functions f and g of M such that $mf + ng = 0$. Equation (3.7) plainly corresponds to the special case $m = 1$, $n = 2$.

From now on, we will call *dynamical exponent* the quantity

$$z = \frac{2}{q} \tag{3.10}$$

where q is as in (3.9). This quantity will be shown to match the ordinary notion of dynamical exponent; see, e.g., [29, 2].

We will, hence, introduce the Galilean avatars, $\mathfrak{cgal}_z(M, \gamma, \theta)$, of the Lie algebra $\mathfrak{so}(d+1, 2)$ of conformal vector fields of a pseudo-Riemannian structure of signature $(d, 1)$ as the Lie algebras spanned by the vector fields X of M satisfying (3.1), and (3.8) — or (3.9) for some rational number z. We will call $\mathfrak{cgal}_z(M, \gamma, \theta)$ the *conformal Galilei Lie algebra with dynamical exponent* z (see (3.10)).

The Lie algebra

$$\mathfrak{sv}(M, \gamma, \theta) = \mathfrak{cgal}_2(M, \gamma, \theta) \tag{3.11}$$

is the obvious generalization to Galilei spacetimes of the *Schrödinger-Virasoro* Lie algebra $\mathfrak{sv}(d) = \mathfrak{sv}(\mathbb{R} \times \mathbb{R}^d, \gamma, \theta)$ introduced in [29] (see also [2]) from a different viewpoint in the case of a flat NC-structure. The representations of the Schrödinger-Virasoro group and of its Lie algebra, $\mathfrak{sv}(d)$, as well as the deformations of the latter have been thoroughly studied and investigated in [37].

An easy calculation using (3.2), (3.4), the new constraint (3.9), and (3.10) shows that $X \in \mathfrak{cgal}_z(d)$ iff

$$X = \xi(t)\frac{\partial}{\partial t} + \Big(\omega^A_B(t)x^B + \frac{1}{z}\xi'(t)x^A + \eta^A(t)\Big)\frac{\partial}{\partial x^A} \tag{3.12}$$

where $\boldsymbol{\omega}(t) \in \mathfrak{so}(d)$, $\boldsymbol{\eta}(t)$, and $\xi(t)$ depend arbitrarily on time, t. Equation (3.12) generalizes (1.4) from $z = 1$ to any z.

The Lie algebra $\mathfrak{cgal}_\infty(M, \gamma, \theta)$ corresponding to the case $q = 0$ is interesting (see below, Section 5.3)). We have, indeed, $X \in \mathfrak{cgal}_\infty(M, \gamma, \theta)$ iff

$$L_X\gamma = 0. \tag{3.13}$$

In the case of a flat NC-structure, $\mathfrak{cgal}_\infty(d)$ is spanned by the vector fields

$$X = \xi(t)\frac{\partial}{\partial t} + \Big(\omega^A_B(t)x^B + \eta^A(t)\Big)\frac{\partial}{\partial x^A} \tag{3.14}$$

where, again, $\boldsymbol{\omega}(t) \in \mathfrak{so}(d)$, $\boldsymbol{\eta}(t)$, and $\xi(t)$ depend arbitrarily on time, t.

4 Conformal Newton-Cartan transformations

As previously emphasized, NC-connections are quite independent geometric objects; they, hence, deserve a special treatment. The idea pervading earlier work [21, 23, 25] on non-relativistic symmetries is that specifying explicitly the transformation law of the NC-connection is mandatory in a number of cases, e.g., those relevant to geometric mechanics and non-relativistic physical theories.

We will, henceforth, focus attention on the notion of Newtonian geodesics; more particularly, we will insist that the above-mentioned Galilean conformal transformations should, in addition, permute the NC-geodesics.

The geodesics of a NC-structure $(M, \gamma, \theta, \Gamma)$ are plainly geodesics of (M, Γ), i.e., the solutions of the differential equations

$$\ddot{x}^c + \Gamma^c_{ab}\,\dot{x}^a\dot{x}^b = \mu\,\dot{x}^c \tag{4.1}$$

for all $c = 0, \ldots, d$, where μ is some smooth (fiberwise linear) function of TM; here, we have put $\dot{x}^a = dx^a/d\tau$, where τ is an otherwise arbitrary curve-parameter.

Let us remind that Equation (4.1) models *free fall* in NC theory [30, 31, 32], just as it does in general relativity. By putting $\dot{t} = \theta_a \dot{x}^a$, we characterize[8]

$$\textit{timelike} \text{ geodesics by:} \quad \dot{t} \neq 0 \tag{4.2}$$

$$\textit{lightlike} \text{ geodesics by:} \quad \dot{t} = 0. \tag{4.3}$$

Spacetime transformations which permute the geodesics of (M, Γ), i.e., preserve the form of the geodesic equation (4.1), are *projective* transformations; they form the projective group of the affine structure. Infinitesimal projective transformations generate a Lie algebra which, hence, consists of vector fields, X, of M satisfying

$$L_X \Gamma^c_{ab} = \delta^c_a \varphi_b + \delta^c_b \varphi_a \tag{4.4}$$

for a certain one-form $\varphi = \varphi_a dx^a$ of M depending on X.

4.1 The Schrödinger Lie algebra

Let us first cope with generic, *timelike*, geodesics of (M, Γ) defined by $\dot{t} \neq 0$, cf. Equation (4.2), and representing the worldlines of *massive non-relativistic test particles*. From now on, we choose to enforce preservation of their equations (4.1), in addition to that, (3.1), of the direction of the Galilei structure (γ, θ).

4.1.1 The expanded Schrödinger Lie algebra, $\widetilde{\mathfrak{sch}}$, of projective Galilei conformal transformations

The Lie-transport (4.4) of the NC-connection, compatible with the conformal rescalings (3.1) of the Galilei structure (γ, θ), must preserve the first constraint $\nabla\theta = 0$, i.e., $L_X \nabla_a \theta_b = \nabla_a L_X \theta_b - \theta_c L_X \Gamma^c_{ab} = 0$; this yields $g' \theta_a \theta_b - 2\theta_{(a}\varphi_{b)} = 0$, or $\varphi_a = \frac{1}{2} g' \theta_a$. The infinitesimal projective transformations to consider are thus given by

$$L_X \Gamma^c_{ab} = g' \delta^c_{(a} \theta_{b)}. \tag{4.5}$$

Likewise, preservation of the second constraint, viz., $\nabla\gamma = 0$, necessarily implies $L_X \nabla_c \gamma^{ab} = \nabla_c L_X \gamma^{ab} + 2 L_X \Gamma^{(a}_{ck} \gamma^{b)k} = 0$; we thus find $(\partial_c f + g' \theta_c)\gamma^{ab} = 0$, and f is therefore a function of T such that

$$f' + g' = 0. \tag{4.6}$$

[8] The condition $\dot{t} = 0$ is clearly a first-integral of Equation (4.1). Lightlike — or *null* — geodesics are, hence, spacelike; the origin of the terminology will be explained later, in Section 5.

We will, hence, define a new Lie algebra, denoted $\widetilde{\mathfrak{sch}}(M,\gamma,\theta,\Gamma)$, as the Lie algebra of those vector fields that are infinitesimal (i) conformal Galilei transformations of (M,γ,θ), and (ii) projective transformations of (M,Γ). We call $\widetilde{\mathfrak{sch}}(M,\gamma,\theta,\Gamma)$ the *expanded Schrödinger Lie algebra*, which is therefore spanned by the vector fields, X, of M such that [21, 22, 23]

$$L_X\gamma^{ab} = f\gamma^{ab}, \qquad L_X\theta_a = g\,\theta_a \qquad \& \qquad L_X\Gamma^c_{ab} = g'\delta^c_{(a}\theta_{b)} \tag{4.7}$$

for all $a,b,c = 0,1,\ldots,d$, and subject to Condition (4.6).

Let us now work out the form of the Schrödinger Lie algebra in the flat case. We will thus determine the generators of the Lie algebra $\widetilde{\mathfrak{sch}}(d) = \widetilde{\mathfrak{sch}}(\mathbb{R}\times\mathbb{R}^d,\gamma,\theta,\Gamma)$ in the special case (2.1). The system (4.7) to solve for $X = X^0\partial_0 + X^A\partial_A$ reads[9]

$$\partial_A X_B + \partial_B X_A = -f\delta_{AB} \tag{4.8}$$
$$\partial_A X^0 = 0 \tag{4.9}$$
$$\partial_0 X^0 = g \tag{4.10}$$
$$\partial_0\partial_0 X^A = 0 \tag{4.11}$$
$$\partial_0\partial_B X^A = \frac{1}{2}g'\delta^A_B \tag{4.12}$$
$$\partial_A\partial_B X^C = 0 \tag{4.13}$$

for all $A,B,C = 1,\ldots,d$.

We deduce, from (4.13) that $X^A = M^A_B(t)x^B + \eta^A(t)$, and, using (4.8), we find $M^A_B(t) = \omega^A_B - \frac{1}{2}f(t)\delta^A_B$, where the $\omega_{AB} = -\omega_{BA}$ are independent of t. Then (4.11) leaves us with $f''(t) = 0$, and $(\eta^A)''(t) = 0$, i.e., with $f(t) = -2(\kappa t + \lambda)$, and $\eta^A(t) = \beta^A t + \gamma^A$, where κ,λ,β^A, and γ^A are constant coefficients. At last, using Equations (4.6) and (4.10), we conclude that $X^0 = \kappa t^2 + \mu t + \varepsilon$, with μ,ε new constants of integration.

We can therefore affirm that $X \in \widetilde{\mathfrak{sch}}(d)$ iff

$$X = \left(\kappa t^2 + \mu t + \varepsilon\right)\frac{\partial}{\partial t} + \left(\omega^A_B\,x^B + \kappa t x^A + \lambda x^A + \beta^A t + \gamma^A\right)\frac{\partial}{\partial x^A} \tag{4.14}$$

where $\boldsymbol{\omega} \in \mathfrak{so}(d)$, $\boldsymbol{\beta},\boldsymbol{\gamma} \in \mathbb{R}^d$, and $\kappa,\mu,\lambda,\varepsilon \in \mathbb{R}$ are respectively infinitesimal rotations, boosts, spatial translations, inversions, time dilations, space dilations, and time translations. We observe in (4.14) that time is dilated *independently* of space [21, 23]. The expanded Schrödinger Lie algebra, $\widetilde{\mathfrak{sch}}(d)$, is a finite-dimensional Lie subalgebra of $\mathfrak{cgal}(d)$.

[9] Let us recall that the Lie derivative of a linear connection, Γ, along the vector field, X, is given by $L_X\Gamma^c_{ab} = \partial_a\partial_b X^c$ in the *flat* case, and in a coordinate system where $\Gamma^c_{ab} = 0$.

Let us recall that the *Galilei Lie algebra* $\mathfrak{gal}(M,\gamma,\theta,\Gamma) \subset \mathfrak{cgal}(M,\gamma,\theta,\Gamma)$ of a NC-structure is plainly defined as its Lie algebra of infinitesimal automorphisms. Thus, $X \in \mathfrak{gal}(M,\gamma,\theta,\Gamma)$ iff [31, 25]

$$L_X\gamma^{ab} = 0, \qquad L_X\theta_a = 0 \qquad \& \qquad L_X\Gamma^c_{ab} = 0 \tag{4.15}$$

for all $a,b,c = 0,1,\ldots,d$, i.e., if (4.7) holds with $f = 0$, and $g = 0$. In the flat case, $\mathfrak{gal}(d) = \mathfrak{gal}(\mathbb{R}\times\mathbb{R}^d,\gamma,\theta,\Gamma)$ is clearly spanned by the vector fields (4.14) with $\kappa = 0$, and $\lambda = \mu = 0$.

4.1.2 The Schrödinger Lie algebras, $\mathfrak{sch}_z$, with dynamical exponent z

Just as in Section 3.2, we define *Schrödinger Lie algebra with dynamical exponent* z as the Lie subalgebra $\mathfrak{sch}_z(M,\gamma,\theta,\Gamma) \subset \widetilde{\mathfrak{sch}}(M,\gamma,\theta,\Gamma)$ defined by the supplementary condition (3.9), i.e.,

$$f + \frac{2}{z}g = 0 \tag{4.16}$$

where z is given by (3.10). This entails, via Equation (4.6), that

$$\Big(\frac{2}{z} - 1\Big)g'(t) = 0. \tag{4.17}$$

- We, hence, find

$$z = 2 \tag{4.18}$$

since $g' \neq 0$, *generically*. For the flat NC-structure, see (4.14), this implies that time is dilated *twice* as much as space [14], a specific property of the (centerless) *Schrödinger Lie algebra*

$$\mathfrak{sch}(d) = \mathfrak{sch}_2(d) \tag{4.19}$$

for which

$$\mu = 2\lambda. \tag{4.20}$$

We therefore contend that $X \in \mathfrak{sch}_2(d)$ iff

$$X = \big(\kappa t^2 + 2\lambda t + \varepsilon\big)\frac{\partial}{\partial t} + \big(\omega^A_B x^B + \kappa t x^A + \lambda x^A + \beta^A t + \gamma^A\big)\frac{\partial}{\partial x^A} \tag{4.21}$$

where $\boldsymbol{\omega} \in \mathfrak{so}(d)$, $\boldsymbol{\beta},\boldsymbol{\gamma} \in \mathbb{R}^d$, and $\kappa,\lambda,\varepsilon \in \mathbb{R}$. The Schrödinger *dynamical exponent* is $z = 2$; see, e.g., [29].

The Lie algebra $\mathfrak{sch}(d)$ admits the faithful $(d+2)$-dimensional representation $X \mapsto Z$ where

$$Z = \begin{pmatrix} \boldsymbol{\omega} & \boldsymbol{\beta} & \boldsymbol{\gamma} \\ 0 & \lambda & \varepsilon \\ 0 & -\kappa & -\lambda \end{pmatrix} \tag{4.22}$$

with the same notation as above.

We therefore have the Levi decomposition

$$\mathfrak{sch}(d) \cong (\mathfrak{so}(d) \times \mathfrak{sl}(2,\mathbb{R})) \ltimes (\mathbb{R}^d \times \mathbb{R}^d). \tag{4.23}$$

The Schrödinger Lie algebra is, indeed, a *finite-dimensional* Lie subalgebra of the Schrödinger-Virasoro Lie algebra (3.11), viz.,

$$\mathfrak{sch}_2(M,\gamma,\theta,\Gamma) \subset \mathfrak{sv}(M,\gamma,\theta). \tag{4.24}$$

• Returning to Equation (4.17) we get, in the special case $g' = 0$, a family of Lie subalgebras algebras $\mathfrak{sch}_z(M,\gamma,\theta,\Gamma) \subset \widetilde{\mathfrak{sch}}(M,\gamma,\theta,\Gamma)$ parametrized by a (rational) dynamical exponent, z. In the flat case, $\mathfrak{sch}_{z\neq 2}(d)$ is spanned by the vector fields (4.14) with $\kappa = 0$, and $\mu = z\lambda$.

In the limit $z \to \infty$, where $f = g' = 0$ in view of (4.6), we obtain the Lie algebra $\mathfrak{sch}_\infty(M,\gamma,\theta,\Gamma)$. For flat NC-spacetime, $\mathfrak{sch}_\infty(d)$ is generated by the vector fields (4.14) with $\kappa = \lambda = 0$.

In both cases we get the Lie algebra of vector fields of the form

$$X = (\mu t + \varepsilon)\,\frac{\partial}{\partial t} + \left(\omega^A_B\, x^B + \frac{\mu}{z}\, x^A + \beta^A t + \gamma^A\right)\frac{\partial}{\partial x^A} \tag{4.25}$$

with the same notation as above.

4.2 Transformation law of NC-connections under conformal Galilei rescalings

The rest of the section will be devoted to the specialization of projective transformations to the specific case of lightlike (4.3) NC-geodesics.

Let us now work out the general form of the variation, $\delta\Gamma$, of a NC-connection, Γ, under infinitesimal conformal rescalings of the Galilei structure (γ,θ) of M, namely

$$\delta\gamma = f\gamma \qquad \text{hence} \qquad \delta\theta = g\,\theta \tag{4.26}$$

where f is an arbitrary function of M, and g an arbitrary function of T (compare Equation (3.1)). We will furthermore put, in full generality,

$$\delta U = -gU + \gamma(\psi) \tag{4.27}$$

in order to comply with the constraint $\theta_a U^a = 1$, where ψ is an arbitrary one-form of M interpreted as an infinitesimal Milne boost (cf. (2.7)).

Starting from (2.2), we get $\delta\Gamma^c_{ab} = (\delta^U\Gamma)^c_{ab} + \delta\theta_{(a}F_{b)k}\gamma^{ck} + \theta_{(a}\delta F_{b)k}\gamma^{ck} + \theta_{(a}F_{b)k}\delta\gamma^{ck}$. Then, using (4.26) and (4.27) applied to the expression (2.3) of the NC-connection ${}^U\Gamma$, we find $\delta({}^U\Gamma)^c_{ab} = -\delta^c_{(a}\partial_{b)}f + U^c\theta_{(a}\partial_{b)}(f+g) + \frac{1}{2}\left(\gamma^{ck}\partial_k f\right){}^U\gamma_{ab} - \theta_{(a}d\phi_{b)k}\gamma^{ck}$, where $\phi = {}^U\gamma(\delta U)$ is the one-form (2.8) associated with the "Milne" variation (4.27) of the observer U. Then, with the help of Equation (4.26), and of general result $\delta F = d\phi$, see (2.7), we can finally claim that

$$\delta\Gamma^c_{ab} = -\delta^c_{(a}\partial_{b)}f + U^c\theta_{(a}\partial_{b)}(f+g) + \frac{1}{2}\left(\gamma^{ck}\partial_k f\right){}^U\gamma_{ab} + (f+g)\gamma^{ck}\theta_{(a}F_{b)k} \tag{4.28}$$

for all $a, b, c = 0, \ldots, d$.

Equation (4.28) is of central importance in our study; it yields the general form of the variations of the NC-connection compatible with the constraints $\nabla\gamma = 0$, and $\nabla\theta = 0$, and induced by the conformal Galilei rescalings (4.26) and the Milne boosts (4.27).

4.3 Conformal NC transformations: lightlike geodesics

So far, we have been dealing with the Galilei-conformal symmetries of the equations of generic, i.e., timelike geodesics. What about those of the equations of *lightlike geodesics* (4.3) that model the worldlines of *massless non-relativistic particles* [39]?

Let us now determine the variations (4.28) of the NC-connection, Γ, that preserve the equations of lightlike geodesics, namely Equation (4.1) supplemented by $\dot{t} = 0$.

We thus must have $\delta\Gamma^c_{ab}\dot{x}^a\dot{x}^b = \delta\mu\,\dot{x}^c$, so that, necessarily,

$$\gamma(df) = 0 \tag{4.29}$$

since $\theta_a\dot{x}^a = 0$ (the third term in the right-hand side of (4.28) has to vanish); this implies $df = f'\theta$, hence that f is, along with g, a function of the time axis, T. We also find that $\delta\mu = 0$. At last, the resulting variation of the NC-connection appears in the new guise

$$\delta\Gamma^c_{ab} = -f'\delta^c_{(a}\theta_{b)} + (f'+g')U^c\theta_a\theta_b + (f+g)\gamma^{ck}\theta_{(a}F_{b)k} \tag{4.30}$$

for all $a, b, c = 0, \ldots, d$, where the unit vector field U, i.e., $\theta_a U^a = 1$, and the two-form F are as in (2.2).

We note that the constraint (4.6), obtained in the "massive" case, does *not* show up in the "massless" case.

Just as in Section 4.1, we will assume that the variations (4.26) of the Galilei structure, and those (4.30) of the NC-connection, are generated by infinitesimal spacetime transformations.

4.3.1 The conformal Newton-Cartan Lie algebra, $\mathfrak{cnc}$, of null-projective conformal Galilei transformations

The next natural step consists in demanding that the variations (4.26) of the Galilei structure, and those (4.30) of the NC-connection are, indeed, generated by infinitesimal spacetime transformations.

We will thus define a new Lie algebra, called the *conformal Newton-Cartan Lie algebra*, and denoted by $\mathfrak{cnc}(M,\gamma,\theta,\Gamma)$, as the Lie algebra of those vector fields that are infinitesimal (i) conformal-Galilei transformation of (M,γ,θ), and (ii) transformations which permute *lightlike* geodesics of (M,γ,θ,Γ). The *conformal Newton-Cartan Lie algebra*, $\mathfrak{cnc}(M,\gamma,\theta,\Gamma)$, is thus spanned by the vector fields, X, of M such that

$$L_X\gamma^{ab} = f\gamma^{ab} \tag{4.31}$$

$$L_X\theta_a = g\,\theta_a \tag{4.32}$$

$$L_X\Gamma^c_{ab} = -f'\delta^c_{(a}\theta_{b)} + (f'+g')U^c\theta_a\theta_b + (f+g)\gamma^{ck}\theta_{(a}F_{b)k} \tag{4.33}$$

for all $a,b,c=0,1,\ldots,d$, where f and g are functions of the time axis, T, while U, and F are as in (2.2) and (2.3).

It is worth noticing that the Lie-transport (4.33) of the NC-connection satisfies the very simple condition, viz.,

$$L_X\Gamma^{abc} = 0 \tag{4.34}$$

where $L_X\Gamma^{abc} = (L_X\Gamma^c_{k\ell})\gamma^{ak}\gamma^{b\ell}$. Interestingly, Equation (4.34) is specific to the so-called *Coriolis* Lie algebra of Galilei isometries of (M,γ,θ); see [25].

Let us emphasize, at this stage, that the Schrödinger Lie algebra we have already been dealing with in Section 4.1.2, is clearly a Lie subalgebra of the conformal Newton-Cartan Lie algebra, viz.,

$$\mathfrak{sch}_2(M,\gamma,\theta,\Gamma) \subset \mathfrak{cnc}(M,\gamma,\theta,\Gamma) \tag{4.35}$$

corresponding to the constraint

$$f+g=0 \tag{4.36}$$

associated with the dynamical exponent $z=2$; see (4.18). We will thus ignore, in the sequel, this special solution, and concentrate on the maximal solutions of Equations (4.31)–(4.33) with $f+g\neq 0$.

We will now determine the conformal Newton-Cartan Lie algebra in the flat case, i.e., the Lie algebra $\mathfrak{cnc}(d) = \mathfrak{cnc}(\mathbb{R}\times\mathbb{R}^d,\gamma,\theta,\Gamma)$ where γ^{ab} and θ_a are as in (2.1), as well as $\Gamma^c_{ab}=0$.

Let us put, in full generality, $U=\partial_0+U^A\partial_A$, where the U^A are smooth functions of spacetime.

The system (4.31)–(4.33) to solve for $X = X^0\partial_0 + X^A\partial_A$ reads then

$$\begin{aligned}
\partial_A X_B + \partial_B X_A &= -f\delta_{AB} && (4.37)\\
\partial_A X^0 &= 0 && (4.38)\\
\partial_0 X^0 &= g && (4.39)\\
\partial_0\partial_0 X^A &= (f' + g')U^A + (f+g)F_{0A} && (4.40)\\
\partial_0\partial_B X^A &= -\frac{1}{2}f'\delta^A_B - \frac{1}{2}(f+g)F_{AB} && (4.41)\\
\partial_A\partial_B X^C &= 0 && (4.42)
\end{aligned}$$

for all $A, B, C = 1, \ldots, d$.

Straightforward computation provides the general solution of that system. We find that $X^0 = \xi(t)$, hence $g(t) = \xi'(t)$, remains arbitrary; Equations (4.37), and (4.42) yield $X^A = \omega^A_B(t)x^B - \frac{1}{2}f(t)x^A + \eta^A(t)$, the functions $\omega_{AB}(t) = -\omega_{BA}(t)$, $f(t)$, and $\eta^A(t)$ being unspecified. Conspicuously, Equations (4.40), and (4.41), bring no further restriction to the spatial components, X^A, as long as the two-form F, in the right-hand side of these equations, is not constrained whatsoever. Indeed, we can easily deduce from (2.10), and (2.11) that

$$F_{AB} = 2\partial_{[A}U_{B]}. \tag{4.43}$$

and

$$F_{0A} = \partial_0 U_A + U^B\partial_A U_B. \tag{4.44}$$

Using then (4.41), and (4.40), we find that $\omega'_{AB}(t)x^B = (f+g)U_A + \partial_A\psi$, as well as $-\frac{1}{2}f''(t)\delta_{AB}\,x^B + \eta''_A(t) = \partial_A\chi$, for some functions ψ, and χ, and some unit vector field, U, of spacetime. Our claim is, hence, justified.

We contend that $X \in \mathfrak{cnc}(d)$ iff

$$X = \xi(t)\frac{\partial}{\partial t} + \Big(\omega^A_B(t)x^B - \frac{1}{2}f(t)x^A + \eta^A(t)\Big)\frac{\partial}{\partial x^A}, \tag{4.45}$$

where, $\boldsymbol{\omega}(t) \in \mathfrak{so}(d)$, $\boldsymbol{\eta}(t)$, $\xi(t)$ and $f(t)$ depend smoothly on time, t, in an arbitrary fashion.

If $\mathfrak{cnc}_z(d)$ denotes the Lie subalgebra with dynamical exponent z, i.e., defined by Equation (4.16), we trivially have

$$\mathfrak{cnc}_z(d) \cong \mathfrak{cgal}_z(d) \tag{4.46}$$

in view of (3.12). Let us emphasize that dealing with *rational* dynamical exponents, z, introduced in (3.10), is clearly allowed by the novel geometric definition (3.8) of conformal Galilei transformations.

4.3.2 The Lie algebra, $\mathfrak{cmil}$, of null-projective conformal transformations of NC-Milne spacetime

We will now confine considerations to the case where the NC-spacetime admits a preferred geodesic and irrotational observer (an "ether"), i.e., a unit vector field, U, such that, Condition (2.12) holds true. With the convention of Section 2.2, we denote such a "NC-Milne-structure" by $(M, \gamma, \theta, {}^U\Gamma)$.

Specializing the system (4.31)–(4.33) to the case $F = 0$, we thereby define the *conformal Milne Lie algebra*, $\mathfrak{cmil}(M, \gamma, \theta, {}^U\Gamma)$, as the *maximal* Lie algebra of vector fields, X, of M such that

$$L_X \gamma^{ab} = f\gamma^{ab} \tag{4.47}$$

$$L_X \theta_a = g\,\theta_a \tag{4.48}$$

$$L_X {}^U\Gamma^c_{ab} = -f'\delta^c_{(a}\theta_{b)} + (f' + g')U^c\theta_a\theta_b \tag{4.49}$$

for all $a, b, c = 0, 1, \ldots, d$, where f and g are functions of the time axis, T.

We now determine the Lie algebra $\mathfrak{cmil}(d) = \mathfrak{cmil}(\mathbb{R}\times\mathbb{R}^d, \gamma, \theta, {}^U\Gamma)$, in the special case of flat NC-Milne spacetime specified by Equations (2.1), where $\Gamma^c_{ab} = {}^U\Gamma^c_{ab} = 0$ for all $a, b, c = 0, 1, \ldots, d$, in a chosen inertial coordinate system — we have, in particular, $U = \partial_0 + U^A\partial_A$ where

$$U^A = \text{const.} \tag{4.50}$$

for all $A = 1, \ldots, d$. Indeed, an ether in flat NC-spacetime is a solution, U, of the PDE (4.43) and (4.44) with $F = 0$. We get $U_A = \partial_A\psi$ and $\partial_A(\partial_0\psi + \frac{1}{2}U_B U^B) = 0$. One can thus choose ψ to be a solution of the free Hamilton-Jacobi equation

$$\partial_t\psi + \frac{1}{2}\delta^{AB}\partial_A\psi\partial_B\psi = 0 \tag{4.51}$$

whose general solution, ψ, is well-known and leads to $\partial_A\psi = U_A$ where (4.50) holds.

The system (4.47)–(4.49) to solve for $X = X^0\partial_0 + X^A\partial_A$ is now given by

$$\partial_A X_B + \partial_B X_A = -f\delta_{AB} \tag{4.52}$$

$$\partial_A X^0 = 0 \tag{4.53}$$

$$\partial_0 X^0 = g \tag{4.54}$$

$$\partial_0\partial_0 X^A = (f' + g')U^A \tag{4.55}$$

$$\partial_0\partial_B X^A = -\frac{1}{2}f'\delta^A_B \tag{4.56}$$

$$\partial_A\partial_B X^C = 0 \tag{4.57}$$

for all $A, B, C = 1, \ldots, d$.

Returning to the calculation done in Section 4.3.1, we get $X^0 = \xi(t)$, with $g(t) = \xi'(t)$, and $X^A = \omega^A_B(t)x^B - \frac{1}{2}f(t)x^A + \eta^A(t)$, with the same notation as before. Equation (4.56) readily implies

$$\omega'_{AB}(t) = 0 \tag{4.58}$$

while Equation (4.55) yields

$$(f' + g')U^A = -\frac{1}{2}f''\,x^A + (\eta^A)''. \tag{4.59}$$

This entails that $f'' = 0$, i.e.

$$f(t) = -2(\kappa t + \lambda). \tag{4.60}$$

The latter equations therefore imply $(f' + g')U^A = (\eta^A)''$, leading us to the expression

$$\eta^A(t) = \alpha^A(\xi(t) - \kappa t^2) + \widetilde{\beta}^A t + \widetilde{\gamma}^A \tag{4.61}$$

where we have put

$$\alpha^A = U^A \tag{4.62}$$

and where the coefficients $\widetilde{\beta}^A$, and $\widetilde{\gamma}^A$ are integration constants.

Now, if X_1, X_2 are solutions of the system (4.47)–(4.49), so is their Lie bracket $X_{12} = [X_1, X_2]$. This yields the consistency relation $f_{12} = X_1 f_2 - X_2 f_1$ which reads here $\kappa_{12}t + \lambda_{12} = \xi_1(t)\kappa_2 - \xi_2(t)\kappa_1$. Thus $\xi(t) = \kappa\, u(t) + \mu t + \varepsilon$, for some function u, the coefficients μ, and ε being constant. Exploiting the fact that $X \mapsto X^0\partial_0$ is a Lie algebra homomorphism into $\mathrm{Vect}(\mathbb{R})$, we write $\xi_{12}(t) = \xi_1(t)\xi_2'(t) - \xi_2(t)\xi_1'(t)$; straightforward calculation then shows that u is a polynomial of degree 2, which, up to lower degree terms, is given by

$$u(t) = \frac{1}{2}c\,t^2 \tag{4.63}$$

with

$$(c-1)(c-2) = 0. \tag{4.64}$$

We thus find $X^0 = \xi(t)$ where $\xi(t) = \frac{1}{2}\kappa c\,t^2 + \mu t + \varepsilon$, and $X^A = \omega^A_B\,x^B + \kappa t x^A + \lambda x^A + \frac{1}{2}(c-2)\kappa t^2\alpha^A + \beta^A t + \gamma^A$, with β^A, and γ^A new integration constants.

The case $c = 2$ gives back the expanded Schrödinger Lie algebra, $\widetilde{\mathfrak{sch}}(d)$, see (4.14), already studied since $f'(t) + g'(t) = 0$. It is not the only possibility, though.[10]

[10]The Schrödinger Lie algebra is the Lie algebra of a group of spacetime transformations that actually permute all geodesics, in particular lightlike geodesics.

Consider then the new case $c = 1$. We claim that $X \in \mathfrak{cmil}(d)$ iff

$$\begin{aligned} X &= \left(\frac{1}{2}\kappa t^2 + \mu t + \varepsilon\right)\frac{\partial}{\partial t} \\ &\quad + \left(\omega^A_B x^B + \kappa t x^A + \lambda x^A - \frac{1}{2}t^2\alpha^A + \beta^A t + \gamma^A\right)\frac{\partial}{\partial x^A} \end{aligned} \tag{4.65}$$

where $\boldsymbol{\omega} \in \mathfrak{so}(d)$, $\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \in \mathbb{R}^d$, and $\kappa, \lambda, \mu, \varepsilon \in \mathbb{R}$.

Let us highlight that, just as in the case of $\widetilde{\mathfrak{sch}}(d)$, time and space dilations are independent within $\mathfrak{cmil}(d)$. As for the parameter, $\boldsymbol{\alpha}$, in (4.65) it serves as a novel *acceleration* generator [18, 1].

4.3.3 Conformal NC-Milne Lie algebras, $\mathfrak{cmil}_z$, with dynamical exponent z; the CGA Lie algebra

Much in the same way than in Section 3.2, we will now introduce subalgebras of the conformal NC-Milne Lie algebra with prescribed dynamical exponent, z.

We will define $\mathfrak{cmil}_z(M, \gamma, \theta, {}^U\Gamma)$ as the Lie subalgebra of $\mathfrak{cmil}(M, \gamma, \theta, {}^U\Gamma)$ defined by Equation (4.16); we will call it the *conformal NC-Milne Lie algebra with dynamical exponent z*.

Let us lastly establish, in the case of a flat NC-Milne structure, the expression of the generators of the Lie algebra $\mathfrak{cmil}_z(d) = \mathfrak{cmil}_z(\mathbb{R} \times \mathbb{R}^d, \gamma, \theta, {}^U\Gamma)$. Those retain the form (4.65) where, in view of (4.54), and (4.60), Equation (4.16) writes

$$\frac{1}{z}\left((1-z)\kappa t + (\mu - z\lambda)\right) = 0. \tag{4.66}$$

- In the generic case, $f' \neq 0$, one ends up with

$$z = 1 \tag{4.67}$$

and

$$\mu = \lambda \tag{4.68}$$

which entails that time and space are related in the same way. Therefore $\mathfrak{cmil}_1(d)$ is spanned by the vector fields (4.65) for which (4.68) holds. It is isomorphic to the CGA, namely, the *Conformal Galilean Algebra* (1.3) of Lukierski, Stichel and Zakrzewski [1], i.e.,

$$X = \left(\frac{1}{2}\kappa t^2 + \lambda t + \varepsilon\right)\frac{\partial}{\partial t} + \left(\omega^A_B x^B + \lambda x^A + \kappa t x^A - \frac{1}{2}\alpha^A t^2 + \beta^A t + \gamma^A\right)\frac{\partial}{\partial x^A}. \tag{4.69}$$

where $\boldsymbol{\omega} \in \mathfrak{so}(d)$, $\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \in \mathbb{R}^d$, and $\kappa, \lambda, \varepsilon \in \mathbb{R}$.

The Lie algebra $\mathfrak{cmil}_1(d)$ admits the faithful $(d+3)$-dimensional representation $X \mapsto Z$, where

$$Z = \begin{pmatrix} \boldsymbol{\omega} & -\frac{1}{2}\boldsymbol{\alpha} & \boldsymbol{\beta} & \boldsymbol{\gamma} \\ 0 & \lambda & 2\varepsilon & 0 \\ 0 & \frac{1}{2}\kappa & 0 & \varepsilon \\ 0 & 0 & -\kappa & -\lambda \end{pmatrix} \tag{4.70}$$

with the same notation as before.

Again, the following decomposition holds

$$\mathfrak{cmil}_1(d) \cong (\mathfrak{so}(d) \times \mathfrak{so}(2,1)) \ltimes (\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d). \tag{4.71}$$

• In the special case, $f' = 0$, i.e., $\kappa = 0$ (no expansions), and $\mu = z\lambda$, we discover a whole family of Lie subalgebras $\mathfrak{cmil}_z(d) \subset \mathfrak{cmil}(d)$ parametrized by an arbitrary dynamical exponent, z.

In the limit $z \to \infty$, where $f = 0$, the Lie subalgebra $\mathfrak{cmil}_\infty(d) \subset \mathfrak{cmil}(d)$ is spanned by the vector fields (4.65) with $\kappa = \lambda = 0$. Let us stress that, in this limiting case, space dilations are ruled out (compare Equation (3.14)).

In both cases we obtain the Lie algebra of vector fields

$$X = \Big(\mu t + \varepsilon\Big)\frac{\partial}{\partial t} + \Big(\omega^A_B\, x^B + \frac{\mu}{z}\, x^A - \frac{1}{2}\alpha^A t^2 + \beta^A t + \gamma^A\Big)\frac{\partial}{\partial x^A}. \tag{4.72}$$

with the same notation as before.

4.3.4 The finite-dimensional conformal Galilei Lie algebras, $\mathfrak{alt}_{2/N}(d)$

Our formalism leads thus to an intrinsic definition of distinguished finite-dimensional subalgebras of the conformal Galilei Lie algebra $\mathfrak{cgal}(d)$, namely $\mathfrak{sch}_2(d)$, and $\mathfrak{cmil}_1(d)$ with dynamical exponents $z = 2$, and $z = 1$ respectively (see (4.23), and (4.71)). Restricting, here, considerations to the very special case of *flat* NC structures (expressed in a given Cartesian coordinate system), one might search for other finite-dimensional Lie subalgebras of the conformal Galilei Lie algebras, $\mathfrak{cgal}_z(d) \cong \mathfrak{cnc}_z(d)$, with prescribed dynamical exponent z; see (4.46).

Recall (see (3.12)) that $\mathfrak{cgal}_z(d)$ is generated by those $X \in \mathrm{Vect}(\mathbb{R} \times \mathbb{R}^d)$ of the form

$$X = \xi(t)\frac{\partial}{\partial t} + \Big(\omega^A_B(t)x^B + \frac{1}{z}\xi'(t)x^A + \eta^A(t)\Big)\frac{\partial}{\partial x^A} \tag{4.73}$$

where $\boldsymbol{\omega}(t) \in \mathfrak{so}(d)$, $\boldsymbol{\eta}(t)$, and $\xi(t)$ depend *smoothly* on on time, t.

Previous experience with the above-mentioned Lie algebras prompts us to look for Lie algebras of *polynomial* — not merely smooth — vector fields of $\mathfrak{cgal}_z(d)$.

Consider, hence, vector fields, $X \in \mathfrak{cgal}_z(d)$, that are polynomials of fixed degree $N > 0$ in the variables $t = x^0, x^1, \ldots, x^d$. This entails the following decompositions $\boldsymbol{\omega}(t) = \sum_{n=0}^{N} \omega_n t^n$, $\boldsymbol{\eta}(t) = \sum_{n=0}^{N} \boldsymbol{\eta}_n t^n$, and $\xi(t) = \sum_{n=0}^{N} \xi_n t^n$, since the spatial components X^A are already of first order in $x^1, \ldots, x^d$. Bearing in mind that $X \mapsto \xi$ is a Lie algebra homomorphism, we claim that the $\xi = \xi(t)\partial/\partial t$ do span a polynomial Lie subalgebra of $\mathrm{Vect}(\mathbb{R})$, hence a Lie subalgebra of $\mathfrak{sl}(2,\mathbb{R})$ since the latter is maximal in the Lie algebra, $\mathrm{Vect}^{\mathrm{Pol}}(\mathbb{R})$, of polynomial vector fields of $\mathbb{R}$. We therefore find $\xi_n = 0$ for all $n \geq 3$, so that

$$\xi(t) = \frac{1}{2}\kappa t^2 + \mu t + \varepsilon \tag{4.74}$$

with $\kappa, \mu, \varepsilon \in \mathbb{R}$.

Let now us seek under which condition (if any) the Lie bracket $X_{12} = [X_1, X_2]$ of two such polynomial vector fields X_1 and X_2 is, itself, polynomial of degree N, Condition (4.74) being granted. Straightforward calculation yields

$$\xi_{12} = \xi_1\xi_2' - \xi_2\xi_1' \tag{4.75}$$

$$\boldsymbol{\omega}_{12} = [\boldsymbol{\omega}_2, \boldsymbol{\omega}_1] + \xi_1\boldsymbol{\omega}_2' - \xi_1\boldsymbol{\omega}_1' \tag{4.76}$$

$$\boldsymbol{\eta}_{12} = \boldsymbol{\omega}_2\boldsymbol{\eta}_1 - \boldsymbol{\omega}_1\boldsymbol{\eta}_2 + \xi_1\boldsymbol{\eta}_2' - \xi_2\boldsymbol{\eta}_1' - \frac{1}{z}\left(\xi_1'\boldsymbol{\eta}_2 - \xi_2'\boldsymbol{\eta}_1\right). \tag{4.77}$$

Condition (4.75) brings no further restriction in view of (4.74). From (4.76), we discover that, necessarily, $\boldsymbol{\omega}_1' = \boldsymbol{\omega}_2' = 0$; this entails that

$$\boldsymbol{\omega} \in \mathfrak{so}(d) \tag{4.78}$$

in (4.73). At last, we readily find that the right hand-side of Equation (4.77) turns out to be a polynomial of degree $N+1$ in t, namely $\boldsymbol{\eta}_{12} = \sum_{n=0}^{N+1} (\boldsymbol{\eta}_{12})_n t^n$ with $(\boldsymbol{\eta}_{12})_{N+1} = \left(\frac{1}{2}N - z\right)(\kappa_1(\boldsymbol{\eta}_2)_N - \kappa_2(\boldsymbol{\eta}_1)_N)$. In order to acquire a Lie algebra of polynomial vector fields of degree $N > 0$, we must simply impose the constraint

$$z = \frac{2}{N} \tag{4.79}$$

on the dynamical exponent. At last, we have shown that, in Equation (4.73),

$$\boldsymbol{\eta}(t) = \boldsymbol{\eta}_N t^N + \cdots + \boldsymbol{\eta}_1 t + \boldsymbol{\eta}_0 \tag{4.80}$$

with $\boldsymbol{\eta}_n \in \mathbb{R}^d$ for all $n = 0, 1, \ldots, N = 2/z$.

We claim that the finite-dimensional Lie subalgebras of $\mathfrak{cgal}_{2/N}(d)$ defined by (4.74), (4.78), and (4.80) together with (4.79) are isomorphic with the so-called

$\mathfrak{alt}_{2/N}(d)$ Lie algebras discovered by Henkel [29] in the study of scale invariance for strongly anisotropic critical systems (with $d = 1$).[11] We have thus proved that

$$\mathfrak{cgal}^{\mathrm{Pol}}_{2/N}(d) \cong \mathfrak{alt}_{2/N}(d). \tag{4.81}$$

Note the special cases $\mathfrak{cgal}^{\mathrm{Pol}}_2(d) = \mathfrak{sch}_2(d)$, and $\mathfrak{cgal}^{\mathrm{Pol}}_1(d) = \mathfrak{cmil}_1(d)$ corresponding to $N = 1$, and $N = 2$ respectively.

It would be desirable to find a truly geometric definition of such Lie subalgebras of the Lie algebra of conformal Galilei Lie algebras, $\mathfrak{cgal}(M, \gamma, \theta)$, in the case of an arbitrary Galilei (or Newton-Cartan) structure.

5 Conformal Galilean symmetries of physical systems

In order to illustrate our general formalism, we first present a framework, due originally to Souriau [39], which allows us to describe, in particular, both massive and massless Galilean elementary systems in a unified way.

Consider a Hamiltonian system with d degrees of freedom, whose *phase space* is a $2d$-dimensional symplectic manifold $(\mathcal{M}, \Omega)$, and whose Hamiltonian is a smooth function, H, of $\mathcal{M}$. The two-form $\Omega = \frac{1}{2}\Omega_{\alpha\beta}\, dx^\alpha \wedge dx^\beta$ of $\mathcal{M}$ is closed, $d\Omega = 0$, and non-degenerate, $\det(\Omega_{\alpha\beta}) \neq 0$. Using its inverse, $\Omega^{-1} = \frac{1}{2}\Omega^{\alpha\beta}\partial_\alpha \wedge \partial_\beta$, we get the Poisson bracket $\{F, G\} = \Omega^{\alpha\beta}\partial_\alpha F\, \partial_\beta G$ of two observables F, and G. (The Jacobi identity is equivalent to $d\Omega = 0$.) Then Hamilton's equations read

$$\frac{dx^\alpha}{dt} = \{H, x^\alpha\} \tag{5.1}$$

where $\alpha = 1, \ldots, 2d$, the parameter t being interpreted as "time".

If $X_H = \{H, \cdot\}$ is the associated Hamiltonian vector field, we see that (5.1) can also be written as[12]

$$\frac{dx}{dt} = X_H \qquad \text{where} \qquad \Omega(X_H) = -dH \tag{5.2}$$

or, using coordinates, $\Omega_{\alpha\beta}\, X_H^\alpha = -\partial_\beta H$, for all $\beta = 1, \ldots, 2d$.

[11] The definition of these Lie algebras clearly involves constraints given by differential operators of higher order, which go, strictly speaking, beyond our formalism relying essentially on *second order* PDE associated with transport equations of NC-structures.

[12] The form (5.2) of Hamilton's equations allows for a variational interpretation; see, e.g., [39, 44, 45].

One can go one step farther [41] and, promoting time as a new coordinate, consider the $(2d+1)$-dimensional "evolution space" $\mathcal{V} = \mathcal{M} \times \mathbb{R}$ endowed with the following closed two-form which we write, with some abuse of notation, as

$$\sigma = \Omega - dH \wedge dt. \tag{5.3}$$

The vector fields $Y = \lambda(X_H + \partial/\partial t)$ of $\mathcal{V}$, with $\lambda \in \mathbb{R}$, clearly satisfy $\sigma(Y) = 0$ since $X_H(H) = 0$, and $Yt = \lambda$. Denoting $y = (x, t)$ points of $\mathcal{V}$, we see that the equations of motion (5.1) admit the alternative form

$$\frac{dy}{d\tau} = Y \qquad \text{with} \qquad \sigma(Y) = 0 \tag{5.4}$$

where τ is now an arbitrary curve-parameter.

Conversely, let us consider a closed two-form, σ, on some general *evolution space*, $\mathcal{V}$, whose kernel, $K = \ker(\sigma)$, has a (nonzero) constant dimension.[13] Then K (see (5.4)) is an integrable distribution. So, there exists, passing through each point $y \in \mathcal{V}$, a submanifold whose tangent space is spanned by those vectors in K. Each leaf (or characteristic) of K is a *classical motion*. The set of these motions (which is assumed to be a well-behaved manifold) is Souriau's *space of motions*, $\mathcal{U} = \mathcal{V}/K$, of the system. The two-form σ passes to the quotient, $\mathcal{U}$, which becomes, hence, a symplectic manifold. Espousing this point of view, one regards the evolution space as fundamental since it hosts the dynamics in a purely intrinsic way. See also the recent essay [42] supporting this standpoint in the classical and quantum context.

A *symmetry* of the evolution space $(\mathcal{V}, \sigma)$ is given by a vector field Z which Lie-transports the two-form σ, namely such that

$$L_Z \sigma = 0. \tag{5.5}$$

A symmetry is called *Hamiltonian* if there exists a function $\mathcal{J}_Z$ of $\mathcal{V}$ such that, globally,

$$\sigma(Z) = -d\mathcal{J}_Z. \tag{5.6}$$

Then, one readily finds that $Y\mathcal{J}_Z = 0$ for all $Y \in K$, i.e., that $\mathcal{J}_Z$ (determined by (5.6) up to an overall constant) is a conserved quantity. See [39] for an account on this formulation of Noether's theorem.

Conversely, symplectic manifolds upon which a given group of Hamiltonian symmetries acts transitively can be constructed in a systematic fashion [39]. For example, the homogeneous symplectic manifolds of the Galilei group will represent the spaces of motions of classical, non-relativistic, elementary particles. Skipping the details, here we simply list the results which are important for our purposes.

[13] One says that the two-form is *presymplectic*.

5.1 Galilean massive particles

Generic elementary systems of the Galilei group (whose Lie algebra has been defined in (4.15)) in four-dimensional, flat, NC-spacetime are classified by the *mass*, m, and *spin*, s, invariants. In the "massive" case, $m > 0$, the evolution space of a spinning particle, with $s > 0$, is $\mathcal{V} = \mathbb{R} \times \mathbb{R}^3 \times \mathbb{R}^3 \times S^2$ parametrized by the quadruples $y = (t, \mathbf{x}, \boldsymbol{v}, \boldsymbol{u})$, and endowed with the two-form

$$\sigma = m\, dv_A \wedge (dx^A - v^A dt) - \frac{s}{2}\epsilon_{ABC}\, u^A du^B \wedge du^C \tag{5.7}$$

where ϵ_{ABC} is the Levi-Civita symbol with $\epsilon_{123} = 1$.

Equation (5.7) happens to be of the form (5.3) that unifies the symplectic structure of phase space $\mathcal{M} = \mathbb{R}^3 \times \mathbb{R}^3 \times S^2$ and the Hamiltonian, namely

$$\sigma = dp_A \wedge dx^A - \frac{1}{2s^2}\epsilon_{ABC}\, s^A ds^B \wedge ds^C - d\left(\frac{\boldsymbol{p}^2}{2m}\right) \wedge dt \tag{5.8}$$

where the vector $\boldsymbol{p} = m\boldsymbol{v}$ stands for the linear momentum, and $\mathbf{s} = s\boldsymbol{u}$ for the *classical spin*.[14] Ordinary phase space has been extended by the sphere S^2, endowed with its canonical surface element. The two-form (5.7) is closed and has a one-dimensional kernel; the characteristic curves, which are solutions of the free equations of motion (5.4), namely

$$\dot{t} = 1, \qquad \dot{\mathbf{x}} = \boldsymbol{v}, \qquad \dot{\boldsymbol{v}} = 0, \qquad \dot{\boldsymbol{u}} = 0 \tag{5.9}$$

project on spacetime as usual straight worldlines. Those are independent of spin, which is itself a constant of the motion.

These worldlines are, in fact, timelike geodesics since

$$\dot{t} \neq 0. \tag{5.10}$$

As for the symmetries of the model coming from conformal Galilean transformations of flat spacetime, one shows [21, 23] that the only vector fields $X \in \mathfrak{cgal}(3)$ that admit a lift, $\widetilde{X}$, to $(\mathcal{V}, \sigma)$ verifying $L_{\widetilde{X}}\sigma = 0$ (see (5.5)) are necessarily Schrödinger vector fields, $X \in \mathfrak{sch}(3)$. The explicit expression is

$$\begin{aligned} \widetilde{X} \;=\; & \big(\kappa t^2 + 2\lambda t + \varepsilon\big)\frac{\partial}{\partial t} + \big(\omega^A_B\, x^B + \kappa t x^A + \lambda x^A + \beta^A t + \gamma^A\big)\frac{\partial}{\partial x^A} \\ & + \big(\omega^A_B\, v^B + \beta^A - \lambda v^A + \kappa(x^A - v^A t)\big)\frac{\partial}{\partial v^A} \\ & + \omega^A_B\, u^B \frac{\partial}{\partial u^A} \end{aligned} \tag{5.11}$$

with the notation of (4.21).

[14] At the purely classical level studied here, s is an arbitrary positive number; the "prequantizability" [39] requires it to be a half-integral multiple of $\hbar$.

We, hence, recover the results of Section 4.1 dealing with the symmetries of the equations of timelike NC-geodesics. Notice that the Schrödinger symmetry still holds in the presence of spin.

The action (5.11) is Hamiltonian, and the conserved quantities, calculated using Noether's theorem (see Equation (5.6)) read[15]

$$\mathcal{J}_{\widetilde{X}} = \boldsymbol{J}\cdot\boldsymbol{\omega} - \mathbf{G}\cdot\boldsymbol{\beta} + \boldsymbol{P}\cdot\boldsymbol{\gamma} - H\varepsilon - K\kappa + D\lambda \tag{5.12}$$

where

$$\begin{array}{rcll} \boldsymbol{P} & = & \boldsymbol{p} & \text{Linear momentum} \\ \boldsymbol{G} & = & m\boldsymbol{q} & \text{Galilean boost} \\ \boldsymbol{J} & = & \mathbf{x}\times\boldsymbol{p} + s\boldsymbol{u} & \text{Angular momentum} \\ H & = & \dfrac{\boldsymbol{p}^2}{2m} & \text{Energy} \\ K & = & \dfrac{m\boldsymbol{q}^2}{2} & \text{Schrödinger expansions} \\ D & = & \boldsymbol{p}\cdot\boldsymbol{q} & \text{Schrödinger dilations} \end{array} \tag{5.13}$$

together with

$$\boldsymbol{q} = \mathbf{x} - \boldsymbol{v}t. \tag{5.14}$$

Note that the spin enters the angular momentum only and is, in fact, separately conserved. The space of motions, $\mathcal{U} = \mathbb{R}^3 \times \mathbb{R}^3 \times S^2$, therefore inherits from σ the symplectic two-form $\Omega = dp_A \wedge dq^A - (s/2)\epsilon_{ABC}\, u^A du^B \wedge du^C$. The associated Poisson brackets of the components (5.13) of the *moment map* [39] then realize the *one-parameter central extension* of the Schrödinger group, via

$$\{P_A, G_B\} = m\,\delta_{AB}. \tag{5.15}$$

As a further example of massive Schrodinger symmetry, we mention non-relativistic Chern-Simons vortices [53].

So far, we have only studied free particles. Let us mention that the $\mathfrak{so}(2,1)$ symmetry would survive, if $d = 3$, the addition of a Dirac monopole [46, 21, 47], and, if $d = 2$, that of a "magnetic vortex" [48, 49].

5.2 Galilean symmetry in hydrodynamics

Another example with Schrödinger symmetry involves hydrodynamics [50, 51, 52]. To shed a new light on the problem, we present our results in a way complementary to the geometric approach followed in the previous sections.

[15] We have put $\omega_A = -\frac{1}{2}\epsilon_{ABC}\omega^{BC}$ for all $A = 1, 2, 3$.

The equations of motion of an isentropic and dissipationless fluid, in flat $(d+1)$-dimensional non-relativistic spacetime, read [51]

$$\partial_t \rho + \boldsymbol{\nabla} \cdot (\rho\, \boldsymbol{v}) = 0, \tag{5.16}$$

$$\partial_t \boldsymbol{v} + \boldsymbol{v} \cdot \boldsymbol{\nabla} \boldsymbol{v} = -\boldsymbol{\nabla} V'(\rho), \tag{5.17}$$

where $\rho(t, \mathbf{x})$ is the density and $\boldsymbol{v}(t, \mathbf{x})$ the velocity field.

The enthalpy, $V'(\rho)$, is related to the pressure, P, via $\rho V'(\rho) - V(\rho) = P$. For simplicity, we focus our attention to the irrotational case, $\boldsymbol{v} = \boldsymbol{\nabla}\theta$, — where $\theta(t, \mathbf{x})$ is a potential for the velocity field — when the system can be derived from a variational principle using the Lagrangian

$$L = L_0 - V(\rho) = -\rho\big(\partial_t \theta + \frac{1}{2}(\boldsymbol{\nabla}\theta)^2\big) - V(\rho). \tag{5.18}$$

Varying L in (5.18) with respect to θ yields the continuity equation (5.16), and varying it with respect to ρ yields the Bernoulli equation

$$\partial_t \theta + \frac{1}{2}(\boldsymbol{\nabla}\theta)^2 = -V'(\rho) \tag{5.19}$$

whose gradient is the Euler equation (5.17).

The system is plainly Galilei-invariant: a boost implemented by $\theta \mapsto \theta^*$, and $\rho \mapsto \rho^*$, where

$$\theta^*(t, \mathbf{x}) = \theta(t, \mathbf{x} + \mathbf{b}t) - \mathbf{b} \cdot \mathbf{x} - \frac{1}{2}\mathbf{b}^2 t \tag{5.20}$$

$$\rho^*(t, \mathbf{x}) = \rho(t, \mathbf{x} + \mathbf{b}t) \tag{5.21}$$

leaves the Lagrangian (5.18) invariant. Routine calculation proves the invariance against space and time translations, as well as rotations [50, 51], proving the full Galilean invariance of the model.

Now we inquire about the conformal symmetries.

Scale invariance

• Consider a dilation with dynamical exponent, z, namely

$$t^* = \lambda^z t, \qquad \mathbf{x}^* = \lambda\, \mathbf{x} \tag{5.22}$$

and attempt to implement it as

$$\theta^* = \lambda^a \theta(t^*, \mathbf{x}^*), \qquad \rho^* = \lambda^b \rho(t^*, \mathbf{x}^*)$$

where a and b have to be determined.

The two terms in the free Lagrangian, L_0, are seen to scale in the same way when $a = z - 2$, and then the entire expression scales by λ^{b+z-2}. However, the measure of integration scales as $dtd\mathbf{x} = \lambda^{-(d+z)} dt^* d\mathbf{x}^*$, where $d\mathbf{x} = dx^1 \dots dx^d$. Invariance of the free Lagrangian density requires therefore $b = d + (2 - z)$. Thus, for *any dynamical exponent*, z, the free Lagrangian density is scale-invariant, whenever

$$\theta^*(t, \mathbf{x}) = \lambda^{z-2} \theta(t^*, \mathbf{x}^*) \tag{5.23}$$

$$\rho^*(t, \mathbf{x}) = \lambda^{d-z+2} \rho(t^*, \mathbf{x}^*). \tag{5.24}$$

Which potential can be added? Restricting ourselves to the polytropic expression $V(\rho) = c\rho^\gamma$, we find $V(\rho^*) = \lambda^{\gamma(d-z+2)} V(\rho)$, and also $V(\rho^*) = \lambda^{d+z} V(\rho)$ to match the free case. Therefore, to preserve the symmetry with respect to (5.22), the polytropic exponent must be

$$\gamma = \frac{d+z}{d+2-z}\,. \tag{5.25}$$

Conversely, to deal with a potential $V(\rho) = c\rho^\gamma$ having dilations as symmetries requires to choosing the dynamical exponent as

$$z = \frac{\gamma(d+2) - d}{\gamma + 1}\,. \tag{5.26}$$

In other words, the potential breaks to (5.26) the freedom of choosing z.

• For $z = 2$, in particular, when time is twice-dilated with respect to space,

$$t^* = \lambda^2 t, \qquad \mathbf{x}^* = \lambda \mathbf{x} \tag{5.27}$$

we recover the known results [50]

$$\theta^*(t, \mathbf{x}) = \theta(t^*, \mathbf{x}^*) \tag{5.28}$$

$$\rho^*(t, \mathbf{x}) = \lambda^d \rho(t^*, \mathbf{x}^*) \tag{5.29}$$

$$\gamma = 1 + \frac{2}{d}\,. \tag{5.30}$$

Expansions

Schrödinger expansions, viz.,

$$t^* = \frac{t}{1 - \kappa t}, \qquad \mathbf{x}^* = \frac{\mathbf{x}}{1 - \kappa t} \tag{5.31}$$

implemented as

$$\rho^*(t, \mathbf{x}) = (1 - \kappa t)^{-d} \rho(t^*, \mathbf{x}^*) \tag{5.32}$$

$$\theta^*(t, \mathbf{x}) = \theta(t^*, \mathbf{x}^*) - \frac{\kappa\, \mathbf{x}^2}{2(1 - \kappa t)} \tag{5.33}$$

are readily seen to be symmetries for the free fluid system.

Let us attempt to generalize (5.31) as

$$t^* = \Omega\, t, \qquad \mathbf{x}^* = \Omega^\alpha\, \mathbf{x} \tag{5.34}$$

and (5.32), (5.33) as

$$\rho^*(t,\mathbf{x}) = \Omega^\delta \rho(t^*,\mathbf{x}^*) \tag{5.35}$$

$$\theta^*(t,\mathbf{x}) = \theta(t^*,\mathbf{x}^*) - \beta\,\kappa\,\Omega^\gamma\,\mathbf{x}^{*2} \tag{5.36}$$

where $\Omega = (1-\kappa t)^{-1}$, and α, β, γ, δ are to be determined. Then the θ-part of the free Lagrangian transforms according to

$$\begin{aligned} \partial_t\theta^* + \frac{1}{2}(\boldsymbol{\nabla}\theta^*)^2 &= \Omega^2\partial_{t^*}\theta + \frac{1}{2}\Omega^{2\alpha}(\boldsymbol{\nabla}^*\theta)^2 \\ &\quad +\kappa\,\mathbf{x}^*\cdot\boldsymbol{\nabla}^*\theta\left(\alpha\Omega - 2\beta\Omega^{2\alpha+\gamma}\right) \\ &\quad +\beta\kappa^2(\mathbf{x}^*)^2\left(2\beta\Omega^{2\alpha+2\gamma} - (2\alpha+\gamma)\Omega^{\gamma+1}\right). \end{aligned} \tag{5.37}$$

Getting a symmetry requires, therefore, $\alpha = 1, \beta = \frac{1}{2}, \gamma = -1$, and $\delta = d$, leading to the above expressions (5.32), and (5.33). This is, hence, the only case allowed by the expansion-symmetry in fluid mechanics.

On the other hand, dilations and Schrödinger expansions generate, along with time translations, the (neutral component of the) group SO(2, 1), only when the dynamical exponent is $z = 2$. The only consistent way to combine dilations and expansions is, hence, when the system carries a full Schrödinger symmetry [50, 51].

Conserved quantities

Noether's theorem associates conserved quantities to symmetries. In the present field-theoretic context, it goes as follows. Let ϕ be any field. An infinitesimal transformation, $\delta\phi$, is a symmetry if it changes the Lagrange density by a "surface term", $\delta L = \partial_a C^a$, for some quantities C^a. Then

$$J^a = \frac{\delta L}{\delta(\partial_a\phi)}\delta\phi - C^a \tag{5.38}$$

is a conserved current, $\partial_\alpha J^\alpha = 0$, so that the integral

$$Q = \int_{t=t_0} d\mathbf{x}\left(\frac{\delta L}{\delta(\partial_t\phi)}\delta\phi - C^t\right) \tag{5.39}$$

is a constant of the motion, i.e., is independent of t_0.

Returning to the Schrödinger case, and using the Noether theorem one finds the conserved quantities,

$$
\begin{aligned}
\boldsymbol{P} &= \int d\mathbf{x}\,\rho\,\boldsymbol{\nabla}\theta && \text{Linear momentum} \\
\mathbf{G} &= \int d\mathbf{x}\,\rho\big(\mathbf{x}-\boldsymbol{\nabla}\theta\,t\big) && \text{Galilean boosts} \\
\boldsymbol{J} &= \int d\mathbf{x}\,\rho\,\mathbf{x}\wedge\boldsymbol{\nabla}\theta && \text{Angular momentum} \\
H &= \int d\mathbf{x}\left(\frac{1}{2}\rho(\boldsymbol{\nabla}\theta)^2+V(\rho)\right) && \text{Energy} \\
K &= -t^2H+2tD+\frac{1}{2}\int d\mathbf{x}\,\rho\,\mathbf{x}^2 && \text{Schrödinger expansion} \\
D &= tH-\frac{1}{2}\int d\mathbf{x}\,\rho\,(\mathbf{x}\cdot\boldsymbol{\nabla}\theta), && \text{Schrödinger dilation} \\
M &= \int d\mathbf{x}\,\rho && \text{Mass}
\end{aligned}
\tag{5.40}
$$

where we have also added the total Galilean mass. Under (suitably defined) Poisson brackets, we get the generators of the *one-parameter centrally extended Schrödinger algebra* [50].

In conclusion, the free system admits, for any $z \neq 2$, the expansion-less and dilations-only Lie subalgebra of $\mathfrak{sch}_z(d)$ as a Lie algebra of symmetries. It possesses the full Schrödinger Lie algebra of symmetries (including expansions) when $z = 2$.

The symmetry is preserved when the polytropic exponent is chosen suitably, namely as in (5.30).

Accelerations

It is worth mentioning that accelerations,

$$
t^* = t, \qquad \mathbf{x}^* = \mathbf{x} - \frac{1}{2}\mathbf{a}t^2, \tag{5.41}
$$

implemented as

$$
\theta^*(t,\mathbf{x}) = \theta(t^*,\mathbf{x}^*) + (\mathbf{a}\cdot\mathbf{x}^*)t^*, \qquad \rho^*(t,\mathbf{x}) = \rho(t^*,\mathbf{x}^*) \tag{5.42}
$$

change the Lagrangian as,

$$
\rho^*\Big(\partial_t\theta^* + \frac{1}{2}(\boldsymbol{\nabla}\theta^*)^2\Big) = \rho\Big(\partial_{t^*}\theta + \frac{1}{2}(\boldsymbol{\nabla}^*\theta)^2\Big) + \rho\,\big(\mathbf{a}\cdot\mathbf{x}^* - \frac{1}{2}\mathbf{a}^2t^{*2}\big). \tag{5.43}
$$

The extra term, here, is not a total divergence. Accelerations are, therefore, not symmetries for the fluid equations. In fact, they carry the system into an accelerated one [8, 12].

This is consistent with the fact that CGA-type symmetries require masslessness, while fluid mechanics has nonzero mass; see (5.40).

Time-dilations: $z = \infty$

Recall that the free system has actually an $\mathrm{SO}(d+1,2)$ *dynamical relativistic conformal symmetry* — see [50] —, which is broken to its Poincaré subgroup in $d+1$ dimensions in the Chaplygin case [51, 50, 52],

$$V(\rho) = \frac{c}{\rho}. \tag{5.44}$$

The Poincaré group in $d+1$ dimensions contains the one-parameter centrally extended Galilei group in d dimensions, augmented with time-dilations

$$t^* = \lambda t, \qquad \mathbf{x}^* = \mathbf{x} \tag{5.45}$$

as a subgroup.[16] Implemented as

$$\theta^*(t,\mathbf{x}) = \lambda\theta(t^*,\mathbf{x}^*), \tag{5.46}$$

$$\rho^*(t,\mathbf{x}) = \lambda^{-1}\rho(t^*,\mathbf{x}^*) \tag{5.47}$$

they provide a symmetry for the free system: indeed, $L \mapsto \lambda L$ is compensated for by the transformation law $dtd\mathbf{x} = \lambda^{-1}dt^*d\mathbf{x}^*$. Space dilations and expansions are broken.

Moreover, the only potential consistent with (5.45) is (5.44), that of the Chaplygin gas [52, 50, 51].

Time dilations (5.45) act infinitesimally on the fields according to

$$\delta\rho = -\rho + t\,\partial_t\rho, \qquad \delta\theta = \theta + t\,\partial_t\theta. \tag{5.48}$$

We find $\delta L = \partial_t(tL)$, so that the conserved quantity (5.39) associated with (5.45), found by the Noether theorem, is therefore

$$\Delta = tH - \int d\mathbf{x}\,\rho\,\theta \tag{5.49}$$

where H is the energy in (5.40), with $V(\rho)$ as in (5.44). The conservation of (5.49) can also be checked directly, using the equations of motion.

[16] The transformation (5.45) can be viewed as the limiting case, $z \to \infty$, of z-dilation.

5.3 Galilean massless particles

Concerning the second, "Conformal Galilean (CGA)-type" symmetries, the situation is more subtle. The natural candidates are the massless Galilean systems, studied by Souriau forty years ago [39]. In geometrical optics, a classical "light ray" can, in fact, be identified with an *oriented straight line*, $\mathcal{D}$, in Euclidean space $\mathbb{R}^3$. Such a line is characterized by an arbitrary point $\mathbf{x} \in \mathcal{D}$, and its direction, i.e., a unit vector, $\boldsymbol{u}$, along $\mathcal{D}$. The manifold of light rays is readily identified with the (co)tangent bundle TS^2 endowed with its canonical symplectic structure, or a twisted symplectic structure if spin is admitted. This model is based on the Euclidean group.

There exists, indeed, a Galilean version of Euclidean "spinoptics". The homogeneous symplectic manifolds of the Galilei group to consider are "massless", i.e., defined by the invariants $m = 0$, $s \neq 0$, and $k > 0$, a new Galilei-invariant [39, 40].

A natural "evolution space" for these massless models is $\mathcal{V} = \mathbb{R} \times \mathbb{R}^3 \times \mathbb{R} \times S^2$ described by the quadruples $y = (t, \mathbf{x}, E, \boldsymbol{u})$ and endowed with the closed two-form

$$\sigma = k\, du_A \wedge dx^A - dE \wedge dt - \frac{s}{2}\epsilon_{ABC}\, u^A du^B \wedge du^C \tag{5.50}$$

where the constants $k > 0$, and s, are the *color* and the *spin*, respectively.[17]

The motions of these massless particles, e.g., the classical "light rays", identified with the characteristic curves of the two-form (5.50), project onto spacetime as oriented *lightlike* straight worldlines directed along $\boldsymbol{u}$, viz.,

$$\dot{t} = 0, \qquad \dot{\mathbf{x}} = \boldsymbol{u}, \qquad \dot{E} = 0, \qquad \dot{\boldsymbol{u}} = 0 \tag{5.51}$$

where we have chosen the parameter τ as the arc-length along the straight line $\mathcal{D}$.

Such a "motion" is *instantaneous*,

$$t = \text{const.} \tag{5.52}$$

and, hence, projects as a lightlike geodesic (4.3) of flat Newton-Cartan spacetime (massless particles have "infinite speed").

By the very construction of the model, the Galilei Lie algebra, $\mathfrak{gal}(3)$, see (4.15), acts in a Hamiltonian way (5.6) on the evolution space $\mathcal{V}$ according to

$$\widetilde{X} = \varepsilon\frac{\partial}{\partial t} + (\omega^A_B\, x^B + \beta^A t + \gamma^A)\frac{\partial}{\partial x^A} + k u_A \beta^A \frac{\partial}{\partial E} + \omega^A_B\, u^B \frac{\partial}{\partial u^A}, \tag{5.53}$$

where $\boldsymbol{\omega} \in \mathfrak{so}(3)$, $\boldsymbol{\beta}, \boldsymbol{\gamma} \in \mathbb{R}^3$, and $\varepsilon \in \mathbb{R}$.

[17] For $s = 0$ we get a "Fermat particle", i.e., "spinless light," described by the Fermat principle [39, 54]. Quantization requires that $s/\hbar$ be a half-integer, and the color becomes $k = 2\pi\hbar/\lambda$, where λ is the wavelength [39, 40].

Using the definition (5.12) of the Hamiltonian, $\mathcal{J}_{\widetilde{X}}$, of this action, we find the associated conserved quantities, namely

$$\begin{array}{rcll} \boldsymbol{P} & = & k\boldsymbol{u} & \text{Linear momentum} \\ \boldsymbol{G} & = & -\boldsymbol{P}t & \text{Galilean boost} \\ \boldsymbol{J} & = & \mathbf{x}\times\boldsymbol{P}+s\boldsymbol{u} & \text{Angular momentum} \\ H & = & E & \text{Energy} \end{array} \tag{5.54}$$

Let us emphasize that the time-dependent Galilei boost, $\mathbf{G}$, *is* a constant of the "motion", since the latter takes place at constant time, cf. (5.52). Under the Poisson bracket defined by the induced symplectic two-form $\Omega = k\, du_A \wedge dq^A - dE \wedge dt$, where $\boldsymbol{q} = -\boldsymbol{u}\times(\boldsymbol{u}\times\mathbf{x})$, on the space of motions $\mathcal{U} = TS^2 \times T\mathbb{R}$, the components (5.54) of the moment map close into the *centerless* Galilei group. In particular, translations and Galilean boosts commute: our "photon" is massless. Curiously, the energy, E, remains arbitrary, and determined by the initial conditions.

What about our conformal extensions? Are they symmetries? For the *trajectories*, the answer is positive: the lightlike "instantaneous" geodesics are permuted by construction, see Section 4.3.

Concerning the *dynamics*, the answer is more subtle though: for any finite dynamical exponent z, *none* of the additional geometric symmetries leaves the dynamics invariant. Consider, for example, a dilation: while $\mathbf{x} \mapsto e^{\lambda}\mathbf{x}$, the unit vector, $\boldsymbol{u}$, cannot be dilated, $\boldsymbol{u} \mapsto \boldsymbol{u}$. Therefore, a "photon" of color k is carried into one with color $ke^{-\lambda}$ (which, in empty space, follows the same trajectories).

There is, however, a way to escape this obstruction: it is enough ... *not to dilate* $\mathbf{x}$! To see this, consider first the spinless "Fermat" case. Then the evolution space can be viewed as the submanifold $\mathcal{V} \subset T^*M$ of the cotangent bundle of spacetime $M = \mathbb{R}\times\mathbb{R}^3$ defined by the equation

$$\gamma^{ab}p_a p_b - k^2 = 0. \tag{5.55}$$

Its presymplectic two-form, given by Equation (5.50) with $s = 0$, is just $\sigma = d\varpi$, where $\varpi = p_a dx^a$ is the restriction to $\mathcal{V}$ of the canonical one-form of T^*M. Recall that a vector field, X, on space-time is *canonically* lifted to T^*M as

$$\widetilde{X} = X^a \frac{\partial}{\partial x^a} - p_b \frac{\partial X^b}{\partial x^a}\frac{\partial}{\partial p_a}. \tag{5.56}$$

One easily sees that this lift is tangent to the submanifold $\mathcal{V}$, cf. (5.55), iff one has identically $\widetilde{X}(\gamma^{ab}p_a p_b - k^2) = (L_X\gamma)^{ab}p_a p_b = 0$, i.e., iff the vector field X leaves γ invariant, viz.,[18]

$$L_X\gamma = 0 \tag{5.57}$$

[18]This construction is general, and can be extended to the case of any NC-structure [21].

which is the Galilei-conformal condition (3.8) with $m = 1$ and $n = 0$; the dynamical exponent is therefore $z = \infty$.

In the flat case under study, the general solution of Equation (5.57) is given by (3.14), i.e., by

$$X = \xi(t)\frac{\partial}{\partial t} + \left(\omega^A_B(t)x^B + \eta^A(t)\right)\frac{\partial}{\partial x^A} \tag{5.58}$$

where $\boldsymbol{\omega}(t) \in \mathfrak{so}(3)$, $\boldsymbol{\eta}(t)$, and $\xi(t)$ depend arbitrarily on time. At last, the maximal Hamiltonian symmetries of the "Fermat" particle model constitute the Lie algebra $\mathfrak{cgal}_\infty(3)$, i.e., an infinite-dimensional conformal extension of the (centerless) Galilei group.

Let us compute the explicit form of the canonical lift, $\widetilde{X}$, of $X \in \mathfrak{cgal}_\infty(3)$ to $\mathcal{V}$. Using Equation (5.56), we end up with

$$\begin{aligned}\widetilde{X} \;=\; & \xi(t)\frac{\partial}{\partial t} + \left(\omega^A_B(t)x^B + \eta^A(t)\right)\frac{\partial}{\partial x^A} \\ & + \left(k(\omega'_{AB}(t)u^A x^B + \eta'_A(t)u^A) - \xi'(t)E\right)\frac{\partial}{\partial E} \\ & + \omega^A_B(t)u^B\frac{\partial}{\partial u^A}\end{aligned} \tag{5.59}$$

with the same notation as before. As previously mentioned, the $\mathfrak{cgal}_\infty(3)$-action on $\mathcal{V}$ is Hamiltonian if $s = 0$. Using Equation (5.6), one finds the conserved Hamiltonian

$$\mathcal{J}_{\widetilde{X}} = (\mathbf{x} \times k\boldsymbol{u}) \cdot \boldsymbol{\omega}(t) + k\boldsymbol{u} \cdot \boldsymbol{\eta}(t) - \xi(t)E. \tag{5.60}$$

What about spin? One easily checks that, in the case $s \neq 0$, the presymplectic two-form (5.50) is no longer $\mathfrak{cgal}_\infty(3)$-invariant. In fact, elementary calculation shows that any $X \in \mathfrak{cgal}_\infty(3)$ such that $L_{\widetilde{X}}\sigma = 0$ is of the form (5.58) with

$$\boldsymbol{\omega}'(t) = 0. \tag{5.61}$$

Thus, in the general case of massless, spinning Galilean particles, the associated constants of the "motion" retain the final form

$$\mathcal{J}_{\widetilde{X}} = (\mathbf{x} \times k\boldsymbol{u} + s\boldsymbol{u}) \cdot \boldsymbol{\omega} + k\boldsymbol{u} \cdot \boldsymbol{\eta}(t) - \xi(t)E \tag{5.62}$$

with $\boldsymbol{\omega} \in \mathfrak{so}(3)$, $\boldsymbol{\eta}(t)$, and $\xi(t)$ remaining arbitrary functions of time. Note that the conservation of these quantities is related to the fact that the "motions" are instantaneous (5.52).

In conclusion, Souriau's "classical photon" admits an infinite-dimensional conformal extension of the (centerless) Galilei group; see (5.54).

Let us mention, for completeness, another type of massless Galilean particle, introduced by Stichel and Zakrzewski [5]. It is described by an extended phase space and, unlike Souriau's photon, has *finite* velocity. It realizes dynamically the Conformal Galilean (CG) symmetry.

5.4 Galilean Electromagnetism

Le Bellac and Lévy-Leblond (LBLL) [56] have discovered, in the early seventies, a full-fledged theory of non-relativistic electromagnetism. They have, actually, highlighted the existence of *two* quite distinct Galilean electromagnetisms, namely a *magnetic-like* and an *electric-like* theory that stem from different non-relativistic limits of Maxwell's theory. The LBLL theories have been, since then, cast into the geometric structure of NC-spacetime [35]. They have, likewise, been formulated in the "null Kaluza-Klein" (or Bargmann) framework of non-relativistic spacetime [27].

Let us, here, confine considerations to the magnetic-like LBLL theory along the lines of [35]. Given a (d+1)-dimensional NC-spacetime structure $(M, \gamma, \theta, \Gamma)$, it is described by the following couple of PDE, namely

$$d\mathcal{F} = 0 \tag{5.63}$$

$$\mathrm{div}\mathcal{F} = J \tag{5.64}$$

involving a two-form, $\mathcal{F} = \frac{1}{2}\mathcal{F}_{ab}\, dx^a \wedge dx^b$, of M interpreted as the electromagnetic field, and a one-form, J, the current density of the sources. In Equation (5.64), one must read

$$\mathrm{div}\mathcal{F}_c = \gamma^{ab}\nabla_a\mathcal{F}_{bc} \tag{5.65}$$

for all $c = 0, \ldots, d$.

Note that $d = 3$ in the original formulation of LBLL theory where equations (5.63), and (5.64) retain the form

$$\boldsymbol{\nabla} \cdot \mathbf{B} = 0, \qquad \boldsymbol{\nabla} \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$$

and

$$\boldsymbol{\nabla} \cdot \mathbf{E} = \varrho, \qquad \boldsymbol{\nabla} \times \mathbf{B} = \mathbf{j}$$

respectively, once we posit $E_A = \mathcal{F}_{A0}$, and $B^A = \frac{1}{2}\epsilon^{ABC}\mathcal{F}_{BC}$, for the components of the electromagnetic field, as well as $\varrho = J_0$, and $j_A = J_A$ for the those of the current density, with $A = 1, 2, 3$.

Notice the absence of the displacement current in Ampère's law: its presence would, clearly, break the Galilean symmetry (Maxwell's equations are relativistic).

Let us show that, much in the same way as Maxwell's sourcefree electromagnetism, the maximal symmetries of the sourcefree LBLL magnetic theory are actually richer than those expected from the original spacetime structure. More specifically, let us look at all conformal Galilei transformations that preserve the LBLL Equations (5.63) and (5.64), with $J = 0$. We will thus seek the maximal Lie

algebra of Galilei conformal vector fields X of (M, γ, θ), i.e., satisfying (3.1), and such that

$$L_X d\mathcal{F} = dL_X\mathcal{F} \tag{5.66}$$

$$L_X \mathrm{div}\mathcal{F} = \mathrm{div} L_X\mathcal{F} \tag{5.67}$$

for all solutions, $\mathcal{F}$, of the above sourcefree LBLL equations.

Equation (5.66) is trivially satisfied (as a consequence of the general fact that the Lie and exterior derivatives commute). As to Equation (5.67), one finds

$$\begin{aligned} 0 &= L_X\gamma^{ab}\nabla_a\mathcal{F}_{bc} - \gamma^{ab}\nabla_a L_X\mathcal{F}_{bc} \\ &= (L_X\gamma)^{ab}\nabla_a\mathcal{F}_{bc} - \gamma^{ab}((L_X\Gamma)^k_{ab}\mathcal{F}_{kc} + (L_X\Gamma)^k_{ac}\mathcal{F}_{bk}) \\ &= f\gamma^{ab}\nabla_a\mathcal{F}_{bc} + 2\gamma^{ab}(L_X\Gamma)^k_{a[b}\mathcal{F}_{c]k} \\ &= -2\gamma^{ab}(L_X\Gamma)^k_{a[b}\delta^\ell_{c]}\mathcal{F}_{k\ell} \end{aligned} \tag{5.68}$$

since $L_X\gamma^{ab} = f\gamma^{ab}$, and $\gamma^{ab}\nabla_a\mathcal{F}_{bc} = 0$. This readily entails

$$\gamma^{ab}(L_X\Gamma)^{[k}_{a[b}\delta^{\ell]}_{c]} = 0 \tag{5.69}$$

for all $c, k, \ell = 0, \ldots, d$.

Utilizing Equation (4.28), giving the most general form of the variations of the NC-connection compatible with Galilei conformal rescalings, we will now put $\delta\Gamma^c_{ab} = L_X\Gamma^c_{ab}$, and easily show that Equation (5.69) writes now

$$\left((4-d)\delta^{[k}_c - 2\theta_c U^{[k}\right)\gamma^{\ell]a}\partial_a f + (f+g)\theta_c F_{ab}\gamma^{ka}\gamma^{\ell b} = 0. \tag{5.70}$$

Taking traces, we readily deduce that $\gamma^{\ell a}\partial_a f = 0$, hence that Equation (5.70) reads

$$(f+g)F_{ab}\gamma^{ka}\gamma^{\ell b} = 0. \tag{5.71}$$

On the one hand, we can have the case $f + g = 0$ (with $z = 2$) leading us to the Schrödinger Lie algebra, $\mathfrak{sch}(M, \gamma, \theta, \Gamma)$, for a general NC-structure.

On the other hand, considering a NC-Milne structure $(M, \gamma, \theta, {}^U\Gamma)$, characterized by $F = 0$, already enables us to satisfy Equation (5.71), hence the full system (4.47)–(4.49) providing us with a higher-dimensional symmetry algebra.

We have just proved that the maximal Lie algebra of symmetries of the Le Bellac-Lévy-Leblond equations in vacuum is isomorphic to $\mathfrak{cmil}(M, \gamma, \theta, {}^U\Gamma)$. In the flat case this is the Lie algebra $\mathfrak{cmil}(d)$ — see (4.65) —, containing the CGA (1.3).

6 Conclusion

In this paper, we have presented a systematic way to derive all types of "non-relativistic conformal transformations" of spacetime M. Due to the degeneracy of the Galilei "metric" (γ, θ), and to the relative independence of Newton-Cartan connections, Γ, there are quite a large number of candidates.

Firstly, the conformal transformations of the "metric" structure alone, (3.1), yield the conformal Galilei Lie algebra $\mathfrak{cgal}(M, \gamma, \theta)$. In the flat case, and in d space dimensions, it is the infinite-dimensional Lie algebra (3.5). Fixing the dynamical exponent, z, via the geometric definition (3.8), yields a family of (still infinite-dimensional) Lie subalgebras, $\mathfrak{cgal}_z(d)$. For $z = 2$, we get the Schrödinger-Virasoro Lie algebra (3.11) of Henkel et al. [29, 2, 3], and for $z = \infty$ we get $\mathfrak{cgal}_\infty(d)$ in (3.14).

Secondly, the Newton-Cartan structure also involves the choice of a connection, which allows us to consider the symmetries of the equations of geodesics, identified with worldlines of test particles.

- The conformal Galilei transformations which permute geodesics, which are generically *timelike*, are, in fact, Schrödinger transformations. Those with dynamical exponent $z = 2$ constitute the Schrödinger Lie algebra, $\mathfrak{sch}(d)$, see (4.21), in the case of flat NC-spacetime.

- Those which exchange *lightlike* geodesics have a richer structure, though. The resulting infinite-dimensional algebra, $\mathfrak{cnc}(d)$, is given by (4.45) in the flat case. There is no *a priori* restriction on the dynamical exponent, z. The infinite-dimensional Lie algebra $\mathfrak{cnc}(d)$ admits a finite-dimensional Lie subalgebra, $\mathfrak{cmil}(d)$ — related to a flat *NC-Milne structure* — featuring independent space and time dilations, as well as new "acceleration" generators; see (4.65). The Lie subalgebras associated with a dynamical exponent, z, are respectively (i) the Schrödinger algebra, for $z = 2$, and (ii) the CGA (1.3) of Henkel [29] and Lukierski et al. [1], for $z = 1$. The infinite-dimensional Lie algebra $\mathfrak{cnc}_\infty(d)$ completes our classification.

Our geometric-algebraic framework, dealing with vector fields on NC-spacetime, leaves no place to central extensions; the latter only arise when conserved quantities — or (pre)quantization — of concrete physical systems are considered.

All these symmetries were derived by considering as fundamental the NC-structure of non-relativistic spacetime. What about concrete physical systems? In Section 5 we study two such systems. Souriau's (pre)symplectic framework [39] allows us to present them in a unified manner.

The dynamics of physical systems reduce further the geometric symmetries to some of their subgroups. This is understood if we think of free fall: massive particles fall in the same way, independently of their respective masses. The trajectory of a particle can be carried therefore into another one by a geometric transformation.

However, such a transformation can change the mass — so it is *not* necessarily a symmetry of the dynamical system.

Firstly, for a massive Galilean particle with spin, we recover the well-known Schrödinger symmetry associated with $\mathfrak{sch}(3)$.

Secondly, the hydrodynamics of irrotational fluids turns out to provide a special instance of a classical field theory invariant under the Schrödinger Lie algebra, $\mathfrak{sch}(d)$, yielding new conserved quantities, apart from the standard Galilean constants of the motion.

As to the GCA-type symmetries, they require to have no mass [1]. The natural candidates are, therefore, Souriau's "Galilean photons" [39], which are associated with the coadjoint orbits of the (centerless) Galilei group. These "particles" have an "instantaneous motion" — they have "infinite velocity". They can carry spin, generalizing "spinless light", described by the Fermat principle [39, 54].

Can one take such models seriously? The answer is positive since they can be obtained as suitable non-relativistic limits of relativistic massless particles, i.e., those associated with the mass zero coadjoint orbits of the Poincaré group [39]. Even more importantly, the model of "Galilean photon" ($s = \pm\hbar$) is a trivial extension of the Euclidean model presented in [54], which has been used to explain the recently observed spin-Hall effect for light [55].

Let us emphasize that Souriau's model of massless Galilean particles carry an infinite-dimensional Lie algebra of symmetries, namely $\mathfrak{cgal}_\infty(3)$.

As a provisionally last illustration of our formalism, we show that the maximal symmetry Lie algebra of the Le Bellac-Lévy-Leblond equations of (magnetic-like) Galilean electromagnetism in vacuum turns out to be the conformal NC-Milne algebra, i.e., $\mathfrak{cmil}(d)$, in flat spacetime, with the CGA as a Lie subalgebra.

Let us also refer to [38] where the space of periodic time-dependent Schrödinger operators, in the case $d = 1$, has been shown to be naturally $\mathfrak{sv}(1)$-invariant.

Recently, a supersymmetric extension of the CGA has been found [57].

Let us end our paper with some historical remarks, cf. [58].

The first person to consider the CGA seems to be Barut [59], in 1973, who derived it by contraction from the relativistic conformal group. But then he discarded it, however, arguing that it is not a symmetry of the Schrödinger equation. In 1978, Havas and Plebański generalized both the Schrödinger and CG groups to an infinite-dimensional group [60].

Even more astonishingly, the Schrödinger symmetry has already be known to Jacobi [60]. In his 1842/43 lectures delivered at the University of Königsberg [61], he studied indeed the dynamics of a particle in a homogeneous potential, U, of degree k. Using a scaling argument reminiscent of the proof of the virial theorem

(see, e.g., [27]), he proved that

$$\frac{d^2}{dt^2}\Big(\frac{m\mathbf{x}^2}{2}\Big) = -(k+2)U + 2E \tag{6.1}$$

where E is, in modern terms, the conserved energy. Then he observed that for the *inverse-square potential*, $k = -2$, Equation (6.1) can be rewritten as

$$\frac{d}{dt}\,(m\mathbf{x}\cdot\dot{\mathbf{x}} - 2Et) = 0.$$

Putting $\boldsymbol{p} = m\dot{\mathbf{x}}$, the quantity

$$D = \boldsymbol{p}\cdot\mathbf{x} - 2Et \tag{6.2}$$

is therefore conserved. Then

$$\frac{d}{dt}\left(\frac{m\mathbf{x}^2}{2} - tD - Et^2\right) = 0$$

so that

$$K = \frac{m\mathbf{x}^2}{2} - tD - Et^2 \tag{6.3}$$

is also conserved. But E, D, K are precisely the conserved quantities which stem, through the Noether theorem, from the conformal, $\mathrm{O}(2,1)$, subgroup of the Schrödinger group, cf. (5.13).

Acknowledgement. We are greatly indebted to M. Henkel, J. Lukierski, P. Stichel, and S. Rouhani for useful correspondence.

References

[1] J. Lukierski, P. C. Stichel, and W. J. Zakrzewski, "Exotic Galilean conformal symmetry and its dynamical realizations," Phys. Lett. A **357** (2006), 1. [arXiv:hep-th/0511259].
J. Lukierski, P. C. Stichel, and W. J. Zakrzewski, "Acceleration-Extended Galilean Symmetries with Central Charges and their Dynamical Realizations," Phys. Lett. B **650** (2007), 203. [arXiv:hep-th/0702179].
J. Lukierski, P. C. Stichel, and W. J. Zakrzewski, "Acceleration-Enlarged Symmetries in Nonrelativistic Space-Time with a Cosmological Constant," arXiv:0710.3093 [hep-th].
J. Gomis, and J. Lukierski, "Enlarged NH symmetries: particle dynamics and gauge symmetries," Phys. Lett. **B 664** (2008), 107.

[2] M. Henkel, "Phenomenology of local scale invariance: from conformal invariance to dynamical scaling," Nucl. Phys. B **641** (2002), 405.
M. Henkel and J. Unterberger, "Schroedinger invariance and space-time symmetries," Nucl. Phys. B **660** (2003) 407. [arXiv:hep-th/0302187].

[3] M. Henkel, R. Schott, S. Stoimenov, and J. Unterberger, "The Poincaré algebra in the context of ageing systems: Lie structure, representations, Appell systems and coherent states," arXiv:math-ph/0601028. See also in *Quantum probability and infinite-dimensional analysis*, L. Accardi, W. Freudenberg and M. Schürmann (eds), World Scientific (Singapour 2007), pp. 233–240.

[4] K. Balasubramanian, and J. McGreevy, "Gravity duals for non-relativistic CFTs," Phys. Rev. Lett. **101** (2008), 061601.
D. T. Son, "Toward and AdS/cold atom correspondence: a geometric realization of the Schrödinger symmetry," Phys. Rev. **D78** (2008), 046003.
The number of papers published on the subject is rapidly growing. It is, therefore, impossible to give full credit to all important contributions. For a relation to earlier work and an (incomplete) list of references, see, e.g., C. Duval, M. Hassaïne, and P. A. Horváthy, "The geometry of Schrödinger symmetry in non-relativistic CFT," arXiv:0809.3128 [hep-th]. Ann. Phys. (N.Y.) **324** (2009), 1158.

[5] P. C. Stichel, and W. J. Zakrzewski, "Nonrelativistic Dark-Energy Fluid in a Baby Universe," arXiv:0807.3660 [gr-qc].
P. C. Stichel, and W. J. Zakrzewski, "Can cosmic acceleration be caused by exotic massless particles?" arXiv:0904.1375v1 [astro-ph.CO]

[6] A. V. Galajinsky, "Remark on quantum mechanics with conformal Galilean symmetry," Phys. Rev. D **78** (2008), 087701. arXiv:0808.1553 [hep-th].

[7] J. Negro, M. A. del Olmo, An A. Rodríguez-Marco, "Nonrelativistic conformal groups," J. Math. Phys. **38** (1997), 3786, and *ibid.* 3810.

[8] I. Fouxon, and Y. Oz, "CFT Hydrodynamics: symmetries, exact solutions and gravity," arXiv: 0812.1266 [hep-th].

[9] A. Bagchi, and R. Gopakumar, "Galilean Conformal Algebras and AdS/CFT," arXiv:0902.1385 [hep-th].

[10] M. Alishahiha, A. Davody, and A. Vahedi, "On AdS/CFT of Galilean Conformal Field Theories," arXiv:0903.3953 [hep-th].

[11] D. Martelli, and Y. Tachikawa, "Comments on Galilean conformal field theories and their geometric realization," arXiv:0903.5184 [hep-th].

[12] P. A. Horváthy, and P. M. Zhang, "Non-relativistic conformal symmetries in fluid mechanics," arXiv:0906.3594 [physics.flu-dyn].

[13] J. Gomis, and J. Lukierski, "Enlarged NH symmetries: particle dynamics and gauge symmetries," Phys. Lett. **B 664** (2008), 107.

[14] R. Jackiw, "Introducing scaling symmetry," Phys. Today, **25** (1972), 23.
C. R. Hagen, "Scale and conformal transformations in Galilean-covariant field theory," Phys. Rev. **D5** (1972), 377.
U. Niederer, "The maximal kinematical symmetry group of the free Schrödinger equation," Helv. Phys. Acta **45** (1972), 802.

[15] G. Burdet, M. Perrin, and P. Sorba, "About the Non-Relativistic Structure of the Conformal Algebra," Commun. Math. Phys. **34** (1973), 85.

[16] C. Duval, and H.P. Künzle, "Minimal Gravitational Coupling in the Newtonian Theory and the Covariant Schrödinger Equation," Gen. Rel. Grav. **16** (1984), 333.

[17] J.-M. Lévy-Leblond, "Galilei group and galilean invariance," in *Group Theory and Applications* (Loebl Ed.), **II**, Acad. Press, New York, p. 222 (1972).
A. Ballesteros, M. Gadella, and M. del Olmo, "Moyal quantization of 2+1 dimensional Galilean systems," Journ. Math. Phys. **33** (1992), 3379.
Y. Brihaye, C. Gonera, S. Giller, and P. Kosiński, "Galilean invariance in $2+1$ dimensions," [hep-th/9503046] (unpublished).
D. R. Grigore, "Transitive symplectic manifolds in $1+2$ dimensions," Journ. Math. Phys. **37** (1996), 240.
D. R. Grigore, "The projective unitary irreducible representations of the Galilei group in $1+2$ dimensions," Journ. Math. Phys. **37** (1996), 460.

[18] J. Lukierski, P. C. Stichel, and W. J. Zakrzewski, "Galilean-invariant $(2+1)$-dimensional models with a Chern-Simons-like term and $d=2$ noncommutative geometry," Annals of Physics (N. Y.) **260** (1997), 224.

[19] C. Duval, and P. A. Horváthy, "The "Peierls substitution" and the exotic Galilei group," Phys. Lett. **B 479** (2000), 284. [hep-th/0002233].
C. Duval, and P. A. Horváthy, "Exotic galilean symmetry in the non-commutative plane, and the Hall effect," Journ. Phys. **A 34** (2001), 10097. [hep-th/0106089].

[20] P. A. Horváthy, L. Martina, and P. Stichel: "Galilean symmetry in noncommutative field theory," Phys. Lett. **B 564** (2003), 149. [hep-th/0304215].
P. A. Horváthy, and M. S. Plyushchay: "Anyon wave equations and the noncommutative plane," Phys. Lett. **B 595** (2004), 547. [hep-th/0404137].

[21] C. Duval, "Quelques procédures géométriques en dynamique des particules," Thèse d'Etat (1982), unpublished.

[22] G. Burdet, C. Duval, and M. Perrin, "Cartan Structures on Galilean Manifolds: the Chronoprojective Geometry," J. Math. Phys. **24** (1983), 1752.
G. Burdet, C. Duval, and M. Perrin, "Chronoprojective Cartan Structures on Four-Dimensional Manifolds," Publ. RIMS, Kyoto Univ. **19** (1983), 813.

[23] C. Duval, "Nonrelativistic Conformal Symmetries and Bargmann Structures," in *Conformal Groups and Related Symmetries. Physical Results and Mathematical Background*, Clausthal 1985, (A.O. Barut & H.D. Doebner Eds), pp 162–182, Lecture Notes in Physics **261**, Springer-Verlag (1986).

[24] B. Carter, and I.M. Khalatnikov, "Canonically covariant formulation of Landau's Newtonian superfluid dynamics," Rev. Math. Phys. **6** (1994), 277.

[25] C. Duval, "Galilean isometries," Class. Quantum Grav. **10** (1993), 2217. arXiv:0903.1641.

[26] C. Duval, G. Burdet, H.P. Künzle, and M. Perrin, "Bargmann Structures And Newton-Cartan Theory," Phys. Rev. **D31**, 1841 (1985).

[27] C. Duval, G. W. Gibbons, and P. A. Horváthy, "Celestial Mechanics, Conformal Structures and Gravitational Waves," Phys. Rev. **D43**, 3907 (1991).

[28] J. Gomis, and J.M. Pons, "Poincaré Transformations and Galilei Transformations," Phys.Lett. **A66** (1978), 463.
J. Gomis, A. Poch, and J. M. Pons, "Poincaré wave equations as Fourier transforms of Galilei wave equations," J. Math. Phys. **21** (1980), 2682.

[29] M. Henkel, "Schrödinger invariance and strongly anisotropic critical systems," J. Stat. Phys. **75** (1994), 1023.
M. Henkel, "Local Scale Invariance and Strongly Anisotropic Equilibrium Critical Systems," Phys Rev. Lett. **78** (1997), 1940.

[30] E. Cartan, "Sur les variétés à connexion affine et la théorie de la relativité généralisée," Ann. Sci. Ecole Norm. Sup. (4) **40** (1923), 325.

[31] A. Trautman, "Sur la théorie newtonienne de la gravitation," C.R. Acad. Sci. Paris **257** (1963), 617; "Comparison of Newtonian and relativistic theories of space time," pp. 413–425 in *Perspectives in Geometry and Relativity*, (B. Hoffmann, Ed.), Indiana University Press, Bloomington, 1964.

[32] H.P. Künzle, "Galilei and Lorentz structures on spacetime: Comparison of the corresponding geometry and physics," Ann. Inst. H. Poincaré, Phys. Théor **17** (1972), 337, and references therein.

[33] C. Duval, and H.P. Künzle, "Sur les connexions newtoniennes et l'extension non triviale du groupe de Galilée," C.R. Acad. Sci. Paris **285 A** (1977), 813

[34] C. Duval, and H.P. Künzle, "Dynamics of continua and particles from general covariance of Newtonian gravitation theory," Rep. Math. Phys. **13** (1978), 351.

[35] H.P. Künzle, "Covariant Newtonian limit of Lorentz space-times," General Relativity and Gravitation **7** (1976), 445.

[36] F. L. Lin, and S. Y. Wu, "Non-relativistic Holography and Singular Black Hole," arXiv:0810.0227 [hep-th].

[37] C. Roger, and J. Unterberger, "The Schrödinger-Virasoro Lie group and algebra: from geometry to representation theory," Ann. Inst. H. Poincaré **7** (2006), 1477.

[38] C. Roger, and J. Unterberger, "A Hamiltonian action of the Schrödinger-Virasoro algebra on a space of periodic time-dependent Schrödinger operators in $(1+1)$-dimensions," arXiv:0810.0902v1 [math-ph].
J. Unterberger, "A classification of periodic time-dependent generalized harmonic oscillators using a Hamiltonian action of the Schrödinger-Virasoro group," arXiv:0806.1185v2 [math-ph].

[39] J.-M. Souriau, *Structure des systèmes dynamiques*, Dunod (1970, ©1969); *Structure of Dynamical Systems. A Symplectic View of Physics*, translated by C.H. Cushman-de Vries (R.H. Cushman and G.M. Tuynman, Translation Editors), Birkhäuser, 1997.

[40] V. Guillemin, and S. Sternberg, *Symplectic techniques in physics*, Cambridge University Press, Cambridge (1984).

[41] R. Abraham, and J.E. Marsden, *Foundations of Mechanics*, Second Edition, Addison-Wesley Publishing Company, Inc (1987).

[42] C. Rovelli, "Forget time", Essay written for the FQXi contest on the Nature of Time, arXiv:0903.3832v3 [gr-qc].

[43] M. Trümper, "Lagrangian Mechanics and the Geometry of Configuration Spacetime," Ann. Phys. (N.Y.) **149** (1983), 203.

[44] P. A. Horváthy, and L. Úry, "Analogy between statics and dynamics – related to variational mechanics," Acta Phys. Hung. **42** (1977), 251.
P. A. Horváthy, "Variational formalism for spinning particles," Journ. Math. Phys. **20** (1979), 49.
P. A. Horváthy, "The non-commutative Landau problem," Ann. Phys. (N.Y.) **299** (2002), 128. [hep-th/0201007].

[45] G. Tuynman, "Un principe variationnel pour les variétés symplectiques," C. R. Acad. Sci. Paris Sér. I Math. **326**:3 (1998), 339.

[46] R. Jackiw, "Dynamical symmetry of the magnetic monopole," Ann. Phys. (N.Y.) **129**, 183 (1980).

[47] P. A. Horváthy, "The dynamical symmetries of the magnetic monopole in geometric quantization," Lett. Math. Phys. **7**, 353 (1983).

[48] R. Jackiw, "Dynamical symmetry of the magnetic vortex," Ann. Phys. (N.Y.) **201**, 83 (1990).

[49] C. Duval and P. A. Horváthy, "Supersymmetry of the magnetic vortex," Tours preprint No-60-93 (unpublished). Available as arXiv:0807.0569 [hep-th].

[50] M. Hassaïne, and P. A. Horváthy, "Field-dependent symmetries of a non-relativistic fluid model," Annals Phys. **282** (2000), 218. [arXiv:math-ph/9904022].
M. Hassaïne, and P. A. Horváthy, "Symmetries of fluid dynamics with polytropic exponent," Phys. Lett. A **279** (2001), 215. [arXiv:hep-th/0009092].
L. O'Raifeartaigh, and V. V. Sreedhar, "The maximal kinematical invariance group of fluid dynamics and explosion-implosion duality," Annals Phys. **293** (2001), 215. [arXiv:hep-th/0007199].

[51] R. Jackiw, *Lectures on fluid dynamics*. CRM series in mathematical Physics, Springer (2002).

[52] D. Bazeia and R. Jackiw, "Nonlinear realization of a dynamical Poincare symmetry by a field-dependent diffeomorphism," Annals Phys. **270** (1998) 246.

[53] R. Jackiw, and S-Y. Pi, "Classical and quantum nonrelativistic Chern-Simons Theory," Phys. Rev. **D42** (1990), 3500.
C. Duval, P. Horváthy, and L. Palla: "Conformal symmetry of the coupled Chern-Simons and gauged non-linear Schrödinger equations," Phys. Lett. **B325** (1994), 39. [hep-th/9401065].
C. Duval, P. Horváthy, and L. Palla, "Spinors in non-relativistic Chern-Simons electromagnetism," Ann. Phys. (N. Y.) **249** (1996), 265. [hep-th/9510114].
B. Chakraborty, and A. J. Majumdar, "On Galilean covariance in a nonrelativistic model involving a Chern-Simons term," Ann. Phys. (N.Y.) **250** (1996), 112.
R. Banerjee, and P. Mukherjee, "Galilean symmetry in a nonabelian Chern-Simons matter system," Ann. Phys. (N.Y.) **264** (1998), 30.

[54] C. Duval, Z. Horvath, and P. Horváthy, "Geometrical Spinoptics and the Optical Hall Effect," J. Geom. Phys. **57** (2007), 925. [arXiv:math-ph/0509031].
C. Duval, Z. Horvath, and P. A. Horváthy, "Fermat principle for spinning light," Phys. Rev. D **74** (2006), 021701. [arXiv:cond-mat/0509636].

[55] K. Yu. Bliokh, and Yu. P. Bliokh, "Topological spin transport of photons: the optical Magnus effect and Berry Phase," *Phys. Lett.* **A333** (2004), 181. [physics/0402110].
M. Onoda, S. Murakami, and N. Nagaosa, "Hall effect for light," *Phys. Rev. Lett.* **93** (2004), 083901. [cond-mat/0405129].
K. Y. Bliokh, A. Niv, V. Kleiner, and E. Hasman, "Geometrodynamics of Spinning Light," arXiv:0810.2136 [physics.optics]. Nature Photon. **2** (2008), 748.
K. Y. Bliokh, "Geometrodynamics of polarized light: Berry phase and spin Hall effect in a gradient-index medium," arXiv:0903.1910 [physics.optics].

[56] M. Le Bellac, and J.-M. Lévy-Leblond, "Galilean Electromagnetism," Nuovo Cimento **14** B (1973), 217.

[57] J. A. de Azcarraga, and J. Lukierski, "Galilean Superconformal Symmetries," arXiv:0905.0141 [math-ph].
M. Sakaguchi, "Super Galilean conformal algebra in AdS/CFT," arXiv:0905.0188 [hep-th].
A. Bagchi, and I. Mandal, "Supersymmetric Extension of Galilean Conformal Algebras," arXiv:0905.0580 [hep-th].

[58] A. Hosseiny, and S. Rouhani, "Affine Extension of Galilean Conformal Algebra in 2+1 Dimensions," arXiv:0909.1203 [hep-th].

[59] A. O. Barut, "Conformal Group → Schrödinger Group → Dynamical Group — The Maximal Kinematical Group of the Massive Schrödinger Particle," *Helv. Phys. Acta* **46**, 496 (1973).
U. Niederer, "The connections between the Schrödinger group and the conformal group," *Helv. Phys. Acta* **47**, 119 (1974).

[60] P. Havas, and J. Plebański, "Conformal extensions of the Galilei group and their relation to the Schrödinger group," *J. Math. Phys.* **19**, 482 (1978).

[61] C. G. J. Jacobi, "Vorlesungen über Dynamik." Univ. Königsberg 1842-43. Herausg. A. Clebsch. Vierte Vorlesung: Das Princip der Erhaltung der lebendigen Kraft. Zweite ausg. C. G. J. Jacobi's Gesammelte Werke. Supplementband. Herausg. E. Lottner. Berlin Reimer (1884).