



HAL
open science

Elliptic Integrable Systems: a Comprehensive Geometric Interpretation

Idrisse Khemar

► **To cite this version:**

| Idrisse Khemar. Elliptic Integrable Systems: a Comprehensive Geometric Interpretation. 2009. hal-00374546v5

HAL Id: hal-00374546

<https://hal.science/hal-00374546v5>

Preprint submitted on 24 Mar 2010 (v5), last revised 15 Apr 2011 (v7)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Elliptic Integrable Systems: a Comprehensive Geometric Interpretation

Idrisse Khemar

Abstract.

In this paper, we study all the elliptic integrable systems, in the sense of C.L. Terng [47]. That is to say the family of all the m -th elliptic integrable systems associated to a k' -symmetric space $N = G/G_0$. Here $m \in \mathbb{N}$ and $k' \in \mathbb{N}^*$ are integers. For example, it is known that the first elliptic integrable system associated to a symmetric space (resp. to a Lie group) is the equation for harmonic maps into this symmetric space (resp. this Lie group). Indeed it is well known that this harmonic maps equation can be written as a zero curvature equation: $d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0$, $\forall \lambda \in \mathbb{C}^*$, where $\alpha_\lambda = \lambda^{-1}\alpha'_1 + \alpha_0 + \lambda\alpha''_1$ is a 1-form on a Riemann surface L taking values in the Lie algebra \mathfrak{g} . This 1-form α_λ is obtained as follows. Let $f: L \rightarrow N = G/G_0$ be a map from the Riemann surface L into the symmetric space G/G_0 . Then let $F: L \rightarrow G$ be a lift of f , and consider $\alpha = F^{-1}.dF$ its Maurer-Cartan form. Then decompose α according to the symmetric decomposition $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ of \mathfrak{g} : $\alpha = \alpha_0 + \alpha_1$. Finally, we define $\alpha_\lambda := \lambda^{-1}\alpha'_1 + \alpha_0 + \lambda\alpha''_1$, $\forall \lambda \in \mathbb{C}^*$. Then the zero curvature equation for this α_λ , for all $\lambda \in \mathbb{C}^*$, is equivalent to the harmonic maps equation for $f: L \rightarrow N = G/G_0$, and is by definition the first elliptic integrable system associated to the symmetric space G/G_0 . Thus the methods of integrable system theory apply to give generalised Weierstrass representations, algebro-geometric solutions, spectral deformations and so on. In particular, we can apply the DPW method [16] to obtain a generalised Weierstrass representation. More precisely, we have a Maurer-Cartan equation in some loop Lie algebra $\Lambda\mathfrak{g}_\tau = \{\xi: S^1 \rightarrow \mathfrak{g} | \xi(-\lambda) = \tau(\xi(\lambda))\}$, then we can integrate it in the corresponding loop group and finally apply some factorizations theorems in loop groups to obtain a generalised Weierstrass representation: this is the DPW method. Moreover, these methods of integrable system theory hold for all the systems written in the forms of a zero curvature equation for some $\alpha_\lambda = \lambda^{-m}\hat{\alpha}_{-m} + \dots + \hat{\alpha}_0 + \dots + \lambda^m\hat{\alpha}_m$. Namely, these method apply to construct the solutions of all the m -th elliptic integrable systems. So it is natural to ask what is the geometric interpretation of these systems. Do they correspond to some generalisations of harmonic maps? This is the problem that we solve in this paper: to describe the geometry behind this family of integrable systems whose we know how to construct (at least locally) all the solutions. The introduction below gives an overview of all the results, as well as some related subjects and works, and some additional motivations.

Contents

• Introduction	5
0.1 The primitive systems	5
0.2 The determined case	6
0.2.1 The minimal determined system	6
0.2.2 The general structure of the maximal determined case.	7
0.2.3 The model system in the even case	7
0.2.4 The model system in the odd case	7

0.2.5	The coupled model system	10
0.2.6	The General maximal determined odd system. ($k' = 2k + 1, m = 2k$)	11
0.2.7	General maximal determined even system. ($k' = 2k, m = 2k - 1$)	12
0.2.8	The intermediate determined systems	14
0.3	The underdetermined case	14
0.4	In the twistor space.	14
0.5	Related subjects and works, and motivations	15
0.5.1	Relations with surface theory.	15
0.5.2	Relations with mathematical physics.	15
0.5.3	Relations of F -stringy harmonicity and supersymmetry	16
0	Notations, Conventions and general definitions	16
0.6	Generalities	16
0.7	Almost complex geometry	17
1	Invariant connections on reductive homogeneous spaces	18
1.1	Linear isotropy representation	19
1.2	Reductive homogeneous space	19
1.3	The (canonical) invariant connection	20
1.4	Associated covariant derivative	20
1.5	G -invariant linear connections in terms of equivariant bilinear maps	21
1.6	A Family of connections on the reductive space M	22
1.7	Differentiation in $\text{End}(T(G/H))$	23
2	m-th elliptic integrable system associated to a k'-symmetric space	25
2.0.1	Definition of G^τ (even when τ does not integrate in G)	25
2.1	Finite order Lie algebra automorphisms	26
2.1.1	The even case: $k' = 2k$	26
2.1.2	the odd case: $k' = 2k + 1$	28
2.2	Definitions and general properties of the m -th elliptic system.	28
2.2.1	Definitions	28
2.2.2	The geometric solution	31
2.2.3	The increasing sequence of spaces of solutions: $(\mathcal{S}(m))_{m \in \mathbb{N}}$	33
2.2.4	The decreasing sequence $(\text{Syst}(m, \tau^p))_{p/k'}$	35
2.3	The minimal determined case	35
2.3.1	The even minimal determined case: $k' = 2k$ and $m = k$	35
2.3.2	The minimal determined odd case	38
2.4	The maximal determined case	42
2.5	The underdetermined case	43
2.6	Examples	43
2.6.1	The trivial case: the 0-th elliptic system associated to a Lie group.	43
2.6.2	Even determined case	44
2.6.3	Primitive case	44
2.6.4	Underdetermined case	44

3	Finite order isometries and Twistor spaces	46
3.1	Isometries of order $2k$ with no eigenvalues $= \pm 1$	47
3.1.1	The set of connected components in the general case	47
3.1.2	Study of $\text{Ad}J$, for $J \in \mathcal{Z}_{2k}^a(\mathbb{R}^{2n})$	49
3.1.3	Study of $\text{Ad}J^j$	51
3.2	Isometries of order $2k + 1$ with no eigenvalue $= 1$	53
3.3	The effect of the power maps on the finite order isometries	53
3.4	The Twistor spaces of a Riemannian manifolds and its reductions	55
3.5	Return to an order $2k$ automorphism $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$.	55
3.5.1	Case $r = k$	55
3.5.2	Action of $\text{Ad}\tau _{\mathfrak{m}}$ on $\text{ad}\mathfrak{g}_j^{\mathbb{C}}$	56
3.6	The canonical section in $(\mathcal{Z}_{2k}(G/H))^2$, the canonical embedding, and the Twistor lifts	57
3.6.1	The canonical embedding	57
3.6.2	The twistor lifts	58
4	Vertically Harmonic maps and Harmonic sections of submersions	60
4.1	Definitions, general properties and examples	60
4.1.1	The vertical energy functional	60
4.1.2	Examples	60
4.1.3	Ψ -torsion, Ψ -difference tensor, and curvature of a Pfaffian system	65
4.2	Harmonic sections of homogeneous fibre bundles	67
4.2.1	Definitions and Geometric properties	68
4.2.2	Vertical harmonicity equation	71
4.2.3	Reductions of homogeneous fibre bundles	73
4.3	Examples of Homogeneous fibre bundles	75
4.3.1	Homogeneous spaces fibration	75
4.3.2	The twistor bundle of almost complex structures $\Sigma(E)$	80
4.3.3	The twistor bundle $\mathcal{Z}_{2k}(E)$ of a Riemannian vector bundle	83
4.3.4	The Twistor subbundle $\mathcal{Z}_{2k,j}^{\alpha}(E)$	86
4.4	Geometric interpretation of the even minimal determined system	92
4.4.1	The injective morphism of homogeneous fibre bundle $\mathcal{J}_{J_0}: G/G_0 \hookrightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$.	92
4.4.2	Conclusion	96
5	Generalized harmonic maps	98
5.1	Affine harmonic maps and holomorphically harmonic maps	98
5.1.1	Affine harmonic maps: general properties	98
5.1.2	Holomorphically harmonic maps	99
5.2	The sigma model with a Wess-Zumino term in Nearly Kähler manifolds	105
5.2.1	Totally skew-symmetric torsion	105
5.2.2	The general case of an almost Hermitian manifold	107
5.2.3	The example of a 3-symmetric space	109
5.2.4	The good geometric context/setting	110
5.2.5	J -twisted harmonic maps	112
5.3	The sigma model with a Wess-Zumino term in \mathcal{G}_1 -manifolds	113
5.3.1	TN -valued 2-forms	113
5.3.2	Stringy Harmonic maps	114
5.3.3	Almost Hermitian \mathcal{G}_1 -manifolds	116
5.3.4	Characterisation of Hermitian connections in terms of their torsion	119

5.3.5	The example of a naturally reductive homogeneous space	120
5.3.6	Geometric interpretation of the maximal determined odd case.	121
5.4	Stringy harmonicity and Holomorphic harmonicity	123
6	Generalized Harmonic maps into f-manifolds.	125
6.1	f -structures: General definitions and properties.	125
6.1.1	f -structures, Nijenhuis tensor and natural action on the space of torsions \mathcal{T}	125
6.1.2	Introducing a linear connection.	126
6.2	The f -connections and their torsion.	127
6.2.1	Definition, notations and first properties.	127
6.2.2	Characterisation of metric connections preserving the splitting.	128
6.2.3	Characterisation of metric f -connections. Existence of a characteristic connection.	132
6.2.4	Precharacteristic and paracheracteristic connections.	138
6.2.5	Reductions of f -manifolds	140
6.3	f -connections on fibre bundles	143
6.3.1	Riemannian submersion and metric f -manifolds of global type \mathcal{G}_1	143
6.3.2	Reductions of f -submersions	147
6.3.3	Horizontally Kähler f -manifolds and horizontally projectible f -submersions.	148
6.3.4	The example of a naturally reductive homogeneous space	150
6.3.5	The example of the twistor space $\mathcal{Z}_{2k}^\alpha(M)$	150
6.3.6	The example of the twistor space $\mathcal{Z}_{2k,j}^\alpha(M, J_j)$	151
6.3.7	The reduction of homogeneous fibre bundle $\mathfrak{I}_{J_0} : G/G_0 \hookrightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$	152
6.4	Stringy Harmonic maps in f -manifolds.	152
6.4.1	Definitions	152
6.4.2	The closeness of the 3-forms $F \bullet T$ and $F \star T$	153
6.4.3	The sigma model with a Wess-Zumino term in reductive metric f -manifold of global type \mathcal{G}_1	158
6.4.4	The example of a naturally reductive homogeneous space	158
6.4.5	Geometric interpretation of the maximal determined even case.	159
6.4.6	Twistorial Geometric interpretation of the maximal determined even case.	161
7	Generalized harmonic maps into reductive homogeneous spaces	162
7.1	Affine harmonic maps into reductive homogeneous spaces.	162
7.2	Affine/holomorphically harmonic maps into 3-symmetric spaces	164
7.3	(Affine) vertically (holomorphically) harmonic maps	166
7.3.1	Affine vertically harmonic maps: general properties	166
7.3.2	Affine vertically holomorphically harmonic maps	166
7.4	Affine vertically harmonic maps into reductive homogeneous space	167
7.5	(Affine) vertically (holomorphically) harmonic maps into reductive homogeneous space with an invariant Pfaffian structure	170
7.6	The intermediate determined systems.	177
7.6.1	The odd case.	177
7.6.2	The even case.	178
7.6.3	Sigma model with a Wess-Zumino term.	179
7.7	Some remarks about the twistorial interpretation.	179
7.7.1	The even case.	179
7.7.2	The odd case.	180

8 Appendix	181
8.1 Vertical harmonicity	181
8.2 G -invariant metrics	183
8.2.1 Existence of an $\text{Ad}H$ -invariant inner product on \mathfrak{m} for which $\tau _{\mathfrak{m}}$ is an isometry.	183
8.2.2 Existence of a naturally reductive metric for which \underline{J} is an isometry, resp. F is metric.	183

Introduction

In this paper, we give a geometric interpretation of all the m -th elliptic integrable systems associated to a k' -symmetric space $N = G/G_0$ (in the sense of C.L. Terng [47]).

Let \mathfrak{g} be a real Lie algebra and $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of finite order k' . This automorphism admits an eigenspace decomposition $\mathfrak{g}^{\mathbb{C}} = \bigoplus_{j \in \mathbb{Z}_{k'}} \mathfrak{g}_j^{\mathbb{C}}$, where $\mathfrak{g}_j^{\mathbb{C}}$ is the eigenspace of τ with respect to the eigenvalue $\omega_{k'}^j$. We denote by $\omega_{k'}$ a k' -th primitive root of unity. Moreover the automorphism τ defines a k' -symmetric space $N = G/G_0$. Furthermore let L be a Riemann surface.

The m -th elliptic integrable system associated to (\mathfrak{g}, τ) can be written as a zero curvature equation

$$d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0, \quad \forall \lambda \in \mathbb{C}^*,$$

where $\alpha_\lambda = \sum_{j=0}^m \lambda^{-j} u_j + \lambda^j \bar{u}_j = \sum_{j=-m}^m \lambda^j \hat{\alpha}_j$ is a 1-form on the Riemann surface L taking values in the Lie algebra \mathfrak{g} . The "coefficient" u_j is a $(1,0)$ -type 1-form on L with values in the eigenspace $\mathfrak{g}_{-j}^{\mathbb{C}}$.

Moreover, we call the integer m the *order* of the system. Let us make precise that the order m has nothing to do with the order of any PDE, but this is only the maximal power on λ in the (finite) Fourier decomposition of α_λ w.r.t. λ .

First, we remark that any solution of the system of order m is a solution of the system of order m' , if $m \leq m'$ (and the automorphism τ is fixed). In other words, the system of order m is a reduction of the system of order m' , if $m \leq m'$.

Moreover, it turns out that we have to introduce the integer $m_{k'}$ defined by

$$m_{k'} = \left\lfloor \frac{k'+1}{2} \right\rfloor = \begin{cases} k & \text{if } k' = 2k \\ k+1 & \text{if } k' = 2k+1 \end{cases} \quad \text{if } k' > 1, \text{ and } m_1 = 0.$$

Then the general problem splits into three cases : the primitive case ($m < m_{k'}$), the determined case ($m_{k'} \leq m \leq k' - 1$) and the underdetermined case ($m \geq k'$).

0.1 The primitive systems

The primitive systems have an interpretation in terms of F -holomorphic maps, with respect to a f -structure F on the target space $N = G/G_0$ (i.e. an endomorphism F satisfying $F^3 + F = 0$). More precisely:

- In the even case ($k' = 2k$), we have a fibration $\pi: G/G_0 \rightarrow G/H$ over a k -symmetric space $M = G/H$ (defined by the square τ^2 of the automorphism τ of order k' defining $N = G/G_0$). We also have a G -invariant splitting $TN = \mathcal{H} \oplus \mathcal{V}$ corresponding to this fibration¹, where $\mathcal{V} = \ker d\pi$.

¹i.e. \mathcal{H} is a connection on this fibration.

Moreover N is naturally endowed with a f -structure F which defines a complex structure on the horizontal subbundle \mathcal{H} and vanishes on the vertical subbundle \mathcal{V} . Furthermore, the eigenspace decomposition of the order k' automorphism τ gives us some G -invariant decomposition $\mathcal{H} = \bigoplus_{j=1}^{k'-1} [\mathfrak{m}_j]$, where $\mathfrak{m}_j \subset \mathfrak{g}$ is defined by $\mathfrak{m}_j^{\mathbb{C}} = \mathfrak{g}_{-j}^{\mathbb{C}} \oplus \mathfrak{g}_j^{\mathbb{C}}$, and $[\mathfrak{m}_j] \subset TN$ is the corresponding G -invariant subbundle. This allows to define, by multiplying F on the left by the projections on the subbundles $\mathcal{H}^m = \bigoplus_{j=1}^m [\mathfrak{m}_j]$, a family of f -structures $F^{[m]}$, $1 \leq m \leq k-1$. Then the primitive system of order m ($m < m_{k'} = k$) associated to G/G_0 is exactly the equation for $F^{[m]}$ -holomorphic maps. Therefore the solutions of the primitive systems are exactly the F -holomorphic maps.

- *In the odd case* ($k' = 2k+1$), $N = G/G_0$ is naturally endowed with an almost complex structure \underline{J} . Then the solutions of the primitive systems are exactly the \underline{J} -holomorphic curves. Moreover, in the same way as for the even case, the eigenspace decomposition of τ provides a G -invariant decomposition $TN = \bigoplus_{j=1}^k [\mathfrak{m}_j]$, which allows to define a family of f -structures $F^{[m]}$, $1 \leq m \leq k$, with $F^{[k]} = \underline{J}$. Then the primitive system of order m ($m < m_{k'} = k+1$) associated to G/G_0 is exactly the equation for $F^{[m]}$ -holomorphic maps. In other words, the solutions of the primitive system of order m are exactly the integral holomorphic curves of the complex Pfaffian system $\bigoplus_{j=1}^m [\mathfrak{m}_j] \subset TN$ in the almost complex manifold (N, \underline{J}) .

0.2 The determined case

We call "the minimal determined system" the determined system of minimal order $m_{k'}$, and "the maximal determined system" the determined system of maximal order $k' - 1$.

Any solution of a determined system is solution of the corresponding maximal determined system. More precisely, a map $f: L \rightarrow G/G_0$ is solution of a determined system (associated to G/G_0) if and only if it is solution of the maximal determined system (associated to G/G_0) and satisfies an additional holomorphicity condition. When this holomorphicity condition is maximal, then we obtain the minimal determined system.

0.2.1 The minimal determined system

The minimal determined system has an interpretation in terms of horizontally holomorphic and vertically harmonic maps $f: L \rightarrow N = G/G_0$. It also has an equivalent interpretation in terms of vertically harmonic twistor lifts in some twistor space. Let us make precise this point.

In the even case. As we have seen in the subsection 0.1 below, the homogeneous space $N = G/G_0$ admits a G -invariant splitting $TN = \mathcal{H} \oplus \mathcal{V}$ corresponding to the fibration $\pi: N \rightarrow M$ and N is naturally endowed with a f -structure F which defines a complex structure on the horizontal subbundle \mathcal{H} and vanishes on the vertical subbundle \mathcal{V} . Then we say that a map $f: L \rightarrow N$ is horizontally holomorphic if

$$(df \circ j_L)^{\mathcal{H}} = F \circ df.$$

Then we prove that *the even minimal determined system* ($\text{Syst}(k, \tau)$) means that the geometric map f is horizontally holomorphic and vertically harmonic, i.e.

$$\tau^v(f) := \text{Tr}_g(\nabla^v d^v f) = 0$$

(for any hermitian metric g on the Riemann surface L). Here ∇^v is the vertical component of the Levi-Civita connection ∇ (of some G -invariant metric on N). Moreover, vertically harmonic

maps are exactly the critical points (w.r.t. vertical variations) of the vertical energy functional:

$$E^v(u) = \frac{1}{2} \int_L |d^v u|^2 d\text{vol}_g.$$

We prove also that this system also has an equivalent interpretation in terms of vertically harmonic twistor lifts in the twistor space $\mathcal{Z}_{2k,j}(M, J_2)$ which is a subbundle of $\mathcal{Z}_{2k}(M)$, where

$$\mathcal{Z}_{k'}(M) = \{J \in SO(TM) \mid J^{k'} = \text{Id}, J^p \neq \text{Id} \text{ if } p < k', \ker(J \pm \text{Id}) = \{0\}\}$$

is the bundle of isometries of TM with finite order k' and with no eigenvalues $= \pm 1$. More precisely denoting by J_2 the order k isometry of M defined by τ_m^2 , then we define $\mathcal{Z}_{2k,j}(M, J_2) = \{J \in \mathcal{Z}_{2k}(M) \mid J^2 = J_2\}$. Then we prove that $N = G/G_0$ can be embedded into the twistor space $\mathcal{Z}_{2k,j}(M, J_2)$ via a natural morphism of bundle over $M = G/H$. We prove that $f: L \rightarrow N$ is solution of the system if and only if the corresponding map $J^f: L \rightarrow \mathcal{Z}_{2k,j}(M, J_2)$ is a vertically harmonic twistor lift.

In the odd case. We obtain an analogous interpretation as in the even case. An interpretation in terms of horizontally holomorphic and vertically harmonic maps $f: L \rightarrow N = G/G_0$. And by embedding G/G_0 into the twistor space $\mathcal{Z}_{2k+1}(N)$ of order $2k+1$ isometries in N , we obtain an interpretation in terms of vertically harmonic twistor lift.

0.2.2 The general structure of the maximal determined case.

First, the maximal determined system has 3 model cases. This means that there are 3 maximal determined systems, namely the three maximal determined systems with lowest order of symmetry (2,3,4), of which the corresponding geometric equations (when put all together) contain already all the structure terms - in a simple form- that will appear in the further maximal determined systems in a more complex and general form due to the more complex geometric structure in the further maximal determined systems. That is in this sense that we can say that all the further determined systems associated to target spaces N with higher order of symmetry will be modeled on these model systems.

0.2.3 The model system in the even case

In the even case, this model is the first elliptic integrable system associated to a symmetric space ($m = 1, k' = 2$) which is - as it is well known - exactly the equation of harmonic maps from the Riemann surface L into the symmetric space under consideration. This is the "smallest" determined system, i.e. with lowest order of symmetry in the target space $N = G/G_0$. In this case - N is symmetric- the determined case is reduced to one system, the one of order 1.

0.2.4 The model system in the odd case

In the odd case, this model is the second elliptic integrable system associated to a 3-symmetric space. This is the "smallest" determined system in the odd case, i.e. with lowest odd order of symmetry in the target space $N = G/G_0$. We prove that this system is exactly the equation for *holomorphically harmonic maps* into the almost complex manifold (N, \underline{J}) with respect to the anticanonical connection $\nabla^1 = \nabla^0 + [,]_{[\underline{m}]}$, where ∇^0 is the canonical connection. Or equivalently this is the equation for holomorphically harmonic maps into the almost complex manifold $(N, -\underline{J})$ with respect to the canonical connection ∇^0 .

Holomorphically harmonic maps. Given a general almost complex manifold (N, J) with a connection ∇ , we define holomorphically harmonic maps $f: L \rightarrow N$ as the solutions of the equation

$$[\bar{\partial}^\nabla \partial f]^{1,0} = 0 \quad (1)$$

where $[\]^{1,0}$ denotes the $(1,0)$ -component according to the splitting $TN^\mathbb{C} = T^{1,0}N \oplus T^{0,1}N$ defined by J . This equation is equivalent to

$$d^\nabla df + Jd^\nabla * df = 0$$

or, equivalently, using any Hermitian metric g on L

$$T_g(f) + J\tau_g(f) = 0$$

where $T_g(f) = *f^*T = f^*T(e_1, e_2)$, with (e_1, e_2) an orthonormal basis of TL , and $\tau_g(f) = *d^\nabla * df = \text{Tr}_g(\nabla df)$ is the tension field of f . Of course Tr_g denotes the trace with respect to g , and the expression ∇df denotes the covariant derivative of df with respect to the connection induced in $T^*L \otimes f^*TN$ by ∇ and the Levi-Civita connection in L .

In particular, we see that if ∇ is torsion free or more generally if f is torsion free, i.e. $f^*T = 0$, then holomorphic harmonicity is equivalent to harmonicity. Therefore, this new notion is interesting only in the case of a non torsion free connection ∇ .

The vanishing of some $\bar{\partial}\partial$ -derivative. Now, let us suppose that the connection ∇ on N is *almost complex*, i.e. $\nabla J = 0$. Then, according to equation (1), we see that any holomorphic curve $f: (L, j_L) \rightarrow (N, J)$ is anti-holomorphically harmonic, i.e. holomorphically harmonic with respect to $-J$. In particular, this allows to recover that a 1-primitive solution (i.e. of order $m = 1$) of the elliptic system associated to a 3-symmetric space is also solution of the second elliptic system associated to this space.

Moreover, the holomorphically harmonic maps admit a formulation very analogous to that of harmonic maps in term of the vanishing of some $\bar{\partial}\partial$ -derivative, which implies a well known characterisation in term of holomorphic 1-forms. Indeed we prove that $f: (L, j_L) \rightarrow (N, J, \nabla)$ is holomorphically harmonic if and only if

$$\widehat{\bar{\partial}} \widehat{\partial} f = 0, \quad (2)$$

i.e. $\widehat{\partial} f$ is a holomorphic section of $T_{1,0}^*L \otimes_{\mathbb{C}} f^*TN$. Here the hat " ^ " means that we extend a 1-form on TL , like d or ∇ , by \mathbb{C} -linearity as a linear map from $TL^\mathbb{C}$ into the complex bundle (TN, J) . In other words instead of extending these 1-forms as \mathbb{C} -linear maps from $TL^\mathbb{C}$ into $TN^\mathbb{C}$ as it is usual, we use the already existing structure of complex vector bundle in (TN, J) and extend these very naturally as \mathbb{C} -linear map from $TL^\mathbb{C}$ into the complex bundle (TN, J) . Therefore we can conclude that holomorphically harmonic maps have the same formulation as harmonic maps with the difference that instead of working in the complex vector bundle $TN^\mathbb{C}$, we stay in TN which is already a complex vector bundle in which we work.

The sigma model with a Wess-Zumino term. Finally, let us suppose that N is endowed with a ∇ -parallel Hermitian metric h . Therefore (N, J, h) is an almost Hermitian manifold with a Hermitian connection ∇ . Suppose also that J anticommutes with the torsion T of ∇ i.e.

$$T(X, JY) = -JT(X, Y)$$

which is equivalent to

$$T = \frac{1}{4}N_J$$

where N_J denotes the torsion of J i.e its Nijenhuis tensor.

Suppose also that the torsion of ∇ is totally skew-symmetric i.e. the trilinear map

$$T^*(X, Y, Z) = \langle T(X, Y), Z \rangle$$

is a 3-form. Lastly, we suppose that the torsion is ∇ -parallel, i.e. $\nabla T^* = 0$ which is equivalent to $\nabla T = 0$. Then we prove that this implies that the 3-form

$$H(X, Y, Z) = -T^*(X, Y, JZ) = \langle JT(X, Y), Z \rangle$$

is closed $dH = 0$.²

Then the equation for holomorphically harmonic maps $f: L \rightarrow N$ is the equation of motion (i.e. Euler-Lagrange equation) for the sigma model in N with the Wess-Zumino term defined by the closed 3-form H . The action functional is given by

$$S(f) = E(f) + S^{WZ}(f) = \frac{1}{2} \int_L |df|^2 d\text{vol}_g + \int_B H,$$

where B is 3-submanifold (or indeed a 3-chain) in N whose boundary is $f(L)$.

Then since $dH = 0$, the variation of the Wess-Zumino term is a boundary term

$$\delta S^{WZ} = \int_B L_{\delta f} H = \int_B d\iota_{\delta f} H = \int_{f(L)} \iota_{\delta f} H,$$

whence its contribution to the Euler-Lagrange equation involves only the original map $f: L \rightarrow N$. In particular, applying this result to the case we are interested in, i.e. N is 3-symmetric, we obtain:

*The second elliptic system associated to 3-symmetric space $N = G/G_0$ is the equation of motion for the sigma model in N with the Wess-Zumino term defined by the closed 3-form H , corresponding to the canonical almost complex structure $-\underline{J}$ and the canonical connection ∇^0 .*³

The good geometric context/setting In the previous variational interpretation, we need to make 3 hypothesis on the torsion of the Hermitian connection: T anticommutes with J , is totally skew-symmetric and ∇ -parallel. It is natural to ask ourself what do these hypothesis mean geometrically and what is the good geometric context in which these take place. It turns out that the good geometric context is the one of *Nearly Kähler manifold*.

An almost Hermitian manifold (N, J, h) is Nearly Kähler if and only if $(\nabla_X^h J)X = 0$, for all $X \in TN$, where ∇^h is the Levi-Civita connection. Then we prove that the almost Hermitian

²Let us point out that in general T^* is not closed even if it is ∇ -parallel. For example, in a Riemannian naturally reductive homogeneous space G/H , endowed with its canonical connection ∇^0 , we have $\nabla^0 T = 0$ but $dT^*(X, Y, Z, V) = -2\langle \text{Jac}_m(X, Y, Z), V \rangle$ where Jac_m is the m -component of the Jacobi identity (i.e. the sum of the circular permutations of $[X, [Y, Z]_m]_m$). Of course m denotes the $\text{Ad}H$ -invariant summand in the reductive decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$.

³In fact, we need a naturally reductive metric on N to ensure that T^* is a 3-form. But if we allow Pseudo-Riemannian metrics and if \mathfrak{g} is semisimple then the metric defined by the Killing form is naturally reductive. In fact, the elliptic integrable system is a priori written in an affine context, i.e. its natural - in the sense of initial-geometric interpretation takes place in the context of affine geometry in terms of the linear connections $\nabla^t = \nabla^0 + t[\cdot, \cdot]_{[m]}$. If we want that this interpretation takes place in the context of Riemannian geometry we need, of course, to add some hypothesis of compactness, like the compactness of $\text{Ad}_m G_0$ and the natural reductivity. But we do not need these hypothesis if we work in the Pseudo-Riemannian context.

manifolds for which there exists an Hermitian connection satisfying the three hypothesis above are exactly Nearly Kähler manifolds. Then the variational interpretation can be rewritten as follows:

Theorem 0.1 *Let (N, h, J) be a nearly Kähler manifold then the equation of holomorphic harmonicity for maps $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the 3-form:*

$$H = \frac{1}{3}d\Omega_J$$

where $\Omega_J = \langle J\cdot, \cdot \rangle$ is the Kähler form.

Therefore: the second elliptic system associated to 3-symmetric space $N = G/G_0$, endowed with its canonical almost complex structure \underline{J} , is the equation of motion for the sigma model in N with the Wess-Zumino term defined by the closed 3-form $H = -\frac{1}{3}d\Omega_J$.

J -twisted harmonic maps. We prove that we can also interpret the holomorphic harmonicity in terms of J -twisted harmonic maps (w.r.t. the Levi-Civita connection). Let us define this notion. Let (E, J) be a complex vector bundle over an almost complex manifold (M, j_M) . Then let $\bar{\nabla}$ be a connection on E . Then we can decompose it in an unique way as the sum of a J -commuting and a J -anticommuting part, i.e. in the form

$$\bar{\nabla} = \nabla^0 + A$$

where $\nabla^0 J = 0$ and $A \in \mathcal{C}(T^*M \otimes \text{End}(E))$, $AJ = -JA$. More precisely, we have $A = \frac{1}{2}J\bar{\nabla}J$. Then we set

$$\bar{\nabla}^J = \nabla^0 - (A \circ j_M)J = \bar{\nabla} - \frac{1}{2}J\bar{\nabla}J - \frac{1}{2}\bar{\nabla}J \circ j_M.$$

Now let $f: (L, j_L) \rightarrow (N, J)$ be a map from a Riemann surface into the almost complex manifold (N, J) endowed with a connection ∇ . Then let us take in what precede $(M, j_M) = (L, j_L)$ and $(E, \bar{\nabla}) = (f^*TN, f^*\nabla)$. Then we say that the map $f: (L, j_L) \rightarrow (N, J, \nabla)$ is J -twisted harmonic if and only if

$$\text{Tr}_g(\bar{\nabla}^J df) = 0$$

(for any hermitian metric g on the Riemann surface L).

0.2.5 The coupled model system

This is the third elliptic integrable system associated to a 4-symmetric space. The corresponding geometric equation in the 4-symmetric space G/G_0 can be viewed as a coupling between the equation of harmonic maps into the symmetric fibre H/G_0 and the equation for harmonic maps into the symmetric space G/H . In other words, it can be viewed as a coupling between the even model system associated to the symmetric space G/H and the even model system associated to the symmetric space H/G_0 . In particular, when harmonicity is replaced by holomorphicity "in the basis G/H ", then we recover the horizontally holomorphic and vertically harmonic maps into G/G_0 , that is to say the solutions of the second elliptic integrable system associated to G/G_0 . We will come back to this in the end of section 0.2.7 in this introduction.

0.2.6 The General maximal determined odd system. ($k' = 2k + 1, m = 2k$)

The maximal determined odd system has a geometric interpretation in terms of *stringy harmonic* maps $f: L \rightarrow (G/G_0, \underline{\mathbb{J}})$, with respect to the canonical connection and the canonical almost complex structure.

Stringy harmonic maps. Let (N, J) be an almost complex manifold with ∇ an linear connection then we will say that a map $f: L \rightarrow N$ from a Riemann surface into N is *stringy harmonic* if it is solution of *the harmonic map equation with a JT-term*:

$$-\tau_g(f) + (J \cdot T)_g(f) = 0.$$

We have used the notation $J \cdot B = -JB(J \cdot, \cdot)$, $\forall B \in \mathcal{C}(\Lambda^2 T^*N \otimes TN)$. This action of J on $B \in \mathcal{C}(\Lambda^2 T^*N \otimes TN)$ can be written more naturally is (N, J) is endowed with a Hermitian metric h . Indeed, in this case, we have an identification, $\mathcal{C}(\Lambda^2 T^*N \otimes TN) = \mathcal{C}(\Lambda^2 T^*N \otimes T^*N) \subset \mathcal{C}(\otimes^3 T^*N)$, between TN -valued 2-forms on N and trilinear forms on N skew-symmetric w.r.t. the 2 first variables: $B(X, Y, Z) := \langle B(X, Y), Z \rangle$. Then $J \cdot B$ is written:

$$J \cdot B = B(J \cdot, J \cdot, J \cdot) =: -B^c.$$

We remark that if T anticommutes with J then stringy harmonicity coincides with holomorphic harmonicity (since in this case $J \cdot T = JT$). More particularly, if $T = 0$, then the stringy harmonicity coincide with the harmonicity.

Furthermore, we look for a general geometric setting in which the stringy harmonicity has an interesting interpretation. First of all, let us remark that in the context of homogeneous reductive space, in which our system takes places, we have a canonical connection, with respect to which the stringy harmonicity can be written "canonically". But in general we do not have a "special" connection with respect to which one can consider the stringy harmonicity. Therefore, if one wants to place stringy harmonicity in a more meaningful, interesting and fruitful context (than the general context of almost complex manifolds endowed with some linear connection) and, in so doing, obtain a better understanding of our elliptic integrable system by writting its geometric interpretation in the best geometric context, a first problem - that we solved - is to find a general class of (almost complex) manifold in which there exists some unique "canonical" connection, with respect to which we then could consider the stringy harmonicity. This provides us, firstly, some special connection (in the same sense that the Levi-Civita connection is special in Riemannian geometry), which solves the problem of the choice of the connection, but secondly it turns out that it provides also a variational interpretation of the stringy harmonicity.

Best geometric setting It turns out that the more rich geometric context in which stringy harmonicity admits interesting properties is the one of \mathcal{G}_1 -manifolds, more precisely \mathcal{G}_1 -manifolds whose the characteristic connection has a parallel torsion. Making systematic use of the covariant derivative of the Kähler form, A. Gray and L. M. Hervella, in the late seventies, classified almost Hermitian structures into sixteen classes. Denote by \mathcal{W} the space of all trilinear forms (on some Hermitian vector spaces, say $T_{y_0}N$ for some reference point $y_0 \in N$) having the same algebraic properties as $\nabla^h \Omega_J$. Then they proved that we have a $U(n)$ -irreducible decomposition $\mathcal{W} = \mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$. The sixteen classes are then respectively the classes of almost Hermitian manifolds for which $\nabla^h \Omega_J$ 'lies in' the $U(n)$ -invariant subspaces $\mathcal{W}_I = \oplus_{i \in I} \mathcal{W}_i$, $I \subset \{1, \dots, 4\}$, respectively. In particular, if we take as invariant subspace $\{0\}$, we obtain the Kähler manifolds, if we take \mathcal{W}_1 , we obtain the class of nearly Kähler manifolds. Moreover the class of \mathcal{G}_1 -manifolds is the one defined by $\mathcal{G}_1 = \mathcal{W}_1 \oplus \mathcal{W}_3 \oplus \mathcal{W}_4$. It is characterised by (N, J, h) is of type \mathcal{G}_1 if and

only if the Nijenhuis tensor N_J is totally skew-symmetric (i.e. a 3-form). In this paper, we prove the following theorem⁴:

Theorem 0.2 *An almost Hermitian manifold (N, J, h) admits a Hermitian connection with totally skew-symmetric torsion if and only if the Nijenhuis tensor N_J is itself totally skew-symmetric. In this case, the connection is unique and determined by its torsion which is given by*

$$T = -d^c\Omega_J + N_J.$$

The characteristic connection is then given by $\nabla = \nabla^h - \frac{1}{2}T$.

Then we prove:

Proposition 0.1 *Let us suppose that the almost Hermitian manifold (N, J, h) is a \mathcal{G}_1 -manifold. Let us suppose that its characteristic connection ∇ has a parallel torsion $\nabla T = 0$. Then the 3-form*

$$H(X, Y, Z) = T(JX, JY, JZ) = \langle (J \cdot T)(X, Y), Z \rangle$$

is closed $dH = 0$.

Which then gives us the following variational interpretation

Theorem 0.3 *Let us suppose that the almost Hermitian manifold (N, J, h) is a \mathcal{G}_1 -manifold. Let us suppose that its characteristic connection ∇ has a parallel torsion $\nabla T = 0$. Then the equation for stringy harmonic maps $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H = -d\Omega_J + JN_J.$$

Moreover any $(2k+1)$ -symmetric space $(G/G_0, \underline{J}, h)$ endowed with its canonical complex structure and a naturally reductive G -invariant metric h (for which \underline{J} is orthogonal) is a \mathcal{G}_1 -manifold and moreover its characteristic connection coincides with its canonical connection ∇^0 . Finally, the torsion of the canonical connection is obviously parallel. Therefore we obtain an interpretation of the maximal determined system associated to a $(2k+1)$ -symmetric space in terms of a sigma models with a Wess-Zumino term.

0.2.7 General maximal determined even system. $(k' = 2k, m = 2k - 1)$

In the even case, the geometric structure of the target space G/G_0 is more complex (and more rich): as we have already seen, there is a fibration $\pi: N = G/G_0 \rightarrow M = G/H$, a splitting $TN = \mathcal{H} \oplus \mathcal{V}$ with $\mathcal{V} = \ker \pi$, and a f -structure F such that $\ker F = \mathcal{V}$ and $\text{Im } F = \mathcal{H}$ (in particular $\bar{J} := F_{\mathcal{H}}$ is a complex structure on \mathcal{H}). Moreover the geometric PDE obtained from our elliptic integrable system calls for this geometric structure. In particular, this geometric PDE splits into its horizontal and vertical parts and can be viewed as a coupling between the equation of \bar{J} -stringy harmonicity and the equation of vertical harmonicity, the coupling terms calling out the curvature of \mathcal{H} .

The maximal determined even system has a geometric interpretation in terms of *stringy harmonic* maps $f: L \rightarrow (G/G_0, F)$, F being the canonical f -structure on G/G_0 .

⁴After the fact, we realized that it has already been stated by Friedrich-Ivanov[17]. However we give a different and more complete proof. See remark 5.14

Stringy harmonic maps w.r.t. a f -structure. Let (N, F) be a f -manifold with ∇ a linear connection. Then we will say that a map $f: L \rightarrow N$ from a Riemann surface into N is *stringy harmonic* if it is solution of the *stringy harmonic maps equation*:

$$-\tau_g(f) + (F \bullet T)_g(f) = 0.$$

where $F \bullet B$, for $B \in \mathcal{C}(\Lambda^2 T^* N \otimes TN)$, denotes some natural (linear) action of F on $\mathcal{C}(\Lambda^2 T^* N \otimes TN)$. For more simplicity, let us write it in the case (N, F) is endowed with a compatible metric h (i.e. $\mathcal{V} \perp \mathcal{H}$ and \bar{J} is orthogonal with respect to $h|_{\mathcal{H} \times \mathcal{H}}$):

$$\begin{aligned} F \bullet B &= B(F \cdot, F \cdot, F \cdot) + \frac{1}{2} F \oslash (B - B_{\mathcal{H}^3}) \\ F \oslash A &= A(F \cdot, \cdot, \cdot) + A(\cdot, F \cdot, \cdot) + A(\cdot, \cdot, F \cdot) \end{aligned}$$

for all $B, A \in \mathcal{C}(\Lambda^2 T^* N \otimes TN)$.

Now, we want to proceed as in the case of stringy harmonicity with respect to a complex structure. That is to say find a class of f -manifolds for which there exists some unique characteristic connection which preserves the structure and then look for a variational interpretation of the stringy harmonicity with respect to this connection.

Best Geometric context. We look for metric f -manifolds (N, F, h) for which there exists metric f -connection ∇ (i.e. $\nabla F = 0$ and $\nabla h = 0$) with skew-symmetric torsion T . In a first step, we consider metric connections which preserve the splitting $TN = \mathcal{V} \oplus \mathcal{H}$ (i.e. $\nabla q = 0$, where q is the projection on \mathcal{V}) and characterize the manifolds (N, h, q) for which there exists such a connection with skew-symmetric torsion, and call these *reductive* metric f -manifolds. Then saying about a metric f -manifolds (N, F, h) that it is of global type \mathcal{G}_1 if its extended Nijenhuis tensor \tilde{N}_F is skew-symmetric, we prove the following theorem:

Theorem 0.4 *A metric f -manifold (N, F, h) admits a metric f -connection ∇ with skew-symmetric torsion if and only if it is reductive and of global type \mathcal{G}_1 . Moreover, in this case, for any $\alpha \in \mathcal{C}(\Lambda^3 \mathcal{V}^*)$, there exists a unique metric connection ∇ with skew-symmetric torsion such that $T|_{\Lambda^3 \mathcal{V}} = \alpha$. This unique connection is given by*

$$T = (-d^c \Omega_F + N_{F|_{\mathcal{H}^3}}) + \text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}}) + \alpha.$$

where $\Omega_F = \langle F \cdot, \cdot \rangle$, Φ and $R_{\mathcal{V}}$ are resp. the curvature of \mathcal{H} and \mathcal{V} resp., and Skew is the sum of all the circular permutations on the three variables.

On a metric f -manifold (N, F, h) , a metric f -connection ∇ with skew-symmetric torsion will be called a *characteristic connection*.

Moreover, we prove that for any reductive metric f -manifold of global type \mathcal{G}_1 , the closure of $H = F \bullet T$ is equivalent to the closure of the 3-form $F \cdot N_F - \frac{1}{2} F \oslash (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}}))$, so that:

Theorem 0.5 *Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . Let us suppose that the 3-form $F \cdot N_F - \frac{1}{2} F \oslash (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}}))$ is closed, where Φ and $R_{\mathcal{V}}$ are respectively the curvatures of the horizontal and vertical subbundles. Let ∇ be one characteristic connection. Then the equation for stringy harmonic maps (w.r.t. ∇) $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H = -d\Omega_F + F \cdot N_F - \frac{1}{2} F \oslash (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}})).$$

Contrary to the case of stringy harmonic maps into an almost Hermitian \mathcal{G}_1 -manifolds, in the present case, the hypothesis that the torsion of one characteristic connection is parallel $\nabla T = 0$ does not imply the closure of the 3-form $H = F \bullet T$. However, we characterize this closure under the hypothesis $\nabla T = 0$ and $R_{\mathcal{V}} = 0$, by some 2 conditions that we will not explain in this introduction (see section 6.4.2): the horizontal complex structure \bar{J} is a cyclic permutation of the horizontal curvature, and the 2-forms $N_{\bar{J}}$ and Φ have orthogonal supports.

Moreover any $2k$ -symmetric space $(G/G_0, F, h)$ endowed with its canonical f -structure and a naturally reductive G -invariant metric h (compatible with F) is reductive and of global type \mathcal{G}_1 , and moreover its canonical connection ∇^0 is a characteristic connection. Furthermore, the torsion of the canonical connection is obviously parallel. Finally, we prove that any $2k$ -symmetric space $(G/G_0, F, h)$ satisfies the two hypothesis above. Therefore we obtain an interpretation of the maximal determined even system associated to a $2k$ -symmetric space in terms of a sigma models with a Wess-Zumino term.

A particular case: Horizontally Kähler f -manifolds. Let (N, F, h) be a metric f -manifold. We will say that (N, F, h) is horizontally Kähler if $DF|_{\mathcal{H}^3} = 0$.

Then, we prove that in this case, the two hypothesis above (which characterise the closure of the 3-form H) are satisfied. Moreover, any characteristic connection ∇ satisfies $T_{\mathcal{H}^3} = 0$, which leads to special properties.

A example of this situation is given by any 4-symmetric space, endowed with its canonical f -structure and a naturally reductive G -invariant metric (compatible with F).

0.2.8 The intermediate determined systems

For the intermediate determined systems ($m_{k'} < m < k' - 1$), these are obtained from the maximal determined case by adding holomorphicity in the subbundle $\mathcal{H}^m = \bigoplus_{j=1}^{\underline{m}} [\mathfrak{m}_j] \subset \mathcal{H}$, where $\underline{m} = k' - 1 - m$. It means that the m -th determined system has a geometric interpretation in terms of stringy harmonic maps which are \mathcal{H}^m -holomorphic:

$$(df \circ j_L)^{\mathcal{H}^m} = F^{[\underline{m}]} \circ df.$$

0.3 The underdetermined case

We prove that the m -th underdetermined system ($m > k' - 1$) is in fact equivalent to some m -th determined or primitive system associated to some new automorphism $\tilde{\tau}$ defined in a product \mathfrak{g}^{q+1} , of the initial Lie algebra \mathfrak{g} . More precisely, we write

$$m = qk' + r, \quad 0 \leq r \leq k' - 1$$

the Euclidean division of m by k' . Then we consider the automorphism in \mathfrak{g}^{q+1} defined by

$$\tilde{\tau}: (a_0, a_1, \dots, a_q) \in \mathfrak{g}^{q+1} \mapsto (a_1, \dots, a_q, \tau(a_0)) \in \mathfrak{g}^{q+1}.$$

Then $\tilde{\tau}$ is of order $(q+1)k'$. We prove that the initial m -th (underdetermined) system associated to (\mathfrak{g}, τ) is in fact equivalent to the m -th (determined) system associated to $(\mathfrak{g}^{q+1}, \tilde{\tau})$.

0.4 In the twistor space.

For each previous geometric interpretation in the target space $N = G/G_0$, there is a corresponding geometric interpretation in the twistor space. Indeed, in the odd case we have a section defined by the embedding

$$G/G_0 \hookrightarrow \mathcal{Z}_{2k+1}(G/G_0)$$

whereas in the even case we have a morphism of bundle over $M = G/H$ defined by the embedding

$$G/G_0 \hookrightarrow \mathcal{Z}_{2k}(G/H).$$

The geometric interpretations in the twistor spaces are universal since these twistor space are defined for any Riemannian manifold and are endowed canonically with the different geometric structures that we need to suppose the target space N to be endowed with, in our previous geometric interpretations.

0.5 Related subjects and works, and motivations

0.5.1 Relations with surface theory.

The theory of harmonic maps of surfaces has been greatly enriched by ideas and methods from integrable systems [7, 9, 8, 11, 16, 23, 24, 48]. In particular, these ideas have revolutionised the theory of harmonic maps from a surface into a symmetric space and so, via an appropriate Gauss map, the theory of constant mean curvature surfaces and Willmore surfaces among others. For example, Pinkall and Sterling [41] were able to give an algebro-geometric construction of all constant mean curvature tori while Dorfmeister-Pedit-Wu [16] gave a Weierstrass formula for all constant mean curvature immersions of any (simply connected) surface in terms of holomorphic data. These advances were taken up by Hélein and Romon [25, 26, 27] who showed that similar ideas could be applied to the study of Hamiltonian stationary Lagrangian surfaces in a 4-dimensional Hermitian symmetric space. It was the first example of second elliptic integrable system associated to a 4-symmetric space. In [32], we presented a new class of isotropic surfaces in the Euclidean space of dimension 8 by identifying \mathbb{R}^8 with the set of octonions \mathbb{O} , and we proved that these surfaces are solutions of a second elliptic integrable system associated to a 4-symmetric space. By restriction to $\mathbb{R}^4 = \mathbb{H}$, we obtained the Hamiltonian stationary Lagrangian surfaces and by restriction to $\mathbb{R}^3 = \text{Im } \mathbb{H}$, we obtained the CMC surfaces. Furthermore, in [34], we presented a geometric interpretation of all the second elliptic integrable systems associated to a 4-symmetric space in terms of vertically harmonic twistor lifts of conformal immersions into the Riemannian symmetric space (associated to our 4-symmetric space) (see also [10]). When the previous Riemannian symmetric space is 4-dimensional, then any conformal immersion admits a unique twistor lift and the vertical harmonicity of this twistor lift is equivalent to the holomorphicity of the mean curvature vector of the conformal immersion (see [10]). In particular, when the Riemannian symmetric space is Hermitian, we obtain a conceptual explanation of the result of Hélein-Romon.

0.5.2 Relations with mathematical physics.

Metric connections with totally skew-symmetric torsion. We refer the reader to [2, 17] about connections with skew-symmetric torsions and their relations to physics and more particularly string theory. Linear metric connections with totally skew-symmetric torsion recently became a subject of interest in theoretical and mathematical physics. Let us give here some examples (taken from [17]).

- The target space of supersymmetric sigma models with Wess-Zumino term carries a geometry of a metric connection with skew-symmetric torsion.
- In supergravity theories, the geometry of the moduli space of a class of black holes is carried out by a metric connection with skew-symmetric torsion.
- The geometry of NS-5 brane solutions of type II supergravity theories is generated by a metric connection with skew-symmetric torsion.

- The existence of parallel spinors with respect to a metric connection with skew-symmetric torsion on a Riemannian spin manifold is of importance in string theory, since they are associated with some string solitons (BPS solitons).

The sigma-models Nonlinear sigma-models provide a much-studied class of field theories of both phenomenological and theoretical interest. The chiral model for example summarizes many low energy QCD interactions while 2-dimensional sigma-models may possess nontrivial classical field configurations and have analogies with 4-dimensional Yang-Mills equations but are simpler to handle. It was in the study of the chiral model that Wess and Zumino introduced their parity violating term satisfying anomalous Ward-identities. This term now has a far broader interpretation in terms of anomalies. Sigma-models also have connections with string theories.

Two dimensional sigma-models have already proven a fertile arena for the interplay of topology, geometry, and physics. The (supersymmetric) sigma-models have been generalized by introducing a Wess-Zumino term into the Lagrangian. This term may be interpreted as adding torsion to the canonical Levi-Civita connection of the earlier models. The addition of such torsion can have a marked effect and imposes constraints on the possible geometries of the target.

0.5.3 Relations of F -stringy harmonicity and supersymmetry

The P.D.E of F -stringy harmonicity splits following the splitting $TN = \mathcal{H} \oplus \mathcal{V}$ defined by the f -structure F .

More precisely, this equation is a coupling between the equation of \bar{J} -stringy harmonicity and the equation of vertical harmonicity, the coupling terms calling out the curvatures of \mathcal{H} and \mathcal{V} , and the component $N_{F|\mathcal{V} \times \mathcal{H}}$ of the Nijenhuis tensor.

Moreover, let us suppose that we have a fibration $\pi: NtoM$. Then we have a supersymmetric interpretation of the F -stringy harmonicity: F -stringy harmonicity can be viewed as a supersymmetric extension of the J -stringy harmonicity. In the splitting $TN = \mathcal{H} \oplus \mathcal{V}$, the horizontal subbundle played the role of the odd part and the vertical subbundle plays the role of the even part. In other words, the bosonic equation is a harmonic map equation (the vertical harmonicity) and the fermionic equation is the \bar{J} -stringy harmonic map equation.

Let us also mention that, in [33], we obtained a supersymmetric interpretation of all the second elliptic integrable systems associated to a 4-symmetric space in terms of superharmonic maps.

Acknowledgements The author wishes to thank Josef Dorfmeister for his useful comments on the first parts of this paper. He is also grateful to him for his interest in the present work, his encouragements as well as his support during the preparation of this paper.

Notations, Conventions and general definitions.

0.6 Generalities

- Let $k \in \mathbb{N}^*$. Then we will often confuse - when it is convenient to do it- an element in \mathbb{Z}_k with one of its representants. For example, let $(a_i)_{i \in \mathbb{Z}_k}$ be a family of elements in some vector space E , and $0 \leq m < k/2$ an integer. Then we will write

$$a_i = a_{-i} \quad 1 \leq i \leq m$$

to say that this equality holds for all $i \in \{1 + k\mathbb{Z}, \dots, m + k\mathbb{Z}\} \subset \mathbb{Z}_k$.

- Let us suppose that a vector space E admits some decomposition $E = \oplus_{i \in I} E_i$. Then, for any

vector $v \in E$ we denote by $[v]_{E_i}$ its component in E_i .

- We denote $E^{\mathbb{C}} := E \otimes \mathbb{C}$ the complexification of a real vector space.
- Furthermore let $A \in \text{End}(E)$ be an endomorphism of a finite dimensional real vector space that we suppose to be diagonalisable. Then if its complex spectrum is $\{\lambda_i, i \in I\}$, we will write its eigenspace decomposition in the form $E^{\mathbb{C}} = \oplus_{i \in I} E_i^{\mathbb{C}}$, where $E_i^{\mathbb{C}} = \ker(A - \lambda_i \text{Id})$. Moreover, if $\lambda_i \in \mathbb{R}$ for some $i \in I$, then we set $E_i := E_i^{\mathbb{C}} \cap E$ so that $E_i^{\mathbb{C}} = (E_i)^{\mathbb{C}}$, for these particular $i \in I$.
- We denote by ω_p a p -th primitive root of unity, which will be often chosen equal to $e^{2i\pi/p}$.
- We denote by $C^\infty(M, N)$ the set of smooth maps from a manifold M into a manifold N . Now, let $\pi: N \rightarrow M$ be a surjection, then we denote by $\mathcal{C}(\pi)$ the set of sections of π , i.e. the maps s such that $\pi \circ s = \text{Id}_M$. If there is no risk of confusion we will also use the notation $\mathcal{C}(N)$. Furthermore, let $p: E \rightarrow M$ be a vector bundle. Then we will denote by the same letter p its tensorial extensions: $p: \text{End}(E) \rightarrow M$ and so on.
- Let (M, g) be a Riemannian manifold. Then we denote by $*$ its Hodge operator (and $*_g$ if the metric need to be precised). Let $(E, h) \rightarrow M$ be a Riemannian vector bundle over a manifold M , we denote by $\Sigma(E)$ the bundle of orthogonal almost complex structures in E . In particular if $E = TM$ then we set $\Sigma(M) := \Sigma(TM)$. More generally, let $\mathcal{Z}(\mathbb{R}^n) \subset \text{End}(\mathbb{R}^n)$ be some submanifold defined for any $n \in \mathbb{N}^*$. This allows us to define $\mathcal{Z}(E)$ for any (Riemannian) vector bundle E and we set in particular $\mathcal{Z}(M) = \mathcal{Z}(TM)$ for any (Riemannian) manifold.
- Let $(A, +, \times)$ be an associative \mathbb{K} -algebra over the field \mathbb{K} . Then for any $a \in A$, we set $\text{Com}(a) = \{b \in A | ab = ba\}$ and $\text{Ant}(a) = \{b \in A | ab = -ba\}$.
- w.r.t. : with respect to.

0.7 Almost complex geometry

Let E be a real vector space endowed with a complex structure: $J \in \text{End}(E)$, $J^2 = -\text{Id}$. Then we denote by $E^{1,0}$ and $E^{0,1}$ respectively the eigenspaces of J associated to the eigenvalues $\pm i$ respectively. Then we have the following eigenspace decomposition

$$E^{\mathbb{C}} = E^{1,0} \oplus E^{0,1} \quad (3)$$

and the following equalities

$$\begin{aligned} E^{1,0} &= \ker(J - i\text{Id}) = (J + i\text{Id})E^{\mathbb{C}} \\ E^{0,1} &= \ker(J + i\text{Id}) = (J - i\text{Id})E^{\mathbb{C}} \end{aligned} \quad (4)$$

so that remarking that $(J \pm i\text{Id})iE = (\text{Id} \mp iJ)E = (\text{Id} \mp iJ)JE = (J \pm i\text{Id})E$, we can also write

$$\begin{aligned} E^{1,0} &= (J + i\text{Id})E = (\text{Id} - iJ)E = \{X - iJX, X \in E\} \\ E^{0,1} &= (J - i\text{Id})E = (\text{Id} + iJ)E = \{X + iJX, X \in E\} \end{aligned} \quad (5)$$

In the same way we denote by

$$(E^*)^{\mathbb{C}} = E_{1,0}^* \oplus E_{0,1}^*$$

the decomposition induced on the dual E^* by the complex structure $J^*: \eta \in E^* \rightarrow \eta J \in E^*$. Besides, given a vector $Z \in E^{\mathbb{C}}$, we denote by

$$Z = [Z]^{1,0} + [Z]^{0,1}$$

its decomposition according to (3). Let us remark that

$$[Z]^{1,0} = (\text{Id} - iJ)Z \quad \text{and} \quad [Z]^{0,1} = (\text{Id} + iJ)Z.$$

Moreover, given η a n -form on E , we denote by $\eta^{(p,q)}$ its component in $\Lambda^{p,q}E^*$ according to the decomposition

$$\Lambda^n E^* = \bigoplus_{p+q=n} \Lambda^{p,q} E^*,$$

where $\Lambda^{p,q}E^* = (\Lambda^p E_{1,0}^*) \wedge (\Lambda^q E_{0,1}^*)$. However for 1-forms, we will often prefer the notation $\eta = \eta' + \eta''$, where η' and η'' denote respectively $\eta^{(1,0)}$ and $\eta^{(0,1)}$.

More generally, all what precedes holds naturally when E is a real vector bundle over a manifold M , endowed with a complex structure J .

We will write

$$d = \partial + \bar{\partial}$$

the decomposition of the exterior derivative of differential forms on an almost complex manifold (M, J) , according to the decomposition $TM^{\mathbb{C}} = T^{1,0}M \oplus T^{0,1}M$.

We will denote by $\text{Hol}((M, J^M), (N, J^N)) := \{f \in C^\infty(M, N) \mid df \circ J^M = J^N \circ df\}$ the set of holomorphic maps between two almost complex manifolds (M, J^M) and (N, J^N) .

In this paper, we will use the following definitions.

Definition 0.1 *Let E be a real vector bundle. A f -structure in E is an endomorphism $F \in \mathcal{C}(\text{End}E)$ such that $F^3 + F = 0$. An f -structure on a manifold M is a f -structure in TM .*

An f -structure F in a vector bundle E is determined by its eigenspaces decomposition that we will denote by

$$E^{\mathbb{C}} = E^+ \oplus E^- \oplus E^0$$

where $E^\pm = \ker(F \mp i\text{Id})$ and $E^0 = \ker F$. In particular if $E = TM$, then we will set $T^i M = (TM)^i$, $\forall i \in \{0, \pm 1\}$.

Definition 0.2 *Let (M, F^M) and (N, F^N) be f -manifolds. Then a map $f: (M, F^M) \rightarrow (N, F^N)$ is said to be **f -holomorphic** if it satisfies*

$$df \circ F^M = F^N \circ df$$

Definition 0.3 *Let (M, J^M) be an almost complex manifold and N a manifold with a splitting $TN = \mathcal{H} \oplus \mathcal{V}$. Let us suppose that the subbundle \mathcal{H} is naturally endowed with a complex structure $J^{\mathcal{H}}$. Then we will say that a map $f: (M, J^M) \rightarrow N$ is **\mathcal{H} -holomorphic** if it satisfies the equation*

$$[df]^{\mathcal{H}} \circ J^M = J^{\mathcal{H}} \circ [df]^{\mathcal{H}},$$

where $[df]^{\mathcal{H}}$ is the projection of df on \mathcal{H} along \mathcal{V} . Moreover, if for some reason, \mathcal{H} inherits the name of horizontal subbundle, then we will say that f is **horizontally holomorphic**.

Remark 0.1 Let us remark that a f -structure in a manifold N is equivalent to a splitting $TN = \mathcal{V} \oplus \mathcal{H}$ together with a complex structure $J^{\mathcal{H}}$ on \mathcal{H} .

Definition 0.4 *An affine manifold (N, ∇) is a manifold endowed with a linear connection. An almost complex affine manifold (N, J, ∇) is an almost complex manifold endowed with an almost complex connection: $\nabla J = 0$.*

1 Invariant connections on reductive homogeneous spaces

The references for this section where we recall some results that we will need in this paper, are [40], [43], [13], and to a lesser extent [29] and [1].

1.1 Linear isotropy representation

Let $M = G/H$ be a homogeneous space with G a real Lie group and H a closed subgroup of G . G acts transitively on M in a natural manner which defines a natural representation: $\phi: g \in G \mapsto (\phi_g: x \in M \mapsto g.x) \in \text{Diff}(M)$. Then $\ker \phi$ is the maximal normal subgroup of G contained in H . Further, let us consider the linear isotropy representation:

$$\rho: h \in H \mapsto d\phi_h(x_0) \in GL(T_{x_0}M)$$

where $x_0 = 1.H$ is the reference point in M . Then we have $\ker \rho \supset \ker \phi$. Moreover the linear isotropy representation is faithful (i.e. ρ is injective) if and only if G acts freely on the bundle of linear frames $L(M)$.

We can always suppose without loss of generality that the action of G on M is effective (i.e. $\ker \phi = \{1\}$) but it does not imply in general that the linear isotropy representation is faithful. However if there exists on M a G -invariant linear connection, then the linear isotropy representation is faithful provided that G acts effectively on M . (Indeed, given a manifold M with a linear connection, and $x \in M$, an affine transformation f of M is determined by $(f(x), df(x))$, i.e. f is the identity if and only if it leaves one linear frame fixed).

1.2 Reductive homogeneous space

Let us suppose now that G/H is reductive, i.e. there exists a decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ such that \mathfrak{m} is $\text{Ad}H$ -invariant: $\forall h \in H, \text{Ad}h(\mathfrak{m}) = \mathfrak{m}$. Then the surjective map $\xi \in \mathfrak{g} \mapsto \xi.x_0 \in T_{x_0}M$ has \mathfrak{h} as kernel and so its restriction to \mathfrak{m} is an isomorphism $\mathfrak{m} \cong T_{x_0}M$. This provides an isomorphism of the associated bundle $G \times_H \mathfrak{m}$ with TM by:

$$[g, \xi] \mapsto g.(\xi.x_0) = \text{Ad}g(\xi).x \quad (6)$$

where $x = \pi(g) = g.x_0$.

Moreover, we have a natural inclusion $G \times_H \mathfrak{m} \mapsto G \times_H \mathfrak{g}$ and the associated bundle $G \times_H \mathfrak{g}$ is canonically identified with the trivial bundle $M \times \mathfrak{g}$ via

$$[g, \xi] \mapsto (\pi(g), \text{Ad}g(\xi)). \quad (7)$$

Thus we have an identification of TM with a subbundle $[\mathfrak{m}]$ of $M \times \mathfrak{g}$, which we may view as a \mathfrak{g} -valued 1-form β on M given by:

$$\beta_x(\xi.x) = \text{Ad}g[\text{Ad}g^{-1}(\xi)]_{\mathfrak{m}},$$

where $\pi(g) = x, \xi \in \mathfrak{g}$ and $[\]_{\mathfrak{m}}$ is the projection on \mathfrak{m} along \mathfrak{h} . Equivalently, for all $X \in T_xM$, $\beta(X)$ is the unique element $\xi \in [\mathfrak{m}]_x (= \text{Ad}g(\mathfrak{m}))$, with $\pi(g) = x$ such that $X = \xi.p$, in other words $\beta(X)$ is characterized by

$$\beta(X) \in [\mathfrak{m}]_x \subset \mathfrak{g} \quad \text{and} \quad X = \beta(X).x .$$

In fact, β is nothing but the projection on M of the H -equivariant 1-form, $\theta_{\mathfrak{m}}$, on G , i.e. $\theta_{\mathfrak{m}}$ is the H -equivariant lift of β . Here, $\theta_{\mathfrak{m}}$ is defined as the \mathfrak{m} -component of the left invariant Maurer-Cartan form θ of G . This can be written as follows

$$(\pi^*\beta)_g = \text{Ad}g(\theta_{\mathfrak{m}}) \quad \forall g \in G \quad (8)$$

with $\theta_g(\xi_g) = g^{-1}.\xi_g$ for all $g \in G, \xi_g \in T_gG$.

Notation For any $\text{Ad}H$ -invariant subspace $\mathfrak{l} \subset \mathfrak{m}$, we will denote by $[\mathfrak{l}]$ the subbundle of $[\mathfrak{m}] \subset M \times \mathfrak{g}$ defined by $[\mathfrak{l}]_{g.x_0} = \text{Ad}g(\mathfrak{l})$.

1.3 The (canonical) invariant connection

On a reductive homogeneous space $M = G/H$, the $\text{Ad}(H)$ -invariant summand \mathfrak{m} provides by left translation in G , a G -invariant distribution $\mathcal{H}(\mathfrak{m})$, given by $\mathcal{H}(\mathfrak{m})_g = g.\mathfrak{m}$ which is horizontal for $\pi: G \rightarrow M$ and right H -invariant and thus defines a G -invariant connection in the principal bundle $\pi: G \rightarrow M$. In fact this procedure defines a bijective correspondance between reductive summands \mathfrak{m} and G -invariant connections in $\pi: G \rightarrow M$ (see [40], chap. 2, Th 11.1). Then the corresponding \mathfrak{h} -valued connection 1-form ω on G (of this G -invariant connection) is the \mathfrak{h} -component of the left invariant Maurer-Cartan form of G :

$$\omega = \theta_{\mathfrak{h}}.$$

1.4 Associated covariant derivative

The connection 1-form ω induces a covariant derivative in the associated bundle $G \times_H \mathfrak{m} \cong TM$ and thus a G -invariant covariant derivative ∇^0 in the tangent bundle TM . In particular, we can conclude according to section 1.1 that if G/H is reductive then the linear isotropy representation is faithful (provided that G acts effectively) or equivalently that $\ker \text{Ad}_{\mathfrak{m}} = \ker \rho = \ker \phi$. We will suppose in the following that, without explicit reference to the contrary, the action of G is effective and (thus) the linear isotropy representation is faithful.

One can compute explicitly ∇^0 .

Lemma 1.1 [13]

$$\beta(\nabla_X^0 Y) = X.\beta(Y) - [\beta(X), \beta(Y)], \quad X, Y \in \Gamma(TM).$$

Let us write (locally) $\beta(X) = \text{Ad}U(X_{\mathfrak{m}})$, $\beta(Y) = \text{Ad}U(Y_{\mathfrak{m}})$ where U is a (local) section of π and $X_{\mathfrak{m}}, Y_{\mathfrak{m}} \in C^\infty(M, \mathfrak{m})$ then we have (using the previous lemma)

$$\begin{aligned} \beta(\nabla_X^0 Y) &= \text{Ad}U(dY_{\mathfrak{m}}(X) + [\alpha(X), Y_{\mathfrak{m}}] - [X_{\mathfrak{m}}, Y_{\mathfrak{m}}]) \\ &= \text{Ad}U(dY_{\mathfrak{m}}(X) + [\alpha_{\mathfrak{h}}(X), Y_{\mathfrak{m}}] + [\alpha_{\mathfrak{m}}(X) - X_{\mathfrak{m}}, Y_{\mathfrak{m}}]) \end{aligned}$$

where $\alpha = U^{-1}.dU$. Besides since U is a section of π ($\pi \circ U = \text{Id}$), then pulling back (8) by U , we obtain $\beta = \text{Ad}U(\alpha_{\mathfrak{m}})$ and then $\alpha_{\mathfrak{m}}(X) = X_{\mathfrak{m}}$, so that

$$\beta(\nabla_X^0 Y) = \text{Ad}U(dY_{\mathfrak{m}}(X) + [\alpha_{\mathfrak{h}}(X), Y_{\mathfrak{m}}]) \quad (9)$$

Remark 1.1 We could also say that $X_{\mathfrak{m}}, Y_{\mathfrak{m}}$ are respectively the pullback by U of the H -equivariant lifts \tilde{X}, \tilde{Y} of X, Y (given by $\beta(X_{\pi(g)}) = \text{Ad}g(\tilde{X}(g))$).

Then $\nabla_X^0 Y$ lifts as the \mathfrak{m} -valued H -equivariant map on G :

$$\widetilde{\nabla_X^0 Y} = d\tilde{Y}(\tilde{X}) + [\theta_{\mathfrak{h}}(\tilde{X}), \tilde{Y}]$$

and then taking the U -pullback we obtain the previous result (without using lemma 1.1).

Moreover, we can express ∇^0 in terms of the flat differentiation in the trivial bundle $M \times \mathfrak{g} (\supset [\mathfrak{m}])$. Let us differentiate the equation $Y = \text{Ad}U(Y_{\mathfrak{m}})$ (we use the identification $TM = [\mathfrak{m}] \subset M \times \mathfrak{g}$)

$$dY = \text{Ad}U(dY_{\mathfrak{m}} + [\alpha, Y_{\mathfrak{m}}]) = \text{Ad}U(dY_{\mathfrak{m}} + [\alpha_{\mathfrak{h}}, Y_{\mathfrak{m}}]) + \text{Ad}U([\alpha_{\mathfrak{m}}, Y_{\mathfrak{m}}]) = \nabla^0 Y + [\beta, Y].$$

Finally, we obtain

$$dY = \nabla^0 Y + [\beta, Y] \quad (10)$$

and we recover lemma 1.1.

1.5 G -invariant linear connections in terms of equivariant bilinear maps

Now let us recall the following results about invariant connections on reductive homogeneous spaces.

Theorem 1.1 [40] *Let $\pi_P: P \rightarrow M$ be a K -principal bundle over the reductive homogeneous space $M = G/H$ and suppose that G acts on P as a group of automorphisms and let $u_0 \in P$ be a fixed point in the fibre of $x_0 \in M$ ($\pi_P(u_0) = x_0$). There is a bijective correspondance between the set of G -invariant connections ω in P and the set of linear maps $\Lambda_{\mathfrak{m}}: \mathfrak{m} \rightarrow \mathfrak{k}$ such that*

$$\Lambda_{\mathfrak{m}}(hXh^{-1}) = \lambda(h)\Lambda_{\mathfrak{m}}(X)\lambda(h)^{-1} \quad \text{for } X \in \mathfrak{m} \text{ and } h \in H \quad (11)$$

where $\lambda: H \rightarrow K$ is the morphism defined by $hu_0 = u_0\lambda(h)$ (H stabilizes the fibre $P_{x_0} = u_0.K$). The correspondance is given by

$$\Lambda(X) = \omega_{u_0}(\tilde{X}), \quad \forall X \in \mathfrak{g} \quad (12)$$

where \tilde{X} is the vector field on P induced by X (i.e. $\forall u \in P, \tilde{X}(u) = \frac{d}{dt}|_{t=0} \exp(tX).u$) and $\Lambda: \mathfrak{g} \rightarrow \mathfrak{k}$ is defined by $\Lambda|_{\mathfrak{m}} = \Lambda_{\mathfrak{m}}$ and $\Lambda|_{\mathfrak{h}} = \lambda$ (hence completely determined by $\Lambda_{\mathfrak{m}}$).

Corollary 1.1 *In the previous theorem, let us suppose that P is a K -structure on $M = G/H$, i.e. P is a subbundle of the bundle $L(M)$ of linear frames on M with structure group $K \subset GL(n, \mathbb{R}) = GL(\mathfrak{m})$ (we identify as usual \mathfrak{m} with $T_{x_0}M$ by $\xi \mapsto \xi.x_0$, and $T_{x_0}M$ to \mathbb{R}^n via the linear frame $u_0 \in P \subset L(M)$). Then in terms of the G -invariant covariant derivative ∇ corresponding to ω , the G -invariant linear connection in P , the previous bijective correspondance may be given by*

$$\Lambda(X)(Y) = \nabla_{\tilde{X}}\tilde{Y}$$

where \tilde{X}, \tilde{Y} are any (local) left G -invariant vector fields extending X, Y i.e. there exists a local section, $g: U \subset M \rightarrow G$, of $\pi: G \rightarrow M$, such that $\tilde{X}_x = \text{Ad}_{g(x)}(X).p$, for all $x \in M$.

Remark 1.2 In theorem 1.1, the G -invariant connection in P defined by $\Lambda_{\mathfrak{m}} = 0$ is called the canonical connection (with respect to the decomposition $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$). If we set $P(M, K) = G(G/H, H)$ with group of automorphisms G , the G -invariant connection defined by the horizontal distribution $\mathcal{H}(\mathfrak{m})$ is the canonical connection.

Now, let P be a G -invariant K -structure on $M = G/H$ as in corollary 1.1. Let P' be a G -invariant subbundle of P with structure group $K' \subset K$, then the canonical connection in P' defined by $\Lambda_{\mathfrak{m}} = 0$ is (the restriction of) the canonical connection in P which is itself the restriction to P of the canonical connection in $L(M)$. In particular, if we set $P' = G.u_0$, this is a subbundle of P with group H , which is isomorphic to the bundle $G(G/H, H)$. Then the canonical linear connection in P' corresponds to the invariant connection in $G(G/H, H)$ defined by the distribution $\mathcal{H}(\mathfrak{m})$.

Theorem 1.2 *Let $P \subset L(M)$ be a K -structure on $M = G/H$. Then the canonical linear connection ($\Lambda_{\mathfrak{m}} = 0$) in P defines the covariant derivative ∇^0 in TM (obtained from $\mathcal{H}(\mathfrak{m})$ in the associated bundle $G \times_H \mathfrak{m} \cong TM$). Moreover there is a bijective correspondance between the set of G -invariant linear connections ∇ , on M , determined by a connection in P , and the set of linear maps $\Lambda_{\mathfrak{m}}: \mathfrak{m} \rightarrow \mathfrak{k} \subset \mathfrak{gl}(\mathfrak{m})$ such that*

$$\Lambda_{\mathfrak{m}}(hXh^{-1}) = \text{Ad}_{\mathfrak{m}}(h)\Lambda_{\mathfrak{m}}(X)\text{Ad}_{\mathfrak{m}}(h)^{-1} \quad \forall X \in \mathfrak{m}, \forall h \in H, \quad (13)$$

given by

$$\nabla = \nabla^0 + \bar{\Lambda}_m$$

i.e. $\nabla_X Y = \nabla_X^0 Y + \bar{\Lambda}_m(X)Y$ for any vector fields X, Y on M , where with the help of (13) we extended the $\text{Ad}(H)$ -equivariant map $\Lambda_m: \mathfrak{m} \times \mathfrak{m} \rightarrow \mathfrak{m}$ to the bundle $G \times_H \mathfrak{m} = TM$ to obtain a map $\bar{\Lambda}_m: TM \times TM \rightarrow TM$.

Example 1.1 Let us suppose that M is Riemannian (i.e. $\text{Ad}_m H$ is compact and \mathfrak{m} is endowed with an $\text{Ad}H$ invariant inner product which defines a G -invariant metric on M) and let us take $P = O(M)$ the bundle of orthonormal frames on M , the previous correspondance is between the set of G -invariant metric linear connections and the set of $\text{Ad}(H)$ -equivariant linear maps $\Lambda_m: \mathfrak{m} \rightarrow \mathfrak{so}(\mathfrak{m})$.

In particular the canonical connection ∇^0 is metric (for any G -invariant metric on M).

Theorem 1.3 • G -invariant tensors on the reductive homogeneous space $M = G/H$ (or more generally G -invariant sections of associated bundles) are parallel with respect to the canonical connection.

- The canonical connection is complete (the geodesics are exactly the curves $x_t = \exp(tX).x_0$, for $X \in \mathfrak{m}$).
- Let P be a G -invariant K -structure on $M = G/H$, then the G -invariant connection defined by $\Lambda: \mathfrak{m} \rightarrow \mathfrak{k}$ has the same geodesics as the canonical connection if and only if

$$\Lambda_m(X)X = 0, \quad \forall X \in \mathfrak{m}$$

Theorem 1.4 The torsion tensor T and the curvature tensor R of the G -invariant connection corresponding to Λ_m is given at the origin point x_0 as follows:

1. $T(X, Y) = \Lambda_m(X)Y - \Lambda_m(Y)X - [X, Y]_m$,
2. $R(X, Y) = [\Lambda_m(X), \Lambda_m(Y)] - \Lambda_m([X, Y]_m) - \text{ad}_m([X, Y]_h)$,

for $X, Y \in \mathfrak{m}$.

In particular, for the canonical connection we have $T(X, Y) = -[X, Y]_m$ and $R(X, Y) = -\text{ad}_m([X, Y]_h)$, for $X, Y \in \mathfrak{m}$; moreover we have $\nabla T = 0$, $\nabla R = 0$.

1.6 A Family of connections on the reductive space M

We take in what precede (i.e. in section 1.5) $P = L(M)$. Then let us consider the one parameter family of connections ∇^t , $0 \leq t \leq 1$ defined by

$$\Lambda_m^t(X)Y = t[X, Y]_m, \quad 0 \leq t \leq 1.$$

For $t = 0$, we obtain the canonical connection ∇^0 . Since for any $t \in [0, 1]$, $\Lambda_m^t(X)X = 0$, $\forall X \in \mathfrak{m}$, ∇^t has the same geodesics as ∇^0 and in particular is complete. The torsion tensor is given (at x_0) by

$$T^t(X, Y) = (2t - 1)[X, Y]_m. \quad (14)$$

In particular $\nabla^{\frac{1}{2}}$ is the unique torsion free G -invariant linear connection having the same geodesics as the canonical connection (according to theorems 1.3 and 1.4).

Assume now that M is Riemannian, and let us take $P = O(M)$. Then ∇^t is metric if and only if Λ_m^t takes values in $\mathfrak{k} = \mathfrak{so}(\mathfrak{m})$ which is equivalent (for $t \neq 0$) to say that M is naturally

reductive (which means by definition that $\forall X \in \mathfrak{m}$, $[X, \cdot]_{\mathfrak{m}}$ is skew symmetric). Now (still in the Riemannian case) let us construct a family of linear connections, $\overset{\text{met}}{\nabla}^t$, $0 \leq t \leq 1$, which are always metric:

$$\overset{\text{met}}{\nabla}^t = \nabla^0 + t([\cdot, \cdot]_{\mathfrak{m}} + U^M)$$

where $U^M: TM \oplus TM \rightarrow TM$ is the "natural reductivity term" which is the symmetric bilinear map defined by⁵

$$\langle U^M(X, Y), Z \rangle = \langle [Z, X]_{\mathfrak{m}}, Y \rangle + \langle X, [Z, Y]_{\mathfrak{m}} \rangle \quad (15)$$

for all $X, Y, Z \in \mathfrak{m}$. Since U^M is symmetric, the torsion of $\overset{\text{met}}{\nabla}^t$ is once again given by

$$T^t(X, Y) = (2t - 1)[X, Y]_{\mathfrak{m}}$$

and thus $\overset{\text{met}}{\nabla}^{\frac{1}{2}}$ is torsion free and metric and we recover that $\overset{\text{met}}{\nabla}^{\frac{1}{2}}$ is the Levi-Civita connection

$$\overset{\text{met}}{\nabla}^{\frac{1}{2}} = \nabla^{\text{L.C.}}$$

Obviously if M is naturally reductive then $\overset{\text{met}}{\nabla}^t = \nabla^t$, $\forall t \in [0, 1]$. Moreover if M is (locally) symmetric, i.e. $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}$, then all these connections coincide and are equal to the Levi-Civita connection: $\overset{\text{met}}{\nabla}^t = \nabla^t = \nabla^0 = \nabla^{\text{L.C.}}$.

Remark 1.3 ∇^1 is interesting since it is nothing but the flat differentiation in the trivial bundle $M \times \mathfrak{g}$ followed by the projection onto $[\mathfrak{m}]$ (along $[\mathfrak{h}]$) (see remark 1.1). So this connection is very natural and following [1], we will call it **the anticanonical connection**.

1.7 Differentiation in $\text{End}(T(G/H))$

According to section 1.2, we have

$$\text{End}(T(G/H)) = G \times_H \text{End}(\mathfrak{m}) \subset (G/H) \times \text{End}(\mathfrak{g}),$$

the previous inclusion being given by $[g, A] \mapsto (\pi(g), \text{Ad}_g \circ A \circ \text{Ad}_g^{-1})$ and we embed $\text{End}(\mathfrak{m})$ in \mathfrak{g} by extending any endomorphism of \mathfrak{m} to the corresponding endomorphism of \mathfrak{g} which vanishes on \mathfrak{h} . In other words $\text{End}(T(G/H))$ can be identified to the subbundle $[\text{End}(\mathfrak{m})]$ of the trivial bundle $(G/H) \times \text{End}(\mathfrak{g})$, with fibers $[\text{End}(\mathfrak{m})]_{g.x_0} = \text{End}(\text{Ad}_g(\mathfrak{m})) = \text{Ad}_g(\text{End}(\mathfrak{m}))\text{Ad}_g^{-1} = \text{Ad}_g(\text{End}(\mathfrak{m}) \oplus \{0\})\text{Ad}_g^{-1}$.

Now, let us compute in terms of the Lie algebra setting, the derivative of the inclusion map $\mathfrak{J}: \text{End}(T(G/H)) \rightarrow M \times \text{End}(\mathfrak{g})$ or more concretely the flat derivative in $M \times \text{End}(\mathfrak{g})$ of any section of $\text{End}(T(G/H))$. To do that, we compute the derivative of

$$\tilde{\mathfrak{J}}: (g, A_{\mathfrak{m}}) \in G \times \text{End}(\mathfrak{m}) \mapsto (g.x_0, \text{Ad}_g \circ A_{\mathfrak{m}} \circ \text{Ad}_g^{-1}) \in M \times \text{End}(\mathfrak{g}).$$

We obtain

$$d\tilde{\mathfrak{J}}(g, A_{\mathfrak{m}}) = (\text{Ad}_g(\theta_{\mathfrak{m}}) \cdot \pi(g), \text{Ad}_g(dA_{\mathfrak{m}} + [\text{ad}\theta, A_{\mathfrak{m}}])\text{Ad}_g^{-1}).$$

Next, let us decompose the endomorphisms in \mathfrak{g} into blocs according to the vector space decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$:

$$\text{End}(\mathfrak{g}) = \begin{pmatrix} \text{End}(\mathfrak{h}) & \text{End}(\mathfrak{m}, \mathfrak{h}) \\ \text{End}(\mathfrak{h}, \mathfrak{m}) & \text{End}(\mathfrak{m}) \end{pmatrix}. \quad (16)$$

⁵ U^M is the G -invariant extension of $U^{\mathfrak{m}}: \mathfrak{m} \oplus \mathfrak{m} \rightarrow \mathfrak{m}$, its restriction to $\mathfrak{m} \oplus \mathfrak{m}$.

By regrouping terms, we obtain the following splitting

$$\text{End}(\mathfrak{g}) = \text{End}(\mathfrak{m}) \oplus (\text{End}(\mathfrak{m}, \mathfrak{h}) \oplus \text{End}(\mathfrak{h}, \mathfrak{m}) \oplus \text{End}(\mathfrak{h})),$$

which applied to $d\tilde{\mathcal{J}}(g, A_{\mathfrak{m}})$, gives us the decomposition

$$\begin{aligned} d\tilde{\mathcal{J}}(g, A_{\mathfrak{m}}) &= (0, \text{Ad}g(dA_{\mathfrak{m}} + [\text{ad}_{\mathfrak{m}}\theta_{\mathfrak{h}}, A_{\mathfrak{m}}] + [[\text{ad}_{\mathfrak{m}}\theta_{\mathfrak{m}}]_{\mathfrak{m}}, A_{\mathfrak{m}}]) \text{Ad}g^{-1}) \\ &\quad + (\text{Ad}g(\theta_{\mathfrak{m}}) \cdot \pi(g), \text{Ad}g([\text{ad}_{\mathfrak{m}}\theta_{\mathfrak{m}}]_{\mathfrak{h}} \circ A_{\mathfrak{m}} - A_{\mathfrak{m}} \circ \text{ad}_{\mathfrak{h}}\theta_{\mathfrak{m}}) \text{Ad}g^{-1}). \end{aligned} \quad (17)$$

The first term is in the vertical space $\mathcal{V}_{\tilde{\mathcal{J}}(g, A_{\mathfrak{m}})} = \text{Ad}g(\text{End}(\mathfrak{m}))\text{Ad}g^{-1} = \text{End}(T_{\pi(g)}M)$ and the previous decomposition (17) provides us with a splitting $T\text{End}(M) = \mathcal{V} \oplus \mathcal{H} = \pi_M^*(\text{End}(M)) \oplus \mathcal{H}$, i.e. a connection on $\text{End}(M)$. Let us determine this connection: we see that the projection on the vertical space (along the horizontal space) corresponds to the projection on $[\text{End}(\mathfrak{m})]$ following (16) so that according to remark 1.3, we can conclude that the horizontal distribution \mathcal{H} defines the connection ∇^1 on $\text{End}(TM) = TM^* \otimes TM$.

Remark 1.4 a) We can recover this previous fact directly from the first term of (17) and the definition of ∇^1 . Indeed, first recall that given two linear connections ∇, ∇' on M , we can write $\nabla' = \nabla + F$, where F is a section of $TM^* \otimes \text{End}(TM)$, and then for any section A in $\text{End}(TM)$,

$$\nabla' A = \nabla A + [F, A].$$

Besides, $\nabla^1 = \nabla^0 + [\cdot, \cdot]_{[\mathfrak{m}]}$, and moreover, if we write (locally) $A = (\pi(U), \text{Ad}U \circ A_{\mathfrak{m}} \circ \text{Ad}U^{-1})$ where U is a local section of π and $A_{\mathfrak{m}} \in C^\infty(M, \text{End}(\mathfrak{m}))$, then according to (9),

$$\nabla^0 A = \text{Ad}U(dA_{\mathfrak{m}} + [\text{ad}_{\mathfrak{m}}\alpha_{\mathfrak{h}}, A_{\mathfrak{m}}]), \quad (18)$$

so that we conclude that

$$\nabla^1 A = \text{Ad}U(dA_{\mathfrak{m}} + [\text{ad}_{\mathfrak{m}}\alpha_{\mathfrak{h}}, A_{\mathfrak{m}}] + [[\text{ad}_{\mathfrak{m}}\alpha_{\mathfrak{m}}]_{\mathfrak{m}}, A_{\mathfrak{m}}]) \text{Ad}U^{-1}$$

which is the (pullback of) the first term of (17).

b) Furthermore, if G/H is (locally) symmetric (i.e. $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}$), then $\nabla^{L.C.} = \nabla^0 = \nabla^1$ and in particular

$$\nabla^{L.C.} A = \text{Ad}U(dA_{\mathfrak{m}} + [\text{ad}_{\mathfrak{m}}\alpha_{\mathfrak{h}}, A_{\mathfrak{m}}]). \quad (19)$$

2 m -th elliptic integrable system associated to a k' -symmetric space

2.0.1 Definition of G^τ (even when τ does not integrate in G)

Here, we will extend the notion of subgroup fixed by an automorphism of Lie group to the situation where only a Lie algebra automorphism is provided. Indeed, let $\tau: G \rightarrow G$ be a Lie group automorphism, then usually one can define $G^\tau = \{g \in G \mid \tau(g) = g\}$ the subgroup fixed by τ . Now, we want to extend this definition to the situation where we only have a Lie algebra automorphism, and so that the two definitions coincide when the Lie algebra automorphism integrates in G .

Let \mathfrak{g} be a real Lie algebra and $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism. Then let us denote by

$$\mathfrak{g}_0 = \mathfrak{g}^\tau := \{\xi \in \mathfrak{g} \mid \tau(\xi) = \xi\} \quad (20)$$

the subalgebra of \mathfrak{g} fixed by τ . Let us assume that τ defines in \mathfrak{g} a τ -invariant reductive decomposition

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{n}, \quad [\mathfrak{g}_0, \mathfrak{n}] \subset \mathfrak{n}, \quad \tau(\mathfrak{n}) = \mathfrak{n}.$$

Without loss of generality, we assume that \mathfrak{g}_0 does not contain non-trivial ideal of \mathfrak{g} , i.e. that $\text{ad}_{\mathfrak{n}}: \mathfrak{g}_0 \rightarrow \mathfrak{gl}(\mathfrak{n})$ is injective (the kernel is a τ -invariant ideal of \mathfrak{g} that we factor out). We then have

$$\mathfrak{g}_0 = \{\xi \in \mathfrak{g} \mid \tau|_{\mathfrak{n}} \circ \text{ad}_{\mathfrak{n}} \xi \circ \tau|_{\mathfrak{n}}^{-1} = \text{ad}_{\mathfrak{n}} \xi\} \quad (21)$$

Let G be a Lie group with Lie algebra \mathfrak{g} . Then since \mathfrak{g}_0 satisfies: $\forall \xi \in \mathfrak{g}_0, \text{ad}_{\xi}(\mathfrak{n}) = \mathfrak{n}$ and $\tau|_{\mathfrak{n}} \circ \text{ad}_{\mathfrak{n}} \xi \circ \tau|_{\mathfrak{n}}^{-1} = \text{ad}_{\mathfrak{n}} \xi$, then the (connected) subgroup G_0^0 generated in G by \mathfrak{g}_0 satisfies: $\forall g \in G_0^0, \text{Ad}_g(\mathfrak{n}) = \mathfrak{n}$ and $\tau|_{\mathfrak{n}} \circ \text{Ad}_{\mathfrak{n}} g \circ \tau|_{\mathfrak{n}}^{-1} = \text{Ad}_{\mathfrak{n}} g$.

Now, let us consider the subgroup

$$G_0 = \{g \in G \mid \text{Ad}_g(\mathfrak{n}) = \mathfrak{n} \text{ and } \tau|_{\mathfrak{n}} \circ \text{Ad}_{\mathfrak{n}} g \circ \tau|_{\mathfrak{n}}^{-1} = \text{Ad}_{\mathfrak{n}} g\}.$$

Then G_0 is a closed subgroup of G and $G_0 \supset G_0^0$, so that $\text{Lie } G_0 \supset \mathfrak{g}_0$. Conversely, differentiating the second relation defining G_0 , we obtain, according to (21), $\text{Lie } G_0 \subset \mathfrak{g}_0$ and thus $\text{Lie } G_0 = \mathfrak{g}_0$, and $G_0^0 = (G_0)^0$.

Moreover, without loss of generality, we will suppose that G_0 does not contain non-trivial normal subgroup of G , i.e. that $\text{Ad}_{\mathfrak{n}} G_0 \rightarrow GL(\mathfrak{n})$ is injective (see section 1). Now, we want to prove that if τ integrates in G , then we have $G_0 = G^\tau$, where G^τ is the subgroup fixed by $\tau: G \rightarrow G$. First, we have $\forall g \in G^\tau, \text{Ad}_g(\mathfrak{n}) = \mathfrak{n}$ and $\tau|_{\mathfrak{n}} \circ \text{Ad}_{\mathfrak{n}} g \circ \tau|_{\mathfrak{n}}^{-1} = \text{Ad}_{\mathfrak{n}} g$, thus $G^\tau \subset G_0$. Conversely, $\forall g \in G_0, \text{Ad}_{\mathfrak{n}} g = \tau|_{\mathfrak{n}} \circ \text{Ad}_{\mathfrak{n}} g \circ \tau|_{\mathfrak{n}}^{-1} = \text{Ad}_{\mathfrak{n}} \tau(g)$ and thus $g = \tau(g)$ since $\text{Ad}_{\mathfrak{n}}: G_0 \rightarrow GL(\mathfrak{n})$ is injective. We have proved $G^\tau = G_0$. This allows us to make the following:

Definition 2.1 *Let \mathfrak{g} be a real Lie algebra and $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism, and G a Lie group with Lie algebra \mathfrak{g} . Let us assume that τ defines in \mathfrak{g} a τ -invariant reductive decomposition: $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{n}$. Then we will set*

$$G^\tau := \{g \in G \mid \text{Ad}_g(\mathfrak{n}) = \mathfrak{n} \text{ and } \tau|_{\mathfrak{n}} \circ \text{Ad}_{\mathfrak{n}} g \circ \tau|_{\mathfrak{n}}^{-1} = \text{Ad}_{\mathfrak{n}} g\}.$$

Let us conclude this subsection by some notations:

Notations and conventions In all the paper, when a real Lie algebra \mathfrak{g} and an automorphism

τ will be given, then \mathfrak{g}_0 will denote the Lie subalgebra defined by (20), G will denote a connected Lie group with Lie algebra \mathfrak{g} and $G_0 \subset G$ a closed subgroup such that $(G^\tau)^0 \subset G_0 \subset G^\tau$ (which implies that its Lie algebra is \mathfrak{g}_0).

Moreover, without loss of generality, we will always suppose that \mathfrak{g}_0 does not contain non-trivial ideal of \mathfrak{g} - we will then say that $(\mathfrak{g}, \mathfrak{g}_0)$ is effective - and also suppose that G^τ does not contain non-trivial normal subgroup of G (by factoring out, if needed, by some discrete subgroup of G). Consequently, when τ can be integrated in G , then G^τ will coincide with the subgroup of G fixed by $\tau: G \rightarrow G$.

2.1 Finite order Lie algebra automorphisms

Let \mathfrak{g} be a real Lie algebra and $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of order k' . Then we have the following eigenspace decomposition:

$$\mathfrak{g}^{\mathbb{C}} = \bigoplus_{j \in \mathbb{Z}/k'\mathbb{Z}} \mathfrak{g}_j^{\mathbb{C}}, \quad [\mathfrak{g}_j^{\mathbb{C}}, \mathfrak{g}_l^{\mathbb{C}}] \subset \mathfrak{g}_{j+l}^{\mathbb{C}}$$

where $\mathfrak{g}_j^{\mathbb{C}}$ is the $e^{2ij\pi/k'}$ -eigenspace of τ .

We then have to distinguish two cases.

2.1.1 The even case: $k' = 2k$

Then we have $\mathfrak{g}_0^{\mathbb{C}} = (\mathfrak{g}_0)^{\mathbb{C}}$. Moreover let us remark that

$$\overline{\mathfrak{g}_j^{\mathbb{C}}} = \mathfrak{g}_{-j}^{\mathbb{C}}, \quad \forall j \in \mathbb{Z}/k'\mathbb{Z}. \quad (22)$$

Therefore $\overline{\mathfrak{g}_k^{\mathbb{C}}} = \mathfrak{g}_{-k}^{\mathbb{C}} = \mathfrak{g}_k^{\mathbb{C}}$ so that we can set $\mathfrak{g}_k^{\mathbb{C}} = (\mathfrak{g}_k)^{\mathbb{C}}$ with

$$\mathfrak{g}_k = \{\xi \in \mathfrak{g} \mid \tau(\xi) = -\xi\}.$$

Moreover, owing to (22), we can define \mathfrak{m}_j as the unique real subspace in \mathfrak{g} such that its complexified is given by

$$\mathfrak{m}_j^{\mathbb{C}} = \mathfrak{g}_j^{\mathbb{C}} \oplus \mathfrak{g}_{-j}^{\mathbb{C}} \quad \text{for } j \neq 0, k,$$

and \mathfrak{n} as the unique real subspace such that

$$\mathfrak{n}^{\mathbb{C}} = \bigoplus_{j \in \mathbb{Z}'_k \setminus \{0\}} \mathfrak{g}_j^{\mathbb{C}},$$

that is $\mathfrak{n} = (\bigoplus_{j=1}^{k-1} \mathfrak{m}_j) \oplus \mathfrak{g}_k$. In particular τ defines a τ -invariant reductive decomposition $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{n}$.

Hence the eigenspace decomposition is written:

$$\mathfrak{g}^{\mathbb{C}} = \left(\mathfrak{g}_{-(k-1)}^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_{-1}^{\mathbb{C}} \right) \oplus \mathfrak{g}_0^{\mathbb{C}} \oplus \left(\mathfrak{g}_1^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_{k-1}^{\mathbb{C}} \right) \oplus \mathfrak{g}_k^{\mathbb{C}}$$

so that by grouping

$$\begin{aligned} \mathfrak{g}^{\mathbb{C}} &= \mathfrak{g}_0^{\mathbb{C}} \oplus \mathfrak{g}_k^{\mathbb{C}} \oplus \left[\bigoplus_{j=1}^{k-1} \mathfrak{m}_j^{\mathbb{C}} \right] \\ &= \mathfrak{h}^{\mathbb{C}} \oplus \mathfrak{m}^{\mathbb{C}} \end{aligned}$$

where $\mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_k$ and $\mathfrak{m} = \bigoplus_{j=1}^{k-1} \mathfrak{m}_j$. Considering the automorphism $\sigma = \tau^2$, we have $\mathfrak{h} = \mathfrak{g}^\sigma$ and $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ is the reductive decomposition defined by the order k automorphism σ . Without

loss of generality, and according to our convention applied to \mathfrak{g} and σ , we will suppose in the following that $(\mathfrak{g}, \mathfrak{h})$ is effective i.e. \mathfrak{h} does not contain non trivial ideal of \mathfrak{g} . This implies in particular that $(\mathfrak{g}, \mathfrak{g}_0)$ is also effective.

Now let us integrate our setting: let G be a Lie group with Lie algebra \mathfrak{g} and we choose G_0 such that $(G^\tau)^0 \subset G_0 \subset G^\tau$. Then G/G_0 is a (locally) $2k$ -symmetric space (it is globally $2k$ -symmetric if τ integrates in G) and is in particular a reductive homogeneous space (reductive decomposition $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{n}$).

Moreover since $\sigma = \tau^2$ is an order k automorphism, then for any subgroup H , such that $(G^\sigma)^0 \subset H \subset G^\sigma$, G/H is a (locally) k -symmetric space. In all the following we will always do this choice for H and suppose that $H \supset G_0$ (it is already true up to covering since $\mathfrak{h} \supset \mathfrak{g}_0$) so that $N = G/G_0$ has a structure of associated bundle over $M = G/H$ with fibre H/G_0 : $G/G_0 \cong G \times_H H/G_0$. We can add that on \mathfrak{h} , τ is an involution: $(\tau|_{\mathfrak{h}})^2 = \text{Id}_{\mathfrak{h}}$, whose symmetric decomposition is $\mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_k$, and gives rise to the (locally) symmetric space H/G_0 . The fibre H/G_0 is thus (locally) symmetric (and globally symmetric if the inner automorphism $\text{Int}\tau|_{\mathfrak{m}}$ stabilizes $\text{Ad}_{\mathfrak{m}}H$). Owing to the effectivity of $(\mathfrak{g}, \mathfrak{h})$, we have the following characterization:

$$\mathfrak{g}_0 = \{\xi \in \mathfrak{h} | [\text{ad}_{\mathfrak{m}}\xi, \tau|_{\mathfrak{m}}] = 0\} \quad (23)$$

$$\mathfrak{g}_k = \{\xi \in \mathfrak{h} | \{\text{ad}_{\mathfrak{m}}\xi, \tau|_{\mathfrak{m}}\} = 0\} \quad (24)$$

$\{\}$ being the anticommutator.

Two different types of $2k$ -symmetric spaces. Since $(\tau|_{\mathfrak{m}})^k$ is an involution, there exist two invariant subspaces \mathfrak{m}' and \mathfrak{m}'' , of \mathfrak{m} , each sum of certain \mathfrak{m}_j 's, such that

$$(\tau|_{\mathfrak{m}})^k = -\text{Id}_{\mathfrak{m}'} \oplus \text{Id}_{\mathfrak{m}''}.$$

These subspaces \mathfrak{m}' and \mathfrak{m}'' can be computed easily : $\mathfrak{m}' = \bigoplus_{j=0}^{\lfloor \frac{k-2}{2} \rfloor} \mathfrak{m}_{2j+1}$ and $\mathfrak{m}'' = \bigoplus_{j=1}^{\lfloor \frac{k-1}{2} \rfloor} \mathfrak{m}_{2j}$. In other words, their complexifications are given by

$$\mathfrak{m}'^{\mathbb{C}} = \bigoplus_{\substack{z^k = -1 \\ z \neq -1}} \ker(\tau - z\text{Id}), \quad \mathfrak{m}''^{\mathbb{C}} = \bigoplus_{\substack{z^k = 1 \\ z \neq \pm 1}} \ker(\tau - z\text{Id}).$$

At this stage, there are two possibilities:

- if $\mathfrak{m}'' = 0$ then $(\tau|_{\mathfrak{m}})^k = -\text{Id}_{\mathfrak{m}}$ and $\tau|_{\mathfrak{m}}$ admits eigenvalues only in the set $\{z^k = -1, z \neq -1\}$.
- if $\mathfrak{m}'' \neq 0$ then $(\tau|_{\mathfrak{m}})^k \neq -\text{Id}_{\mathfrak{m}}$ and $\tau|_{\mathfrak{m}}$ admits eigenvalues in each the sets $\{z^k = 1, z \neq \pm 1\}$ and $\{z^k = -1, z \neq -1\}$.

These two cases give rise to two different types of $2k$ -symmetric spaces (see section 3.5)

G -invariant metrics. Now, let us suppose that $M = G/H$ is Riemannian (i.e. $\text{Ad}_{\mathfrak{m}}H$ is compact) then we can choose an $\text{Ad}H$ -invariant inner product on \mathfrak{m} for which $\tau|_{\mathfrak{m}}$ is an isometry⁶. In the next of the paper, we will always choose this kind of inner product on \mathfrak{m} . Therefore, $\tau|_{\mathfrak{m}}$ is an order $2k$ isometry. We will study this kind of endomorphism in section 3.

Moreover, let us remark that if G/H is Riemannian then so is G/G_0 . Further, since the elliptic system we will study in this paper is given in the Lie algebra setting it is useful to know how the fact that G/H is Riemannian can be read in the Lie algebra setting. In fact, under our hypothesis of effectivity, G/H is Riemannian if and only if \mathfrak{h} is compactly embedded in \mathfrak{g} .

⁶See the Appendice, theorem 8.3, for the proof of the existence of such a inner product

2.1.2 the odd case: $k' = 2k + 1$

As in the even case we have $\mathfrak{g}_0^{\mathbb{C}} = (\mathfrak{g}_0)^{\mathbb{C}}$ and $\overline{\mathfrak{g}_j^{\mathbb{C}}} = \mathfrak{g}_{-j}^{\mathbb{C}}, \forall j \in \mathbb{Z}/k'\mathbb{Z}$. Then we obtain the following eigenspace decomposition:

$$\mathfrak{g}^{\mathbb{C}} = (\mathfrak{g}_{-k}^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_{-1}^{\mathbb{C}}) \oplus \mathfrak{g}_0^{\mathbb{C}} \oplus (\mathfrak{g}_1^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_k^{\mathbb{C}}), \quad (25)$$

which provides in particular the following reductive decomposition:

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{m}$$

with $\mathfrak{m} = \bigoplus_{j=1}^k \mathfrak{m}_j$ and \mathfrak{m}_j is the real subspace whose the complexification is $\mathfrak{m}_j^{\mathbb{C}} = \mathfrak{g}_{-j}^{\mathbb{C}} \oplus \mathfrak{g}_j^{\mathbb{C}}$. According to our convention, we suppose that $(\mathfrak{g}, \mathfrak{g}_0)$ is effective.

Then, as in the even case, integrating our setting and choosing G_0 such that $(G^\tau)^0 \subset G_0 \subset G^\tau$, we consider $N = G/G_0$ which is a locally $(2k + 1)$ -symmetric space and in particular a reductive homogeneous space. Moreover, the decomposition (25) gives rises to a splitting $TN^{\mathbb{C}} = T^{1,0}N \oplus T^{0,1}N$ defined by

$$\begin{aligned} TN^{\mathbb{C}} &= (\bigoplus_{j=1}^k [\mathfrak{g}_{-j}^{\mathbb{C}}]) \oplus (\bigoplus_{j=1}^k [\mathfrak{g}_j^{\mathbb{C}}]) \\ &= T^{1,0}N \oplus T^{0,1}N \end{aligned} \quad (26)$$

This splitting defines a canonical almost complex structure on G/G_0 , that we will denote by \underline{J} . Moreover, we have the following characterization

$$\mathfrak{g}_0 = \{\xi \in \mathfrak{g} \mid [\text{ad}_m \xi, \tau_m] = 0\}.$$

Let us suppose that $N = G/G_0$ is Riemannian then the subgroup generated by $\text{Ad}_m G_0$ and τ_m is compact (because $\text{Ad}_m G_0$ is compact and $\tau_m \text{Ad}_m g \tau_m^{-1} = \text{Ad}_m g, \forall g \in G_0$, and τ_m is of finite order). Therefore there exists an $\text{Ad}G_0$ -invariant inner product on \mathfrak{m} for which τ_m is an isometry. In the next of the paper, we will always choose this kind of inner product on \mathfrak{m} (when N is Riemannian).

2.2 Definitions and general properties of the m -th elliptic system.

2.2.1 Definitions

Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an order k' automorphism with $k' \in \mathbb{N}^*$ (if $k' = 1$ then $\tau = \text{Id}$). We use the notations of 2.1. Let us begin by defining some useful notations.

Notation and convention Given $I \subset \mathbb{N}$, we denote by $\prod_{j \in I} \mathfrak{g}_j^{\mathbb{C}}$, the product $\prod_{j \in I} \mathfrak{g}_{j \bmod k'}^{\mathbb{C}}$. In the case $\sum_{j \in I} \mathfrak{g}_{j \bmod k'}^{\mathbb{C}}$ is a direct sum in $\mathfrak{g}^{\mathbb{C}}$, we will identify it with the previous product via the canonical isomorphism

$$(a_j)_{j \in I} \longmapsto \sum_{j \in I} a_j, \quad (27)$$

and we will denote these two subspaces by the same notation $\bigoplus_{j \in I} \mathfrak{g}_j^{\mathbb{C}}$.

Now, let us define the m -th elliptic integrable system associated to a k' -symmetric space, in the sense of Terng [47].

Definition 2.2 Let L be a Riemann surface. The m -th (\mathfrak{g}, τ) -system (with the $(-)$ -convention) on L is the equation for (u_0, \dots, u_m) , $(1, 0)$ -type 1-form on L with values in $\prod_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}$:

$$\begin{cases} \bar{\partial}u_j + \sum_{i=0}^{m-j} [\bar{u}_i \wedge u_{i+j}] = 0 & (S_j), \quad \text{if } 1 \leq j \leq m, \\ \bar{\partial}u_0 + \partial\bar{u}_0 + \sum_{j=0}^m [u_j \wedge \bar{u}_j] = 0 & (S_0) \end{cases} \quad (\text{Syst})$$

It is equivalent to say that the 1-form

$$\alpha_\lambda = \sum_{j=0}^m \lambda^{-j} u_j + \lambda^j \bar{u}_j = \sum_{j=-m}^m \lambda^j \hat{\alpha}_j \quad (28)$$

satisfies the zero curvature equation:

$$d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0, \quad \forall \lambda \in \mathbb{C}^*. \quad (29)$$

Definition 2.3 Let L be a Riemann surface. The m -th (\mathfrak{g}, τ) -system (with the $(+)$ -convention) on L is the equation (Syst) as in definition 2.2 but for (u_0, \dots, u_m) , $(1, 0)$ -type 1-form on L with values in $\prod_{j=0}^m \mathfrak{g}_j^{\mathbb{C}}$.

It is equivalent to say that the 1-form

$$\alpha_\lambda = \sum_{j=0}^m \lambda^j u_j + \lambda^{-j} \bar{u}_j = \sum_{j=-m}^m \lambda^j \hat{\alpha}_j \quad (30)$$

satisfies the zero curvature equation (29).

Remark 2.1 The difference between the two conventions is that in the first one $\alpha'_\lambda = \sum_{j=0}^m \lambda^{-j} u_j$ involves negative powers of λ whereas in the second one α'_λ involves positive powers of λ (in other words $\hat{\alpha}''_{-j} = 0$, for $j \geq 1$ in the first one whereas $\hat{\alpha}''_j = 0$, for $j \geq 1$ in the second one). In fact the second system is the first system associated to τ^{-1} and vice versa.

The first convention is the traditional one: it was used for harmonic maps into symmetric space (see [16]) and by Hélein-Romon [25, 26, 27] for Hamiltonian stationary Lagrangian surfaces in Hermitian symmetric space – first example of second elliptic integrable system associated to a 4-symmetric space. Then this convention was used in [32, 33, 34]. Terng [47], herself, in her definition of the elliptic integrable system uses also this convention. However in [10], it is the second convention which is used.

The $(+)$ -convention is in fact the most natural, as we will see, since in the $(-)$ -convention, there is a minus sign which appears when we pass from the Lie algebra setting to the geometric setting. This is what happens for example when we will associate to the automorphism τ , a almost complex structure in the target space (see also [34], remark 13). Namely, we will consider that the eigenspace spaces \mathfrak{g}_{-j} will define the $(1, 0)$ -part and the subspaces \mathfrak{g}_j the $(0, 1)$ -part of the tangent space of the target. But the $(+)$ -convention leads to several changes in the traditional conventions, like for example in the DPW method [16], we must use the Iwasawa decomposition $\Lambda G_\tau^{\mathbb{C}} = \Lambda G_\tau \cdot \Lambda_{\mathcal{B}}^- G_\tau^{\mathbb{C}}$ instead of $\Lambda G_\tau^{\mathbb{C}} = \Lambda G_\tau \cdot \Lambda_{\mathcal{B}}^+ G_\tau^{\mathbb{C}}$ and in particular the holomorphic potential involves positive power of λ instead of negative one as it is the case traditionally. We decided

⁷instead of $\prod_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}$.

here to continue to perpetuate the tradition as in [34] and to use the first convention. So in the following **when we will speak about the m -th elliptic integrable system, it will be according to the definition 2.2.**

Notation Sometimes, to avoid confusion we will denote (Syst) either by $(\text{Syst}(m, \mathfrak{g}, \tau))$, $(\text{Syst}(m, \tau))$ or simply by $(\text{Syst}(m))$ depending on the context and the needs.

For shortness we will also often say the (m, \mathfrak{g}, τ) -system instead of the m -th (\mathfrak{g}, τ) -system. We will also say the m -th elliptic (integrable) system associated to (the k' -symmetric space) G/G_0 . We will say that a family of 1-forms $(\alpha_\lambda)_{\lambda \in \mathbb{C}^*}$ (denoted by abuse of notation, simply by α_λ) is solution of the (m, \mathfrak{g}, τ) -system (or of (Syst)) if it corresponds to some solution u of this system, according to (28). Therefore α_λ is solution of the (m, \mathfrak{g}, τ) -system if and only if it can be written in the form (28), for some $(1, 0)$ -type 1-form u on L with values in $\prod_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}$, and satisfies the zero curvature equation (29).

It will turn out that the (m, \mathfrak{g}, τ) -system has distinctively different behaviour if k' is even and if k' is odd. Moreover, for every k' there are three different types of behaviour, according to the size of m relatively to k' .

Definition 2.4 *If $k' = 1$, set $m_{k'} = 0$. If $k' > 1$, we set $m_{k'} = \left\lfloor \frac{k'+1}{2} \right\rfloor = \begin{cases} k & \text{if } k' = 2k \\ k+1 & \text{if } k' = 2k+1 \end{cases}$.*

We will say that the m -th (\mathfrak{g}, τ) -system is:

- in the **primitive case** (or that the system is primitive) if $0 \leq m < m_{k'}$,
 - in the **determined case** (or that the system is determined) if $m_{k'} \leq m \leq k' - 1$,
 - and in the **underdetermined case** (or that the system is underdetermined) if $m > k' - 1$.
- Moreover, the determined system of minimal order $m_{k'}$ will be called "the minimal determined system", and the one of maximal order $k' - 1$ will be called "the maximal determined system".

Let us consider the \mathfrak{g} -valued 1-form $\alpha := \alpha_{\lambda=1}$. Then we have $\alpha = \sum_{j=0}^m u_j + \bar{u}_j$ according to (28) which is equivalent to $\alpha' = \sum_{j=0}^m u_j$, since α is \mathfrak{g} -valued.

- In the primitive and determined cases ($m \leq k' - 1$), $\sum_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}$ is a direct sum so that $u = (u_0, \dots, u_m)$ can be identified with $\sum_{j=0}^m u_j = \alpha'$ via (27) and according to our convention. We will then write simply $u = \alpha'$. In particular we have

$$u_j = \alpha'_{-j} \quad \forall j, 0 \leq j \leq m$$

with $\alpha_j := [\alpha]_{\mathfrak{g}_j} \forall j \in \mathbb{Z}/k'\mathbb{Z}$. Hence in the primitive and determined cases the m -th (\mathfrak{g}, τ) -system can be considered as a system on α . Moreover, we can recover α_λ from α and we will speak about the "extended Maurer Cartan form" α_λ which is then associated to α by

$$\alpha_\lambda = \sum_{j=1}^m \lambda^{-j} \alpha'_{-j} + \alpha_0 + \sum_{j=1}^m \lambda^j \alpha'_j$$

according to (28).

- In the underdetermined case, $\sum_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}$ is not a direct sum so that to a given α (coming from some solution α_λ of the m -th (\mathfrak{g}, τ) -system, according to $\alpha = \alpha_{\lambda=1}$) there are a priori many (other) corresponding solutions $u = (u_0, \dots, u_m)$ since

$$\forall j \in \mathbb{Z}/k'\mathbb{Z}, \alpha'_{-j} = \sum_{i \equiv j[k']} u_i.$$

In fact, we will prove that there are effectively an infinity of other solutions α_λ satisfying the condition $\alpha_{\lambda=1} = \alpha$ (see proposition 2.4; see also section 2.5 for a conceptual explanation).

2.2.2 The geometric solution

Convention. Our study, in the present paper, is local therefore we will suppose (when it is necessary to do so) either that L is simply connected or that all lifts (of maps defined on L) and integrations (of 1-forms on L) are made locally. We consider that these considerations are implicit and will not precise these most of the time.

The equations (29) and (28) are invariant by gauge transformations by the group $C^\infty(L, G_0)$:

$$U_0 \cdot \alpha_\lambda = \text{Ad}U_0(\alpha_\lambda) - dU_0 \cdot U_0^{-1}.$$

where $U_0 \in C^\infty(L, G_0)$. This means that if α_λ satisfies (29) then so is $U_0 \cdot \alpha_\lambda$ and if α_λ can be written in the form (28) then so is $U_0 \cdot \alpha_\lambda$. Therefore if α_λ is a solution of (Syst) then so is $U_0 \cdot \alpha_\lambda$. This allows us to define a **geometric solution** of (Syst) as follows:

Definition 2.5 *A map $f: L \rightarrow G/G_0$ is a **geometric solution** of (Syst) if for any (local) lift U , to G , of f , there exists a (local) solution α_λ of (Syst) such that $U^{-1} \cdot dU = \alpha_{\lambda=1}$.*

In other words, we obtain the set of geometric solutions as follows: for each solution α_λ of (Syst), consider the \mathfrak{g} -valued 1-form $\alpha := \alpha_{\lambda=1}$, then integrate it by $U: L \rightarrow G$, $U^{-1} \cdot dU = \alpha$, and finally project U on G/G_0 to obtain the map $f: L \rightarrow G/G_0$.

Now, to simplify the exposition, let us suppose that L is simply connected (until the end of 2.2.2). Then $(\alpha_\lambda)_{\lambda \in \mathbb{C}^*} \mapsto \alpha = \alpha_{\lambda=1}$ is a surjective map from the set of solution of (Syst) to the set of Maurer-Cartan forms of lifts of geometric solutions. According to the discussion at the end of subsection 2.2.1, this map is bijective in the primitive and determined case ($m \leq k' - 1$) and not injective in the underdetermined case ($m > k' - 1$). By quotienting by $C^\infty(L, G_0)$, we obtain a surjective map π_m with the same properties, taking values in the set of geometric solutions.

Let us make more precise all that. We suppose, until the end of this subsection 2.2.2, that the automorphism $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ is fixed (so that the only data which varies in the (m, \mathfrak{g}, τ) -system is the order m). First, let us give an explicit expression of the space $\mathcal{S}(m)$ of solutions α_λ of the system (Syst(m)), i.e. the solutions of the zero curvature equation (29), which satisfies the equality (28) for some (1, 0)-type 1-form u on L with values in $\prod_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}$. To do that, we want to express the condition to be written in the form (28) as a condition on the loop $\alpha_\bullet: \lambda \in S^1 \mapsto \alpha_\lambda \in \mathcal{C}(T^*L \otimes \mathfrak{g})$. The "•" means of course functions on the parameter⁸ $\lambda \in S^1$. More precisely, we will consider α_\bullet as a 1-form on L with values in the loop Lie algebra $C(S^1, \mathfrak{g})$. Then the condition to be written in the form (28) means:

$$(28) \iff (\alpha_\bullet \in \Lambda_m \mathfrak{g}_\tau \quad \text{and} \quad \alpha'_\bullet \in \Lambda^- \mathfrak{g}_\tau^{\mathbb{C}}) \tag{31}$$

where

$$\begin{aligned} \Lambda \mathfrak{g}_\tau &= \{\eta_\bullet: S^1 \rightarrow \mathfrak{g} \mid \eta_{\omega\lambda} = \tau(\eta_\lambda), \forall \lambda \in S^1\} \\ \Lambda_m \mathfrak{g}_\tau &= \{\eta_\bullet \in \Lambda \mathfrak{g}_\tau \mid \eta_\lambda = \sum_{|j| \leq m} \lambda^j \hat{\eta}_j\} \\ \Lambda^- \mathfrak{g}_\tau^{\mathbb{C}} &= \{\eta_\bullet \in \Lambda \mathfrak{g}_\tau^{\mathbb{C}} \mid \eta_\lambda = \sum_{j \leq 0} \lambda^j \hat{\eta}_j\} \end{aligned}$$

and ω is a k' -th root of unity. We refer the reader to [44] for more details about loop groups and their Lie algebras (in particular about the possible choices of topology which makes $\Lambda \mathfrak{g}_\tau$ be

⁸Remark that α_\bullet determines $(\alpha_\lambda)_{\lambda \in \mathbb{C}^*}$, when this latter satisfies (28).

a Banach Lie algebra). Therefore the space of solutions of $(\text{Syst}(m))$ is given by

$$\mathcal{S}(m) = \{\alpha_\bullet \in \mathcal{C}(T^*L \otimes \Lambda_m \mathfrak{g}_\tau) \mid \alpha'_\bullet \in \Lambda^- \mathfrak{g}_\tau^{\mathbb{C}} \text{ and } d\alpha_\bullet + \frac{1}{2}[\alpha_\bullet \wedge \alpha_\bullet] = 0\}. \quad (32)$$

Let us remark that the condition $\alpha'_\bullet \in \Lambda^- \mathfrak{g}_\tau^{\mathbb{C}}$ can be interpreted as a condition of \mathbb{C} -linearity. Indeed, the Banach vector space $\Lambda \mathfrak{g}_\tau / \mathfrak{g}_0$ is naturally endowed with the complex structure defined by the following decomposition

$$(\Lambda \mathfrak{g}_\tau / \mathfrak{g}_0)^{\mathbb{C}} = \Lambda \mathfrak{g}_\tau^{\mathbb{C}} / \mathfrak{g}_0^{\mathbb{C}} = \Lambda_*^- \mathfrak{g}_\tau \oplus \Lambda_*^+ \mathfrak{g}_\tau, \quad (33)$$

where $\Lambda_*^\pm \mathfrak{g}_\tau = \{\eta_\bullet \in \Lambda \mathfrak{g}_\tau^{\mathbb{C}} \mid \eta_\lambda = \sum_{j \geq 0} \lambda^j \hat{\eta}_j\}$. Then the condition $\alpha'_\bullet \in \Lambda^- \mathfrak{g}_\tau^{\mathbb{C}}$ means that $[\alpha'_\bullet]_* : TL \rightarrow (\Lambda \mathfrak{g}_\tau / \mathfrak{g}_0)^{\mathbb{C}}$ is \mathbb{C} -linear, where $[\]_*$ denotes the component in $\Lambda_* \mathfrak{g}_\tau = \{\eta_\bullet \in \Lambda \mathfrak{g}_\tau \mid \eta_\lambda = \sum_{j \neq 0} \lambda^j \hat{\eta}_j\} \cong \Lambda \mathfrak{g}_\tau / \mathfrak{g}_0$.

Now let us integrate our setting. Firstly, let us define the twisted loop group ([44])

$$\Lambda G_\tau = \{U_\bullet : S^1 \rightarrow G \mid U_{\omega\lambda} = \tau(U_\lambda)\}.$$

Then, let us set

$$\begin{aligned} \mathcal{E}^m &= \{U_\bullet : L \rightarrow \Lambda G_\tau \mid U_\lambda(0) = 1, \forall \lambda \in S^1; \alpha_\lambda := U_\lambda^{-1}.dU_\lambda \text{ is a solution of } (\text{Syst}(m))\} \\ \mathcal{E}_1^m &= \{U : L \rightarrow G \mid \exists U_\bullet \in \mathcal{E}^m, U = U_1\} \\ \mathcal{G}_1^m &= \{f : L \rightarrow G/G_0 \text{ geometric solution of } (\text{Syst}(m)), f(0) = 1.G_0\} \\ \mathcal{G}^m &= \{f_\bullet = \pi_{G/G_0} \circ U_\bullet, U_\bullet \in \mathcal{E}^m\} \end{aligned}$$

Remark that because of the gauge invariance: $\mathcal{E}(m).\mathcal{K} \subset \mathcal{E}(m)$ where $\mathcal{K} = C_*^\infty(L, G_0) = \{U \in C^\infty(L, G_0) \mid U(0) = 1\}$, any lift⁹ $U_\bullet : L \rightarrow \Lambda G_\tau$ of an element $f_\bullet \in \mathcal{G}^m$ belongs to \mathcal{E}^m .

Definition 2.6 *An element $f_\bullet \in \mathcal{G}^m$ will be called an extended geometric solution of $(\text{Syst}(m))$.*

The space of geometric solutions is obviously obtained from the space of extended geometric solutions \mathcal{G}^m by the evaluation at $\lambda = 1$. Moreover $\mathcal{S}(m) \simeq \mathcal{E}^m$ is determined by \mathcal{G}_\bullet^m because of the gauge invariance: $\mathcal{E}(m).\mathcal{K} \subset \mathcal{E}(m)$, so that we can write $\mathcal{G}^m = \mathcal{E}(m)/\mathcal{K}$. Consequently, we have also $\mathcal{G}_1^m = \mathcal{E}_1^m/\mathcal{K}$.

Finally, we obtain the following diagram

$$\begin{array}{ccccccc} \mathcal{S}(m) & \xrightarrow[\cong]{\text{int}} & \mathcal{E}^m & \xrightarrow{\pi_{\mathcal{K}}} & \mathcal{E}^m/\mathcal{K} & \xlongequal{\quad} & \mathcal{G}_\bullet^m \\ \text{ev}_1 \downarrow & & \downarrow & & \downarrow & & \pi_m \downarrow \\ \mathcal{S}(m)_1 & \xrightarrow[\cong]{\text{int}} & \mathcal{E}_1^m & \longrightarrow & \mathcal{E}_1^m/\mathcal{K} & \xlongequal{\quad} & \mathcal{G}^m \end{array} .$$

Therefore, the surjective map π_m is bijective for $m \leq k' - 1$ and not injective for $m > k' - 1$ (because so is ev_1). We will need the following definition:

Definition 2.7 *Given a \mathfrak{g} -valued Maurer-Cartan 1-form α on L , we define **the geometric map corresponding to α** , as $f = \pi_{G/G_0} \circ U$, where U integrates α : $U^{-1}.dU = \alpha$, $U(0) = 1$.*

⁹With initial condition $U(0) = 1$.

We have seen that in the primitive and determined cases, we can consider $(\text{Syst}(m))$ as a system on the \mathfrak{g} -valued 1-form $\alpha := \alpha_{\lambda=1}$. Since the Maurer-Cartan equation for α is always contained in $(\text{Syst}(m))$ according to (29), this systems on α is itself equivalent to a system on the geometric map f corresponding to α . This system on f is then a G -invariant elliptic PDE on f of order ≤ 2 .

Let us summarize:

Proposition 2.1 *The natural surjective map $\pi_m: \mathcal{G}^m \rightarrow \mathcal{G}_1^m$ from the set of extended geometric solutions of the (m, \mathfrak{g}, τ) -system into the set of geometric solutions is bijective in the primitive and determined cases ($m \leq k' - 1$) and not injective in the underdetermined case ($m > k' - 1$). Moreover, in the primitive and determined cases, the (m, \mathfrak{g}, τ) -system - which is initially a system on the $\Lambda_m \mathfrak{g}_\tau$ -valued 1-form α_λ - is in fact a system on the 1-form $\alpha := \alpha_{\lambda=1}$, itself equivalent to an elliptic PDE of order ≤ 2 on the corresponding geometric map $f: L \rightarrow G/G_0$.*

Furthermore, let us interpret the \mathbb{C} -linearity of $[\alpha'_\bullet]_*: TL \rightarrow (\Lambda \mathfrak{g}_\tau / \mathfrak{g}_0)^\mathbb{C}$ in terms of the corresponding extended geometric solution $f_\bullet: L \rightarrow \Lambda G_\tau / G_0$, defined by $f_\bullet = \pi_{G/G_0} \circ U_\bullet$ where U_\bullet integrates α_\bullet . Firstly, the complex structure defined in $\Lambda \mathfrak{g}_\tau / \mathfrak{g}_0$ by (33) is $\text{Ad}G_0$ -invariant so that it defines a ΛG_τ -invariant complex structure on the homogeneous space $\Lambda G_\tau / G_0$. Therefore the \mathbb{C} -linearity of $[\alpha'_\bullet]_*$ means exactly that $f_\bullet: L \rightarrow \Lambda G_\tau / G_0$ is holomorphic. Now, let us interpret the condition $\alpha_\bullet \in \Lambda_m \mathfrak{g}_\tau$ in terms of the map f_\bullet . Let us consider the following $\text{Ad}G_0$ -invariant decomposition

$$\Lambda \mathfrak{g}_\tau / \mathfrak{g}_0 = \Lambda_{m*} \mathfrak{g}_\tau \oplus \Lambda_{>m} \mathfrak{g}_\tau$$

where $\Lambda_{m*} \mathfrak{g}_\tau = \Lambda_m \mathfrak{g}_\tau \cap \Lambda_* \mathfrak{g}_\tau$ and $\Lambda_{>m} \mathfrak{g}_\tau = \{\eta_\bullet \in \Lambda \mathfrak{g}_\tau \mid \eta_\lambda = \sum_{|j|>m} \lambda^j \hat{\eta}_j\}$, which gives rise respectively to some ΛG_τ -invariant splitting

$$T(\Lambda G_\tau / G_0) = \mathcal{H}_m^\Lambda \oplus \mathcal{V}_m^\Lambda.$$

Then \mathcal{H}_m^Λ and \mathcal{V}_m^Λ inherit respectively the qualificatifs horizontal and vertical subbundle respectively. Therefore, in the same spirit as [16] (remark 2.5 and proposition 2.6), the equation (32) gives us the following familiar twistorial characterization

Proposition 2.2 *A map $f_\bullet: L \rightarrow \Lambda G_\tau / G_0$ is an extended geometric solution of the (m, \mathfrak{g}, τ) -system if and only if it is holomorphic and horizontal.*

2.2.3 The increasing sequence of spaces of solutions: $(\mathcal{S}(m))_{m \in \mathbb{N}}$

Again, we suppose in all 2.2.3 that the automorphism τ is fixed and that L is simply connected. Then according to the realisation of $(\text{Syst}(m))$ in the forms (29) and (28), we see that any solution of $(\text{Syst}(m))$ is solution of $(\text{Syst}(m'))$ for $m \leq m'$ (take $u_j = 0$ for $m < j \leq m'$). More precisely, $(\text{Syst}(m))$ is a reduction of $(\text{Syst}(m'))$: $(\text{Syst}(m))$ is obtained from $(\text{Syst}(m'))$ by putting $u_j = 0$, $m < j \leq m'$, in $(\text{Syst}(m'))$. In particular, $\mathcal{S}(m) \subset \mathcal{S}(m')$ for $m \leq m'$; so that any solution in the primitive case ($m < m_{k'}$) is solution of any determined system ($m_{k'} \leq m \leq k' - 1$), and any solution of a determined system is solution of any underdetermined system ($m > k' - 1$).

$$\{\text{Primitive case}\} \subset \{\text{determined case}\} \subset \{\text{underdetermined case}\}.$$

Remark 2.2 We have $\pi_{m'}|_{\mathcal{G}^m} = \pi_m$ if $m \leq m'$. In particular, $\pi_{m'}(\mathcal{G}^m) = \mathcal{G}_1^m$. We can set $\mathcal{S}(\infty) = \cup_{m \in \mathbb{N}} \mathcal{S}(m)$, $\mathcal{E}^\infty = \cup_{m \in \mathbb{N}} \mathcal{E}^m$ and $\mathcal{G}^\infty = \cup_{m \in \mathbb{N}} \mathcal{G}^m$. Then we have $\mathcal{G}^\infty = \mathcal{E}^\infty / \mathcal{K}$. Moreover we can define the surjective map $\pi_\infty: \mathcal{G}^\infty \rightarrow \mathcal{G}_1^\infty$ such that $\pi_\infty|_{\mathcal{G}^m} = \pi_m$, $\forall m \in \mathbb{N}$. Then $\pi_\infty|_{\mathcal{G}^m}$ is a bijection onto \mathcal{G}_1^m for each $m \leq k' - 1$.

We can call $\mathcal{S}(\infty)$ the (\mathfrak{g}, τ) -system, and then we can speak about its subsystem of order m , namely $\mathcal{S}(m)$. In particular, we have the following characterization:

$$\mathcal{S}(\infty) = \{\alpha_{\bullet} \in \mathcal{C}(T^*L \otimes \Lambda_{(\infty)}\mathfrak{g}_{\tau}) \mid \alpha'_{\bullet} \in \Lambda^{-}\mathfrak{g}_{\tau} \quad \text{and} \quad d\alpha_{\bullet} + \frac{1}{2}[\alpha_{\bullet} \wedge \alpha_{\bullet}] = 0\}$$

where $\Lambda_{(\infty)}\mathfrak{g}_{\tau} = \cup_{m \in \mathbb{N}} \Lambda_m\mathfrak{g}_{\tau}$.

The primitive and determined cases ($m \leq k' - 1$) Now, let us apply the previous discussion (about the increasing sequence $(\mathcal{S}(m))_{m \in \mathbb{N}}$) to the study of the determined case. Let us recall that in this case, we can consider that the system $(\text{Syst}(m))$ deals only with \mathfrak{g} -valued 1-forms α . Let us also keep in mind that, in this case, we have $\alpha' = u \in \oplus_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}$ (see the discussion in the end of 2.2.1, after definition 2.4). Then we obtain immediately:

Proposition 2.3 *The solutions of a determined system $(\text{Syst}(m))$, $m_{k'} \leq m \leq k' - 1$, are exactly the solutions of the maximal determined system, i.e. $(\text{Syst}(k' - 1))$, which satisfy the holomorphicity conditions:*

$$\alpha''_{-j} = 0, \quad 1 \leq j \leq k' - 1 - m.$$

Moreover, the solutions of a primitive system $(\text{Syst}(m))$, $1 \leq m \leq m_{k'} - 1$, are the solutions of the minimal determined system, i.e. $(\text{Syst}(m_{k'}))$, which satisfy

(i) if $k' = 2k$ is even, the horizontality conditions:

$$\alpha_k = \alpha_{\pm(k-1)} = \dots \alpha_{\pm(m+1)} = 0$$

(ii) if $k' = 2k + 1$ is odd,

- the holomorphicity condition : $\alpha''_{-k} = 0$ if $m = k$,
- the horizontality conditions : $\alpha_{\pm k} = \alpha_{\pm(k-1)} = \dots \alpha_{\pm(m+1)} = 0$ if $m \leq k - 1$.

The non injectivity of π_m in the underdetermined case Now, let us turn ourself to the underdetermined case.

Proposition 2.4 *In the underdetermined case, $m > k' - 1$, the map $\text{ev}_1 : \alpha_{\bullet} \in \mathcal{S}(m) \rightarrow \alpha \in \mathcal{S}(m)_1$ is not injective.*

Proof. It follows from proposition 2.2. Indeed the projection $\Lambda G_{\tau}/G_0 \rightarrow G/G_0$ is a fibration which comes from the natural morphism of group (itself a fibration) $\Lambda G_{\tau} \rightarrow G$. We want to prove that the surjection

$$\pi_m : f_{\bullet} \in \text{Hol}^m(L, \Lambda G_{\tau}/G_0) \mapsto f_1 \in \mathcal{G}_1^m \subset C^{\infty}(L, G/G_0) \quad (34)$$

is non injective, where $\text{Hol}^m(L, \Lambda G_{\tau}/G_0)$ is the set of holomorphic maps such that $df \in \mathcal{H}_m^{\Lambda}$, and $f(0) = 1.G_0$. Then denoting by $(\mathcal{H}_m^{\Lambda})_*$ the ΛG_{τ} -invariant subbundle defined by $\{\eta \in \Lambda_{m*}\mathfrak{g}_{\tau} \mid \eta_1 = 0\}$, we remark that

$$\{f \in \text{Hol}(L, \Lambda G_{\tau}/G_0) \mid df \in (\mathcal{H}_m^{\Lambda})_*\}$$

is sent on the constant map egal to $1.G_0$ by π_m . This proves the non injectivity. In fact, we have more: the map (34) is a submersion and the tangent space of the fibre is (isomorphic to)

$$\{\beta \in \text{Hol}(L, \Lambda_{m*}\mathfrak{g}_{\tau}) \mid \beta_1 = 0\}.$$

This completes the proof. □

2.2.4 The decreasing sequence $(\text{Syst}(m, \tau^p))_{p/k'}$

We will call m -th \mathfrak{g} -system, the m -th $(\mathfrak{g}, \text{Id})$ -system (i.e. $u = (u_0, \dots, u_m)$ takes values in $(\mathfrak{g}^{\mathbb{C}})^{m+1}$, in definition 2.2).

Any solution of the m -th (\mathfrak{g}, τ) -system is solution of the m -th \mathfrak{g} -system. More precisely, the m -th (\mathfrak{g}, τ) -system is the restriction to $\oplus_{j=0}^m \mathfrak{g}_{-j}^{\mathbb{C}}(\tau)$ of the m -th \mathfrak{g} -system.

More generally, for any $p \in \mathbb{N}^*$ such that p divides k' , the m -th (\mathfrak{g}, τ) -system is the m -th (\mathfrak{g}, τ^p) -system restricted to $\oplus_{j=0}^m \mathfrak{g}_{-j}(\tau)$, or equivalently - in terms of $\alpha_\lambda \in \Lambda \mathfrak{g}_{\tau^p}^{\mathbb{C}}$ - restricted to $\Lambda \mathfrak{g}_\tau^{\mathbb{C}}$.

2.3 The minimal determined case

We study here the elliptic system $(\text{Syst}(m))$ in the minimal determined case and by the way its subcase the primitive case. Let us recall again that in this case, we can consider that the system $(\text{Syst}(m))$ deals only with Maurer-cartan forms α and consequently also with geometric maps f . Then we have to translate the equations on α into geometric conditions on f . This is what we will begin to do now.

The minimal determined case splits into two cases.

2.3.1 The even minimal determined case: $k' = 2k$ and $m = k$

Let us recall the following decomposition

$$\mathfrak{g}^{\mathbb{C}} = \left(\mathfrak{g}_{-(k-1)}^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_{-1}^{\mathbb{C}} \right) \oplus \mathfrak{g}_0^{\mathbb{C}} \oplus \left(\mathfrak{g}_1^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_{k-1}^{\mathbb{C}} \right) \oplus \mathfrak{g}_k^{\mathbb{C}}.$$

It is useful for the following to keep in mind that $k = -k \pmod{2k}$.

Proposition 2.5 *The system $(\text{Syst}(k, \tau))$ can be written*

$$\begin{cases} \alpha''_{-j} = 0, \quad 1 \leq j \leq k-1 & (H_j) \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 & (MC) \\ \bar{\partial}\alpha'_{-k} + [\alpha''_0 \wedge \alpha'_{-k}] = 0 & (S_k) \end{cases} . \quad (35)$$

More precisely the equations (S_j) , $0 \leq j \leq k-1$, of $(\text{Syst}(k, \tau))$ are respectively the projection on $\mathfrak{g}_{-j}^{\mathbb{C}}$, $0 \leq j \leq k-1$, of (MC) (owing to the holomorphicity conditions (H_j) given by proposition 2.3).

Proof. The equation (S_j) , for $1 \leq j \leq k-1$, is written in terms of α :

$$\bar{\partial}\alpha'_{-j} + \sum_{i=0}^{k-j} [\alpha''_i \wedge \alpha'_{-j-i}] = 0,$$

according to definition 2.2. Since we have $\alpha''_{-j} = 0$, $1 \leq j \leq k-1$, according to proposition 2.3, this equation is equivalent to

$$d\alpha_{-j} + \sum_{i=0}^{k-j} [\alpha_i \wedge \alpha_{-j-i}] = 0 \quad (36)$$

which is nothing but the projection of (MC) on $\mathfrak{g}_{-j}^{\mathbb{C}}$. Indeed, we have

$$\begin{aligned} \frac{1}{2}[\alpha \wedge \alpha]_{\mathfrak{g}_{-j}^{\mathbb{C}}} &= \frac{1}{2} \left(\sum_{i+l=-j} [\alpha_i'' \wedge \alpha_l'] + \sum_{i+l=-j} [\alpha_i' \wedge \alpha_l''] \right) = \sum_{i+l=-j} [\alpha_i'' \wedge \alpha_l'] = \sum_{i=-k+1}^k [\alpha_i'' \wedge \alpha'_{-j-i}] \\ &= \sum_{i=0}^{k-j} [\alpha_i'' \wedge \alpha'_{-j-i}] \end{aligned}$$

where we have used, in the last line, the holomorphicity conditions: $\alpha_i'' = 0$ if $-k+1 \leq i \leq -1$ and $\alpha'_{-j-i} = 0$ if $k-j \leq i \leq k$.

Now, since α is real (i.e. is \mathfrak{g} -valued), this equation (36) is also equivalent to its complex conjugate i.e. the projection of (MC) on $\mathfrak{g}_j^{\mathbb{C}}$. Moreover, the equation (S_0) is written in terms of α :

$$\bar{\partial}\alpha'_0 + \partial\alpha''_0 + \sum_{j=0}^k [\alpha'_{-j} \wedge \alpha''_j] = 0$$

and moreover, using the holomorphicity conditions, we have

$$\sum_{j=0}^k [\alpha'_{-j} \wedge \alpha''_j] = \sum_{j=0}^{k-1} [\alpha_{-j} \wedge \alpha_j] + \frac{1}{2}[\alpha_{-k} \wedge \alpha_k] = \frac{1}{2}[\alpha \wedge \alpha]_{\mathfrak{g}_0}$$

which proves that (S_0) is equivalent to the projection of (MC) on \mathfrak{g}_0 . Finally, the equation (S_k) is written in terms of α as in (35). This completes the proof. \square

Moreover, always owing to the holomorphicity conditions, the projection of (MC) on \mathfrak{g}_k gives us

$$d\alpha_k + [\alpha_0 \wedge \alpha_k] = 0$$

which is the real part of (S_k). Hence the only new information (in addition to (MC) and (H)) given by the minimal determined elliptic integrable system in the even case is the imaginary part of (S_k):

$$d(*\alpha_k) + [\alpha_0 \wedge (*\alpha_k)] = 0 \quad (E_k)$$

which is as we will see the vertical part of a harmonic map equation. Hence (Syst(k, τ)) is equivalent to

$$\begin{cases} \alpha''_{-j} = 0, \quad 1 \leq j \leq k-1 & (H_j) \\ d(*\alpha_k) + [\alpha_0 \wedge (*\alpha_k)] = 0 & (E_k) \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 & (MC) \end{cases} .$$

Besides (S_k) can be written in the form:

$$[\bar{\partial}(\text{Ad}U(\alpha'_k))]_{\mathfrak{g}_k} = 0.$$

where U integrates α . In term of $f: L \rightarrow G/G_0$, the projection of U , this last equation means (as we will see in section 4)

$$\bar{\partial}^{\nabla^v} \partial^v f = 0$$

where ∇^v is the vertical part of the Levi-Civita connection ∇ on the Riemannian homogeneous space G/G_0 , the vertical and horizontal spaces are defined by $\mathcal{V} = [\mathfrak{g}_k]$ and $\mathcal{H} = [\mathfrak{m}]$ since we

can do the splitting: $T(G/G_0) = [\mathfrak{m}] \oplus [\mathfrak{g}_k]$.

Then the equation (E_k) is equivalent to (see section 4)

$$d^{\nabla^v}(*d^v f) = 0 \iff \tau^v(f) := \text{Tr}_g(\nabla^v d^v f) = 0$$

(for any hermitian metric g on the Riemann surface L). It is easy to see that the equation (E_k) is a vertically harmonic map equation ($\tau^v(f) = 0$) for the canonical connection $\nabla^0 = d + \alpha_0$. In fact, we will see that in the vertical subbundle \mathcal{V} the vertical component of Levi-Civita ∇^v coincides with ∇^0 (see section 4).

The primitive case The m -primitive case is obtained by putting $\alpha_k = \alpha_{\pm(k-1)} = \dots \alpha_{\pm(m+1)} = 0$ in the minimal determined case (35). In particular $\alpha_k = 0$ and (S_k) is trivial so that the only conditions on the geometric map $f: L \rightarrow G/G_0$ (whose existence is guaranteed by (MC)) are the equations (H_j) : $\alpha''_{-j} = \alpha'_j = 0$, $1 \leq j \leq m$, and $\alpha_k = \alpha_{\pm(k-1)} = \dots \alpha_{\pm(m+1)} = 0$, according to proposition 2.3 .

Proposition 2.6 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an order $2k$ automorphism, and an integer $m < k$ then the m -th elliptic integrable system $(\text{Syst}(m, \tau))$ means that the geometric map $f: L \rightarrow G/G_0$ satisfies*

$$\partial f \in \oplus_{j=1}^m [\mathfrak{g}_{-j}^{\mathbb{C}}] \subset T(G/G_0)^{\mathbb{C}}.$$

Proof. Let f be the geometric map corresponding to the Maurer-Cartan form α , that we integrate by $U: L \rightarrow G$, then we have $\partial f = \text{Ad}U(\alpha'_n)$ and α is solution of $(\text{Syst}(m, \tau))$ if and only if

$$\alpha'_n = \alpha'_{-1} + \dots + \alpha'_{-m} \in \oplus_{j=1}^m \mathfrak{g}_{-j}^{\mathbb{C}} \iff \partial f \in \oplus_{j=1}^m [\mathfrak{g}_{-j}^{\mathbb{C}}].$$

This completes the proof. □

Remark 2.3 In particular, in the primitive case f is horizontal ($\alpha_k = 0$). Therefore (S_k) is trivial and (owing to the holomorphicity conditions (H_j) , $1 \leq j \leq k-1$) the free curvature equation (29) is equivalent to (MC) in the primitive case.

Definition 2.8 *Let $m < k$ be an integer. We will call m -primitive map (into the locally $(2k)$ -symmetric space G/G_0) a geometric solution of the system $(\text{Syst}(m, \tau))$.*

Geometric interpretation of the equations (H_j) . For $m < k$, let $F^{[m]}$ be the f -structure on $N = G/G_0$ defined by the following (eigenspace) decomposition:

$$\begin{aligned} TN^{\mathbb{C}} &= (\oplus_{j=1}^m [\mathfrak{g}_{-j}^{\mathbb{C}}]) \oplus (\oplus_{|j|>m} [\mathfrak{g}_j^{\mathbb{C}}]) \oplus (\oplus_{j=1}^m [\mathfrak{g}_j^{\mathbb{C}}]) \\ &= T^+N \oplus T^0N \oplus T^-N \end{aligned} \quad (37)$$

We will set $F = F^{[k-1]}$, and we will call F the canonical f -structure on N . Then according to proposition 2.6 we have

Theorem 2.1 *A map $f: L \rightarrow G/G_0$ is m -primitive if and only if it is $F^{[m]}$ -holomorphic.*

Remark 2.4 The equations (H_j) : $\alpha''_{-j} = 0$, $1 \leq j \leq m$, on a Maurer-Cartan 1-form α means that the corresponding geometric map $f: L \rightarrow G/G_0$ satisfies $\text{pr}_m \circ (df \circ j_L) = F^{[m]} \circ df$ where j_L is the complex structure in L , and $\text{pr}_m: TN \rightarrow \oplus_{j=1}^m [\mathfrak{m}_j]$ is the projection on $\oplus_{j=1}^m [\mathfrak{m}_j]$ along $(\oplus_{j=m+1}^{k-1} [\mathfrak{m}_j]) \oplus [\mathfrak{g}_k]$. This means that the projection $\text{pr}_m \circ df: TL \rightarrow \oplus_{j=1}^m [\mathfrak{m}_j]$ is a morphism

of complex vector bundle. Let us denote $C_m^\infty(L, G/G_0) = \{f \in C^\infty(L, G/G_0) \mid df \in \oplus_{j=1}^m [\mathfrak{m}_j]\}$. Then we have the following equivalences between the Maurer-Cartan 1-form α and its geometric map:

$$(H_j) : \begin{aligned} \alpha \in \mathfrak{g}_0 \oplus (\oplus_{j=1}^m \mathfrak{m}_j) &\iff f \in C_m^\infty(L, G/G_0) \\ \alpha''_{-j} = 0, 1 \leq j \leq m &\iff \text{pr}_m \circ (df \circ j_L) = F^{[m]} \circ df \end{aligned} \quad (38)$$

Then additionning these two equivalences, we recover the equivalence: " α solves $(\text{Syst}(m, \tau))$ " \iff " f is $F^{[m]}$ -holomorphic". Moreover, the equations $\alpha''_{-j} = 0, 1 \leq j \leq k-1$, mean that f is horizontally holomorphic.

Theorem 2.2 *Let α be a \mathfrak{g} -valued 1-form on L and f its geometric map. The following statements are equivalent:*

- (i) $\alpha''_{-j} = 0, 1 \leq j \leq k-1$
- (ii) f is horizontally holomorphic: $(df \circ j_L)^{\mathcal{H}} = F^{[k-1]} \circ df, \mathcal{H} = [\mathfrak{m}]$ being the horizontal space and $F^{[k-1]}|_{\mathcal{H}}$ defining a complex structure on \mathcal{H} .

So that we can conclude:

Theorem 2.3 *The even minimal determined system $(\text{Syst}(k, \tau))$ means that the geometric map f is horizontally holomorphic and vertically harmonic.*

Remark 2.5 We can express what precedes in terms of the projection map $\bar{\pi}_{G/G_0} : \alpha \rightarrow f$ defined as follows. Let \mathcal{MC} be the set of \mathfrak{g} -valued integrable 1-form on L and for $m < k$, \mathcal{MC}^m the subset of interable 1-form taking values in $\mathfrak{g}_0 \oplus (\oplus_{j=1}^m \mathfrak{m}_j)$, then $\bar{\pi}_{G/G_0} : \mathcal{MC} \rightarrow C^\infty(L, G/G_0)$ is defined by:

$$\bar{\pi}_{G/G_0} : \alpha \in \mathcal{MC} \xrightarrow{\text{int}} U \in C_*^\infty(L, G) \xrightarrow{\pi_{G/G_0}} f = \pi_{G/G_0} \circ U \in C^\infty(L, G/G_0).$$

The preceding results can be summarized as follows: for any $m < k$

$$\bar{\pi}_{G/G_0}(\mathcal{MC}^m) = C_m^\infty(L, G/G_0) \quad \text{and} \quad \bar{\pi}_{G/G_0}(\mathcal{S}(m)) = \text{Hol}((L, j), (G/G_0, F^{[m]})),$$

the set of $F^{[m]}$ -holomorphic maps; and the equations $(H_j), 1 \leq j \leq m$, in \mathcal{MC} are transformed by $\bar{\pi}_{G/G_0}$ into the equation $\text{pr}_m \circ (df \circ j_L) = F^{[m]} \circ df$ in $C^\infty(L, G/G_0)$.

2.3.2 The minimal determined odd case

The order of the automorphism τ is odd: $k' = 2k + 1$, and $m = k + 1$. Let us recall the following decomposition

$$\mathfrak{g}^{\mathbb{C}} = (\mathfrak{g}_{-k}^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_{-1}^{\mathbb{C}}) \oplus \mathfrak{g}_0^{\mathbb{C}} \oplus (\mathfrak{g}_1^{\mathbb{C}} \oplus \dots \oplus \mathfrak{g}_k^{\mathbb{C}}).$$

Proposition 2.7 *The equations $(S_j), 0 \leq j \leq k-1$, of $(\text{Syst}(k+1, \tau))$ are respectively the projection on $\mathfrak{g}_{-j}^{\mathbb{C}}, 0 \leq j \leq k-1$, of the Maurer-Cartan equation (MC) (owing to the holomorphicity conditions given by proposition 2.3). Hence the elliptic system $(\text{Syst}(k+1, \tau))$ can be written:*

$$\begin{cases} \alpha''_{-j} = 0, 1 \leq j \leq k-1 & (H_j) \\ \bar{\partial}\alpha'_k + [\alpha''_0 \wedge \alpha'_k] = 0 & (S_{k+1}) \\ \bar{\partial}\alpha'_{-k} + [\alpha''_0 \wedge \alpha'_{-k}] + [\alpha''_1 \wedge \alpha'_k] = 0 & (S_k) \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 & (\text{MC}) \end{cases} \quad (39)$$

Proof. Proceed as for proposition 2.5. □

Then we see that the projection on $\mathfrak{g}_{-k}^{\mathbb{C}}$ of (MC):

$$d\alpha_{-k} + [\alpha_0 \wedge \alpha_{-k}] + [\alpha_1 \wedge \alpha_k] = 0 \quad (40)$$

is nothing but $(S_k) + (\overline{S_{k+1}})$.

Now we have to distinguish two cases.

• **The strictly minimal determined case**¹⁰ Let us suppose that $k \geq 2$, then we have

$$(S_k) \vee (S_{k+1}) \equiv (S_k) + (S_{k+1}) \iff [\bar{\partial}(\text{Ad}U(\alpha'_{\mathfrak{m}_k}))]_{\mathfrak{m}_k} = 0$$

where U integrates α . For the last equivalence, just do the computation:

$$\begin{aligned} [\bar{\partial}(\text{Ad}U(\alpha'_{\mathfrak{m}_k}))]_{\mathfrak{m}_k} &= \bar{\partial}\alpha'_{\mathfrak{m}_k} + [\alpha'' \wedge \alpha'_{\mathfrak{m}_k}]_{\mathfrak{m}_k} \\ &= \bar{\partial}\alpha'_{\mathfrak{m}_k} + [\alpha''_0 \wedge \alpha'_{\mathfrak{m}_k}] + [\alpha''_1 \wedge \alpha'_k] + [\alpha''_{-1} \wedge \alpha'_{-k}] \\ &= (S_k) + (S_{k+1}) \end{aligned}$$

since $\alpha''_{-1} = 0$. Hence we obtain that

$$\begin{aligned} (\text{Syst}) \iff & \begin{cases} \alpha''_{-j} = 0, 1 \leq j \leq k-1 & (H_j) \\ \bar{\partial}\alpha'_k + [\alpha''_0 \wedge \alpha'_k] = 0 & (S_{k+1}) \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 & (\text{MC}) \end{cases} \iff \\ \begin{cases} \alpha''_{-j} = 0, 1 \leq j \leq k-1 & (H_j) \\ \bar{\partial}\alpha'_{-k} + [\alpha''_0 \wedge \alpha'_{-k}] + [\alpha''_1 \wedge \alpha'_k] = 0 & (S_k) \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 & (\text{MC}) \end{cases} & \iff \begin{cases} \alpha''_{-j} = 0, 1 \leq j \leq k-1 & (H_j) \\ [\bar{\partial}(\text{Ad}U(\alpha'_{\mathfrak{m}_k}))]_{\mathfrak{m}_k} = 0 & (S_{\mathfrak{m}_k}) \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 & (\text{MC}) \end{cases} \end{aligned}$$

In terms of the geometric map $f: L \rightarrow G/G_0$, we have according to remark 1.3 (see also section 5) the following geometric interpretation:

$$(S_{\mathfrak{m}_k}) \iff \bar{\partial}^{(\nabla^1)^v} \partial^v f = 0,$$

where the splitting $TN = \mathcal{H} \oplus \mathcal{V}$ is defined by $\mathcal{H} = [\mathfrak{m}']$, $\mathcal{V} = [\mathfrak{m}_k]$ and $\mathfrak{m}' = \bigoplus_{j=1}^{k-1} \mathfrak{m}_j$. Moreover since $2 \text{Re}(S_{\mathfrak{m}_k})$ is

$$d\alpha_{\mathfrak{m}_k} + [\alpha_0 \wedge \alpha_{\mathfrak{m}_k}] + [\alpha_{\mathfrak{m}_1} \wedge \alpha_{\mathfrak{m}_k}]_{\mathfrak{m}_k} = 0$$

which is nothing but $[\text{MC}]_{\mathfrak{m}_k}$ (in the presence of (H)), the projection of (MC) on \mathfrak{m}_k , then the only new information (in addition to (MC) and (H)) given by the determined elliptic integrable system in the odd case is the imaginary part of $(S_{\mathfrak{m}_k})$ which means that f is vertically harmonic (with respect to ∇^1):

$$2 \text{Im}(S_{\mathfrak{m}_k}) : d * \alpha_{\mathfrak{m}_k} + [\alpha_0 \wedge * \alpha_{\mathfrak{m}_k}] + [\alpha_{\mathfrak{m}_1} \wedge * \alpha_{\mathfrak{m}_k}]_{\mathfrak{m}_k} = 0 \iff \tau_1^v(f) := \text{Tr}((\nabla^1)^v d^v f) = 0.$$

¹⁰That is to say the minimal, but non maximal, determined cases.

• **The model case.** Let us suppose that $k = 1$. Let us remark that, in this situation, the determined case reduces to the (model) system $(\text{Syst}(2, \tau))$ which is then simultaneously minimal and maximal. Furthermore, coming back to (39), we have

$$(S_1) \iff [\bar{\partial}(\text{Ad}U(\alpha'_m))]_{\mathfrak{m}}^{1,0} = 0$$

where U integrates α and $[\]^{(1,0)}$ denotes the $(1, 0)$ -component with respect to the canonical complex structure \underline{J} in N defined by the decomposition (26), i.e. in our case $TN^{\mathbb{C}} = [\mathfrak{g}_{-1}^{\mathbb{C}}] \oplus [\mathfrak{g}_1^{\mathbb{C}}]$. Indeed we have

$$\begin{aligned} [\bar{\partial}(\text{Ad}U(\alpha'_m))]_{\mathfrak{m}} &= \bar{\partial}\alpha'_m + [\alpha'' \wedge \alpha'_m]_{\mathfrak{m}} \\ &= \bar{\partial}\alpha'_m + [\alpha''_0 \wedge \alpha'_m] + [\alpha''_m \wedge \alpha'_m]_{\mathfrak{m}} \\ &= \bar{\partial}\alpha'_{-1} + [\alpha''_0 \wedge \alpha'_{-1}] + [\alpha''_1 \wedge \alpha'_1] \\ &\quad + \bar{\partial}\alpha'_1 + [\alpha''_0 \wedge \alpha'_1] + [\alpha''_{-1} \wedge \alpha'_{-1}] \end{aligned}$$

the up term being the $(1, 0)$ -component and the down one, the $(0, 1)$ -component. Then recalling that $(S_1) + \overline{(S_2)}$ is the projection on $\mathfrak{g}_{-1}^{\mathbb{C}}$ of (MC), we obtain that

$$(\text{Syst}) \iff \left\{ \begin{array}{l} (S_2) \\ (\text{MC}) \end{array} \right\} \iff \left\{ \begin{array}{l} (S_1) \\ (\text{MC}) \end{array} \right\} \iff \left\{ \begin{array}{l} [\bar{\partial}(\text{Ad}U(\alpha'_m))]_{\mathfrak{m}}^{1,0} = 0 \quad (S_m^{(1,0)}) \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 \quad (\text{MC}) \end{array} \right.$$

and the only new information (in addition to (MC)) given by the determined elliptic integrable system in this case is $(S_m^{(1,0)})$.

In terms of the geometric map, $f: L \rightarrow G/G_0$, we have according to remark 1.3 the following geometric interpretation:

$$(S_1) \equiv (S_m^{(1,0)}) \iff [\bar{\partial}^{\nabla^1} \partial f]^{1,0} = 0,$$

we will say that f is holomorphically harmonic w.r.t. ∇^1 and \underline{J} (see section 5, definition 5.2, for a precise definition).

In the same way, we also have the following (equivalent) geometric interpretation

$$(S_2) \iff [\bar{\partial}^{\nabla^0} \partial f]^{0,1} = 0,$$

we will say that f is holomorphically harmonic¹¹ w.r.t. ∇^0 and $-\underline{J}$.

Moreover, let us write the equations of the system as a real equation and then write the corresponding geometric equation (which will then take place in TN and not in $TN^{\mathbb{C}}$). We have that the real equation $(S_1) + \overline{(S_2)}$ is the projection on $\mathfrak{g}_{-1}^{\mathbb{C}}$ of (MC). Now, let us write the equation

$$(E_{-1}) \equiv \frac{(S_1) - \overline{(S_2)}}{i} \equiv \left(d * \alpha_{-1} + [\alpha_0 \wedge * \alpha_{-1}] + \frac{1}{2i}[\alpha_1 \wedge \alpha_1] = 0 \right),$$

then taking the sum with its complex conjugate, we obtain

$$\begin{aligned} (E_{-1}) + \overline{(E_{-1})} &\equiv \left(d * \alpha_m + [\alpha_0 \wedge * \alpha_m] + \frac{1}{2i}([\alpha_1 \wedge \alpha_1] - [\alpha_{-1} \wedge \alpha_{-1}]) = 0 \right) \\ &\equiv \left(d * \alpha_m + [\alpha_0 \wedge * \alpha_m] - \frac{1}{2} \underline{J}_0[\alpha_m \wedge \alpha_m] = 0 \right), \end{aligned}$$

¹¹See also section 7, theorem 7.3.

since $\frac{1}{2^i}([\alpha_1 \wedge \alpha_1] - [\alpha_{-1} \wedge \alpha_{-1}]) = \frac{1}{2}[\underline{\mathbf{J}}_0 \alpha_{\mathbf{m}} \wedge \alpha_{\mathbf{m}}] = -\frac{1}{2}\underline{\mathbf{J}}_0[\alpha_{\mathbf{m}} \wedge \alpha_{\mathbf{m}}]$. Then written in terms of the geometric maps $f: L \rightarrow N$, the last equation means

$$\tau^0(f) + \underline{\mathbf{J}}T^0(f) = 0,$$

where $\tau^0(f) = \text{Tr}_g(\nabla^0 df)$ is the tension field of f w.r.t. ∇^0 (and g is some Hermitian metric on L), and $T^0(f) = *(f^*T^0)$, T^0 being the torsion of ∇^0 . As we will see in section 5.1.2, this equation is in fact a general characterization for holomorphically harmonic maps.

The primitive case. The m -primitive case is obtained by putting, in the minimal determined case (39), $\alpha'_k = 0$, if $m = k$, and $\alpha_j = 0$, $m + 1 \leq |j| \leq k$, if $m \leq k - 1$. As in the even case we obtain:

Proposition 2.8 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an order $2k+1$ automorphism, and an interger $m \leq k$ then the m -elliptic integrable system $(\text{Syst}(m, \tau))$ means that the geometric map $f: L \rightarrow G/G_0$ satisfies:*

$$\partial f \in \oplus_{j=1}^m [\mathfrak{g}_{-j}^{\mathbb{C}}] \subset T(G/G_0)^{\mathbb{C}}.$$

Geometric interpretation of the equations (H_j). Recall that $\underline{\mathbf{J}}$ denotes the canonical complex structure on $N = G/G_0$ (see (26)) and set $F^{[m]} := \text{pr}_m \circ \underline{\mathbf{J}} = \underline{\mathbf{J}} \circ \text{pr}_m$ for $m \leq k$, where $\text{pr}_m: TN \rightarrow \oplus_{j=1}^m [\mathfrak{m}_j]$ is the projection on $\oplus_{j=1}^m [\mathfrak{m}_j]$ along $\oplus_{j \geq m+1}^k [\mathfrak{m}_j]$ (remark that $\text{pr}_k = \text{Id}$). Then $F^{[m]}$ is a f -structure on N (remark that $F^{[k]} = \underline{\mathbf{J}}$ is a complex structure). Then we have:

Theorem 2.4 *A map $f: L \rightarrow G/G_0$ is m -primitive if and only if it is $F^{[m]}$ -holomorphic. In particular, f is k -primitive if and only if it is holomorphic (with respect to the canonical complex structure on G/G_0), and thus any m -primitive map is in particular a holomorphic curve in G/G_0 . More precisely, m -primitive maps are exactly the integral holomorphic curves of the complex Pfaff system $\oplus_{j=1}^m [\mathfrak{m}_j] \subset TN$.*

Remark 2.6 The equivalences (38) hold also in the odd case. However for $m = k$, the first equivalence of (38) is trivial: $\alpha \in \mathfrak{g} \iff f \in C^\infty(L, G/G_0)$. There is no restriction (in the form " α takes values in a subspace of \mathfrak{g}^n ") in the highest primitive case.

Theorem 2.5 *Let α be a \mathfrak{g} -valued 1-form on L and f its geometric map. The following statements are equivalent:*

- (i) $\alpha''_{-j} = 0$, $1 \leq j \leq k - 1$
- (ii) f is horizontally holomorphic: $(df \circ j_L)^{\mathcal{H}} = F^{[k-1]} \circ df$, where $\mathcal{H} = [\mathfrak{m}']$ is the horizontal space, and $F^{[k-1]}|_{\mathcal{H}} = \underline{\mathbf{J}}|_{\mathcal{H}}$ defines a complex structure on \mathcal{H} .

We can conclude:

Theorem 2.6 *The odd minimal determined system $(\text{Syst}(k + 1, \tau))$ means that the geometric map f is horizontally holomorphic and vertically harmonic if $k \geq 2$, and if $k = 1$, it means that f is holomorphically harmonic.*

2.4 The maximal determined case

Let us see how can be rewritten the elliptic system in this case in more geometric terms. Let us recall that the determined system can be considered as a system on the 1-form α .

Theorem 2.7 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of odd order $k' = 2k + 1$. Let us set $J_0 = \tau_{\mathfrak{m}}$ and let \underline{J}_0 be the corresponding complex structure on \mathfrak{m} i.e. the value of \underline{J} at the reference point $y_0 = 1.G_0 \in N$ (see equation (26)). Then the associated maximal determined system, $\text{Syst}(k' - 1, \tau)$, is equivalent to*

$$\begin{cases} d * \alpha_{\mathfrak{m}} + [\alpha_0 \wedge * \alpha_{\mathfrak{m}}] + \frac{1}{2} [\underline{J}_0 \alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} + \sum_{1 \leq i < j \leq k} [\underline{J}_0 \alpha_{\mathfrak{m}_i} \wedge \alpha_{\mathfrak{m}_j}]_{\mathfrak{m}_{j-i}} = 0 & (E_{\mathfrak{m}}) \\ d\alpha + \frac{1}{2} [\alpha \wedge \alpha] = 0 & (\text{MC}) \end{cases}$$

Theorem 2.8 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of even order $k' = 2k$. Let us set $J_0 = \tau_{\mathfrak{m}}$ and let \underline{J}_0 be the corresponding complex structure on \mathfrak{m} i.e. the value of $F_{|\mathcal{H}}$ at the reference point $y_0 = 1.G_0 \in N$ (see equation (37)). Then the associated maximal determined system, $\text{Syst}(k' - 1, \tau)$, is equivalent to*

$$\begin{cases} d * \alpha_k + [\alpha_0 \wedge * \alpha_k] + \frac{1}{2} [\underline{J}_0 \alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{g}_k} = 0 & (E_k) \\ d * \alpha_{\mathfrak{m}} + [\alpha_0 \wedge * \alpha_{\mathfrak{m}}] + \frac{1}{2} [\underline{J}_0 \alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} + \sum_{1 \leq i < j \leq k} [\underline{J}_0 \alpha_{\mathfrak{m}_i} \wedge \alpha_{\mathfrak{m}_j}]_{\mathfrak{m}_{j-i}} + [\alpha_k \wedge \underline{J}_0 \alpha_{\mathfrak{m}}] = 0 & (E_{\mathfrak{m}}) \\ d\alpha + \frac{1}{2} [\alpha \wedge \alpha] = 0 & (\text{MC}) \end{cases}$$

Proof of theorems 2.7 and 2.8. This is a straightforward computation. Indeed, it suffices to check that we have

$$\begin{cases} [(\text{MC})]_{\mathfrak{g}_0} = (S_0) \\ [(\text{MC})]_{\mathfrak{g}_{-j}} = (S_j) + \overline{(S_{k'-j})}, \quad 1 \leq j \leq k' - 1 \\ [(E)]_{\mathfrak{g}_{-j}} = \frac{(S_j) - \overline{(S_{k'-j})}}{i}, \quad 1 \leq j \leq k' - 1 \end{cases}$$

where $(E) = (E_{\mathfrak{m}})$ in the odd case and $(E) = (E_{\mathfrak{m}}) + (E_k)$ in the even case. This completes the proof. \square

Remark 2.7 We will see, in sections 5 and 6 respectively, that the equation (E) means that the geometric map f is stringy harmonic.

Adding holomorphicity conditions; the intermediate determined systems.

According to proposition 2.3, we know that the determined system $(\text{Syst}(m))$, $m_{k'} \leq m \leq k' - 1$, is obtained by adding the holomorphicity conditions, $\alpha''_{-j} = 0$, $1 \leq j \leq k' - 1 - m$, to the maximal determined system, whose equations are given respectively by theorems 2.7 and 2.8. Therefore, using the notations defined in the proof of theorems 2.7 and 2.8, we have the following.

Proposition 2.9 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of order k' . Let m be an integer such that $m_{k'} \leq m \leq k' - 1$, and let us set $\underline{m} = k' - 1 - m$. Then the determined system $(\text{Syst}(m))$ is*

equivalent to

$$\left\{ \begin{array}{l} \bar{\partial}\alpha'_{-j} + \sum_{i=0}^{m-j} [\alpha''_i \wedge \alpha'_{-i-j}] = 0 \quad (S_j), \quad \underline{m} + 1 \leq j \leq m \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 \quad (\text{MC}) \\ \alpha''_{-j} = 0 \quad (H_j), \quad 1 \leq j \leq \underline{m} \end{array} \right. \iff \left\{ \begin{array}{l} [(E)]_{\mathfrak{p}} \\ d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 \quad (\text{MC}) \\ \alpha''_{-j} = 0 \quad (H_j), \quad 1 \leq j \leq \underline{m} \end{array} \right.$$

We have set $\mathfrak{m}' = \bigoplus_{j=1}^{\underline{m}} \mathfrak{m}_j$ and $\mathfrak{p} = \bigoplus_{j=\underline{m}+1}^k \mathfrak{m}_j$, where in the even case $\mathfrak{m}_k := \mathfrak{g}_k$. Then $[(E)]_{\mathfrak{p}}$ denotes the projection of (E) on \mathfrak{p} according to the decomposition $\mathfrak{m} = \mathfrak{m}' \oplus \mathfrak{p}$ in the odd case, and $\mathfrak{n} = \mathfrak{m}' \oplus \mathfrak{p}$ in the even case.

Proof. Consider $(\text{Syst}(m))$ as a subsystem of $(\text{Syst}(k' - 1))$. Then we already have seen in the proof of theorems 2.7 and 2.8 that $[(\text{MC})]_{\mathfrak{g}_0} = (S_0)$, $[(\text{MC})]_{\mathfrak{g}_{-j}} = (S_j) + \overline{(S_{k'-j})}$, $1 \leq j \leq m$ and $[(E)]_{\mathfrak{g}_{-j}} = \frac{(S_j) - \overline{(S_{k'-j})}}{i}$, $1 \leq j \leq m$. Then it suffices to check that the holomorphicity conditions, i.e. $\alpha''_{-j} = 0$, $1 \leq j \leq \underline{m}$, imply that the equation (S_j) , for $1 \leq j \leq \underline{m}$, is nothing but the projection on \mathfrak{g}_{-j} of the Maurer-Cartan equation. This can be made by computation, with the same method as in the proof of proposition 2.5. This completes the proof. \square

2.5 The underdetermined case

Here we study the underdetermined case.

Theorem 2.9 *Let us consider an underdetermined system $(\text{Syst}(m, \mathfrak{g}, \tau))$, $m \geq k'$. Let us write*

$$m = qk' + r, \quad 0 \leq r \leq k' - 1$$

the Euclidean division of m by k' . Then let us consider the automorphism in \mathfrak{g}^{q+1} defined by

$$\tilde{\tau}: (a_0, a_1, \dots, a_q) \in \mathfrak{g}^{q+1} \mapsto (a_1, \dots, a_q, \tau(a_0)) \in \mathfrak{g}^{q+1}.$$

Then $\tilde{\tau}$ is of order $(q+1)k'$. Moreover the m -th system associated to (\mathfrak{g}, τ) is in fact equivalent to the m -th system associated to $(\mathfrak{g}^{q+1}, \tilde{\tau})$. More precisely, denoting by $\tilde{\omega}$ a $(q+1)k'$ -th primitive root of unity¹², then the map

$$\alpha_\lambda \mapsto (\alpha_\lambda, \alpha_{\tilde{\omega}\lambda}, \dots, \alpha_{\tilde{\omega}^q\lambda})$$

*is a bijection from the set of solutions of the **underdetermined** (m, \mathfrak{g}, τ) -system into the set of solutions of the **determined** $(m, \mathfrak{g}^{q+1}, \tilde{\tau})$ -system.*

Proof. It suffices to check that α_λ satisfies the properties in equation (32) if and only if $\tilde{\alpha}_\lambda := (\alpha_\lambda, \alpha_{\tilde{\omega}\lambda}, \dots, \alpha_{\tilde{\omega}^q\lambda})$ do so (in the setting $(\mathfrak{g}^{q+1}, \tilde{\tau})$), which is immediate. \square

2.6 Examples

2.6.1 The trivial case: the 0-th elliptic system associated to a Lie group.

We consider the determined system $(\text{Syst}(m, \tau))$ with $\tau = \text{Id}$ and (thus) $k' = 1$ so that $m'_k = m_1 = 0 = k' - 1$. Then the determined system $(\text{Syst}(0, \text{Id}))$ is nothing but the Maurer-Cartan equation for \mathfrak{g} -valued 1-form α (i.e. in other words the "equation" for the trivial geometric map $f: L \rightarrow G/G = \{1\}$).

¹²Chosen such that $\tilde{\omega}^{q+1} = \omega_{k'}$

2.6.2 Even determined case

The first elliptic system associated to a symmetric space [16]. We consider the even determined system $(\text{Syst}(k, \tau))$, with $k = 1$ and τ an involution. Then the horizontal subbundle is trivial $\mathcal{H} = [\mathfrak{m}] = \{0\}$ and $TN = [\mathcal{V}] = [\mathfrak{g}_1]$ so that the horizontal holomorphicity is trivial and vertical harmonicity means harmonicity. Hence the first elliptic system associated to a symmetric space, $(\text{Syst}(1, \tau))$, **is the equation for harmonic maps** $f: L \rightarrow G/G_0$.

The second elliptic system associated to a 4-symmetric space ([34, 10]). Here τ is an order four automorphism and (thus) $k = 2$. Then we consider the even determined system $(\text{Syst}(2, \tau))$. A geometric interpretation in terms of vertically harmonic twistor lift of this system is given in [34] (see also [10]). Let us give some examples.

- Surfaces with holomorphic mean curvature vector in 4-dimensional symmetric spaces ([10]).
- Hamiltonian stationary Lagrangian surfaces in Hermitian symmetric spaces ([27, 25, 26]).
- Surfaces with holomorphic mean curvature vector in 4-dimensional spaces forms ([10]).
- surfaces with anti-holomorphic mean curvature vector in $\mathbb{C}P^2$ ([10]).
- ρ -harmonic surfaces in \mathbb{O} [32].

The second, third and fourth examples are particular cases of the first one (see [10]).

2.6.3 Primitive case

Let us give some examples where the automorphism τ is of order $k' \geq 3$, and $m = 1$.

- Minimal surfaces in $\mathbb{C}P^2$. ($m = 1, k' = 3, N = SU(3)/S(U(1)^3)$). ([7],[8]).
- Minimal Lagrangian surfaces in $\mathbb{C}P^2$. ($m = 1, k' = 6, N = SU(3)/S(U(1)^2)$). ([47]).
- Special Lagrangian surfaces in 4-dim Hermitian symmetric spaces. ($m = 1, k' = 4$). ([27, 25, 26])
- Constant mean curvature (CMC) surfaces in H^3 : their Gauss maps are primitive maps into the unit tangent bundle which is a 4-symmetric space. ([15])
- Affine Toda fields or Toda lattice. ($m = 1, k' \geq 3$). ([7],[11]).
- Non-superminimal (weakly) conformal harmonic maps into S^n . ($m = 1, k' = 2r + 2, N = F^r(S^n)$). ([8]).
- (Weakly) conformal non-isotropic harmonic maps into $\mathbb{C}P^n$. ($m = 1, k' = r + 2, N = F^r(\mathbb{C}P^n)$). ([8]).

$F^r(S^n)$ denotes some bundle of isotropic flags over S^n , and $F^r(\mathbb{C}P^n)$ denotes some bundle of flags over $\mathbb{C}P^n$ (see [8]).

2.6.4 Underdetermined case

First elliptic integrable system associated to a Lie group ([48, 47]). We consider the system $(\text{Syst}(m, \tau))$ with $m = 1$ and $\tau = \text{Id}$. Therefore $k' = 1$ and $m_1 = 0 < m$, and thus this is an underdetermined system. Then $(\text{Syst}(1, \text{Id}))$ **is the equation for harmonic maps into the Lie group** G , $f: L \rightarrow G$.

Second elliptic integrable system associated to the symmetric space $Gr_{3,1}(\mathbb{R}^{n+1,1})$. Constrained Willmore surfaces in S^n corresponds to particular solutions of this system. (See [12], [45]).

3 Finite order isometries and Twistor spaces

The aim of this section is to realise any k' -symmetric space as the subbundle of some bundle of endomorphisms. To make things more precise we need first to introduce some notations.

Let E be an Euclidean space and let us define (for $p \in \mathbb{N}^*$)

$$\mathcal{U}_p(E) = \{A \in SO(E), A^p = \text{Id}, A^i \neq \text{Id} \text{ if } 1 \leq i < p\}$$

$$\mathcal{U}_p^*(E) = \{A \in \mathcal{U}_p(E) | 1 \notin \text{Spect}(A)\}, \quad \mathcal{U}_p^{**}(E) = \{A \in \mathcal{U}_p(E) | \pm 1 \notin \text{Spect}(A)\}.$$

Then for $k \in \mathbb{N}^*$ we set

$$\mathcal{Z}_{2k}(E) = \mathcal{U}_{2k}^{**}(E) \quad \text{and} \quad \mathcal{Z}_{2k+1}(E) = \mathcal{U}_{2k+1}^*(E) = \mathcal{U}_{2k+1}^{**}(E).$$

Let us now explain the general spirit of this section. There are two main ideas which will allow us to provide a twistor interpretation of our elliptic integrable systems. These are the two ideas that we will follow in our exposition.

1st idea. A $2k$ -symmetric homogenous space has a very particular geometry. Then we want to write it more universally by embedding it in a more universal object. To do that, we prove that any $2k$ -symmetric space G/G_0 can be embedded canonically in the twistor bundle $\mathcal{Z}_{2k}(M) \rightarrow M$ of the associated k -symmetric $M = G/H$ via a morphism of bundles over M (recall that $G/G_0 = G \times_H (H/G_0) \rightarrow G/H$ is a bundle over G/H with a symmetric fibre H/G_0). Therefore to any geometric map $f: L \rightarrow G/G_0$ corresponds a map in the twistor space $J: L \rightarrow \mathcal{Z}_{2k}(G/H)$.

2nd idea. Conversely, we write each connected component of $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$ as a $2k$ -symmetric homogeneous space $\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}) = SO(2n)/K$, where K is some Lie subgroup of $SO(2n)$. Indeed for all $J \in \mathcal{Z}_{2k}(\mathbb{R}^{2n})$, the inner automorphism $\text{Int}J$ is of finite order $2k$ and hence defines a $2k$ -symmetric space which is nothing but the connected component of J in $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$. In particular the corresponding Lie algebra automorphism $\text{Ad}J: \mathfrak{so}(2n) \rightarrow \mathfrak{so}(2n)$ gives rise to an eigenspace decomposition $\mathfrak{so}(2n)^\mathbb{C} = \bigoplus_{j \in \mathbb{Z}_{2k}} \mathfrak{so}_j^\mathbb{C}(2n)$ analogous to the decomposition $\mathfrak{g}^\mathbb{C} = \bigoplus_{j \in \mathbb{Z}_{2k}} \mathfrak{g}_j^\mathbb{C}$. Therefore, the geometric properties of a map $f: L \rightarrow G/G_0$ can be translated in terms of the corresponding twistor map $J: L \rightarrow \mathcal{Z}_{2k}(G/H)$.

We are then led to study the space of endomorphisms $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$ which we do in the subsection 3.1. The directing idea of this study is to generalize the space $\Sigma(\mathbb{R}^{2n}) := \mathcal{Z}_4(\mathbb{R}^{2n})$ of (orthogonal) almost complex structure. This is a homogeneous symmetric space of which the tangent space at an element J is the subspace of $\mathfrak{so}(2n)$ which anticommutes with J . We will prove analogous properties in $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$.

In a second time (section 3.6), we will prove the embedding $G/G_0 \hookrightarrow \mathcal{Z}_{2k}(G/H)$.

Even if we will use the twistor space and the embedding described above especially in the even case, however we will sometimes need that in the odd case $k' = 2k + 1$. Moreover, it will also happen in our study that we will need to consider the spaces $(\mathcal{Z}_{2k}(M))^j = \{J^j, J \in \mathcal{Z}_{2k}(M)\}$ which are in fact general twistor spaces $\mathcal{U}_p(M)$. Therefore, we will also study these spaces. Nevertheless, the even case, $\mathcal{Z}_{2k}(M)$, is the most rich and most complex. Indeed it satisfies all the properties which hold in the general case $\mathcal{U}_p(M)$ (in particular in the odd case $\mathcal{Z}_{2k+1}(M)$), so that the study of $\mathcal{Z}_{2k}(M)$ contains the one of $\mathcal{U}_p(M)$. Moreover, the even case $\mathcal{Z}_{2k}(M)$ contains special properties and has additional behaviors that need to be studied separately. This is why we will concentrate our study on this case, in a first time. Then in a second time we will precise which properties hold in general and which ones are particular to the even case.

3.1 Isometries of order $2k$ with no eigenvalues $= \pm 1$

In this section 3.1, we will study of $\mathcal{Z}_{2k}(E)$, where again E is an Euclidean space. For each $A \in \mathcal{Z}_{2k}(E)$, we have the following eigenspace decomposition:

$$E^{\mathbb{C}} = \bigoplus_{j=1}^{k-1} \left(E_A(\omega_{2k}^j) \oplus E_A(\omega_{2k}^{-j}) \right)$$

with $E_A(\lambda) = \ker(A - \lambda \text{Id})$ and $\omega_{2k} = e^{i\pi/k}$.

Let us set $\mathfrak{m}_j^{\mathbb{C}} = E_A(\omega_{2k}^j) \oplus E_A(\omega_{2k}^{-j})$ for $j \geq 0$. Then we have $\dim_{\mathbb{R}} \mathfrak{m}_j = \frac{1}{2} \left(\dim_{\mathbb{R}} E_A(\omega_{2k}^j) + \dim_{\mathbb{R}} E_A(\omega_{2k}^{-j}) \right) = \dim_{\mathbb{R}} E_A(\omega_{2k}^j) = 2 \dim_{\mathbb{C}} E_A(\omega_{2k}^j)$. Hence $\dim_{\mathbb{R}} \mathfrak{m}_j$ is even and hence we will suppose now that $E = \mathbb{R}^{2n}$ (in all the section 3.1).

Example 3.1 We have $\mathcal{Z}_2(E) = \emptyset$, and $\mathcal{Z}_4(E) = \Sigma(E)$ the set of almost complex structures in E .

Situation in the Euclidean plane Here $E = \mathbb{R}^2$, and any element of $A \in \mathcal{Z}_{2k}(E)$ is written in the form $A = R(\frac{l\pi}{k})$, with $(l, 2k) = 1$, $R(\theta)$ being the rotation of angle $\theta \in \mathbb{R}/2\pi\mathbb{Z}$. In other words, considered as a complex number, A is a primitive $(2k)$ -th root of the unity. Hence $\text{card}(\mathcal{Z}_{2k}(\mathbb{R}^2)) = \phi(2k)$, ϕ being the Euler phi-function.

3.1.1 The set of connected components in the general case

Theorem 3.1 $\pi_0(\mathcal{Z}_{2k}(\mathbb{R}^{2n}))$, the set of connected components of $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$, is (in one to one correspondance with):

$$X_{2k} := \left\{ (\varepsilon, p) \in \mathbb{Z}_2 \times \mathbb{N}^{k-1} \left| \sum_{j=1}^{k-1} p_j = n \text{ and } \text{lcm} \left(\left\{ \frac{2k}{2k \wedge j}, p_j \neq 0 \right\} \right) = 2k \right. \right\}$$

Proof. Let $A \in \mathcal{Z}_{2k}(\mathbb{R}^{2n})$, then $A|_{\mathfrak{m}_j^{\mathbb{C}}} = \omega_{2k}^j \text{Id}_{E_A(\omega_{2k}^j)} \oplus \omega_{2k}^{-j} \text{Id}_{E_A(\omega_{2k}^{-j})}$. We choose an orientation on each \mathfrak{m}_j (such that the induced orientation on $\bigoplus_{l=1}^{k-1} \mathfrak{m}_j$ is the one of \mathbb{R}^{2n}). Then there exist oriented planes P_j^l such that $\mathfrak{m}_j = \bigoplus_{l=1}^{p_j} P_j^l$ (sum of non oriented spaces), where $p_j = \frac{\dim \mathfrak{m}_j}{2}$, and

$$A|_{\mathfrak{m}_j} = \bigoplus_{l=1}^{p_j} R_{P_j^l}(\theta_j)$$

where $R_{P_j^l}(\theta_j)$ is the rotation on P_j^l of angle $\theta_j = \frac{j\pi}{k}$. Let ε_j be the sign of the orientation¹³ of $\bigoplus_{l=1}^{p_j} P_j^l$ (sum of oriented spaces) w.r.t. the one of \mathfrak{m}_j . Now let us consider the map

$$f: A \in \mathcal{Z}_{2k}(\mathbb{R}^{2n}) \mapsto (\prod_{j=1}^{k-1} \varepsilon_j, (p_j)_{1 \leq j \leq k-1}) \in X_{2k}.$$

Then it is a continuous¹⁴ surjection and $f^{-1}(\{(\varepsilon, p)\})$ is an $SO(2n)$ -orbit of the $SO(2n)$ -adjoint action on $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$. This completes the proof. \square

Remark 3.1 Let P be an oriented plane and $A = R_P(\theta) \setminus \{\pm \text{Id}_P\}$ a rotation of this plane. Then we can define $\varepsilon_P(A) := \text{sign} \det_P(x, Ax)$ which is well defined because independent of $x \in P$. We have set $\det_P := \det_{(e_1, e_2)}$, for any direct orthonormal basis (e_1, e_2) of P .

¹³see remark 3.1

¹⁴By lower semicontinuity of the rank, and the fact that $\sum_{l=1}^{k-1} p_j$ is constant.

Now, let E be an oriented Euclidean vector space of dimension $2n$ and let $E = \bigoplus_{l=1}^q P_l^0$ be a decomposition of E as a sum of oriented planes (endowed with the orientation induced by E). Let $A \in \bigoplus_{l=1}^q R_{P_l^0}(\theta_l) \in SO(E) \setminus \{\bigoplus_{l=1}^q (\pm \text{Id}_{P_l^0})\}$. We can define

$$\varepsilon(A) = \varepsilon_{P_1^0}(A|_{P_1^0}) \times \cdots \times \varepsilon_{P_q^0}(A|_{P_q^0}) = \text{sign} \det_{\mathcal{B}}(x_1, Ax_1, \dots, x_q, Ax_q)$$

where \mathcal{B} is any direct orthonormal basis of E , and $x_l \in P_l^0$.

In the other hand let P_l be the plane P_l^0 endowed with the unique orientation such that $\theta_l \in]0, \pi[$. Then we have

$$\varepsilon(A) = \text{orient}_E(\bigoplus_{l=1}^q P_l),$$

the right hand side meaning the sign of the orientation of $\bigoplus_{l=1}^q P_l$ w.r.t. to the one of E .

Each connected component is a $SO(2n)$ -orbit and thus is compact, and consequently closed. Hence its complementary which is an finite union of closed subset is closed: each connected component is an open and closed submanifold of $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$ (which is itself a compact submanifold in $SO(2n)$).

Definition 3.1 We will denote by $\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$ (and sometimes only by \mathcal{Z}_{2k}^α) the connected component $f^{-1}(\{\alpha\})$, for $\alpha = (\varepsilon, \underline{p}) \in X_{2k}$. We define

$$\begin{aligned} \mathcal{Z}_{2k}^0(\mathbb{R}^{2n}) &= \{A \in \mathcal{Z}_{2k}(\mathbb{R}^{2n}) | A^k = -\text{Id}\} = \bigsqcup_{\{\alpha | \forall j, p_{2j}=0\}} \mathcal{Z}_{2k}^\alpha \\ \mathcal{Z}_{2k}^*(\mathbb{R}^{2n}) &= \{A \in \mathcal{Z}_{2k}(\mathbb{R}^{2n}) | A^k \neq -\text{Id}\} = \bigsqcup_{\{\alpha | \exists j, p_{2j} \neq 0\}} \mathcal{Z}_{2k}^\alpha \end{aligned}$$

$\mathcal{Z}_{2k}^0(\mathbb{R}^{2n})$ is the union of **order k components** in $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$, and $\mathcal{Z}_{2k}^*(\mathbb{R}^{2n})$ is the union of **order $2k$ components** in $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$ (see below for the meaning of this terms).

In the following we will denote by $\mathcal{Z}_{2k}^a(\mathbb{R}^{2n})$, for $a \in \{0, *\}$, any of the two spaces $\mathcal{Z}_{2k}^0(\mathbb{R}^{2n})$ and $\mathcal{Z}_{2k}^*(\mathbb{R}^{2n})$, and r the order of these two spaces i.e. $r = \begin{cases} 2k & \text{in } \mathcal{Z}_{2k}^*(\mathbb{R}^{2n}) \\ k & \text{in } \mathcal{Z}_{2k}^0(\mathbb{R}^{2n}) \end{cases}$. r is in fact the order of $\text{Ad}J$, for $J \in \mathcal{Z}_{2k}(\mathbb{R}^{2n})$ (see 3.1.2 below). Let us compute the tangent space of $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$: $\forall J \in \mathcal{Z}_{2k}(\mathbb{R}^{2n})$,

$$T_J \mathcal{Z}_{2k}(\mathbb{R}^{2n}) = \left\{ A \in J.\mathfrak{so}(2n) \left| \sum_{p+l=2k-1} J^p A J^l = 0 \right. \right\} \quad (41)$$

and for $J \in \mathcal{Z}_{2k}^0(\mathbb{R}^{2n})$, we have in addition

$$T_J \mathcal{Z}_{2k}(\mathbb{R}^{2n}) = T_J \mathcal{Z}_{2k}^0(\mathbb{R}^{2n}) = \left\{ A \in J.\mathfrak{so}(2n) \left| \sum_{p+l=k-1} J^p A J^l = 0 \right. \right\} \quad (42)$$

It could seem strange that the two expressions (41) and (42) are equal for $J \in \mathcal{Z}_{2k}^0(\mathbb{R}^{2n})$, but as we will see below, it comes from the fact that the "even" eigenspaces of $\text{Ad}J$ vanish, for $J \in \mathcal{Z}_{2k}^0(\mathbb{R}^{2n})$, which leads to this last equality (which is in general an inclusion " \supset ")

Example 3.2 If $k = 2$, then $X_{2k} = \{\pm 1\} = \mathbb{Z}_2$ and $\mathcal{Z}_4(\mathbb{R}^{2n}) = \mathcal{Z}_4^0(\mathbb{R}^{2n}) = \Sigma(\mathbb{R}^{2n}) = \{J \in SO(\mathbb{R}^{2n}) | J^2 = -\text{Id}\} = \Sigma^+(\mathbb{R}^{2n}) \sqcup \Sigma^-(\mathbb{R}^{2n})$ (resp. the positive and negative components of $\Sigma(\mathbb{R}^{2n})$), whereas $\mathcal{Z}_4^*(\mathbb{R}^{2n}) = \emptyset$.

3.1.2 Study of $\text{Ad}J$, for $J \in \mathcal{Z}_{2k}^a(\mathbb{R}^{2n})$

Let $J \in \mathcal{Z}_{2k}^a(\mathbb{R}^{2n})$. $\text{Ad}J$ is then an order r automorphism of $\text{End}(\mathbb{R}^{2n})$ (since $(\text{Ad}J)^p = \text{Id} \Leftrightarrow J^p = \pm \text{Id}$) thus we have the following eigenspaces decomposition:

$$\text{End}(\mathbb{R}^{2n})^{\mathbb{C}} = \bigoplus_{j \in \mathbb{Z}/r\mathbb{Z}} \ker(\text{Ad}J - \omega_r^j \text{Id})$$

with $\omega_r = e^{2i\pi/r}$. Let us set

$$\mathcal{A}_j^{\mathbb{C}}(J) = \ker(\text{Ad}J - \omega_r^j \text{Id}).$$

Then $\mathcal{A}_0(J) = \text{Com}(J) := \{A \in \text{End}(\mathbb{R}^{2n}) \mid [A, J] = 0\}$ and for $j \neq 0$ we have: $\forall A \in \mathcal{A}_j^{\mathbb{C}}$,

$$\sum_{l+p=r-1} J^l A J^p = \sum_{l+p=r-1} (\omega_r^j)^l A J^{p+l} = \left[\sum_0^{l-1} (\omega_r^j)^l \right] J^{r-1} = 0.$$

Hence

$$\bigoplus_{j \in \mathbb{Z}/r\mathbb{Z} \setminus \{0\}} \mathcal{A}_j^{\mathbb{C}}(J) \subset \ker \left(\sum_{l+p=r-1} L(J^l) \circ R(J^p) \right)$$

where L, R denotes respectively the left and right multiplications. This inclusion is in fact an equality. Indeed, let $A \in \text{End}(\mathbb{R}^{2n})^{\mathbb{C}}$, then $A = \sum_{j=0}^{r-1} A_j$, with $A_j \in \mathcal{A}_j^{\mathbb{C}}(J)$, thus $\sum_{j=0}^{r-1} J^l A J^{r-1-l} = r A_0 J^{r-1} + 0 = r A_0 J^{r-1}$ which vanishes if and only if $A_0 = 0$. This proves:

Proposition 3.1 *The following equality holds*

$$\bigoplus_{j \in \mathbb{Z}/r\mathbb{Z} \setminus \{0\}} \mathcal{A}_j^{\mathbb{C}}(J) = \ker \left(\sum_{l+p=r-1} L(J^l) \circ R(J^p) \right).$$

Now, let us restrict ourself to $J \cdot \mathfrak{so}(2n)$, resp. to $\mathfrak{so}(2n)$, (which does not change the order of $\text{Ad}J|_{J \cdot \mathfrak{so}(2n)}$, resp. $\text{Ad}J|_{\mathfrak{so}(2n)}$) and set¹⁵

$$\mathcal{B}_j^{\mathbb{C}}(J) = \mathcal{A}_j^{\mathbb{C}}(J) \cap (J \cdot \mathfrak{so}(2n))^{\mathbb{C}}, \quad \text{resp.} \quad \mathfrak{so}_j^{\mathbb{C}}(J) = \mathcal{A}_j^{\mathbb{C}}(J) \cap \mathfrak{so}(2n)^{\mathbb{C}}.$$

Then we have, according to (41)-(42),

Proposition 3.2 *The tangent space of $\mathcal{Z}_{2k}^a(\mathbb{R}^{2n})$ at one element J is given by*

$$T_J \mathcal{Z}_{2k}^a(\mathbb{R}^{2n}) = \left(\bigoplus_{j=1}^{r-1} \mathcal{B}_j^{\mathbb{C}}(J) \right) \cap \text{End}(\mathbb{R}^{2n}). \quad (43)$$

The inner automorphism¹⁶ $T = \text{Int}J|_{\mathfrak{so}(2n)}$ gives rise to the r -symmetric space $SO(2n)/\mathbb{U}_0(J)$, where $\mathbb{U}_0(J) = SO(2n)^T = \text{Com}(J) \cap SO(2n)$, which is nothing but the connected component \mathcal{Z}_{2k}^a of J (which is also the orbit $SO(2n) \cdot J = \text{Int}(SO(2n))(J)$):

$$\mathcal{Z}_{2k}^a(\mathbb{R}^{2n}) = SO(2n)/\mathbb{U}_0(J).$$

¹⁵ $\mathcal{B}_j^{\mathbb{C}}(J)$ is stable by $\text{Ad}J$ and we have $\mathcal{B}_j^{\mathbb{C}}(J) = J \cdot \mathfrak{so}_j^{\mathbb{C}}(J) = \mathfrak{so}_j^{\mathbb{C}}(J) \cdot J$. Besides we have more generally $J \cdot \mathcal{A}_j^{\mathbb{C}}(J) = \mathcal{A}_j^{\mathbb{C}}(J) \cdot J = \mathcal{A}_j^{\mathbb{C}}(J)$.

¹⁶ The conjugaison by J is denoted by $\text{Int}J: GL_n(\mathbb{R}) \rightarrow GL_n(\mathbb{R})$ when the domain of definition is a Lie subgroup and by $\text{Ad}J: \mathfrak{gl}_n(\mathbb{R}) \rightarrow \mathfrak{gl}_n(\mathbb{R})$ when it is a Lie subalgebra.

Consider now

$$\mathbb{U}_{j-1}(J) := \text{Com}(J^j) \cap SO(2n) = SO(2n)^{T^j},$$

Then T is an order j automorphism¹⁷ on $\mathbb{U}_{j-1}(J)$ and gives rises to the j -symmetric space $\mathbb{U}_{j-1}(J)/\mathbb{U}_0(J)$ which is in fact equal to

$$\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) := \{J' \in \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}) \mid (J')^j = J^j\}.$$

Indeed let $J' \in \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$, then there exists $g \in SO(2n)$ such that $J' = gJg^{-1}$, then $(J')^j = J^j$ if and only if $gJ^jg^{-1} = J^j$ i.e. $g \in \mathbb{U}_{j-1}(J)$, which proves that $\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) = \text{Int}(\mathbb{U}_{j-1}(J))(J)$ i.e.

$$\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) = \mathbb{U}_{j-1}(J)/\mathbb{U}_0(J).$$

Let us recapitulate what precedes.

Proposition 3.3 *Let $J \in \mathcal{Z}_{2k}(\mathbb{R}^{2n})$. The connected component $\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$ of J is a $SO(2n)$ -orbit (via the adjoint action):*

$$\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}) = SO(2n)/\mathbb{U}_0(J).$$

Moreover, denoting by r the order of this component (i.e. the order of $\text{Ad}J$), then it is also a r -symmetric space defined by the inner automorphism $T = \text{Int}J|_{SO(2n)}$.

Furthermore, for any $j \in \{1, \dots, r\}$, let us consider the subgroup $\mathbb{U}_{j-1}(J) := \{g \in SO(2n), gJ^jg^{-1} = J^j\}$. Then the submanifold of $\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$, defined by

$$\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) := \{J' \in \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}) \mid (J')^j = J^j\}$$

is the $\mathbb{U}_{j-1}(J)$ -orbit of J . Moreover, the restriction of T to $\mathbb{U}_{j-1}(J)$ is an order j automorphism of which the associated j -symmetric space is

$$\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) = \mathbb{U}_{j-1}(J)/\mathbb{U}_0(J).$$

We have then a increasing sequence (N_j) indexed by $j \in \{1, \dots, r\}$, of j -symmetric spaces, all included in $N_r = \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$.

Remark 3.2 Obviously, in this equation J can be replaced by any $J' \in \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j)$.

Example 3.3 If $k = 2$, then we have $\mathcal{Z}_{4,2}^\alpha(\mathbb{R}^{2n}, J^2) = \mathcal{Z}_{4,2}^\alpha(\mathbb{R}^{2n}, -\text{Id}) = \mathcal{Z}_4^\alpha(\mathbb{R}^{2n}) = \Sigma^\alpha(\mathbb{R}^{2n}) = SO(2n)/U(n)$, and the other values of j are trivial $\mathcal{Z}_{4,\pm 1}^\alpha(\mathbb{R}^{2n}, J^{\pm 1}) = \{J\}$.

Remark 3.3 Sometimes, we will need to precise clearly what is the eigenvalues of the eigenspaces $\mathcal{A}_i^{\mathbb{C}}(J)$ and $\mathfrak{so}_i^{\mathbb{C}}(J)$, then we will simply use the notation

$$\mathcal{A}_{(\omega)}^{\mathbb{C}}(J) = \ker(\text{Ad}J - \omega \text{Id})$$

and idem for $\mathfrak{so}_{(\omega)}^{\mathbb{C}}(J)$ and $\mathcal{B}_{(\omega)}^{\mathbb{C}}(J)$.

Besides, sometimes for the homogeneity of the equations, we will extend the notations $\mathcal{A}_i^{\mathbb{C}}(J)$ for real index and set for $t \in \mathbb{R}$

$$\mathcal{A}_t^{\mathbb{C}}(J) = \ker(\text{Ad}J - \omega_r^t \text{Id}).$$

¹⁷We confuse $j \in \mathbb{Z}_r$ and its representant in $\{1, \dots, r\}$.

3.1.3 Study of $\text{Ad}J^j$

Expression of the eigenspaces of $\text{Ad}J^j$ in terms of those of $\text{Ad}J$. Let $j \in \mathbb{Z}^*$. Then we have

$$\text{Ad}J^j = (\text{Ad}J)^j = \bigoplus_{l=0}^{r-1} (\omega_r^l)^j \text{Id}_{\mathcal{A}_l^{\mathbb{C}}(J)},$$

ω_r^j is of order $p = \frac{r}{(r,j)}$, i.e. it is in $\hat{U}_p = \{z \in S^1 | z^p = 1\} = \exp\left(\frac{2i\pi}{p}(\mathbb{Z}/p\mathbb{Z})\right)$. Then we have

$$\text{Ad}J^j = \bigoplus_{q=0}^{(r,j)-1} \left[\bigoplus_{l=0}^{p-1} (\omega_r^j)^l \text{Id}_{\mathcal{A}_{qp+l}^{\mathbb{C}}(J)} \right],$$

hence writing (that $\text{Ad}J^j$ is of order p):

$$\text{Ad}J^j = \bigoplus_{l=0}^{p-1} \omega_p^l \text{Id}_{\mathcal{A}_l^{\mathbb{C}}(J^j)},$$

we obtain

Lemma 3.1 *We have the following relation between the eigenspaces of $\text{Ad}J^j$ and those of $\text{Ad}J$:*

$$\mathcal{A}_l^{\mathbb{C}}(J^j) = \bigoplus_{q=0}^{(r,j)-1} \mathcal{A}_{qp+l'}^{\mathbb{C}}(J) \quad (44)$$

where $l' = (j')^{-1}l$ in the ring $\mathbb{Z}/p\mathbb{Z}$, and $j' = \left[\frac{j}{(j,r)} \right]_{\text{mod } p}$ (j' is inversible in the ring $\mathbb{Z}/p\mathbb{Z}$, since $(j',p) = 1$ by definition of (r,j)).

In particular,

$$\text{Com}(J^j)^{\mathbb{C}} = \mathcal{A}_0^{\mathbb{C}}(J^j) = \bigoplus_{q=0}^{(r,j)-1} \mathcal{A}_{qp}^{\mathbb{C}}(J). \quad (45)$$

More particularly,

$$\text{Com}(J^k)^{\mathbb{C}} = \begin{cases} \bigoplus_{q=0}^{k-1} \mathcal{A}_q^{\mathbb{C}}(J) = \text{End}(\mathbb{R}^{2n})^{\mathbb{C}} & \text{if } r = k \\ \bigoplus_{q=0}^{k-1} \mathcal{A}_{2q}^{\mathbb{C}}(J) & \text{if } r = 2k \end{cases}$$

and

$$\text{Com}(J^2) = \begin{cases} \mathcal{A}_0(J) \oplus \mathcal{A}_k(J) & \text{if } r = 2k \\ \mathcal{A}_0(J) \oplus \mathcal{A}_{\frac{k}{2}}(J) & \text{if } r = k \in 2\mathbb{Z} \\ \mathcal{A}_0(J) & \text{if } r = k \text{ is odd} \end{cases} = \mathcal{A}_0(J) \oplus \mathcal{A}_{\frac{r}{2}}(J) \quad .$$

Decomposition of the tangent space $T_J \mathcal{Z}_{2k,j}^{\alpha}(\mathbb{R}^{2n}, J^j)$ in terms the eigenspaces $\mathcal{B}_l^{\mathbb{C}}(J)$ of $\text{Ad}J$. We can rewrite all what precedes in $J.\mathfrak{so}(2n)$ (resp. in $\mathfrak{so}(2n)$) by replacing $\mathcal{A}_l^{\mathbb{C}}$ by $\mathcal{B}_l^{\mathbb{C}}$ (resp. $\mathfrak{so}_l^{\mathbb{C}}$). In particular we have, according to (45),

$$\mathfrak{u}_{j-1}(J) := \text{Lie}(\mathbb{U}_{j-1}(J)) = \mathfrak{so}_0(J^j) = \left(\bigoplus_{q=0}^{(r,j)-1} \mathfrak{so}_{qp}^{\mathbb{C}}(J) \right) \cap \mathfrak{so}(2n),$$

this¹⁸ is the eigenspace decomposition of the order j automorphism obtained by restricting $T = \text{Ad}J$ to $\mathfrak{u}_{j-1}(J)$. Therefore we have

¹⁸We mean $\mathfrak{u}_{j-1}(J)^{\mathbb{C}} = \bigoplus_{q=0}^{(r,j)-1} \mathfrak{so}_{qp}^{\mathbb{C}}(J)$

Proposition 3.4 *The tangent space of $\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j)$ is given by*

$$T_J \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) = \left(\bigoplus_{q=1}^{(r,j)-1} \mathcal{B}_{qp}(J) \right) \cap \text{End}(\mathbb{R}^{2n}). \quad (46)$$

Proof. Indeed, $g \in \mathbb{U}_{j-1}(J) \mapsto gJg^{-1} \in \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j)$ is a surjective submersion whose the (surjective) derivative at $g = 1$,

$$A \in \mathfrak{u}_{j-1}(J) \mapsto [A, J] = \sum_{q=0}^{(r,j)-1} [A_{qp}, J] = \sum_{q=0}^{(r,j)-1} (1 - \omega_r^{qp}) A_{qp} J \in T_J \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j)$$

has $\left(\bigoplus_{q=1}^{(r,j)-1} \mathcal{B}_{qp}(J) \right) \cap \text{End}(\mathbb{R}^{2n})$ as image, which proves the equality (46). \square

More simply by differentiating the definition equation of $\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j)$ we obtain

$$T_J \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) = \left\{ A \in J \cdot \mathfrak{so}(2n) \left| \sum_{p+l=j-1} J^p A J^l = 0 \right. \right\} \quad (47)$$

Now, let us apply (46) for $j = 2$:

$$T_J \mathcal{Z}_{2k,2}^\alpha(\mathbb{R}^{2n}, J^2) = \begin{cases} \mathcal{B}_{\frac{r}{2}}(J) = \{A \in J \cdot \mathfrak{so}(2n) \mid AJ + JA = 0\} & \text{if } r \text{ is even} \\ 0 & \text{if } r \text{ is odd.} \end{cases}$$

This can be recovered from (47) by remarking that if r is odd then -1 is not a r -th root of unity (and thus not an eigenvalue of $\text{Ad}J$).

Remark 3.4 If $(j, 2k) = 1$ (so that $(j, r) = 1$ also) then $J^j \in \mathcal{Z}_{2k}(\mathbb{R}^{2n})$ and T^j is of order r and we have, according to (44)

$$\mathcal{A}_l^{\mathbb{C}}(J^j) = \mathcal{A}_{[j]_r^{-1} \cdot l}^{\mathbb{C}}(J), \quad \forall l \in \mathbb{Z}/r\mathbb{Z},$$

in others words $\mathcal{A}_l^{\mathbb{C}}(J) = \mathcal{A}_{j \cdot l}^{\mathbb{C}}(J^j)$, $\forall l \in \mathbb{Z}/r\mathbb{Z}$. In particular

$$\mathbb{U}_{j-1}(J) = \mathcal{A}_0(J^j) \cap SO(2n) = \mathcal{A}_0(J) \cap SO(2n) = \mathbb{U}_0(J).$$

Hence

$$\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) = \{J\}.$$

More generally, we have, according to (45), since $(jl, r) = (l, r)$,

$$\text{Com}((J^j)^l)^{\mathbb{C}} = \text{Com}(J^{jl})^{\mathbb{C}} = \bigoplus_{q=0}^{(l,r)-1} \mathcal{A}_{qp}^{\mathbb{C}}(J) = \text{Com}(J^l)^{\mathbb{C}}$$

with $p = \frac{r}{(l,r)}$, and thus

$$\text{Com}((J^j)^l) = \text{Com}(J^l) \quad \forall l \in \mathbb{Z}/r\mathbb{Z}.$$

In particular, $\mathbb{U}_{l-1}(J^j) = \mathbb{U}_{l-1}(J) \quad \forall l \in \mathbb{Z}/r\mathbb{Z}$ and thus

$$\mathcal{Z}_{2k,l}^\alpha(\mathbb{R}^{2n}, (J^j)^l) = \mathcal{Z}_{2k,l}^{[j]_{2k}^{-1} \cdot \alpha}(\mathbb{R}^{2n}, J^l)$$

where $[j]_{2k}^{-1} \cdot \alpha$ is the action of $[j]_{2k}^{-1}$ on $\alpha \in X_{2k}$, the action of $l \in (\mathbb{Z}/r\mathbb{Z})^*$ on X_{2k} being defined by the bijective map

$$J \in \mathcal{Z}_{2k}(\mathbb{R}^{2n}) \mapsto J^l \in (\mathcal{Z}_{2k}(\mathbb{R}^{2n}))^l = \mathcal{Z}_{2k}(\mathbb{R}^{2n})$$

which sends a connected component onto another one

$$(\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}))^l =: \mathcal{Z}_{2k}^{l \cdot \alpha}(\mathbb{R}^{2n}).$$

In particular, for $j = 1$, we have

$$\mathbb{U}_{l-1}(J^{-1}) = \mathbb{U}_{l-1}(J) \quad \forall l \in \mathbb{Z}/r\mathbb{Z}$$

and thus

$$\mathcal{Z}_{2k,l}^\alpha(\mathbb{R}^{2n}, (J^{-1})^l) = \mathcal{Z}_{2k,l}^{-\alpha}(\mathbb{R}^{2n}, J^l)$$

where $-(\varepsilon, \underline{p}) = ((-1)^n \varepsilon, \underline{p})$ in X_{2k} . Hence $\mathcal{Z}_{2k,l}^\alpha(\mathbb{R}^{2n}, (J^{-1})^l) = \mathcal{Z}_{2k,l}^\alpha(\mathbb{R}^{2n}, J^l)$ if and only if n is even (i.e. J and J^{-1} are in the same connected component).

3.2 Isometries of order $2k + 1$ with no eigenvalue $= 1$

We can do exactly the same study for $\mathcal{Z}_{2k+1}(E)$ as we did for $\mathcal{Z}_{2k}(E)$, with however the following simplification: all the connected components have the same order $r = 2k + 1$ and we do not have to distinguish two types of orbits as previously. Let us review all the results obtained in 3.1 to see those which hold and for these latter, see if some modifications are necessary.

- The space $\mathcal{Z}_{2k+1}(\mathbb{R}^{2n+1})$ is empty.
- Theorem 3.1 holds:

$$\pi_0(\mathcal{Z}_{2k+1}(\mathbb{R}^{2n})) = X_{2k+1} := \left\{ (\varepsilon, p) \in \mathbb{Z}_2 \times \mathbb{N}^k \left| \sum_{j=1}^k p_j = n \text{ and } \text{lcm} \left(\left\{ \frac{2k+1}{(2k+1) \wedge j}, p_j \neq 0 \right\} \right) = 2k+1 \right. \right\}$$

i.e. more generally

$$\pi_0(\mathcal{Z}_{k'}(\mathbb{R}^{2n})) = X_{k'} := \left\{ (\varepsilon, p) \in \mathbb{Z}_2 \times \mathbb{N}^{m_{k'}-1} \left| \sum_{j=1}^{m_{k'}-1} p_j = n \text{ and } \text{lcm} \left(\left\{ \frac{k'}{k' \wedge j}, p_j \neq 0 \right\} \right) = k' \right. \right\}$$

- Equation (41) holds in general for any order
- Propositions 3.1 and 3.2 holds also for any order k' .
- Propositions 3.3 holds for $\mathcal{Z}_{2k+1}(\mathbb{R}^{2n})$ (replace everywhere $2k$ and r by $2k + 1$).

More generally, let us consider the family of spaces $\mathcal{U}_{k'}(\mathbb{R}^{2n})$. Then we could write explicitly $\pi_0(\mathcal{U}_{k'})$. Moreover equation (41) holds in general. Propositions 3.1 and 3.2 holds also in general. Finally, Propositions 3.3 holds also in general. Let us precise that then r is always the order of $\text{Ad}J$, in the considered connected component.

3.3 The effect of the power maps on the finite order isometries

Let $J \in \mathcal{U}_{k'}(\mathbb{R}^{2n})$ then $J^j \in \mathcal{U}_p(\mathbb{R}^{2n})$ with $p = \frac{k'}{(k',j)}$. Moreover it is easy to see (from the diagonalisation) that the power map

$$J \mapsto J^j$$

is surjective from $\mathcal{U}_{k'}(\mathbb{R}^{2n})$ onto $\mathcal{U}_p(\mathbb{R}^{2n})$ (since $z \in \hat{U}_{k'}(\mathbb{R}^{2n}) \mapsto z^j \in \hat{U}_p(\mathbb{R}^{2n})$ is surjective). Besides, since each connected component in $\mathcal{U}_{k'}(\mathbb{R}^{2n})$ (and in $\mathcal{U}_p(\mathbb{R}^{2n})$) is a $SO(2n)$ -orbit, then the power map $J \mapsto J^j$ sends one component in $\mathcal{U}_{k'}(\mathbb{R}^{2n})$ onto another one in $\mathcal{U}_p(\mathbb{R}^{2n})$ so that it induces a map:

$$\alpha \in \pi_0(\mathcal{U}_{k'}(\mathbb{R}^{2n})) \longrightarrow j \cdot \alpha \in \pi_0(\mathcal{U}_p(\mathbb{R}^{2n}))$$

such that

$$(\mathcal{U}_{k'}^\alpha)^j(\mathbb{R}^{2n}) =: \mathcal{U}_p^{j \cdot \alpha}(\mathbb{R}^{2n}), \quad \forall \alpha \in \pi_0(\mathcal{U}_{k'}(\mathbb{R}^{2n})). \quad (48)$$

Remark 3.5 In general we have $(\mathcal{Z}_{2k})^j(\mathbb{R}^{2n}) \not\subseteq \mathcal{Z}_p(\mathbb{R}^{2n})$. For example, for $j = 2$, we have

$$(\mathcal{Z}_{2k}(\mathbb{R}^{2n}))^2 = \begin{cases} \mathcal{U}_k^*(\mathbb{R}^{2n}) & \text{if } k \text{ is even} \\ \mathcal{Z}_k(\mathbb{R}^{2n}) & \text{if } k \text{ is odd.} \end{cases}$$

Besides, given $J \in \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$, then $\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j)$ is the inverse image of J^j by the map

$$J' \in \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}) \longmapsto (J')^j \in (\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}))^j = \mathcal{U}_p^{j \cdot \alpha}(\mathbb{R}^{2n}).$$

Since $(J')^j$ is constant in $\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j)$, we can denote it by J_j and then

$$\mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J^j) = \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, J_j). \quad \blacksquare$$

Furthermore, we have also for any $J \in \mathcal{U}_{k'}^\alpha(\mathbb{R}^{2n})$, $\mathbb{U}_0(J^j) = \mathbb{U}_{j-1}(J)$ so that, according to (48), the component of J^j in $\mathcal{U}_p(\mathbb{R}^{2n})$ satisfies

$$\mathcal{U}_p^{j \cdot \alpha}(\mathbb{R}^{2n}) = SO(2n)/\mathbb{U}_0(J^j) = SO(2n)/\mathbb{U}_{j-1}(J),$$

hence

$$(\mathcal{U}_{k'}^\alpha(\mathbb{R}^{2n}))^j = SO(2n)/\mathbb{U}_{j-1}(J) \quad (49)$$

which we can recover directly by taking the power j in the equality $\mathcal{U}_{k'}^\alpha(\mathbb{R}^{2n}) = \{gJg^{-1}, g \in SO(2n)\}$.

Convention: for each $\alpha \in \pi_0(\mathcal{Z}_{2k}(\mathbb{R}^{2n}))$, we will choose (and fix) a canonical representant in $\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$. For example, let $(\epsilon_1, \dots, \epsilon_{2n})$ be the canonical basis in \mathbb{R}^{2n} , and

$$\begin{aligned} e_{2l+1} &= \frac{\epsilon_{2l+1} + i\epsilon_{2l+2}}{\sqrt{2}}, \quad 0 \leq l \leq n-1, \\ e_{2l} &= \bar{\epsilon}_{2l-1}, \quad 1 \leq l \leq n. \end{aligned}$$

Then $e = (e_1, \dots, e_{2n})$ is a hermitian basis in \mathbb{C}^{2n} and we can take J_0^α such that

$$\text{Mat}_e(J_0^\alpha) = \text{Diag} \left(\begin{pmatrix} e^{i\theta_j} \text{Id}_{p_j} & 0 \\ 0 & e^{-i\theta_j} \text{Id}_{p_j} \end{pmatrix}, 1 \leq j \leq n-1 \right)$$

where $\underline{p} = (p_1, \dots, p_{n-1})$ is determined by $\alpha = (\varepsilon, \underline{p}) \in \pi_0(\mathcal{Z}_{2k}(\mathbb{R}^{2n}))$ (see section 3.1.1), $\theta_j = \frac{j\pi}{k}$, and $\text{Mat}_e(\cdot)$ means "the matrice in the basis e of".

3.4 The Twistor spaces of a Riemannian manifolds and its reductions

Let M be an oriented (even dimensional) Riemannian manifolds and let us consider the bundle of order $2k$ isometries $\mathcal{U}_{2k}(M)$ as well as its subbundles $\mathcal{U}_{2k}^*(M)$ and $\mathcal{Z}_{2k}(\mathbb{R}^{2n})$. Let us fix $\alpha \in \pi_0(\mathcal{Z}_{2k}(\mathbb{R}^{2n}))$ and consider the component $\mathcal{Z}_{2k}^\alpha(M)$. Then denoting by $\mathcal{SO}(M)$ the $SO(2n)$ -bundle of positively oriented orthonormal frames on M , we have¹⁹

$$\mathcal{Z}_{2k}^\alpha(M) = \mathcal{SO}(M)/\mathbb{U}_0(J_0^\alpha).$$

We want to ask the following question: does $\mathcal{SO}(M)$ admit a $\mathbb{U}_{j-1}(J_0^\alpha)$ -reduction for $1 \leq j \leq r$. We know (according to [40]) that $\mathcal{SO}(M)$ admits a $\mathbb{U}_{j-1}(J_0^\alpha)$ -reduction if and only if the associated bundle $\mathcal{SO}(M)/\mathbb{U}_{j-1}(J_0^\alpha)$ ($= \mathcal{SO}(M) \times_{SO(2n)} SO(2n)/\mathbb{U}_{j-1}(J_0^\alpha)$) admits a global section $J_j: M \rightarrow \mathcal{SO}(M)/\mathbb{U}_{j-1}(J_0^\alpha)$.

Besides, according to (49) and (48) applied to $\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$, we have

$$\mathcal{SO}(M)/\mathbb{U}_{j-1}(J_0^\alpha) = (\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}))^j = \mathcal{U}_p^{j,\alpha}(M)$$

with $p = \frac{2k}{(2k, j)}$. Hence J_j (when it exists) is a global section of $(\mathcal{Z}_{2k}^\alpha(M))^j$ and then the $\mathbb{U}_{j-1}(J_0^\alpha)$ -reduction of $\mathcal{SO}(M)$ is given in terms of J_j by:

$$\mathfrak{U}_{j-1}^\alpha(M) := \{e = (e_1, \dots, e_{2n}) \in \mathcal{SO}(M) \mid \text{Mat}_e(J_j) = (J_0^\alpha)^j\}.$$

Then we have

$$\mathfrak{U}_{j-1}^\alpha(M)/\mathbb{U}_0(J_0^\alpha) = \mathcal{Z}_{2k,j}^\alpha(M, J_j).$$

In particular, since $(J_0^\alpha)^r = \pm \text{Id}$, we have $\mathbb{U}_{r-1}(J_0^\alpha) = SO(2n)$ and $\mathcal{SO}(M)$ has always an (unique and trivial) $SO(2n)$ -reduction for which $J_r = \pm \text{Id}_{TM}$ and thus $\mathfrak{U}_{r-1}^\alpha(M) = \mathfrak{U}_0^\alpha(M) = \mathcal{SO}(M)$ and $\mathcal{Z}_{2k,r}^\alpha(M, J_r) = \mathcal{Z}_{2k}^\alpha(M)$.

Example 3.4 If $k = 2$, and thus $r = 2$, then $J_0 = -\text{Id}$ defines the trivial reduction. Moreover, for $j = 1$, a global section J_1 (when it exists) defines on M an almost complex structure and $\mathfrak{U}_{j-1}^\alpha(M) = \mathfrak{U}_0^\alpha(M)$ is then the subbundle of hermitian frames on M (with respect to this almost complex structure).

3.5 Return to an order $2k$ automorphism $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$.

We give ourself the same ingredients as in section 2.1.1 and we use the same notations. In particular, we suppose that the subgroup H is chosen such that $(G^\sigma)^0 \subset H \subset G^\sigma$. In addition to that we suppose G/H Riemannian.

3.5.1 Case $r = k$

Suppose that we have $\tau|_{\mathfrak{m}} = -\text{Id}$ i.e. $\tau|_{\mathfrak{m}} \in \mathcal{Z}_{2k}^0(\mathfrak{m})$. Then $\mathfrak{g}_{2j}^{\mathbb{C}} = 0$ for all $2j \in \mathbb{Z}/(2k)\mathbb{Z} \setminus \{0, k\}$. Hence we have

$$[\mathfrak{g}_p^{\mathbb{C}}, \mathfrak{g}_l^{\mathbb{C}}] = \{0\} \quad \text{if } p+l \neq 0, k.$$

Indeed, if p or l is even then the corresponding eigenspace vanishes. If p and l are odd then $[\mathfrak{g}_p^{\mathbb{C}}, \mathfrak{g}_l^{\mathbb{C}}] \subset \mathfrak{g}_{p+l}^{\mathbb{C}}$ and $p+l$ is even, thus $\mathfrak{g}_{p+l}^{\mathbb{C}} = \{0\}$ except if $p+l = 0$ or k . Consequently, we have $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}$ and thus G/H is a (locally) symmetric space. Let us distinguish the following two cases.

¹⁹The canonical isomorphism which allows to write this equality will be explicitely given in section 4.3.

k is odd. Then $[\mathfrak{g}_k, \mathfrak{g}_j^{\mathbb{C}}] \subset \mathfrak{g}_{k+j}^{\mathbb{C}} = \{0\}$ for all j odd $\neq 0, k$. Hence $[\mathfrak{g}_k, \mathfrak{m}] = \{0\}$ i.e. $\text{ad}_{\mathfrak{m}} \mathfrak{g}_k = 0$ so that $\mathfrak{g}_k = 0$ and thus this case is trivial because $H = G_0$ up to covering and thus the fibre H/G_0 is trivial (i.e. a discrete set). Moreover we have $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h} = \mathfrak{g}_0$ and $G/H = G/G_0$ (up to covering) is the (locally) symmetric space associated to the involution τ^k .

k is even. Then the symmetric decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ is the eigenspace decomposition of τ^k , and G/H is the (locally) symmetric space corresponding to this involution τ^k .

In conclusion, if $r = k$, then G/H is the (locally) symmetric space corresponding to τ^k .

Example 3.5 If $2k = 4$, then we always have $r = k = 2$ (since $\tau_{\mathfrak{m}}^2 = -\text{Id}$) and G/H is the symmetric space corresponding to $\sigma = \tau^2$.

3.5.2 Action of $\text{Ad}\tau_{\mathfrak{m}}$ on $\text{ad}\mathfrak{g}_j^{\mathbb{C}}$

We have $\tau \circ \text{ad}X \circ \tau^{-1} = \text{ad}\tau(X)$, $\forall X \in \mathfrak{g}$. In particular, for all $j \in \mathbb{Z}/(2k)\mathbb{Z}$ we have

$$\forall X_j \in \mathfrak{g}_j^{\mathbb{C}}, \tau \circ \text{ad}X_j \circ \tau^{-1} = \omega_{2k}^j \text{ad}X_j.$$

Hence by taking the restriction to \mathfrak{m} and projecting on \mathfrak{m} :

$$\tau_{\mathfrak{m}} \circ [\text{ad}_{\mathfrak{m}} X_j]_{\mathfrak{m}} \circ \tau_{\mathfrak{m}}^{-1} = \omega_{2k}^j [\text{ad}_{\mathfrak{m}} X_j]_{\mathfrak{m}} \quad (50)$$

so that²⁰

$$[\text{ad}_{\mathfrak{m}} \mathfrak{g}_j^{\mathbb{C}}]_{\mathfrak{m}} \subset \mathcal{A}_j^{\mathbb{C}}(\tau_{\mathfrak{m}}) \quad \forall j \in \mathbb{Z}/(2k)\mathbb{Z}.$$

If $r = k$ then $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}$, hence $[\text{ad}_{\mathfrak{m}} \mathfrak{g}_j^{\mathbb{C}}]_{\mathfrak{m}} = 0$, for all $j \in \mathbb{Z}/(2k)\mathbb{Z} \setminus \{0, k\}$.

Moreover, let us recall that we always have ($r = 2k$ or k)

$$\begin{aligned} [\text{ad}_{\mathfrak{m}} \mathfrak{g}_0]_{\mathfrak{m}} &= \text{ad}_{\mathfrak{m}} \mathfrak{g}_0 \subset \mathfrak{so}_0(\tau_{\mathfrak{m}}) = \text{Com}(\tau_{\mathfrak{m}}) \cap \mathfrak{so}(\mathfrak{m}) \\ [\text{ad}_{\mathfrak{m}} \mathfrak{g}_k]_{\mathfrak{m}} &= \text{ad}_{\mathfrak{m}} \mathfrak{g}_k \subset \begin{cases} \mathfrak{so}_{(-1)}(\tau_{\mathfrak{m}}) = \text{Ant}(\tau_{\mathfrak{m}}) \cap \mathfrak{so}(\mathfrak{m}) = \mathfrak{so}_{\frac{\pi}{2}}(\tau_{\mathfrak{m}}) & \text{if } r \text{ is even} \\ 0 & \text{if } r = k \text{ is odd (trivial case)} \end{cases} \end{aligned}$$

where²¹ $\text{Ant}(\tau_{\mathfrak{m}}) = \{A \in \text{End}(\mathfrak{m}) \mid A\tau_{\mathfrak{m}} + \tau_{\mathfrak{m}}A = 0\}$, and we used the notation defined in remark 3.3 for $\mathfrak{so}_{(-1)}(\tau_{\mathfrak{m}})$.

Remark 3.6 In general, we do not have $[\text{ad}_{\mathfrak{m}} \mathfrak{g}_j^{\mathbb{C}}]_{\mathfrak{m}} \subset \mathfrak{so}_j^{\mathbb{C}}(\tau_{\mathfrak{m}})$ and we also do not have $[\text{ad}_{\mathfrak{m}} \mathfrak{g}_j^{\mathbb{C}}]_{\mathfrak{m}} \subset \mathcal{B}_j(\tau_{\mathfrak{m}})$. However, if the metric is naturally reductive, then these inclusions hold.

We can easily generalize (50) as follows. We keep in mind the conventions and notations defined in the subsection 2.0.1. We use also the notations of the beginning of subsection 2.1.

Proposition 3.5 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of Lie algebra of finite order k' . Let us consider some decomposition $k' = pq$, with $p, q \in \mathbb{N}^*$. Then let us consider the order p automorphism $\hat{\sigma} = \tau^q$, and set $\hat{\mathfrak{h}} = \mathfrak{g}^{\hat{\sigma}}$. We denote by $\mathfrak{g} = \hat{\mathfrak{h}} \oplus \hat{\mathfrak{m}}$ the corresponding $\hat{\sigma}$ -invariant reductive decomposition of the Lie algebra \mathfrak{g} . Then we have*

$$\tau_{\hat{\mathfrak{m}}} \circ [\text{ad}_{\hat{\mathfrak{m}}} X_j]_{\hat{\mathfrak{m}}} \circ \tau_{\hat{\mathfrak{m}}}^{-1} = \omega_{k'}^j [\text{ad}_{\hat{\mathfrak{m}}} X_j]_{\hat{\mathfrak{m}}}, \quad \forall X_j \in \mathfrak{g}_j^{\mathbb{C}}.$$

Remark 3.7 In particular, we can apply the previous proposition in the case $k' = 2k + 1$ and $q = 1$.

²⁰See remark 3.6.

²¹According to the notation defined in 0.6.

3.6 The canonical section in $(\mathcal{Z}_{2k}(G/H))^2$, the canonical embedding, and the Twistor lifts

3.6.1 The canonical embedding

Once more, we give ourself the same ingredients and notations as in section 2.1.1. We suppose that $(G^\sigma)^0 \subset H \subset G^\sigma$ and that G/H is Riemannian. We denote by $p_0 := 1.H$ the reference point in G/H . According to the definition of H , we have

Lemma 3.2 *Let J_0 be the element in $\mathcal{Z}_{2k}^{\alpha_0}(T_{p_0}M)$ corresponding²² to $\tau_{|\mathfrak{m}}$ (or more generally to $\tau_{|\mathfrak{m}}^j$ with $(j, 2k) = 1$) under the identification $T_{p_0}M = \mathfrak{m}$. Then we have $\forall g \in H, gJ_0^2g^{-1} = J_0^2$. Hence there exists a unique section*

$$J_2: G/H \mapsto (\mathcal{Z}_{2k}^{\alpha_0})^2 = \mathcal{U}_k^{\alpha_0^2}(G/H)$$

defined by

$$g.p_0 \in G/H \mapsto gJ_0^2g^{-1} \in (\mathcal{Z}_{2k}^{\alpha_0})^2.$$

Proceeding as in [34], Theorem 3, we obtain:

Theorem 3.2 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an order $2k$ automorphism and $M = G/H$ a (locally) Riemannian k -symmetric space corresponding to $\sigma = \tau^2$. Let us make G acting on $\mathcal{Z}_{2k}(M)$: $g \cdot J = gJg^{-1}$. Let $J_0 \in \mathcal{Z}_{2k}^{\alpha_0}(T_{p_0}M)$ be the finite order isometry corresponding to $\tau_{|\mathfrak{m}}$ under the identification $T_{p_0}M = \mathfrak{m}$. Then the orbit of J_0 under the action of G is an immersed submanifold in $\mathcal{Z}_{2k}^{\alpha_0}(M)$. Denoting by G_0 the stabilizer of J_0 , then $G_0 = G^\tau \cap H$ and thus $N = G/G_0$ is a locally $2k$ -symmetric bundle over M and the natural map:*

$$\begin{aligned} \mathfrak{J}_{J_0}: G/G_0 &\longrightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2) \\ g.G_0 &\longmapsto gJ_0g^{-1} \end{aligned}$$

is an injective immersion and a morphism of bundle. Moreover, if the image of G in $\text{Is}(M)$ (the group of isometry of M) is closed, then \mathfrak{J}_{J_0} is an embedding.

Remark 3.8 Remark that we could also have chosen $J_0 = \tau_{|\mathfrak{m}}^j$ with $(j, 2k) = 1$, since in this case τ^j is an order $2k$ -isomorphism. See [34], theorem 3 and remark 13.

Notation For a geometric map $f: L \rightarrow G/G_0$, we will denote by J the corresponding map $\mathfrak{J}_{J_0} \circ f: L \rightarrow \mathcal{Z}_{2k,2}(G/H, J_2)$ under the previous inclusion $G/G_0 \hookrightarrow \mathcal{Z}_{2k,2}(G/H, J_2)$.

A remark about the canonical section In fact the lemma 3.2 can be written more generally as follows:

Lemma 3.3 *Let $N = G/G_0$ be a (locally) k' -symmetric space. We denote by $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{n}$ the reductive decomposition of the Lie algebra. Let $y_0 = 1.G_0 \in N$ and let J_0 be the element in $\mathcal{U}_p^*(T_{y_0}N)$ corresponding to $\tau_{|\mathfrak{n}}$ under²³ the identification $T_{y_0}N = \mathfrak{n}$. Then we have $\forall g \in G_0, gJ_0g^{-1} = J_0$. Hence there exists a canonical section $J_1: G/G_0 \mapsto \mathcal{U}_p^*(T_{y_0}N)$ defined by*

$$g.y_0 \in G/G_0 \mapsto gJ_0g^{-1} \in \mathcal{U}_p^*(T_{y_0}N).$$

²² α_0 denotes of course the connected component of J_0 in $\mathcal{Z}_{2k}(T_{p_0}M)$.

²³Or more generally corresponding to $\tau_{|\mathfrak{n}}^j$ with $(j, k') = 1$.

3.6.2 The twistor lifts

Definition 3.2 An isometry $A \in SO(\mathbb{R}^{2n})$ will be called an $e^{i\theta}$ -structure if $\text{Spect}(A) = \{e^{i\theta}, e^{-i\theta}\}$. An isometry $A \in SO(\mathbb{R}^{2n})$ will be called a $2k$ -structure if $A \in \mathcal{Z}_{2k}(\mathbb{R}^{2n})$.

Definition 3.3 Let $(E, h) \rightarrow M$ be a Riemannian vector bundle over a manifold M . Then for each $2k$ -structure $J \in \mathcal{Z}_{2k}(E)$, we denote by $\underline{\mathbf{J}}$ the complex structure in E defined by

$$\begin{aligned}\ker(\underline{\mathbf{J}} - i\text{Id}) &= \bigoplus_{j=1}^{k-1} \ker(J - \omega_{2k}^{-j}\text{Id}) \\ \ker(\underline{\mathbf{J}} + i\text{Id}) &= \bigoplus_{j=1}^{k-1} \ker(J - \omega_{2k}^j\text{Id})\end{aligned}$$

Remark 3.9 Let us remark that if $J \in \Sigma(E)$ is complex structure then $\underline{\mathbf{J}} = -J$. This sign is needed because of our convention chosen in remark 2.1. See also [34], remark 13.

Remark 3.10 We see that definition 3.3 defines a $SO(2n)$ -invariant map

$$P: \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}) \longrightarrow \Sigma(\mathbb{R}^{2n})$$

which is nothing but the $SO(2n)$ -invariant projection

$$SO(2n)/\mathbb{U}_0(J_0^\alpha) \longrightarrow SO(2n)/\mathbb{U}(\underline{\mathbf{J}}_0^\alpha),$$

since $\mathbb{U}_0(J_0^\alpha) \subset \mathbb{U}(\underline{\mathbf{J}}_0^\alpha)$. Indeed by definition $\underline{\mathbf{J}}_0^\alpha$ stabilizes the eigenspaces of J_0^α :

$$\underline{\mathbf{J}}_0^\alpha = \left(\bigoplus_{j=1}^{k-1} i\text{Id}_{[\mathfrak{g}_{-j}^c]} \right) \oplus \left(\bigoplus_{j=1}^{k-1} -i\text{Id}_{[\mathfrak{g}_j^c]} \right).$$

Moreover, the restriction of P to $\mathcal{Z}_{2k,j}(\mathbb{R}^{2n}, (J_0^\alpha)^j)$ is the $\mathbb{U}_{j-1}(J_0^\alpha)$ -invariant projection

$$\mathbb{U}_{j-1}(J_0^\alpha)/\mathbb{U}_0(J_0^\alpha) \longrightarrow \mathbb{U}_{j-1}(J_0^\alpha)/\mathbb{U}(\underline{\mathbf{J}}_0^\alpha).$$

Definition 3.4 Let $(E, h) \rightarrow M$ be a Riemannian vector bundle over a manifold M . Then for each element $J \in \mathcal{Z}_{2k+1}(E)$, we denote by $\underline{\mathbf{J}} \in \Sigma(E)$ the complex structure in E defined by

$$\begin{aligned}\ker(\underline{\mathbf{J}} - i\text{Id}) &= \bigoplus_{j=1}^k \ker(J - \omega_{2k+1}^{-j}\text{Id}) \\ \ker(\underline{\mathbf{J}} + i\text{Id}) &= \bigoplus_{j=1}^k \ker(J - \omega_{2k+1}^j\text{Id})\end{aligned}$$

Remark 3.11 We see then that the canonical almost complex structure $\underline{\mathbf{J}}$ defined in a $(2k+1)$ -symmetric space $N = G/G_0$ (see (26)), is the almost complex structure associated to the element $J_1 \in \mathcal{Z}_{2k+1}(N)$ defined by lemma 3.3, according to definition 3.4.

Definition 3.5 Let (L, i) be a complex manifold (of dimension $d \geq 1$), M an oriented Riemannian manifold and $u: L \rightarrow M$ a immersion. Then an element $J: L \rightarrow u^*(\mathcal{Z}_{2k}(M))$ is an admissible twistor lift of u if one of the following equivalent statements holds:

(i) $[\bar{\partial}u]_{\mathfrak{m}_j(J)^c} \in \ker(J - \omega_{2k}^j\text{Id})$ for all $1 \leq j \leq k-1$, where $\mathfrak{m}_j(J) \subset u^*TM$ is defined by

$$\mathfrak{m}_j(J)^c = \ker(J - \omega_{2k}^{-j}\text{Id}) \oplus \ker(J - \omega_{2k}^j\text{Id}).$$

(ii) Let $\underline{\mathbf{J}}$ be the complex structure on $u^*(TM)$ defined by the $2k$ -structure $J \in \mathcal{C}(u^*(\mathcal{Z}_{2k}(M)))$, then u is $\underline{\mathbf{J}}$ -holomorphic: $du \circ i = \underline{\mathbf{J}} \circ du$

In particular, if (L, i) is a Riemann surface, then we can add that the existence of an admissible twistor lift J of u implies in particular that u is a conformal immersion.

Remark 3.12 Let E_j be the orthogonal projection of the tangent subbundle $u_*(TL)$ on the subbundle $\mathfrak{m}_j(J)$. Then J is an admissible twistor lift means also : for all $j \in \{1, \dots, k-1\}$, J stabilizes E_j and $J|_{E_j}$ is a ω_{2k}^j -structure, i.e. a rotation of the plan E_j , and moreover this rotation defined by $J|_{E_j}$ and the rotation defined by i have opposite sens of rotation.

Theorem 3.3 In the situation described in theorem 3.2, let α be a \mathfrak{g} -valued Maurer-Cartan 1-form on a Riemann surface L and $f: L \rightarrow G/G_0$ its geometric map and $J = \mathfrak{I}_{J_0} \circ f$. The the following statements are equivalent:

- (i) $\alpha''_{-j} = 0, 1 \leq j \leq k-1$
- (ii) $J: L \rightarrow \mathcal{Z}_{2k,2}(G/H, J_2)$ is an admissible twistor lift.

Proof. This follows immediately from definition 3.5-(i). □

4 Vertically Harmonic maps and Harmonic sections of submersions

We will first recall some definitions and properties about vertical harmonicity and homogeneous fibre bundles (sections 4.1 and 4.2). We refer to [50, 51] for proofs. Then in section 4.3 we will apply this to the study of the examples which interest us, examples that we will already introduce and start to study in 4.1.2 : homogeneous spaces and twistor spaces. Finally, we will conclude with a geometric interpretation of the even minimal determined elliptic integrable system in terms of vertically harmonic twistor lifts (section 4.4).

4.1 Definitions, general properties and examples

4.1.1 The vertical energy functional

Let $(M, g), (N, h)$ be Riemannian manifolds and $\pi: N \rightarrow M$ a surjective submersion. We can do the splitting $TN = \mathcal{V} \oplus \mathcal{H}$, where the vertical and horizontal subbundles are defined by $\mathcal{V} = \ker d\pi$ and $\mathcal{H} = (\ker d\pi)^\perp = \mathcal{V}^\perp$.

For any map $u: M \rightarrow N$, we denote by $d^v u = (du)^v$ the vertical component of du . Following [50], this allows us to define the vertical energy density of u , $e^v(u) = \frac{1}{2}|d^v u|^2$, and the associated vertical energy functional:

$$E^v(u) = \frac{1}{2} \int_M |d^v u|^2 d\text{vol}_g.$$

Let us define the vertical tension field of $u: M \rightarrow N$ by

$$\tau^v(u) = \text{Tr}_g(\nabla^v d^v u)$$

where ∇^v denotes the vertical component of the Levi-Civita connection (of N) in TN , and Tr_g the trace with respect to g . Then we have

Theorem 4.1 [50] *The map $u: M \rightarrow N$ is a critical point of E^v with respect to vertical variations if and only if $\tau^v(u) = 0$. In particular, if u is a section, i.e. $\pi \circ u = \text{Id}_M$, then it is a critical point of E^v with respect to variations through sections if and only if $\tau^v(u) = 0$.*

Definition 4.1 *A map $u: M \rightarrow N$ is vertically harmonic if $\tau^v(u) = 0$. If moreover u is a section we will say that it is a harmonic section.*

4.1.2 Examples

Example 4.1 Let $\pi: N \rightarrow M$ be like above. Let (L, b) be a Riemannian manifold and $f: L \rightarrow N$ a map. Then we can consider the projection $u = \pi \circ f: L \rightarrow M$ and the manifold

$$u^*N := \{(z, n) \in L \times N, n \in \pi^{-1}(\{u(z)\})\}.$$

Then we have the submersion $u^*\pi: (z, n) \in u^*N \mapsto z \in L$. Furthermore, u^*N can be endowed canonically with a Riemannian metric: take the metric induced by the product metric

$$|(dz, dn)|^2 = |dz|^2 + |dn|^2$$

in $L \times N \supset u^*N$.

Definition 4.2 We will say that $f: L \rightarrow N$ is vertically harmonic if

$$\text{Tr}_b(\nabla^v d^v f) = 0.$$

When $u = \pi \circ f$ is an isometry and π a Riemannian submersion **this is equivalent to say that the corresponding section $\tilde{f}: L \rightarrow u^*N$ is a harmonic section** (see the Appendix, theorem 8.1).

Example 4.2 The twistor bundle of almost complex structures $\Sigma(E)$.

Let $p: (E, \nabla, \langle \cdot, \cdot \rangle) \mapsto (M, g)$ be a Riemannian vector bundle of rank $2n$ (in particular $\langle \cdot, \cdot \rangle$ is ∇ -parallel). Then we consider the bundle of orthogonal almost complex structure: $N_\Sigma = \Sigma(E) = \{(x, J_x), J_x \in \Sigma(E_x)\}$, where $\Sigma(E_x) = \{J \in \mathfrak{so}(E_x) | J^2 = -\text{Id}\}$. We have a fibration $\pi_\Sigma: N_\Sigma \rightarrow M$. The vertical space is given by: $\forall J \in N_\Sigma$,

$$\mathcal{V}_J := T_J \Sigma(E_x) = \{A \in \mathfrak{so}(E_x) | AJ + JA = 0\}$$

where $x = \pi_\Sigma(J)$.

The metric connection ∇ gives us a splitting: $T\Sigma(E) = \mathcal{V}^\Sigma \oplus \mathcal{H}^\Sigma$. Indeed we have the following splitting (coming from ∇)

$$T\mathfrak{so}(E) = p^*\mathfrak{so}(E) \oplus \mathcal{H} \quad (51)$$

where $p: \mathfrak{so}(E) \rightarrow M$ is the natural fibration²⁴. Then for any (local) section $J: U \subset M \rightarrow \Sigma(E)$, we have

$$0 = \nabla J^2 = (\nabla J)J + J(\nabla J)$$

so that $\nabla J \in \mathcal{V}^\Sigma$ and thus in the decomposition (51): $[dJ]_{\mathfrak{so}(E)} = \nabla J \in \mathcal{V}^\Sigma$ and thus $[dJ]_{\mathcal{H}} = dJ - \nabla J \in TN_\Sigma$ which allows us to conclude that

$$T\Sigma(E) = \mathcal{V}^\Sigma \oplus \mathcal{H}_{|\Sigma(E)}.$$

Then we can endow N_Σ with the metric

$$h = \pi^*g + \langle \cdot, \cdot \rangle_{\mathcal{V}^\Sigma} \quad (52)$$

where $\langle \cdot, \cdot \rangle_{\mathcal{V}^\Sigma}$ is the fibre metric in \mathcal{V}^Σ induced by the metric in $\mathfrak{so}(E)$:

$$\langle A, B \rangle = \text{Tr}(A^t.B). \quad (53)$$

With this metric we have obviously $\mathcal{H}^\Sigma = \mathcal{V}^{\Sigma^\perp}$.

Furthermore, let us remark that $T\Sigma(E)$ is a subbundle of $T\mathfrak{so}(E)|_{\Sigma(E)}$ and that we have

$$T\mathfrak{so}(E)|_{\Sigma(E)} = \pi_\Sigma^*\mathfrak{so}(E) \oplus \mathcal{H}_{|\Sigma(E)} = \mathfrak{so}_-(\pi_\Sigma^*E) \oplus \mathfrak{so}_+(\pi_\Sigma^*E) \oplus \mathcal{H}_{|\Sigma(E)} \quad (54)$$

$$= T\Sigma(E) \oplus \mathfrak{so}_+(\pi_\Sigma^*E) \quad (55)$$

with²⁵

$$\mathfrak{so}_+(\pi_\Sigma^*E)_J = \mathfrak{so}_+(E_x, J) := \{A \in \mathfrak{so}(E_x) | [A, J] = 0\}$$

$$\mathfrak{so}_-(\pi_\Sigma^*E)_J = \mathfrak{so}_-(E_x, J) := \{A \in \mathfrak{so}(E_x) | AJ + JA = 0\} = \mathcal{V}_J^\Sigma$$

²⁴we denote by the same letter the fibration $p: E \rightarrow M$ and all its "tensorial extensions": $p: \text{End}(E) \rightarrow M$, $p: \mathfrak{so}(E) \rightarrow M$, etc..

²⁵using the notations defined in section 3.1.2 (i.e. the definition of $\mathfrak{so}_j^{\mathbb{C}}(J)$ for $j \in \mathbb{Z}/2\mathbb{Z}$).

for all $J \in \Sigma(E)$ (and where $x = \pi_\Sigma(J)$). In other words, π_Σ^*E is canonically endowed with a complex structure: $\mathcal{I}_J = J, \forall J \in N_\Sigma$, and this complex structure defines the two spaces $\mathfrak{so}_\pm(\pi_\Sigma^*E)$ by

$$\mathfrak{so}_\pm(\pi_\Sigma^*E) = \mathfrak{so}_\pm(\pi_\Sigma^*E, \mathcal{I}).$$

Now given a section $J \in \mathcal{C}(\pi_\Sigma)$, then we consider the vertical part of the rough Laplacian $\nabla^* \nabla J$, in the decomposition (54): $(\nabla^* \nabla J)^{\mathcal{V}^\Sigma} = \frac{1}{2} J[J, \nabla^* \nabla J]$. We will see in section 4.3.2 that this is in fact exactly the vertical tension field of J in N_Σ :

$$\tau^v(J) = \frac{1}{2} J[J, \nabla^* \nabla J].$$

In particular, we recover the definition of vertical harmonicity used in [34] and [10].

Example 4.3 The twistor bundle $\mathcal{Z}_{2k}(E)$ of a Riemannian vector bundle.

Let $p: (E, \nabla, \langle \cdot, \cdot \rangle) \mapsto (M, g)$ be a Riemannian vector bundle of rank $2n$. Then we consider more generally the bundle of order $2k$ isometries $\mathcal{U}_{2k}(E)$ as well as its subbundles $\mathcal{U}_{2k}^*(E)$ and $\mathcal{Z}_{2k}(E)$. Let us fix $\alpha \in \pi_0(\mathcal{Z}_{2k}(\mathbb{R}^{2n}))$ and consider the component $\mathcal{Z}_{2k}^\alpha(E) := N_{\mathcal{Z}}$. We have a natural fibration $\pi_{\mathcal{Z}}: \mathcal{Z}_{2k}^\alpha(E) \rightarrow M$. The vertical space is given by

$$\forall J \in N_{\mathcal{Z}}, \quad \mathcal{V}_J^{\mathcal{Z}} = T\mathcal{Z}_{2k}^\alpha(E_x) = \left(\bigoplus_{j \in \mathbb{Z}/r\mathbb{Z} \setminus \{0\}} \mathcal{B}_j^{\mathbb{C}}(E_x, J) \right) \cap \text{End}(E_x) = J \cdot \mathfrak{so}_*(E_x, J) \quad (56)$$

according to section 3.1.2 (more particularly equation (43)) and where

$$\mathfrak{so}_*(E_x, J) := \left(\bigoplus_{j \in \mathbb{Z}/r\mathbb{Z} \setminus \{0\}} \mathfrak{so}_j^{\mathbb{C}}(E_x, J) \right) \cap \mathfrak{so}(E_x).$$

The metric connection ∇ gives us a splitting: $T\mathcal{Z}_{2k}^\alpha(E) = \mathcal{V}^{\mathcal{Z}} \oplus \mathcal{H}^{\mathcal{Z}}$. Indeed we have the following splitting (coming from ∇)

$$TSO(E) = \mathcal{V}^{SO(E)} \oplus \mathcal{H} \quad (57)$$

where $\mathcal{V}_J^{SO(E)} = T_J SO(E_x) = J \cdot \mathfrak{so}(E_x)$ (since $0 = \nabla(J^t J) = (\nabla J)^t J + J^t (\nabla J) \implies \nabla J \in T_J SO(E_x)$). Then for all (local) section $J: U \subset M \rightarrow N_{\mathcal{Z}}$, we have

$$0 = \nabla J^{2k} = \sum_{p+l=2k-1} J^p (\nabla J) J^l$$

so that according to (41), $\nabla J \in \mathcal{V}^{\mathcal{Z}}$ and thus in the decomposition (57), we have $[dJ]_{\mathcal{V}^{so(E)}} \in \mathcal{V}^{\mathcal{Z}}$ and hence $[dJ]_{\mathcal{H}} = dJ - [dJ]_{\mathcal{V}^{so(E)}} \in TN_{\mathcal{Z}}$ which leads to

$$T\mathcal{Z}_{2k}^\alpha(E) = \mathcal{V}^{\mathcal{Z}} \oplus \mathcal{H}|_{\mathcal{Z}_{2k}^\alpha(E)}. \quad (58)$$

Then we can endow $N_{\mathcal{Z}}$ with the metric defined as in (52) and where the fibre $\langle \cdot, \cdot \rangle_{\mathcal{V}^{\mathcal{Z}}}$ is induced by the trace metric (53), for which we have $\mathcal{H}^{\mathcal{Z}} = \mathcal{V}^{\mathcal{Z}\perp}$.

Furthermore let us remark that $T\mathcal{Z}_{2k}^\alpha(E)$ is a subbundle of $TSO(E)|_{\mathcal{Z}_{2k}^\alpha(E)}$ and that we have

$$\begin{aligned} TSO(E)|_{\mathcal{Z}_{2k}^\alpha(E)} &= \mathcal{B}_0(\pi_{\mathcal{Z}}^*E) \oplus \mathcal{B}_*(\pi_{\mathcal{Z}}^*E) \oplus \mathcal{H}|_{\mathcal{Z}_{2k}^\alpha(E)} \\ &= \mathcal{B}_0(\pi_{\mathcal{Z}}^*E) \oplus T\mathcal{Z}_{2k}^\alpha(E) \end{aligned}$$

where ²⁶

$$\begin{aligned}\mathcal{B}_0(\pi_{\mathcal{Z}}^*E)_J &= \mathcal{B}_0(E_x, J) \quad \text{and} \\ \mathcal{B}_*(\pi_{\mathcal{Z}}^*E)_J &= \mathcal{B}_*(E_x, J) := \left(\bigoplus_{j \in \mathbb{Z}/r\mathbb{Z} \setminus \{0\}} \mathcal{B}_j^{\mathbb{C}}(E_x, J) \right) \cap \text{End}(E_x) = \mathcal{V}_J^{\mathcal{Z}}\end{aligned}$$

for all $J \in \mathcal{Z}_{2k}^\alpha(E)$. In other words, $\pi_{\mathcal{Z}}^*E$ is canonically endowed with a $2k$ -structure: $\mathcal{I}_J = J$, $\forall J \in N_{\mathcal{Z}}$, and this $2k$ -structure defines the spaces $\mathcal{B}_j^{\mathbb{C}}(\pi_{\mathcal{Z}}^*E) := \mathcal{B}_j^{\mathbb{C}}(\pi_{\mathcal{Z}}^*E, \mathcal{I})$. Now let us precise the relation between $SO(E)$ and $\mathfrak{so}(E)$ and in particular the relation $T_J SO(E_x) = J \cdot \mathfrak{so}(E_x)$. For $J \in SO(E)$, let

$$L_J: A \in \text{End}(E_x) \longmapsto J \cdot A \in \text{End}(E_x)$$

be the left multiplication by J in $\text{End}(E_x)$, with $x = p(J)$. Let us still denote by \mathcal{I} , the tautological section of $p^*SO(E)$ defined by $\mathcal{I}_J = J$, $\forall J \in SO(E)$, and whose restriction to $N_{\mathcal{Z}}$ is our canonical $2k$ -structure \mathcal{I} on $\pi_{\mathcal{Z}}^*E$. Then let $L_{\mathcal{I}}: SO(E) \rightarrow \text{Aut}(\text{End}(p^*E))$ be the section of the bundle of linear automorphism of the vector bundle $\text{End}(p^*E)$ defined by

$$L_{\mathcal{I}}: J \in SO(E) \longmapsto L_J \in \text{Aut}(\text{End}(E_{p(J)}))$$

or more concretely

$$L_{\mathcal{I}}: (J, A) \in \text{End}(p^*E) \longmapsto (J, J \cdot A) \in \text{End}(p^*E).$$

Then we have

$$\mathcal{V}^{SO(E)} = L_{\mathcal{I}}(\mathfrak{so}(p^*E)) \quad \text{and} \quad \mathcal{B}_j^{\mathbb{C}}(\pi_{\mathcal{Z}}^*E) = L_{\mathcal{I}}(\mathfrak{so}_j^{\mathbb{C}}(\pi_{\mathcal{Z}}^*E))$$

which we will denote more simply by

$$\mathcal{V}^{SO(E)} = \mathcal{I} \cdot \mathfrak{so}(p^*E) \quad \text{and} \quad \mathcal{B}_j^{\mathbb{C}}(\pi_{\mathcal{Z}}^*E) = \mathcal{I} \cdot \mathfrak{so}_j^{\mathbb{C}}(\pi_{\mathcal{Z}}^*E).$$

Example 4.4 The Twistor subbundle $\mathcal{Z}_{2k,j}^\alpha(E)$.

Let us consider the previous example and let us suppose that there exists a (global) section J_j of $(\mathcal{Z}_{2k}(E))^j = \mathcal{U}_{p'}^{j,\alpha}(E)$ for some $j \in \mathbb{Z}$ and $p' = \frac{2k}{(2k,j)}$. Let us consider the subbundle

$$N_{\mathcal{Z}}^j := \mathcal{Z}_{2k,j}^\alpha(E, J_j) = \{J \in \mathcal{Z}_{2k}^\alpha(E) \mid J^j = J_j\}$$

for which we have the natural fibration $\pi_{\mathcal{Z}}^j: \mathcal{Z}_{2k,j}^\alpha(E, J_j) \rightarrow M$. The vertical space is given by

$$\forall J \in N_{\mathcal{Z}}^j, \quad \mathcal{V}_J^{\mathcal{Z},j} = T_J \mathcal{Z}_{2k,j}^\alpha(E_x, J_j) = \left(\bigoplus_{q=1}^{(r,j)-1} \mathcal{B}_{pq}^{\mathbb{C}}(E_x, J) \right) \cap \text{End}(E_x) = J \cdot \mathfrak{u}_{j-1}^*(E_x, J)$$

according to (46)²⁷, where

$$\mathfrak{u}_{j-1}^*(E_x, J) = \left(\bigoplus_{q=1}^{(r,j)-1} \mathfrak{so}_{pq}^{\mathbb{C}}(E_x, J) \right) \cap \mathfrak{so}(E_x) = \mathfrak{u}_{j-1}(E_x, J) / \mathfrak{u}_0(E_x, J) = \mathfrak{so}_0(E_x, J^j) / \mathfrak{so}_0(E_x, J).$$

Furthermore, differentiating the definition equation of $\mathcal{Z}_{2k,j}^\alpha(E, J_j): J^j = J_j$, we obtain: for all (local) section J of $\pi_{\mathcal{Z}}^j$,

$$\nabla J^j = \sum_{l+q=j-1} J^l \nabla J J^q = \nabla J_j \tag{59}$$

²⁶still with the notation defined in section 3.1.2

²⁷and with the notations of section 3.1.3, in particular $p = \frac{r}{(r,j)}$.

so that

$$\nabla J \in \mathcal{V}^{\mathcal{Z},j} \iff \nabla J_j = 0,$$

therefore in general, we have $\nabla J \notin \mathcal{V}^{\mathcal{Z},j}$. We will simply set

$$\mathcal{H}^{\mathcal{Z},j} = \mathcal{V}^{\mathcal{Z},j\perp} \cap T\mathcal{Z}_{2k,j}^\alpha(E, J_j).$$

Then the splitting

$$T\mathcal{Z}_{2k,j}^\alpha(E, J_j) = \mathcal{V}^{\mathcal{Z},j} \oplus \mathcal{H}^{\mathcal{Z},j}$$

do not correspond to the splitting (57) or equivalently to (58), in general. In other words, the connection in $\pi_{\mathcal{Z}}$ defined by the horizontal distribution $\mathcal{H}|_{N_{\mathcal{Z}}^j}$ is not reducible to a connection in $\pi_{\mathcal{Z}}^j$ (which could only be $\mathcal{H}^{\mathcal{Z},j}$): it happens if and only if \mathcal{H} is tangent to $N_{\mathcal{Z}}^j$. Besides we have two different ways to decompose the orthogonal of $\mathcal{V}^{\mathcal{Z},j}$ in $TN_{\mathcal{Z}}$, using the decompositions $TN_{\mathcal{Z}} = \mathcal{V}^{\mathcal{Z}} \oplus \mathcal{H}^{\mathcal{Z}}$ or $TN_{\mathcal{Z}}|_{N_{\mathcal{Z}}^j} = TN_{\mathcal{Z}}^j \oplus TN_{\mathcal{Z}}^{j\perp}$:

$$\begin{aligned} TN_{\mathcal{Z}}|_{N_{\mathcal{Z}}^j} &= \mathcal{V}^{\mathcal{Z},j} \oplus \mathcal{V}^{\mathcal{Z},j\perp} \\ &= \mathcal{V}^{\mathcal{Z},j} \oplus \mathcal{V}^{\mathcal{Z},j\perp} \cap \mathcal{V}^{\mathcal{Z}} \oplus \mathcal{H}|_{N_{\mathcal{Z}}^j} \\ &= \underbrace{\mathcal{V}^{\mathcal{Z},j} \oplus \mathcal{H}_{\mathcal{Z}}^j}_{TN_{\mathcal{Z}}^j} \oplus TN_{\mathcal{Z}}^{j\perp} \end{aligned}$$

In particular, we have for any (local) section $J: U \subset M \rightarrow N_{\mathcal{Z}}^j$

$$[dJ]_{\mathcal{V}^{\mathcal{Z},j}} = [\nabla J]_{\mathcal{V}^{\mathcal{Z},j}} = \text{pr}_{\mathcal{V}^{\mathcal{Z},j}}^{\mathcal{V}^{\mathcal{Z}}}(\nabla J)$$

where $[\]_{\mathcal{V}^{\mathcal{Z},j}}: TN_{\mathcal{Z}} \rightarrow \mathcal{V}^{\mathcal{Z},j}$ and $\text{pr}_{\mathcal{V}^{\mathcal{Z},j}}^{\mathcal{V}^{\mathcal{Z}}}: \mathcal{V}^{\mathcal{Z}} \rightarrow \mathcal{V}^{\mathcal{Z},j}$ are resp. the orthogonal projections. Moreover, let us decompose $T\mathcal{S}\mathcal{O}(E)|_{\mathcal{Z}_{2k,j}^\alpha(E)}$ into an orthogonal sum making appear the vertical subbundle $\mathcal{V}^{\mathcal{Z},j}$ of $N_{\mathcal{Z}}^j$:

$$\begin{aligned} T\mathcal{S}\mathcal{O}(E)|_{N_{\mathcal{Z}}^j} &= \mathcal{B}_0(\pi_{\mathcal{Z}}^{j*}E, \mathcal{I}) \oplus \mathcal{B}_*(\pi_{\mathcal{Z}}^{j*}E, \mathcal{I}) \oplus \mathcal{H}|_{N_{\mathcal{Z}}^j} \\ &= \underbrace{\mathcal{B}_0(\pi_{\mathcal{Z}}^{j*}E, \mathcal{I}) \oplus \mathcal{V}^{\mathcal{Z},j}}_{\mathcal{I}\cdot\text{so}_*(\pi_{\mathcal{Z}}^{j*}E, \mathcal{I}^j)} \oplus \underbrace{\mathcal{B}_*(\pi_{\mathcal{Z}}^{j*}E, \mathcal{I})}_{\mathcal{V}^{\mathcal{Z},j\perp} \cap \mathcal{V}^{\mathcal{Z}}} \oplus \mathcal{H}|_{N_{\mathcal{Z}}^j}. \end{aligned}$$

Now let us see how we can determine $\mathcal{H}^{\mathcal{Z},j}$ from the section J_j . First we remark that $\mathcal{H}^{\mathcal{Z},j} \cap \mathcal{V}^{\mathcal{Z}} = \{0\}$ (indeed $\ker d\pi_{\mathcal{Z}} \cap \mathcal{H}^{\mathcal{Z},j} \subset \ker d\pi_{\mathcal{Z}} \cap TN_{\mathcal{Z}}^j = \mathcal{V}^{\mathcal{Z},j}$ and of course $\mathcal{V}^{\mathcal{Z},j} \cap \mathcal{H}^{\mathcal{Z},j} = \{0\}$). Therefore $\mathcal{H}^{\mathcal{Z},j}$ is a vector subbundle of $(\mathcal{V}^{\mathcal{Z},j\perp} \cap \mathcal{V}^{\mathcal{Z}}) \oplus \mathcal{H}$ which satisfies $(\mathcal{V}^{\mathcal{Z},j\perp} \cap \mathcal{V}^{\mathcal{Z}}) \cap \mathcal{H}^{\mathcal{Z},j} = \{0\}$. Thus $\mathcal{H}^{\mathcal{Z},j}$ is the graph of some linear map²⁸ $\Gamma: \mathcal{H} \rightarrow \mathcal{V}^{\mathcal{Z},j\perp} \cap \mathcal{V}^{\mathcal{Z}}$,²⁹

$$\text{Id} + \Gamma: W \in \mathcal{H} \mapsto W + \Gamma(W) \in \mathcal{H} \oplus (\mathcal{V}^{\mathcal{Z},j\perp} \cap \mathcal{V}^{\mathcal{Z}})$$

has $\mathcal{H}^{\mathcal{Z},j}$ as image.

Let us concentrate ourself on (59). ∇J is in $\mathcal{V}^{\mathcal{Z}}$ so that we can write it $\nabla J = \sum_{i=1}^{r-1} JA_i$ with $A_i \in \mathfrak{so}_i^{\mathbb{C}}(E, J)$, according to (56). Then we have $\forall i \in \{1, \dots, r-1\}$,

$$\sum_{l+q=j-1} J^q (JA_i) J^l = \sum_{l=0}^{j-1} J^{j-1-l} \omega_r^l JA_i = \frac{1 - (\omega_r^i)^j}{1 - \omega_r^i} J^j A_i$$

²⁸i.e. a morphism of vector bundle

²⁹In the following reasoning, we will forget the index " $N_{\mathcal{Z}}^j$ " in $\mathcal{H}|_{N_{\mathcal{Z}}^j}$ to do not weigh down the equations. The right notation will reappear in the final equation.

so that

$$\sum_{l+q=j-1} J^q(\nabla J)J^l = J^j \left(\sum_{\substack{i=1 \\ i \notin p.\mathbb{Z}}}^{r-1} \frac{1 - (\omega_r^i)^j}{1 - \omega_r^i} A_i \right)$$

where as usual $p = \frac{r}{(r, j)}$ is the order of ω_r^j . In particular, we remark that (with obvious notation)³⁰

$$\sum_{l+q=j-1} L(J^l) \circ R(J^q)^{-1} : \mathcal{B}_*(E, J) \longmapsto \mathcal{B}_*(E, J_j) \quad (60)$$

is a surjective map with kernel³¹

$$\left(\bigoplus_{i \in p.\mathbb{Z}_r \setminus \{0\}} \mathcal{B}_i^{\mathbb{C}}(E, J) \right) \cap \text{End}(E) = J^* \mathcal{V}^{\mathbb{Z}, j},$$

so that it induces an isomorphism from

$$J.\mathfrak{so}_*(E, J_j) = J^*(\mathcal{V}^{\mathbb{Z}, j^\perp} \cap \mathcal{V}^{\mathbb{Z}})$$

onto $\mathcal{B}_*(E, J_j)$. Let us denote by $P^j(J)$ the surjective map (60) and by $P^j(J)^{-1}$ the inverse map of the isomorphism induced on $J.\mathfrak{so}_*(E, J_j)$. Then we have

$$P^j(J)(\nabla J) = \nabla J_j$$

so that $[\nabla J]^{\mathcal{V}^{\mathbb{Z}, j^\perp} \cap \mathcal{V}^{\mathbb{Z}}} = P^j(J)^{-1}(\nabla J_j)$, but we have $\nabla J = [dJ]^{\mathcal{V}^{\mathbb{Z}}}$, and therefore

$$[dJ]^{\mathcal{V}^{\mathbb{Z}, j^\perp} \cap \mathcal{V}^{\mathbb{Z}}} = P^j(J)^{-1}(\nabla J_j). \quad (61)$$

On the other hand, $d\pi^{\mathbb{Z}} \circ dJ = \text{Id}_{TM}$ so that $d\pi^{\mathbb{Z}} \circ [dJ]^{\mathcal{H}} = \text{Id}_{TM}$, which with (61) allows to conclude that

$$\Gamma = P^j(J)^{-1} \circ (\nabla J_j) \circ d\pi_{\mathcal{H}}^{\mathbb{Z}}$$

that is to say, for all $W \in \mathcal{H}_{|N_{\mathbb{Z}}^j}$

$$\Gamma(W) = P^j(J_0)^{-1} \cdot \nabla_{(\pi^{\mathbb{Z}})_*} W J_j$$

where $W = (J_0, W_{J_0})$, $J_0 \in N_{\mathbb{Z}}^j$, $W_{J_0} \in \mathcal{H}_{J_0}$.

4.1.3 Ψ -torsion, Ψ -difference tensor, and curvature of a Pfaffian system

Ψ -torsion, Ψ -difference. Let us consider a vector bundle morphism

$$\begin{array}{ccc} TM & \xrightarrow{\Psi} & (E, \nabla) \\ \downarrow & & \downarrow \\ M & \xrightarrow{\psi} & N \end{array},$$

³⁰remark that $\mathcal{B}_*(E, J_j) = J^j.\mathfrak{so}_*(E, J^j) = J^j. \left(\left(\bigoplus_{i \in \mathbb{Z}_r \setminus p.\mathbb{Z}_r} \mathfrak{so}_i^{\mathbb{C}}(E, J) \right) \cap \mathfrak{so}(E, J) \right)$

³¹Obviously, since J is a local section, everything is local here and E must be replaced by $E_U := p^{-1}(U)$, but we do not want to weigh down the notations.

∇ being a connections on the vector bundle E . Then the Ψ -torsion of ∇ is the ψ^*E -valued 2-form on M ,

$$T^\Psi(X, Y) = \nabla_X(\Psi Y) - \nabla_Y(\Psi X) - \Psi[X, Y] = d^\nabla \Psi(X, Y) \quad \forall X, Y \in \mathcal{C}(TM).$$

Let us give now some examples.

Example 4.5 Let N be a manifold and suppose that we have a splitting $TN = \mathcal{V} \oplus \mathcal{H}$ and suppose also that the vertical bundle \mathcal{V} is endowed with a covariant derivative ∇^c and let $\phi: TN \rightarrow \mathcal{V}$ be the projection (morphism) on \mathcal{V} along \mathcal{H} , then we can speak about the ϕ -torsion of ∇^c , $T^\phi = d^{\nabla^c} \phi$.

$$\begin{array}{ccc} TN & \xrightarrow{\phi} & (\mathcal{V}, \nabla^c) \\ \downarrow & & \downarrow \\ N & \xrightarrow{\text{Id}} & N \end{array},$$

Example 4.6 Let $s: M \rightarrow (N, \nabla)$ be a map from a manifold M into an affine manifold (N, ∇) and suppose that we have a splitting $TN = \mathcal{V} \oplus \mathcal{H}$, then let us consider the morphism of bundle

$$\begin{array}{ccc} TM & \xrightarrow{d^v s} & (\mathcal{V}, \nabla^v) \\ \downarrow & & \downarrow \\ M & \xrightarrow{s} & N \end{array} \quad \text{where } \nabla^v \text{ is the vertical part} \\ \text{of the linear connection } \nabla.$$

Then the *vertical s -torsion* of N is $T^s := T^{d^v s} = d^{\nabla^v} d^v s$.

Example 4.7 In particular let us take $s = \text{Id}_N$ (in the previous example) and thus $d^v s = \phi$ the projection on \mathcal{V} and then the ϕ -torsion of ∇^v (or Id_N -torsion of N) is the *vertical torsion* in \mathcal{V} : $T^v = d^{\nabla^v} \phi$.

Now for any map $s: M \rightarrow N$ we have

$$T^s = s^* T^v.$$

We will say that s is *vertically torsion free* if $T^s = 0$.

Example 4.8 Let $E \rightarrow N$ be a vector bundle and let us suppose that we have a isomorphism of bundle $\psi: TN \rightarrow E$ (over Id_N). Let ∇ be a connection in E . Then we have $T^\psi = \psi \circ T$, where T is the torsion of the linear connection $\psi^{-1} \circ \nabla \circ \psi$ on N . ■

Now, we define the ψ -difference and the ψ -equivalence.

Definition 4.3 Let $E \rightarrow N$ be a vector bundle and let us suppose that we have a morphism of bundle $\psi: TN \rightarrow E$ (over Id_N). Given two connections ∇ and ∇' in E , the ψ -**difference tensor** S^ψ for the pair (∇, ∇') is defined by

$$S^\psi(X, Y) = \nabla_X(\psi Y) - \nabla'_X(\psi Y) = (\nabla - \nabla')_X(\psi Y).$$

Then S^ψ is symmetric precisely when ∇ and ∇' have the same ψ -torsion. On the other hand, if S^ψ is skew-symmetric we will say (following [51]) that ∇ and ∇' are ψ -**equivalent**: it means that these have the same ψ -geodesics, a ψ -geodesic of ∇ being a path $y(t)$ in N solution of the equation

$$\nabla_{y'(t)}(\psi y'(t)) = 0$$

(if ψ is an isomorphism then ψ -geodesics of ∇ are precisely the geodesics of $\psi^{-1} \circ \nabla \circ \psi$).

Curvature of a Pfaffian system

Definition 4.4 Let \mathcal{P} be a Pfaffian system on the manifold N . Then for any local sections of \mathcal{P} , $X, Y: (N, a) \rightarrow \mathcal{P}$, defined in the neighbourhood of $a \in N$, the image $([X, Y]_a)_{\mathcal{V}}$ of $[X, Y]_a$ by the canonical projection $T_a N \rightarrow \mathcal{V}_a = T_a N / \mathcal{P}_a$, depends only on the values X_a, Y_a at $a \in N$, of the vector fields X, Y . We define the curvature of \mathcal{P} as the tensor $R \in \Lambda^2 \mathcal{P}^* \otimes \mathcal{V}$,

$$R_a(X_a, Y_a) := -([X, Y]_a)_{\mathcal{V}}.$$

Definition 4.5 Let N be a manifold endowed with a Pfaffian system \mathcal{V} ("vertical subbundle") and let us suppose that \mathcal{V} admits a connection i.e. a complement \mathcal{H} ("horizontal subbundle"): $TN = \mathcal{V} \oplus \mathcal{H}$. Then \mathcal{V} is identified to TN/\mathcal{P} so that the curvature of the connection \mathcal{H} becomes the tensor $R \in \Lambda^2 \mathcal{H}^* \otimes \mathcal{V}$ defined by

$$R(X, Y) = -[X, Y]_{\mathcal{V}} \quad \forall X, Y \in \mathcal{C}(\mathcal{H})$$

the subscripts " \mathcal{V} " designing the \mathcal{V} -component along \mathcal{H} .

Convention We will often extend R to the corresponding horizontal 2-form on N , still denoted by R : $R \in \Lambda^2 T^* N \otimes \mathcal{V}$ such that $R(X, Y) = 0$ if X or $Y \in \mathcal{V}$.

Proposition 4.1 Let N be a manifold and suppose that we have a splitting $TN = \mathcal{V} \oplus \mathcal{H}$ and suppose also that the vertical bundle \mathcal{V} is endowed with a covariant derivative ∇^c , then we have

$$T_{|\mathcal{H} \times \mathcal{H}}^c = R_{\mathcal{H}},$$

$R_{\mathcal{H}}$ being the curvature of \mathcal{H} .

Proof. For any $H_1, H_2 \in \mathcal{H}$, we have $T^c(H_1, H_2) = \nabla_{H_1}^c H_2^v - \nabla_{H_2}^c H_1^v - [H_1, H_2]^v = -[H_1, H_2]^v = R_{\mathcal{H}}(H_1, H_2)$. \square

Proposition 4.2 Let $\pi: Q \rightarrow Q/H = M$ be a H -principal bundle endowed with a connection 1-form $\omega: TQ \rightarrow \mathfrak{h}$. Let $\mathcal{H} = \ker \omega \subset TQ$ be the corresponding horizontal subbundle. Let be $\Omega = d\omega + \frac{1}{2}[\omega \wedge \omega]$ the curvature 2-form. Then we have

$$(R_{\mathcal{H}})_q(X, Y) = q.\Omega_q(X, Y) \quad \forall q \in Q, \forall X, Y \in \mathcal{H}_q$$

$R_{\mathcal{H}}$ being the curvature of the connection \mathcal{H} . In other words, we have

$$R_{\mathcal{H}} = \Omega^*$$

where $\Omega_q^* = q.\Omega_q$.

4.2 Harmonic sections of homogeneous fibre bundles

In this section, we study fibre bundles $\pi: N \rightarrow M$ for which the fibre is a homogeneous space H/K . To do that, we follow, in subsections 4.2.1 and 4.2.2, the exposition of [51]. Next, we present a generalisation of the results (of [51]) to non section maps in the end of 4.2.2. Finally we study the homogeneous fibre bundle reductions in 4.2.3.

4.2.1 Definitions and Geometric properties

Let $\pi_M: Q \rightarrow M$ be a principal H -bundle, with H a Lie group. Let K be a Lie subgroup of H and $N = Q/K$. Then the map $\pi_N: Q \rightarrow N$ is a principal K -bundle and we have $\pi_M = \pi \circ \pi_N$ where $\pi: N \rightarrow M$ is a fibre bundle with fibre H/K , which is naturally isomorphic to the associated bundle $Q \times_H H/K$. We assume the following hypothesis

- (i) H/K is reductive: $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{p}$, and $\text{Ad}K(\mathfrak{p}) \subset \mathfrak{p}$, where \mathfrak{h} and \mathfrak{k} are respectively the Lie algebras of H and K .
- (ii) M is endowed with a Riemannian metric g
- (iii) H/K is Riemannian: there exists a H -invariant Riemannian metric on H/K (equivalently an $\text{Ad}K$ -invariant (positive definite) inner product on \mathfrak{p}). Equivalently $\text{Ad}_\mathfrak{p}K$ is compact.
- (iv) The principal H -bundle $\pi_M: Q \rightarrow M$ is endowed with a connection. We denote by ω the corresponding \mathfrak{h} -valued connection form on Q .

Then the splitting $TQ = \mathcal{V}_0 \oplus \mathcal{H}_0$ defined by ω ($\mathcal{V}_0 = \ker d\pi_M$, $\mathcal{H}_0 = \ker \omega$) gives rise by $d\pi_N$, to the following decomposition $TN = \mathcal{V} \oplus \mathcal{H}$, where $\mathcal{V} = \ker d\pi = d\pi_N(\mathcal{V}_0)$ and $\mathcal{H} = d\pi_N(\mathcal{H}_0)$. Let $\mathfrak{p}_Q := Q \times_K \mathfrak{p} \rightarrow N$ be the vector bundle associated to $\pi_N: Q \rightarrow N$ with fibre \mathfrak{p} . Let us denote by $[q, a] \in \mathfrak{p}_Q$ the element defined by $(q, a) \in Q \times \mathfrak{p}$. Then we have the following vector bundle isomorphism

$$I: \begin{array}{ccc} \mathcal{V} & \longrightarrow & \mathfrak{p}_Q \\ d\pi_N(q.a) & \longmapsto & [q, a] \end{array}$$

where $q \in Q$, $a \in \mathfrak{p}$ and as usual $q.a = \frac{d}{dt}|_{t=0} q \cdot \exp(ta) \in T_q Q$. Decomposing $\omega = \omega_{\mathfrak{h}} + \omega_{\mathfrak{p}}$ following $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{p}$, then since H/K is reductive, $\omega_{\mathfrak{p}}$ is a K -equivariant ($\omega_{\mathfrak{p}}(X.h) = \text{Ad}h \omega_{\mathfrak{p}}(X)$) and π_N -horizontal ($\omega_{\mathfrak{p}}|_{\mathcal{V}_0} = 0$) \mathfrak{p} -valued 1-form on Q and hence projects to a \mathfrak{p} -valued 1-form ϕ on N :

$$\phi(d\pi_N(X)) = [q, \omega_{\mathfrak{p}}(X)].$$

Then we have

$$\phi|_{\mathcal{V}} = I \quad \text{and} \quad \ker \phi = \mathcal{H}.$$

We can now construct a Riemannian metric h on N :

$$h = \pi^*g + \langle \phi, \phi \rangle \tag{62}$$

where $\langle \cdot, \cdot \rangle$ is the fibre metric induced on \mathfrak{p}_Q by the inner product on \mathfrak{p} .

In the same way, let Φ be the \mathfrak{p}_Q -valued 2-form on N defined by the component $\Omega_{\mathfrak{p}}$ of the curvature form Ω of ω . Since $\Omega_{\mathfrak{p}}$ is π_M -horizontal ($\Omega(X, Y) = 0$ if $X \in \mathcal{V}_0$ or $Y \in \mathcal{V}_0$), then Φ is π -horizontal: $\Phi(X, Y) = 0$, if $X \in \mathcal{V}$ or $Y \in \mathcal{V}$.

Remark 4.1 In [51], \mathfrak{p}_Q is called the *canonical bundle*, I the *canonical isomorphism*, ϕ the *homogeneous connection form*, and Φ the *homogeneous curvature form*.

Definition 4.6 We will call the of the following datas (Q, H, K, ω) a *homogeneous fibre bundle structure* on $\pi: N \rightarrow M$ (or on N , when the fibration π is considered as implicitly given). Moreover $\pi: N \rightarrow M$ will then be sayed to be a *homogeneous fibre bundle*.

The 1-form $\omega_{\mathfrak{k}}$ (which is a connection form in π_N because H/K is reductive) defines a connection in π_N called the *canonical connection*. This connection induces a covariant derivative ∇^c in the associated bundle \mathfrak{p}_Q , with respect to which the fibre metric is parallel. ∇^c defines a exterior derivative d^c on the space of \mathfrak{p}_Q -valued differential forms on N . This allows us to define the *canonical torsion* T^c which is nothing but the ϕ -torsion of ∇^c (see section 4.1.3)

$$T^c(A, B) = d^c\phi(A, B) = \nabla_A^c(\phi B) - \nabla_B^c(\phi A) - \phi[A, B], \quad \forall A, B \in \mathcal{C}(TN) \quad (63)$$

Let $\mathfrak{h}_Q := Q \times_H \mathfrak{h} \rightarrow M$ be the vector bundle associated to π_M with fibre \mathfrak{h} , and in the same way $\mathfrak{k}_Q := Q \times_K \mathfrak{k} \rightarrow N$ the bundle associated to π_N with fibre \mathfrak{h} . Then we have

$$\pi^*\mathfrak{h}_Q = \mathfrak{k}_Q \oplus \mathfrak{p}_Q.$$

The Lie bracket of \mathfrak{h} induces a bracket on the fibres of \mathfrak{h}_Q , and those of $\pi^*\mathfrak{h}_Q$, which we continue to denote by $[\cdot, \cdot]$, and we denote also its \mathfrak{p}_Q -component (when there is no risk of confusion) by $[\cdot, \cdot]_{\mathfrak{p}}$ (otherwise we denote it by $[\cdot, \cdot]_{\mathfrak{p}_Q}$). Taking the \mathfrak{p} -component of the structure equation $d\omega = \Omega - \frac{1}{2}[\omega \wedge \omega]$ and then projecting on N , we obtain *the homogeneous structure equation*:

$$T^c = \Phi - \frac{1}{2}[\phi \wedge \phi]_{\mathfrak{p}} \quad (64)$$

and thus

$$\begin{aligned} T^c_{|\mathcal{V} \times \mathcal{V}} &= -[I \cdot, I \cdot]_{\mathfrak{p}}, & T^c_{|\mathcal{V} \times \mathcal{H}} &= 0 \\ T^c_{|\mathcal{H} \times \mathcal{H}} &= \Phi_{|\mathcal{H} \times \mathcal{H}}. \end{aligned}$$

In particular, T^c is horizontal if and only if H/K is a (locally) symmetric space, and in this case

$$T^c = \Phi \quad (65)$$

Remark 4.2 According to (64) and (63), for all $X, Y \in \mathcal{H}$, (extended to vector fields in N denoted by the same letters), we have

$$\Phi(X, Y) = T^c(X, Y) = -\phi([X, Y])$$

so that

$$\Phi = R_{\mathcal{H}},$$

according to definition 4.5. The homogeneous curvature form is nothing but the curvature of the connection \mathcal{H} . •

Now, let U be the \mathfrak{p}_Q -valued symmetric bilinear form defined on \mathfrak{p}_Q by:

$$\langle U(a, b), c \rangle = \langle [c, a]_{\mathfrak{p}}, b \rangle + \langle a, [c, b]_{\mathfrak{p}} \rangle \quad (66)$$

where $\langle \cdot, \cdot \rangle$ is the fibre metric, and $a, b, c \in \mathfrak{p}_Q$. Let us set

$$B = U + [\cdot, \cdot]_{\mathfrak{p}} \quad (67)$$

which is a \mathfrak{p}_Q -valued bilinear form on \mathfrak{p}_Q , whose the symmetric and skew symmetric components are respectively U and $[\cdot, \cdot]_{\mathfrak{p}}$. U vanishes if and only if H/K is naturally reductive and B if and only if H/K is (locally) symmetric. Then denoting by ∇ the Levi-Civita connection on N , we have:

Theorem 4.2 [51] *Let us consider the difference tensor:*

$$S(A, B) = \phi(\nabla_A B) - \nabla_A^c(\phi B)$$

then we have

$$2S = \phi^*U - T^c = \phi^*B - \Phi.$$

Consequently, $\forall V \in \mathcal{C}(\mathcal{V})$

$$I(\nabla_A^v V) = \nabla_A^c(IV) + \frac{1}{2}B(\phi A, IV). \quad (68)$$

In particular, if H/K is a (locally) symmetric space, we have

$$I\nabla_A^v V = \nabla_A^c(IV).$$

Remark 4.3 If H/K is a symmetric space, under the canonical identification $I: \mathcal{V} \xrightarrow{\cong} \mathfrak{p}_Q$, we have $\nabla^v = \nabla^c$ on \mathcal{V} . More generally the difference between ∇^v and ∇^c looks like to the difference between the Levi-Civita and canonical connections of a reductive Riemannian homogeneous space (see section 1.6).

Moreover, ∇^v is ϕ -equivalent to ∇^c when H/K is naturally reductive, according to (68).

Let ∇^ω be the covariant derivative in the vector bundle \mathfrak{h}_Q (associated to π_M), defined by the connection form ω . Let us decompose (the π -pullback of) ∇^ω following the decomposition $\pi^*\mathfrak{h}_Q = \mathfrak{k}_Q \oplus \mathfrak{p}_Q$, and the \mathfrak{p}_Q -component gives us a connection ∇^p in \mathfrak{p}_Q .

Theorem 4.3 [51] *For all $V \in \mathcal{C}(\mathfrak{p}_Q)$,*

$$\nabla^p V = \nabla^\omega V - [\phi, \alpha]_{\mathfrak{h}}$$

and

$$\nabla^c V = \nabla^p V - [\phi, V]_{\mathfrak{p}} = \nabla^\omega V - [\phi, V].$$

Consequently, ∇^p and ∇^c are ϕ -equivalent (since their ϕ -difference is $[\phi, \phi]_{\mathfrak{p}}$). In particular $\nabla^c = \nabla^p$ if H/K is a (locally) symmetric space.

Example 4.9 Let us consider the situation described by example 4.1 and suppose that $u: L \rightarrow M$ is an isometry. Then if $\pi: N \rightarrow M$ is a homogeneous fibre bundle like above then this is also the case for $u^*\pi: u^*N \rightarrow L$.

Indeed let us set

$$u^*Q = \{(z, q) \in L \times Q, q \in \pi_M^{-1}(\{u(z)\})\} = \bigsqcup_{z \in L} \{z\} \times f(z).H$$

then $u^*\pi_M: (z, q) \in u^*Q \mapsto z \in L$ is a principal H -bundle over L . Then we have $u^*N = u^*Q/K$, and $u^*\pi: u^*N \rightarrow L$ is a fibre bundle with fibre H/K .

Finally we have to define a connection on $u^*\pi_M: u^*Q \rightarrow L$. Let us extend the connection ω , to a connection on $\text{Id}_L \times \pi_M: L \times Q \rightarrow L \times M$ by $\tilde{\omega}_{(z,q)}(dz + dq) = \omega_q(dq)$ and then let us set

$$u^*\omega := \tilde{\omega}|_{T(u^*Q)}.$$

In the same way, the homogeneous connection and curvature forms on u^*N are given respectively by

$$u^*\phi := \tilde{\phi}|_{T(u^*N)} \quad \text{and} \quad u^*\Phi := \tilde{\Phi}|_{T(u^*N) \oplus T(u^*N)}.$$

The canonical torsion T^c on u^*N is also given by $u^*T^c := \tilde{T}^c|_{T(u^*N) \oplus T(u^*N)}$.

Definition 4.7 *Let (L, b) be a Riemannian manifold. We will then say that $f: L \rightarrow N$ is vertically geodesic if $\phi(\nabla df) = 0$, horizontally geodesic if $\pi_*(\nabla df) = 0$, and superflat if its vertical second fundamental form vanishes $\Pi^v(f) := \nabla^v d^v f = 0$.*

4.2.2 Vertical harmonicity equation

We know that the structure group H of $\pi_M: Q \rightarrow M$ is reducible to K (i.e. there exists an K -bundle $\pi'_M: Q' \rightarrow M$) if and only if the associated bundle $\pi: N \rightarrow M$ admits a (global) section $s: M \rightarrow N$ (see [40]) so that there is a one to one correspondance between the K -reductions of π_M and the space of sections $\mathcal{C}(\pi)$.

Let $\omega' = \omega_{\mathfrak{h}|_{TQ'}}$. Then ω' is a connection in π'_M , and ω is reducible if and only if $\omega|_{TQ'} = \omega'$ (see [40]). The reducibility of ω can be characterized as follows.

Proposition 4.3 *The following statements are equivalent:*

- (i) s is horizontal;
- (ii) $s^*\phi = 0$;
- (iii) s is an isometric immersion;
- (iv) ω is reducible

Now we have the following expression of the tension field for sections $s: M \rightarrow N$.

Theorem 4.4 [51] *For all $s \in \mathcal{C}(\pi)$,*

$$I(\tau^v(s)) = -d^*(s^*\phi) + \frac{1}{2}\text{Tr}(s^*\phi^*U)$$

where d^* is the coderivative for $s^*\mathfrak{p}_Q$ -valued differential forms on M relative to the s -pullback of any connection in \mathfrak{p}_Q which is ϕ -equivalent to ∇^c . In particular, if H/K is naturally reductive then s is an harmonic section if and only if $s^*\phi$ is coclosed.

Remark 4.4 If H/K is naturally reductive, to compute the vertical tension field $\tau^v(s) = \text{Tr}(\nabla^v d^v s)$, we can use instead of ∇^v any connection in $\mathcal{V} \cong \mathfrak{p}_Q$ which is ϕ -equivalent to ∇^c .

From the homogeneous structure equation (64), we obtain

$$s^*\Phi = d^c(s^*\phi) + \frac{1}{2}[s^*\phi \wedge s^*\phi]_{\mathfrak{p}},$$

hence every horizontal section is flat (i.e. $s^*\Phi = 0$). Let us introduce the following 3-covariant tensor $\langle s^*\phi \otimes s^*\Phi \rangle$ on M :

$$\langle s^*\phi \otimes s^*\Phi \rangle(X, Y, Z) = \langle s^*\phi(X), s^*\Phi(Y, Z) \rangle,$$

then we have

Theorem 4.5 [51] *For all $s \in \mathcal{C}(\pi)$ we have*

- (i) $\phi(\nabla ds) = \nabla^c(s^*\phi) + \frac{1}{2}s^*\phi^*B - \frac{1}{2}s^*\Phi$.

In particular, if s is vertically geodesic then s is a harmonic section.

- (ii) $2g(\pi_*\nabla ds(X, Y), Z) = \langle s^*\phi \otimes s^*\Phi \rangle(X, Y, Z) + \langle s^*\phi \otimes s^*\Phi \rangle(Y, X, Z)$.

Therefore s is horizontally geodesic if and only if $\langle s^\phi \otimes s^*\Phi \rangle$ is a 3-form on M . In particular, if s is flat then s is horizontally geodesic.*

Theorem 4.6 [51]

(i) The symmetric and skew symmetric components of $\Pi^v s := \nabla^v d^v s$ are given by:

$$I(\Pi^v s) = \phi \circ \nabla ds + \frac{1}{2} s^* \Phi.$$

- (ii) The section s is superflat if and only if s is flat and totally geodesic. In particular, if s is flat then s is totally geodesic if and only if s is super-flat.
- (iii) Moreover $\tau^v(s)$ is the vertical component of the tension field $\tau(s)$. So if s is an harmonic map, then it is certainly a harmonic section.

Theorem 4.7 [51] An harmonic section s is a harmonic map if and only if $\langle s^* \phi, s^* \Phi \rangle = 0$ where

$$\langle s^* \phi, s^* \Phi \rangle(X) = \sum_i \langle s^* \phi \otimes s^* \Phi \rangle(E_i, E_i, X)$$

for any orthonormal tangent frame (E_i) of M .

In particular, if s is flat ($s^* \Phi = 0$) then s is a harmonic map if and only if s is a harmonic section.

Remark 4.5 Let us consider the situation described by examples 4.1 and 4.9. Then if $f^* \Phi = 0$, $f: L \rightarrow N$ is vertically harmonic if and only if $\tilde{f}: L \rightarrow u^* N$ is an harmonic section if and only if $\tilde{f}: L \rightarrow u^* N$ is an harmonic map. But it does not imply that $f: L \rightarrow N$ is harmonic! (See the Appendix.) Indeed in the previous theorem it is essential that s be a section: $\pi \circ s = \text{Id}$.

In fact the previous theorems can be easily generalized for non section map. The proofs in [51] holds without any change for theorems 4.4, 4.5-(i), 4.6-(i, iii), while for theorems 4.5-(ii), 4.6-(ii), 4.7: follow the proof of [51], just replace the starting equation $\pi \circ s = \text{Id}$ by $\pi \circ s = u$. Then we obtain

Theorem 4.8 For all $s \in C^\infty(M, N)$, we have

(i) $I(\tau^v(s)) = -d^*(s^* \phi) + \frac{1}{2} \text{Tr}(s^* \phi^* U)$

(ii) $\phi(\nabla ds) = \nabla^c(s^* \phi) + \frac{1}{2} s^* \phi^* B - \frac{1}{2} s^* \Phi.$

In particular if s is vertically geodesic then s is a harmonic section.

(iii) $I(\Pi^v s) = \phi \circ \nabla ds + \frac{1}{2} s^* \Phi.$

The map s is superflat if and only if s is flat and vertically geodesic. Moreover $\tau^v(s)$ is the vertical component of the tension field $\tau(s)$. So if s is an harmonic map, then it is certainly vertically harmonic.

(iv) $2g(\pi_* \nabla ds(X, Y), u_* Z) = \langle s^* \phi \otimes s^* \Phi \rangle(X, Y, Z) + \langle s^* \phi \otimes s^* \Phi \rangle(Y, X, Z) + 2g(\nabla du(X, Y), u_* Z).$

Let us suppose now that u is an immersion, then this equation determines the horizontal part of ∇ds . In particular, if s is flat then s is horizontally geodesic if and only if u is totally geodesic; and s is totally geodesic if and only if s is superflat and u is totally geodesic.

(v) A vertically harmonic map s is a harmonic map if and only if

$$g(\tau(u), \cdot) + \langle s^* \phi, s^* \Phi \rangle = 0.$$

In particular if s is flat, then s is a harmonic map if and only if s is vertically harmonic and $u = \pi \circ s$ is harmonic.

We could also deduce this generalisation from the previous theorems 4.4-4.7 themselves. Indeed we can apply these to the section $\tilde{s} \in \mathcal{C}(u^*N)$ corresponding to s and use theorems 8.1 and 8.2 in the Appendix, but we must suppose in addition that u is an isometry.

Let us go further in the generalisation and consider maps $f \in \mathcal{C}^\infty(L, N)$ with (L, b) a Riemannian manifold (see examples 4.1 and 4.9). Then the proofs in [51] holds for theorems 4.4, 4.5-(i), 4.6-(i, iii), whereas theorems 4.5-(ii), 4.6-(ii), 4.7 are no longer valid. Indeed the equation in theorem 4.8-(iii) holds, but it gives us only $[\pi_* \nabla df]_{u_* TL}$, the component of $[\pi_* \nabla df]$ in the tangent bundle $u_* TL$. So if we want $[\pi_* \nabla df]_{(u_* TL)^\perp}$ we must introduce the 3-linear form $\langle f^* \phi \otimes \Phi(f_* \cdot, \cdot) \rangle \in \mathcal{C}(T^*L \otimes T^*L \otimes f^* \mathcal{H})$ defined by :

$$\langle f^* \phi \otimes \Phi(f_* \cdot, \cdot) \rangle(a, b, Z) = \langle f^* \phi(a), \Phi(f_* a, Z) \rangle.$$

Then we have

Theorem 4.9 *For all $f \in \mathcal{C}^\infty(L, N)$, we have*

(i) $g(\pi_* \nabla df(a, b), \pi^* Z) = \langle f^* \phi \otimes \Phi(f_* \cdot, \cdot) \rangle(a, b, Z) + g(\nabla du(a, b), \pi_* Z).$

In particular, if f is strongly flat i.e. $(\Phi(f_ \cdot, \cdot) = 0)$ then:*

- *f is horizontally geodesic if and only if u is totally geodesic.*

- *f is totally geodesic if and only if f is superflat and u is totally geodesic.*

(ii) *A vertically harmonic map f is a harmonic map if and only if*

$$g(\tau(u), \cdot) + \langle f^* \phi, \Phi(f_* \cdot, \cdot) \rangle = 0$$

where $\langle f^ \phi, \Phi(f_* \cdot, \cdot) \rangle(X) = \sum_i \langle f^* \phi \otimes \Phi(f_* \cdot, \cdot) \rangle(e_i, e_i, X)$ for any tangent frame (e_i) of L . In particular if f is strongly flat $(\Phi(f_* \cdot, \cdot) = 0)$ then f is a harmonic map if and only if f is vertically harmonic and u is harmonic.*

4.2.3 Reductions of homogeneous fibre bundles

Let us suppose now that the structure group H of $\pi_M: Q \rightarrow M$ is reducible to a (closed) subgroup $H^\vee \supset K$. That is to say, there exists a principal H^\vee -subbundle $\pi_M^\vee: Q^\vee \rightarrow M$. Let us suppose in addition to that, that H^\vee/K is reductive: $\mathfrak{h}^\vee = \mathfrak{k} \oplus \mathfrak{p}^\vee$ and $\text{Ad}k(\mathfrak{p}^\vee) = \mathfrak{p}^\vee, \forall k \in K$. The restriction to \mathfrak{p}^\vee of the $\text{Ad}K$ -invariant inner product on \mathfrak{p} defines a H^\vee -invariant metric on H^\vee/K which is nothing but the metric induced by the H -invariant metric on H/K , so that the inclusion $H^\vee/K \rightarrow H/K$ is an isometric embedding. Let $\mathfrak{p}' = (\mathfrak{p}^\vee)^\perp$ in \mathfrak{p} and let us suppose that \mathfrak{p}' is $\text{Ad}H^\vee$ -invariant, so that $\mathfrak{h} = \mathfrak{h}^\vee \oplus \mathfrak{p}'$ is a reductive decomposition and H/H^\vee is reductive. Conversely if H/H^\vee is reductive: $\mathfrak{h} = \mathfrak{h}^\vee \oplus \mathfrak{p}'$ with \mathfrak{p}' $\text{Ad}H^\vee$ -invariant, then we can always complete any $\text{Ad}H^\vee$ -inner product in \mathfrak{p}' by an $\text{Ad}K$ -invariant inner product $\mathfrak{p} = \mathfrak{p}^\vee \oplus \mathfrak{p}'$:

$$\langle \cdot, \cdot \rangle_{\mathfrak{p}} = \langle \cdot, \cdot \rangle_{\mathfrak{p}^\vee} + \langle \cdot, \cdot \rangle_{\mathfrak{p}'}$$

for which $\mathfrak{p}' = (\mathfrak{p}^\vee)^\perp$ in \mathfrak{p} .

In the following we suppose that H/H^\vee is reductive and that the inner product in \mathfrak{p} is chosen as described above.

Now let us turn toward the connection 1-form ω . Its restriction $\omega^\vee := \omega|_{TQ^\vee}$ defines a connection on $\pi_M^\vee: Q^\vee \rightarrow M$. We endow Q^\vee with ω^\vee and $(Q^\vee, H^\vee, K, \omega^\vee)$ is then a homogeneous fibre bundle as defined in the beginning of 4.2.

Definition 4.8 *We will then say that $\pi: N \rightarrow M$ is a homogeneous fibre bundle of $\pi: N \rightarrow M$.*

Moreover ω is reducible (to ω^\vee) in Q^\vee if and only if one of the following equivalent statements holds ([40])

- $\forall q \in Q^\vee, (\mathcal{H}_0)_q$ is tangent to Q^\vee .
- $\omega|_{TQ^\vee} = \omega^\vee$ (i.e. $\omega|_{TQ^\vee}$ is \mathfrak{h}^\vee -valued).
- The canonical cross section s^\vee of the associated bundle $E^\vee := Q/H^\vee = Q \times_H (H/H^\vee)$, which defines the H^\vee -reduction Q^\vee is horizontal.

Definition 4.9 Denoting by $i_\vee: N^\vee \rightarrow N$ the natural inclusion, we will say that $i_\vee: N^\vee \rightarrow N$ is a reduction of homogeneous fibre bundles if ω is reducible in Q^\vee (being implicit that the homogeneous fibre structure are respectively (Q, H, K, ω) and $(Q^\vee, H^\vee, K^\vee, \omega^\vee)$).

The vertical bundle (in TQ), \mathcal{V}_0 , splits as follows

$$\mathcal{V}_0 = \mathcal{V}'_0 \oplus \mathcal{V}^\vee_0$$

where $(\mathcal{V}'_0)_q = q \cdot \mathfrak{h}^\vee = T_q(q \cdot H^\vee)$ and $(\mathcal{V}^\vee_0)_q = q \cdot \mathfrak{p}'$, and quotienting by \mathfrak{k} , i.e. by applying $d\pi_N$ we obtain the following decomposition of \mathcal{V} :

$$\mathcal{V} = \mathcal{V}' \oplus \mathcal{V}^\vee$$

with $\mathcal{V}' = d\pi_N(\mathcal{V}'_0)$ and $\mathcal{V}^\vee = d\pi_N(\mathcal{V}^\vee_0)$.

Then the canonical isomorphism $I: \mathcal{V} \rightarrow \mathfrak{p}_Q$ sends the previous decomposition onto the following $\mathfrak{p}_Q = \mathfrak{p}'_Q \oplus \mathfrak{p}^\vee_Q$ (i.e. $\mathcal{V}', \mathcal{V}^\vee$ are sent resp. onto \mathfrak{p}'_Q and \mathfrak{p}^\vee_Q).

Then the vertical space in TN^\vee is $\mathcal{V}|_{N^\vee}$ that we will also denote by \mathcal{V}^\vee when there is no possibilities of confusion. The splitting of TN^\vee defined by ω^\vee is then

$$TN^\vee = \mathcal{V}|_{N^\vee} \oplus \mathcal{H}|_{N^\vee}$$

where $\mathcal{H}^\vee = d\pi_N(\mathcal{H}_0^\vee)$ and $\mathcal{H}_0^\vee = \ker \omega_{\mathfrak{h}^\vee}$. Let us remark that ω is reducible if and only if one of the following equivalent statements holds

- $\mathcal{H}|_{N^\vee} = \mathcal{H}|_{N^\vee}$,
- the inclusion $i_\vee: (N^\vee, h^\vee) \rightarrow (N, h)$ is an isometry, where h^\vee is the metric defined by ω^\vee and $\langle \cdot, \cdot \rangle_{\mathfrak{p}^\vee}$, according to (62).

The canonical bundle on N^\vee , $\mathfrak{p}^\vee_{Q^\vee} = Q^\vee \times_{K^\vee} \mathfrak{p}^\vee \rightarrow N^\vee$ is the restriction to N^\vee of $\mathfrak{p}^\vee_Q \rightarrow N$, and the canonical isomorphism $I^\vee: \mathcal{V}|_{N^\vee} \rightarrow \mathfrak{p}^\vee_{Q^\vee}$ is the restriction to $\mathcal{V}|_{N^\vee}$ of $I: \mathcal{V} \rightarrow \mathfrak{p}_Q$.

Since $\omega_{\mathfrak{p}} = \omega_{\mathfrak{p}^\vee} + \omega_{\mathfrak{p}'}$, the homogeneous connection form on N^\vee , ϕ^\vee (the $\mathfrak{p}^\vee_{Q^\vee}$ -valued 1-form on N^\vee defined by $\omega_{\mathfrak{p}^\vee}$) is the restriction to N^\vee of the \mathfrak{p}^\vee_Q -component of ϕ :

$$\phi^\vee = (i_\vee)^* [\phi]_{\mathfrak{p}^\vee_Q} = [(i_\vee)^* \phi]_{\mathfrak{p}^\vee_{Q^\vee}},$$

where $i_\vee: N^\vee \rightarrow N$ is the natural inclusion. The homogeneous curvature form Φ^\vee (defined by $\Omega_{\mathfrak{p}^\vee}$, with $\Omega^\vee = d\omega^\vee + \frac{1}{2}[\omega^\vee \wedge \omega^\vee]$) is given by

$$\Phi^\vee = (i_\vee)^* [\Phi]_{\mathfrak{p}^\vee_Q} = [(i_\vee)^* \Phi]_{\mathfrak{p}^\vee_{Q^\vee}}.$$

Furthermore, $\mathfrak{p}^\vee_{Q^\vee}$ is ∇^c -parallel: the covariant derivative on $\mathfrak{p}^\vee_{Q^\vee}$ defined by $\omega_{\mathfrak{k}|_{TQ^\vee}}$ is the restriction of ∇^c to $\mathfrak{p}^\vee_{Q^\vee}$. In other words ∇^c commutes with the projection on \mathfrak{p}^\vee_Q . The canonical torsion on N^\vee is given by

$$T^{c,\vee} = d^c \phi^\vee = [(i_\vee)^* T^c]_{\mathfrak{p}^\vee_Q}.$$

In particular, the results of what precedes is:

Proposition 4.4 *Let us suppose that ω is reducible in Q^\vee . Then we have*

- (i) $\phi^\vee = (i_\vee)^*\phi$,
- (ii) $\Phi^\vee = (i_\vee)^*\Phi$,
- (iii) $T^{c,\vee} = (i_\vee)^*T^c$.

Now, let us remark that

Lemma 4.1 *We have the following identity*

$$B^\vee = (i_\vee)^*B$$

Proof. Let $a_{\mathfrak{p}^\vee}, b_{\mathfrak{p}^\vee} \in \mathfrak{p}^\vee$ and $c_{\mathfrak{p}} \in \mathfrak{p}$. Then we have

$$[a_{\mathfrak{p}^\vee}, b_{\mathfrak{p}^\vee}]_{\mathfrak{p}} = [a_{\mathfrak{p}^\vee}, b_{\mathfrak{p}^\vee}]_{\mathfrak{p}^\vee}$$

since $\mathfrak{h}^\vee = \mathfrak{k} \oplus \mathfrak{p}^\vee$ is a Lie subalgebra of $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{p}$. Moreover, we have also

$$\begin{aligned} \langle U(a_{\mathfrak{p}^\vee}, b_{\mathfrak{p}^\vee}), c_{\mathfrak{p}} \rangle &= \langle [c_{\mathfrak{p}}, a_{\mathfrak{p}^\vee}]_{\mathfrak{p}}, b_{\mathfrak{p}^\vee} \rangle + \langle [c_{\mathfrak{p}}, a_{\mathfrak{p}^\vee}]_{\mathfrak{p}}, b_{\mathfrak{p}^\vee} \rangle \\ &= \langle [c_{\mathfrak{p}^\vee}, a_{\mathfrak{p}^\vee}]_{\mathfrak{p}^\vee}, b_{\mathfrak{p}^\vee} \rangle + \langle [c_{\mathfrak{p}^\vee}, a_{\mathfrak{p}^\vee}]_{\mathfrak{p}^\vee}, b_{\mathfrak{p}^\vee} \rangle = \langle U^\vee(a_{\mathfrak{p}^\vee}, b_{\mathfrak{p}^\vee}), c_{\mathfrak{p}^\vee} \rangle \end{aligned}$$

since $[\mathfrak{p}', \mathfrak{p}^\vee] \subset \mathfrak{p}'$ because the decomposition $\mathfrak{h} = \mathfrak{h}^\vee \oplus \mathfrak{p}'$ is $\text{Ad}H^\vee$ -invariant. We have denoted by U^\vee the map associated with the bundle $\pi: N^\vee \rightarrow M$ according to equation (67). This completes the proof. \square

Therefore we deduce from what precedes and theorem 4.2, the following.

Proposition 4.5 *Let us suppose that ω is reducible in Q^\vee . Let (L, b) be a Riemannian manifold, and $f: L \rightarrow N^\vee$ be a map. Then we have*

- (i) $\tau^\vee(f) = \tau^v(f)$,
- (ii) $\Pi^\vee(f) = \Pi^v(f)$,

where $\tau^\vee(f)$ is the vertical tension field of f in N^\vee and $\Pi^\vee(f)$ its vertical second fundamental form in N^\vee .

4.3 Examples of Homogeneous fibre bundles

In this section, we give examples and applications for the theory developed in the previous sections whose we use here the same notations.

4.3.1 Homogeneous spaces fibration

Let us take $Q = G$ a Lie group, and $K \subset H \subset G$ subgroups of G , (H, K) satisfying the hypothesis in the begining of section 4.2.1. Let us suppose that $M = G/H$ is reductive and Riemannian: that is to say if $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ is the reductive decomposition, then $\text{Ad}_{\mathfrak{m}}H$ is compact and we choose an $\text{Ad}H$ -invariant inner product $\langle \cdot, \cdot \rangle_{\mathfrak{m}}$ in \mathfrak{m} . For ω , we take the canonical connection on $\pi_M: G \rightarrow G/H$ which is given, let us recall it, by $\omega = \theta_{\mathfrak{h}}$ where θ is the Maurer-Cartan form in G (see section 1.3). Then the corresponding decomposition $TQ = \mathcal{V}_0 \oplus \mathcal{H}_0$ is given by

$$T_g G = g \cdot \mathfrak{g} = \underbrace{g \cdot \mathfrak{h}}_{\mathcal{V}_0} \oplus \underbrace{g \cdot \mathfrak{m}}_{\mathcal{H}_0}.$$

Since $\mathfrak{n} := \mathfrak{p} \oplus \mathfrak{m}$ is $\text{Ad}K$ -invariant, then $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{n}$ is a reductive decomposition and $N = G/K$ is reductive. Let us recall that we have the canonical identification $G \times_K \mathfrak{g} \cong N \times \mathfrak{g}$ given by (7), which gives us an identification $\mathfrak{n}_G = G \times_K \mathfrak{n} \cong [\mathfrak{n}]$. Then under this last identification and under the one given by the Maurer-Cartan form of G/K , $\beta: TN \xrightarrow{\cong} [\mathfrak{n}]$ (see section 1.2), the splitting $TN = \mathcal{V} \oplus \mathcal{H}$ is

$$TN = [\mathfrak{p}] \oplus [\mathfrak{m}],$$

the canonical isomorphism $I: \mathcal{V} \rightarrow \mathfrak{p}_G$ is then the identity, and $\phi: TN \rightarrow \mathfrak{p}_G$ the projection on $[\mathfrak{p}]$ along $[\mathfrak{m}]$. The metric h on G/K is then defined by the $\text{Ad}K$ -invariant inner product:

$$\langle \cdot, \cdot \rangle_{\mathfrak{n}} = \langle \cdot, \cdot \rangle_{\mathfrak{p}} + \langle \cdot, \cdot \rangle_{\mathfrak{m}}.$$

Furthermore, $\Omega = d\omega + \frac{1}{2}[\omega \wedge \omega] = d\theta_{\mathfrak{h}} + \frac{1}{2}[\theta_{\mathfrak{h}} \wedge \theta_{\mathfrak{h}}]$ and thus

$$\Omega_{\mathfrak{p}} = d\theta_{\mathfrak{p}} + [\theta_{\mathfrak{k}} \wedge \theta_{\mathfrak{p}}] + \frac{1}{2}[\theta_{\mathfrak{p}} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}}.$$

Since $d\theta + \frac{1}{2}[\theta \wedge \theta] = 0$, then (projecting on \mathfrak{h}) we have

$$d\theta_{\mathfrak{h}} + \frac{1}{2}[\theta_{\mathfrak{h}} \wedge \theta_{\mathfrak{h}}] + \frac{1}{2}[\theta_{\mathfrak{m}} \wedge \theta_{\mathfrak{m}}]_{\mathfrak{h}} = 0$$

thus

$$\Omega = -\frac{1}{2}[\theta_{\mathfrak{m}} \wedge \theta_{\mathfrak{m}}]_{\mathfrak{h}}$$

so that

$$\Omega_{\mathfrak{p}} = -\frac{1}{2}[\theta_{\mathfrak{m}} \wedge \theta_{\mathfrak{m}}]_{\mathfrak{p}} \tag{69}$$

therefore

$$\Phi = -\frac{1}{2}[\psi \wedge \psi]_{\mathfrak{p}} \tag{70}$$

where $\psi: TN \rightarrow \mathcal{H} = [\mathfrak{m}]$ is the projection on \mathcal{H} along $\mathcal{V} = [\mathfrak{p}]$.

The covariant derivative ∇^c , which lifts into $d + \theta_{\mathfrak{k}}$ in G , is nothing but the canonical linear connection ∇^0 in $N = G/K$ restricted to $[\mathfrak{p}] \subset TN$ (see section 1.4 and 1.5).

The canonical torsion T^c , which lifts in G into

$$d\theta_{\mathfrak{p}} + [\theta_{\mathfrak{k}} \wedge \theta_{\mathfrak{p}}] = -\frac{1}{2}[\theta_{\mathfrak{m}} \wedge \theta_{\mathfrak{m}}]_{\mathfrak{p}} - \frac{1}{2}[\theta_{\mathfrak{p}} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}} \tag{71}$$

is given by ³²

$$\begin{aligned} T^c &= -\frac{1}{2}[\psi \wedge \psi]_{\mathfrak{p}} - \frac{1}{2}[\phi \wedge \phi]_{\mathfrak{p}} \\ &= [\cdot, \cdot]_{\mathfrak{n}} + [\phi \wedge \psi]_{\mathfrak{p}}. \end{aligned}$$

The associated bundle

$$\mathfrak{h}_G = G \times_H \mathfrak{h} \cong [\mathfrak{h}]^M := \{(g.p_0, \text{Ad}g(a)), g \in G, a \in \mathfrak{h}\} \subset M \times \mathfrak{g}$$

can be embedded into $\mathfrak{so}(TM)$ by³³

$$\xi = \text{Ad}g(a) \in \text{Ad}g(\mathfrak{h}) = [\mathfrak{h}]_{g.p_0}^M \mapsto \text{ad}\xi|_{\text{Ad}g(\mathfrak{m})} = \text{Ad}g \circ \text{ad}_{\mathfrak{m}}a \circ \text{Ad}g^{-1} \in \mathfrak{so}(\text{Ad}g(\mathfrak{m})). \tag{72}$$

³²In particular, according to (70), we recover, for this example, the Homogeneous structure equation (64).

³³Or in other words $[g, a] \in \mathfrak{h}_G = G \times_H \mathfrak{h} \mapsto [g, \text{ad}_{\mathfrak{m}}a] \in G \times_H \mathfrak{so}(\mathfrak{m}) \cong \mathfrak{so}(TM)$.

In the same way, $\mathfrak{k}_G = G \times_K \mathfrak{k} \cong [\mathfrak{k}]^N$ embeds in $\mathfrak{so}(N)$. Moreover let us remark that we have

$$\pi^* \mathfrak{h}_G = G \times_K \mathfrak{h} = [\mathfrak{h}]^N = \{(g.n_0, \text{Ad}g(a)), g \in G, a \in \mathfrak{h}\},$$

and that $\pi^* \mathfrak{h}_G$ embeds into $\mathfrak{so}(\pi^* TM)$.

As concerns the covariant derivative ∇^ω , defined in \mathfrak{h}_G , it lifts into $d + \theta_{\mathfrak{h}}$ in G and under the embedding (72), it is nothing but the restriction to the subbundle $\mathfrak{so}(TM)$ of the endomorphism connection on M (i.e. the tensor product connection in $T^*M \otimes T^*M$) defined by the canonical linear connection in M , ∇^0 . Indeed under the embedding (72), ∇^ω lifts to the derivative $d + \text{ad}_{\mathfrak{m}} \theta_{\mathfrak{h}}$ and equation (18) allows to conclude.

Therefore $\nabla^{\mathfrak{p}}$ is given by its lift ³⁴

$$[(d + \theta_{\mathfrak{h}})_{|\mathfrak{p}}]_{\mathfrak{p}} = d + \text{ad}_{\mathfrak{p}} \theta_{\mathfrak{k}} + [\theta_{\mathfrak{p}}, \cdot]_{\mathfrak{p}} \quad (73)$$

that is to say $\nabla^{\mathfrak{p}}$ is the $[\mathfrak{p}]$ -component of the linear connection ∇^1 in G/K (see section 1.6) restricted to $[\mathfrak{p}] \subset TN$. (Indeed we have $[\theta_{\mathfrak{n}}, \cdot]_{|\mathfrak{p}}]_{\mathfrak{p}} = [\theta_{\mathfrak{p}}, \cdot]_{|\mathfrak{p}}]_{\mathfrak{p}} + [\theta_{\mathfrak{m}}, \cdot]_{|\mathfrak{p}}]_{\mathfrak{p}}$, but $[\mathfrak{m}, \mathfrak{p}] \subset [\mathfrak{m}, \mathfrak{h}] \subset \mathfrak{m}$ by reductivity.)

Let us see to which corresponds ∇^c under the embedding (72). The canonical connection ∇^0 gives rise by restriction to a linear connection on the horizontal subbundle $\mathcal{H} = \pi^* TM$. Then ∇^c is nothing but the restriction to $\mathfrak{p}_G \subset \mathfrak{so}(\pi^* TM)$ of the tensor product connection in $\text{End}(\pi^* TM)$ defined by ∇^0 .

Moreover the Levi-Civita connection in N is given by (see section 1.6)

$$\nabla^N = \nabla^{\frac{\text{met}}{2}} = \nabla^0 + \frac{1}{2} B^N$$

where $B^N = [,]_{|\mathfrak{n}} + U^N$ and U^N is defined by equation (15). Then we have by taking the projection on the vertical subbundle $[\mathfrak{p}]$:

$$\phi(\nabla_A^N V) = \nabla_A^0(\phi V) + \frac{1}{2} \phi \circ B^N(A, V)$$

so that we can conclude according to theorem 4.2 that

$$\phi \circ B^N = \phi^* B - \Phi, \quad (74)$$

which can also be verified directly using the expressions of B^N , B and Φ .

If H/K is (locally) symmetric. In this case, we have $T^c = \Phi$ (see (64), or (71) and (69)). Moreover, according to (73), $\nabla^{\mathfrak{p}}$ lifts to $d + \theta_{\mathfrak{k}}$, so that we recover that $\nabla^{\mathfrak{p}} = \nabla^c$ in this case. Now, let us apply the equality $\nabla^v = \nabla^c$ in \mathcal{V} (theorem 4.2)³⁵.

Let $f: (L, b) \rightarrow N$ be a map then we have

$$\tau^v(f) = \text{Tr}_b(\nabla^v d^v f) = *d^{\nabla^v} * d^v u = * \text{Ad}F (d(*\alpha_{\mathfrak{p}}) + [\alpha_{\mathfrak{k}} \wedge (*\alpha_{\mathfrak{p}})]) . f$$

where F lifts f in G and $\alpha = F^{-1} dF$. Then f is vertically harmonic if and only if

$$d(*\alpha_{\mathfrak{p}}) + [\alpha_{\mathfrak{k}} \wedge (*\alpha_{\mathfrak{p}})] = 0$$

Moreover f is flat ($f^* \Phi = 0$) if and only if it is vertically torsion free ($f^* T^c = 0$) if and only if

$$[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{p}} = 0 \iff d\alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge \alpha_{\mathfrak{p}}] = 0.$$

³⁴ $[\cdot]_{\mathfrak{p}}$ denotes as usual the \mathfrak{p} -component.

³⁵ We can also use directly theorem 4.4.

G/K is a (locally) $2k$ -symmetric space Let us suppose that there exists an order $2k$ automorphism $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ such that $K = G_0$ with G_0 such that $(G^\tau)^0 \subset G_0 \subset G^\tau$, and $(G^\sigma)^0 \subset H \subset G^\sigma$ with $\sigma = \tau^2$ (see section 2.1). Then H/K is (locally) symmetric (see section 2.1). We have the following identities (with the notation of section 2.1)

$$\mathfrak{m} = \bigoplus_{j=1}^{k-1} \mathfrak{m}_j \quad \text{and} \quad \mathfrak{k} = \mathfrak{g}_0, \mathfrak{p} = \mathfrak{g}_k.$$

Then we have

$$\Omega_{\mathfrak{p}} = -\frac{1}{2}[\theta_{\mathfrak{m}} \wedge \theta_{\mathfrak{m}}]_{\mathfrak{p}} = -\frac{1}{2} \sum_{\substack{i+j=k \\ i,j \in \mathbb{Z}_{2k} \setminus \{0,k\}}} [\theta_j \wedge \theta_i],$$

so that in particular

Proposition 4.6 Let (L, j) be a Riemann surface. If $f: (L, j) \rightarrow N$ satisfies the equations $\alpha''_{-j} = 0$, $1 \leq j \leq k-1$ then we have $f^*\Phi = 0$. In other words if $f: (L, j) \rightarrow N$ is horizontally holomorphic then it is flat, that is to say f is vertically torsion free or equivalently

$$d\alpha_k + [\alpha_0 \wedge \alpha_k] = 0. \quad (75)$$

Theorem 4.10 In the even minimal determined elliptic integrable system $(\text{Syst}(k, \tau))$, the last equation (S_k) is equivalent to

$$\begin{cases} (\text{Re}(S_k)) & \equiv & d\alpha_k + [\alpha_0 \wedge \alpha_k] = 0 & \iff & f \text{ is vertically torsion free i.e. } f \text{ is flat,} \\ (\text{Im}(S_k)) & \equiv & d(*\alpha_k) + [\alpha_0 \wedge (*\alpha_k)] = 0 & \iff & f \text{ is vertically harmonic.} \end{cases}$$

In conclusion the even minimal determined elliptic system $(\text{Syst}(k, \tau))$ means that the geometric map f is horizontally holomorphic (which implies that f is flat) and vertically harmonic.

Remark 4.6 The vertical torsion free equation (75) is the projection on \mathfrak{p} of the Maurer-Cartan equation provided that we assume the horizontal holomorphicity $\alpha''_{-j} = 0, 1 \leq j \leq k-1$. In the same way, the equations (S_j) , $0 \leq j \leq k-1$, of the elliptic system $(\text{Syst}(k, \tau))$, are the projections on the different spaces \mathfrak{g}_{-j} , of the Maurer-Cartan equation, provided that we assume the horizontal holomorphicity.

Use of the canonical $2k$ -structure \mathfrak{J}_{J_0} . Furthermore the morphism of bundle (over M) $\mathfrak{J}_{J_0}: N \rightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2) \subset \mathcal{Z}_{2k}^{\alpha_0}(M)$ defines a $2k$ -structure on π^*TM (still denoted by \mathfrak{J}_{J_0}), which according to (23) allows to precise the subbundles \mathfrak{k}_G and \mathfrak{p}_G (under the embedding $\pi^*\mathfrak{h}_G \hookrightarrow \mathfrak{so}(\pi^*TM)$)

$$\mathfrak{k}_G = \{A \in \pi^*\mathfrak{h}_G \mid [A, \mathfrak{J}_{J_0}] = 0\} := \mathfrak{so}_{(+1)}(\pi^*TM, \mathfrak{J}_{J_0}) \cap \pi^*\mathfrak{h}_G \quad (76)$$

$$\mathfrak{p}_G = \{A \in \pi^*\mathfrak{h}_G \mid A\mathfrak{J}_{J_0} + \mathfrak{J}_{J_0}A\} := \mathfrak{so}_{(-1)}(\pi^*TM, \mathfrak{J}_{J_0}) \cap \pi^*\mathfrak{h}_G \quad (77)$$

Remark 4.7 The embedding $\mathfrak{h}_G \hookrightarrow \mathfrak{so}(TM)$ is the H -equivariant extension of the map $a \in \mathfrak{h} \mapsto \text{ad}_{\mathfrak{m}}a \in \mathfrak{so}(\mathfrak{m}) \cong \mathfrak{so}(T_{p_0}M)$, and in the same way \mathfrak{J}_{J_0} is the H -equivariant extension of the map $h.G_0 \in H/G_0 \mapsto hJ_0h^{-1} \in \mathcal{Z}(T_{p_0}M, J_0)$, so that the equations (76) are obtained by H -equivariance from (23).

Let us now express the homogeneous fibre bundle tools ϕ , Φ , and ∇^c in terms of the embedding \mathfrak{J}_{J_0} . To do not weigh the notation we will forget the index J_0 in \mathfrak{J}_{J_0} , in the following theorem.

Theorem 4.11 *If $A, B \in TN$, $F \in \mathcal{C}(\mathfrak{p}_G)$ then*

$$(i) \quad \phi A = -\frac{1}{2}\mathcal{J}^{-1}\overset{M}{\nabla}_A^0\mathcal{J}$$

$$(ii) \quad \Phi(A, B) = \frac{1}{2}\mathcal{J}^{-1}[\mathcal{J}, \pi^*R^{\overset{M}{\nabla}^0}(A, B)] \text{ where } R^{\overset{M}{\nabla}^0} \text{ is the curvature of } \overset{M}{\nabla}^0.$$

$$(iii) \quad \nabla_A^c F = \overset{N}{\nabla}_A^0 F = \frac{1}{2}\mathcal{J}^{-1}[\mathcal{J}, \overset{M}{\nabla}_A^0 F].$$

Proof. Lifted in G (i) become

$$\text{ad}_{\mathfrak{m}}\theta_k = -\frac{1}{2}J_0^{-1}(dJ_0 + [\text{ad}_{\mathfrak{m}}\theta_{\mathfrak{h}}, J_0]),$$

keeping in mind that under the embedding $\pi^*\mathfrak{h}_G \hookrightarrow \mathfrak{so}(\pi^*TM)$, θ_k corresponds to $\text{ad}_{\mathfrak{m}}\theta_k$. Moreover this equality holds since $[\text{ad}_{\mathfrak{m}}\theta_{\mathfrak{h}}, J_0] = -2J_0\text{ad}_{\mathfrak{m}}\theta_k$. This proves (i).

For the following, let us keep in mind that \mathfrak{h}_G is embedded into $\mathfrak{so}(\pi^*TM)$.

Then lifted in G , (ii) becomes

$$-\text{ad}_{\mathfrak{m}}([\theta_{\mathfrak{m}}, \theta_{\mathfrak{m}}]_{\mathfrak{p}}) = \frac{1}{2}J_0^{-1}[J_0, -\text{ad}_{\mathfrak{m}}([\theta_{\mathfrak{m}}, \theta_{\mathfrak{m}}]_{\mathfrak{h}})]$$

which holds for the same reason as above.

Finally (iii) lifted in G becomes

$$d + \theta_0 = -\frac{1}{2}J_0^{-1}\left[J_0, d\tilde{F} + [\text{ad}_{\mathfrak{m}}\theta_{\mathfrak{h}}, \tilde{F}]\right]$$

which holds since the right hand side is equal to $d\tilde{F} + [\text{ad}_{\mathfrak{m}}\theta_0, \tilde{F}]$ which lifts $\overset{N}{\nabla}^0 F$, because $[\text{ad}_{\mathfrak{m}}\theta_k, \tilde{F}]$ commutes with J_0 . We have denoted by $\tilde{F}: G \rightarrow \mathfrak{g}_k$ the lift of F . This completes the proof. \square

Theorem 4.12 *Let $s \in \mathcal{C}(\pi)$ and $J = s^*\mathcal{J}_{J_0} \in \mathcal{C}(\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2))$ be the corresponding $2k$ -structure. Then*

$$(i) \quad I(d^v s) = -\frac{1}{2}J^{-1}\overset{M}{\nabla}^0 J. \text{ Thus } s \text{ is horizontal if and only if } J \text{ is } \overset{M}{\nabla}^0\text{-parallel.}$$

$$(ii) \quad I(\Pi^v(s)) = -\frac{1}{4}[J^{-1}, (\overset{M}{\nabla}^0)^2 J].$$

Thus s is superflat if and only if $(\overset{M}{\nabla}^0)^2 J$ commutes with J .

$$(iii) \quad I(\tau^v(s)) = \frac{1}{4}[J^{-1}, (\overset{M}{\nabla}^0)^*\overset{M}{\nabla}^0 J].$$

Thus s is a harmonic section if and only if $(\overset{M}{\nabla}^0)^\overset{M}{\nabla}^0 J$ commutes with J .*

$$(iv) \quad s^*\Phi = \frac{1}{2}J^{-1}[J, R^{\overset{M}{\nabla}^0}].$$

These properties hold also for maps $f \in \mathcal{C}^\infty(L, N)$, (L, b) being a Riemannian manifold: (i), (ii), (iii)

without any change and (iv) becoming $f^\Phi = \frac{1}{2}J^{-1}[J, u^*R^{\overset{M}{\nabla}^0}]$, with $u = \pi \circ f$.*

Proof. (i) and (iv) follows from theorem 4.12 (i) and (iii) respectively. The assertions (iii) is a consequence of the assertion (ii). Finally (iv) follows from theorem 4.12-(iii) and the fact that

$$\left[\begin{matrix} M \\ \nabla^0 J^{-1}, \nabla^0 J \end{matrix} \right] = - \left[\begin{matrix} M \\ J^{-1} \left(\begin{matrix} M \\ \nabla^0 J \end{matrix} \right) J^{-1}, \nabla^0 J \end{matrix} \right]$$

commutes with J . This completes the proof. \square

Now, we can conclude.

Corollary 4.1 *Let (L, j) be a Riemann surface, $f: L \rightarrow N$ a map and $J = f^* \mathcal{J}_{J_0}$ the corresponding map into $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$. Then f is a geometric solution of the even minimal determined system $(\text{Syst}(k, \tau))$ if and only if*

(i) J is an admissible twistor lift ($\Leftrightarrow f$ is horizontally holomorphic).

(ii) J is vertically harmonic³⁶: $\left[\begin{matrix} M \\ (\nabla^0)^* \nabla^0 J, J \end{matrix} \right] = 0$ ($\Leftrightarrow f$ is vertically harmonic).

Moreover the first condition implies that $\left[\begin{matrix} M \\ u^* R^{\nabla^0}, J \end{matrix} \right] = 0$ i.e. that J is a flat section in

$(\text{End}(u^* TM), u^* \nabla^0)$ ($\Leftrightarrow f$ is flat).

Furthermore f is a primitive geometric solution (i.e. there exists $m \leq k$ such that f is m -primitive, which is equivalent to say that f is k -primitive) if and only if

(i) J is an admissible twistor lift

(ii) J is parallel: $\nabla^0 J = 0$ ($\Leftrightarrow f$ is horizontal).

4.3.2 The twistor bundle of almost complex structures $\Sigma(E)$

We give ourself the same ingredients as in example 4.2. Let us suppose that the vector bundle E is oriented. Then the bundle of positive (resp. negative) orthogonal almost complex structure on E (i.e. the component $\Sigma^\varepsilon(E)$ of $\Sigma(E)$ with $\varepsilon = \pm 1$), $\pi_\Sigma: \Sigma^\varepsilon(E) \rightarrow M$ is a homogeneous fibre bundle. Indeed, we take $Q = SO(E)$ the $SO(2n)$ -bundle of positively oriented orthonormal frames of E , $H = SO(2n)$ and $K = U(n)$ (embedded in $SO(2n)$ via $A + iB \mapsto \begin{pmatrix} A & -B \\ B & A \end{pmatrix}$). K

is the subgroup of $SO(2n)$ which commutes with $J_0^\varepsilon = \varepsilon \begin{pmatrix} 0 & -\text{Id} \\ \text{Id} & 0 \end{pmatrix}$. The involution $T = \text{Int} J_0^\varepsilon$ in $SO(2n)$ gives rise to the symmetric space $H/K = \Sigma^\varepsilon(\mathbb{R}^{2n})$, and to the following symmetric decomposition $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{p}$ with

$$\begin{aligned} \mathfrak{k} &= \{A \in \mathfrak{so}(2n) \mid [A, J_0^\varepsilon] = 0\} \\ \mathfrak{p} &= \{A \in \mathfrak{so}(2n) \mid AJ_0^\varepsilon + J_0^\varepsilon A = 0\}. \end{aligned}$$

Concerning ω , we take the $\mathfrak{so}(2n)$ -valued connection 1-form on Q corresponding to the covariant derivative ∇ in E : if $e = (e_1, \dots, e_{2n})$ is a (local) moving frame of E (i.e. a section of Q) then

$$\nabla(e_1, \dots, e_{2n}) = (e_1, \dots, e_{2n})\omega(e; de).$$

³⁶The vertical harmonicity is w.r.t. the splitting defined by example 4.4. See section 4.3.4 and (more precisely) theorem 4.21, for the above characterisation of vertical harmonicity in $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$.

Now, let us consider the isomorphism of bundle:

$$\mathcal{J}: e.U(n) \in \mathcal{SO}(E)/U(n) \xrightarrow{\cong} J \in \Sigma^\varepsilon(E) | \text{Mat}_{e.U(n)}(J) = J_0^\varepsilon.$$

The isomorphism \mathcal{J} defines a bijection between the set of section of $\pi: N \rightarrow M$ ³⁷ and the set of complex structure of E (sections of π_Σ): $s \in \mathcal{C}(N) \rightarrow J = \mathcal{J} \circ s \in \mathcal{C}(\Sigma^+(E))$.

The existence of a (positive) complex structure J in E – i.e. a section of $\pi_\Sigma: \Sigma^+(E) \rightarrow M$ – is equivalent to the existence of an $U(n)$ -reduction of the principal bundle $\mathcal{SO}(E) \rightarrow M$: J defines a Hermitian structure on E and then the $U(n)$ -subbundle of unitary frames for this Hermitian structure, and vice versa.

The isomorphism of bundle over M , $\mathcal{J}: N \rightarrow \Sigma^+(E)$ defines tautologically a canonical complex structure on $\pi^*E \rightarrow N$ (which we still denote by \mathcal{J})³⁸ $\mathcal{J}: N \rightarrow \Sigma^+(\pi^*E)$. Under this identification, let us precise the subbundles \mathfrak{p}_Q and \mathfrak{k}_Q . First, we have $\mathfrak{h}_Q = \mathfrak{so}(E)$ ³⁹, the bundle of skew-symmetric endomorphism of E and then⁴⁰

$$\begin{aligned} (\mathfrak{k}_Q)_y &= \{F \in \mathfrak{so}(E_{\pi(y)}) | [F, \mathcal{J}(y)] = 0\} =: \mathfrak{so}_+(E_{\pi(y)}, \mathcal{J}(y)) =: \mathfrak{so}_+(\pi^*E)_y \\ (\mathfrak{p}_Q)_y &= \{F \in \mathfrak{so}(E_{\pi(y)}) | F\mathcal{J}(y) + \mathcal{J}(y)F = 0\} =: \mathfrak{so}_-(E_{\pi(y)}, \mathcal{J}(y)) =: \mathfrak{so}_-(\pi^*E)_y. \end{aligned}$$

Then the decomposition following $\pi^*\mathfrak{h}_Q = \mathfrak{k}_Q \oplus \mathfrak{p}_Q$ of any element $F \in \pi^*\mathfrak{h}_Q = \mathfrak{so}(\pi^*E)$ is given by

$$F = \frac{1}{2}\mathcal{J}\{F, \mathcal{J}\} + \frac{1}{2}\mathcal{J}[F, \mathcal{J}]$$

where $\{, \}$ is the anticommutator.

Remark 4.8 The canonical complex structure \mathcal{J} is a section of the associated bundle over N : $\Sigma^+(\pi^*E) = \pi^*(Q \times_H \Sigma^\varepsilon(\mathbb{R}^{2n})) = \pi^*Q \times_H \Sigma^\varepsilon(\mathbb{R}^{2n})$, so that it can be lifted to a H -equivariant map $\tilde{\mathcal{J}}: \pi^*Q \rightarrow \Sigma^\varepsilon(\mathbb{R}^{2n}) \subset \mathfrak{h}$, which is given by

$$\tilde{\mathcal{J}}: (e.K, e.h^{-1}) \in \pi^*Q \mapsto hJ_0^\varepsilon h^{-1} \in \Sigma^\varepsilon(\mathbb{R}^{2n}).$$

Remark that the restriction of $\tilde{\mathcal{J}}$ to $Q \subset \pi^*Q$ is the constant map J_0^ε (the inclusion $Q \subset \pi^*Q$ is given by $e \mapsto (e.U(n), e)$), and that $\tilde{\mathcal{J}}: \pi^*Q \rightarrow \Sigma^\varepsilon(\mathbb{R}^{2n})$ is the H -equivariant extension of the K -equivariant constant map J_0^ε on Q . $\tilde{\mathcal{J}}$ can also be given in term of \mathcal{J} by

$$\tilde{\mathcal{J}}: (y; e) \in \pi^*Q \mapsto \text{Mat}_e(\mathcal{J}(y)) \in \Sigma^\varepsilon(\mathbb{R}^{2n}) \subset \mathfrak{h}.$$

Furthermore, we have a canonical identification $N = Q \times_H H/K$ (via $[e, h.K] \mapsto (e.h).K$) and the identification depending on J_0^ε : $H/K = \Sigma^\varepsilon(\mathbb{R}^{2n})$ (via $h.K \mapsto hJ_0^\varepsilon h^{-1}$) so that $N = Q \times_H \Sigma^\varepsilon(\mathbb{R}^{2n})$ (via $e.K \mapsto [e, J_0]$). Then under this last identification, \mathcal{J} is the restriction to N of the canonical identification

$$\begin{aligned} \mathfrak{h}_Q := Q \times_H \mathfrak{h} &\xrightarrow{\cong} \mathfrak{so}(E) \\ [e, a] &\mapsto A | \text{Mat}_e(A) = a. \end{aligned} \tag{78}$$

Therefore

$$\mathfrak{h}_Q = \mathcal{J}^*\mathfrak{so}_+(\pi_\Sigma^*E) \quad \text{and} \quad \mathfrak{p}_Q = \mathcal{J}^*\mathfrak{so}_-(\pi_\Sigma^*E),$$

with the notations of example 4.2.

³⁷In all the section 4.3, as it was the case in all the section 4.2, $N := Q/K$.

³⁸and which is in fact nothing but $\mathcal{J}^*\mathcal{I} = \mathcal{I} \circ \mathcal{J}$, see example 4.2.

³⁹See remark (4.8) (and more precisely equation (78)) for the identification map.

⁴⁰Since π^*E is canonically endowed with the complex structure \mathcal{J} , we need not to precise this latter in the notation $\mathfrak{so}_\pm(\pi^*E)$, whereas E_x , for $x \in M$, could be endowed with any element $J_x \in \Sigma^+(E_x)$, this is why we must precise it in the notation $\mathfrak{so}_\pm(E_x, J_x)$.

Let us now express the homogeneous connection ϕ , the curvature forms Φ and the canonical connection ∇^c in terms of \mathcal{J} (following [51]).

Theorem 4.13 [51] *If $A, B \in TN$, $F \in \mathcal{C}(\mathfrak{p}_Q)$ then:*

- (i) $\phi A = \frac{1}{2} \mathcal{J} \cdot \nabla_A \mathcal{J}$
- (ii) $\Phi(A, B) = \frac{1}{2} \mathcal{J} [\pi^* R(A, B), \mathcal{J}]$, where R is the curvature operator of the ∇ .
- (iii) $\nabla^c F = \frac{1}{2} \mathcal{J} [\nabla_A F, \mathcal{J}]$

Theorem 4.14 [51] *Let $s \in \mathcal{C}(\pi)$ and $J = s^* \mathcal{J}$ be the corresponding complex structure, and $\nabla^* \nabla = -\text{Tr} \nabla^2$, the rough Laplacian of E . Then*

- (i) $I(d^v s) = \frac{1}{2} J \cdot \nabla J = \frac{1}{4} [J, \nabla J]$. Thus s is horizontal if and only if J is parallel.
- (ii) $I(\Pi^v(s)) = \frac{1}{4} [J, \nabla^2 J]$. Thus s is superflat if and only if $\nabla^2 J$ commutes with J .
- (iii) $I(\tau^v(s)) = -\frac{1}{4} [J, \nabla^* \nabla J]$. Thus s is a harmonic section if and only if $\nabla^* \nabla J$ commutes with J .
- (iv) $s^* \Phi = \frac{1}{2} J [R, J]$.

These properties hold also for maps $f \in \mathcal{C}^\infty(L, N)$, (L, b) being a Riemannian manifold: (i), (ii), (iii) without any change and (iv) becoming $f^* \Phi = \frac{1}{2} [u^* R, J]$, with $u = \pi \circ f$.

From theorem 4.13-(i) (or theorem 4.14-(i)) it follows that $d\mathcal{J}$ sends the decomposition $TN = \mathcal{V} \oplus \mathcal{H}$ onto the decomposition $T\Sigma^\varepsilon(E) = \mathcal{V}^\Sigma \oplus \mathcal{H}^\Sigma$ coming from ∇ (see example 4.2) so that we can consider $\pi_\Sigma: \Sigma^\varepsilon(E) \rightarrow M$ as a homogeneous fibre bundle over M with structure group $H = SO(2n)$ and $K = U(n)$. Besides, since the vertical and horizontal subbundles corresponds via \mathcal{J} , then we can conclude according to (52) and (62) that \mathcal{J} is an isometry.

Moreover, we see that s is vertically harmonic in N if and only if the rough Laplacian $\nabla^* \nabla J$ of J in $\mathfrak{so}(E)$ is vertical (i.e. in \mathcal{V}_J^Σ , see example 4.2) so that we recover the definition of vertically harmonic twistor lifts used in [34] and [10]. More precisely, via the isometry \mathcal{J} , **the vertical tension field of s** – which is, let us recall it, defined using the Levi-Civita connection in N which corresponds via the isometry \mathcal{J} to the Levi-Civita connection in $\Sigma^+(E)$ – **is exactly the vertical part in $\pi_\Sigma^* \mathfrak{so}(E)$ of the rough laplacian of J :**

$$d\mathcal{J}(\tau^v(s)) = \nabla_{\tau^v(s)} \mathcal{J} = -2\mathcal{J} \circ \phi(\tau^v(s)) = \frac{1}{2} J [J, \nabla^* \nabla J]$$

according to theorem 4.13-(i) and theorem 4.14-(iii). Concretely, **to compute the vertical tension field in $\Sigma^+(E)$** , instead of using the (abstract) Levi-Civita connection, it is enough to **take the vertical part of the rough Laplacian** (which uses the concrete metric connection ∇).

4.3.3 The twistor bundle $\mathcal{Z}_{2k}(E)$ of a Riemannian vector bundle

We give ourself the same ingredients and notations as in example 4.3. Let us suppose that the vector bundle E is oriented. Then the bundle $\pi_{\mathcal{Z}}: \mathcal{Z}_{2k}^{\alpha}(E) \rightarrow M$ is a homogeneous fibre bundle. Indeed, we take $Q = \mathcal{SO}(E)$, $H = \mathcal{SO}(2n)$ and $K = \mathbb{U}_0(J_0^{\alpha})$. Let us recall that the order r automorphism $T = \text{Int}J_0^{\alpha}$ in $\mathcal{SO}(2n)$ gives rise to the r -symmetric space $H/K = \mathcal{Z}_{2k}^{\alpha}(\mathbb{R}^{2n})$, and to the following reductive decomposition $\mathfrak{h} = \mathfrak{k} \oplus \mathfrak{p}$ with

$$\mathfrak{k} = \mathfrak{so}_0(J_0^{\alpha}) \quad \text{and} \quad \mathfrak{p} = \mathfrak{so}_*(J_0^{\alpha}) := \left(\bigoplus_{j \in \mathbb{Z}/r\mathbb{Z} \setminus \{0\}} \mathfrak{so}_j^{\mathbb{C}}(J_0^{\alpha}) \right) \cap \mathfrak{so}(2n).$$

Concerning ω , we take the same as in the previous example. Now let us consider the isomorphism of bundle:

$$\mathcal{J}: e.\mathbb{U}_0(J_0^{\alpha}) \in \mathcal{SO}(E)/\mathbb{U}_0(J_0^{\alpha}) \xrightarrow{\cong} J \in \mathcal{Z}_{2k}^{\alpha}(E) | \text{Mat}_e(J) = J_0^{\alpha}. \quad (79)$$

The isomorphism \mathcal{J} defines a bijection between the sections of $\pi: N \rightarrow M$ and the set of sections of $\pi_{\mathcal{Z}}: \mathcal{Z}_{2k}^{\alpha}(E) \rightarrow M$, $s \in \mathcal{C}(N) \mapsto J = \mathcal{J} \circ s \in \mathcal{C}(\mathcal{Z}_{2k}^{\alpha}(E))$.

The isomorphism of bundle over M , $\mathcal{J}: N \rightarrow \mathcal{Z}_{2k}^{\alpha}(E)$ defines tautologically a canonical $2k$ -structure⁴¹ on $\pi^*E \rightarrow N$ (still denoted by \mathcal{J}), $\mathcal{J}: N \rightarrow \mathcal{Z}_{2k}^{\alpha}(\pi^*E)$. Under this consideration, we therefore have $\mathfrak{h} = \mathfrak{so}(E)$ and for all $y \in N$,

$$\begin{aligned} \mathfrak{k}_Q &= \mathfrak{so}_0(\pi^*E, \mathcal{J}) \\ \mathfrak{p}_Q &= \mathfrak{so}_*(\pi^*E, \mathcal{J}) = \left(\bigoplus_{j \in \mathbb{Z}/r\mathbb{Z} \setminus \{0\}} \mathfrak{so}_j^{\mathbb{C}}(\pi^*E, \mathcal{J}) \right) \cap \mathfrak{so}(\pi^*E). \end{aligned}$$

Since π^*E is canonically endowed with \mathcal{J} , we will not precise it and use the notation $\mathfrak{so}_j^{\mathbb{C}}(\pi^*E) := \mathfrak{so}_j^{\mathbb{C}}(\pi^*E, \mathcal{J})$.

Let us consider the surjective morphism of vector bundle

$$\begin{aligned} \text{ad}\mathcal{J}: \quad \pi^*\mathfrak{h}_Q = \mathfrak{so}(\pi^*E) &\longrightarrow \mathcal{B}_*(\pi^*E) = \mathcal{J}.\mathfrak{so}_*(\pi^*E) = \mathcal{J}.\mathfrak{p}_Q \\ (J, A) &\longmapsto \text{ad}J(A) = [J, A] = J \sum_{j=1}^r (1 - \omega_r^j) A_j \end{aligned}$$

where $A_j = [A]_{\mathfrak{so}_j^{\mathbb{C}}(E_x)}$ is the $\mathfrak{so}_j^{\mathbb{C}}(E_x)$ -component of $A \in \mathfrak{so}(E_x)$. The kernel of $\text{ad}\mathcal{J}$ is $\mathfrak{k}_Q = \mathfrak{so}_0(\pi^*E)$ so that $\text{ad}\mathcal{J}$ induces an isomorphism from \mathfrak{p}_Q onto $\mathcal{J}.\mathfrak{p}_Q$. We will set

$$(\text{ad}\mathcal{J})^{-1} = (\text{ad}\mathcal{J}|_{\mathcal{J}.\mathfrak{p}_Q})^{-1} \oplus 0_{\mathcal{J}.\mathfrak{k}_Q}$$

so that

$$(\text{ad}\mathcal{J})^{-1} \circ \text{ad}\mathcal{J} = \text{pr}_{\mathfrak{p}_Q}, \quad \text{the projection on } \mathfrak{p}_Q \text{ along } \mathfrak{k}_Q, \quad \text{and} \quad (80)$$

$$\text{ad}\mathcal{J} \circ (\text{ad}\mathcal{J})^{-1} = \text{pr}_{\mathcal{J}.\mathfrak{p}_Q}, \quad \text{the projection on } \mathcal{J}.\mathfrak{p}_Q \text{ along } \mathcal{J}.\mathfrak{k}_Q. \quad (81)$$

Let us remark that $\mathcal{J}.\mathfrak{p}_Q = \mathcal{J}^*\mathcal{V}^{\mathcal{Z}}$ is the (pullback by \mathcal{J} of the) vertical space of $\pi_{\mathcal{Z}}$ (see example 4.3). More precisely the \mathcal{J} -pullback of the decomposition $\mathcal{V}^{SO(E)}|_{N_{\mathcal{Z}}} = \mathcal{B}_0(\pi_{\mathcal{Z}}^*E) \oplus \mathcal{B}_*(\pi_{\mathcal{Z}}^*E)$ (see example 4.3) is the decomposition $\mathcal{J}.\mathfrak{so}(E) = \mathcal{J}.\mathfrak{k}_Q \oplus \mathcal{J}.\mathfrak{p}_Q$.

Let us now express the homogeneous fibre bundle tools ϕ, Φ and $\nabla^{\mathfrak{p}}$ in terms of \mathcal{J} .

Theorem 4.15 *If $A, B \in TN$, $F \in \mathcal{C}(\mathfrak{p}_Q)$ then*

⁴¹and which is in fact nothing but $\mathcal{J}^*\mathcal{I} = \mathcal{I} \circ \mathcal{J}$, see example 4.3.

- (i) $\nabla \mathcal{J} = -\text{ad} \mathcal{J} \circ \phi$ thus $\phi A = -(\text{ad} \mathcal{J})^{-1} \nabla_A \mathcal{J}$
- (ii) $\Phi(A, B) = (\text{ad} \mathcal{J})^{-1} [\mathcal{J}, \pi^* R(A, B)]$
- (iii) $\nabla_A^p F = (\text{ad} \mathcal{J})^{-1} [\mathcal{J}, \nabla_A F]$

Proof. We follow the proof given in [51] of theorem 4.13.

(i) Let D be the exterior covariant derivative for $\mathfrak{gl}(2n)$ -valued differential form on $\pi^* Q$. Let $\tilde{A} \in TQ$ lifting $A \in TN$, then

$$D\tilde{\mathcal{J}}(\tilde{A}) = d\mathcal{J}(\tilde{A}) + [\omega(\tilde{A}), \tilde{\mathcal{J}}] = [\omega_p(\tilde{A}), J_0^\alpha],$$

since $\tilde{\mathcal{J}} = J_0^\alpha$. Projection on N yields

$$\nabla_A \mathcal{J} = -\text{ad} \mathcal{J} \circ \phi(A).$$

(ii) First we have $\Omega_p = (\text{ad} J_0^\alpha)^{-1} [J_0^\alpha, \Omega]$, according to (80). The right hand side is the restriction to Q of the \mathfrak{h} -valued 2-form $(\text{ad} \tilde{\mathcal{J}})^{-1} [\tilde{\mathcal{J}}, \tilde{\Omega}]$ on $\pi^* Q$, where $\tilde{\Omega}$ is the pullback of Ω . Since $\tilde{\Omega}$ is the curvature of the pullback connection, on projection to N we obtain $\Phi = (\text{ad} \mathcal{J})^{-1} [\mathcal{J}, \pi^* R]$ (by definition of ω , Ω is the lift in Q of the curvature operator R of ∇).

(iii) ∇^ω is the restriction of the tensor product connection in $E^* \otimes E = \text{End}(E)$ to $\mathfrak{h}_Q = \mathfrak{so}(E)$, and its \mathfrak{p}_Q -component follows from (80). This completes the proof. \square

Given A, B 1-forms on N with values in some Lie algebra, we define

$$[A \odot B](X, Y) = [A(X), B(Y)] + [A(Y), B(X)], \quad \forall X, Y \in TN$$

Theorem 4.16 *Let $s \in \mathcal{C}(\pi)$ and $J = s^* \mathcal{J}$ be the corresponding $2k$ -structure. Then*

(i) $I(d^v s) = -(\text{ad} J)^{-1} \nabla J$. Thus s is horizontal if and only if J is parallel.

(ii) $I(\Pi^v(s)) = -(\text{ad} J)^{-1} \nabla^2 J + \frac{1}{2} (\text{ad} J)^{-1} [\nabla J \odot (\text{ad} J)^{-1} \nabla J]$.

Thus s is superflat if and only if $\nabla^2 J - \frac{1}{2} [\nabla J \odot (\text{ad} J)^{-1} \nabla J]$ commutes with J .

(iii) $I(\tau^v(s)) = +(\text{ad} J)^{-1} \nabla^* \nabla J + (\text{ad} J)^{-1} \text{Tr}([\nabla J, (\text{ad} J)^{-1} \nabla J])$.

Thus s is a harmonic section if and only if $\nabla^* \nabla J + \text{Tr}([\nabla J, (\text{ad} J)^{-1} \nabla J])$ commutes with J .

(iv) $s^* \Phi = (\text{ad} J)^{-1} [J, R]$.

These properties hold also for maps $f \in \mathcal{C}^\infty(L, N)$, (L, b) being a Riemannian manifold: (i), (ii), (iii) without any change and (iv) becoming $f^* \Phi = (\text{ad} J)^{-1} [J, u^* R]$, with $u = \pi \circ f$.

Proof. (i) Take the pullback by f of theorem 4.15-(i).

(iv) Take the pullback by f of theorem 4.15-(ii).

(ii) According to theorems 4.6 and 4.5-(i), we have

$$\begin{aligned} I(\Pi^v s) &= \nabla^c(f^* \phi) + \frac{1}{2} f^* \phi^* B \\ &= \nabla^c(f^* \phi) + \frac{1}{2} [f^* \phi, f^* \phi]_{\mathfrak{p}} \quad (\text{since } H/K \text{ is naturally reductive}), \\ &= \nabla^p(f^* \phi) - \frac{1}{2} [f^* \phi, f^* \phi]_{\mathfrak{p}} \quad (\text{according to (4.3)}). \end{aligned}$$

Moreover,

$$\begin{aligned}
\nabla^p(f^*\phi) = \text{pr}_{\mathfrak{p}_Q}(\nabla^\omega(f^*\phi)) &= \text{pr}_{\mathfrak{p}_Q}(\nabla(-(\text{ad}J)^{-1}\nabla J)) \\
&= (\text{ad}J)^{-1} \circ \text{ad}(\nabla J) \circ (\text{ad}J)^{-1}(\nabla J) - (\text{ad}J)^{-1}(\nabla^2 J) \\
&= (\text{ad}J)^{-1}([\nabla J, (\text{ad}J)^{-1}\nabla J]) - (\text{ad}J)^{-1}(\nabla^2 J)
\end{aligned}$$

where in the second line we have used the lemma 4.2 below.

Now, by an immediate computation using (80), we obtain that

$$[(\text{ad}J)^{-1}\nabla J, (\text{ad}J)^{-1}\nabla J]_{\mathfrak{p}_Q} = [\nabla J \wedge (\text{ad}J)^{-1}\nabla J]$$

This completes the proof of (ii).

Then (iii) follows immediately from (ii). This completes the proof. \square

Lemma 4.2 *Let E, F be two vector bundles each one endowed with a connection that we denote indistinctly each by ∇ . Let $A \in E^* \otimes F$ be an linear morphism. Let us consider a splitting $E = K_E \oplus \bar{E}$ where $K_E = \ker A$. Then let us set $B = (A_{\bar{E}})^{-1} \oplus 0_{K_F}$, where K_F is some complementary subbundle to $\bar{F} := A(E)$.*

Then we have

$$\text{pr}_{\bar{F}} \circ \nabla B \circ \text{pr}_{\bar{E}} = -B \circ \nabla A \circ B$$

Proof. Differentiating the equations $A \circ B = \text{pr}_{\bar{F}}$ and $\text{pr}_{\bar{E}} \circ B = B$, we obtain

$$\begin{aligned}
\nabla A \circ B + A \circ \nabla B &= \nabla \text{pr}_{\bar{F}} \\
\nabla \text{pr}_{\bar{E}} \circ B + \text{pr}_{\bar{E}} \circ \nabla B &= \nabla B
\end{aligned}$$

so that multiplying the first equation B on the the left, and injecting it in the second equation, we obtain

$$\begin{aligned}
\nabla B &= \nabla \text{pr}_{\bar{E}} \circ B + B \circ \nabla \text{pr}_{\bar{F}} - B \circ \nabla A \circ B \\
&= \text{pr}_{K_E} \circ \nabla B + B \circ \nabla \text{pr}_{\bar{F}} - B \circ \nabla A \circ B
\end{aligned}$$

This completes the proof. \square

As in 4.3.2 above, we conclude from theorem 4.15-(i) that $d\mathcal{J}$ sends the decomposition $TN = \mathcal{V} \oplus \mathcal{H}$ onto the decomposition $TN_{\mathcal{Z}} = \mathcal{V}^{\mathcal{Z}} \oplus \mathcal{H}^{\mathcal{Z}}$ coming from ∇ (see example 4.3) so that we can consider $\pi_{\mathcal{Z}}: \mathcal{Z}_{2k}^\alpha(E) \rightarrow M$ as a homogeneous fibre bundle over M with structure groups $H = SO(2n)$ and $K = \mathbb{U}_0(J_0^\alpha)$. We will call this structure *the homogeneous fibre bundle structure defined in $N_{\mathcal{Z}}$ by ∇ (or by the Riemannian vector bundle (E, ∇))*.

Besides, since the vertical and horizontal subbundles corresponds via \mathcal{J} , then we can conclude according to (52) and (62) that \mathcal{J} is an isometry.

Moreover, the vertical tension field of J in $N_{\mathcal{Z}} = \mathcal{Z}_{2k}^\alpha$ is given by

$$\begin{aligned}
d\mathcal{J}(\tau^v(s)) = \nabla_{\tau^v(s)}\mathcal{J} &= -(\text{ad}\mathcal{J}) \circ \phi(\tau^v(s)) \\
&= -(\text{ad}\mathcal{J}) \circ (\text{ad}\mathcal{J})^{-1}(\nabla^*\nabla J + \text{Tr}([\nabla J, (\text{ad}J)^{-1}\nabla J])) \\
&= -\left[\nabla^*\nabla J + \text{Tr}([\nabla J, (\text{ad}J)^{-1}\nabla J]\right]_{\mathcal{V}^{\mathcal{Z}}}
\end{aligned}$$

By taking $k = 2$ in the two preceding theorems, we recover of course the results of the previous section: just remark that in this case, $\text{ad}\mathcal{J} = 0_{\mathfrak{k}_Q} \oplus 2L_{\mathcal{J}|_{\mathfrak{p}_Q}}$, and that ∇J anticommutes with J .

Remark 4.9 Let us consider the canonical identification

$$H_Q := Q \times_H H \xrightarrow{\cong} SO(E) \quad (82)$$

$$[e, h] \mapsto A | \text{Mat}_e(A) = h.$$

then \mathcal{J} is the restriction to $N \cong Q \times_H \mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n})$ (via $e.K \mapsto [e, J_0^\alpha]$) of (82).

More generally, for $j \in \mathbb{Z}$, we can consider \mathcal{J}_j the restriction of (82) to $Q/\mathbb{U}_{j-1}(J_0^\alpha) = Q \times_{SO(2n)} (\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}))^j$ (via $e.\mathbb{U}_{j-1}(J_0^\alpha) \mapsto [e, (J_0^\alpha)^j]$):

$$\mathcal{J}_j: e.\mathbb{U}_{j-1}(J_0^\alpha) \in \mathcal{SO}(E)/\mathbb{U}_{j-1}(J_0^\alpha) \xrightarrow{\cong} J \in (\mathcal{Z}_{2k}^\alpha(E))^j | \text{Mat}_e(J) = (J_0^\alpha)^j. \quad (83)$$

Remark 4.10 The previous study could have been done (without any change) for any component $\mathcal{U}_{2k}^\alpha(E)$. In particular, by replacing J_0^α by $(J_0^\alpha)^j$ in what precedes, we get the isomorphism (83)

$$\mathcal{J}_j: \mathcal{SO}(E)/\mathbb{U}_{j-1}(J_0^\alpha) \xrightarrow{\cong} (\mathcal{Z}_{2k}^\alpha(E))^j = \mathcal{U}_p^{j,\alpha}(E),$$

where $p = \frac{2k}{(2k, j)}$, and by applying theorem 4.16, we see that a cross section $s_j: M \rightarrow \mathcal{SO}(E)/\mathbb{U}_{j-1}(J_0^\alpha)$ is horizontal if and only if the corresponding section $J_j = \mathcal{J}_j \circ s_j: M \rightarrow (\mathcal{Z}_{2k}^\alpha(E))^j$ is parallel: $\nabla J_j = 0$.

4.3.4 The Twistor subbundle $\mathcal{Z}_{2k,j}^\alpha(E)$

We continue here the study of example 4.4, $\pi_{\mathcal{Z}}^j: \mathcal{Z}_{2k,j}^\alpha(E, J_j) \rightarrow M$, and prove that it defines a homogeneous bundle fibre bundle. Let us recall that we have a bijection between the set of (global) sections J_j in $(\mathcal{Z}_{2k}^\alpha(E))^j = \mathcal{U}_p^{j,\alpha}(E)$ and the set of $\mathbb{U}_{j-1}(J_0^\alpha)$ -reductions $\pi^j: Q^j \rightarrow M$ of $\mathcal{SO}(E)$, which is given by

$$Q^j = \mathfrak{U}_{j-1}^\alpha(E) := \{e \in \mathcal{SO}(E) | \text{Mat}_e(J_j) = (J_0^\alpha)^j\}. \quad (84)$$

Let us consider such a reduction Q^j (defined by some J_j). Then $\pi^j: Q^j \rightarrow M$ is a principal bundle with structural group $H^j = \mathbb{U}_{j-1}(J_0^\alpha)$ and we take for the second structural group $K = \mathbb{U}_0(J_0^\alpha)$ as in the previous example. Let us recall that the order j automorphism $T = \text{Int}_{J_0^\alpha}|_{\mathbb{U}_{j-1}(J_0^\alpha)}$ gives rise to the j -symmetric space $H^j/K = \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, (J_0^\alpha)^j)$, and to the following reductive decomposition $\mathfrak{h}^j = \mathfrak{k} \oplus \mathfrak{p}^j$ where

$$\begin{aligned} \mathfrak{k} &= \mathfrak{so}_0(J_0^\alpha) \quad \text{and} \\ \mathfrak{p}^j &= \mathfrak{u}_{j-1}^*(J_0^\alpha) := \left(\bigoplus_{q=1}^{(r,j)-1} \mathfrak{so}_{pq}^{\mathbb{C}}(J_0^\alpha) \right) \cap \mathfrak{so}(\mathbb{R}^{2n}) = \mathfrak{u}_{j-1}(J_0^\alpha)/\mathfrak{u}_0(J_0^\alpha) = \mathfrak{so}_0((J_0^\alpha)^j)/\mathfrak{so}_0(J_0^\alpha) \\ &\cong T_{J_0^\alpha} \mathcal{Z}_{2k,j}^\alpha(\mathbb{R}^{2n}, (J_0^\alpha)^j), \end{aligned}$$

the last identification is given by

$$A \in \bigoplus_{q=1}^{(r,j)-1} \mathfrak{so}_{pq}^{\mathbb{C}}(J_0^\alpha) \mapsto A \cdot J_0^\alpha = [A, J_0^\alpha] \in \bigoplus_{q=1}^{(r,j)-1} \mathfrak{B}_{pq}^{\mathbb{C}}(J_0^\alpha)$$

(see section 3.1.3).

For the connection form on Q^j we take

$$\omega^j := \omega_{\mathfrak{h}^j}|_{TQ^j}.$$

We set as usual $N^j = Q^j/K$ which is a Homogeneous fibre bundle over M . Moreover the isomorphism of bundle (and isometry) $\mathcal{J}: N \rightarrow \mathcal{Z}_{2k}^\alpha(E)$ satisfies

$$\mathcal{J}(N^j) = \mathcal{Z}_{2k,j}^\alpha(E, J_j)$$

by definition of \mathcal{J} and Q^j (see (79) and (84)), so that it induces an isomorphism of bundle from N^j onto $N_{\mathcal{Z}}^j$.

Let us denote by $TN^j = \mathcal{V}^j \oplus \mathcal{H}^j$ the splitting in terms of vertical and horizontal subbundles given by ω^j . Then denoting by s_j the cross section in the associated bundle $Q/H^j = \mathcal{S}\mathcal{O}(E)/\mathbb{U}_{j-1}(J_0^\alpha)$ defining the H^j -reduction Q^j (i.e. $\mathcal{J}_j \circ s_j = J_j$)⁴², according to section 4.2.3, we have the following equivalences

$$\begin{array}{ccccc} \omega \text{ is reducible in } Q^j \text{ (to } \omega^j) & \xleftrightarrow{4.2.3} & s_j \text{ is horizontal} & \xleftrightarrow{4.2.3} & \mathcal{H}^j = \mathcal{H}|_{N^j} \\ & & \text{Rmk 4.10} \Downarrow & & \Downarrow \\ & & \nabla J_j = 0 & \xleftrightarrow{\text{ex. 4.4}} & \mathcal{H}^{\mathcal{Z},j} = \mathcal{H}^{\mathcal{Z}}|_{N_{\mathcal{Z}}^j} \end{array}$$

Example 4.10 Let $M = G/H$ be the k -symmetric space corresponding to some $2k$ -symmetric space G/G_0 (see section 2.1.1), and take $(E, \nabla) = (TM, \nabla^M)$, $j = 2$ and J_2 given by lemma 3.2. Then we have

$$\nabla^M J_2 = 0.$$

Indeed $\nabla^M J_2$ lifts in G into

$$(d + \theta_{\mathfrak{h}})J_2^0 = dJ_0 + [\theta_{\mathfrak{h}}, J_0^2] = 0$$

(see lemma 3.2). Therefore we can conclude that in this case ω is reducible in Q^2 (to ω^2).

If ω is not reducible in Q^j (to ω^j), then according to (52) and (62), $N^j \hookrightarrow N$ and $N_{\mathcal{Z}}^j \hookrightarrow N_{\mathcal{Z}}$ are not isometries, and thus we can not say directly that \mathcal{J} induces an isometry from N^j onto $N_{\mathcal{Z}}^j$, even if as we will see below it is effectively the case. As above, the result of (52) and (62), and $d\mathcal{J}(\mathcal{V}^j) = \mathcal{V}^{\mathcal{Z},j}$, is that: $\mathcal{J}: N^j \rightarrow N_{\mathcal{Z}}^j$ is an isometry if and only if $d\mathcal{J}(\mathcal{H}^j) = \mathcal{H}^{\mathcal{Z},j}$.

Now let us come back to the connection form $\omega^j: TQ^j \rightarrow \mathfrak{h}^j \subset \mathfrak{so}(2n)$. It defines a metric covariant derivative $\nabla^{[j]}$ in the associated vector bundle E . Then we have

$$\nabla^{[j]}J_j = 0.$$

Indeed J_j lifts into the H^j -equivariant (constant) map $\tilde{J}_j: e \in Q^j \rightarrow (J_0^\alpha)^j \in (\mathcal{Z}_{2k}^\alpha(\mathbb{R}^{2n}))^j \subset \mathfrak{gl}_{2n}(\mathbb{R}^{2n})$ and $\nabla^{[j]}J_j$ lifts into

$$\bar{D}^j \tilde{J}_j = d\tilde{J}_j + [\omega^j, \tilde{J}_j] = 0 + 0 = 0,$$

since by definition $\mathfrak{h}^j = \mathfrak{u}_{j-1}(J_0^\alpha) = \mathfrak{so}_0((J_0^\alpha)^j)$ commutes with $(J_0^\alpha)^j$.

Remark 4.11 We can do the things more concretely by using a (local) moving frame e in Q^j : $\nabla^{[j]}$ is then characterized by

$$\nabla^{[j]}(e_1, \dots, e_{2n}) = (e_1, \dots, e_{2n}).\omega^j(e; de) \quad (85)$$

⁴²See remark 4.9 and 4.10.

Then by definition of Q^j we have

$$J_j e = e \cdot (J_0^\alpha)^j \quad (86)$$

so that

$$(\nabla^{[j]} J_j) e + J_j (\nabla^{[j]} e) = e \cdot \omega^j (J_0^\alpha)^j$$

then using (86) and (85), we obtain

$$(\nabla^{[j]} J_j) e = e \cdot \omega^j (J_0^\alpha)^j - J_j (e \cdot \omega^j) = e \cdot (\omega^j (J_0^\alpha)^j - (J_0^\alpha)^j \omega^j) = 0$$

since ω^j takes values in $\mathfrak{h}_j = \mathfrak{so}_0((J_0^\alpha)^j)$.

In fact we can characterize $\nabla^{[j]}$ in the following more general way, which in particular generalizes a well-known result of Rawnsley [46] about complex structures on vector bundles.

Theorem 4.17 *Let E be a Riemannian vector bundle as above. Let $p' \in \mathbb{N}^*$ and $J \in \mathcal{C}(\mathcal{U}_{p'}(E))$, then $\text{Ad}J$ defines an automorphism of the linear bundle $\text{End}(E)$ (over Id_M), i.e. a section of $\text{End}(\text{End}(E))$. Then the metric covariant derivative ∇ in E admits a unique decomposition in the form:*⁴³

$$\nabla = \overset{J}{\nabla^0} + \sum_{i=1}^{r_{p'}-1} A_i \quad (87)$$

where $\overset{J}{\nabla^0}$ is a metric covariant derivative for which

$$\overset{J}{\nabla^0} J = 0$$

and $A_i \in \mathcal{C}(T^*M \otimes \mathfrak{so}_i^{\mathbb{C}}(E, J))$, i.e. $J A_i J^{-1} = \omega_r^i A_i$ and $A_i \in \mathfrak{so}(E)^{\mathbb{C}}$.

$\overset{J}{\nabla^0}$ will be called the J -commuting component of ∇ , $A_* = \sum_{i=1}^{r_{p'}-1} A_i \in \mathcal{C}(T^*M \otimes \mathfrak{so}_*(E, J))$ the $\mathfrak{so}_*(E, J)$ -component of ∇ , and A_i the $\mathfrak{so}_i^{\mathbb{C}}(E, J)$ -component of ∇ .

Proof. *Unicity.* Let us suppose that (87) exists then we have

$$\nabla J = \sum_{i=1}^{r_{p'}-1} [A_i, J]$$

so that

$$\sum_{i=1}^{r_{p'}-1} A_i = -(\text{ad}J)^{-1}(\nabla J)$$

(see section 4.3.2) which proves the unicity of $(A_i)_{1 \leq i \leq r_{p'}-1}$ (these are determined by ∇ and J , more precisely these are the components of $-(\text{ad}J)^{-1}(\nabla J)$). Now $\overset{J}{\nabla^0} = \nabla - \sum_{i=1}^{r_{p'}-1} A_i$ is also unique.

Existence. Let $\overset{J}{\nabla^0}$ be any metric covariant derivative commuting with J , that is to say $\overset{J}{\nabla^0}$

⁴³As usual $r_{p'}$ is the order of $\text{Ad}J$, i.e. $r_{p'} = p'$ if p' is odd, and if p' is even then $r = p'$ if $J^{\frac{p'}{2}} \neq -\text{Id}$ and $r_{p'} = \frac{p'}{2}$ if $J^{\frac{p'}{2}} = -\text{Id}$.

corresponds to a connection on the principal bundle of Hermitian frames on $(E, \langle \cdot, \cdot \rangle, J)$ (such a connection always exists, see [40]). Then consider

$$A = \nabla - \nabla^0 \in \mathcal{C}(T^*M \otimes \mathfrak{so}(E))$$

and let $A = \sum_{i=0}^{r_{p'}-1} A_i$ be the decomposition of A following $\mathfrak{so}(E, J)^{\mathbb{C}} = \bigoplus_{i=0}^{r_{p'}-1} \mathfrak{so}_i^{\mathbb{C}}(E, J)$. Let us set

$$\nabla^0 = \nabla^0 + A_0$$

then ∇^0 is a J -commuting metric covariant derivative in E and we have

$$\nabla = \nabla^0 + \sum_{i=1}^{r_{p'}-1} A_i$$

which proves the existence. □

Applying this theorem to J_j , we obtain the following.

Corollary 4.2 $\nabla^{[j]}$ is the J_j -commuting component of ∇ .

Proof. The H -equivariant lift of ∇ is the following derivative on Q :⁴⁴

$$d + \omega = (d + \omega_0) + \sum_{i=1}^{r_{p'}-1} \omega_i \quad (88)$$

where $\omega_i = [\omega]_{\mathfrak{so}_i^{\mathbb{C}}((J_0^\alpha)^j)}$, and in particular $\omega_0 = \omega^j$. Then restricting (88) to Q^j , and projecting on M , we obtain the decomposition (87) of ∇ :

$$\nabla = \nabla^{[j]} + \sum_{i=1}^{r_{p'}-1} A_i$$

that is to say $d + \omega_0$ is the H^j -equivariant lift of ∇^0 , which is thus equal to $\nabla^{[j]}$, and ω_i is the H^j -equivariant lift of the $\mathfrak{so}_i^{\mathbb{C}}(E, J_j)$ -valued 1-form on M , A_i . This completes the proof.

Remark 4.12 Moreover ω is reducible in Q^j ($\nabla J_j = 0$) if and only if $\nabla^{[j]} = \nabla$.

Remark 4.13 Under the hypothesis of theorem 4.17 we have

$$\forall F \in \mathcal{C}(\mathcal{A}_0(E, J)), \quad \nabla^0 F = \text{pr}_{\mathcal{A}_0(E, J)} \circ \nabla F$$

where $\text{pr}_{\mathcal{A}_0(E, J)}: \text{End}(E) \rightarrow \mathcal{A}_0(E, J)$ is the orthogonal projection (i.e. along $\mathcal{A}_*(E, J)$) so that in particular

$$\forall F \in \mathcal{C}(\mathfrak{so}_0(E, J)), \quad \nabla^0 F = \text{pr}_{\mathfrak{so}_0(E, J)} \circ \nabla F$$

⁴⁴Remark that here $r_{p'}$ is the order of $\text{Ad}(J_0^\alpha)^j$, so that $r_{p'} = \frac{r}{(r, j)} = p$.

where $\text{pr}_{\mathfrak{so}_0(E,J)}: \mathfrak{so}(E) \rightarrow \mathfrak{so}_0(E, J)$ is the orthogonal projection. Indeed,

$$\nabla F = \nabla^0 F + \sum_{i=1}^{r_{p'}-1} [A_i, F]$$

and J commutes with ∇^0 and F so with $\nabla^0 F: (\nabla^0 F).J = \nabla^0(F.J) - F\nabla^0 J = \nabla^0(J.F) = J\nabla^0 F$. Moreover $[A_i, F] \in [\mathcal{A}_i(J), \mathcal{A}_0(J)] \subset \mathcal{A}_i(J)$, so that we can conclude. •

The canonical $2k$ -structure in π^*E , $\mathcal{J}: N \rightarrow \pi^*E$ induces by restriction a $2k$ -structure in $\pi^{j*}E$, still denoted by $\mathcal{J}: N^j \rightarrow \pi^{j*}E$.

Now, let us precise the subbundles $\mathfrak{p}_{Q^j}^j$ and \mathfrak{k}_{Q^j} . First, we have $\mathfrak{h}_{Q^j}^j = \mathfrak{u}_0(E, J_j)^{45}$ and then

$$\begin{aligned} \mathfrak{k}_{Q^j} &= \mathfrak{k}_{Q|Q^j} = \mathfrak{so}_0(\pi^{j*}E, \mathcal{J}) \\ \mathfrak{p}_{Q^j}^j &= \mathfrak{u}_{j-1}^*(E, \mathcal{J}) = \left(\bigoplus_{i \in p.\mathbb{Z}_r \setminus \{0\}} \mathfrak{so}_i^{\mathbb{C}}(\pi^{j*}E, \mathcal{J}) \right) \cap \mathfrak{so}(\pi^{j*}E). \end{aligned}$$

The morphism of vector bundle $\text{ad}\mathcal{J}: \mathfrak{so}(\pi^*E) \rightarrow \mathcal{J}.p_Q$ induces a surjective morphism from $\pi^{j*}\mathfrak{h}_{Q^j}^j = \mathfrak{u}_{j-1}(\pi^{j*}E, \mathcal{J})$ onto $\mathcal{J}.p_{Q^j}^j$, with kernel \mathfrak{k}_{Q^j} :

$$\begin{aligned} \text{ad}\mathcal{J}: \pi^{j*}\mathfrak{h}_{Q^j}^j = \mathfrak{u}_{j-1}(\pi^{j*}E, \mathcal{J}) &\longrightarrow \mathcal{J}.p_{Q^j}^j = \mathcal{J}.p_{Q^j}^j \\ (J, A) &\longmapsto \text{ad}J(A) = [J, A] = J \sum_{i=1}^{(r,j)-1} (1 - \omega_r^{ip}) A_{ip} \end{aligned}$$

where $A_i = [A]_{\mathfrak{so}_i^{\mathbb{C}}(E_x)}$.

As above, now we express the homogeneous fibre bundle tools ϕ^j , Φ^j and ∇^{p^j} in terms of \mathcal{J} .

Theorem 4.18 *If $A, B \in TN^j$, $F \in \mathcal{C}(p_{Q^j}^j)$ then*

- (i) $\nabla^{[j]}\mathcal{J} = -\text{ad}\mathcal{J} \circ \phi^j$ thus $\phi^j A = -(\text{ad}\mathcal{J})^{-1} \nabla_A^{[j]}\mathcal{J}$
- (ii) $\Phi^j(A, B) = (\text{ad}\mathcal{J})^{-1} \left[\mathcal{J}, \pi^{j*} R^{\nabla^{[j]}}(A, B) \right]$ where $R^{\nabla^{[j]}}$ is the curvature of $\nabla^{[j]}$.
- (iii) $\nabla_A^{p^j} F = (\text{ad}\mathcal{J})^{-1} \left[\mathcal{J}, \nabla_A^{[j]} F \right]$

In the following theorem, we use the notation of 4.2.3. In particular, we denote by " \cdot^v " instead of " \cdot^v " the vertical component in $\mathcal{V}^j \subset TN^j$.

Theorem 4.19 *Let $s \in \mathcal{C}(\pi^j)$ and $J = s^*\mathcal{J}$ be the corresponding $2k$ -structure. Then*

- (i) $I(d^v s) = -(\text{ad}J)^{-1} \nabla^{[j]} J$. Thus s is horizontal if and only if J is $\nabla^{[j]}$ -parallel.
- (ii) $I(\Pi^v(s)) = -(\text{ad}J)^{-1} (\nabla^{[j]})^2 J + \frac{1}{2} (\text{ad}J)^{-1} \left[\nabla^{[j]} J \odot (\text{ad}J)^{-1} \nabla^{[j]} J \right]$.

Thus s is superflat if and only if $(\nabla^{[j]})^2 J - \frac{1}{2} \left[\nabla^{[j]} J \odot (\text{ad}J)^{-1} \nabla^{[j]} J \right]$ commutes with J .

⁴⁵ The restriction of the identification (78), $\mathfrak{h}_Q \cong \mathfrak{so}(E)$ to $\mathfrak{h}_{Q^j}^j$ gives rises to an identification $\mathfrak{h}_{Q^j}^j := Q^j \times_{H^j} \mathfrak{h}^j \cong \mathfrak{so}_0(E, J_j)$.

$$(iii) \quad I(\tau^\vee(s)) = +(\text{ad}J)^{-1}\nabla^{[j]*}\nabla^{[j]}J + (\text{ad}J)^{-1}\text{Tr}\left([\nabla^{[j]}J, (\text{ad}J)^{-1}\nabla^{[j]}J]\right).$$

Thus s is a harmonic section if and only if $\nabla^{[j]*}\nabla^{[j]}J + \text{Tr}\left([\nabla^{[j]}J, (\text{ad}J)^{-1}\nabla^{[j]}J]\right)$ commutes with J .

$$(iv) \quad s^*\Phi^j = (\text{ad}J)^{-1}\left[J, R^{\nabla^{[j]}}\right].$$

These properties hold also for maps $f \in \mathcal{C}^\infty(L, N)$, (L, b) being a Riemannian manifold: (i), (ii), (iii) without any change and (iv) becoming $f^*\Phi^j = (\text{ad}J)^{-1}\left[J, u^*R^{\nabla^{[j]}}\right]$, with $u = \pi \circ f$.

Proof. Let us endow E with $\nabla^{[j]}$ and apply the theorems 4.15 and 4.16 (with the Riemannian vector bundle $(E, \nabla^{[j]})$), then by restriction to N^j , we obtain theorems 4.18 and 4.19. Indeed, in this case ω is reducible in Q^j and then everything corresponds via the reduction $N^j \hookrightarrow N$, according to propositions 4.4 and 4.5. This completes the proof. \square

As above, from theorem 4.18-(i), we conclude that $d\mathcal{J}$ sends the decomposition $TN^j = \mathcal{V}^j \oplus \mathcal{H}^j$ onto the decomposition $TN_{\mathcal{Z}}^j = \mathcal{V}^{\mathcal{Z},j} \oplus \mathcal{H}^{\mathcal{Z},j}$ (see example 4.4) so that we can consider $\pi_{\mathcal{Z}}^j: N_{\mathcal{Z}}^j(E) \rightarrow M$ as a homogeneous fibre bundle over M with structure groups $H^j = \mathbb{U}_{j-1}(J_0^\alpha)$ and $K = \mathbb{U}_0(J_0^\alpha)$. We will call this structure *the homogeneous fibre bundle structure defined by (the J_j -commuting part of) ∇* .

Besides, since the vertical and horizontal subbundles corresponds via \mathcal{J} , then we can conclude according to (52) and (62) that $\mathcal{J}: N^j \rightarrow N_{\mathcal{Z}}^j$ is an isometry.

Moreover, the vertical tension field of J in $N_{\mathcal{Z}}^j = \mathcal{Z}_{2k,2}^\alpha(E, J_j)$ is given by

$$d\mathcal{J}(\tau^\vee(s)) = -\left[\nabla^{[j]*}\nabla^{[j]}J + \text{Tr}\left([\nabla^{[j]}J, (\text{ad}J)^{-1}\nabla^{[j]}J]\right)\right]_{\mathcal{V}^{\mathcal{Z},j}}$$

Remark 4.14 According to 4.2.3, the canonical connection in $\mathfrak{p}_{Q^j}^j \rightarrow N^j$ is the restriction of the canonical connection in $\mathfrak{p}_Q \rightarrow N$, to $\mathfrak{p}_{Q^j}^j$.

Remark 4.15 If we endow E with $\nabla^{[j]}$ and apply the theorems 4.15 and 4.16 (with the Riemannian vector bundle $(E, \nabla^{[j]})$), then by restriction to N^j , we obtain theorems 4.18 and 4.19. In particular, superflatness and vertical harmonicity (for sections in N^j) are the same in N^j and N . This is what happens in particular in example 4.10.

The particular case of $\mathcal{Z}_{2k,2}^\alpha(E, J_2)$ According to theorem 3.2, we will be especially interested by this subcase in our interpretation of the even elliptic integrable system. In this subcase the fibre $H^2/K = \mathcal{Z}_{2k,2}(\mathbb{R}^{2n}, (J_0^\alpha)^2)$ is symmetric so that we obtain simplifications (coming in particular from the facts that $\nabla^c = \nabla^p$ and that any section $J \in \mathcal{C}(\pi_{\mathcal{Z}}^2)$ anticommutes with $\nabla^{[2]}J$) in theorems 4.18 and 4.19 which then take the same forms as theorems 4.13 and 4.14 about the twistor bundle $\Sigma^\varepsilon(E)$, just by doing the change $\nabla \longleftrightarrow \nabla^{[2]}$. Therefore the case $\mathcal{Z}_{2k,2}^\alpha(E, J_2)$ is very similar to that of $\Sigma^\varepsilon(E)$.

Before writing the simplified theorems for $j = 2$, let us do some useful observations.

First, we have⁴⁶

$$\mathfrak{k}_{Q^2} = \mathfrak{so}_0(\pi^{2*}E, \mathcal{J}) = \{A \in \mathfrak{so}(\pi^{2*}E) \mid [A, \mathcal{J}] = 0\} = \mathfrak{so}_{(+1)}(\pi^{2*}E, \mathcal{J}) \quad (89)$$

$$\mathfrak{p}_{Q^2}^2 = \mathfrak{so}_{\frac{2}{2}}(\pi^{2*}E, \mathcal{J}) = \{A \in \mathfrak{so}(\pi^{2*}E) \mid A.\mathcal{J} + \mathcal{J}.A = 0\} = \mathfrak{so}_{(-1)}(\pi^{2*}E, \mathcal{J}). \quad (90)$$

⁴⁶with notation defined in remark 3.3.

Then $\text{ad}\mathcal{J}$ induces a surjective morphism from $\pi^{2*}\mathfrak{h}_{Q^2}^2 = \mathfrak{u}_1(\pi^{2*}E, \mathcal{J})$ onto $\mathcal{J}\cdot\mathfrak{p}_{Q^2}^2 = \mathcal{B}_{\frac{r}{2}}(\pi^{2*}E, \mathcal{J})$ with kernel \mathfrak{k}_{Q^2}

$$\begin{aligned} \text{ad}\mathcal{J}: \quad \mathfrak{u}_1(\pi^{2*}E, \mathcal{J}) = \mathfrak{so}_{(+1)}(\pi^{2*}E, \mathcal{J}) \oplus \mathfrak{so}_{(-1)}(\pi^{2*}E, \mathcal{J}) &\longrightarrow \mathcal{B}_{(-1)}(\pi^{2*}E, \mathcal{J}) = \mathcal{J}\cdot\mathfrak{so}_{(-1)}(\pi^{2*}E, \mathcal{J}) \\ (J, A_0 + A_1) &\longmapsto \text{ad}J(A) = [J, A] = 2JA_1 \end{aligned}$$

where we denote by $A_0 + A_1$ the decomposition following $\mathfrak{so}_{(+1)}(\pi^{2*}E, \mathcal{J}) \oplus \mathfrak{so}_{(-1)}(\pi^{2*}E, \mathcal{J})$ instead of $A_0 + A_{\frac{r}{2}}$.

Theorem 4.20 *If $A, B \in TN^2$, $F \in \mathcal{C}(\mathfrak{p}_{Q^2}^2)$ then*

$$(i) \quad \phi^2 A = -\frac{1}{2}\mathcal{J}^{-1}\nabla^{[2]}\mathcal{J}$$

$$(ii) \quad \Phi^2(A, B) = \frac{1}{2}\mathcal{J}^{-1} \left[\mathcal{J}, \pi^{2*}R^{\nabla^{[2]}}(A, B) \right] \text{ where } R^{\nabla^{[2]}} \text{ is the curvature of } \nabla^{[2]}.$$

$$(iii) \quad \nabla_A^c F = \frac{1}{2}\mathcal{J}^{-1} \left[\mathcal{J}, \nabla_A^{[2]}F \right]^{47}$$

Theorem 4.21 *Let $s \in \mathcal{C}(\pi^2)$ and $J = s^*\mathcal{J}$ be the corresponding $2k$ -structure. Then*

$$(i) \quad I(d^\vee s) = -\frac{1}{2}J^{-1}\nabla_A^{[2]}J. \text{ Thus } s \text{ is horizontal if and only if } J \text{ is } \nabla^{[2]}\text{-parallel.}$$

$$(ii) \quad I(\Pi^\vee(s)) = -(\text{ad}J)^{-1}(\nabla^{[2]})^2 J = -\frac{1}{4} \left[J^{-1}, (\nabla^{[2]})^2 J \right].$$

Thus s is superflat if and only if $(\nabla^{[2]})^2 J$ commutes with J .

$$(iii) \quad I(\tau^\vee(s)) = (\text{ad}J)^{-1}\nabla^{[2]*}\nabla^{[2]}J = \frac{1}{4} \left[J^{-1}, \nabla^{[2]*}\nabla^{[2]}J \right].$$

Thus s is a harmonic section if and only if $\nabla^{[2]}\nabla^{[2]}J$ commutes with J .*

$$(iv) \quad s^*\Phi^2 = \frac{1}{2}J^{-1} \left[J, R^{\nabla^{[2]}} \right].$$

These properties hold also for maps $f \in \mathcal{C}^\infty(L, N)$, (L, b) being a Riemannian manifold: (i), (ii), (iii) without any change and (iv) becoming $f^\Phi^2 = \frac{1}{2}J^{-1} \left[J, u^*R^{\nabla^{[2]}} \right]$, with $u = \pi \circ f$.*

Let us add that the vertical tension field in $N_{\frac{r}{2}}^2$ is given by

$$d\mathcal{J}(\tau^\vee(s)) = - \left[\nabla^{[2]*}\nabla^{[2]}J \right]_{\mathfrak{V}^{\mathbb{Z}, 2}} = -\frac{1}{2}J \left[J^{-1}, \nabla^{[2]*}\nabla^{[2]}J \right] \quad (91)$$

4.4 Geometric interpretation of the even minimal determined system

4.4.1 The injective morphism of homogeneous fibre bundle $\mathfrak{J}_{J_0}: G/G_0 \hookrightarrow \mathcal{Z}_{2k, 2}^{\alpha_0}(G/H, J_2)$.

Here, we want to ask ourself if the inclusion $\mathfrak{J}_{J_0}: G/G_0 \hookrightarrow \mathcal{Z}_{2k, 2}^{\alpha_0}(G/H, J_2)$ given by theorem 3.2 conserves the homogeneous fibre bundle structure, in particular: the vertical harmonicity is it conserved. We use the notations of 4.3.1 and 4.3.4 (with $E = TM$, ∇ a metric connection on

⁴⁷see remark 4.14

M and $j = 2$)⁴⁸. First, we see that \mathfrak{J}_{J_0} is obtained under quotient from the following injective morphism of bundle (which is an embedding if G is closed in $\text{Is}(M)$):

$$\begin{aligned} \mathfrak{J}_{e_0}: \quad G &\hookrightarrow Q^2 = \mathcal{U}_1^{\alpha_0}(G/H, J_2) \subset \mathcal{SO}(M) \\ g &\longmapsto g \cdot e_0 \end{aligned} \quad (92)$$

where $e_0 \in \mathcal{SO}(T_{p_0}M)$ is such that $\text{Mat}_{e_0}(J_0) = J_0^{\alpha_0}$, and $J_0 = \tau_{\mathfrak{m}}$. In other words $G \rightarrow M$ is a reduction of $\mathcal{U}_1^{\alpha_0}(G/H, J_2) \rightarrow M$ itself a reduction of $\mathcal{SO}(M) \rightarrow M$.

Further quotienting in (92) by $\mathbb{U}_0(J_0^\alpha)$ the target space and then by G_0 the domain, we obtain (by definition of $G_0 = G^\tau \cap H$, see theorem 3.2) the injective morphism of bundle

$$\mathfrak{J}_{\bar{e}_0}: g \cdot G_0 \longmapsto (g \cdot e_0) \mathbb{U}_0(J_0^\alpha) \in \mathcal{U}_1^{\alpha_0}(M, J_2) / \mathbb{U}_0(J_0^\alpha) \subset \mathcal{SO}(M) / \mathbb{U}_0(J_0^\alpha)$$

where $\bar{e}_0 = e_0 \mathbb{U}_0(J_0^\alpha) \in N^2$, and finally composing with \mathcal{J} (in the target space) we obtain the map \mathfrak{J}_{J_0} :

$$g \cdot G_0 \longmapsto g \cdot (e_0 \mathbb{U}_0(J_0^\alpha)) \xrightarrow{\mathcal{J}} J = g J_0 g^{-1} \in \mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2).$$

Since \mathfrak{J}_{J_0} (resp. $\mathfrak{J}_{\bar{e}_0}$) is an injective morphism of bundle (and an immersion) $d\mathfrak{J}_{J_0}$ (resp. $d\mathfrak{J}_{\bar{e}_0}$) induces an injective morphism of bundle from the vertical subbundle $\mathcal{V}^{G/G_0} = [\mathfrak{g}_k]$ into the vertical subbundle $\mathcal{V}^{\mathcal{Z},2}$ (resp. \mathcal{V}^2).

\mathfrak{J}_{J_0} is the restriction to G/G_0 of the inclusion map $\mathfrak{J}: \text{End}(G/H) \rightarrow M \times \text{End}(\mathfrak{g})$ (see 1.7). Indeed, we have the inclusion depending on J_0 : $g \cdot G_0 \in G/G_0 \mapsto [g, J_0] \in G \times_H \text{End}(\mathfrak{m}) = \text{End}(G/H)$ which under the inclusion \mathfrak{J} gives $g \cdot G_0 \in G/G_0 \mapsto (g \cdot x_0, \text{Ad}g \circ \tau_{\mathfrak{m}} \circ \text{Ad}g^{-1}) \in M \times \text{End}(\mathfrak{g})$ which is in nothing but \mathfrak{J}_{J_0} (as usual under the identification $TM = [\mathfrak{m}]$). Then under the inclusion $\mathfrak{h}_G \subset \mathfrak{so}(TM)$, we have $\mathfrak{h}_G \subset \mathfrak{h}_{Q^2}^2 = \mathfrak{u}_0(TM, J_2)$. Indeed, under the linear isotropy representation of H in $T_{x_0}M$, we have $H \subset \mathbb{U}_0(T_{x_0}M, J_0^2) = \mathbb{U}_1(T_{x_0}M, J_0)$ so that $\mathfrak{h} \subset \mathfrak{u}_0(T_{p_0}M, J_0^2)$ and thus $\mathfrak{h}_G := G \times_H \mathfrak{h} \subset \mathfrak{u}_0(TM, J_2)$. Moreover let us remark that $\pi^2 \circ \mathfrak{J}_{\bar{e}_0} = \pi$ so that $\pi^* \mathfrak{h}_G \subset \pi^{2*} \mathfrak{h}_{Q^2}^2$ over $\mathfrak{J}_{\bar{e}_0}: N \rightarrow N^2$ (i.e. the inclusion is a morphism of bundle over $\mathfrak{J}_{\bar{e}_0}$).

Furthermore, since $\text{Ad}J_0$ leaves invariant $\mathfrak{h} \subset \mathfrak{u}_1(T_{x_0}M, J_0)$, the restriction to \mathfrak{h} of the symmetric decomposition

$$\mathfrak{u}_1(T_{x_0}M, J_0) = \mathfrak{so}_{(+1)}(T_{x_0}M, J_0) \oplus \mathfrak{so}_{(-1)}(T_{x_0}M, J_0)$$

gives rise to the decomposition $\mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_k$ according to (23), so that the symmetric decomposition given by $\text{Ad}\mathcal{J}$ on $\pi^{2*} \mathfrak{h}_{Q^2}^2 = \mathfrak{u}_1(\pi^{2*}TM, \mathcal{J})$, that is to say

$$\mathfrak{u}_1(TM, \mathcal{J}) = \mathfrak{so}_{(+1)}(\pi^{2*}TM, \mathcal{J}) \oplus \mathfrak{so}_{(-1)}(\pi^{2*}TM, \mathcal{J})$$

gives rise in the $\text{Ad}\mathcal{J}$ -invariant subspace $\pi^* \mathfrak{h}_G \subset \pi^{2*} \mathfrak{h}_{Q^2}^2$ to the symmetric decomposition of $\text{Ad}\mathfrak{J}_{J_0}$ (restricted to $\pi^* \mathfrak{h}_G \subset \mathfrak{so}(\pi^*TM)$)

$$\pi^* \mathfrak{h}_G = \mathfrak{k}_G \oplus \mathfrak{p}_G$$

according to (76). In other words, the decomposition given by (76) injects into the decomposition given by (89) via the inclusion $\pi^* \mathfrak{h}_G \subset \pi^{2*} \mathfrak{h}_{Q^2}^2$.

Now let us interpret theorems 4.11 and 4.12 using the homogeneous fibre bundle structure in $\mathcal{Z}_{2k}^{\alpha_0}(M, J_2)$ defined by the Riemannian vector bundle $(E, \nabla) = (TM, \overset{M}{\nabla^0})$ (in the sense of 4.3.4). We continue to use the same conventions for the notations in N and N^2 (no subscript for N and subscript 2 for N^2 and $N^{\mathcal{Z},2}$). Recall that we have $\mathfrak{J}_{J_0} = \mathcal{J} \circ \mathfrak{J}_{\bar{e}_0} = \mathfrak{J}_{\bar{e}_0}^* \mathcal{J}$ and that $(\overset{M}{\nabla^0})^{[2]} = \overset{M}{\nabla^0}$. Then according to theorems 4.20 and 4.21, theorems 4.11 and 4.12 implies

⁴⁸That is to say the notations for $\mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$ will have the subscript "2" and these of G/G_0 will not have subscript according to 4.3.1 and 4.3.4

Theorem 4.22 *We have the following identities*

(i) $\phi = \mathfrak{I}_{\bar{e}_0}^* \phi^2$

(ii) $\Phi = \mathfrak{I}_{\bar{e}_0}^* \Phi^2$

(iii) $\nabla^c = \nabla^0|[\mathfrak{p}] = \mathfrak{I}_{\bar{e}_0}^* \nabla^{c,2}$, where $\nabla^{c,2}$ is the canonical connection in $\mathfrak{p}_{Q^2}^2$.

Theorem 4.23 *Let $s \in \mathcal{C}(\pi)$ and identify it (temporarily) with $s^* \mathfrak{I}_{\bar{e}_0} \in \mathcal{C}(\pi^2)$. Then under the inclusion $\mathfrak{I}_{\bar{e}_0} : N \rightarrow N^2$, we have:*

(i) $d^v s = d^{v,2} s$

(ii) $\Pi^v s = \Pi^{v,2}$

(iii) $\tau^v s = \tau^{v,2} s$

(iv) $s^* \Phi = s^* \Phi^2$

These properties holds also, without any change, for maps $f \in \mathcal{C}^\infty(L, N)$, (L, b) being a Riemannian manifold.

Let us remark that since the connection form ω , on $Q = \mathcal{SO}(TM)$ defined by ∇^0 is reducible in Q^2 , then in the previous theorems all the "quantities" in N^2 (right handside) can also be computed in $\mathcal{SO}(TM)/\mathbb{U}_0(J_0^{\alpha_0}) \cong \mathcal{Z}_{2k}^{\alpha_0}(M)$, since "everything is reducible" in this case (see remark 4.15).

Now, let us compute the vertical tension field of $J : L \rightarrow N_{\bar{Z}}^2$ for the homogenous fibre bundle structure defined in $N_{\bar{Z}}^2$ by ∇^0 : according to (91) we have

$$\tau^{v,2}(J) = -\frac{1}{2} J \left[J^{-1}, (\nabla^0)^* \nabla^0 J \right]. \quad (93)$$

Then suppose that J takes values in $\mathfrak{I}_{J_0}(G/G_0)$ i.e. $J = f^* \mathfrak{I}_{J_0}$ for some $f \in \mathcal{C}^\infty(L, N)$, then according to theorem 4.23 (and $\mathfrak{I}_{J_0} = \mathcal{J} \circ \mathfrak{I}_{\bar{e}_0}$) we have

$$d\mathfrak{I}_{J_0}(\tau^v(f)) = d\mathcal{J}(\tau^{v,2}(\bar{f})) = \tau^{v,2}(J)$$

where $\bar{f} = \mathcal{J}^{-1} \circ J$ i.e. $J = \bar{f}^* \mathcal{J}$.

The tension fields (and thus vertical harmonicity) correspond (up to multiplicative constant) **via the different inclusions and identifications, in particular via $\mathfrak{I}_{J_0} : N \rightarrow N_{\bar{Z}}^2$.**

The canonical embedding is a reduction of homogeneous fibre bundles. In fact, we can recover theorems 4.22 and 4.23 as well as theorems 4.11, 4.12 by a more conceptual way: just remark that $\mathfrak{I}_{J_0} : N \rightarrow N_{\bar{Z}}^2$ is a reduction of homogeneous fibre bundles.

Theorem 4.24 *The injective morphism of bundle $\mathfrak{I}_{J_0} : G/G_0 \hookrightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$ is a reduction of homogeneous fibre bundles.*

Proof. According to remark 1.2, the canonical connection in the H -principal bundle $G(G/H, H)$ corresponds (via the natural injection \mathfrak{J}_{e_0} defined by equation (92)) to the canonical connection in the G -invariant G -structure $P' = G.e_0$, itself the restriction to P' of the canonical connection in the G -invariant $\mathbb{U}_1(J_0^{\alpha_0})$ -structure $Q^2 = \mathcal{U}_1^{\alpha_0}(G/H, J_2)$, itself the restriction to Q^2 of the canonical connection in $\mathcal{SO}(M)$. This completes the proof. \square

Now, applying propositions 4.4 and 4.5 we obtain:

Corollary 4.3 *The theorems 4.22 and 4.23 as well as theorems 4.11, 4.12 are corollaries of theorems 4.20 and 4.21.*

Using the Levi-Civita connection. In fact in what precedes we can replace the canonical connection in M , $\overset{M}{\nabla}^0$, by (the J_2 commuting part of) the Levi-Civita connection in M .

Proposition 4.7 *The canonical linear connection on M is the J_2 -commuting component of the Levi-Civita connection $\overset{M}{\nabla}$ on M :*

$$\overset{M}{\nabla}^0 = \overset{J_2}{\nabla}^0 = \overset{M}{\nabla}^{[2]}.$$

Proof. According to subsection 1.6, we have

$$\overset{M}{\nabla} = \overset{M}{\nabla}^0 + \frac{1}{2} ([\ ,]_{[m]} + U^M).$$

Then it suffices to apply equation (50) in subsection 3.5.2. This completes the proof. \square

Now, we can rewrite the previous results by replacing the canonical connection by (the J_2 commuting part of) the Levi-Civita connection.

Corollary 4.4 *The homogeneous fibre bundle structures in $N_{\mathbb{Z}}^2$ defined by the canonical affine connection $\overset{M}{\nabla}^0$ and by (the J_2 -commuting part of) the Levi-Civita connection $\overset{M}{\nabla}$, in M , are the same. Therefore theorems 4.11, 4.12 and corollary 4.1 still hold if we replace $\overset{M}{\nabla}^0$ by $\overset{M}{\nabla}^{[2]}$. Moreover theorems 4.22 and 4.23 hold with the homogeneous fibre bundle structure defined in N^2 by the (J_2 -commuting part of) the Levi-Civita connection $\overset{M}{\nabla}$.*

Some additionnal identities Let us conclude this subsection by some additionnal equalities.

Proposition 4.8 *The canonical linear connection on M is the J_2 -commuting component of the connections $\overset{\text{met}}{\nabla}^t$ on M :*

$$\overset{M}{\nabla}^0 = \left(\overset{\text{met}}{\nabla}^t \right)^{[2]}.$$

Proof. Same proof as for proposition 4.7. \square

Proposition 4.9 *Let $J_1 \in \mathcal{C}(\mathcal{U}_{2k}^*(N))$ be the section defined by $\check{J}_0 = \tau_{|\mathfrak{n}}^{-1}$ with, let us recall it, $\mathfrak{n} = \mathfrak{p} \oplus \mathfrak{m} = \mathfrak{g}_k \oplus \mathfrak{m}$, then*

- (i) The J_1 -commuting component of the (π -pullback of the) canonical linear connection in M , $\pi^*\overset{M}{\nabla}^0$, is $\overset{N}{\nabla}^0$ the canonical linear connection in N . This latter is also the J_1 -commuting component of the (π -pullback of the) Levi-Civita connection, and more generally of the connections $\overset{\text{met},M}{\nabla}^t$.
- (ii) The J_1 -commuting component of the Levi-Civita connection in N , $\overset{N}{\nabla}$, is $\overset{N}{\nabla}^0$.
- (iii) More generally, the J_1 -commuting component of $\overset{\text{met},N}{\nabla}^t$ is $\overset{N}{\nabla}^0$.
- (iv) Let $s \in \mathcal{C}(\pi)$ and $J = s^*\mathfrak{J}_{J_0}$ the corresponding $2k$ -structure on M , then $s^*\overset{N}{\nabla}^0$ is the J_1 -commuting component of $\overset{M}{\nabla}^0$, and also the J_1 -commuting component of (the s -pullback of) the Levi-Civita connection on M , $s^*\overset{M}{\nabla}$; and more generally of (the s -pullback of) the connections $\overset{\text{met},N}{\nabla}^t$.

Proof. The first point of (i) becomes obvious when it is written in terms of the lift of the connections: the J_0 -commuting part of $d + \theta_{\mathfrak{h}} = d + \theta_0 + \theta_k$ is $d + \theta_0$ (whereas θ_k is its anticommuting part). For the second point of (i), (ii), (iii), and (iv) use the same method as for proposition 4.7. \square

4.4.2 Conclusion

Now⁴⁹ we can conclude:

Theorem 4.25 *Let (L, j) be a Riemann surface, $f: L \rightarrow N = G/G_0$ be a map and $J = f^*\mathfrak{J}_{J_0}$ the corresponding map into $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$. Then f is a geometric solution of the even minimal determined system $(\text{Syst}(k, \tau))$ if and only if*

- (i) J is an admissible twistor lift ($\Leftrightarrow f$ is horizontally holomorphic)
- (ii) J is vertically harmonic in $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$ endowed with its homogeneous fibre bundle structure defined by the Levi-Civita connection, ∇ , in M :

$$\left[\nabla^{[2]*} \nabla^{[2]} J, J \right] = 0,$$

where $\nabla^{[2]}$ is the J_2 -commuting component of ∇ . ($\Leftrightarrow f$ is vertically harmonic in G/G_0).

Moreover the first condition implies that J is flat in $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$:

$$J^* \Phi^{\mathcal{Z},2} = \left[u^* R^{\nabla^{[2]}}, J \right] = 0,$$

where $\Phi^{\mathcal{Z},2}$ is the homogeneous curvature form in $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$, which means also that J is a flat section in $\text{End}(u^*TM, u^*\nabla^{[2]})$. ($\Leftrightarrow f$ is flat in the homogeneous fibre bundle $N \rightarrow M$).

Furthermore f is a primitive geometric solution (i.e. there exists $m \leq k$ such that f is m -primitive, which is equivalent to say that f is k -primitive) if and only if

- (i) J is an admissible twistor lift

⁴⁹We still consider the same situation as in 4.4.1

(ii) J is parallel: $\nabla^{[2]}J = 0$ ($\Leftrightarrow f$ is horizontal).

Besides in these characterizations, in the points (ii) the Levi-Civita connection can be replaced by any G -invariant metric connection ∇' whose the J_2 -commuting component $\nabla'^{[2]}$ leaves invariant $\mathfrak{h}_G \subset \mathfrak{so}(TM)$. This is the case in particular for the connections

$$\overset{\text{met}}{\nabla}^t = \overset{M}{\nabla}^0 + t([\ , \]_{[\mathfrak{m}]} + U^M), \quad 0 \leq t \leq 1,$$

for which the J_2 -commuting component is the canonical connection on M : $\overset{M}{\nabla}^0$.

5 Generalized harmonic maps

5.1 Affine harmonic maps and holomorphically harmonic maps

A map $u: M \rightarrow N$ between two Riemannian manifolds (M, g) and (N, h) is harmonic if it extremizes the energy functional

$$E(u) = \frac{1}{2} \int_D |du|^2 d\text{vol}_g$$

for all compact subdomains $D \subset M$, where $|du|^2 = \text{Tr}_g(u^*h)$. The associated Euler-Lagrange equation is $\tau(u) := \text{Tr}_g(\nabla du) = 0$, where ∇ is the connection on $T^*M \otimes u^*TN$ induced by the Levi-Civita connections of M and N .

Now, we generalise this definition for maps from a Riemannian manifold into an affine manifold. We present two different ways to do that. The first one is the natural one (see also [29]) and concerns general affine manifolds whereas the second one concerns maps from Riemann surfaces into affine almost complex manifolds.

5.1.1 Affine harmonic maps: general properties

Definition 5.1 Let $s: (M, g) \rightarrow (N, \nabla)$ be a smooth map from a Riemannian manifold (M, g) into an affine manifold (N, ∇) . We set

$$\tau(s) = \text{Tr}_g(\nabla ds) = -\nabla^* ds = *d^\nabla * ds$$

and we say that s is affine harmonic with respect to ∇ or ∇ -harmonic if $\tau(s) = 0$.

Now, let us consider the case where (M, g) is a Riemannian surface i.e. a Riemann surface (L, j) with a Hermitian metric g . Then the action of the Hodge operator $*$ of L , is independent of the metric g on 1-forms ($*\alpha = \alpha \circ j$), whereas in 2-forms (resp. 0-forms) it is multiplied by the factor λ^2 (resp. $\lambda^{-2} > 0$) when the metric is multiplied by the factor $\lambda \in C^\infty(L, \mathbb{R}_+^*)$. Hence the tension field $\tau(f) = *d^\nabla(*df)$ is multiplied by λ^2 , under this last transformation. In particular the affine harmonicity for maps $f: (L, j) \rightarrow (N, \nabla)$ does not depend on the hermitian metric L but only on the conformal structure of (L, j) . Thus we have:

Theorem 5.1 Let (L, j) be a Riemann surface and $f: (L, j) \rightarrow (N, \nabla)$ a smooth map. Let $TL^{\mathbb{C}} = T'L \oplus T''L$ be the decomposition of $TL^{\mathbb{C}}$ into the $(1, 0)$ and $(0, 1)$ -parts, and $d = \partial + \bar{\partial}$ and $\nabla^{f^*(TN)} = \nabla' + \nabla''$ the corresponding splittings. Then we have

$$2\bar{\partial}^\nabla \partial f = d^\nabla df + id^\nabla * df,$$

moreover $d^\nabla df = f^*T$, where T is the torsion of ∇ and $d^\nabla * df = \tau(f)\text{vol}_g$ for any hermitian metric g on L . Therefore the following statements are equivalent:

- (i) $\nabla'' \partial f = 0$
- (ii) $\bar{\partial}^\nabla \partial f = 0$
- (iii) $\nabla_{\frac{\partial}{\partial \bar{z}}} \left(\frac{\partial f}{\partial z} \right) = 0$, for any holomorphic local coordinate $z = x + iy$ (i.e. (x, y) are conformal coordinates for any hermitian metric in L).
- (iv) f is ∇ -harmonic with respect to any hermitian metric in L and torsion free: $f^*T = 0$ (i.e. $T\left(\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}\right) = 0$ for any conformal coordinates (x, y)).

We will say in this case that f is strongly ∇ -harmonic.

Remark 5.1 We remark that the imaginary part (resp. the real part) of equation (ii) (resp. equation (iii)) is the ∇ -harmonic maps equation whereas its real part (resp. imaginary part) is the torsion free equation $f^*T = 0$.

If $T = 0$ or more generally $f^*T = 0$, then f is strongly ∇ -harmonic if and only if it is ∇ -harmonic.

5.1.2 Holomorphically harmonic maps

In the case the target space N is endowed with an almost complex structure J then we have another way to generalise the definition of harmonicity to maps from a Riemann surface into N .

Definition 5.2 Let (L, j) be a Riemann surface and (N, ∇) be an affine manifold endowed with a complex structure J . Let us denote $TN^{\mathbb{C}} = T^{1,0}N \oplus T^{0,1}N$ the corresponding decomposition of $TN^{\mathbb{C}}$. We will say that $f: L \rightarrow N$ is holomorphically harmonic if

$$[\bar{\partial}^{\nabla} \partial f]^{1,0} = 0.$$

Proposition 5.1 Let (L, j) be a Riemann surface and (N, ∇) be an affine manifold endowed with a complex structure J . Then f is holomorphically harmonic if and only if (for any hermitian metric g in M)

$$T_g(f) + J\tau_g(f) = 0$$

where $T_g(f) = *(f^*T) = f^*T(e_1, e_2)$, with (e_1, e_2) an orthonormal basis of TL , or equivalently

$$\tau_g(f)\text{vol}_g = J(f^*T).$$

Therefore f is strongly harmonic if and only if it is torsion free and holomorphically harmonic. In particular, if $T = 0$, or more generally $f^*T = 0$, then f is holomorphically harmonic if and only if it is harmonic. Hence for torsion free connection ∇ harmonicity and holomorphic harmonicity are the same.

Proof. Let $Z = X + iY \in TN^{\mathbb{C}}$ with $X, Y \in TN$, then since $T^{1,0}N$ and $T^{0,1}N$ are given respectively by $\{V \mp iJV, V \in TN\}$, we deduce that

$$[Z]^{1,0} = 0 \Leftrightarrow X + JY = 0 \quad \text{and} \quad [Z]^{0,1} = 0 \Leftrightarrow X - JY = 0.$$

Now, let us apply that to the $TN^{\mathbb{C}}$ -valued 2-form $\bar{\partial}^{\nabla} \partial f$, we obtain

$$[\bar{\partial}^{\nabla} \partial f]^{1,0} = 0 \iff d^{\nabla} df + Jd^{\nabla} * df = 0$$

according to theorem 5.1. This proves the first assertion. Then the assertion concerning strongly harmonicity follows from theorem 5.1(iv). This completes the proof. \square

Let us remark that

Proposition 5.2 In the same situation as in the previous proposition, let us suppose in addition that $\nabla J = 0$. Then if a map $f: L \rightarrow N$ is holomorphic i.e. $df \circ j_L = Jdf$, then f is anti-holomorphically harmonic (i.e. holomorphically harmonic with respect to $-J$).

Proof. f is holomorphic if and only if $df(T^{1,0}L) \subset T^{1,0}N$ i.e. $[\partial f]^{0,1} = 0$. Then we have

$$[\bar{\partial}^\nabla \partial f]^{0,1} = \bar{\partial}^\nabla [\partial f]^{0,1} = 0$$

since ∇ commutes with J . This completes the proof. \square

It can also be useful to observe the following.

Proposition 5.3 *Let (N, J) be an almost complex manifold with an almost complex linear connection that we will denote by ∇^0 . Then let us define a family of connection*

$$\nabla^t = \nabla^0 - tT^0, \quad 0 \leq t \leq 1.$$

Then a map $f: (L, j_L) \rightarrow (N, J)$ from a Riemann surface L into the almost complex manifold N is holomorphically harmonic w.r.t. ∇^1 and J if and only if f is holomorphically harmonic w.r.t. ∇^0 and $-J$. We will say more simply that f is ∇^1 -holomorphically harmonic if and only if it is ∇^0 -anti-holomorphically harmonic.

Holomorphic sections of complex vector bundles Now, we need to do some recalls about complex vector bundles that we will apply in the next paragraph to obtain an interpretation of the holomorphic harmonicity in terms of holomorphic 1-forms.

Let $E \rightarrow M$ be a real vector bundle (over a manifold M) endowed with a complex structure $J \in \text{End}(E)$. Then any frame in the form $(e_x^1, \dots, e_x^r, Je_x^1, \dots, Je_x^r)$ at some point $x \in M$ can be extended to a local frame $(e^1, \dots, e^r, Je^1, \dots, Je^r)$ in the neighbourhood of x . Then there exists an open covering $(U_\alpha)_{\alpha \in I}$ of M and local trivialisations $\Phi_\alpha: (E|_{U_\alpha}, J) \rightarrow U_\alpha \times (\mathbb{C}^r, i\text{Id})$ which are \mathbb{C} -linear isomorphisms ($\Phi_\alpha \circ J = i\Phi_\alpha$), or equivalently of which transition maps take values in the endomorphisms of \mathbb{C}^r : $\phi_{\alpha\beta} = \Phi_\beta \circ \Phi_\alpha^{-1}: U_\alpha \cap U_\beta \rightarrow GL(\mathbb{C}^r)$. Therefore E is a complex vector bundle.

Remark 5.2 Let us set $\hat{\mathbb{C}} = \mathbb{R}[J]$, then $\hat{\mathbb{C}} = \mathbb{R}[J]$ is a vector bundle over M whose fibres are fields isomorphic to $\mathbb{C} = \mathbb{R}[i]$ and each fibre E_x of E is a $\hat{\mathbb{C}}_x$ -vector space. Then $E^{\mathbb{C}}$ is endowed with two different structures of vector bundle: one over the field \mathbb{C} (the tautological one defined by the complexification of E) and another one "over the distribution of field $\hat{\mathbb{C}}$ " (i.e. the one defined by J). In particular, we have two different complex structures in $E^{\mathbb{C}}$.

Now, let us suppose that E is endowed with a complex connection ∇ , i.e. a connection which commutes with J : $\nabla J = 0$. Then for all $X \in TM$, $\nabla_X: \mathcal{C}(E) \rightarrow \mathcal{C}(E)$ is \mathbb{C} -linear with respect to the complex vector space structure defined on $\mathcal{C}(E)$ by the complex vector bundle structure on E . Then we have two different ways to extend ∇ to $TM^{\mathbb{C}}$.

1. The canonical one: for any section $s \in \mathcal{C}(E^{\mathbb{C}})$, we extend ∇s by \mathbb{C} -linearity to a linear morphism from $TM^{\mathbb{C}}$ to $E^{\mathbb{C}}$,

$$\nabla_{iX}s = i\nabla_X s, \quad \forall X \in TM, s \in \mathcal{C}(E^{\mathbb{C}})$$

after, of course, having extended ∇ to a connection on $E^{\mathbb{C}}$ by setting $\nabla i s = i\nabla s, \forall s \in \mathcal{C}(E)$. In conclusion, $\forall s \in \mathcal{C}(E^{\mathbb{C}}), \nabla s \in \mathcal{C}(T^*M^{\mathbb{C}} \otimes E^{\mathbb{C}})$.

2. By using the complex vector bundle structure of E defined by J : for any $s \in \mathcal{C}(E)$, we extend ∇s by \mathbb{C} -linearity to a linear morphism from $TM^{\mathbb{C}}$ to E :

$$\hat{\nabla}_{iX}s = J\hat{\nabla}_X s, \quad \forall X \in TM, s \in \mathcal{C}(E).$$

Let us remark that $\widehat{\nabla}$ depends on J , and since we use the complex vector space structure defined by J , one needs that ∇ and J commute. On the other side the simple canonical complex extension defined in 1 (that we still denote by ∇) is independant of J and one needs not to do any additional hypothesis. Remark that the extension 1 is nothing but the extension $\widehat{\nabla}$ defined by the complex structure $i\text{Id}_{E^{\mathbb{C}}}$ on E (which commutes obviously with ∇).

Now let us suppose that M is an (almost) complex manifold with (almost) complex structure j_M . Then we have the splitting $TM^{\mathbb{C}} = T^{1,0}M \oplus T^{0,1}M$ defined by j_M which gives rise respectively to the following decompositions of ∇ and $\widehat{\nabla}$:

$$\begin{aligned}\widehat{\nabla} &= \nabla^{(1,0)} + \nabla^{(0,1)} \\ \nabla &= \nabla' + \nabla''.\end{aligned}$$

Definition 5.3 $\nabla^{(0,1)}$ is called the *Cauchy-Riemann operator* defined by ∇ and J .

More generally, let $\eta \in \mathcal{C}(T^*M \otimes E)$ be a 1-form on M with values in E . Then we can extend it in two different ways by \mathbb{C} -linearity in $TM^{\mathbb{C}}$ by setting:

$$\begin{aligned}\eta^{\mathbb{C}}(X + iY) &= \eta(X) + i\eta(Y), \quad \forall X, Y \in TM \\ \hat{\eta}(X + iY) &= \eta(X) + J\eta(Y), \quad \forall X, Y \in TM.\end{aligned}$$

Remark that $\eta^{\mathbb{C}} \in \mathcal{C}(T^*M^{\mathbb{C}} \otimes E^{\mathbb{C}})$ whereas $\hat{\eta} \in \mathcal{C}(T^*M^{\mathbb{C}} \otimes E)$. As above we can decompose $\eta^{\mathbb{C}}$ and $\hat{\eta}$ according to the decomposition $TM^{\mathbb{C}} = T^{1,0}M \oplus T^{0,1}M$:

$$\eta^{\mathbb{C}} = \eta' + \eta'' \tag{94}$$

$$\hat{\eta} = \eta^{(1,0)} + \eta^{(0,1)}. \tag{95}$$

Then we have the following relations

Lemma 5.1

$$\begin{aligned}[\eta']^{1,0} &= \eta^{(1,0)} - iJ\eta^{(1,0)} & [\eta'']^{0,1} &= \eta^{(1,0)} + iJ\eta^{(1,0)} \\ [\eta']^{0,1} &= \eta^{(0,1)} + iJ\eta^{(0,1)} & [\eta'']^{1,0} &= \eta^{(0,1)} - iJ\eta^{(0,1)}\end{aligned} \tag{96}$$

Proof. Let $Z = X - ij_M X \in T^{1,0}M$ with $X \in TM$. Then

$$\begin{aligned}[\eta(Z)]^{1,0} &= [\eta(X) - i\eta(j_M X)]^{1,0} = \eta(X) - iJ\eta(X) - i(\eta(j_M X) - iJ\eta(j_M X)) \\ &= \eta(X) - J\eta(j_M X) - iJ(\eta(X) - J\eta(j_M X)) \\ &= \eta^{(1,0)}(Z) - iJ\eta^{(1,0)}(Z).\end{aligned}$$

This gives us $[\eta']^{1,0}$. Then by applying this to $-J$, we obtain $[\eta']^{0,1}$. Finally, the second column of (96) is obtained by \mathbb{C} -conjugaison from the first column. This completes the proof. \square

We can apply what precedes to the flat differentiation d . Let (N, J) be an almost complex manifold and $s: M \rightarrow N$ a map. Then we consider the complex vector bundle $E = s^*TN$ over M . Then applying what precedes to the 1-form $\eta = ds$, we can consider the extensions \widehat{ds} and $(ds)^{\mathbb{C}}$, which then allows us to define the following extension of d to $TM^{\mathbb{C}}$:

$$\widehat{ds} = \widehat{ds} \quad \text{and} \quad d^{\mathbb{C}}s = (ds)^{\mathbb{C}},$$

and by abuse of notation⁵⁰ $d^{\mathbb{C}}$ will be still denoted by d . Then we can write the following decompositions

$$\hat{d} = \hat{\partial} + \overline{\hat{\partial}} \quad \text{and} \quad d = \partial + \overline{\partial}$$

according to the decomposition $TM^{\mathbb{C}} = T^{1,0}M \oplus T^{0,1}M$.

Definition 5.4 $\overline{\hat{\partial}}$ is called the Cauchy-Riemann derivative defined by J .

Now let us come back to the general situation of a complex vector bundle E over an almost complex manifold (M, j_M) , endowed with a complex connection ∇ . Let us set

$$H(M, E) = \{\eta \in T^*M \otimes E \mid \eta j_M = J\eta\}.$$

Then $H(M, E)$ is a vector subbundle of the vector bundle $T^*M \otimes E$ and is naturally endowed with the complex structure defined by

$$I(\eta) = \eta j_M = J\eta, \quad \forall \eta \in T^*M \otimes E, \quad (97)$$

which makes $H(M, E)$ being a complex vector bundle whose the set of sections is

$$\text{Hom}((TM, j_M), (E, J)) = \{\eta \in \mathcal{C}(T^*M \otimes E) \mid \eta \circ j_M = J \circ \eta\}.$$

The sections of $H(M, E)$ can also be characterized by using the splittings (94-95):

Lemma 5.2 We have the following equivalences for 1-forms $\eta \in \mathcal{C}(T^*M \otimes E)$:

$$\eta \circ j_M = J \circ \eta \iff \eta' \in T^*M \otimes E^{1,0} \iff \eta^{(0,1)} = 0.$$

Then we deduce in particular

Lemma 5.3 Let $s \in \mathcal{C}(E)$, then we have the following equivalences:

$$\nabla s \circ j_M = J \circ \nabla s \iff \nabla^{(0,1)} s = 0 \iff [\nabla' s]^{0,1} = 0 \iff \nabla'(s - iJs) = 0.$$

We will say that s is a vertically holomorphic section.

In fact we can say more

Lemma 5.4 Let us consider the splitting $TE = \mathcal{H} \oplus \mathcal{V}$ given by ∇ , where $\mathcal{V} = \ker \pi = \pi^*E$ is the vertical subbundle and \mathcal{H} the horizontal one. Then let us define an almost complex structure \check{J} on the manifold E by setting

$$\check{J} = ((d\pi)^* j_M)|_{\mathcal{H}} \oplus \pi^* J.$$

Then a section $s \in \mathcal{C}(E)$ is \check{J} -holomorphic if and only if it is vertically holomorphic.

Proof. It suffices to prove that any section $s \in \mathcal{C}(E)$ is horizontally holomorphic, i.e. satisfies the horizontal part of the equation $ds \circ j_M = J \circ ds$. We have $d\pi \circ (ds \circ j_M) = j_M$ since s is a section. In the other side we have $d\pi \circ (\check{J} \circ ds) = j_M \circ d\pi \circ ds = j_M$, by definition of \check{J} and using the fact s is a section. In conclusion $d\pi \circ (ds \circ j_M) = d\pi \circ (\check{J} \circ ds)$. This completes the proof. \square

In the following, we will say that a section of a complex vector bundle (E, J, ∇) is holomorphic if it is \check{J} -holomorphic.

⁵⁰and to be coherent with the notation used until now, in the paper.

Now, let us apply the two previous lemmas to the vector bundle $\mathbf{H}(M, E)$. First, let us endow M with an almost complex connection ∇^M (it means $\nabla^M J = 0$; such a connection always exists, see [40]). Then $T^*M \otimes E$ is naturally endowed with the connection $\overline{\nabla}^{\otimes}$ defined by ∇^M and ∇ . Further, we denote by $\overline{\nabla}$ the restriction to $\mathbf{H}(M, E)$ of $\overline{\nabla}^{\otimes}$. Then we remark that $\overline{\nabla}$ commutes with the complex structure I (defined by (97)). Therefore, we can now apply the two previous lemmas to the complex vector bundle $(\mathbf{H}(M, E), I, \overline{\nabla})$:

Proposition 5.4 *A section of $\mathbf{H}(M, E)$, $\eta \in \text{Hom}((TM, j_M), (E, J))$, is holomorphic if and only if it satisfies one of the following equivalent statements*

- (i) $\overline{\nabla}^{(0,1)} \eta = 0$
- (ii) $\overline{\nabla}'' \eta' = 0$
- (iii) $[\overline{\nabla}'' \eta]^{1,0} = 0$.

Moreover if M is a Riemann surface⁵¹, then it is also equivalent to

- (iv) $\overline{\partial}^{\widehat{\nabla}} \widehat{\eta} = \overline{\partial}^{\widehat{\nabla}} \eta^{(1,0)} = 0$, or
- (v) $\overline{\partial}^{\nabla} \eta' = 0$

Moreover, if M is a complex manifold (i.e. j_M is integrable) then we choose for ∇^M the unique torsion free complex connection on M . Then we obtain the following result:

Proposition 5.5 *Let $l \in TM$ be a complex line in the tangent bundle of the complex manifold M . Then for any section $\eta \in \text{Hom}((TM, j_M), (E, J))$ we have the following equality*

$$\overline{\nabla}^{(0,1)} \eta|_{l \times l} = d^{\nabla} \eta|_{l \times l}.$$

Moreover if η is holomorphic then $d^{\nabla} \eta = 0$. More particularly, if M is a Riemann surface then we have the following equivalence

$$\eta \text{ is holomorphic} \iff d^{\nabla} \eta = 0.$$

Remark 5.3 One could directly deduces the case of a Riemann surface by using proposition 5.4. Indeed, the first way to do that is to write $d^{\nabla} \eta = d^{\nabla} \eta' + d^{\nabla} \eta''$. Then remark that η' and η'' takes values in $E^{1,0}$ and $E^{0,1}$ respectively, according to lemma 5.3. Therefore since $E^{1,0}$ and $E^{0,1}$ are ∇ -parallel, we can say that $d^{\nabla} \eta'$ and $d^{\nabla} \eta''$ take values resp. in $E^{1,0}$ and $E^{0,1}$ resp., so that $d^{\nabla} \eta = 0 \iff d^{\nabla} \eta' = 0 \iff d^{\nabla} \eta'' = 0$. Then if M is a Riemann surface $d^{\nabla} \eta' = \overline{\partial}^{\nabla} \eta'$, and we conclude by using proposition 5.4.

The second way to do that is to use the $\widehat{\mathbb{C}}$ -linearity. Indeed, the extension to $TM^{\mathbb{C}}$ by $\widehat{\mathbb{C}}$ -linearity of $d^{\nabla} \eta$ is $\widehat{d^{\nabla} \eta} = d^{\widehat{\nabla}} \widehat{\eta} = d^{\widehat{\nabla}} \eta^{(1,0)} = 0$, since $\eta^{(0,1)} = 0$ (see lemma 5.3). Then if M is a Riemann surface $d^{\widehat{\nabla}} \eta^{(1,0)} = \overline{\partial}^{\widehat{\nabla}} \eta^{(1,0)}$, and we conclude by using proposition 5.4-(iv).

⁵¹and ∇^M the unique torsion free complex connection in M , which coincides also with the Levi-Civita connection of any Hermitian metric on M .

Remark 5.4 Let us consider a 1-form $\beta \in \mathcal{C}(T^*M \otimes E)$, then we can associate to it

$$\eta = \beta - J\beta \circ j_M = \hat{\beta} \circ (\text{Id} - ij_M) = \beta^{(1,0)} \circ (\text{Id} - ij_M).$$

By definition $\eta \in \mathcal{C}(\mathbb{H}(M, E))$, i.e. $\eta \circ j_M = J \circ \eta$. Moreover, still supposing that M is complex and that ∇^M is the unique torsion free complex connection on M , we have

$$\widehat{\nabla} \hat{\eta} = \widehat{\nabla} \beta^{(1,0)} \circ (\text{Id} - ij_M) \quad (98)$$

because $(\text{Id} - ij_M)$ is ∇^M -parallel.

Let us remark that since j_M and the multiplication by i coincide in $T^{1,0}M$, they define the same complex structure, which we will suppose $T^{1,0}M$ to be canonically endowed with. Then, since $\hat{\beta}$ is by definition a complex linear morphism from $TM^{\mathbb{C}}$ to E , $\beta^{(1,0)}$ is also a complex linear morphism from $T^{1,0}M$ to E . Hence $\beta^{(1,0)}$ is a section of the complex vector bundle $T_{1,0}^*M \otimes_{\mathbb{C}} E$. Therefore, from equation (98), we deduce that η is a holomorphic section of $\mathbb{H}(M, E)$ if and only if $\beta^{(1,0)}$ is a holomorphic section of $T_{1,0}^*M \otimes_{\mathbb{C}} E$. In particular if M is a Riemann surface, we deduce from (98), that

$$\overline{\partial} \hat{\eta} = \overline{\partial} \beta^{(1,0)} \circ (\text{Id} - ij_M).$$

Now, we come back to our complex vector bundle (E, J, ∇) and we recall a theorem ([39]) which characterizes when \check{J} is integrable.

Theorem 5.2 *Let $(E, J, \nabla) \rightarrow (M, j_M)$ be a complex vector over a complex manifold, with a complex connection ∇ . Then we will say that a holomorphic structure \mathcal{E} is compatible with ∇ (or that ∇ is adapted to \mathcal{E}) if it is induced by the almost complex structure \check{J} (defined by lemma 5.4). In other words, a section $s \in \mathcal{C}(E)$ is holomorphic with respect to \mathcal{E} if and only if*

$$\forall Z \in T^{1,0}M, \widehat{\nabla}_{\bar{Z}} s = 0.$$

An holomorphic structure \mathcal{E} exists on E if and only if \check{J} is integrable, and in this case \mathcal{E} is unique. Moreover \check{J} is integrable if and only if the $(0, 2)$ -component of the curvature operator⁵² R of ∇ vanishes.

When M is of dimension 2, then the $(0, 2)$ -component of the curvature operator always vanishes so that E always admits a holomorphic structure compatible with ∇ , that we will call, following [13], the Koszul-Malgrange holomorphic structure induced by ∇ . In the following, we suppose that a complex vector bundle (E, J, ∇) over a Riemann surface is always endowed with its Koszul-Malgrange holomorphic structure.

Interpretation of the holomorphic harmonicity in terms of holomorphic 1-forms.

Now we come back to the situation in the beginning of 5.1.2. More precisely, we consider (N, J) an almost complex manifold, with ∇ an almost complex connection, (L, j_L) a Riemann surface and $f: L \rightarrow N$ a map. Then we apply what precedes to the complex vector bundle $E = (f^*TN, f^*\nabla, f^*J)$ over L (i.e. L plays the role of M and f the one of s with respect to the notation of the previous paragraphs). We obtain a first theorem:

Proposition 5.6 *Let $f: (L, j_L) \rightarrow (N, J, \nabla)$ be a map from a Riemann surface into an almost complex affine manifold. Let us set*

$$\eta = df - Jdf \circ j_M.$$

⁵²i.e. the $(0, 2)$ -component of the extension \hat{R} of R to $\Lambda^2 T^*M^{\mathbb{C}}$ by $\hat{\mathbb{C}}$ -linearity.

Then η is a section of $\mathbb{H}(L, f^*TN)$, i.e. $\eta \circ j_M = J \circ \eta$. Moreover f is holomorphically harmonic if and only if η is a holomorphic section of the complex vector bundle $\mathbb{H}(L, f^*TN)$, i.e.

$$\bar{\partial}^\nabla \eta' = 0.$$

Proof. We write

$$d^\nabla \eta = d^\nabla (df - Jdf \circ j_M) = d^\nabla (df + *Jdf) = d^\nabla df + Jd^\nabla * df$$

so that we can conclude according to proposition 5.5 and proposition 5.1. This completes the proof. \square

Now, we can give a characterisation of holomorphic harmonicity which looks like very closely to the one which holds for harmonic maps ([13]):

Theorem 5.3 *A map $f: (L, j_L) \rightarrow (N, J, \nabla)$ from a Riemann surface into an almost complex affine manifold, is holomorphically harmonic if and only if*

$$\bar{\partial}^{\widehat{\nabla}} \hat{\partial} f = 0, \tag{99}$$

i.e. $\hat{\partial} f$ is a holomorphic section of $T_{1,0}^*L \otimes_{\mathbb{C}} f^*TN$.

Proof. Apply remark 5.4 to $\beta = df$ and then use proposition 5.6 to prove that $\hat{\partial} f$ is a holomorphic section and proposition 5.4-(iv) to prove the equation (99). \square

Remark 5.5 Let us derive (99) by a direct computation. Let be $Z = X + ij_L X \in T^{1,0}L$, then we have

$$\begin{aligned} (f^*\nabla)_{\bar{Z}}^{0,1} \hat{\partial} f(Z) &= (\nabla_{f(X)} + J\nabla_{f(j_L X)}) (df(X) - Jdf(j_L X)) \\ &= \nabla_{f(X)} df(X) + \nabla_{f(j_L X)} df(j_L X) + J (\nabla_{f(j_L X)} df(X) - \nabla_{f(X)} df(j_L X)) \\ &= d^\nabla * df(X, j_L X) - Jd^\nabla df(X, j_L X). \end{aligned}$$

5.2 The sigma model with a Wess-Zumino term in Nearly Kähler manifolds

Here we present an interpretation of the holomorphic harmonicity in terms of a sigma model with a Wess-Zumino term.

5.2.1 Totally skew-symmetric torsion

First, let us recall some useful properties about connections ⁵³.

Definition 5.5 *Two linear connection ∇ and ∇' in a manifold N are said to be geodesically equivalent if they have the same geodesics. A connection ∇ on a Riemannian manifold (N, h) is said to be geodesic-preserving if it is geodesically equivalent to the Levi-Civita connection ∇^h of h .*

⁵³See also [2] for a nice presentation about metric connections and their torsion

Proposition 5.7 *Let ∇ be a connection on a manifold N and $A \in \mathcal{C}(T^*N \otimes \text{End}(TN))$. Then the connection*

$$\nabla' = \nabla + A$$

has the same geodesic as ∇ if and only if $A(\cdot, \cdot)$ is skew-symmetric (as a bilinear map). In this case for any map $f: (M, g) \rightarrow N$, from a Riemannian manifold in to N , we have $\tau'_g(f) = \tau_g(f)$, where $\tau'_g(f)$ and $\tau_g(f)$ are the tension fields w.r.t. ∇ and ∇' respectively. Moreover (still in this case), we have

$$T^{\nabla'} = T^{\nabla} + 2A.$$

*Now, let us suppose that ∇ is metric w.r.t. some metric h in N . Then ∇' is metric if and only if A takes values (as a 1-form) in the skew-symmetric endomorphisms of TN : $A \in \mathcal{C}(T^*N \otimes \mathfrak{so}(TN))$. Therefore ∇' is metric and geodesically equivalent to ∇ if and only if A is totally skew-symmetric which means that the associated 3-linear map defined by $A^*(X, Y, Z) = \langle A(X, Y), Z \rangle$ is a 3-form on N .*

Now let us see how we can introduce the Levi-Civita connection starting from a given metric connection.

Proposition 5.8 *Let N be a manifold endowed with some connection that we denote by ∇^0 (for some reason that will appear clearly below). Let us set*

$$\nabla^t = \nabla^0 - tT^0, \quad 0 \leq t \leq 1,$$

where $T^0 = T^{\nabla^0}$ is the torsion of ∇^0 . Then we have

$$T^t := T^{\nabla^t} = -(2t - 1)T^0.$$

In particular, $\nabla^{\frac{1}{2}}$ is torsion free. Moreover all the connections ∇^t , $0 \leq t \leq 1$, are geodesically equivalent.

Now, let h be a metric on N which is ∇^0 -parallel. Then ∇^t , $t \neq 0$, is metric if and only if T^0 is totally skew-symmetric that is to say the 3-linear map defined by

$$(T^0)^*(X, Y, Z) := \langle T^0(X, Y), Z \rangle$$

is a 3-form. In this case, $\nabla^{\frac{1}{2}}$ coincides with the Levi-Civita connection ∇^h of h .

Remark 5.6 *We see that for a map $f: (M, g) \rightarrow N$, the strongly ∇^t -harmonicities are all equivalent for $t \neq \frac{1}{2}$.*

Conversely,

Proposition 5.9 *Let (N, h) be a Riemannian manifold, and let us denote by ∇^h its Levi-Civita connection. Then a metric connection ∇ on N is entirely determined by its torsion T . Moreover a metric connection ∇ on N is geodesic-preserving if and only if its torsion T is totally skew-symmetric. Then in this case we have*

$$\nabla = \nabla^h + \frac{1}{2}T.$$

Proof. For any metric connection $\nabla = \nabla^h + A$, we have

$$T(X, Y) = A(X, Y) - A(Y, X) \quad (100)$$

$$2A^*(X, Y, Z) = T^*(X, Y, Z) + T^*(Z, X, Y) + T^*(Z, Y, X), \quad (101)$$

which proves the first assertion. Concerning the second assertion, we see (according to (100-101)) that A is totally skew-symmetric if and only if T is so, i.e., according to proposition 5.7, ∇ is geodesic preserving if and only if T is totally skew-symmetric. Then in this case $T = 2A$ i.e. $\nabla = \nabla^h + \frac{1}{2}T$. This completes the proof. \square

Remark 5.7 The second equation (101) can be derived directly from the first one (100) (compute the right hand side of the second equation using the first equation which gives $2A^*(X, Y, Z)$). But there is another way (which will be useful in the following) to interpret this second equation. Indeed, first let us identify (via the metric h) in the following of this remark, each TN -valued bilinear form B on N with the corresponding trilinear form B^* . Then let us set $A := \frac{1}{2}(T + U) = \frac{1}{2}(T(X, Y, Z) + T(Z, X, Y) + T(Z, Y, X))$, where $U(X, Y, Z) = \langle U(X, Y), Z \rangle = T(Z, X, Y) + T(Z, Y, X)$. We remark that U is symmetric w.r.t. to the variables X, Y , so that the connection $\nabla - A = \nabla - \frac{1}{2}(T + U)$ is torsion free. Moreover we see that $A(X, Y, Z) = \frac{1}{2}(T(X, Y, Z) + T(Z, X, Y) + T(Z, Y, X))$ is skew symmetric w.r.t. the two last variables Y, Z . Therefore $\nabla - A$ is metric and thus this is the Levi-Civita connection ∇^h :

$$\nabla^h = \nabla - \frac{1}{2}(T + U).$$

Moreover, T is totally skew-symmetric if and only if the "natural reductivity term" $U = 0$. Furthermore, let us remark that the bijective correspondence between T and A comes simply from the isomorphism of vector bundle $T \in \Lambda^2 T^*N \otimes TN \mapsto T + U \in T^*N \otimes \mathfrak{so}(TN)$.

5.2.2 The general case of an almost Hermitian manifold

Let (E, J) be a complex vector space and let us set

$$\text{Bil}(E) = E^* \otimes E^* \otimes E \quad \text{and} \quad \mathcal{T}(E) = (\Lambda^2 E^*) \otimes E \subset \text{Bil}(E).$$

and for $\varepsilon, \varepsilon' \in \mathbb{Z}_2$ we set

$$\text{Bil}^{\varepsilon, \varepsilon'}(E, J) = \{A \in \text{Bil}(E) \mid A(J \cdot, \cdot) = \varepsilon JA, A(\cdot, J \cdot) = \varepsilon' JA\}$$

so that we have the decomposition

$$\text{Bil}(E) = \bigoplus_{(\varepsilon, \varepsilon') \in \mathbb{Z}_2 \times \mathbb{Z}_2} \text{Bil}^{\varepsilon, \varepsilon'}(E, J). \quad (102)$$

Let us remark that for any $A \in \text{Bil}(E)$, its component $A^{\varepsilon, \varepsilon'} \in \text{Bil}^{\varepsilon, \varepsilon'}(E, J)$ is given by

$$A^{\varepsilon, \varepsilon'}(X, Y) = -\frac{1}{4}(\varepsilon \varepsilon' A(JX, JY) + \varepsilon JA(JX, Y) + \varepsilon' JA(X, JY) - A(X, Y)). \quad (103)$$

Moreover we also have the decomposition

$$\mathcal{T}(E) = \mathcal{T}^{2,0} \oplus \mathcal{T}^{0,2} \oplus \mathcal{T}^{1,1},$$

where $\mathcal{T}^{2,0} = (\Lambda^{2,0} E^{*\mathbb{C}}) \otimes_{\mathbb{C}} E = \text{Bil}^{++}(E, J) \cap \mathcal{T}(E)$, $\mathcal{T}^{0,2} = (\Lambda^{0,2} E^{*\mathbb{C}}) \otimes_{\mathbb{C}} E = \text{Bil}^{--}(E, J) \cap \mathcal{T}(E)$ and $\mathcal{T}^{1,1} = (\Lambda^{1,1} E^{*\mathbb{C}}) \otimes_{\mathbb{C}} E = (\text{Bil}^{+-} + \text{Bil}^{-+})(E, J) \cap \mathcal{T}(E)$.

Of course, these notation can be extended to the case (E, J) is a complex vector bundle. In particular, we will use these for the tangent bundle (TN, J) of an almost complex manifold, and will forget in this case the precision of the bundle in the notation and write for example simply \mathcal{T} and Bil .

Given an almost complex manifold (N, J) with a connection ∇ (that we do not suppose to be almost complex) then its torsion T satisfies $T \in \mathcal{T}$ and⁵⁴ we can decompose it following (102): $T = T^{++} + T^{--} + T^{-+} + T^{+-}$. Then since T is skew-symmetric, then so is T^{++} , T^{--} and $T^{+-} + T^{-+}$. In other words, we have $T^{++} = T^{2,0}$, $T^{--} = T^{0,2}$ and $T^{+-} + T^{-+} = T^{1,1}$. In particular we have $T^{+-}(X, Y) = -T^{-+}(Y, X)$. Now, let us see how this decomposition can have a geometric meaning.

Lemma 5.5 *Let (N, J, ∇) be an almost complex manifold with an almost complex connection ∇ . Then we have*

$$N_J = 4T^{--}$$

where N_J denotes the torsion of J i.e its Nijenhuis tensor.

Proof. According to [40], Chap. IX, Prop. 3.6, the torsion N_J of J can be expressed in terms of the torsion T of the almost complex connection ∇ :

$$-N_J(X, Y) = T(JX, JY) - JT(JX, Y) - JT(X, JY) - T(X, Y)$$

which gives us $N_J = 4T^{--}$. This completes the proof. \square

Proposition 5.10 *Let (N, J, ∇) be an almost complex manifold with an almost complex connection ∇ . Then the following statements are equivalent.*

- (i) J anticommutes with the torsion T of ∇ : $T(X, JY) = -JT(X, Y)$.
- (ii) $T = T^{--}$ i.e. $T \in \mathcal{T}^{2,0}$.
- (iii) $T = \frac{1}{4}N_J$.

Proof. (ii) \Leftrightarrow (iii) follows from the previous lemma. Now, we have obviously (ii) \Rightarrow (i). Conversely (i) implies that $T = T^{--} + T^{+-}$ but since T is skew-symmetric this implies $T^{+-} = 0$ and $T = T^{--}$. This completes the proof. \square

From now until the end of this section 5.2.2, we consider (N, J) an almost complex manifold with an almost complex linear connection ∇ and a ∇ -parallel Hermitian metric h . Therefore (N, J, h) is an almost Hermitian manifold with a Hermitian connection ∇ .

Proposition 5.11 *Let (N, J, h) be an almost Hermitian manifold with a Hermitian connection ∇ . Let us suppose that J anticommutes with the torsion T of ∇ . Let us suppose also that the torsion of ∇ is totally skew-symmetric i.e.*

$$T^*(X, Y, Z) = \langle T(X, Y), Z \rangle$$

is a 3-form. Lastly, we suppose that the torsion is ∇ -parallel, i.e. $\nabla T^* = 0$ which is equivalent to $\nabla T = 0$. Then the trilinear map

$$H(X, Y, Z) = -T^*(X, Y, JZ) = \langle JT(X, Y), Z \rangle$$

is 3-form and is closed $dH = 0$.

⁵⁴i.e. $T \in \mathcal{C}(T)$

Proof. Firstly, according to proposition 5.10, we have $T^*(JX, Y, Z) = T^*(X, JY, Z) = T^*(X, Y, JZ)$, which prove that H is a 3-form.

Let us compute the exterior differential of H in terms of the connection (with torsion) ∇ :

$$\begin{aligned} dH(X_0, X_1, X_2, X_3) &= \sum_{i=0}^3 (-1)^i \nabla_{\hat{X}_i}^0 H(X_0, \dots, \hat{X}_i, \dots, X_3) \\ &\quad - \sum_{0 \leq i < j \leq 3} (-1)^{i+j} H(T(X_i, X_j), X_0, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_3) \\ &= \mathcal{S}_{i,j,k} H(T(X_0, X_i), X_j, X_k) + H(T(X_i, X_j), X_0, X_k) \end{aligned}$$

where the last sum is on all the circular permutations of 1,2,3. Moreover, we have

$$\begin{aligned} H(T(X_0, X_i), X_j, X_k) &= -H(X_k, X_j, T(X_0, X_i)) \\ &= \langle T(X_k, X_j), JT(X_0, X_i) \rangle \\ &= \langle T(X_0, X_i), JT(X_j, X_k) \rangle \\ &= -H(X_0, X_i, T(X_j, X_k)) \\ &= -H(T(X_j, X_k), X_0, X_i) \end{aligned}$$

so that we can conclude that $dH = 0$. This completes the proof. \square

Theorem 5.4 *Let (N, J, h) be an almost Hermitian manifold with a Hermitian connection ∇ . Then, under the 3 hypothesis of the previous proposition (T anticommutes with J , is totally skew-symmetric and ∇ -parallel), the equation for holomorphically harmonic maps $f: L \rightarrow N$ is the equation of motion (i.e. the Euler-Lagrange equation) for the sigma model in N with the Wess-Zumino term defined by the closed 3-form H . The action functional is given by*

$$S(f) = E(f) + S^{WZ}(f) = \frac{1}{2} \int_L |df|^2 d\text{vol}_g + \int_B H,$$

where B is 3-submanifold (or indeed a 3-chain) in N whose boundary is $f(L)$.

Proof. Since $dH = 0$ we have

$$\delta S^{WZ} = \int_B L_{\delta f} H = \int_B d\iota_{\delta f} H = \int_{f(L)} \iota_{\delta f} H,$$

therefore the Euler-Lagrange equation is

$$-\tau_g(f) + JT_g(f) = 0$$

which is the equation for holomorphically harmonic maps w.r.t. ∇ , since ∇ is geodesic preserving (see propositions 5.7 and 5.9) (g being as always a Hermitian metric on L). This completes the proof. \square

5.2.3 The example of a 3-symmetric space

Let us suppose now that $N = G/G_0$ is a (locally) 3-symmetric space. We use the notations of subsection 2.1.2. In particular, N is endowed with its canonical almost complex structure \underline{J} defined by (26).

Proposition 5.12 *The canonical connection ∇^0 in N commutes with the canonical almost complex structure \underline{J}*

$$\nabla^0 \underline{J} = 0.$$

Moreover, \underline{J} anticommutes with the torsion T^0 of ∇^0 . Lastly, if N is Riemannian, then ∇^0 is metric and (N, J, h) is almost Hermitian for any G -invariant metric h .⁵⁵ Furthermore, the torsion of ∇^0 is totally skew-symmetric if and only if h is naturally reductive.

Now, we can conclude

Theorem 5.5 *Let $N = G/G_0$ be a (locally) 3-symmetric space, that we suppose to be Riemannian and naturally reductive, and endowed with its canonical almost complex structure \underline{J} and its canonical connection ∇^0 . Let h be a G -invariant naturally reductive metric on N . Then the equation for holomorphically harmonic maps $f: L \rightarrow N$ is the Euler-Lagrange equation for the sigma model in N with the Wess-Zumino term defined by the closed 3-form H defined by*

$$H(X, Y, Z) = \langle JT(X, Y), Z \rangle$$

where T is the torsion of ∇^0 .

Remark 5.8 The hypothesis of natural reductivity is always satisfied if we allow us to use pseudo-Riemannian metrics and if \mathfrak{g} is semi-simple: the metric defined by the Killing form is then naturally reductive. Moreover, let us remark that w.r.t. Riemannian metrics the natural reductivity is in fact an hypothesis of compactness: the Lie subgroup of $GL(\mathfrak{m})$ generated by $\{[\text{ad}_{\mathfrak{m}}(X)]_{\mathfrak{m}}, X \in \mathfrak{m}\}$ must be compact.

5.2.4 The good geometric context/setting

In the previous variational interpretation given by theorem 5.4, we need to make 3 hypothesis on the torsion of the almost Hermitian connection: T anticommutes with J , is totally skew-symmetric and ∇ -parallel. Here, we want to understand what do these hypothesis mean geometrically and what is the good geometric context in which these take place. It will turn out that the good geometric context is the one of *Nearly Kähler manifold*.

Definition 5.6 *An almost Hermitian manifold (N, h, J) is called nearly Kähler if*

$$(\nabla_X^h J)X = 0,$$

where ∇^h is the Levi-Civita connection of h .

We can deduce immediately the following properties.

Proposition 5.13 *Let (N, h, J) be an almost Hermitian manifold. Let us consider its canonical Hermitian connection*

$$\nabla^0 := \nabla^h - \frac{1}{2}J\nabla^h J,$$

the torsion of which is denoted by T^0 . Then the following statements are equivalent:

(i) $T^0(\cdot, J\cdot) = -JT^0(\cdot, \cdot)$ and T^0 is totally skew-symmetric.

⁵⁵chosen according to our convention explained in subsection 2.1.2: that is $\tau_{\mathfrak{m}}$ leaves invariant the inner product defining h .

(ii) $T^0 = -J\nabla^h J$.

(iii) $\frac{1}{2}J\nabla^h J(\cdot, \cdot)$ is skew-symmetric.

(iv) (N, h, J) is nearly Kähler.

(v) $\nabla_{JX}^h J = -J\nabla_X J$ and T^0 is totally skew-symmetric.

Proof. First, we see that the implications (ii) \Rightarrow (i) and (ii) \Rightarrow (iii) \Leftrightarrow (iv) are obvious. Then by definition of ∇^0 we have

$$T^0(X, Y) = -\frac{1}{2}(J(\nabla_X^h J)Y - J(\nabla_Y^h J)X) \quad (104)$$

which gives us the implication (iii) \Rightarrow (ii). Furthermore, according to proposition 5.8, if T^0 is totally skew-symmetric then we have $\nabla^h = \nabla^0 - \frac{1}{2}T^0$ which provides the implication (i) \Rightarrow (ii). Finally the equivalence (i) \Leftrightarrow (v) follows directly from (104). This completes the proof. \square

In particular, a nearly Kähler manifold endowed with its canonical Hermitian connection satisfies 2 of our 3 hypothesis on the torsion (T^0 anticommutes with J and is totally skew-symmetric). Conversely, we have

Theorem 5.6 *Let (N, h, J, ∇^0) be an almost Hermitian manifold with an almost Hermitian connection ∇ . If the torsion T^0 of ∇^0 anticommutes with J and is totally skew-symmetric then (N, J, h) is nearly Kähler. Moreover, in this case, ∇^0 is the canonical Hermitian connection. Therefore the injective map*

$$(h, J) \mapsto (h, J, \nabla^h - \frac{1}{2}J\nabla^h J)$$

is in fact a bijection from the set of nearly Kähler structures on N into the set of almost Hermitian structures, (h, J, ∇^0) , with an almost Hermitian connection whose the torsion is totally skew-symmetric and anticommutes with J .

Remark 5.9 In other words, in an almost Hermitian manifold there exists at most only one Hermitian connection with totally skew-symmetric and J -anticommuting torsion, and if this connection exists then it coincides with the canonical Hermitian connection and the almost Hermitian manifold is nearly Kähler.

Moreover, the third hypothesis (the torsion is parallel) is implied by the first two.

Proposition 5.14 [Kirichenko], [38, 3] *If (N, h, J) is nearly Kähler then the canonical Hermitian connection has a parallel torsion: $\nabla^0 T^0 = 0$.*

Now, we can reformulate our theorem 5.4 by using the right geometric context:

Theorem 5.7 *Let (N, h, J) be a nearly Kähler manifold then the equation of holomorphic harmonicity for maps $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the 3-form:*

$$H = \frac{1}{3}d\Omega_J$$

where $\Omega_J = \langle J, \cdot, \cdot \rangle$ is the Kähler form.

Proof of theorem 5.6. If the torsion of ∇^0 is totally skew-symmetric then we have $\nabla^h = \nabla^0 - \frac{1}{2}T^0$, so that if moreover T^0 anticommutes with J then $-\frac{1}{2}T^0$ is the J -anticommuting part⁵⁶ of ∇^h i.e. $-\frac{1}{2}T^0 = \frac{1}{2}J\nabla^h J$ and ∇^0 is the canonical Hermitian connection. Therefore, we can apply proposition 5.13 which allows us to conclude. This completes the proof. \square

Proof of theorem 5.7 With the notation of proposition 5.11 we have $H(X, Y, Z) = \langle JT^0(X, Y), Z \rangle = \langle (\nabla^h J)(X, Y), Z \rangle = \nabla^h \Omega_J(X, Y, Z)$ according to proposition 5.13, and since $\nabla^h \Omega_J$ is a 3-form $((N, J, h)$ is nearly Kähler), we have $d\Omega_J = 3\nabla^h \Omega_J$. This completes the proof. \square

Remark 5.10 With this new setting, the closeness of $H = \frac{1}{3}d\Omega_J$ is obvious.

Return to the example of a 3-symmetric space According to proposition 5.12, a Riemannian (locally) 3-symmetric space $N = G/G_0$ is nearly Kähler if and only if it is naturally reductive. Then, we can reformulate the theorem 5.5 as follows:

Theorem 5.8 *Let $N = G/G_0$ be a (locally) 3-symmetric space, that we suppose to be Riemannian and naturally reductive, and endowed with its canonical almost complex structure \underline{J} and its canonical connection ∇^0 . Let h be a G -invariant naturally reductive metric on N . Then the equation for holomorphically harmonic maps $f: L \rightarrow N$ is the Euler-Lagrange equation for the sigma model in N with the Wess-Zumino term defined by the closed 3-form $H = \frac{1}{3}d\Omega_{\underline{J}}$.*

5.2.5 J -twisted harmonic maps

Definition 5.7 *Let $f: (M, g) \rightarrow N$ be a map from a Riemannian manifold (M, g) to a manifold N . Let us suppose that the vector bundle f^*TN is naturally endowed with some connection $\overline{\nabla}$. Then we will say that f is roughly harmonic w.r.t. $\overline{\nabla}$ (or $\overline{\nabla}$ -roughly harmonic) if*

$$\text{Tr}_g(\overline{\nabla}df) = 0.$$

This definition is useful in the case there exists a natural mapping which associates to each map $f: (M, g) \rightarrow N$ a connection in the vector bundle f^*TN . For example, we have the following.

Theorem 5.9 *A map $f: (L, j_L) \rightarrow (N, J, \nabla)$ from a Riemann surface into an almost complex manifold with a connection ∇ is holomorphically harmonic if and only if it is roughly harmonic w.r.t.*

$$\overline{\nabla} = f^*\nabla + \frac{1}{2}JT(df \circ j_L, \cdot).$$

Definition 5.8 *Let (N, J) be an almost complex manifold with an arbitrary connection ∇ . Then let us decompose it (in an unique way) as the sum of a J -commuting and respectively J -anticommuting part: $\nabla = \nabla^0 + A$, where $\nabla^0 J = 0$, and $A \in \mathcal{C}(T^*N \otimes \text{End}(TN))$, $AJ = -JA$, i.e. $A = \frac{1}{2}J\nabla J$. Then to any map $f: L \rightarrow N$ let us associate the connection*

$$\overline{\nabla} = f^*\nabla^0 - JA \circ j_L.$$

We will say that $f: L \rightarrow N$ is J -twisted harmonic w.r.t. ∇ if f is roughly harmonic w.r.t. $\overline{\nabla}$.

Now, we can conclude by the following interpretation of holomorphic harmonicity.

Theorem 5.10 *Let (N, J, h) be a Nearly Kähler manifold. Then a map $f: L \rightarrow N$ is holomorphically harmonic w.r.t. the canonical Hermitian connection ∇^0 if and only if it is J -twisted harmonic w.r.t. ∇^h .*

⁵⁶In the sense of theorem 4.17.

5.3 The sigma model with a Wess-Zumino term in \mathcal{G}_1 -manifolds

5.3.1 TN -valued 2-forms

Let (N, J, h) be an almost Hermitian manifold. In all the section 5.3, each TN -valued 2-form on N , $B \in \mathcal{C}(\mathcal{T})$, will be identified (via the metric h) with the corresponding trilinear form, skew-symmetric w.r.t. to the 2 first arguments:

$$B(X, Y, Z) := \langle B(X, Y), Z \rangle.$$

In particular, the left multiplication by J on $\mathcal{C}(\mathcal{T})$ defined a multiplication on the set of corresponding trilinear forms $(JB)(X, Y, Z) = \langle JB(X, Y), Z \rangle$. Moreover, under this identification, the space $\Omega^3(N) := \mathcal{C}(\Lambda^3 T^*N)$ of 3-forms will be considered as a subspace of $\mathcal{C}(\mathcal{T})$. We denote by Skew the following surjective linear map from $\mathcal{C}(\mathcal{T})$ onto $\Omega^3(N)$:

$$\text{Skew}(B)(X, Y, Z) = B(X, Y, Z) + B(Y, Z, X) + B(Z, X, Y).$$

Let us remark that $\frac{1}{3}\text{Skew}(B)$ is the skew-symmetric part of the trilinear form B and $\frac{1}{3}\text{Skew}: \mathcal{C}(\mathcal{T}) \rightarrow \Omega^3(N)$ is a projector (called the *Bianchi projector* in [18]). To any trilinear form $\alpha \in \mathcal{C}(\otimes^3 T^*N)$ will be associated its J -twisted trilinear form

$$\alpha^c = -\alpha(J\cdot, J\cdot, J\cdot).$$

In particular, if $\alpha = d\beta$, with $\beta \in \Omega^2(N) := \mathcal{C}(\Lambda^2 T^*N)$ then we set $d^c\beta := \alpha^c$.

We will also use the following action of the complex structure J on $\mathcal{C}(\mathcal{T})$: for any $B \in \mathcal{C}(\mathcal{T})$

$$J \cdot B := -JB(J\cdot, J\cdot) = J(B^{++} + B^{--}) - J(B^{+-} + B^{-+}) \quad (105)$$

i.e. in term of trilinear forms

$$J \cdot B = B(J\cdot, J\cdot, J\cdot) = -B^c.$$

Let us remark that $J \cdot (J \cdot B) = -B$.

Furthermore, let (L, j_L) be a Riemann surface and $B \in \mathcal{C}(\mathcal{T})$, then for any map $f: L \rightarrow N$ and any Hermitian metric g on (L, j_L) , we set

$$B_g(f) = *f^*B = B(f_*TL). \quad (106)$$

We will use a second natural action of J on $\mathcal{C}(\mathcal{T})$ defined by: for any $B \in \mathcal{C}(\mathcal{T})$,

$$J \circlearrowleft B = B(J\cdot, \cdot, \cdot) + B(\cdot, J\cdot, \cdot) + B(\cdot, \cdot, J\cdot)$$

and in terms of the components $B^{\varepsilon, \varepsilon'}$:

$$J \circlearrowleft B = JB^{++} - 3JB^{--} - J(B^{+-} + B^{-+}).$$

Moreover, by the aid of the two previous natural action, we can define a third action that will turn out to be the relevant one in the interpretation of the maximal odd determined system: for any $B \in \mathcal{C}(\mathcal{T})$,

$$J \star B = \frac{1}{2}(B(J\cdot, J\cdot, J\cdot) + B(J\cdot, \cdot, \cdot) + B(\cdot, J\cdot, \cdot) + B(\cdot, \cdot, J\cdot)) = \frac{1}{2}(J \cdot B + J \circlearrowleft B),$$

and in terms of the components $B^{\varepsilon, \varepsilon'}$:

$$J \star B = JB^{++} - JB^{--} - J(B^{+-} + B^{-+}).$$

Remark 5.11 Let us remark that all the three previous actions are independent of the metric h , as we can see it from the expressions in terms of the components $B^{\varepsilon, \varepsilon'}$, or more simply by writing their definitions using TN -valued 2-forms like in (105): $J \circlearrowright B = B(J\cdot, \cdot) + B(\cdot, J\cdot) - JB(\cdot, \cdot)$ and

$$J \star B = -\frac{1}{2}J(B(J\cdot, J\cdot) + JB(J\cdot, \cdot) + JB(\cdot, J\cdot) + B(\cdot, \cdot)).$$

We remark that this last formula - up to the factor $-\frac{1}{2}J$ and to the signs - makes $J \star B$ look like to some kind of torsion tensor of J w.r.t. B (cf. the definition of N_J). In particular, these three actions are defined in a general almost complex manifold (N, J) .

Remark 5.12 The equations (103) can be rewritten using the metric h as follows

$$B^{\varepsilon, \varepsilon'} = -\frac{1}{4}(\varepsilon \varepsilon' B(J\cdot, J\cdot, \cdot) - \varepsilon B(J\cdot, \cdot, J\cdot) - \varepsilon' B(\cdot, J\cdot, J\cdot) - B(\cdot, \cdot, \cdot)). \quad (107)$$

This leads us to define the following action

$$J \circlearrowright_2 B = B(J\cdot, J\cdot, \cdot) + B(J\cdot, \cdot, J\cdot) + B(\cdot, J\cdot, J\cdot) = B - 4B^{--}.$$

It is also important to remark that $J \cdot B - J \circlearrowright B = 4JB^{--}$ and $J \cdot B - J \star B = 2JB^{--}$, so that

Proposition 5.15 *Let T be the torsion of some Hermitian connection ∇ on (N, J, h) , then we have*

$$J \cdot T - J \circlearrowright T = JN_J \quad \text{and} \quad J \cdot T - J \star T = \frac{1}{2}JN_J.$$

5.3.2 Stringy Harmonic maps

We have seen two different ways to generalise the harmonicity to the case of affine target manifold. The first one is very natural and consist simply to write the harmonic map equation $\text{Tr}_g(\nabla df) = 0$ for a linear connection ∇ . The second one concerns holomorphically harmonic maps (from a Riemann surface into an almost complex manifold) and was dictated to us by the geometric equation of the second elliptic integrable system associated to a 3-symmetric space (section 2.3.2, paragraph: The model case). Furthermore, the preliminary study of the maximal determined system done in section 2.4, leads us to introduce the following generalisation of harmonic maps (which will turn out to be a generalisation of holomorphically harmonic maps for particular target spaces like nearly Kähler manifolds).

Definition 5.9 *Let (N, J) be an almost complex manifold with ∇ a linear connection then we will say that a map $f: L \rightarrow N$ from a Riemann surface into N is stringy harmonic if it is solution of the harmonic map equation with a JT -term:*

$$-\tau_g(f) + (J \cdot T)_g(f) = 0.$$

where g is a Hermitian metric on L .

We remark that if T anticommutes with J then stringy harmonicity coincides with holomorphic harmonicity (since in this case $J \cdot T = JT$). However, even though stringy harmonicity seems to be the more natural generalisation of (holomorphically) harmonic maps - in particular because of the property $J \cdot (J \cdot B) = -B$ which makes the first action look like very closely to the simple multiplication by J , which is not the case for the two other actions- it will turn out that the interpretation of the maximal odd determined system (see section 2.4) will use the action $J \star T$ of J on T . This leads us to the following modified definition.

Definition 5.10 Let (N, J) be an almost complex manifold with ∇ a linear connection then we will say that a map $f: L \rightarrow N$ from a Riemann surface into N is \star -stringy harmonic if it is solution of the modified stringy harmonic maps equation:

$$-\tau_g(f) + (J \star T)_g(f) = 0.$$

where g is a Hermitian metric on L .

We remark that if T anticommutes with J then \star -stringy harmonicity coincides with antiholomorphic harmonicity (since in this case $J \star T = -JT$).

Now, we will see that, under some hypothesis, the two previous definitions are in fact equivalent in the sense that there exists a new almost complex structure J^* such that $J \star T = J^* \cdot T$.

Proposition 5.16 Let (N, J) be an almost complex manifold. Let us suppose that there exists a J -invariant decomposition $TN = E^+ \oplus E^-$ and some $B \in \mathcal{C}(T)$ such that we have

$$\forall \alpha, \alpha' \in \mathbb{Z}_2, \quad B^{**}(E^\alpha, E^{\alpha'}) \subset E^{\alpha\alpha'} \quad \text{and} \quad B^{--}(E^\alpha, E^{\alpha'}) \subset E^{-\alpha\alpha'}. \quad (108)$$

where $B^{**} := B^{++} + B^{+-} + B^{-+} = B - B^{--}$. Let us define $J^* = J|_{E^+} \oplus -J|_{E^-}$, then we have

$$J \star B = J^* \cdot B.$$

Proof. Applying equation (103) to B^{**} , we obtain

$$B^{\varepsilon, \varepsilon'}(E^\alpha, E^{\alpha'}) \subset E^{\alpha\alpha'}, \quad \forall (\varepsilon, \varepsilon') \in \mathbb{Z}_2^2 \setminus \{(-, -)\}.$$

Moreover, we have

$$B = \sum_{\varepsilon, \varepsilon' \in \mathbb{Z}_2, \alpha, \alpha' \in \mathbb{Z}_2} B_{|E^\alpha \times E^{\alpha'}}^{\varepsilon, \varepsilon'}.$$

Besides, since $B_{|E^\alpha \times E^{\alpha'}}^{\varepsilon, \varepsilon'}(J^* \cdot, \cdot) = \varepsilon \alpha J B_{|E^\alpha \times E^{\alpha'}}^{\varepsilon, \varepsilon'} = \varepsilon \alpha (\alpha \alpha') J^* B_{|E^\alpha \times E^{\alpha'}}^{\varepsilon, \varepsilon'}$, it follows that $B_{|E^\alpha \times E^{\alpha'}}^{\varepsilon, \varepsilon'}$ is of type $(\varepsilon \alpha', \varepsilon' \alpha)$ w.r.t. to J^* , if $(\varepsilon, \varepsilon') \neq \{(-, -)\}$. And $B_{|E^\alpha \times E^{\alpha'}}^{--}$ is of type (α', α) . Therefore denoting by $\bar{B}^{\varepsilon, \varepsilon'}$ the $(\varepsilon, \varepsilon')$ -component of B w.r.t. J^* , we have

$$\bar{B}^{\varepsilon, \varepsilon'} = \sum_{(\alpha', \alpha) \neq -(\varepsilon, \varepsilon')} B_{|E^\alpha \times E^{\alpha'}}^{\varepsilon \alpha', \varepsilon' \alpha} + B_{|E^{\varepsilon'} \times E^\varepsilon}^{--}.$$

Finally, using this last equation we check by computation that $J^* \cdot B = J \star B$. This completes the proof. \square

Therefore, this yields the following corollary.

Corollary 5.1 Let (N, J) be an almost complex manifold with ∇ a linear connection. Let us suppose that there exists a J -invariant decomposition $TN = E^+ \oplus E^-$ such that the torsion T of ∇ satisfies the conditions (108). Let $J^* = J|_{E^+} \oplus -J|_{E^-}$. Then the \star -stringy harmonicity with respect to J is exactly the stringy harmonicity with respect to J^* .

Remark 5.13 Let us remark that proposition 5.16 is an algebraic identity, and holds in any complex vector space, or more generally any complex vector bundle (E, J) .

5.3.3 Almost Hermitian \mathcal{G}_1 -manifolds

In all this subsection, we consider (N, J, h) an almost Hermitian manifold with a Hermitian connection ∇ , whose the torsion is denoted by T .

We prove easily the following.

Proposition 5.17 *The components $B^{\varepsilon\varepsilon'}$ of an element $B \in \mathcal{C}(\mathcal{T})$, considered as trilinear forms, satisfy the following properties:*

$$\begin{aligned} B^{++} &\in \mathcal{C}((\Lambda^{2,0} \otimes \Lambda^{0,1}) \oplus (\Lambda^{0,2} \otimes \Lambda^{1,0})) \\ B^{+-} &\in \mathcal{C}((\Lambda^{1,0} \otimes \Lambda^{0,1} \otimes \Lambda^{0,1}) \oplus (\Lambda^{0,1} \otimes \Lambda^{1,0} \otimes \Lambda^{1,0})) \\ B^{-+} &\in \mathcal{C}((\Lambda^{1,0} \otimes \Lambda^{1,0} \otimes \Lambda^{0,1}) \oplus (\Lambda^{0,1} \otimes \Lambda^{0,1} \otimes \Lambda^{1,0})) \\ B^{--} &\in \mathcal{C}((\Lambda^{2,0} \otimes \Lambda^{1,0}) \oplus (\Lambda^{0,2} \otimes \Lambda^{0,1})), \end{aligned}$$

where $\Lambda^{p,q} = \Lambda^{p,q}T^*N$.

The results of this is:

Corollary 5.2 *Let $B \in \Omega^3(N) \subset \mathcal{C}(\mathcal{T})$ be a 3-form on N , then B^{--} is also a 3-form and is of type $(3,0) + (0,3)$. Moreover $B^{++} + B^{+-} + B^{-+}$ is a 3-form of type $(2,1) + (1,2)$ and we have the following relations:*

$$\begin{aligned} B^{++}(X, Y, Z) &= B^{+-}(Z, X, Y) \\ B^{-+}(X, Y, Z) &= B^{+-}(Y, Z, X) \end{aligned}$$

in other words $B^{**} := B^{++} + B^{+-} + B^{-+} = \text{Skew}(B^{\varepsilon\varepsilon'}), \forall(\varepsilon, \varepsilon') \in \mathbb{Z}_2 \setminus \{(-, -)\}$.

In particular, let us suppose that the torsion T of the Hermitian connection ∇ is totally skew-symmetric, then T^{--} is also a 3-form and is of type $(3,0) + (0,3)$, and T^{**} is a 3-form of type $(2,1) + (1,2)$. More particularly, the Nijenhuis tensor N_J is totally skew-symmetric.

Corollary 5.3 *Let us suppose that the torsion T of the Hermitian connection ∇ is totally skew-symmetric, then*

$$d\Omega_J = 3JT^{--} + J(T^{+-} + T^{-+} - T^{++}) = -J \circlearrowright T$$

i.e.

$$d\Omega_J = JN_J - J \cdot T.$$

Proof. Since T is skew-symmetric, we have $\nabla = \nabla^h + \frac{1}{2}T$, so that $\nabla J = 0$ implies

$$\nabla^h \Omega_J = -\frac{1}{2}(T(\cdot, J\cdot, \cdot) + T(\cdot, \cdot, J\cdot))$$

and therefore applying the operator Skew to that and using the fact that T is skew-symmetric we obtain

$$d\Omega_J = -J \circlearrowright T.$$

Then the last assertion follows from proposition 5.15. This completes the proof. \square

Now, we can conclude that

Theorem 5.11 *An almost Hermitian manifold (N, J, h) admits a Hermitian connection with totally skew-symmetric torsion if and only if the Nijenhuis tensor N_J is itself totally skew-symmetric. In this case, the connection is unique and determined by its torsion which is given by*

$$T = -d^c\Omega_J + N_J.$$

Proof. If such a Hermitian connection with skew-symmetric torsion exists, then according to corollary 5.2, N_J is itself skew-symmetric and moreover, according to corollary 5.3, we have $d\Omega_J = JN_J - J \cdot T = J \cdot (N_J - T)$ therefore $T = J \cdot d\Omega_J + N_J$. This proves the unicity.

Conversely, let us suppose that N_J is skew-symmetric and let ∇ be the metric connection defined by the torsion $T = J \cdot d\Omega_J + N_J$, i.e. $\nabla = \nabla^h - \frac{1}{2}T$. We have to check that $\nabla J = 0$. Let us recall ([40], proposition 4.2) that we have

$$2(\nabla_X^h \Omega_J)(Y, Z) = d\Omega_J(X, Y, Z) - d\Omega_J(X, JY, JZ) + N_J(Y, Z, JX). \quad (109)$$

Applying Skew to that, we obtain

$$\begin{aligned} 2d\Omega_J(X, Y, Z) &= 3d\Omega_J(X, Y, Z) - d\Omega_J(X, JY, JZ) - d\Omega_J(JX, Y, JZ) - d\Omega_J(JX, JY, Z) \\ &\quad + N_J(Y, Z, JX) + N_J(X, Y, JZ) + N_J(Z, X, JY) \end{aligned}$$

therefore

$$\begin{aligned} -4(d\Omega_J)^{0,2}(X, Y, Z) &= N_J(Y, Z, JX) + N_J(X, Y, JZ) + N_J(Z, X, JY) \\ &= 3N_J(X, Y, JZ) \text{ (since } N_J \text{ is skew-symmetric (and of type } (0, 2)\text{)).} \end{aligned} \quad (110)$$

Now, we can compute

$$\begin{aligned} \nabla \Omega_J &= \left(\nabla^h + \frac{1}{2}(J \cdot d\Omega_J + N_J) \right) \Omega_J \\ &= \nabla^h \Omega_J + \frac{1}{2}(N_J(X, JY, Z) + N_J(X, Y, JZ) - d\Omega_J(JX, Y, JZ) - d\Omega_J(JX, JY, Z)) \\ &= \frac{1}{2}(d\Omega_J(X, Y, Z) - d\Omega_J(X, Y, JZ) - d\Omega_J(JX, Y, JZ) - d\Omega_J(JX, JY, Z) \\ &\quad + N_J(Y, Z, JX) + N_J(X, JY, Z) + N_J(X, Y, JZ)) \\ &= \frac{1}{2}(3N_J(X, Y, JZ) + N_J(Y, Z, JX) + N_J(X, JY, Z) + N_J(X, Y, JZ)) = 0. \end{aligned}$$

This completes the proof. \square

Remark 5.14 This theorem can also be deduced from a more general result of Gauduchon [18], Proposition 2 (see also section 5.3.4 below). Moreover, it has already been stated by Friedrich-Ivanov [17] (but without writing completely the proof). To our knowledge a complete proof has never been given in the literature⁵⁷. In fact, Friedrich and Ivanov present a unified approach to construct G -connections, for any G -structure on some manifold, with skew-symmetric torsion. For example, they characterise the class of G_2 -structures and the class of contact metric structures for which such a connection exists, and prove the unicity of this connection.

Definition 5.11 *The unique Hermitian connection with skew-symmetric torsion is called the characteristic connection. According to the Gray-Hervella classification [19] of almost Hermitian manifolds, (N, J, h) admits a skew-symmetric Nijenhuis tensor if and only if it is of class $\mathcal{W}_1 \oplus \mathcal{W}_2 \oplus \mathcal{W}_3 =: \mathcal{G}_1$ (see [19]). These manifolds are called \mathcal{G}_1 -manifolds and according to the previous theorem they are exactly the almost Hermitian manifolds which admit a characteristic connection.*

⁵⁷See also [2] for the proof of the fact that the connection given in theorem 5.11 satisfies $\nabla J = 0$.

Proposition 5.18 *Let us suppose that the almost Hermitian manifold (N, J, h) is a \mathcal{G}_1 -manifold. Let us suppose that its characteristic connection ∇ has a parallel torsion $\nabla T = 0$. Then the 3-form*

$$H(X, Y, Z) = T(JX, JY, JZ) = \langle (J \cdot T)(X, Y), Z \rangle$$

is closed $dH = 0$.

Proof. Since, according to corollary 5.3, we have $H = -d\Omega_J + JN_J$, we only have to prove that the 3-form $JN_J = 4JT^{--}$ is closed. Moreover, since J is ∇ -parallel, so is the decomposition $\text{Bil}(E) = \oplus_{\mathbb{Z}_2 \times \mathbb{Z}_2} \text{Bil}^{\varepsilon, \varepsilon'}(E, J)$, so that if T is ∇ -parallel then so are its components $T^{\varepsilon\varepsilon'}$. Then the ∇ -parallel 3-forms, H , $d\Omega_J$ and JT^{--} satisfies the following formula for ∇ -parallel 3-forms α :

$$\begin{aligned} d\alpha(X_0, X_1, X_2, X_3) &= \sum_{i=0}^3 (-1)^i \nabla_{X_i} \alpha(X_0, \dots, \hat{X}_i, \dots, X_3) \\ &\quad - \sum_{0 \leq i < j \leq 3} (-1)^{i+j} \alpha(T(X_i, X_j), X_0, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_3) \\ &= \mathfrak{S}_{i,j,k} \alpha(T(X_0, X_i), X_j, X_k) + \alpha(T(X_i, X_j), X_0, X_k) \\ &= \mathfrak{S}_{i,j,k} \alpha(T(X_0, X_i), X_j, X_k) + \mathfrak{S}_{i,j,k} \alpha(T(X_j, X_k), X_0, X_i) \\ &= \mathfrak{S}_{X,Y,Z} \alpha(T(V, Z), X, Y) + \alpha(T(X, Y), V, Z) \end{aligned}$$

where we have set $(X, Y, V, Z) = (X_2, X_3, X_0, X_1)$. Then applying this formula to $d\Omega_J$, we obtain

$$\begin{aligned} 0 = -d(d\Omega_J) &= -d(3JT^{--} + J(T^{+-} + T^{-+} - T^{++})) \\ &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} \langle (3T^{--} - T^{++} + T^{1,1})(X, Y), J(T^{--} + T^{++} + T^{1,1})(V, Z) \rangle \end{aligned}$$

where $(X, Y) \rightleftharpoons (V, Z)$ means that we sum on the set $\{(X, Y, V, Z), (V, Z, X, Y)\}$. After a straightforward computation, we find

$$\begin{aligned} 0 = -d(d\Omega_J) &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} \{4\langle T^{--}(X, Y), JT^{++}(V, Z) \rangle \\ &\quad + 2\langle T^{--}(X, Y), JT^{1,1}(V, Z) \rangle \\ &\quad - 2\langle T^{++}(X, Y), JT^{1,1}(V, Z) \rangle\} \end{aligned} \quad (111)$$

Now, let us consider 4-linear forms on the variable $(X, Y, V, Z) \in TN^4$ and the associated decomposition $\otimes^4 T^*N^{\mathbb{C}} = \oplus_{\varepsilon \in (\mathbb{Z}_2)^4} \Lambda^{\varepsilon_1} \otimes \Lambda^{\varepsilon_2} \otimes \Lambda^{\varepsilon_3} \otimes \Lambda^{\varepsilon_4}$, where $\Lambda^+ = \Lambda^{1,0}T^*N$ and $\Lambda^- = \Lambda^{0,1}T^*N$. Then the term in the first line of (111) is in $(\otimes^4 \Lambda^+) \oplus (\otimes^4 \Lambda^-)$ whereas the terms in the second and third lines are in $(\otimes^{+++}) \oplus (\otimes^{----}) \oplus (\otimes^{++--}) \oplus (\otimes^{--+-})$, where $\otimes^{\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4} = \Lambda^{\varepsilon_1} \otimes \Lambda^{\varepsilon_2} \otimes \Lambda^{\varepsilon_3} \otimes \Lambda^{\varepsilon_4}$, $\forall \varepsilon \in (\mathbb{Z}_2)^4$. Hence the sums $\mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)}$ of these terms are respectively in $\Lambda^{4,0} \oplus \Lambda^{0,4}$ (first line) and in $\Lambda^{3,1} \oplus \Lambda^{1,3}$ (second and third lines). Therefore we obtain (in particular) that the first line vanishes

$$\mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4\langle T^{--}(X, Y), JT^{++}(V, Z) \rangle = 0.$$

Let us apply this to the computation of $d(JN_J)$:

$$\begin{aligned} d(JN_J) &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4 \langle JT^{--}(X,Y), T^{**}(V,Z) \rangle \\ &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4 \langle JT^{--}(X,Y), T^{1,1}(V,Z) \rangle. \end{aligned}$$

We see that $\langle JT^{--}(X,Y), T^{+-}(V,Z) \rangle$ is in $(\otimes^{+++}) \oplus (\otimes^{---+})$. But since JT^{--} is a 3-form, we have

$$\langle JT^{--}(X,Y), T^{+-}(V,Z) \rangle = -\langle JT^{--}(X, T^{+-}(V,Z)), Y \rangle$$

and this second 4-linear form (in the variable (X,Y,V,Z)) is in $(\otimes^{+++}) \oplus (\otimes^{---+})$, which imposes that $\langle JT^{--}(X,Y), T^{+-}(V,Z) \rangle = 0$, $\forall (X,Y,V,Z) \in TN^4$. We can prove the same result if we replace T^{+-} by T^{-+} . Therefore $d(JN_J) = 0$. This completes the proof. \square

Moreover, according to proposition 5.15, we deduce

Proposition 5.19 *Let us suppose that the almost Hermitian manifold (N, J, h) is a \mathcal{G}_1 -manifold. Let us suppose that its characteristic connection ∇ has a parallel torsion $\nabla T = 0$. Then the 3-form*

$$H^*(X, Y, Z) = \langle (J \star T)(X, Y), Z \rangle$$

is closed $dH^ = 0$.*

Now, we can conclude with the following variational interpretation of the stringy harmonicity.

Theorem 5.12 *Let us suppose that the almost Hermitian manifold (N, J, h) is a \mathcal{G}_1 -manifold. Let us suppose that its characteristic connection ∇ has a parallel torsion $\nabla T = 0$.*

• *Then the equation for stringy harmonic maps $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H = -d\Omega_J + JN_J.$$

• *Moreover the equation for \star -stringy harmonic maps $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H^* = -d\Omega_J + \frac{1}{2}JN_J.$$

Remark 5.15 We remark that the two previous sigma model differ by the Wess-Zumino term defined by the 3-form $\frac{1}{2}JN_J$.

5.3.4 Characterisation of Hermitian connections in terms of their torsion

In this subsection, we will give a result of Gauduchon [18] characterizing the Hermitian connections in terms of their torsion. We need to write with our notations and inside our setting and to write one proof in such a way that it will appear clearly that this result holds as well for Riemannian f -structure (see section 6) so that we will not have to reprove it (at least not entirely) in this more general context.

Theorem 5.13 *Let (N, J, h) be a Hermitian manifold. Then a metric connection ∇ is almost complex if and only if the following statements hold*

$$N_J = 4T^{0,2} \quad \text{and} \quad \text{Skew}(T^{2,0} - T^{1,1}) = (d^c \Omega_J)^{**}.$$

Proof. The metric connection ∇ can be written in the form $\nabla = \nabla^h - \frac{1}{2}(T + U)$, where $U(X, Y, Z) = T(Z, X, Y) + T(Z, Y, X)$. Then $\nabla\Omega_J = 0$ if and only if

$$\nabla^h J = -\frac{1}{2}[(T + U), J] = -((T + U)^{--}J + (T + U)^{+-}J) \quad (112)$$

but⁵⁸

$$U^{--}(X, Y, Z) = U(T^{--}) \quad \text{and} \quad U^{+-}(X, Y, Z) = T^{-+}(Z, X, Y) + T^{++}(Z, Y, X).$$

so that, according to (109)

$$\begin{aligned} (d\Omega_J)^{+-} = (\nabla^h\Omega_J)^{+-} &= -T^{+-}(X, JY, Z) - T^{-+}(Z, X, JY) - T^{++}(Z, JY, X) \\ &= -(\text{Skew}(T)^{+-}(X, JY, Z) - 2T^{++}(JY, Z, X)) \\ &= -J \cdot (\text{Skew}(T)^{+-}(X, Y, Z) - 2T^{++}(Y, Z, X)). \end{aligned}$$

Therefore applying Skew,

$$\begin{aligned} J \cdot (d\Omega_J)^{**} = (J \cdot d\Omega_J)^{**} &= \text{Skew}(T)^{**} - 2\text{Skew}(T^{++}(Y, Z, X)) \\ &= \text{Skew}(T^{**} - 2T^{++}) \\ &= \text{Skew}(-T^{2,0} + T^{1,1}) \end{aligned}$$

Besides, taking the $(\varepsilon, \varepsilon')$ -component of equation (112) for $(\varepsilon, \varepsilon') = (++)$, $(-+)$ instead of $(+-)$ would give the same result. Now, it remains to see what gives us the $(-, -)$ -component of this equation. Equations (109) and then (110) yield

$$\begin{aligned} 2(\nabla^h\Omega_J)^{0,2} &= 2(d\Omega_J)^{0,2} + N_J(Y, Z, JX) = -\frac{1}{2}(N_J(Y, Z, JX) + N_J(JX, Y, Z) + N_J(Z, JX, Y)) + N_J(Y, Z, JX) \\ &= -\frac{1}{2}(N_J(JX, Y, Z) + N_J(Z, JX, Y) + N_J(Z, Y, JX)) \end{aligned}$$

so that the $(0, 2)$ -component of equation (112) is written

$$-\frac{1}{4}(N_J(X, JY, Z) + N_J(Z, X, JY) + N_J(Z, JY, X)) = -(T^{--}(X, JY, Z) + T^{--}(Z, X, JY) + T^{--}(Z, JY, X)).$$

Using the fact that the map $B \in \mathcal{T} \mapsto B + U(B) \in T^*N \otimes \mathfrak{so}(TN)$ is bijective⁵⁹, we obtain

$$T^{--} = \frac{1}{4}N_J.$$

This completes the proof. □

5.3.5 The example of a naturally reductive homogeneous space

In this subsection we consider G/K a reductive homogeneous space and we denote by $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$ a reductive decomposition of the Lie algebra \mathfrak{g} .

⁵⁸with $U(B)(X, Y, Z) = B(Z, X, Y) + B(Z, Y, X)$, for any $B \in \mathcal{T}$.

⁵⁹See remark 5.7.

Theorem 5.14 *Let $N = G/K$ be a Riemannian naturally reductive homogeneous space. Then the canonical connection is a metric connection with skew-symmetric torsion (w.r.t. any naturally reductive G -invariant metric h). Let us suppose also that $N = G/K$ is endowed with some G -invariant complex structure J (i.e. \mathfrak{m} is endowed with some $\text{Ad}K$ -invariant complex structure J_0). If moreover one can choose a naturally reductive G -invariant metric h for which J is orthogonal⁶⁰, then (N, h, J) is an almost Hermitian \mathcal{G}_1 -manifold and its characteristic connection coincides with the canonical connection. Therefore, in this case its characteristic connection ∇ has a parallel torsion $\nabla T = 0$.*

Proof. The naturally reductivity means exactly that the torsion of the canonical connection is skew-symmetric. Then according to theorem 5.11, we deduce that (N, h, J) is \mathcal{G}_1 -manifold. This completes the proof. \square

Remark 5.16 In particular, we see that the Nijenhuis tensor is skew-symmetric. We can recover that by saying that since the G -invariant complex structure is parallel with respect to the canonical connection, then $N_J = 4T^{--}$ and moreover since T is a 3-form so is its component T^{--} .

By definition of (\star) -stringy harmonicity and the expression of the torsion of ∇^0 in terms of the Lie bracket, we have the following.

Proposition 5.20 *Let $N = G/K$ be a Riemannian homogeneous manifold endowed with a G -invariant complex structure J . Let $f: L \rightarrow N$ be a smooth map, $F: L \rightarrow G$ be a (local) lift of f and $\alpha = F^{-1}.dF$ the corresponding Maurer-Cartan form. Then in term of α , the equation of stringy harmonicity (w.r.t. ∇^0) is written*

$$d * \alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{m}}] - \frac{1}{2} J_0 [J_0 \alpha_{\mathfrak{m}} \wedge J_0 \alpha_{\mathfrak{m}}]_{\mathfrak{m}} = 0$$

whereas the equation of \star -stringy harmonicity is written:

$$d * \alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{m}}] + \frac{1}{2} [J_0 \alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} + \frac{1}{4} J_0 ([J_0 \alpha_{\mathfrak{m}} \wedge J_0 \alpha_{\mathfrak{m}}]_{\mathfrak{m}} + [\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}}) = 0$$

where J_0 is the complex structure on \mathfrak{m} corresponding to J .

5.3.6 Geometric interpretation of the maximal determined odd case.

In this subsection, we suppose that $N = G/K$ is a (locally) $(2k+1)$ -symmetric space, and we use the notations and the conventions of 2.1. In particular, we have defined the subspace $\mathfrak{m}_j \subset \mathfrak{m}$, for $1 \leq j \leq k$ (see subsection 2.1.2). We will set $\mathfrak{m}_{-j} = \mathfrak{m}_j$, $\mathfrak{m}_{j+2k+1} = \mathfrak{m}_j$, and $\mathfrak{m}_0 = \{0\}$, so that \mathfrak{m}_j is now defined for all $j \in \mathbb{Z}$.

Proposition 5.21 *Let us suppose that $N = G/K$ is a (locally) $(2k+1)$ -symmetric space endowed with its canonical almost complex structure \underline{J} and its canonical connection ∇^0 . Then the $(\varepsilon, \varepsilon')$ -*

⁶⁰which means that denoting by $G(\mathfrak{m})$, the compact subgroup in $GL(\mathfrak{m})$ generated by $\Lambda_{\mathfrak{m}}(\mathfrak{m}) := \{[\text{ad}_{\mathfrak{m}}(X)]_{\mathfrak{m}}, X \in \mathfrak{m}\} \subset \mathfrak{gl}(\mathfrak{m})$, and by $\langle G(\mathfrak{m}), J_0 \rangle$ the subgroup generated by $G(\mathfrak{m})$ and J_0 , then $\langle G(\mathfrak{m}), J_0 \rangle / G(\mathfrak{m})$ is compact.

component of the torsion T of ∇^0 are given by

$$\begin{aligned} T^{++}(\widetilde{X}, Y) &= - \sum_{\substack{i+j \leq k \\ 1 \leq i \leq j \leq k}} \left(1 - \frac{\delta_{ij}}{2}\right) ([X_{\mathfrak{m}_i}, Y_{\mathfrak{m}_j}]_{\mathfrak{m}_{i+j}} + [X_{\mathfrak{m}_j}, Y_{\mathfrak{m}_i}]_{\mathfrak{m}_{i+j}}) \\ T^{--}(\widetilde{X}, Y) &= - \sum_{\substack{i+j \geq k+1 \\ 1 \leq i \leq j \leq k}} \left(1 - \frac{\delta_{ij}}{2}\right) ([X_{\mathfrak{m}_i}, Y_{\mathfrak{m}_j}]_{\mathfrak{m}_{i+j}} + [X_{\mathfrak{m}_j}, Y_{\mathfrak{m}_i}]_{\mathfrak{m}_{i+j}}). \\ T^{+-}(\widetilde{X}, Y) &= - \sum_{1 \leq i \leq j \leq k} [X_{\mathfrak{m}_j}, Y_{\mathfrak{m}_i}]_{\mathfrak{m}_{j-i}} \\ T^{-+}(\widetilde{X}, Y) &= - \sum_{1 \leq i \leq j \leq k} [X_{\mathfrak{m}_i}, Y_{\mathfrak{m}_j}]_{\mathfrak{m}_{j-i}}. \end{aligned}$$

where $X, Y \in \mathcal{C}(TN)$ with lifts $X_{\mathfrak{m}}, Y_{\mathfrak{m}} \in \mathcal{C}^\infty(G, \mathfrak{m})$, and $T^{\varepsilon, \varepsilon'}(\widetilde{X}, Y) \in \mathcal{C}^\infty(G, \mathfrak{m})$ denotes the lift of $T^{\varepsilon, \varepsilon'}(X, Y) \in \mathcal{C}(TN)$. An other possibility is to consider that $X, Y \in TN$ are tangent vectors at some point $y \in N$ and that we have chosen $g \in G$ such that $g.G_0 = y$ and that we have set $X = \text{Adg}(X_{\mathfrak{m}})$, and $Y = \text{Adg}(Y_{\mathfrak{m}})$. Then the above equation, when written in the form $\widetilde{A} = B_{\mathfrak{m}}$, means in fact that we have $A = \text{Adg}(B_{\mathfrak{m}})$.

Proof. This follows from the fact that the lift in G of the torsion of ∇^0 is given by $\widetilde{T}(X_{\mathfrak{m}}, Y_{\mathfrak{m}}) = -[X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{m}}$, from the fact that the commutation relations $[\mathfrak{g}_i^{\mathbb{C}}, \mathfrak{g}_j^{\mathbb{C}}] \subset \mathfrak{g}_{i+j}^{\mathbb{C}}$ implies the following relations $[\mathfrak{m}_i, \mathfrak{m}_j] \subset \mathfrak{m}_{i+j} \oplus \mathfrak{m}_{i-j}$, and finally from the definition of \underline{J} . \square

Theorem 5.15 *Let us suppose that $N = G/K$ is a (locally) $(2k+1)$ -symmetric space endowed with its canonical almost complex structure \underline{J} and its canonical connection ∇^0 . Then the associated maximal determined system, $\text{Syst}(2k, \tau)$ is the equation of \star -stringy harmonicity for the geometric map $f: L \rightarrow N: (\nabla^0)^* df + (\underline{J} \star T^0)(f) = 0$.⁶¹*

Moreover, if we consider now that $N = G/K$ is endowed with the almost complex structure $\underline{J}^ := \bigoplus_{j=1}^k (-1)^j \underline{J}_{[\mathfrak{m}_j]}$, then this system is the equation of stringy harmonicity for the geometric map $f: L \rightarrow N: (\nabla^0)^* df + (\underline{J}^* \cdot T^0)(f) = 0$.*

Now, Suppose also that $N = G/K$ is naturally reductive⁶². Therefore, the previous system is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form

$$H^* = -d\Omega_{\underline{J}} + \frac{1}{2} \underline{J} N_{\underline{J}}.$$

Moreover, if $N = G/K$ is endowed with the almost complex structure \underline{J}^ , the previous system is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H = -d\Omega_{\underline{J}^*} + \underline{J}^* N_{\underline{J}^*}.$$

Proof. The first point follows from theorem 2.7, proposition 5.20, and proposition 5.21. Then the second point follows from corollary 5.1 and proposition 5.21. Let us precise that we apply corollary 5.1 with the J -invariant decomposition $TN = E^+ \oplus E^-$, defined by $E^+ = \bigoplus_{\substack{j=1 \\ j \text{ even}}}^k [\mathfrak{m}_j]$, and $E^- = \bigoplus_{\substack{j=1 \\ j \text{ odd}}}^k [\mathfrak{m}_j]$. Then according to proposition 5.21, this decomposition satisfies the

⁶¹Where we have removed the index "g" which precises that the previous terms are computed with respect to some Hermitian metric g on L .

⁶²And we choose a naturally reductive G -invariant metric h for which $\tau_{\mathfrak{m}}$ and thus \underline{J} are orthogonal. See the Appendix, lemma 8.2 for the existence of such an metric.

hypothesis of corollary 5.1. Finally the two last points (Wess-Zumino formulations) follow then from theorem 5.12 and 5.14. This completes the proof. \square

5.4 Stringy harmonicity and Holomorphic harmonicity

In this section, we will compare these two notions and prove that in the case of an almost complex affine manifold, these are equivalent. Indeed, the stringy harmonicity w.r.t. an almost complex connection ∇ is equivalent to the holomorphic harmonicity w.r.t. a new almost complex connection ∇^\bullet . In particular, according to subsection 5.1.2, stringy harmonicity has an interpretation in terms of holomorphic 1-forms.

Proposition 5.22 *Let (N, J, ∇) be an almost complex affine manifold (i.e. $\nabla J = 0$) and (L, j_L) a Riemann surface. Let us set*

$$\begin{aligned}\nabla^\bullet &= \nabla - (T^{2,0} + T^{0,2}) \\ \nabla^\star &= \nabla - T^{2,0}\end{aligned}$$

Then

- $f: L \rightarrow N$ is stringy harmonic w.r.t. ∇ if and only if f is anti-holomorphically harmonic w.r.t. to ∇^\bullet .
- f is \star -stringy harmonic w.r.t. ∇ if and only if f is anti-holomorphically harmonic w.r.t. to ∇^\star .

Proof. Since $T^{2,0} + T^{0,2} \in \mathcal{T}$ and $T^{2,0} \in \mathcal{T}$, then $\tau^{\nabla^\bullet}(f) = \tau^{\nabla^\star}(f) = \tau^\nabla(f)$, according to proposition 5.7. Moreover, we have

$$\begin{aligned}T^{\nabla^\star} &= T - 2T^{2,0} = -T^{2,0} + T^{0,2} + T^{1,1} \\ T^{\nabla^\bullet} &= T - 2(T^{2,0} + T^{0,2}) = -(T^{2,0} + T^{0,2}) + T^{1,1}\end{aligned}$$

so that

$$JT^{\nabla^\star} = -J \star T \quad \text{and} \quad JT^{\nabla^\bullet} = -J \cdot T,$$

where, of course, T , T^{∇^\bullet} and T^{∇^\star} are respectively the torsions of ∇ , ∇^\bullet and ∇^\star . We conclude by using proposition 5.1 and definitions 5.9 and 5.10. This completes the proof. \square

Now, since $\nabla^\star J = 0$, and according to theorem 5.3, we have

Corollary 5.4 *Let $f: (L, j_L) \rightarrow (N, J, \nabla)$ be a map from a Riemann surface into an almost complex affine manifold. Then f is \star -stringy harmonic if and only if*

$$\hat{\partial}^{\nabla^\star} \bar{\partial} f = 0,$$

i.e. $\bar{\partial} f$ is an anti-holomorphic section of $T_{1,0}^* L \otimes_{\mathbb{C}} f^* TN$, endowed with the holomorphic structure defined by ∇^\star .

Proof. This follows from proposition 5.22 above, and the theorem 5.3. \square

Remark 5.17 We can also check the previous corollary by direct computations as in remark 5.5.

Remark 5.18 Let us remark that, in general, the metricity of the connection is not preserved when one pass from ∇ to ∇^* (resp. ∇^\bullet). Indeed, if ∇ is metric then ∇^* is metric if and only if $T^{2,0}$ is a 3-form if and only if $T^{2,0} = T^{**} = 0$, and therefore $T = T^{0,2}$. Therefore, if $\nabla J = 0$, then (N, J, h) is nearly kähler.

Let us conclude this subsection by the following:

Proposition 5.23 *Let $f: (L, j_L) \rightarrow (N, J, \nabla)$ be a map from a Riemann surface into an almost complex manifold with a connection ∇ . Then*

- *f is stringy harmonic if and only if it is roughly harmonic w.r.t.*

$$\bar{\nabla} = f^* \nabla + \frac{1}{2}(J \cdot T)(df \circ j_L, \cdot).$$

- *And f is \star -stringy harmonic if and only if it is roughly harmonic w.r.t.*

$$\bar{\nabla}^* = f^* \nabla + \frac{1}{2}(J \star T)(df \circ j_L, \cdot).$$

6 Generalized Harmonic maps into f -manifolds.

6.1 f -structures: General definitions and properties.

6.1.1 f -structures, Nijenhuis tensor and natural action on the space of torsions \mathcal{T} .

Let us consider (N, F) a f -manifold, i.e. a manifold endowed with a f -structure (see definition 0.1). Let us set $\mathcal{H} = \text{Im } F$ and $\mathcal{V} = \ker F$, then we have $TN = \mathcal{H} \oplus \mathcal{V}$. If we put $P = -F^2$, then P is the projector on \mathcal{H} along \mathcal{V} . Moreover $PF = FP = F$ and $F^2P = -P$. In particular, $\bar{J} := F|_{\mathcal{H}}$ is a complex structure in the vector bundle \mathcal{H} .

Let us denote also by $q := \text{Id} - P$ the projector on \mathcal{V} along \mathcal{H} . We denote by $X = X^{\mathcal{V}} + X^{\mathcal{H}}$, or sometimes simply by $X = X^{\mathcal{V}} + X^{\mathcal{H}}$, the decomposition of any element $X \in TN$.

In all the section 6, we will consider the bundles \mathcal{H}^* and \mathcal{V}^* as well as all their tensor products respectively, as subbundles of T^*N and $\otimes^k T^*N$, $k \in \mathbb{N}^*$, respectively. For example, for any trilinear form $B \in \mathcal{C}(\otimes^3 T^*N)$, we will consider $B|_{\mathcal{H} \times \mathcal{V} \times \mathcal{H}}$ as an element of $\mathcal{C}(T^*N^3)$ by identifying it to $B(P \cdot, q \cdot, P \cdot)$.

Moreover, we will often identify a k -linear map with its expression in terms of the vectors $(X_1, \dots, X_k) \in TN^k$. For example, given $B \in \mathcal{C}(\otimes^3 T^*N)$, we will write "let $\beta \in \mathcal{H}^* \times \mathcal{H}^* \times \mathcal{V}^*$ be defined by $\beta = B(Z^{\mathcal{V}}, X^{\mathcal{H}}, Y^{\mathcal{H}})$ " instead of "let $\beta \in \mathcal{H}^* \times \mathcal{H}^* \times \mathcal{V}^*$ be defined by $\beta(X^{\mathcal{H}}, Y^{\mathcal{H}}, Z^{\mathcal{V}}) = B(Z^{\mathcal{V}}, X^{\mathcal{H}}, Y^{\mathcal{H}})$, for all $X, Y, Z \in TN$ ".

Notations We extend the notations and definitions of section 5.3 and the beginning of subsection 5.2.2, concerning there the complex bundle (TN, J) (defined by a complex manifold (N, J)) to the complex bundle (\mathcal{H}, \bar{J}) , defined in the present section by the f -manifold (N, F) . Then all the algebraic results of section 5.3 -like corollary 5.2 - can be extended to the complex bundle (\mathcal{H}, \bar{J}) .⁶³

Definition 6.1 The Nijenhuis tensor N_F of F is defined by

$$N_F(X, Y) = [FX, FY] - F[FX, Y] - F[X, FY] - P[X, Y],$$

where $X, Y \in \mathcal{C}(TN)$.

Then we obtain immediately ([31])

Proposition 6.1 We have the following identities.

$$\begin{aligned} N_F(qX, qY) &= -P[qX, qY] &= PN_F(qX, qY) \\ qN_F(X, Y) &= q[FX, FY] &= qN_F(pX, pY) \\ N_F(qX, PY) &= -F[qX, FY] - P[qX, PY] \end{aligned}$$

so that

$$N|_{\mathcal{V} \times \mathcal{V}} = R_{\mathcal{V}} \quad \text{and} \quad N^{\mathcal{V}} = -R_{\mathcal{H}}(F \cdot, F \cdot),$$

where $R_{\mathcal{V}}$ and $R_{\mathcal{H}}$ are the curvature of \mathcal{V} and \mathcal{H} respectively (in the sense of definition 4.4). In particular, $N^{\mathcal{V}}(\mathcal{V}, \mathcal{V}) = N^{\mathcal{V}}(\mathcal{H}, \mathcal{V}) = \{0\}$ i.e

$$N(\mathcal{H}, \mathcal{V}) \subset \mathcal{H} \quad \text{and} \quad N(\mathcal{V}, \mathcal{V}) \subset \mathcal{H}.$$

Moreover $N_F|_{\mathcal{V} \times \mathcal{H}} = N_F^{\mathcal{H}}|_{\mathcal{V} \times \mathcal{H}}$ satisfies the following property

$$N_F(X^{\mathcal{V}}, \bar{J}Y^{\mathcal{H}}) = -\bar{J}N_F(X^{\mathcal{V}}, Y^{\mathcal{H}})$$

i.e. $N_F(X^{\mathcal{V}}, \cdot)|_{\mathcal{H}}$ anticommutes with \bar{J} .

⁶³Or to the Hermitian bundle $(\mathcal{H}, \bar{J}, h|_{\mathcal{H}})$ if (N, F) is endowed with a (compatible) metric h ; see definition 6.6 below for a precise definition of a compatible metric.

Definition 6.2 Let (N, F) be a f -manifold. Then for any $B \in \mathcal{T}$, we set

$$B^{\varepsilon, \varepsilon'}(X, Y) = -\frac{1}{4}(\varepsilon \varepsilon' B(FX, FY) + \varepsilon FB(FX, Y) + \varepsilon' FB(X, FY) - B(X, Y)).$$

and $B^{2,0} := B^{++}$, $B^{1,1} := B^{+,-} + B^{-+}$ and $B^{0,2} := B^{--}$.

Then, setting $\bar{B} = B|_{\mathcal{H}^2}$, we have $PB^{\varepsilon, \varepsilon'} = \bar{B}^{\varepsilon, \varepsilon'}$ or in other words $B^{\varepsilon, \varepsilon'} = \bar{B}^{\varepsilon, \varepsilon'} - \frac{1}{4}(B^{\mathcal{V}}(F\cdot, F\cdot) - B^{\mathcal{V}})$, where the components $\bar{B}^{\varepsilon, \varepsilon'}$ are computed with respect to the complex structure \bar{J} on \mathcal{H} .

As for the case of an almost complex structure (section 5.3.1), we can define natural actions of F on elements $B \in \mathcal{T}$:

$$\begin{aligned} F \cdot B &:= B(F\cdot, F\cdot, F\cdot) := -B^c \\ F \odot B &:= B(F\cdot, \cdot) + B(\cdot, F\cdot) - FB(\cdot, \cdot) \\ F \bullet B &= F \cdot B + \frac{1}{2}F \odot(B - \bar{B}) \\ F \star B &= \frac{1}{2}(F \cdot B + F \odot B). \end{aligned}$$

It is then important to remark that $F \cdot B = \bar{J} \cdot \bar{B}$, so that

$$\begin{aligned} F \bullet B &= \bar{J} \cdot \bar{B} + \frac{1}{2}F \odot(B - \bar{B}) \\ F \star B &= \bar{J} \star \bar{B} + \frac{1}{2}F \odot(B - \bar{B}). \end{aligned}$$

Moreover, it is also useful to remark that $F \bullet B - F \odot B = 4\bar{J} \cdot \bar{B}^{--} - \frac{1}{2}F \odot(B - \bar{B})$, and $F \bullet B - F \star B = 2\bar{J} \cdot \bar{B}^{--}$.

6.1.2 Introducing a linear connection.

Now, we introduce a linear connection and want to compare the vertical component of the torsion with the vertical torsion.

We obtain immediately the two followings properties.

Proposition 6.2 Let (N, F, ∇) be an affine manifold endowed with a parallel f -structure ($\nabla F = 0$). Then the subbundles $\mathcal{H} = \text{Im } F$ and $\mathcal{V} = \ker F$ are ∇ -parallel.

Proposition 6.3 Let (N, ∇) be an affine manifold. Let us suppose that we have a ∇ -parallel splitting $TN = \mathcal{H} \oplus \mathcal{V}$, where \mathcal{H}, \mathcal{V} inherit the names of horizontal and vertical subbundles respectively. Then the vertical torsion coincides with the vertical component of the torsion:

$$T^v = T^{\mathcal{V}},$$

where we use notations of section 4.1.3 for T^v , and the notations defined above (in the beginning of 6.1.1) for $T^{\mathcal{V}}$.

In a more general context we can relate T^v and $T^{\mathcal{V}}$ as follows.

Proposition 6.4 *Let (N, ∇) be an affine manifold. Let us suppose that we have some splitting $TN = \mathcal{V} \oplus \mathcal{H}$, where \mathcal{H}, \mathcal{V} inherit the names of horizontal and vertical subbundles respectively. Then the vertical torsion and the vertical part of the torsion satisfy the following relations*

$$T^v_{|\mathcal{V} \times \mathcal{V}} = T^{\mathcal{V}}_{|\mathcal{V} \times \mathcal{V}} \quad \text{and} \quad T^v_{|\mathcal{H} \wedge \mathcal{V}} = T^{\mathcal{V}}_{|\mathcal{H} \wedge \mathcal{V}} + \sigma^v.$$

where σ^v is the restriction to $\mathcal{H} \wedge \mathcal{V}$ of the \mathcal{V} -valued 2-form $\nabla^v q(X, Y) - \nabla^v q(Y, X)$.

Definition 6.3 *The term $\mathcal{R} = T^v_{|\mathcal{H} \wedge \mathcal{V}}$ will be called the reductivity term.*

Proposition 6.5 *Furthermore, in the situation off proposition 6.4, we have the following equality:*

$$T^v = \Phi \oplus \mathcal{R} \oplus T^{\mathcal{V}}(q, q),$$

where $\Phi = R_{\mathcal{H}}$ is the curvature of \mathcal{H} .

Proof of proposition 6.4 and 6.4. This is a straightforward computation. \square

Let $X^h, Y^h \in \mathcal{C}(\mathcal{H})$. Then for any $f \in \mathcal{C}^\infty(N)$, we have $\nabla^v_{X^h}(fY^h) = f\nabla^v_{X^h}Y^h + (X^h \cdot f)(Y^h)^v = f\nabla^v_{X^h}Y^h$ so that $\nabla^v_{X^h}Y^h$ defines a bilinear map from $\mathcal{H} \times \mathcal{H}$ into \mathcal{V} . Let Ψ be its skew-symmetric part: $\Psi(H_1, H_2) = \nabla^v_{H_1}H_2 - \nabla^v_{H_2}H_1$. Then we have $T^{\mathcal{V}}(H_1, H_2) = \nabla^v_{H_1}H_2 - \nabla^v_{H_2}H_1 - [H_1, H_2]^v$ i.e.

$$T^{\mathcal{V}}_{|\mathcal{H} \times \mathcal{H}} = \Psi + \Phi.$$

Therefore,

Proposition 6.6 *The following relation holds*

$$T^{\mathcal{V}} = (\Psi + \Phi) \oplus (\mathcal{R} - \sigma^v) \oplus T^{\mathcal{V}}(q, q).$$

Therefore $T^{\mathcal{V}} = T^v$ if and only if $\Psi = 0$ and $\sigma^v = 0$, which happens in particular if \mathcal{H} is ∇ -parallel.

6.2 The f -connections and their torsion.

6.2.1 Definition, notations and first properties.

Let us come back to the case of a f -manifold (N, F) .

Definition 6.4 *A linear connection ∇ , on a f -manifold (N, F) , which preserves the f -structure, i.e $\nabla F = 0$, is called a f -connection.*

Then we obtain easily:

Proposition 6.7 *Let (N, F, ∇) be a f -manifold endowed with an affine f -connection. Then the torsion T satisfies the following identity*

$$T(FX, FY) - FT(FX, Y) - FT(X, FY) - PT(X, Y) = -N_F(X, Y)$$

Therefore, we deduce the following.

Corollary 6.1 *Setting $N_{\bar{J}} = N_F^{\mathcal{H}}|_{\mathcal{H} \times \mathcal{H}}$, the torsion T (of a f -connection ∇ on a f -manifolds (N, F)) satisfies the following identities:*

$$\begin{aligned} N_{\bar{J}} &= 4 (T_{\mathcal{H} \times \mathcal{H}}^{\mathcal{H}})_{\bar{J}}^{0,2} \\ T^{\mathcal{H}}(\bar{J}X^h, Y^v) - \bar{J}T^{\mathcal{H}}(X^h, Y^v) &= -\bar{J}N_F^{\mathcal{H}}(X^h, Y^v) = -([\bar{J}X^h, Y^v]^{\mathcal{H}} - \bar{J}[X^h, Y^v]^{\mathcal{H}}) \\ T^{\mathcal{H}}(X^v, \bar{J}Y^h) - \bar{J}T^{\mathcal{H}}(X^v, Y^h) &= -\bar{J}N_F^{\mathcal{H}}(X^v, Y^h) = -([X^v, \bar{J}Y^h]^{\mathcal{H}} - \bar{J}[X^v, Y^h]^{\mathcal{H}}) \\ T^{\mathcal{V}}(X^h, Y^h) &= -N_F^{\mathcal{V}}(FX^h, FY^h) = \Phi(X, Y) \\ T^{\mathcal{H}}(X^v, Y^v) &= N_F^{\mathcal{H}}(X^v, Y^v) = R_{\mathcal{V}}(X^v, Y^v) \end{aligned}$$

where $X, Y \in \mathcal{C}(TN)$. Consequently, the following component of the torsion $T_{|\mathcal{H} \times \mathcal{H}}^{\mathcal{V}}$, $T_{|\mathcal{V} \times \mathcal{V}}^{\mathcal{H}}$, $(T_{|\mathcal{H} \times \mathcal{H}}^{\mathcal{H}})^{0,2}$ and $[T_{|\mathcal{V} \times \mathcal{H}}^{\mathcal{H}}, \bar{J}]$ are independent of ∇ .

Introducing a metric After having introduced a metric h on N , we want to characterize the metric connections ∇ which preserves F . More precisely, we want to find a necessary and sufficient condition on the torsion T for ∇ metric to be a f -connection. In a first time, we will begin by characterizing the metric connections which preserve the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$, then in a second time we will introduce the additionnal condition that the induced connection on \mathcal{H} preserves the complex structure \bar{J} .

Let us define some notations. In the following, since a metric is given we use the convention defined in section 5.3.1: each TN -valued bilinear form on N , $B \in \mathcal{C}(T^*N \otimes T^*N \otimes TN)$, will be identified (via the metric h) with the corresponding trilinear form. Moreover, we denote by Ω_A the bilinear form associated (via the metric h) to an endomorphism $A \in \mathcal{C}(\text{End}(TN))$:

$$\Omega_A(X, Y) = \langle A(X), Y \rangle, \quad \forall X, Y \in TN.$$

Then, under our convention, for any endomorphism $A \in \mathcal{C}(TN)$, $\nabla^h A$ is identified to $\nabla^h \Omega_A$. Moreover, we set

$$\text{Sym}(B)(X, Y) = B(X, Y) + B(Y, X), \quad \forall X, Y \in TN,$$

for all $B \in \mathcal{C}(T^*N \otimes T^*N \otimes TN)$.

Furthermore, let E_1, E_2, E_3 be vector bundles over N , then we set also $\mathcal{S}(E_1 \times E_2 \times E_3) = \mathcal{S}_{i,j,k} E_i \otimes E_j \otimes E_k$, where we do a direct sum on the circular permutation of 1,2,3.

Finally, to avoid any risk of confusion of the index "h" denoting the metric in the notation of the Levi-Civita connection ∇^h , with the same index in the notation of the horizontal component X^h of a vector $X \in TN$, we will denote in all this section 6 the Levi-Civita connection by D :

$$D := \nabla^h.$$

6.2.2 Characterisation of metric connections preserving the splitting.

Theorem 6.1 *Let (N, h) be a Riemannian manifold with an orthogonal decomposition $TN = \mathcal{H} \oplus \mathcal{V}$. Then a metric connection ∇ leaves invariant this decomposition (i.e. \mathcal{H} and \mathcal{V} are ∇ -parallel) if and only if its torsion T satisfies*

$$T_{|\mathcal{H} \times \mathcal{H} \times \mathcal{V}} = \Phi, \quad T_{|\mathcal{V} \times \mathcal{V} \times \mathcal{H}} = R_{\mathcal{V}}$$

and

$$\begin{aligned}\mathrm{Sym}_{\mathcal{H}\times\mathcal{H}}(T_{|\mathcal{V}\times\mathcal{H}\times\mathcal{H}})(X^v, Y^h, Z^h) &= \mathrm{Sym}_{\mathcal{H}\times\mathcal{H}}\left(D\Omega_{q|\mathcal{H}\times\mathcal{H}\times\mathcal{V}}\right)(Y^h, Z^h, X^v) \\ \mathrm{Sym}_{\mathcal{V}\times\mathcal{V}}(T_{|\mathcal{H}\times\mathcal{V}\times\mathcal{V}})(X^h, Y^v, Z^v) &= \mathrm{Sym}_{\mathcal{V}\times\mathcal{V}}\left(D\Omega_{q|\mathcal{V}\times\mathcal{V}\times\mathcal{H}}\right)(Y^v, Z^v, X^h)\end{aligned}$$

In particular, the components $\mathrm{Sym}_{\mathcal{H}\times\mathcal{H}}(T_{|\mathcal{V}\times\mathcal{H}\times\mathcal{H}})$ and $\mathrm{Sym}_{\mathcal{V}\times\mathcal{V}}(T_{|\mathcal{H}\times\mathcal{V}\times\mathcal{V}})$ are independent of ∇ .

Proof. According to remark 5.7, we have $D = \nabla - A = \nabla - \frac{1}{2}(T + U)$. Therefore $\nabla q = 0$ if and only if

$$Dq = -[A, q]. \quad (113)$$

But $\langle [A, q](X, Y), Z \rangle = A(X, Y^v, Z) - A(X, Y, Z^v) = A(X^h, Y^v, Z^h) + A(X^v, Y^v, Z^h) - A(X^h, Y^h, Z^v) - A(X^v, Y^h, Z^v)$, and according to the characterization of the Levi-Civita connection:

$$\begin{aligned}-2\langle (D_{X^h}q)Y^h, Z^v \rangle &= 2\langle D_{X^h}Y^h, Z^v \rangle = -Z^v \cdot \langle X^h, Y^h \rangle + \langle [X^h, Y^h], Z^v \rangle + \langle [Z^v, X^h], Y^h \rangle + \langle X^h, [Z^v, Y^h] \rangle \\ &= -\Phi(X^h, Y^h, Z^v) + \langle -D_{Z^v}X^h + [Z^v, X^h], Y^h \rangle + \langle -D_{Z^v}Y^h + [Z^v, Y^h], X^h \rangle \\ &= -\Phi(X^h, Y^h, Z^v) - \langle D_{X^h}Z^v, Y^h \rangle - \langle D_{Y^h}Z^v, X^h \rangle \\ &= -\Phi(X^h, Y^h, Z^v) - D\Omega_q(X^h, Z^v, Y^h) - D\Omega_q(Y^h, Z^v, X^h) \\ &= -\Phi(X^h, Y^h, Z^v) - \mathrm{Sym}_{\mathcal{H}\times\mathcal{H}}(D\Omega_q(X^h, Y^h, Z^v)).\end{aligned}$$

In the last line we have used the fact that $D\Omega_q$ is symmetric w.r.t. the two last variables (since q is a symmetric projector). Therefore the condition (113) restricted to $\mathcal{H} \times \mathcal{H} \times \mathcal{V}$ is written $(T + U)|_{\mathcal{H}\times\mathcal{H}\times\mathcal{V}} = \Phi + \mathrm{Sym}_{\mathcal{H}\times\mathcal{H}}(D\Omega_{q|\mathcal{H}\times\mathcal{H}\times\mathcal{V}})$, that is to say by identifying respectively the symmetric and skew-symmetric part (w.r.t. the two first variables) of the two hand sides of this equality respectively, we obtain

$$\begin{aligned}T_{|\mathcal{H}\times\mathcal{H}\times\mathcal{V}} = \Phi \quad \text{and} \quad \mathrm{Sym}_{\mathcal{H}\times\mathcal{H}}(T_{|\mathcal{V}\times\mathcal{H}\times\mathcal{H}})(X^v, Y^h, Z^h) &=: U_{|\mathcal{H}\times\mathcal{H}\times\mathcal{V}}(Y^h, Z^h, X^v) \\ &= \mathrm{Sym}_{\mathcal{H}\times\mathcal{H}}\left(D\Omega_{q|\mathcal{H}\times\mathcal{H}\times\mathcal{V}}\right)(Y^h, Z^h, X^v).\end{aligned} \quad (114)$$

Moreover, since $D\Omega_q$ is symmetric w.r.t. the two last variables and A is skew-symmetric w.r.t. these two last variables, we see that the restriction to $\mathcal{H} \times \mathcal{H} \times \mathcal{V}$ and to $\mathcal{H} \times \mathcal{V} \times \mathcal{H}$ of the condition (113) are in fact equivalent. Furthermore, we have $\langle (D_{X^v}q)Y^h, Z^h \rangle = 0$ and $\langle [A, q](X^v, Y^h), Z^h \rangle = 0$. Therefore, the restriction to $\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})$ of the condition (113) is equivalent to (114).

Proceeding in the same way as above, we obtain that the restriction to $\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})$ of (113) is equivalent to

$$T_{|\mathcal{V}\times\mathcal{V}\times\mathcal{H}} = R_{\mathcal{V}} \quad \text{and} \quad \mathrm{Sym}_{\mathcal{V}\times\mathcal{V}}(T_{|\mathcal{H}\times\mathcal{V}\times\mathcal{V}})(X^h, Y^v, Z^v) = \mathrm{Sym}_{\mathcal{V}\times\mathcal{V}}\left(D\Omega_{q|\mathcal{H}\times\mathcal{V}\times\mathcal{V}}\right)(Y^v, Z^v, X^h).$$

Finally, we have $\langle (D_{X^h}q)Y^h, Z^h \rangle = 0 = \langle [A, q](X^h, Y^h), Z^h \rangle$ and $\langle (D_{X^v}q)Y^v, Z^v \rangle = 0 = \langle [A, q](X^v, Y^v), Z^v \rangle$. This completes the proof. \square

Skew-symmetric torsion. Now, let us see under which condition on the Riemannian manifold, there exists a connection preserving the splitting and with skew-symmetric torsion. It will turn out that the existence of a connection preserving the splitting and of which the horizontal component of the torsion $T_{|\mathcal{H}^3}$ is skew-symmetric does not imposes any condition on the Riemannian manifold (N, h) , but the skew-symmetry of the other components $T_{|\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ and $T_{|\mathcal{S}(\mathcal{V}\times\mathcal{V}\times\mathcal{H})}$ imposes constraints on the Riemannian manifold.

Corollary 6.2 *Let (N, h) be a Riemannian manifold with an orthogonal decomposition $TN = \mathcal{H} \oplus \mathcal{V}$. Then, the following statements are equivalent*

- (i) *There exists a metric connection ∇ leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$, such that the following component of the torsion $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ is skew-symmetric, i.e. $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = \text{Skew}(\Phi)$.*
- (ii) *For any metric connection ∇ leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$, the component of the torsion $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ is skew-symmetric.*
- (iii) *$D\Omega_{q|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}}}$ is skew-symmetric w.r.t. the two first variables, i.e. $Dq|_{\mathcal{H} \times \mathcal{H}} \in \mathcal{C}((\Lambda^2 \mathcal{H}^*) \otimes \mathcal{V})$, or equivalently $D\Omega_q(P \cdot, P \cdot, q \cdot) = \frac{1}{2}\Phi$.*
- (iv) *$D\Omega_{q|_{\mathcal{H} \times \mathcal{V} \times \mathcal{H}}}$ is skew-symmetric w.r.t. the first and third variables, i.e. $D\Omega_q(PX, qY, PZ) = -\frac{1}{2}\Phi(Z, X, Y)$.*
- (v) $\text{Skew} \left(D\Omega_{q|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}} \right) = 0$.

We will then say that (N, q, h) is of type $\mathcal{H}^2\mathcal{V}$.

Corollary 6.3 *Let (N, h) be a Riemannian manifold with an orthogonal decomposition $TN = \mathcal{H} \oplus \mathcal{V}$. Then, the following statements are equivalent.*

- (i) *There exists a metric connection ∇ leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$, such that the following component of the torsion $T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}$ is skew-symmetric, i.e. $T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})} = \text{Skew}(R_{\mathcal{V}})$.*
- (ii) *For any metric connection ∇ leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$, the component of the torsion $T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}$ is skew-symmetric.*
- (iii) *$D\Omega_{q|_{\mathcal{V} \times \mathcal{V} \times \mathcal{H}}}$ is skew-symmetric w.r.t. the two first variables, i.e. $Dq|_{\mathcal{V} \times \mathcal{V}} \in \mathcal{C}((\Lambda^2 \mathcal{V}^*) \otimes \mathcal{H})$, or equivalently $D\Omega_q(q \cdot, q \cdot, P \cdot) = -\frac{1}{2}R_{\mathcal{V}}$.*
- (iv) *$D\Omega_{q|_{\mathcal{V} \times \mathcal{H} \times \mathcal{V}}}$ is skew-symmetric w.r.t. the first and third variables, i.e. $D\Omega_q(qX, PY, qZ) = \frac{1}{2}R_{\mathcal{V}}(Z, X, Y)$.*
- (v) $\text{Skew} \left(D\Omega_{q|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}} \right) = 0$.

We will then say that (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$.

Proof of corollary 6.2 and 6.3. We have seen in the proof of theorem 6.1 that for any metric connection ∇ , we have $\nabla\Omega_{q|_{\mathcal{H}^3}} = 0$ and $\nabla\Omega_{q|_{\mathcal{V}^3}} = 0$, so that we have $\nabla q = 0$ if and only if $\nabla q|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = 0$ and $\nabla q|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})} = 0$, which is equivalent respectively to the conditions on $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = 0$ and $T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})} = 0$ respectively, described by theorem 6.1. In particular, we see that there always exists metric connection ∇ leaving invariant the decomposition of TN , which provides us the implication (ii) \Rightarrow (i). Moreover, a necessary condition for (i) is $\text{Sym}_{\mathcal{H} \times \mathcal{H}} \left(D\Omega_{q|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}}} \right) = 0$, and respectively $\text{Sym}_{\mathcal{V} \times \mathcal{V}} \left(D\Omega_{q|_{\mathcal{V} \times \mathcal{V} \times \mathcal{H}}} \right) = 0$. Conversely if this condition is satisfied, then according to theorem 6.1, (ii) is also satisfied (since $T \in \mathcal{T}$, then the skew-symmetry of $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ (resp. $T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}$) is equivalent to $\text{Sym}_{\mathcal{H} \times \mathcal{H}} (T|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}) = 0$, resp. $\text{Sym}_{\mathcal{V} \times \mathcal{V}} (T|_{\mathcal{H} \times \mathcal{V} \times \mathcal{V}}) = 0$). Therefore we have proved the sequence of implications (ii) \Rightarrow (i) \Rightarrow (iii) \Rightarrow (ii), i.e. (i), (ii) and (iii) are equivalent. Concerning the equivalent reformulation

of (i) and (iii), the former follows from theorem 6.1 and the latter from the fact that according to the proof of theorem 6.1, the skew-symmetric part (w.r.t. the two first variables X, Y) of $D\Omega_{q|\mathcal{H}\times\mathcal{H}\times\mathcal{V}}$ (resp. $D\Omega_{q|\mathcal{V}\times\mathcal{V}\times\mathcal{H}}$) is $\frac{1}{2}\Phi$ (resp. $-\frac{1}{2}R_{\mathcal{V}}$). The equivalence between (iii) and (iv) follows from the symmetry of $D\Omega_q$ w.r.t. the two last variables. Finally, using the computation done in the proof of compute theorem 6.1, we can compute

$$\begin{aligned}\text{Skew}\left(D\Omega_{q|\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}\right) &= \frac{1}{2}(\text{Skew}(\Phi) + \text{Skew}(U(P\cdot, P\cdot, q\cdot))) - \frac{1}{2}(\text{Skew}(\Phi) - \text{Skew}(U(P\cdot, P\cdot, q\cdot))) \\ &= \text{Skew}(U(P\cdot, P\cdot, q\cdot))\end{aligned}$$

and idem for $\mathcal{S}(\mathcal{V}\times\mathcal{V}\times\mathcal{H})$. This completes the proof. \square

Remark 6.1 According to the previous proof, we see that if (N, q, h) is of type $\mathcal{H}^2\mathcal{V}$, then for any extension $T \in \mathcal{C}(\mathcal{T})$ of the skew-symmetric trilinear form $\text{Skew}(\Phi) \in \mathcal{C}(\mathcal{S}(\mathcal{H}^* \times \mathcal{H}^* \times \mathcal{V}^*))$, the corresponding metric connection ∇ satisfies $\nabla q_{|\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} = 0$. In the same way, if (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$, then for any extension $T \in \mathcal{C}(\mathcal{T})$ of the skew-symmetric trilinear form $\text{Skew}(R_{\mathcal{V}}) \in \mathcal{C}(\mathcal{S}(\mathcal{V}^* \times \mathcal{V}^* \times \mathcal{H}^*))$, the corresponding metric connection ∇ satisfies $\nabla q_{|\mathcal{S}(\mathcal{V}\times\mathcal{V}\times\mathcal{H})} = 0$.

Definition 6.5 We will say that the (orthogonal) decomposition on the Riemannian manifold (N, h) is **reductive** (w.r.t. the metric h) or that (N, q, h) is **reductive** if (N, q, h) is of type $\mathcal{H}^2\mathcal{V}$ and of type $\mathcal{V}^2\mathcal{H}$. This is equivalent to say that there exists a metric connection ∇ leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$, and with skew-symmetric torsion.

Proposition 6.8 (N, q, h) is reductive if and only if the trilinear map

$$\text{Skew}(D\Omega_{q|\mathcal{H}\times\mathcal{H}\times\mathcal{V}}) \oplus \text{Skew}(D\Omega_{q|\mathcal{V}\times\mathcal{V}\times\mathcal{H}})$$

is skew-symmetric.

Proof. An element $\alpha \in \mathcal{C}(\mathcal{H}^* \otimes \mathcal{H}^* \otimes \mathcal{V}^*)$ satisfies $\text{Skew}(\alpha) \in \mathcal{C}(\Lambda^2\mathcal{H}^*) \wedge \mathcal{V}^*$ if and only if $\alpha(X^h, Y^h, Z^v)$ is skew-symmetric w.r.t. (X^h, Y^h) . \square

Proposition 6.9 Let (N, q, h) be a reductive Riemannian manifold. Let us suppose that is given some metric connection ∇^c on \mathcal{V} . Then there exists a metric connection ∇ on N preserving the decomposition $TN = \mathcal{V} \oplus \mathcal{H}$, with skew-symmetric torsion, and which coincides with ∇^c on \mathcal{V} if and only if

$$(D^v - \nabla^c)(X^h, Y^v, Z^v) = -\frac{1}{2}R_{\mathcal{V}}(Y^v, Z^v, X^h) \quad \text{and} \quad (D^v - \nabla^c)|_{\mathcal{V}^3} \in \mathcal{C}(\Lambda^3\mathcal{V}^*).$$

Proof. Let us suppose that such a metric connection ∇ exists. Then we have $D = \nabla - \frac{1}{2}T$ and thus $(D^v - \nabla^c)|_{\mathcal{V}^3} \in \mathcal{C}(\Lambda^3\mathcal{V}^*)$ and $(D^v - \nabla^c)(X^h, Y^v, Z^v) = -\frac{1}{2}T(X^h, Y^v, Z^v) = -\frac{1}{2}T(Y^v, Z^v, X^h) = -\frac{1}{2}R_{\mathcal{V}}(Y^v, Z^v, X^h)$, since T is skew-symmetric and ∇ leaves invariant the decomposition of TN .

Conversely, if ∇^c satisfies the above conditions, then let $\beta \in \mathcal{C}(\Lambda^3\mathcal{H}^*)$ and $\alpha = (D^v - \nabla^c)|_{\mathcal{V}^3} \in \mathcal{C}(\Lambda^3\mathcal{V}^*)$. Let us consider the 3-form

$$T = \beta \oplus \text{Skew}(\Phi) \oplus \text{Skew}(R_{\mathcal{V}}) \oplus \alpha,$$

as well the corresponding metric connection $\nabla = D + \frac{1}{2}T$. Then, since (N, q, h) is reductive, according to theorem 6.1, ∇ preserves the decomposition of TN . Moreover, by definition of ∇ , we have $(\nabla - \nabla^c)_{TN\times\mathcal{V}\times\mathcal{V}} = 0$. This completes the proof. \square

A useful additional property. Let us add the following characterization of the type $\mathcal{V}^2\mathcal{H}$ in term of the vertical torsion of the Levi-Civita connection.

Proposition 6.10 *Let (N, h) be a Riemannian manifold with an orthogonal decomposition $TN = \mathcal{H} \oplus \mathcal{V}$. Let T^v be the vertical torsion of the Levi-Civita connection. Then, (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$ if and only if*

$$\text{Sym}_{\mathcal{V} \times \mathcal{V}}(T^v_{|\mathcal{H} \times \mathcal{V} \times \mathcal{V}}) = 0.$$

Proof Let $H \in \mathcal{H}$, $V_1, V_2 \in \mathcal{V}$. Then

$$\begin{aligned} T^v(H, V_1, V_2) + T^v(H, V_2, V_1) &= \langle D_H^v V_1 - [H, V_1]^v, V_2 \rangle + \langle D_H^v V_2 - [H, V_2]^v, V_1 \rangle \\ &= \langle D_H V_1 - [H, V_1], V_2 \rangle + \langle D_H V_2 - [H, V_2], V_1 \rangle \\ &= \langle D_{V_1} H, V_2 \rangle + \langle D_{V_2} H, V_1 \rangle \\ &= \text{Sym}_{\mathcal{V} \times \mathcal{V}}(DP)(V_1, H, V_2) = \text{Sym}_{\mathcal{V} \times \mathcal{V}}(DP)(V_1, V_2, H) \\ &= -\text{Sym}_{\mathcal{V} \times \mathcal{V}}(Dq)(V_1, V_2, H). \end{aligned}$$

Then we completes the proof by applying corollary 6.3. \square

6.2.3 Characterisation of metric f -connections. Existence of a characteristic connection.

Now, let us come back to the case of a f -manifolds. Then the condition $\nabla F = 0$ is equivalent of the fact that ∇ leaves invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$ and moreover $\nabla^{\mathcal{H}} \bar{J} = 0$, where $\nabla^{\mathcal{H}}$ is the connection induced by ∇ on \mathcal{H} . In some sense, we have to add to the conditions of theorem 6.1 those of theorem as well as the condition $\nabla^{\mathcal{H}} \bar{J} = 0$.

Definition 6.6 *We will say that a f -structure F and a metric h on a manifold N are compatible if $\mathcal{H} \perp \mathcal{V}$ and if \bar{J} is an orthogonal complex structure on \mathcal{H} endowed with the metric induced by h . This is equivalent to say that F is skew-symmetric w.r.t. the metric h : $F \in \mathfrak{so}(TN)$, or equivalently that $I = \bar{J} \oplus \text{Id}_{\mathcal{V}}$ is orthogonal: $I^*h = h$. We will then say that (N, F, h) is a metric f -manifold.*

Theorem 6.2 *Let (N, F, h) be a metric f -manifold. Then a metric connection ∇ preserves the f -structure F if and only if all the following statements hold:*

$$\begin{aligned} \nabla \Omega_F|_{\mathcal{H}^3} = 0 &\iff N_{\bar{J}} = 4 (T|_{\mathcal{H}^3})^{0,2} \text{ and } \text{Skew} \left((T|_{\mathcal{H}^3})^{2,0} - (T|_{\mathcal{H}^3})^{1,1} \right) = (d^c \Omega_F|_{\mathcal{H}^3})^{**} \\ \nabla \Omega_F|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = 0 &\iff \left\{ \begin{array}{l} \nabla \Omega_F|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}} = 0 \\ \nabla \Omega_F|_{\mathcal{H} \times \mathcal{V} \times \mathcal{H}} = 0 \\ \nabla \Omega_F|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}} = 0 \end{array} \right\} \iff \left\{ \begin{array}{l} \nabla \Omega_q|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = 0 \\ T(X^v, FY^h, Z^h) + T(X^v, Y^h, FZ^h) = \\ N(X^v, Y^h, FZ^h) \end{array} \right. \\ \nabla \Omega_F|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})} = 0 &\iff \nabla \Omega_q|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})} = 0 \end{aligned}$$

Proof. We first notice that $\nabla \Omega_F(X^h, Y^h, Z^v) = \langle (\nabla_{X^h} F)Y^h, Z^v \rangle = \langle \nabla_{X^h}(FY^h), Z^v \rangle = -\nabla \Omega_q(X^h, FY^h, Z^v)$. Therefore $\nabla \Omega_F|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}} = 0 \Leftrightarrow \nabla \Omega_q|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}} = 0$. In the same way, $\nabla \Omega_F|_{\mathcal{H} \times \mathcal{V} \times \mathcal{H}} = 0 \Leftrightarrow \nabla \Omega_q|_{\mathcal{H} \times \mathcal{V} \times \mathcal{H}} = 0$. Therefore, according to the proof of corollary 6.2, we have proved the equivalence between the two boxes.

Now, let us compute

$$\begin{aligned} T(X^v, FY^h, Z^h) + T(X^v, Y^h, FZ^h) &= \langle \nabla_{X^v}(FY^h) - \nabla_{FY^h}X^v - [X^v, FY^h], Z^h \rangle \\ &\quad + \langle \nabla_{X^v}Y^h - \nabla_{Y^h}X^v - [X^v, Y^h], FZ^h \rangle \\ &= \nabla_{\Omega_F}(X^v, Y^h, Z^h) - \nabla_{\Omega_q}(FY^h, X^v, Z^h) - \nabla_{\Omega_q}(Y^h, X^v, FZ^h) + N_F(X^v, Y^h, FZ^h) \end{aligned}$$

so that if $\nabla_{\Omega_q|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}} = 0$ then we obtain the equivalence $(\nabla_{\Omega_F|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}} = 0) \Leftrightarrow (T(X^v, FY^h, Z^h) + T(X^v, Y^h, FZ^h) = N(X^v, Y^h, FZ^h))$.

We have also to compute that $\nabla_{\Omega_F}(X^v, Y^v, Z^h) = \langle (\nabla_{X^v}F)Y^v, Z^h \rangle = -\langle F\nabla_{X^v}Y^v, Z^h \rangle = \nabla q(X^v, Y^v, FZ^h)$, also that $\nabla_{\Omega_F}(X^v, Y^h, Z^v) = -\nabla q(X^v, FY^h, Z^v)$, and that $\nabla_{\Omega_F}(X^h, Y^v, Z^v) = 0$. Therefore, according to the proof of corollary 6.3, we have proved the equivalence $(\nabla_{\Omega_F|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}} = 0) \Leftrightarrow (\nabla_{\Omega_q|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}} = 0)$.

Furthermore, we can prove (by a straightforward but a little bit long computation) the following formula generalizing equation (109):

$$\begin{aligned} 2D\Omega_F(X, Y, Z) &= d\Omega_F(X, Y, Z) - d\Omega_F(X, FY, FZ) + N(Y, Z, FX) \\ &\quad + \Phi(Y, FZ, X) + \Phi(FY, Z, X) - \Phi(X, FY, Z) - \Phi(FZ, X, Y) \\ &\quad + \text{Sym}_{\mathcal{V} \times \mathcal{V}}(\nabla_{\Omega_F|_{\mathcal{V} \times \mathcal{H} \times \mathcal{V}}}) + \text{Sym}_{\mathcal{V} \times \mathcal{V}}(\nabla_{\Omega_F|_{\mathcal{V} \times \mathcal{V} \times \mathcal{H}}}). \end{aligned} \quad (115)$$

We deduce from this, that

$$2D\Omega_F(X^h, Y^h, Z^h) = d\Omega_F(X^h, Y^h, Z^h) - d\Omega_F(X^h, FY^h, FZ^h) + N(Y^h, Z^h, FX^h).$$

Moreover, $\nabla_{\Omega|_{\mathcal{H}^3}} = 0$ if and only if

$$D\Omega_F|_{\mathcal{H}^3} = -\frac{1}{2}(T + U)|_{\mathcal{H}^3}.$$

Therefore, we can proceed as in the proof of theorem 5.13 to prove the equivalence concerning the restriction to \mathcal{H}^3 . This completes the proof. \square

Remark 6.2 A priori, we could think that we can deduce from this theorem that in a metric f -manifold, it could not exist any metric f -connection. Indeed, we see that the condition $T(X^v, FY^h, Z^h) + T(X^v, Y^h, FZ^h) = N_F(X^v, Y^h, FZ^h)$ is compatible with the condition $\nabla_{\Omega_q|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}} = 0$ if and only if

$$\text{Sym}_{\mathcal{H} \times \mathcal{H}}(N_F|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}) = 2 \left[\text{Sym}_{\mathcal{H} \times \mathcal{H}}(D\Omega_q|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}}) \right]^{(2,0)+(0,2)}(Y^h, Z^h, X^v) \quad (116)$$

according to theorem 6.1. In other words, the existence of some metric connection ∇ such that $\nabla_{\Omega_F|_{\mathcal{S}(\mathcal{V} \times \mathcal{H} \times \mathcal{H})}} = 0$ is possible if and only if equation (116) holds. Nevertheless, the existence of some metric connection ∇ such that $\nabla F|_{\mathcal{H}^3} = 0$ and $\nabla F|_{\mathcal{S}(\mathcal{H} \times \mathcal{V} \times \mathcal{V})} = 0$ always holds without condition. But in fact, equation (116) also always holds.

Lemma 6.1 *In a metric f -manifold (N, F, h) , the following identity holds*

$$\text{Sym}_{\mathcal{H} \times \mathcal{H}}(N_F|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}) = 2 \left[\text{Sym}_{\mathcal{H} \times \mathcal{H}}(D\Omega_q|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}}) \right]^{(2,0)+(0,2)}(Y^h, Z^h, X^v).$$

Any metric f -manifold (N, F, h) admits a metric f -connection.

Proof. This can be checked directly by computation (use the characterization of the Levi-Civita connection, as in the proof of theorem 6.1). \square

Remark 6.3 In a metric f -manifold, there are several constraints on the component $T_{|\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ of the torsion of some metric f -connection ∇ : the components $(T_{|\mathcal{V} \times \mathcal{H} \times \mathcal{H}})^{(2,0)+(0,2)}$ and $\text{Sym}_{\mathcal{H} \times \mathcal{H}} \left([T_{|\mathcal{V} \times \mathcal{H} \times \mathcal{H}}]^{1,1} \right)$ are determined. The only degree of freedom is on $\text{Skew}_{\mathcal{H} \times \mathcal{H}} \left([T_{|\mathcal{V} \times \mathcal{H} \times \mathcal{H}}]^{1,1} \right)$ which can be chosen freely. Therefore, this is also the only degree of freedom of $T_{|\mathcal{S}(\mathcal{V} \times \mathcal{H} \times \mathcal{H})}$, according to theorem 6.1.

Remark 6.4 A compatible f -structure F on a Riemannian manifold (N, h) defines a reduction of the orthogonal frame bundle of TN (which is of course a $O(n)$ -bundle) to some $U(2p) \times O(n-2p)$ -bundle, where $n = \dim N$ and $2p$ is the rank of F . Therefore a metric f -connection is nothing but a connection on this $U(2p) \times O(n-2p)$ -bundle, which always exists. This provides a new proof of lemma 6.1.

Remark 6.5 If (N, F, h) is of type $\mathcal{H}^2\mathcal{V}$, then $N_{F|\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ is skew-symmetric w.r.t. the two last variables.

Skew-symmetric torsion. Further, we are interested by metric f -connections with skew-symmetric torsion. As we have done above we have to study first the condition of skew-symmetry on each component of the torsion and then to group all the obtained conditions to obtain a global condition on the metric f -manifold for the existence of metric f -connection with skew-symmetric torsion.

Definition 6.7 Let (N, F, h) be a metric f -manifold. We define the **extended Nijenhuis tensor** \tilde{N}_F as the TN -valued 2-form on N (whose corresponding trilinear map is) defined by

$$\tilde{N}_F := N_F + \Phi + R_{\mathcal{V}}(Z^v, X^v, Y^h) + R_{\mathcal{V}}(Y^v, Z^v, X^h).$$

We remark that $\tilde{N}_{F|\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})} = \text{Skew}(R_{\mathcal{V}})$ is always skew-symmetric.

Proposition 6.11 Let (N, F, h) be a metric f -manifold. Then the following statements are equivalent.

- (i) There exists a metric f -connection ∇ (satisfying then $\nabla F = 0$) with a torsion T such that $T^{0,2}_{|\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ is skew-symmetric.
- (ii) There exists a metric connection ∇ , satisfying $\nabla F_{|\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = 0$, with a torsion T such that $T^{0,2}_{|\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ is skew-symmetric.
- (iii) $N_F(FY^h, Z^h, X^v) + N_F(Y^h, FZ^h, X^v) = N_F(X^v, Y^h, FZ^h)$.
- (iv) The extended Nijenhuis tensor \tilde{N}_F satisfies: $\tilde{N}_{F|\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ is skew-symmetric.

Proof. Since the condition " $T^{0,2}_{|\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ is skew-symmetric" concerns only the subspace $\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})$, then (i) and (ii) are equivalent. Moreover, we have

$$\tilde{N}_{F|\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = N_{F|\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} - N_F(F \cdot, F \cdot, q \cdot)$$

(according to the definition of \tilde{N}_F and proposition 6.1), and this equality gives easily the equivalence between (iii) and (iv). Now, it remains to prove the equivalence between (i) and (iv).

Let us first recall that

$$-4(T^{0,2}) := T(F \cdot, F \cdot, q \cdot) + T(F \cdot, q \cdot, F \cdot) + T(q \cdot, F \cdot, F \cdot) - T$$

Let us suppose (i). Then we have according to proposition 6.7, that $T(F\cdot, F\cdot, q\cdot) + T(F\cdot, q\cdot, F\cdot) + T(q\cdot, F\cdot, F\cdot) - T(\cdot, \cdot, P\cdot) = -N_F$ and thus

$$-4T^{0,2} = -N_F - T(\cdot, \cdot, q\cdot) = -N_F - \Phi - T|_{\mathcal{V}^3} - T|_{(\mathcal{V}\wedge\mathcal{H})\otimes\mathcal{V}}$$

and hence $-4T^{0,2}|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} = -\tilde{N}_F|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$, which implies (iv).

Conversely, let us suppose (iv). Then we have to construct a trilinear map $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$, on $\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})$, such that

(C₁): $-4T^{0,2}|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ is a 3-form, and

(C₂): the following conditions of theorem 6.1 hold

$$T|_{\mathcal{H}\times\mathcal{H}\times\mathcal{V}} = \Phi, \quad \text{Sym}_{\mathcal{H}\times\mathcal{H}}(T|_{\mathcal{V}\times\mathcal{H}\times\mathcal{H}}) = \text{Sym}_{\mathcal{H}\times\mathcal{H}}(D\Omega_{q|_{\mathcal{V}\times\mathcal{H}\times\mathcal{H}}})(Y^h, Z^h, X^v).$$

Indeed, for any trilinear map $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} \in \mathcal{C}(\mathcal{T})$, let us set $T|_{\mathcal{V}\times\mathcal{H}\times\mathcal{H}} = S + A$, where S, A are resp. symmetric and skew-symmetric w.r.t. the two last variables. Then we can prove easily that a necessary condition on S, A so that $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ satisfies the required conditions (C₁)-(C₂) above, is

$$S = \text{Sym}_{\mathcal{H}\times\mathcal{H}}(D\Omega_{q|_{\mathcal{H}\times\mathcal{H}\times\mathcal{V}}})(Y^h, Z^h, X^v) \quad (117a)$$

$$S^{(2,0)+(0,2)} = 0 \quad (117b)$$

$$A^{(2,0)+(0,2)}(X^v, Y^h, Z^h) = \Phi^{(2,0)+(0,2)}(Y^h, Z^h, X^v) \quad (117c)$$

We then see that the two equations on S are compatible according to (iv) and lemma 6.1, and defines then uniquely S . Moreover the equation on A implies $T(X^v, FY^h, Z^h) + T(X^v, Y^h, FZ^h) = N(X^v, Y^h, FZ^h)$, according to (iv) (written in the form of (iii)) and (117b). Therefore, we see that, according to theorem 6.2, the conditions (C₁)-(C₂) on $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ implies that $\nabla F|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} = 0$, for any metric connection ∇ whose the torsion $T \in \mathcal{C}(\mathcal{T})$ is an extension of $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$.

Now, we can verify easily that any trilinear map $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} \in \mathcal{T}$ defined by $T|_{\mathcal{H}\times\mathcal{H}\times\mathcal{V}} = \Phi$ and $T|_{\mathcal{V}\times\mathcal{H}\times\mathcal{H}} = S + A$ with A, S satisfying (117), satisfies the required conditions (C₁)-(C₂). We have then proved (ii). This completes the proof. \square

Proposition 6.12 *Let (N, F, h) be a metric f -manifold. Then the following statements are equivalent.*

(i) *There exists a metric f -connection ∇ (satisfying then $\nabla F = 0$) with a torsion T such that $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ is skew-symmetric.*

(ii) *There exists a metric connection ∇ , satisfying $\nabla F|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} = 0$, with a torsion T such that $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ is skew-symmetric.*

(iii) *(N, q, h) is of type $\mathcal{H}^2\mathcal{V}$, and $\tilde{N}_F|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ is skew-symmetric.*

Furthermore, under these statements, for any such connection satisfying (i) or (ii), then $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})}$ is unique (i.e. uniquely determined by the metric f -manifold (N, F, h)) and equal to $\text{Skew}(\Phi)$. Conversely, any extension $T \in \mathcal{C}(\mathcal{T})$ of this unique skew-symmetric trilinear form $\text{Skew}(\Phi)$ defines a metric connection ∇ , satisfying $\nabla F|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} = 0$.

Proof. (i) and (ii) are equivalent for the same reason as in the proof of the previous proposition. Moreover, (i) and (iii) are equivalent according to theorem 6.2 and corollary 6.2. Moreover, by skew-symmetry and theorem 6.1, we have $T|_{\mathcal{S}(\mathcal{H}\times\mathcal{H}\times\mathcal{V})} = \text{Skew}(\Phi)$. This completes the proof. \square

We are led to the following definition.

Definition 6.8 We will say that a metric f -manifold (N, F, h) is reductive if (N, q, h) is reductive, where q is defined by F .

We will say that a metric f -manifold (N, F, h) is reductively of type \mathcal{G}_1 if $\tilde{N}_{F|S(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ is skew-symmetric.

Now, let us turn ourself on the horizontal component \mathcal{H}^3 of TN^3 .

Definition 6.9 We will say that a metric f -manifold is horizontally of type \mathcal{G}_1 or that it is of horizontal type \mathcal{G}_1 if one the following equivalent statements holds.

- (i) The horizontal Nijenhuis tensor $N_{\bar{J}}$ is skew-symmetric.
- (ii) There exists a metric f -connection ∇ , such that $(T|_{\mathcal{H}^3})^{0,2}$ is skew-symmetric.
- (iii) There exists a metric connection ∇ , satisfying $\nabla F|_{\mathcal{H}^3} = 0$, such that $(T|_{\mathcal{H}^3})^{0,2}$ is skew-symmetric.

Proposition 6.13 Let (N, F, h) be a metric f -manifold. Then the following statements are equivalent.

- (i) (N, F, h) is horizontally of type \mathcal{G}_1 .
- (ii) There exists a metric f -connection ∇ , such that $T|_{\mathcal{H}^3}$ is skew-symmetric.
- (iii) There exists a metric connection ∇ , satisfying $\nabla F|_{\mathcal{H}^3} = 0$, such that $T|_{\mathcal{H}^3}$ is skew-symmetric.

In this case, for any such connection satisfying (i) or (ii), then $T|_{\mathcal{H}^3}$ is unique (i.e. uniquely determined by the metric f -manifold (N, F, h)). Conversely any extension $T \in \mathcal{C}(\mathcal{T})$ of this unique skew-symmetric trilinear form $T|_{\mathcal{H}^3}$ defines a metric connection ∇ , satisfying $\nabla F|_{\mathcal{H}^3} = 0$.

Proof. (ii) and (iii) are equivalent for the same reason as above. Furthermore, according to theorem 6.2, if $\nabla F|_{\mathcal{H}^3} = 0$ and $T|_{\mathcal{H}^3}$ is skew-symmetric, then $N_{\bar{J}} = 4(T|_{\mathcal{H}^3})^{0,2}$ is also skew-symmetric (according to corollary 5.2 applied to the Hermitian bundle $(\mathcal{H}, \bar{J}, h_{\mathcal{H}})$) and moreover $-(T|_{\mathcal{H}^3})^{**} = (d^c \Omega_{F|_{\mathcal{H}^3}})^{**}$, which proves the unicity of $T|_{\mathcal{H}^3}$.

Conversely, if (i) is satisfied, then let $T \in \mathcal{C}(\mathcal{T})$ such that $(T|_{\mathcal{H}^3})^{0,2} = \frac{1}{4}N_{\bar{J}}$, $(T|_{\mathcal{H}^3})^{**} = -(d^c \Omega_{F|_{\mathcal{H}^3}})^{**}$, and the other components being arbitrary. Then $T|_{\mathcal{H}^3}$ is skew-symmetric, and the corresponding metric connection ∇ satisfies $\nabla \Omega_{F|_{\mathcal{H}^3}} = 0$, according to theorem 6.2. This completes the proof. \square

Now, let us regroup the previous results to conclude.

Definition 6.10 A metric f -manifold (N, F, h) with skew-symmetric extended Nijenhuis tensor \tilde{N}_F will be said of **global type \mathcal{G}_1** or globally of type \mathcal{G}_1 .

Theorem 6.3 A metric f -manifold (N, F, h) admits a metric f -connection ∇ with skew-symmetric torsion if and only if it is reductive and of global type \mathcal{G}_1 . Moreover, in this case, for any $\alpha \in \mathcal{C}(\Lambda^3 \mathcal{V}^*)$, there exists a unique metric connection ∇ with skew-symmetric torsion such that $T|_{\Lambda^3 \mathcal{V}} = \alpha$. This unique connection is given by

$$T = (-d^c \Omega_F + N_{F|_{\mathcal{H}^3}}) + \text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}}) + \alpha.$$

Proof. The first assertion follows from propositions 6.13, 6.12, theorem 6.2, corollary 6.3 and remark 6.1. Then in this case, $T|_{\mathcal{H}^3}$ is entirely determined, according to proposition 6.13. Moreover, by skew-symmetry and theorem 6.1, we have $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = \text{Skew}(\Phi)$ and $T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})} = \text{Skew}(\mathbf{R}_{\mathcal{V}})$. Now, let us determine $T|_{\mathcal{H}^3}$. Since, $\nabla = D + \frac{1}{2}T$, the equation $\nabla F = 0$ can be written

$$D\Omega_F + \frac{1}{2}(T(\cdot, F\cdot, \cdot) + T(\cdot, \cdot, F\cdot)) = 0$$

so that

$$d\Omega_F = -F \circlearrowleft T$$

and thus $d\Omega_F|_{\mathcal{H}^3} = -\bar{J} \circlearrowleft T|_{\mathcal{H}^3} = 4\bar{J} \cdot (T|_{\mathcal{H}^3})^{--} - \bar{J} \cdot T|_{\mathcal{H}^3} = \bar{J} \cdot N_{\bar{J}} - \bar{J} \cdot T|_{\mathcal{H}^3}$. This completes the proof. \square

Remark 6.6 *In the particular case where \mathcal{V} is a line bundle, (N, F, h) is an almost contact metric manifold. Then in this particular case, the theorem 6.3 above allows us to recover the result of [17], theorem 8.2.*

Definition 6.11 *On a metric f -manifold (N, F, h) , a metric f -connection ∇ with skew-symmetric torsion will be called characteristic connection.*

Corollary 6.4 *Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . Let us suppose that is given some metric connection ∇^c on the vertical subbundle \mathcal{V} . There exists a metric f -connection ∇ on N with skew symmetric torsion, which coincides with ∇^c on \mathcal{V} if and only if*

$$(D^v - \nabla^c)(X^h, Y^v, Z^v) = -\frac{1}{2}\mathbf{R}_{\mathcal{V}}(Y^v, Z^v, X^h) \quad \text{and} \quad (D^v - \nabla^c)|_{\mathcal{V}^3} \in \mathcal{C}(\Lambda^3\mathcal{V}^*).$$

In this case this connection ∇ is unique and will be called the characteristic connection extending or defined by ∇^c .

Remark 6.7 In other words, in a reductive metric f -manifold of global type \mathcal{G}_1 , the set of metric connection ∇^c on the vertical subbundle \mathcal{V} which can be extended to a characteristic connection, is the affine space

$$D^v - \frac{1}{2}\mathbf{R}_{\mathcal{V}}(Y^v, Z^v, X^h) + \mathcal{C}(\Lambda^3\mathcal{V}^*).$$

Proof of corollary 6.4. This follows immediately from the theorem 6.3, the proposition 6.9 and the theorem 6.2. \square

Proposition 6.14 *Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . Let ∇ be some characteristic connection on N . Then the we have*

$$\begin{aligned} d\Omega_F = -F \circlearrowleft T &= F \cdot N_F - F \cdot T - F \circlearrowleft (T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} + T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}) \\ &= F \cdot N_F - F \bullet T - \frac{1}{2}F \circlearrowleft (T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} + T|_{\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})}). \end{aligned}$$

Proof. We have seen (in the proof of theorem 6.3) that $d\Omega_F = -F \circlearrowleft T$, and that $(-F \circlearrowleft T)|_{\mathcal{H}^3} = F \cdot N_F - F \cdot T$, moreover we have $(-F \circlearrowleft T)|_{\mathcal{V}^3} = 0$, by definition of the action $F \circlearrowleft$. This completes the proof. \square

A useful expression of the Nijenhuis tensor in term of the Levi-Civita connection.

By a direct computation, we prove:

Proposition 6.15 *Let (N, F, h) be a metric Riemannian f -manifold. Then we have*

$$N_F(X, Y) + d^D q(X, Y) = (D_{FX}F)Y - (D_{FY}F)X + (D_XF)(FY) - (D_YF)(FX).$$

6.2.4 Precharacteristic and paracheracteristic connections.

Sometimes the condition of global type G_1 could be too much strong and it could happen that one needs the existence (and unicity up the \mathcal{V}^3 -component of the torsion) of some characteristic connection by supposing weaker conditions on the metric f -manifold (N, F, h) .

Definition 6.12 *Let (N, F, h) be a metric f -manifold of horizontal type \mathcal{G}_1 . Then any metric f -connection ∇ with a skew-symmetric component $T|_{\mathcal{H}^3}$ of its torsion, will be called a horizontal-characteristic connection.*

Moreover, if we suppose that (N, F, h) is also of type $\mathcal{V}^2\mathcal{H}$, then a metric f -connection ∇ with skew-symmetric components $T|_{\mathcal{H}^3}$, $T|_{\mathcal{S}(\mathcal{V}\times\mathcal{V}\times\mathcal{H})}$ and $T|_{\mathcal{V}^3}$ of its torsion, will be called a precharacteristic connection.

Remark 6.8 Let us remark that in a metric f -manifold of horizontal type \mathcal{G}_1 , horizontal-characteristic connections always exist, and the component $T|_{\mathcal{H}^3}$ of the torsion is unique. Moreover, if we suppose that (N, F, h) is also of type $\mathcal{V}^2\mathcal{H}$, then precharacteristic connections always exist and the components $T|_{\mathcal{H}^3}$ and $T|_{\mathcal{S}(\mathcal{V}\times\mathcal{V}\times\mathcal{H})}$ are unique.

The following properties will hold for the horizontal curvature in all the examples of interest for us.

Definition 6.13 *Let (N, F) be a f -manifold. Let $A \in \mathcal{C}(\mathcal{H}^* \otimes \mathcal{H}^* \otimes \mathcal{V})$, $B \in \mathcal{C}(\mathcal{H}^* \otimes \mathcal{V}^* \otimes \mathcal{H})$, and $C \in \mathcal{C}(\mathcal{V}^* \otimes \mathcal{H}^* \otimes \mathcal{H})$. Then we will say respectively that A , B or C is pure if respectively*

- (i) $A(\bar{J}X, Y) = A(X, \bar{J}Y)$, $\forall X, Y \in \mathcal{H}$.
- (ii) $B(\bar{J}X, Y) = -\bar{J}B(X, Y)$, $\forall X \in \mathcal{H}, Y \in \mathcal{V}$.
- (iii) $C(X, \bar{J}Y) = -\bar{J}C(X, Y)$, $\forall X \in \mathcal{V}, Y \in \mathcal{H}$.

If (N, F) is endowed with a compatible metric h , then this means that A , B or C considered as element of $\mathcal{C}(\mathcal{H}^ \otimes \mathcal{H}^* \otimes \mathcal{V})$, satisfies respectively $A^{1,1} = 0$, $B^{1,1} = 0$, $C^{1,1} = 0$. Moreover, we will say that A , B or C resp. (considered as trilinear forms) is skew-symmetric in $\mathcal{H} \times \mathcal{H}$ if resp. A is skew-symmetric w.r.t. the 2 first variables, B w.r.t. the first and third variables, and C w.r.t. the 2 last variables.*

Let us remark that $N_F|_{\mathcal{V}\times\mathcal{H}\times\mathcal{H}}$ is pure by definition of N_F (see proposition 6.1), and skew-symmetric in $\mathcal{H} \times \mathcal{H}$ if (N, F, h) is of type $\mathcal{H}^2\mathcal{V}$ (see remark 6.5).

Definition 6.14 \bullet *(N, F, h) will be called almost of type $\mathcal{H}^2\mathcal{V}$ if one of the following equivalent statements holds*

- (i) $\text{Sym}_{\mathcal{H}\times\mathcal{H}} \left(D\Omega_q|_{\mathcal{H}\times\mathcal{H}\times\mathcal{V}} \right)$ is pure.

$$(ii) \text{Sym}_{\mathcal{H} \times \mathcal{H}} (N_{F|\mathcal{V} \times \mathcal{H} \times \mathcal{H}}) = 2 \text{Sym}_{\mathcal{H} \times \mathcal{H}} \left(D\Omega_{q|\mathcal{H} \times \mathcal{H} \times \mathcal{V}} \right) (Y^h, Z^h, X^v).$$

• If moreover (N, F, h) is of type $\mathcal{V}^2\mathcal{H}$, then we will say that it is almost reductive.

Remark 6.9 We remark that if (N, F, h) is of type $\mathcal{H}^2\mathcal{V}$ then it is almost of type $\mathcal{H}^2\mathcal{V}$, and therefore if it is reductive then it is, in particular, almost reductive.

Theorem 6.4 A metric f -manifold (N, F, h) admits a precharacteristic connection ∇ such that the component $T|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ of its torsion is pure if and only if it is almost reductive and horizontally of type \mathcal{G}_1 . Moreover, in this case, for any $\alpha \in \mathcal{C}(\Lambda^3\mathcal{V}^*)$, there exists a unique precharacteristic connection ∇ such that the component $T|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ of its torsion is pure, and such that $T|_{\Lambda^3\mathcal{V}} = \alpha$. This unique connection is given by

$$T = (-d^c\Omega_F + N_{F|\mathcal{H}^3}) + T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} + \text{Skew}(\mathbf{R}_{\mathcal{V}}) + \alpha.$$

where $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{V} \times \mathcal{V})} = \Phi + \frac{1}{2}(N_F(X^v, Y^h, Z^h) - N_F(Y^v, X^h, Z^h))$.

Moreover if we impose also the component $T|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ to be skew-symmetric in $\mathcal{H} \times \mathcal{H}$, then this is possible if and only if (N, F, h) is reductive and of horizontal type \mathcal{G}_1 .

Proof. Concerning the components in \mathcal{H}^3 , $\mathcal{S}(\mathcal{V} \times \mathcal{V} \times \mathcal{H})$ and \mathcal{V}^3 , we can proceed as in the proof of theorem 6.3. Concerning the component in $\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})$, use the identity $T(X^v, FY^h, Z^h) + T(X^v, Y^h, FZ^h) = N(X^v, Y^h, FZ^h)$ from theorem 6.2 and the identity $T|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}} = \Phi$ from theorem 6.1. This completes the proof. \square

Definition 6.15 On a metric f -manifold (N, F, h) , a precharacteristic connection ∇ such that the component $T|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ of the torsion is pure, will be called a **paracharacteristic** connection.

Proposition 6.16 Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . Then the paracharacteristic connection (defined by some $\alpha \in \mathcal{C}(\Lambda^3\mathcal{V}^*)$) coincide with the characteristic connection (defined by the same $\alpha \in \mathcal{C}(\Lambda^3\mathcal{V}^*)$) if and only if the horizontal curvature Φ is pure.

Proof. According to theorems 6.3 and 6.4, we only have to prove that the components $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})}$ of the torsions of the two connections coincide if Φ is pure. But, since Φ is pure, the component $T|_{\mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V})} = \text{Skew}(\Phi)$ of the characteristic connection is such that $T|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ is pure, and therefore by unicity of the paracharacteristic connection, the two connections coincide. Conversely, if they coincide, then $\text{Skew}(\Phi)|_{\mathcal{V} \times \mathcal{H} \times \mathcal{H}}$ is pure, i.e. Φ is pure. This completes the proof. \square

The Linear representations of the curvatures. Using the metric h , we have canonical isomorphisms $\Lambda^2\mathcal{H}^* \cong \mathfrak{so}(\mathcal{H})$ and $\Lambda^2\mathcal{V}^* \cong \mathfrak{so}(\mathcal{V})$. Let us denote by $\rho \in \mathfrak{so}(\mathcal{H}) \otimes \mathcal{V}^*$ and $\sigma \in \mathfrak{so}(\mathcal{V}) \otimes \mathcal{H}^*$ respectively, the elements corresponding to Φ and $\mathbf{R}_{\mathcal{V}}$ respectively under these isomorphisms:

$$\langle \rho(V)H_1, H_2 \rangle = \langle \Phi(H_1, H_2), V \rangle, \quad H_1, H_2 \in \mathcal{H}, V \in \mathcal{V}, \quad (118)$$

and

$$\langle \sigma(H)V_1, V_2 \rangle = \langle \mathbf{R}_{\mathcal{V}}(V_1, V_2), H \rangle, \quad V_1, V_2 \in \mathcal{V}, H \in \mathcal{H}. \quad (119)$$

To do not weigh down the notation, we denote $\bar{J} = \bar{J}$. Let us introduce the horizontal curvature operator :

$$\bar{R}(X, Y)Z = \rho(\Phi(X, Y))Z.$$

as well as its derivation term

$$\bar{A}(X, Y) = \bar{R}(\bar{J}X, Y) + \bar{R}(X, \bar{J}Y) - [\bar{J}, \bar{R}(X, Y)].$$

(which vanishes if and only if \bar{J} is a derivation of \bar{R} hence its name). We denote by $\Phi = \Phi^{(+)} + \Phi^{(-)}$ the splitting of Φ according to the eigenspace decomposition of the endomorphism of $\Lambda^2\mathcal{H}^* \otimes \mathcal{V}$ defined by $B \mapsto B(\bar{J}\cdot, \bar{J}\cdot)$, i.e.

$$\Phi^{(\varepsilon)}(\bar{J}\cdot, \bar{J}\cdot) = \varepsilon\Phi^{(\varepsilon)}.$$

In other words, $\Phi^{(+)}$ is the $(1, 1)$ -type part of Φ whereas $\Phi^{(-)}$ is the part of Φ of type $(2, 0) + (0, 2)$. Under the isomorphism $\Lambda^2\mathcal{H}^* \cong \mathfrak{so}(\mathcal{H})$, to this, corresponds the decomposition $\rho = \rho^+ + \rho^-$, where $\bar{J}\rho^{\varepsilon}\bar{J}^{-1} = \varepsilon\rho^{\varepsilon}$, according to the splitting $\mathfrak{so}(\mathcal{H}) = \mathfrak{so}_+(\mathcal{H}) \oplus \mathfrak{so}_-(\mathcal{H})$ of $\mathfrak{so}(\mathcal{H})$ following its \bar{J} -commuting and \bar{J} -anticommuting parts. Then, this being done, we can define the corresponding curvature operator and antiderivation terms:

$$\bar{R}^{(\varepsilon)} = \rho^{\varepsilon}(\Phi^{(\varepsilon)}) \quad \text{and} \quad \bar{A}^{(\varepsilon)}(X, Y) = \bar{R}^{(\varepsilon)}(\bar{J}X, Y) + \bar{R}^{(\varepsilon)}(X, \bar{J}Y) - [\bar{J}, \bar{R}^{(\varepsilon)}(X, Y)].$$

We remark that $\bar{A}^{(+)} = 0$ and $\bar{A}^{(-)}(X, Y) = 2\bar{R}^{(-)}(X, \bar{J}Y) - 2\bar{J} \circ \bar{R}^{(-)}(X, Y)$.

6.2.5 Reductions of f -manifolds

Definition 6.16 A q -manifold (N, q) is a manifold endowed with a linear projector $q \in \text{End}(TN)$, i.e. with a splitting $TN = \mathcal{V} \oplus \mathcal{H}$. We will say that q is a q -structure. Moreover, we will say that (N, q) is associated to the f -manifold (N, F) if q is associated to F : i.e. $-F^2 = \text{Id} - q$, or equivalently $\ker F = \mathcal{V}$ and $\text{Im } F = \mathcal{H}$.

A metric q -manifold (N, q, h) is a q -manifold endowed with a metric h for which q is orthogonal, or equivalently the splitting $TN = \mathcal{V} \oplus \mathcal{H}$ is orthogonal.

Definition 6.17 A map $i_{\mathcal{V}}: (N^{\mathcal{V}}, q^{\mathcal{V}}) \rightarrow (N, q)$ from a q -manifold to another one is called a q -immersion if it is an immersion which satisfies

$$(i_{\mathcal{V}})^*q = q^{\mathcal{V}}. \tag{120}$$

Then $(N^{\mathcal{V}}, q^{\mathcal{V}})$ is q -immersed in (N, q) . If $i_{\mathcal{V}}$ is injective we will say that $(N^{\mathcal{V}}, q^{\mathcal{V}})$ is a q -submanifold of the q -manifold (N, q) , and that $i_{\mathcal{V}}$ is a q -embedding. Moreover we will say that $i_{\mathcal{V}}: (N^{\mathcal{V}}, q^{\mathcal{V}}) \rightarrow (N, q)$ is a **reduction** of q -manifolds, or that $(N^{\mathcal{V}}, q^{\mathcal{V}})$ is a reduction of (N, q) , if $i_{\mathcal{V}}$ is a q -embedding and the vector bundles $\mathcal{H}^{\mathcal{V}}$ and \mathcal{H} have the same rank that is to say

$$\mathcal{H}|_{N^{\mathcal{V}}} = \mathcal{H}^{\mathcal{V}}. \tag{121}$$

A map $i_{\mathcal{V}}: (N^{\mathcal{V}}, q^{\mathcal{V}}, h^{\mathcal{V}}) \rightarrow (N, q, h)$ between two metric q -manifolds is called a q -isometry if it is a q -immersion and an isometry. If $i_{\mathcal{V}}$ is injective we will say that $(N^{\mathcal{V}}, q^{\mathcal{V}})$ is a metric q -submanifold of the metric q -manifold (N, q) , and that $i_{\mathcal{V}}$ is a isometric q -embedding. If $i_{\mathcal{V}}$ is moreover a reduction of q -manifolds, we will say that $i_{\mathcal{V}}$ is a **reduction** of metric q -manifolds. A map $i_{\mathcal{V}}: (N^{\mathcal{V}}, F^{\mathcal{V}}) \rightarrow (N, F)$ from a f -manifold $(N^{\mathcal{V}}, F^{\mathcal{V}})$ into another one (N, F) is called a f -immersion if it is an immersion which satisfies

$$(i_{\mathcal{V}})^*F = F^{\mathcal{V}}. \tag{122}$$

When $i_{\mathcal{V}}$ is also injective, we will say that $(N^{\mathcal{V}}, F^{\mathcal{V}})$ is a f -submanifold of (N, F) and that $i_{\mathcal{V}}$ is a f -embedding. Moreover we will then say that $i_{\mathcal{V}}: (N^{\mathcal{V}}, F^{\mathcal{V}}) \rightarrow (N, F)$ is a reduction of f -manifolds or that $(N^{\mathcal{V}}, F^{\mathcal{V}})$ is a reduction of (N, F) if the vector bundles $\mathcal{H}^{\mathcal{V}} = \text{Im } F^{\mathcal{V}}$ and $\mathcal{H} = \text{Im } F$ have

the same rank that is to say $\mathcal{H}|_{N^\vee} = \mathcal{H}^\vee$.

A map $i_\vee: (N^\vee, F^\vee, h^\vee) \rightarrow (N, F, h)$ between two metric f -manifolds is called a **f -isometry** if it is a f -immersion and an isometry. If i_\vee is injective we will say that (N^\vee, F^\vee) is a metric **f -submanifold** of the metric f -manifold (N, F) , and that i_\vee is a **isometric f -embedding**. If i_\vee is moreover a reduction of f -manifolds, we will say that i_\vee is a **reduction** of metric f -manifolds.

Remark 6.10 (120) means that i_\vee sends the splitting $TN^\vee = \mathcal{H}^\vee \oplus \mathcal{V}^\vee$ into the splitting $TN = \mathcal{H} \oplus \mathcal{V}$. In other words, the former is the trace on TN^\vee of the latter. It can be written also in the following form (which suggests the definition of holomorphicity)

$$di_\vee \circ q^\vee = q \circ di_\vee.$$

(122) implies (120), so that a f -immersion is in particular a q -immersion. Moreover a reduction of (metric) f -manifolds is a reduction of the corresponding (metric) q -manifolds. Remark also that (122) means simply that i_\vee is a **f -holomorphic map** from (N^\vee, F^\vee) to (N, F) : $di_\vee \circ F^\vee = F \circ di_\vee$.

Lemma 6.2 • Let $i_\vee: (N^\vee, q^\vee) \rightarrow (N, q)$ be a q -immersion. Then we have

$$(i_\vee)^* \Phi = \Phi^\vee \quad \text{and} \quad (i_\vee)^* R_{\mathcal{V}} = R_{\mathcal{V}^\vee}$$

where Φ^\vee is of course the curvature of the horizontal subbundle \mathcal{H}^\vee .

• Let $i_\vee: (N^\vee, F^\vee) \rightarrow (N, F)$ be a f -immersion. Then we have

$$(i_\vee)^* N_F = N_{F^\vee}$$

Definition 6.18 Let $i_\vee: (N^\vee, h^\vee) \rightarrow (N, h)$ be a isometry between two Riemannian manifolds. Let ∇ be a linear connection on N . The projection on N^\vee of ∇ is the connection ∇^\vee on N^\vee defined by

$$\langle \nabla_X^\vee Y, Z \rangle = \langle \nabla_X \tilde{Y}, Z \rangle \quad \forall X, Z \in TN^\vee, Y \in \mathcal{C}(TN^\vee)$$

where $\tilde{Y} \in \mathcal{C}(TN)$ is some extension of Y .

Let $(E, h) \rightarrow N$ be a Riemannian vector bundle endowed with a linear connection ∇ . Let $N^\vee \subset N$ be an immersed submanifold and $E^\vee \rightarrow N^\vee$ be a vector subbundle of $E|_{N^\vee}$. Then we define analogously the projection of ∇ on E^\vee .

Lemma 6.3 If ∇^\vee is the projection of ∇ then their respective torsion 3-forms are related by

$$\langle T^{\nabla^\vee}(\cdot, \cdot), \cdot \rangle = (i_\vee)^* \langle T^\nabla(\cdot, \cdot), \cdot \rangle$$

Lemma 6.4 • Let $i_\vee: (N^\vee, h^\vee, q^\vee) \rightarrow (N, h, q)$ be a q -isometry.

(i) Then the Levi-Civita connection D^\vee of (N, h^\vee) is the projection on N^\vee of the Levi-Civita connection of (N, h) .

(ii) The following identity holds:

$$D^\vee \Omega_{q^\vee} = (i_\vee)^*(D\Omega_q)$$

• Let $i_\vee: (N^\vee, h^\vee, F^\vee) \rightarrow (N, h, F)$ be a f -isometry. Then $(i_\vee)^* \Omega_F = \Omega_{F^\vee}$.

Theorem 6.5 Let $i_v: (N^v, h^v, F^v) \rightarrow (N, h, F)$ be a f -isometry.

- Then if (N, F, h) is horizontally of type \mathcal{G}_1 , resp. of global type \mathcal{G}_1 , resp. almost reductive, resp. reductive, then so is (N, F^v, h^v) .
- Moreover, if (N, F, h) is horizontally of type \mathcal{G}_1 and almost reductive, then the projection on N^v of its paracharacteristic connection (defined by some $\alpha \in \mathcal{C}(\Lambda^3 \mathcal{V}^*)$) coincides with the paracharacteristic connection of (N^v, F^v, h^v) (defined by $(i_v)^* \alpha \in \mathcal{C}(\Lambda^3 (\mathcal{V}^v)^*)$). In the same way, if (N, F, h) is reductive and of global type \mathcal{G}_1 , then the projection on N^v of its characteristic connection (defined by some $\alpha \in \mathcal{C}(\Lambda^3 \mathcal{V}^*)$) coincides with the characteristic connection of (N^v, F^v, h^v) (defined by $(i_v)^* \alpha$).

Proof. The first assertion follows from lemmas 6.2 and 6.4. The second assertion follows from theorems 6.4 and 6.3, and lemmas 6.2 and 6.4. This completes the proof. \square

Definition 6.19 Let $E \rightarrow N$ be a vector bundle endowed with a linear connection ∇ . Let $N^v \subset N$ be an immersed submanifold and $E^v \rightarrow N^v$ be a vector subbundle of $E|_{N^v}$. We will say that ∇ is reducible in E^v if E^v is ∇ -parallel, that is to say parallel w.r.t. the induced connection on $E|_{N^v}$. Then the induced connection on E^v will be called the reduction of ∇ in E^v .

Remark 6.11 If $E = TN$ and $E^v = TN^v$ then we recover the usual definition of reducibility of linear connection.

Lemma 6.5 If $i_v: L \rightarrow N$ is an immersion and ∇^v the reduction of ∇ then their torsion 2-forms are related by

$$T^{\nabla^v} = (i_v)^* T^\nabla$$

Proposition 6.17 Let $i_v: (N^v, h^v, F^v) \rightarrow (N, h, F)$ be a reduction of metric f -manifolds. Let us suppose that the vertical component D^v of the Levi-Civita connection of (N, h) is reducible in \mathcal{V}^v . Then the Levi-Civita connection of (N, h) is reducible in N^v and its reduction is the Levi-Civita connection D^v of (N, h^v) .

Definition 6.20 Let (N, q, h) be a metric q -manifold. We define the affine space of compatible vertical connection by

$$\text{Con}(\mathcal{V}) = \left\{ D^v - \frac{1}{2} R_{\mathcal{V}}(Y^v, Z^v, X^h) + \mathcal{C}(\Lambda^3 \mathcal{V}^*) \right\}.$$

If (N, q, h) is associated to a metric f -manifold (N, F, h) which is horizontally of type \mathcal{G}_1 and almost reductive (respectively reductive and of global type \mathcal{G}_1) then this set corresponds to the set of metric connection ∇^c on the vertical subbundle \mathcal{V} which can be extended to a paracharacteristic connection (respectively a characteristic connection), (see remark 6.7).

Remark 6.12 If $R_{\mathcal{V}} = 0$, then $D^v \in \text{Con}(\mathcal{V})$.

Definition 6.21 Let $i_v: (N^v, h^v, F^v) \rightarrow (N, h, F)$ be a reduction of metric f -manifolds. We will say that $i_v: (N^v, h^v, F^v) \rightarrow (N, h, F)$ is a **complete reduction** of metric f -manifold if there exists $\nabla^c \in \text{Con}(\mathcal{V})$ which is reducible in \mathcal{V}^v .

More generally, we define analogously a complete reduction of metric q -submersions.

Proposition 6.18 Let $i_v: (N^v, q^v, h^v) \rightarrow (N, q, h)$ be a reduction of metric q -manifolds. If $\nabla^c \in \text{Con}(\mathcal{V})$ is reducible in \mathcal{V}^v , then its reduction is also in $\text{Con}(\mathcal{V}^v)$.

Proof. According to lemmas 6.2 and 6.4 and definition 6.20, the projection of $\text{Con}(\mathcal{V})$ on \mathcal{V}^\vee is $\text{Con}(\mathcal{V}^\vee)$. This completes the proof. \square

Proposition 6.19 *Let (N, q, h) be a metric q -manifold, and $f: L \rightarrow N$ be a map from a Riemann surface in N . Then $\text{Tr}_g(\nabla^c d^v f)$ does not depend on the choice of $\nabla^c \in \text{Con}(\mathcal{V})$.*

Proof. The difference between two elements of $\text{Con}(\mathcal{V})$ is in $\mathcal{C}(\Lambda^3 \mathcal{V}^*)$.

Proposition 6.20 *Let $i_v: (N^\vee, h^\vee, F^\vee) \rightarrow (N, F, h)$ be a complete reduction of metric f -manifolds. Let $\nabla^c \in \text{Con}(\mathcal{V})$ reducible in \mathcal{V}^\vee , and $\nabla^{c,\vee}$ its reduction. Then if (N, F, h) is horizontally of type \mathcal{G}_1 and almost reductive (respectively reductive and of global type \mathcal{G}_1), its paracharacteristic connection (respectively its characteristic connection) extending ∇^c is reducible to the paracharacteristic connection (respectively the characteristic connection) in (N^\vee, h^\vee, F^\vee) extending $\nabla^{c,\vee}$.*

Proof. This follows from theorem 6.5, equation (121), and proposition 6.18. This completes the proof. \square

Therefore, according to proposition 6.19, we deduce that

Proposition 6.21 *Let $i_v: (N^\vee, h^\vee, F^\vee) \rightarrow (N, h, F)$ be a complete reduction of metric f -manifolds. Let us suppose that (N, F, h) is horizontally of type \mathcal{G}_1 and almost reductive. Let $f: L \rightarrow N^\vee$ be a map.*

- *The tension field of f w.r.t. to one paracharacteristic connection ∇ of (N, h, F) coincides with the tension field of f w.r.t. to any paracharacteristic connection of (N^\vee, h^\vee, F^\vee) .*
- *In the same way, if (N, F, h) is reductive and of global type \mathcal{G}_1 , then the tension field of f w.r.t. to one characteristic connection ∇ of (N, h, F) coincides with the tension field of f w.r.t. to any characteristic connection of (N^\vee, h^\vee, F^\vee) .*

6.3 f -connections on fibre bundles

Here, we consider the case where the vertical subbundle is the tangent space of the fibres of a fibration (or more generally a submersion) $\pi: N \rightarrow M$, i.e. $\mathcal{V} = \ker d\pi$. Let us first remark that in this case $R_{\mathcal{V}} = 0$, which leads to immediate simplifications in the preceding results (obtained in section 6.2).

Convention In all the next of the paper, all the submersions $\pi: N \rightarrow M$ are supposed to be surjective (i.e. the open set $\pi(N)$ coincides with M).

6.3.1 Riemannian submersion and metric f -manifolds of global type \mathcal{G}_1 .

Proposition 6.22 *Let $\pi: (N, h) \mapsto (M, g)$ be a Riemannian submersion, over which we consider the natural orthogonal decomposition: $TN = \mathcal{V} \oplus \mathcal{H}$, where $\mathcal{V} = \ker d\pi$ and $\mathcal{H} = \mathcal{V}^\perp$. Denote by D and D^g respectively the Levi-Civita connections of (N, h) and (M, g) , respectively. Let \widetilde{D}^g be the connection on \mathcal{H} defined by the lift of D^g : $\widetilde{D}_A^g B = (d\pi|_{\mathcal{H}})^{-1}(D_A^g \pi_*(B)) \in \mathcal{H}$ for all $A, B \in \mathcal{C}(TN)$.*

Let us suppose that (N, q, h) is of type $\mathcal{V}^2 \mathcal{H}$. Then the horizontal component of the Levi-Civita connection in N is related to \widetilde{D}^g by the following formula:

$$\langle D_A B, H \rangle = \langle \widetilde{D}_A^g B, H \rangle + \frac{1}{2} (\Phi(A, H, B^v) + \Phi(B, H, A^v))$$

for all $A, B \in \mathcal{C}(TN)$ and $H \in \mathcal{C}(\mathcal{H})$.

Proof. Let us set

$$S^M(A, B) = \pi_*(D_A B) - D_A^g(\pi_* B), \quad \forall A, B \in \mathcal{C}(TN).$$

Then it is easy to see that S^M is in fact a tensor, i.e. $S^M \in \mathcal{C}(T^*N \otimes T^*N \otimes \pi^*TM)$. Let $A, B \in TN$ and $H \in \mathcal{H}$, and let us extend these to vector fields, denoted by the same notations, such that the horizontal components of these extension are projectible: there exist vector fields $\bar{A}, \bar{B}, \bar{H}$ on M such that

$$\pi_* A = \bar{A} \circ \pi, \pi_* B = \bar{B} \circ \pi, \pi_* H = \bar{H} \circ \pi.$$

Using the fact that $h|_{\mathcal{H} \times \mathcal{H}} = \pi^*$, the characterization of Levi-Civita yields:

$$\begin{aligned} 2\langle S^M(A, B), \pi_* H \rangle &= 2h(D_A B, H) - 2g(D_A^g(\bar{B}), \bar{H}) \circ \pi \\ &= A.h(B^{\mathcal{H}}, H) + B.h(A^{\mathcal{H}}, H) - H.h(A, B) + h([A, B], H) + h([H, A], B) + h([H, B], A) \\ &\quad - (\bar{A}.g(\bar{B}, \bar{H}) + \bar{B}.g(\bar{A}, \bar{H}) - \bar{H}.g(\bar{A}, \bar{B}) + g([\bar{A}, \bar{B}], H) + g([\bar{H}, \bar{A}], \bar{B}) + g([\bar{H}, \bar{B}], \bar{A})) \circ \pi \\ &= -H \cdot \langle qA, qB \rangle + \langle q[H, A], qB \rangle + \langle q[H, B], qA \rangle \\ &= -\langle D_H^v(qA), qB \rangle - \langle qA, D_H^v(qB) \rangle + \langle q[H, A], qB \rangle + \langle q[H, B], qA \rangle \\ &= \langle T^v(A, H), B^v \rangle + \langle T^v(B, H), A^v \rangle \\ &= -\text{Sym}_{\mathcal{V} \times \mathcal{V}}(T_{|\mathcal{H} \times \mathcal{V} \times \mathcal{V}}^v)(H, A^v, B^v) + \Phi(A^h, H, B^v) + \Phi(B^h, H, A^v) \end{aligned}$$

where we have used in the last line $T_{|\mathcal{H} \times \mathcal{H}}^v = \Phi$ (see proposition 6.5). We conclude by using proposition 6.10. This completes the proof. \square

Remark 6.13 By a direct computation using the characterisation of Levi-Civita, we can prove

$$\langle D_{X^v} Y^h, Z^h \rangle = \frac{1}{2} \Phi(Y^h, Z^h, X^v) + \frac{1}{2} \text{Sym}_{\mathcal{H} \times \mathcal{H}}(\nabla \Omega_{q|\mathcal{H} \times \mathcal{V} \times \mathcal{H}})(Z^h, X^v, Y^h) + \langle [X^v, Y^h], Z^h \rangle$$

Then according to the previous proposition, we obtain

$$\langle \widetilde{D_{X^v} Y^h}, Z^h \rangle = \frac{1}{2} \text{Sym}_{\mathcal{H} \times \mathcal{H}}(\nabla \Omega_{q|\mathcal{H} \times \mathcal{V} \times \mathcal{H}})(Z^h, X^v, Y^h) + \langle [X^v, Y^h], Z^h \rangle$$

Moreover, this equation still holds if we replace D^g by any metric connection ∇^M on M , since then $(D^g - \nabla^M)$ is a horizontal trilinear form.

Proposition 6.23 *Let $\pi: (N, h) \mapsto (M, g)$ be a Riemannian submersion, with the same notations and definitions as in the previous proposition. Let us suppose that some metric connection ∇^c on \mathcal{V} is given, and denote by T^c its vertical connection.*

Then the vertical component of the Levi-Civita connection on N is given by

$$\langle D_A B, V \rangle = \langle \nabla_A^c B^v, V \rangle + \frac{1}{2} (B(A^v, B^v, V) - \Phi(A^h, B^h, V) - R_a^c(A^h, B^v, V) - R_s^c(B^h, V, A^v))$$

where

$$\begin{aligned} B(V_1, V_2, V_3) &= -T^c(V_1, V_2, V_3) - U^c(V_1, V_2, V_3) = -T^c(V_1, V_2, V_3) + T^c(V_1, V_3, V_2) + T^c(V_2, V_3, V_1) \\ R_a^c &= \text{Skew}_{\mathcal{V} \times \mathcal{V}}(T_{|\mathcal{H} \times \mathcal{V} \times \mathcal{V}}^c) \quad \text{and} \quad R_s^c = \text{Sym}_{\mathcal{V} \times \mathcal{V}}(T_{|\mathcal{H} \times \mathcal{V} \times \mathcal{V}}^c). \end{aligned}$$

Proof. Let us set $S^\mathcal{V}(A, B) = q(D_A B) - \nabla_A^c(qB)$ for all $A, B \in \mathcal{C}(TN)$, this defines a element $S^\mathcal{V} \in \mathcal{C}(T^*N \otimes T^*N \otimes \mathcal{V})$. Then let $A, B \in TN$ and $V \in \mathcal{V}$, and let us extend these to vector fields, denoted by the same notations, such that the horizontal components of these extension are projectible: there exist vector fields \bar{A}, \bar{B} on M such that

$$\pi_* A = \bar{A} \circ \pi, \pi_* B = \bar{B} \circ \pi, \pi_* V = 0.$$

Then using the characterisation of the Levi-Civita connection, we have

$$\begin{aligned} 2\langle S^\mathcal{V}(A, B), V \rangle &= 2\langle D_A B, V \rangle - 2\langle \nabla_A^c B^v, V \rangle \\ &= A \cdot \langle B^v, V \rangle + B \cdot \langle A^v, V \rangle - V \cdot \langle A^v, B^v \rangle \\ &\quad - \langle A^v, [B^v, V]^v \rangle + \langle B^v, [V, A]^v \rangle + \langle V, [A, B]^v \rangle - 2\langle \nabla_A^c B^v, V \rangle \\ &\quad - V \cdot g(\pi_* A, \pi_* B) - g(\pi_* A, \pi_* [B, V]) + g(\pi_* B, \pi_* [V, A]) \\ &= \langle \nabla_A^c B^v + \nabla_B^c A^v - 2\nabla_A^c B^v + [A, B]^v, V \rangle + \langle B^v, \nabla_A^c V - \nabla_V^c A^v + [V, A]^v \rangle \\ &\quad + \langle \nabla_B^c V - \nabla_V^c B^v - [B, V]^v, A^v \rangle \\ &= \langle T^c(B, A), V \rangle + \langle T^c(A, V), B^v \rangle + \langle T^c(B, V), A^v \rangle. \end{aligned}$$

Then we complete the proof by using proposition 4.1. \square

Remark 6.14 The results of that is that

$$\begin{aligned} 2(D - \nabla^c)|_{\mathcal{H} \times \mathcal{V} \times \mathcal{V}} &= R_a^c \\ 2(D - \nabla^c)|_{\mathcal{V} \times \mathcal{H} \times \mathcal{V}} &= R_s^c(B^h, V, A^v) \end{aligned}$$

so that $(D - \nabla^c)|_{S(\mathcal{V} \times \mathcal{H})} = 0$ if and only if the reductivity term $T_{|\mathcal{V} \times \mathcal{H}}^c$ vanishes, and in this case the vertical component of the Levi-Civita connection D is given by

$$\langle D_A B, V \rangle = \langle \nabla_A^c B^v, V \rangle + \frac{1}{2} (B(A^v, B^v, V) - \Phi(A^h, B^h, V))$$

and we recover the corresponding relation in homogeneous fibre bundles of theorem 4.2, since in a homogeneous fibre bundle we have $T_{|\mathcal{V} \times \mathcal{H}}^c = 0$. Moreover, from these considerations and proposition 6.10, we deduce the following.

Corollary 6.5 *The symmetric component of the reductivity term, $\text{Sym}_{\mathcal{V} \times \mathcal{V}}(T_{|\mathcal{H} \times \mathcal{V} \times \mathcal{V}}^c)$, is independent of ∇^c and vanishes if and only if (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$. Moreover, the reductivity term $T_{|\mathcal{V} \times \mathcal{H}}^c$ vanishes if and only if (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$ and $(D - \nabla^c)|_{\mathcal{H} \times \mathcal{V} \times \mathcal{V}} = 0$. We will then say that (N, q) is ∇^c -reductive. In particular, if we take $\nabla^c = D^v$, the restriction to \mathcal{V} of the vertical component of Levi-Civita, then the D^v -reductivity means that (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$.*

Applying propositions 6.22, 6.23, we obtain:

Proposition 6.24 *Let $\pi: (N, h) \mapsto (M, g)$ be a Riemannian submersion. Let us suppose that there exists some metric connection ∇^c on \mathcal{V} for which (N, q) is ∇^c -reductive. Then, w.r.t. the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$, the Levi-Civita connection in N admits the following decomposition:*

$$D_A = \begin{pmatrix} \nabla_A^c & 0 \\ 0 & \widetilde{D}_A^g \end{pmatrix} + \frac{1}{2} \begin{pmatrix} B(A^v, \cdot) & -\Phi(A^h, \cdot) \\ \rho(\cdot)A^h & \rho(A^v) \end{pmatrix}.$$

where, let us recall it, ρ is defined by (118). In particular, this decomposition holds for homogeneous fibre bundles.

Remark 6.15 We see that according to this decomposition of Levi-Civita, we have $Dq|_{\mathcal{H} \times \mathcal{H} \times \mathcal{V}} = \frac{1}{2}\Phi$, so that the ∇^c -reductivity implies that (N, q, h) is of type $\mathcal{H}^2\mathcal{V}$. In particular let us take $\bar{\nabla}^c = D^v$ restricted to \mathcal{V} , then the D^v -reductivity means that (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$, and thus we see that the type $\mathcal{V}^2\mathcal{H}$ implies the type $\mathcal{H}^2\mathcal{V}$.

Corollary 6.6 *Let $\pi: (N, h) \mapsto (M, g)$ be a Riemannian submersion, endowed with its canonical orthogonal splitting $TN = \mathcal{V} \oplus \mathcal{H}$. Then if (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$ then it is also of type $\mathcal{H}^2\mathcal{V}$ and thus it is reductive. In particular, if \mathcal{V} can be endowed with a metric connection ∇^c with a vanishing reductivity term $T_{|\mathcal{V} \times \mathcal{H}}^c$, then (N, q, h) is reductive. In particular, a homogeneous fibre bundle is reductive.*

Corollary 6.7 *Let $\pi: (N, h) \mapsto (M, g)$ be a Riemannian submersion, endowed with its canonical orthogonal splitting $TN = \mathcal{V} \oplus \mathcal{H}$. Let us suppose that \mathcal{H} is endowed with an orthogonal complex structure, that is to say N is endowed with a metric f -structure compatible⁶⁴ with the previous splitting.*

Let us suppose that there exists some metric connection ∇^c on \mathcal{V} for which (N, q) is ∇^c -reductive, and that $T_{|\mathcal{V}^3}^c$ is skew-symmetric. Then the following statements are equivalent

- (i) *There exists a characteristic connection on (N, F, h) .*
- (ii) *(N, F, h) is of global type \mathcal{G}_1 .*
- (iii) *The canonical connection ∇^c can be extended to a characteristic connection.*
- (iv) *There exists a Hermitian connection $\nabla^{\mathcal{H}}$ on \mathcal{H} such that $\nabla := \nabla^c \oplus \nabla^{\mathcal{H}}$ has a skew-symmetric torsion.*

In particular, these equivalences hold when $\pi: (N, h) \mapsto (M, g)$ is a homogeneous fibre bundle with a naturally reductive fibre H/K .

Remark 6.16 In other words, if (N, q, h) is of type $\mathcal{V}^2\mathcal{H}$, then the existence of a characteristic connection is equivalent to the global type \mathcal{G}_1 , and in this case, the set of metric connections ∇^c on the vertical subbundle \mathcal{V} which can be extended to a characteristic connection, is the affine space

$$D^v + \mathcal{C}(\Lambda^3\mathcal{V}^*).$$

Proof of corollary 6.7. Since, according to corollary 6.6, the ∇^c -reductivity implies the reductivity, then (i) \Leftrightarrow (ii) according to theorem 6.3. Moreover the equivalence (ii) \Leftrightarrow (iii) follows from corollary 6.4 and remark 6.14. Finally, the equivalence (iii) \Leftrightarrow (iv) is obvious. This completes the proof. \square

We can rewrite the proposition 6.7 for paracharacteristic connections.

Corollary 6.8 *Let $\pi: (N, h) \mapsto (M, g)$ be a Riemannian submersion, endowed with its canonical orthogonal splitting $TN = \mathcal{V} \oplus \mathcal{H}$. Let us suppose that \mathcal{H} is endowed with an orthogonal complex structure, that is to say N is endowed with a metric f -structure compatible⁶⁵ with the previous splitting.*

Let us suppose that there exists some metric connection ∇^c on \mathcal{V} for which (N, q) is ∇^c -reductive, and that $T_{|\mathcal{V}^3}^c$ is skew-symmetric. Then the following statements are equivalent

⁶⁴i.e. $\ker F = \mathcal{V}$ and $\text{Im } F = \mathcal{H}$.

⁶⁵i.e. $\ker F = \mathcal{V}$ and $\text{Im } F = \mathcal{H}$.

(i) There exists a paracharacteristic connection on (N, F, h) .

(ii) (N, F, h) is of horizontal type \mathcal{G}_1 .

(iii) The canonical connection ∇^c can be extended to a paracharacteristic connection.

In particular, these equivalences hold when $\pi: (N, h) \mapsto (M, g)$ is a homogeneous fibre bundle with a naturally reductive fibre H/K .

6.3.2 Reductions of f -submersions

Definition 6.22 We will say that $\pi: (N, q, h) \rightarrow (M, g)$ is a metric q -submersion if $\pi: (N, h) \rightarrow (M, g)$ is a Riemannian submersion and if (N, q, h) is the metric q -manifold defined by the orthogonal spitting $TN = \mathcal{H} \oplus \mathcal{V}$, with $\mathcal{V} = \ker d\pi$ and $\mathcal{H} = \mathcal{V}^\perp$.

Definition 6.23 • A map $\pi: (N, F) \rightarrow M$ from a f -manifold (N, F) to a manifold M is called a **f -submersion** if it is a submersion and $\ker F = \ker d\pi$.

• A map $\pi: (N, F, h) \rightarrow (M, g)$ from a metric f -manifold to a Riemannian manifold is called a **metric f -submersion** if $\pi: (N, h) \rightarrow (M, g)$ is a Riemannian submersion and if $\pi: (N, F) \rightarrow M$ is a f -submersion.

Remark 6.17 Since F is compatible with h , a metric f -submersion is also a metric q -submersion: the splitting defined by F coincides with the orthogonal splitting defined by the Riemannian submersion.

Remark 6.18 A metric f -submersion $\pi: (N, F, h) \rightarrow (M, g)$ is a metric q -submersion $\pi: (N, h) \rightarrow (M, g)$ with a orthogonal complex structure $\bar{J} \in \mathcal{C}(\sigma(\pi^*(TM))) \simeq \mathcal{C}(\Sigma(\mathcal{H}))$.

Definition 6.24 Let $\pi_1: (N_1, F_1) \rightarrow M_1$ and $\pi_2: (N_2, F_2) \rightarrow M_2$ be two f -submersions. Then a map $\Psi: (N_1, F_1) \rightarrow (N_2, F_2)$ is a morphism of f -submersions if it satisfies the following conditions

(i) Ψ is f -holomorphic: $d\Psi \circ F_1 = F_2 \circ d\Psi$

(ii) Ψ is a morphism of submersion: there exists a map $\psi: M_1 \rightarrow M_2$ such that $\pi_2 \circ \Psi = \psi \circ \pi_1$.

If $M_1 = M_2$ and $\psi = \text{Id}$, we will say that Ψ is a morphism of f -submersion over M .

Definition 6.25 • A morphism of metric f -submersion is a morphism of f -submersion which is also an isometry.

• A reduction of f -submersion is a morphism of f -submersion which is also an injective immersion.

• A reduction of metric f -submersion is a morphism of metric f -submersion which is injective. We will then say that it is a complete reduction of metric f -submersion if it is a complete reduction of metric f -manifolds.

Definition 6.26 • We will say that $\pi: (N, F) \rightarrow M$ is a f -fibration if it is a f -submersion and a fibration.

• We will say that $\pi: (N, F, h) \rightarrow (M, g)$ is a homogeneous fibre f -bundle if $\pi: (N, h) \rightarrow (M, g)$ is a homogeneous fibre bundle and $\pi: (N, F, h) \rightarrow (M, g)$ is a metric f -submersion.

Proposition 6.25 *Let $\pi: N \rightarrow M$ be a homogeneous fibre bundle. Then a homogeneous fibre subbundle $\pi: N^\vee \rightarrow M$, of $\pi: N \rightarrow M$ defines a reduction of homogeneous fibre bundle⁶⁶ if and only if the inclusion $i_\vee: N^\vee \rightarrow N$ is a reduction of q -manifolds. Moreover, a reduction of homogenous fibre bundle is always a complete reduction.*

Proof. We have seen in subsection 4.2.3 that $\pi: N^\vee \rightarrow M$ is a reduction of homogeneous fibre bundle if and only if the equation (121) holds. This proves the first assertion. The second one follows from the fact that according to subsection 4.2.3, the connection ∇^c always leaves invariant the subbundle \mathcal{V}^\vee . This completes the proof. \square

Therefore, we obtain, more particularly,

Proposition 6.26 *In the same situation as in proposition 6.25, let us suppose that the homogeneous fibre bundle $\pi: N \rightarrow M$ admits an orthogonal complex structure $\bar{J} \in \mathcal{C}(\Sigma(\pi^*TM))$ defining then a metric f -structure and therefore a structure of homogeneous fibre f -bundle. Then the homogeneous fibre subbundle $\pi^\vee: N^\vee \rightarrow M$ inherits also a structure of homogeneous fibre f -subbundle defined by $\bar{J}^\vee = (i_\vee)^*\bar{J} \in \mathcal{C}(\Sigma((\pi^\vee)^*TM))$. If this subbundle defines a reduction of homogeneous fibre bundle, then it defines also a complete reduction of homogeneous fibre f -subbundle.*

Definition 6.27 *Let $\pi: (N, F, h) \mapsto (M, g)$ be a homogeneous fibre f -bundle with a naturally reductive fibre H/K . Suppose that⁶⁷ (N, F, h) is horizontally \mathcal{G}_1 (respectively of global type \mathcal{G}_1). Then we call the canonical paracharacteristic (respectively characteristic) connection of the homogeneous fibre f -bundle (N, F, h) its paracharacteristic (respectively characteristic) connection extending its vertical canonical connection⁶⁸ ∇^c .*

Proposition 6.20 yields the following.

Proposition 6.27 *Let $\pi: (N, h) \rightarrow (M, g)$ be a homogeneous fibre bundle with a naturally reductive fibre H/K , and $\pi^\vee: N^\vee \rightarrow M$ a homogeneous fibre subbundle such that the inclusion $i_\vee: N^\vee \rightarrow N$ is a reduction of homogeneous fibre bundle⁶⁹. Suppose also that is given an orthogonal complex structure $\bar{J} \in \mathcal{C}(\Sigma(\pi^*TM))$ defining then structures of homogeneous fibre f -bundles: $\pi: (N, F, h) \rightarrow (M, g)$ and $\pi^\vee: (N^\vee, F^\vee, h^\vee) \rightarrow (M, g)$. Suppose that (N, F, h) is horizontally of type \mathcal{G}_1 (of global type \mathcal{G}_1) then the canonical paracharacteristic (respectively characteristic) connection of (N, F, h) is reducible in (N^\vee, F^\vee, h^\vee) to the canonical paracharacteristic (respectively characteristic) connection of (N^\vee, F^\vee, h^\vee) .*

6.3.3 Horizontally Kähler f -manifolds and horizontally projectible f -submersions.

Definition 6.28 *Let (N, F, h) be a metric f -manifold. We will say that (N, F, h) is*

- *horizontally Hermitian if $N_{\bar{J}} = 0$.*
- *horizontally Kähler if $DF|_{\mathcal{H}^3} = 0$.*

Lemma 6.6 *Let $\pi: (N, F, h) \rightarrow (M, g)$ be a metric f -submersion. Let us suppose that (N, F, h) is reductive. Then it is horizontally Kähler if and only if*

$$D_H^g \bar{J} = 0, \forall H \in \mathcal{H}.$$

⁶⁶See definition 4.9

⁶⁷Keep in mind that a homogeneous fibre bundle is always reductive, according to corollary 6.6.

⁶⁸See section 4.2.1.

⁶⁹in the sens of definition 4.9

where, as usual D^g denotes the Levi-Civita connections of g .

Proof. This follows from proposition 6.22. \square

Theorem 6.6 *Let $\pi: (N, F, h) \rightarrow (M, g)$ be a metric f -submersion. Let us suppose that (N, F, h) is reductive and horizontally Kähler. Then (N, F, h) is horizontally Hermitian: $N_{\bar{J}} = 0$. In particular, (N, F, h) is horizontally of type \mathcal{G}_1 . Therefore (N, F, h) admits a paracharacteristic connection (unique up to $\mathcal{C}(\Lambda^3\mathcal{V}^*)$).*

Furthermore, let $\bar{T} \in \mathcal{C}(\Lambda^3\mathcal{H}^)$ be the unique horizontal 3-form, such that the torsion T of any metric f -connection ∇ in N with a skew-symmetric component $T_{\mathcal{H}^3}$, satisfies $T_{\mathcal{H}^3} = \bar{T}$ (see proposition 6.13). Then we have $\bar{T} = 0$, so that for any such connection ∇ , we have $\langle \nabla_{X^h} Y^h, Z^h \rangle = \langle \widetilde{D^g_{X^h}} Y^h, Z^h \rangle$, $\forall X^h, Y^h, Z^h \in \mathcal{C}(\mathcal{H})$.*

Moreover, we have

$$N(X^v, Y^h, Z^h) = \langle (D^g_{X^v} \bar{J}) \bar{J} Y^h, Z^h \rangle$$

where $X, Y, Z \in TN$.

Proof. According to proposition 6.15, $DF|_{\mathcal{H}^3} = 0$ implies $N_{F|_{\mathcal{H}^3}} = 0$. Moreover, $DF|_{\mathcal{H}^3} = 0$ implies also $d\Omega_{F|_{\mathcal{H}^3}} = 0$ which implies $\bar{T} = 0$ since $d\Omega_{F|_{\mathcal{H}^3}} = \bar{J} \odot \bar{T}$.

Concerning the last assertion, it can be proved by applying propositions 6.15 and 6.24. Or more simply by using directly remark 6.13. This completes the proof. \square

Definition 6.29 *Let $\pi: (N, F, h) \rightarrow (M, g)$ be a metric f -submersion. Let us suppose that (N, F, h) is a reductive and horizontally \mathcal{G}_1 . A metric connection $\bar{\nabla}$ will be said to be the **canonical connection** on M (w.r.t. the f -submersion) if its torsion \bar{T} satisfies the equation $\widetilde{\bar{T}} = T|_{\mathcal{H}^3}$ where T is the torsion of any paracharacteristic connection (see proposition 6.13) and $\widetilde{\bar{T}}$ is the lift in \mathcal{H} of \bar{T} . If such a connection exists, we will say that the metric f -submersion is **horizontally projectible**.*

Remark 6.19 The canonical connection is unique when it exists (since it is metric and its torsion is given). Moreover it has a skew-symmetric torsion. The metric f -submersion is **horizontally projectible** if and only if the horizontal 3-form $T|_{\mathcal{H}^3}$ is projectible to a 3-form on N .

Proposition 6.28 *Let $\pi: (N, F, h) \rightarrow (M, g)$ be a metric f -submersion. Let us suppose that there exists a metric connection $\bar{\nabla}$ geodesically equivalent to the Levi-Civita connection, i.e. with a skew-symmetric torsion, such that*

$$\bar{\nabla}_H \bar{J} = 0, \quad \forall H \in \mathcal{H}.$$

Then (N, F, h) is horizontally of type \mathcal{G}_1 , and horizontally projectible, $\bar{\nabla}$ being then the canonical connection of M .

Proof. Any metric connection ∇ on N which satisfies

$$\langle \nabla_{H_1} H_2, H_3 \rangle = \langle \widetilde{\bar{\nabla}}_{H_1} H_2, H_3 \rangle,$$

satisfies $\nabla F|_{\mathcal{H}^3} = 0$ and $T|_{\mathcal{H}^3} = \widetilde{\bar{T}}$. Then proposition 6.13 allows us to conclude. This completes the proof. \square

In particular, we have

Corollary 6.9 *Let $\pi: (N, F, h) \rightarrow (M, g)$ be a metric f -submersion. Let us suppose that (N, F, h) is a reductive and horizontally kähler. Then it is horizontally projectible and the canonical connection on M is its Levi-Civita connection D^g .*

Proof. This follows immediately from lemma 6.6 and proposition 6.28. More directly, according to theorem 6.6, we have $T|_{\mathcal{H}^3} = 0$ and therefore this 3-form is projectible. This completes the proof. \square

6.3.4 The example of a naturally reductive homogeneous space

Proposition 6.29 *Let $N = G/K$ be a Riemannian homogeneous space and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{n}$ a reductive decomposition of \mathfrak{g} . Let us suppose that \mathfrak{n} admits an $\text{Ad}K$ -invariant decomposition $\mathfrak{n} = \mathfrak{m} \oplus \mathfrak{p}$, defining then a splitting $TN = \mathcal{H} \oplus \mathcal{V}$, where $\mathcal{H} = [\mathfrak{m}]$ and $\mathcal{V} = [\mathfrak{p}]$. Let us suppose that there exists on \mathfrak{m} an $\text{Ad}K$ -invariant complex structure \bar{J}_0 , defining then a f -structure F on N . Then for any G -invariant metric h for which \bar{J}_0 and the decomposition $\mathfrak{n} = \mathfrak{m} \oplus \mathfrak{p}$ are orthogonal (such a metric always exists), (N, F, h) is a metric f -manifold.*

Furthermore, let us suppose that $N = G/K$ is naturally reductive, and that one can choose a naturally reductive metric h as above⁷⁰, then (N, F, h) is reductive and of global type \mathcal{G}_1 . Moreover, the canonical connection ∇^0 is a characteristic connection. Therefore, in this case its characteristic connection ∇ has a parallel torsion $\nabla T = 0$.

Proof. The naturally reductivity means exactly that the torsion of the canonical connection is skew-symmetric. This completes the proof. \square

Proposition 6.30 *Let us consider the situation described by proposition 6.29. Then the horizontal curvature and its linear representation, and the horizontal curvature operator are given (in terms of their lifts in G) by*

$$\begin{aligned}\widetilde{\Phi}(X, Y) &= -[X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}} \\ \widetilde{\rho}(V) &= -\text{ad}_{\mathfrak{m}} V_{\mathfrak{p}} \\ \widetilde{R}(X, Y)Z &= \text{ad}_{\mathfrak{m}} [X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}}\end{aligned}$$

6.3.5 The example of the twistor space $\mathcal{Z}_{2k}^{\alpha}(M)$.

Let (M, g) be a Riemannian manifold endowed with a metric connection $\bar{\nabla}$. Then we consider the homogeneous fibre bundle $\pi: (\mathcal{Z}_{2k}^{\alpha}(M), h) \rightarrow (M, g)$ defined by the Riemannian vector bundle $(TM, g, \bar{\nabla})$. (See sections 4.1.2 (example 4.3) and 4.3.3.) Let us consider also its canonical $2k$ -structure $\mathcal{J} \in \mathcal{C}(\mathcal{Z}_{2k}^{\alpha}(\pi^*TM))$, to which corresponds the orthogonal complex structure $\underline{\mathcal{J}} \in \mathcal{C}(\Sigma(\pi^*TM)) \cong \mathcal{C}(\Sigma(\pi^*\mathcal{H}))$. This complex structure defines then a metric f -structure \mathcal{F} on $\mathcal{Z}_{2k}^{\alpha}(M)$. Then the twistor bundle $(\mathcal{Z}_{2k}^{\alpha}(M), \mathcal{F}, h)$ is a reductive metric f -manifold, more precisely a homogeneous fibre f -bundle.

Proposition 6.31 *Let (M, g) be a Riemannian manifold endowed with a metric connection $\bar{\nabla}$. Then we have*

$$\bar{\nabla}_H \underline{\mathcal{J}} = 0, \quad \forall H \in \mathcal{H}.$$

⁷⁰which means that denoting by $G(\mathfrak{n})$, the compact subgroup in $GL(\mathfrak{n})$ generated by $\Lambda_{\mathfrak{n}}(\mathfrak{n}) := \{[\text{ad}_{\mathfrak{n}}(X)]_{\mathfrak{n}}, X \in \mathfrak{n}\} \subset \mathfrak{gl}(\mathfrak{n})$, and by $\langle G(\mathfrak{n}), I_0 \rangle$ the subgroup generated by $G(\mathfrak{n})$ and $I_0 := \bar{J}_0 \oplus -\text{Id}_{\mathfrak{p}}$, then $\langle G(\mathfrak{n}), I_0 \rangle / G(\mathfrak{n})$ is compact.

Proof. We have the following sequence of morphisms of bundle over M which are $SO(2n)$ -invariant projections on the fibres:

$$Q = \mathcal{SO}(TM) \xrightarrow{\pi_{\mathcal{Z}}} Q/\mathbb{U}_0(J_0^\alpha) = \mathcal{Z}_{2k}^\alpha(M) \xrightarrow{\mathcal{P}} \Sigma(M) = Q/\mathbb{U}(J_0^\alpha)$$

In particular, the structure of homogeneous fibre bundle of $N_\Sigma = \Sigma(M)$ (defined by $\overline{\nabla}$) if the image by the projection \mathcal{P} of the structure of homogeneous fibre bundle of $N_{\mathcal{Z}} = \mathcal{Z}_{2k}^\alpha(M)$ (defined by $\overline{\nabla}$):

$$TQ = \mathcal{H}^Q \oplus \mathcal{V}^Q \xrightarrow{(\pi_{\mathcal{Z}})_*} TN_{\mathcal{Z}} = \mathcal{H}^{\mathcal{Z}} \oplus \mathcal{V}^{\mathcal{Z}} \xrightarrow{(\mathcal{P})_*} TN_\Sigma = \mathcal{H}^\Sigma \oplus \mathcal{V}^\Sigma.$$

In particular we have $\overline{\nabla} \underline{\mathcal{J}} = \mathcal{P}_*(\overline{\nabla} \mathcal{J})$ and therefore

$$\overline{\nabla}_H \underline{\mathcal{J}} = 0, \quad \forall H \in \mathcal{H},$$

according to theorem 4.15 (i). This completes the proof. \square

Theorem 6.7 *Let (M, g) be a Riemannian manifold endowed with a metric connection $\overline{\nabla}$. Let us suppose that $\overline{\nabla}$ is geodesically equivalent to the Levi-Civita connection, i.e. it admits a skew-symmetric torsion. Then the twistor bundle $(\mathcal{Z}_{2k}^\alpha(M), \mathcal{F}, h)$ is horizontally of type \mathcal{G}_1 , and horizontally projectible, $\overline{\nabla}$ being the canonical connection of M .*

Proof. This follows immediately from proposition 6.31 and proposition 6.28. \square

Corollary 6.10 *Let (M, g) be a Riemannian manifold endowed with a metric connection $\overline{\nabla}$. If $\overline{\nabla} = D^g$ is the Levi-Civita connection, then $(\mathcal{Z}_{2k}^\alpha(M), \mathcal{F}, h)$ is horizontally Kähler.*

Proof. This follows immediately from proposition 6.31 and lemma 6.6. \square

6.3.6 The example of the twistor space $\mathcal{Z}_{2k,j}^\alpha(M, J_j)$.

Let (M, g) be a Riemannian manifold endowed with a metric connection $\overline{\nabla}$ and J_j a section of $(\mathcal{Z}_{2k}^\alpha(M))^j$. Then we consider the homogeneous fibre bundle $\pi: (\mathcal{Z}_{2k,j}^\alpha(M, J_j), h) \rightarrow (M, g)$ defined by the Riemannian vector bundle $(TM, g, \overline{\nabla}^{[j]})$. We denote by \mathcal{F}_j the canonical f -structure defined as above in 6.3.5. Then the twistor bundle $(\mathcal{Z}_{2k,j}^\alpha(M, J_j), \mathcal{F}_j, h)$ is a reductive metric f -manifold, more precisely a homogeneous fibre f -bundle.

Theorem 6.8 *Let (M, g) be a Riemannian manifold endowed with a metric connection $\overline{\nabla}$ and J_j a section of $(\mathcal{Z}_{2k}^\alpha(M))^j$. Let us suppose that $\overline{\nabla}^{[j]}$ is geodesically equivalent to the Levi-Civita connection, i.e. it admits a skew-symmetric torsion.*

• *Then the twistor bundle $(\mathcal{Z}_{2k,j}^\alpha(M, J_j), \mathcal{F}_j, h)$ is horizontally of type \mathcal{G}_1 , and horizontally projectible, $\overline{\nabla}^{[j]}$ being the canonical connection of M .*

• *Furthermore, if $\overline{\nabla} J_j = 0$ so that $(\mathcal{Z}_{2k,j}^\alpha(M, J_j), \mathcal{F}_j, h) \rightarrow (M, g)$ is a reduction of $(\mathcal{Z}_{2k}^\alpha(M), \mathcal{F}, h) \rightarrow (M, g)$, the paracharacteristic connection of $(\mathcal{Z}_{2k}^\alpha(M), \mathcal{F}, h)$ is reducible in $(\mathcal{Z}_{2k,j}^\alpha(M, J_j), \mathcal{F}_j, h)$ and its reduction is the paracharacteristic connection of $(\mathcal{Z}_{2k,j}^\alpha(M, J_j), \mathcal{F}_j, h)$.*

Proof. For the first point, proceed as in 6.3.5. For the second point, apply proposition 6.27. \square

Corollary 6.11 *Let (M, g) be a Riemannian manifold endowed with a metric connection $\overline{\nabla}$ and J_j a section of $(\mathcal{Z}_{2k}^\alpha(M))^j$. If in particular $\overline{\nabla} = D^g$ is the Levi-Civita connection, then $(\mathcal{Z}_{2k,j}^\alpha(M, J_j), \mathcal{F}_j, h)$ horizontally Kähler.*

Proof. Proceed as in 6.3.5. \square

6.3.7 The reduction of homogeneous fibre bundle $\mathfrak{J}_{J_0} : G/G_0 \hookrightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$.

We consider the situation described by 4.4.1 and use the same notations.

We have seen in that $\mathfrak{J}_{J_0} : G/G_0 \longrightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$ is a reduction of homogeneous fibre bundles. Moreover by definition of \mathfrak{J}_{J_0} and \mathcal{J} , this is also an f -immersion. Therefore, since $(G/G_0, F, h)$ and $\mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2), \mathcal{F}_2, h_2$ are both metric f -manifolds, it follows that \mathfrak{J}_{J_0} is a reduction of homogeneous fibre f -bundle. Moreover, we suppose that the G -invariant metric g on G/H , induced⁷² by the metric h on G/G_0 , is naturally reductive. This is the case in particular if h is naturally reductive.

Therefore according to proposition 6.27, we have

Proposition 6.32 *The canonical paracharacteristic connection of $\mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$ is reducible in G/G_0 and its reduction is the canonical connection of G/G_0 .*

6.4 Stringy Harmonic maps in f -manifolds.

6.4.1 Definitions

We have defined the notion of stringy harmonic maps in the context of almost complex manifolds (endowed with a linear connection) and we have seen that it corresponds to a generalisation of harmonic maps. Now, we will extend this notion of stringy harmonicity to f -manifolds endowed with a linear connection. Indeed, the preliminary study of the maximal determined system done in section 2.4, leads us to introduce the following generalisation of (stringy) harmonic maps.

Definition 6.30 *Let (N, F) be a f -manifold with ∇ a linear connection. Then we will say that a map $f : L \rightarrow N$ from a Riemann surface into N is stringy harmonic if it is solution of the stringy harmonic maps equation:*

$$-\tau_g(f) + (F \bullet T)_g(f) = 0$$

Definition 6.31 *Let (N, F) be a f -manifold with ∇ a linear connection. Then we will say that a map $f : L \rightarrow N$ from a Riemann surface into N is \star -stringy harmonic if it is solution of the modified stringy harmonic maps equation:*

$$-\tau_g(f) + (F \star T)_g(f) = 0.$$

Proposition 6.33 *Let (N, F) be a f -manifold with ∇ a linear f -connection. Let us suppose that $R_{\mathcal{V}} = 0$. Then, setting $\bar{T} = T_{\mathcal{H}^2}^{\mathcal{H}}$, the equation of stringy harmonicity, w.r.t. ∇ , for maps $f : L \rightarrow N$ is written*

$$\begin{cases} - * \tau^v(f) + \frac{1}{2} \Phi(df \wedge \bar{J}df) = 0 \\ - * \tau^h(f) + \frac{1}{2} (\bar{J} \cdot \bar{T})(df \wedge df) + \frac{1}{2} N_F(d^v f \wedge \bar{J}d^h f) = 0 \end{cases}$$

whereas the equation of \star -stringy harmonicity is written:

$$\begin{cases} - * \tau^v(f) + \frac{1}{2} \Phi(df \wedge \bar{J}df) = 0 \\ - * \tau^h(f) + \frac{1}{2} (\bar{J} \star \bar{T})(df \wedge df) + \frac{1}{2} N_F(d^v f \wedge \bar{J}d^h f) = 0 \end{cases}$$

⁷¹We denote the metric in $\mathcal{Z}_{2k,2}^{\alpha_0}(G/H, J_2)$ by h_2 to differentiate it from the metric in G/G_0 . See the convention of notations in 4.4.1.

⁷²I.e. denoting by $\langle \cdot, \cdot \rangle$ the inner product on \mathfrak{n} defining h , then g is defined by $\langle \cdot, \cdot \rangle|_{\mathfrak{m} \times \mathfrak{m}}$, see 4.3.1.

where $\tau^v(f)$, $\tau^h(f)$ are respectively the vertical and horizontal component of the tension field w.r.t. ∇ (which coincide also with the vertical and horizontal tension fields respectively, since ∇ preserves the splitting $TN = \mathcal{V} \oplus \mathcal{H}$). Moreover, as usual, $*$ is the Hodge operator for some Hermitian metric on L .

Proof. Use theorem 6.2. □

Remark 6.20 We see that stringy harmonicity in (N, F) can be viewed as a kind of coupling between some equation of \bar{J} -stringy harmonicity and the equation of vertical harmonicity. We will come back to this in details elsewhere [36, 37].

Proposition 6.34 *Let $i_v: (N^v, h^v, F^v) \rightarrow (N, F, h)$ be a complete reduction of metric f -manifolds. Let $\nabla^c \in \text{Con}(\mathcal{V})$ reducible in \mathcal{V}^v , and $\nabla^{c,v}$ its reduction. Let us suppose that (N, F, h) is horizontally of type \mathcal{G}_1 and almost reductive. Let $f: L \rightarrow N^v$ be a map.*

- *The (\star) -stringy harmonicity of f w.r.t. to the paracharacteristic connection ∇ of (N, h, F) extending ∇^c , is equivalent to the (\star) -stringy harmonicity of f w.r.t. to the paracharacteristic connection of (N^v, h^v, F^v) extending $\nabla^{c,v}$.*

- *In the same way, if (N, F, h) is reductive and of global type \mathcal{G}_1 , then the (\star) -stringy harmonicity of f w.r.t. to the characteristic connection ∇ of (N, h, F) extending ∇^c , is equivalent to the (\star) -stringy harmonicity of f w.r.t. to the characteristic connection of (N^v, h^v, F^v) extending $\nabla^{c,v}$.*

In particular, these properties hold when $i_v: (N^v, h^v, F^v) \rightarrow (N, F, h)$ is a reduction of homogeneous fibre bundle as described in proposition 6.27.

Proof. This results immediately from proposition 6.20 and lemma 6.5. This completes the proof. □

Proposition 6.35 *Let $i_v: (N^v, h^v, F^v) \rightarrow (N, h, F)$ be a complete reduction of metric f -manifolds. Let us suppose that (N, F, h) is horizontally of type \mathcal{G}_1 and almost reductive. Let $f: L \rightarrow N^v$ be a map.*

- *The (\star) -stringy harmonicity of f w.r.t. to one paracharacteristic connection ∇ of (N, h, F) is equivalent to the (\star) -stringy harmonicity of f w.r.t. to any paracharacteristic connection of (N^v, h^v, F^v) .*

- *In the same way, if (N, F, h) is reductive and of global type \mathcal{G}_1 , then the (\star) -stringy harmonicity of f w.r.t. to one characteristic connection ∇ of (N, h, F) is equivalent to the (\star) -stringy harmonicity of f w.r.t. to any characteristic connection of (N^v, h^v, F^v) .*

Proof. We deduce from proposition 6.21, theorem 6.5 and equation (121), that the terms in $\nabla^* df + (F \bullet T)_g(f)$ which could be no reducible in N^v are only those corresponding to the vertical terms $(F \bullet T)_{\mathcal{H} \times \mathcal{V} \times \mathcal{V}}$ and $(F \bullet T)_{\mathcal{V} \times \mathcal{H} \times \mathcal{V}}$ equal to $R_{\mathcal{V}}(Y^v, Z^v, \bar{J}X^h)$ and $R_{\mathcal{V}}(Z^v, X^v, \bar{J}Y^h)$ respectively. These terms do not depend on the choice of ∇ (and ∇^v for the corresponding terms in N^v) and from proposition 6.34, we know that for a convenient choice of ∇ and ∇^v , these terms are reducible, i.e. if B (resp. B^v) is one of these terms then $B_g^v(f) = B_g(f)$. This completes the proof. □

6.4.2 The closeness of the 3-forms $F \bullet T$ and $F \star T$.

Let us see under which conditions on a reductive metric f -manifold of global type \mathcal{G}_1 , the 3-forms $F \bullet T$ and $F \star T$ defined by one characteristic connection are closed.

Proposition 6.36 *Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . Let ∇ be some characteristic connection on N . Let us consider the following statements:*

- (i) The 3-forms $H = F \bullet T$ is closed.
- (ii) The 3-forms $H^* = F \star T$ is closed.
- (iii) The horizontal 3-form $F \cdot N_F$ is closed.
- (iv) The 3-form $F \cdot N_F - \frac{1}{2}F \odot (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}}))$ is closed.
- (v) The 3-form $\frac{1}{2}(F \cdot N_F - F \odot (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}})))$ is closed.
- (vi) The 3-form $F \odot (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}}))$ is closed.

Then we have the following equivalences: (i) \Leftrightarrow (iv) and (ii) \Leftrightarrow (v). Moreover any couple of statements in $\{(i), (ii), (iii)\}$ imply the third one. Any couple of statements in $\{(iii), (iv), (v), (vi)\}$ imply the two others. In other words identifying (i) with (iv) and (ii) with (v) (since they are respectively equivalent) in the set of six elements $\{(i), \dots, (vi)\}$, then in the obtained set of four elements, any couple of two statements is equivalent to the couple of the two others.

Proof. This follows from the following identities:

$$\begin{aligned} \frac{1}{2}F \cdot N_F &= F \bullet T - F \star T \\ F \bullet T &= F \cdot N_F - \frac{1}{2}F \odot (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}})) - d\Omega_F \quad (\text{proposition 6.14}) \\ F \star T &= F \bullet T - \frac{1}{2}F \cdot N_F = \frac{1}{2}F \cdot N_F - \frac{1}{2}F \odot (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}})) - d\Omega_F. \end{aligned}$$

□

Definition 6.32 Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . We will say that (N, F, h) **has a closed stringy structure** if the 3-forms $F \cdot N_F$ and $F \odot (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}}))$ are closed. This is equivalent to say that the two 3-forms H and H^* are closed.

Characteristic connections with parallel torsion. In this paragraph, (N, F, h) is a reductive metric f -manifold of global type \mathcal{G}_1 . Let us suppose that one of its characteristic connections, ∇ , has a parallel torsion $\nabla T = 0$.

Now, let us compute all the components in each space $(\Lambda^p \mathcal{H}^*) \wedge (\Lambda^q \mathcal{V}^*)$, $p + q = 4$, of the exterior differential $d\alpha$ of the different 3-forms α we have considered in proposition 6.36.

Lemma 6.7 Let α be a ∇ -parallel 3-form. Then

$$d\alpha(X, Y, V, Z) = \mathfrak{S}_{X, Y, Z} \alpha(T(V, Z), X, Y) + \alpha(T(X, Y), V, Z)$$

In particular, if α is horizontal then

$$\begin{aligned} d\alpha|_{\Lambda^4 \mathcal{V}} &= 0 \\ d\alpha|_{(\Lambda^3 \mathcal{V}) \wedge \mathcal{H}} &= 0 \\ d\alpha|_{(\Lambda^2 \mathcal{V}) \wedge \mathcal{H}^2}(V_0, V_1, H_2, H_3) &= \alpha(R_{\mathcal{V}}(V_0, V_1), H_2, H_3) \\ d\alpha|_{\mathcal{V} \wedge (\Lambda^3 \mathcal{H})}(V_0, H_1, H_2, H_3) &= \mathfrak{S}_{i, j, k} \alpha(T^{\mathcal{H}}(V_0, H_i), H_j, H_k) \end{aligned}$$

Notation We set $\mathcal{S}(\mathcal{H}, \mathcal{V}) = \mathcal{S}(\mathcal{H} \times \mathcal{H} \times \mathcal{V}) \oplus \mathcal{S}(\mathcal{H} \times \mathcal{V} \times \mathcal{V})$. Then for any $B \in \mathcal{T}$, we set $\dot{B} = B|_{\mathcal{S}(\mathcal{H}, \mathcal{V})}$. Let us remark that we have

$$-F \circlearrowleft \dot{T} = d\Omega_{F|_{\mathcal{S}(\mathcal{H}, \mathcal{V})}} \quad \text{and} \quad -\bar{J} \circlearrowleft \bar{T} = d\Omega_{F|_{\mathcal{H}^3}}.$$

Lemma 6.8 *We have the following identity:*

$$-d(\bar{J} \circlearrowleft \bar{T}) = d(F \circlearrowleft \dot{T})$$

Proof. We have

$$0 = -d(d\Omega_F) = d(\bar{J} \circlearrowleft \bar{T}) + d(F \circlearrowleft \dot{T}).$$

This completes the proof. \square

Proposition 6.37 *The following identity holds:*

$$d(F \cdot N_F)|_{\mathcal{H}^4} = d(F \circlearrowleft (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}})))|_{\mathcal{H}^4}$$

Therefore the following statements are equivalent

- (i) $dH|_{\mathcal{H}^4} = 0$
- (ii) $dH^*|_{\mathcal{H}^4} = 0$
- (iii) $d(F \cdot N_F)|_{\mathcal{H}^4} = 0$
- (iv) $\mathfrak{S}_{X,Y,Z} \bar{A}(X, Y)Z = 0$
- (v) $\mathfrak{S}_{X,Y,Z} \bar{A}^{(-)}(X, Y)Z = 0$.

We will say that \bar{J} is a cyclic derivation of the horizontal curvature when (iv) holds. \bullet

Proof. We consider 4-forms on \mathcal{H} and denote by (X, Y, V, Z) the variable in \mathcal{H}^4 . Proceeding as in the proof of proposition 5.18, we obtain according to lemma 6.7

$$\begin{aligned} d(\bar{J} \circlearrowleft \bar{T}) &= \mathfrak{S}_{X,Y,Z} (\bar{J} \circlearrowleft \bar{T})(X, Y, \bar{T}(V, Z)) + (\bar{J} \circlearrowleft \bar{T})(V, Z, \bar{T}(X, Y)) \\ &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} \langle (\bar{J} \circlearrowleft \bar{T})(X, Y), \bar{T}(V, Z) \rangle \\ &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4\langle \bar{T}^{--}(X, Y), \bar{J}\bar{T}^{++}(V, Z) \rangle + 2\langle \bar{T}^{--}(X, Y), \bar{J}\bar{T}^{1,1}(V, Z) \rangle - 2\langle \bar{T}^{++}(X, Y), \bar{J}\bar{T}^{1,1}(V, Z) \rangle. \end{aligned}$$

Moreover, using the same arguments as in the end of the proof of proposition 5.18, we can conclude that the two last terms inside the sum vanish. More precisely, using the fact that $\bar{J}\bar{T}^{--}$ is a 3-form, we can write the second term inside the sum in two different forms which have different types in $\otimes^4 \mathcal{H}^*$, therefore, this terms vanishes. Idem for the third term, by using the corollary 5.2 applied to \bar{T}^{++} . Therefore

$$d(\bar{J} \circlearrowleft \bar{T}) = \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4\langle \bar{T}^{--}(X, Y), \bar{J}\bar{T}^{++}(V, Z) \rangle. \quad (123)$$

Furthermore, we also have according to lemma 6.7

$$\begin{aligned} d(F \cdot N_F) &= \mathfrak{S}_{X,Y,Z} (F \cdot N_F)(X, Y, \bar{T}(V, Z)) + (F \cdot N_F)(V, Z, \bar{T}(X, Y)) \\ &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4\langle \bar{J}\bar{T}^{--}(X, Y), \bar{T}(V, Z) \rangle \\ &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4\langle \bar{J}\bar{T}^{--}(X, Y), \bar{T}^{**}(V, Z) \rangle \\ &= \mathfrak{S}_{X,Y,Z} \sum_{(X,Y) \rightleftharpoons (V,Z)} 4\langle \bar{J}\bar{T}^{--}(X, Y), \bar{T}^{++}(V, Z) \rangle \end{aligned}$$

Hence, we obtain

$$d(F \cdot N_F) = -d(\bar{J} \oslash \bar{T}).$$

This proves the first assertion according to lemma 6.8. Moreover, using lemma 6.8, we compute that

$$d(F \oslash \bar{T})|_{\mathcal{H}^4} = \mathfrak{S}_{X,Y,Z} \langle (\bar{J} \oslash \Phi)(X, Y), \Phi(W, Z) \rangle + \langle (\bar{J} \oslash \Phi)(W, Z), \Phi(X, Y) \rangle. \quad (124)$$

Furthermore, introducing the linear representation $\rho: \mathcal{V} \rightarrow \mathfrak{so}(\mathcal{H})$, we have

$$\begin{aligned} \langle (\bar{J} \oslash \Phi)(X, Y), \Phi(W, Z) \rangle + \langle (\bar{J} \oslash \Phi)(W, Z), \Phi(X, Y) \rangle &= \langle \rho(\bar{J} \oslash \Phi(X, Y)) W, Z \rangle - \langle [J, \rho(\Phi(X, Y))] W, Z \rangle \\ &= \langle \bar{A}(X, Y) W, Z \rangle = -\langle \bar{A}(X, Y) Z, W \rangle \end{aligned} \quad (125)$$

Finally, according to (123), we see that the 4-form $d(\bar{J} \oslash \bar{T})$ is of type $(4, 0) + (0, 4)$, and thus

$$\mathfrak{S}_{X,Y,Z} \langle \bar{A}(X, Y) Z, W \rangle = \mathfrak{S}_{X,Y,Z} \langle \bar{A}^{(-)}(X, Y) Z, W \rangle.$$

This completes the proof. \square

Notation Given $B \in \mathcal{C}(\Lambda^2 T^* N \otimes \mathcal{H})$, we denote simply $\text{Im } B = \{B(X, Y) \in \mathcal{H}, X, Y \in TN\} \subset \mathcal{H}$. In particular, we have $\text{Im } N_{\bar{J}} = N_{\bar{J}}(\mathcal{H}, \mathcal{H}) \subset \mathcal{H}$ and $\text{Im } R_{\mathcal{V}} = R_{\mathcal{V}}(\mathcal{V}, \mathcal{V}) \subset \mathcal{H}$. Moreover, we will also use the notations $\ker C = \{X \in \mathcal{H} \mid C(X, \cdot) = 0\}$ and $\text{Supp}(C) = (\ker C)^\perp$, for any $C \in \mathcal{C}(\Lambda^2 \mathcal{H}^* \otimes TN)$.

Proposition 6.38 *Let us suppose that $R_{\mathcal{V}} = 0$. Then the following identities hold:*

$$\begin{array}{ll} \text{(i)} & dH|_{\mathcal{H}^2 \times \mathcal{V}^2} = 0 \\ \text{(ii)} & dH|_{\mathcal{H}^2 \times \mathcal{V}^2}^* = 0 \end{array} \quad \begin{array}{ll} \text{(iii)} & d(F \cdot N_F)|_{\mathcal{H}^2 \times \mathcal{V}^2} = 0 \\ \text{(iv)} & d(F \oslash (\text{Skew}(\Phi) + \text{Skew}(R_{\mathcal{V}})))|_{\mathcal{H}^2 \times \mathcal{V}^2} = 0. \end{array}$$

Proof. It results from lemma 6.7 applied to the horizontal 3-forms $F \cdot N_F$ and $\bar{J} \oslash \bar{T}$, and then from lemma 6.8.

Proposition 6.39 *Let us suppose that $R_{\mathcal{V}} = 0$ and that the horizontal curvature Φ is pure. Then the following statements are equivalent*

$$\begin{array}{ll} \text{(i)} & dH|_{\mathcal{H}^3 \times \mathcal{V}} = 0 \\ \text{(ii)} & dH|_{\mathcal{H}^3 \times \mathcal{V}}^* = 0 \end{array} \quad \begin{array}{ll} \text{(iii)} & d(F \cdot N_F)|_{\mathcal{H}^3 \times \mathcal{V}} = 0 \\ \text{(iv)} & \mathfrak{S}_{X,Y,Z} \bar{J} N_{\bar{J}}(\rho(V) X, Y, Z) = 0. \end{array}$$

Moreover these later are also equivalent to the following equivalent statements

$$\text{(iv)} \quad N_{\bar{J}}(X, Y, \rho(V) Z) = 0.$$

$$\text{(v)} \quad N_{\bar{J}}(\mathcal{H}, \mathcal{H}) \perp \rho(\mathcal{V})(\mathcal{H}), \text{ or equivalently } \ker N_{\bar{J}} \perp \ker \Phi, \text{ i.e. } \text{Supp}(N_{\bar{J}}) \perp \text{Supp}(\Phi).$$

We will say that the 2-forms $N_{\bar{J}}$ and Φ have orthogonal supports, when (v) is satisfied.

Proof. According to lemma 6.7, we have

$$\begin{aligned} d(\bar{J} \oslash \bar{T})(V_0, H_1, H_2, H_3) &= \mathfrak{S}_{i,j,k} (\bar{J} \oslash \bar{T})(\rho(V_0) H_i, H_j, H_k) \\ &= \mathfrak{S}_{i,j,k} \langle \bar{J} (-3\bar{T}^{--} + \bar{T}^{++} - \bar{T}^{1,1}) (H_i, H_j), \rho(V_0) H_k \rangle \\ &= \mathfrak{S}_{i,j,k} \langle \bar{J} (-3\bar{T}^{--} + \bar{T}^{++} - \bar{T}^{1,1}) (H_i, H_j), \rho^{(-)}(V_0) H_k \rangle \end{aligned} \quad (126)$$

In the same way, we have

$$d(F \cdot N_F)(V_0, H_1, H_2, H_3) = \mathfrak{S}_{i,j,k} \langle (4\bar{J}\bar{T}^{--})(H_i, H_j), \rho^{(-)}(V_0)H_k \rangle.$$

Moreover, for any ∇ -parallel 3-form $\alpha \in \mathcal{C}((\Lambda^2\mathcal{H}^*) \wedge \mathcal{V}^*)$, we have

$$\begin{aligned} d\alpha((V_0, H_1, H_2, H_3)) &= \mathfrak{S}_{i,j,k} \alpha(T(V_0, H_i), H_j, H_k) + \alpha(T(H_i, H_j), V_0, H_k) \\ &= \mathfrak{S}_{i,j,k} \alpha(T^\vee(V_0, H_i), H_j, H_k) + \alpha(\bar{T}(H_i, H_j), V_0, H_k) \\ &= \mathfrak{S}_{i,j,k} \alpha(\bar{T}(H_i, H_j), V_0, H_k) \end{aligned}$$

since $R^\vee = 0$ and hence $T^\vee(V_0, H_i) = 0$. Therefore let us apply this:

$$\begin{aligned} d(F \odot \bar{T})(V_0, H_1, H_2, H_3) &= d(F \odot \text{Skew}(\Phi))(V_0, H_1, H_2, H_3) = \mathfrak{S}_{i,j,k} (F \odot \text{Skew}(\Phi))(H_k, \bar{T}(H_i, H_j), V_0) \\ &= \mathfrak{S}_{i,j,k} (\bar{J} \odot \Phi)(H_k, \bar{T}(H_i, H_j), V_0) = \mathfrak{S}_{i,j,k} \langle [\rho(V_0), \bar{J}] H_k, \bar{T}(H_i, H_j) \rangle \\ &= 2 \mathfrak{S}_{i,j,k} \langle \rho^{(-)}(V_0)H_k, \bar{J}\bar{T}(H_i, H_j) \rangle \quad (127) \end{aligned}$$

Then applying lemma 6.8 to (126) and (127), we obtain

$$\mathfrak{S}_{i,j,k} \langle \bar{J}(-\bar{T}^{--} + \bar{T}^{1,1} + 3\bar{T}^{++})(H_i, H_j), \rho^{(-)}(V_0)H_k \rangle = 0$$

and then taking respectively the $(3, 0) + (0, 3)$ -type and the $(2, 1) + (1, 2)$ -type part w.r.t. to the horizontal variables $(H_1, H_2, H_3) \in \mathcal{H}^3$, of this equation we obtain

$$\langle \bar{J}\bar{T}^{++}(H_i, H_j), \rho^{(-)}(V_0)H_k \rangle = 0 \quad (128)$$

$$\langle \bar{J}(-\bar{T}^{--} + \bar{T}^{1,1})(H_i, H_j), \rho^{(-)}(V_0)H_k \rangle = 0 \quad (129)$$

This proves the equivalence between the four first statements.

The equivalence (v) \Leftrightarrow (vi) is obvious. It remains to prove the equivalence (iv) \Leftrightarrow (v). But this results immediately from the fact that $N_{\bar{J}} = 4\bar{T}^{--}$ is a 3-form of type $(3, 0) + (0, 3)$ and $\rho = \rho^{(-)}$. This completes the proof. \square

Finally,

Lemma 6.9 *The components in $\Lambda^4\mathcal{V}^*$ and $(\Lambda^3\mathcal{V}^*) \wedge \mathcal{H}^*$ of the following exterior derivatives: dH , dH^* , $d(F \cdot N_F)$ and $d(F \odot (\text{Skew}(\Phi) + \text{Skew}(R_\gamma)))$, vanishes.*

Proof. Apply lemma 6.7 to the horizontal 3-forms $\bar{J} \odot \bar{T}$ and $F \cdot N_F$. Then use lemma 6.8. This completes the proof. \square

Let us summarize:

Theorem 6.9 *Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . Let us suppose that one of its characteristic connections, ∇ , has a parallel torsion $\nabla T = 0$. Let us suppose that $R_\gamma = 0$ and that the horizontal curvature Φ is pure.*

- *Then (N, F, h) has a closed stringy structure if and only if the horizontal 3-form $F \cdot N_F$ is closed.*
- *Moreover, this happens if and only if the horizontal complex structure \bar{J} is a cyclic permutation of the horizontal curvature, and the 2-forms $N_{\bar{J}}$ and Φ have orthogonal supports.*

The particular case of Horizontally Hermitian f -manifolds. In horizontally Hermitian f -manifolds, we have $F \cdot N_F = 0$ and therefore $F \cdot N_F = 0$ is a closed 3-form. Therefore we can apply theorem 6.9.

Corollary 6.12 *Let $\pi: (N, F, h) \rightarrow (M, g)$ be a Riemannian submersion from a metric f -manifold into a Riemannian manifold. Let us suppose that (N, F, h) is reductive and horizontally Hermitian, so that it is in particular horizontally of type \mathcal{G}_1 . Let us suppose also that one of its characteristic connections, ∇ , has a parallel torsion $\nabla T = 0$ and that the horizontal curvature Φ is pure. Then (N, F, h) has a closed stringy structure.*

6.4.3 The sigma model with a Wess-Zumino term in reductive metric f -manifold of global type \mathcal{G}_1 .

Now, we can conclude with the following variational interpretation of the stringy harmonicity.

Theorem 6.10 *Let (N, F, h) be a reductive metric f -manifold of global type \mathcal{G}_1 . Let us suppose that (N, F, h) is stringy closed. Let ∇ be one characteristic connection.*

• *Then the equation for stringy harmonic maps (w.r.t. ∇) $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H = -d\Omega_F + F \cdot N_F - \frac{1}{2}F \odot (\text{Skew}(\Phi)).$$

• *Moreover the equation for \star -stringy harmonic maps $f: L \rightarrow N$ is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H^* = -d\Omega_F + \frac{1}{2}F \cdot N_F - \frac{1}{2}F \odot (\text{Skew}(\Phi)).$$

6.4.4 The example of a naturally reductive homogeneous space

By definition of (\star) -stringy harmonicity and the expression of the torsion of ∇^0 in term of the Lie bracket, we have the following.

Proposition 6.40 *Let $N = G/K$ be a Riemannian homogeneous manifold endowed with a G -invariant complex structure F . Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{n}$ be a reductive decomposition of \mathfrak{g} , and $\mathfrak{n} = \mathfrak{m} \oplus \mathfrak{p}$ the $\text{Ad}K$ -invariant decomposition defined by F . Let $f: L \rightarrow N$ be a smooth map, $U: L \rightarrow G$ be a (local) lift of f and $\alpha = U^{-1}.dU$ the corresponding Maurer-Cartan form. Then in term of α , the equation of stringy harmonicity (w.r.t. ∇^0) is written*

$$\begin{cases} d * \alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{p}}] + \frac{1}{2}[\bar{J}_0 \alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{p}} = 0 \\ d * \alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{m}}] - \frac{1}{2}\bar{J}_0 [\bar{J}_0 \alpha_{\mathfrak{m}} \wedge \bar{J}_0 \alpha_{\mathfrak{m}}]_{\mathfrak{m}} + \frac{1}{2}([\alpha_{\mathfrak{p}} \wedge \bar{J}_0 \alpha_{\mathfrak{m}}] - \bar{J}_0[\alpha_{\mathfrak{p}} \wedge \alpha_{\mathfrak{m}}]) = 0 \end{cases}$$

whereas the equation of \star -stringy harmonicity is written:

$$\begin{cases} d * \alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{p}}] + \frac{1}{2}[\bar{J}_0 \alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{p}} = 0 \\ d * \alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{m}}] + \frac{1}{2}[\bar{J}_0 \alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} + \frac{1}{4}J_0([\bar{J}_0 \alpha_{\mathfrak{m}} \wedge \bar{J}_0 \alpha_{\mathfrak{m}}]_{\mathfrak{m}} + [\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}}) \\ \quad + \frac{1}{2}([\alpha_{\mathfrak{p}} \wedge \bar{J}_0 \alpha_{\mathfrak{m}}] - \bar{J}_0[\alpha_{\mathfrak{p}} \wedge \alpha_{\mathfrak{m}}]) = 0. \end{cases}$$

where \bar{J}_0 is the complex structure on \mathfrak{m} corresponding to F .

6.4.5 Geometric interpretation of the maximal determined even case.

In this subsection, we suppose that $N = G/K$ is a (locally) $(2k + 1)$ -symmetric space, and we use the notations and the conventions of 2.1.

Theorem 6.11 *Let us suppose that $N = G/K$ is a (locally) $2k$ -symmetric space endowed with its canonical f -structure⁷³ F and its canonical connection ∇^0 . Then the associated maximal determined system, $\text{Syst}(2k - 1, \tau)$ is the equation of \star -stringy harmonicity for the geometric map $f: L \rightarrow N: (\nabla^0)^* df + (F \star T)(f) = 0$.⁷⁴*

Moreover, if we consider now that $N = G/K$ is endowed with the f -structure $F^ := \bigoplus_{j=1}^{k-1} (-1)^j F_{[\mathfrak{m}_j]} \oplus 0_{[\mathfrak{g}_k]}$, then this system is the equation of stringy harmonicity for the geometric map $f: L \rightarrow N: (\nabla^0)^* df + (F^* \cdot T)(f) = 0$.*

Proof. The first point follows from theorem 2.8, proposition 6.40 and the fact that the component $\bar{T} = T_{|\mathcal{H}^2}^{\mathcal{H}}$ of the torsion T of ∇^0 satisfies the same kind of identities as in proposition 5.21. The second point follows from proposition 5.16 and remark 5.13 applied to the complex vector bundle (\mathcal{H}, \bar{J}) , the 2-form $\bar{T} \in \mathcal{T}(\mathcal{H})$ and the \bar{J} -invariant decomposition $TN = E^+ \oplus E^-$, defined by $E^+ = \bigoplus_{\substack{j=1 \\ j \text{ even}}}^{k-1} [\mathfrak{m}_j]$, and $E^- = \bigoplus_{\substack{j=1 \\ j \text{ odd}}}^{k-1} [\mathfrak{m}_j]$. This completes the proof. \square

Theorem 6.12 *Let us suppose that $N = G/K$ is a (locally) $2k$ -symmetric space endowed with its canonical f -structure F and its canonical connection ∇^0 . Let us suppose that $N = G/K$ is naturally reductive and we choose a naturally reductive G -invariant metric h for which $\tau_{|\mathfrak{n}}$ is orthogonal⁷⁵ and thus which is compatible with F . Then (N, F, h) is a reductive metric f -manifold of global type \mathcal{G}_1 . Moreover, (N, F, h) has a closed stringy structure.*

Therefore, the associated maximal determined system, $\text{Syst}(2k - 1, \tau)$, is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form

$$H^* = -d\Omega_F + \frac{1}{2}F \cdot N_F - \frac{1}{2}F \odot (\text{Skew}(\Phi)).$$

Moreover, if we consider now that $N = G/K$ is endowed with the f -structure F^ , then the previous system is exactly the Euler-Lagrange equation for the sigma model in N with a Wess-Zumino term defined by the closed 3-form*

$$H = -d\Omega_{F^*} + F^* \cdot N_{F^*} - \frac{1}{2}F \odot (\text{Skew}(\Phi)).$$

The theorem will follow from the two following lemmas.

Lemma 6.10 *In the metric f -manifold (N, F, h) of theorem 6.12, \bar{J} is a cyclic derivation of the horizontal curvature.*

Proof. According to subsection 6.3.4, we have

$$\begin{aligned} \bar{A}(\widetilde{X, Y})Z &= [[\bar{J}_0 X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}}, Z_{\mathfrak{m}}] + [[X_{\mathfrak{m}}, \bar{J}_0 Y_{\mathfrak{m}}]_{\mathfrak{p}}, Z_{\mathfrak{m}}] - \bar{J}_0 [[X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}}, Z_{\mathfrak{m}}] + [[X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}}, \bar{J}_0 Z_{\mathfrak{m}}] \\ &= 2([[\bar{J}_0 X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}}, Z_{\mathfrak{m}}] + [[X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}}, \bar{J}_0 Z_{\mathfrak{m}}]) \end{aligned} \quad (130)$$

⁷³defined by (37)

⁷⁴Where we have removed the index "g" which precises that the previous terms are computed with respect to some Hermitian metric g on L .

⁷⁵See the Appendix, lemma 8.3.

In this equation, $X, Y, Z \in \mathcal{C}(\mathcal{H})$ with lifts $X_m, Y_m, Z_m \in \mathcal{C}^\infty(G, \mathfrak{m})$, and $\widetilde{\bar{A}(X, Y)Z} \in \mathcal{C}^\infty(G, \mathfrak{m})$ denotes the lift of $\bar{A}(X, Y)Z \in \mathcal{C}(\mathcal{H})$. An other possibility is to consider that $X, Y, Z \in \mathcal{H}$ are horizontal vectors at some point $y \in N$ and that we have chosen $g \in G$ such that $g.G_0 = y$ and that we have set $X = \text{Ad}g(X_m)$, and idem for Y and Z . Then the equation (130), when written in the form $\widetilde{\bar{A}} = B_m$, means in fact that we have $A = \text{Ad}g(B_m)$.

In the following, we will use this kind of notation without recalling these precisions.

Moreover, since we have $[\bar{J}_0 X_m, Y_m]_{\mathfrak{p}} = [X_m, \bar{J}_0 Y_m]_{\mathfrak{p}}$ and $(\text{ad}_m \mathfrak{g}_k) \bar{J}_0 = -\bar{J}_0 (\text{ad}_m \mathfrak{g}_k)$, then we obtain

$$\begin{aligned} \mathfrak{S}_{X, Y, Z} \widetilde{\bar{A}(X, Y)Z} &= 2i \mathfrak{S}_{X, Y, Z} \left(\sum_{1 \leq |j|, |l| \leq k-1} s(-j) [[X_j, Y_{-j+k}], Z_l] + \sum_{1 \leq |j|, |l| \leq k-1} s(-l) [[X_j, Y_{-j+k}], Z_l] \right) \\ &= 2i \mathfrak{S}_{X, Y, Z} \sum_{1 \leq |l| \leq k-1} s(-l) \sum_{\substack{1 \leq |j| \leq k-1 \\ j_l > 0}} [[X_j, Y_{-j+k}], Z_l] \end{aligned}$$

where $s(n)$ is the sign of $n \in \mathbb{Z}$. $\alpha := \mathfrak{S}_{X, Y, Z} \widetilde{\bar{A}(X, Y)Z}$ defines a element of $\Lambda^3 \mathfrak{g}^* \otimes \mathfrak{g}$, which in fact corresponds to a 4-form of type $(4, 0) + (0, 4)$ according to the proof of proposition 6.37 (see in particular, equations (124) and (125)). Let us consider the component in $(\mathfrak{g}_j^* \wedge \mathfrak{g}_{-j+k}^* \wedge \mathfrak{g}_l^*) \otimes \mathfrak{g}$ of this the element of this element of $\Lambda^3 \mathfrak{g}^* \otimes \mathfrak{g}$. We remark first that this component is in fact in $(\mathfrak{g}_j^* \wedge \mathfrak{g}_{-j+k}^* \wedge \mathfrak{g}_l^*) \otimes \mathfrak{g}_{k+l}$. Let us set

$$\begin{aligned} I^+ &= \{ \{j, -j+k, l\}, 1 \leq j, l \leq k-1 \} \\ I^- &= \{ \{j, -j+k, l\}, -(k-1) \leq j, l \leq -1 \} = -I^+ \\ I &= I^+ \sqcup I^-. \end{aligned}$$

Then, denoting by $\overline{k+l}$ the representant modulo $2k$ of $k+l$ in $\{-(k-1), \dots, -1\} \cup \{1, \dots, k-1\}$, we have $\{j, \overline{k+l}, l\} \notin I$ because $\overline{k+l}$ has a opposite sign to j and l . Therefore, the component in $(\mathfrak{g}_j^* \wedge \mathfrak{g}_{\overline{k+l}}^* \wedge \mathfrak{g}_l^*) \otimes \mathfrak{g}_{-j+k}$ of our element α vanishes, which since α is a 4-form implies that $\alpha = 0$. This completes the proof. \square

Lemma 6.11 *In the metric f -manifold (N, F, h) of theorem 6.12, the 2-forms $N_{\bar{j}}$ and Φ have orthogonal supports.*

Proof. Denoting by T the torsion of the canonical connection ∇^0 , we have

$$T^{--}(\widetilde{X}, Y) = - \sum_{\substack{k+1 \leq i+j \leq 2(k-1) \\ 1 \leq i \leq j \leq k-1}} \left(1 - \frac{\delta_{ij}}{2} \right) ([X_{m_i}, Y_{m_j}]_{m_{i+j}} + [X_{m_j}, Y_{m_i}]_{m_{i+j}}).$$

Therefore

$$\langle T^{--}(X, Y), \rho(V)Z \rangle = - \sum_{\substack{k+1 \leq i+j \leq 2(k-1) \\ 1 \leq i \leq j \leq k-1}} \left(1 - \frac{\delta_{ij}}{2} \right) \langle [X_{m_i}, Y_{m_j}]_{m_{i+j}}, [Z_{m_{i+j-k}}, V_k] \rangle$$

but we have

$$\begin{aligned} \langle [X_{m_i}, Y_{m_j}]_{m_{i+j}}, [Z_{m_{i+j-k}}, V_k] \rangle &= \langle [X_{m_i}, Y_{m_j}]_{i+j}, [Z_{i+j-k}, V_k] \rangle + \langle [X_{m_i}, Y_{m_j}]_{-(i+j)}, [Z_{-(i+j)-k}, V_k] \rangle \\ &= \langle [[X_{m_i}, Y_{m_j}]_{i+j}, Z_{i+j-k}], V_k \rangle + \langle [[X_{m_i}, Y_{m_j}]_{-(i+j)}, Z_{-(i+j)-k}], V_k \rangle \end{aligned}$$

and these two scalar products vanishes, because $2(i+j) - k \neq k \pmod{2k}$ and $-2(i+j) + k \neq k \pmod{2k}$. Indeed $2(i+j) \neq 0 \pmod{2k}$, since $k+1 \leq i+j \leq 2(k-1)$. This completes the proof. \square

Proof of theorem 6.12 It follows from proposition 6.29, lemmas 6.10 and 6.11, theorems 6.9 and 6.10. This completes the proof. \square

Remark 6.21 Let us remark that a 4-symmetric space, endowed with its canonical f -structure F and a naturally reductive metric is horizontally Kähler.

6.4.6 Twistorial Geometric interpretation of the maximal determined even case.

We come back to the situation of 6.3.7. We use also the notation of definition 3.5. Then according to proposition 6.32⁷⁶ and theorem 6.11, we obtain

Theorem 6.13 *Let (L, j) be a Riemann surface, $f: L \rightarrow N = G/G_0$ be a map and $J = f^*\mathcal{J}_{J_0}$ the corresponding map into $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$. Then f is a geometric solution of the even maximal determined system $(\text{Syst}(2k-1, \tau))$ if and only if $J: L \rightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$ is \star -stringy harmonic w.r.t. the canonical paracharacteristic connection in the twistor space $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$.*

Moreover, let us consider now that $N = G/G_0$ is endowed with the f -structure $F^ := \bigoplus_{j=1}^{k-1} (-1)^j F_{[\mathfrak{m}_j]} \oplus 0_{[\mathfrak{g}_k]}$, and that $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$ is endowed with the f -structure $\mathcal{F}^* := \bigoplus_{j=1}^{k-1} (-1)^j \underline{\mathcal{J}}_{\mathfrak{m}_j(\mathcal{J})} \oplus 0_{\mathfrak{m}_k(\mathcal{J})}$, where $\mathfrak{m}_j(\mathcal{J})$ is defined as in definition 3.5, and $\mathfrak{m}_k(\mathcal{J}) = \ker(\mathcal{J} + \text{Id})$.*

Then f is a geometric solution of this system if and only if $J: L \rightarrow \mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$ is stringy harmonic w.r.t. the canonical paracharacteristic connection in the twistor space $\mathcal{Z}_{2k,2}^{\alpha_0}(M, J_2)$.

⁷⁶Or we can use proposition 6.34 as well

7 Generalized harmonic maps into reductive homogeneous spaces

This section has 3 objectives.

1. We want to precise clearly and concretely the geometric meaning of each component of the zero curvature equation on a 1-form α_λ taking values in a finite dimensional vector subspace of a loop Lie algebra. Indeed, we would like to interpret each component of this equation with a geometric property, as we have already done that, for example for the even minimal determined system, in section 4.3.1. But here, we would like to keep our study in the Lie algebra setting and in the corresponding homogeneous space for the geometric interpretation. Moreover, we would like to see how one can derive naively some of the results obtained in sections 4, 5, and 6. In other words, we would like to continue and complete, in the light of the studies done in sections 4, 5, and 6, the preliminary and naive study begun in section 2, keeping the same naive spirit. Then, over and above recovering by a different way, precisising, or completing results of the previous sections, as well as outlining results already implicitly contained in the previous more general studies but which can be precisised in more particular contexts, we also prove some new results. For example, we prove that the strongly harmonicity in a reductive homogeneous space has a formulation in terms of a zero curvature equation (that we call elliptic integrable system associated to a reductive homogeneous space). Moreover, we complete the study of the odd minimal determined system begun in section 2.3.2.
2. Moreover, we want to precise the geometric interpretation of the intermediate determined systems ($m_{k'} < m < k' - 1$). We already know that these are the equations of stringy harmonic maps which satisfies some additional holomorphicity conditions. We would like to make this more precise.
3. Finally, we conclude with some remarks about the twistorial interpretation.

7.1 Affine harmonic maps into reductive homogeneous spaces.

Let $N = G/K$ be a reductive homogeneous space and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$ a reductive decomposition of the Lie algebra \mathfrak{g} . We use the notations of section 1 (applied to $N = G/K$ instead of $M = G/H$).

Theorem 7.1 *Let (L, j) be a Riemann surface and $f: (L, j) \rightarrow N$ be a smooth map, let $F: L \rightarrow G$ be a (local) lift of f and $\alpha = F^{-1}.dF$. Then the following statements are equivalent:*

- (i) f is ∇^t -harmonic for one $t \in [0, 1]$.
- (ii) f is ∇^t -harmonic for every $t \in [0, 1]$.
- (iii) $d * \alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{m}}] = 0$.
- (iv) $\text{Im} (\bar{\partial} \alpha'_{\mathfrak{m}} + [\alpha''_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{m}}] + t[\alpha''_{\mathfrak{m}} \wedge \alpha'_{\mathfrak{m}}]_{\mathfrak{m}}) = 0, \forall t \in [0, 1]$.

In fact, the tension field $\tau^t(f)$ of f with respect to ∇^t is independent of $t \in [0, 1]$.

Theorem 7.2 *In the same situation as above, the following statements are equivalent:*

- (i) f is strongly ∇^t -harmonic for one $t \in [0, 1] \setminus \{\frac{1}{2}\}$.
- (ii) f is strongly ∇^t -harmonic for every $t \in [0, 1] \setminus \{\frac{1}{2}\}$.
- (iii) $\bar{\partial} \alpha'_{\mathfrak{m}} + [\alpha''_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{m}}] + t[\alpha''_{\mathfrak{m}} \wedge \alpha'_{\mathfrak{m}}]_{\mathfrak{m}} = 0, \forall t \in [0, 1] \setminus \{\frac{1}{2}\}$.

(iv) f is ∇^t -harmonic for one $t \in [0, 1]$ and $[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} = 0$.

(iv) $d\alpha_{\lambda} + \frac{1}{2}[\alpha_{\lambda} \wedge \alpha_{\lambda}] = 0, \forall \lambda \in S^1$, with $\alpha_{\lambda} = \lambda^{-1}\alpha'_{\mathfrak{m}} + \alpha_{\mathfrak{k}} + \lambda\alpha''_{\mathfrak{m}}$.

Furthermore f is strongly $\nabla^{\frac{1}{2}}$ -harmonic if and only if it is $\nabla^{\frac{1}{2}}$ -harmonic: indeed $\nabla^{\frac{1}{2}}$ is torsion free.

Proof of theorem 7.1 The tension field $\tau^t(f)$ of f with respect to ∇^t is given by

$$\begin{aligned} \tau^t(f) &= *d^{\nabla^t} *df = *\text{Ad}F(d*\alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge *\alpha_{\mathfrak{m}}] + t[\alpha_{\mathfrak{m}} \wedge *\alpha_{\mathfrak{m}}]_{\mathfrak{m}}) \\ &= *\text{Ad}F(d*\alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge *\alpha_{\mathfrak{m}}]) \end{aligned}$$

(see section 1.4 (especially equation (9)) and section 1.6). This proves the equivalence between (i), (ii) and (iii). Then we conclude by remarking that $2\text{Im}(\bar{\partial}\alpha'_{\mathfrak{m}} + [\alpha''_{\mathfrak{k}} \wedge \alpha''_{\mathfrak{m}}]) = d*\alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge *\alpha_{\mathfrak{m}}]$ and that $[\alpha'_{\mathfrak{m}} \wedge \alpha''_{\mathfrak{m}}]_{\mathfrak{m}} = \frac{1}{2}[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}}$ is real. This completes the proof. \square

Proof of theorem 7.2 We have for all $t \in [0, 1]$

$$\bar{\partial}^{\nabla^t} \partial f = \text{Ad}F(\bar{\partial}\alpha'_{\mathfrak{m}} + [\alpha''_{\mathfrak{k}} \wedge \alpha''_{\mathfrak{m}}] + t[\alpha''_{\mathfrak{m}} \wedge \alpha'_{\mathfrak{m}}]_{\mathfrak{m}}) \quad (131)$$

so that the ∇^t -strongly harmonicity of f is written:

$$\bar{\partial}\alpha'_{\mathfrak{m}} + [\alpha''_{\mathfrak{k}} \wedge \alpha''_{\mathfrak{m}}] + t[\alpha''_{\mathfrak{m}} \wedge \alpha'_{\mathfrak{m}}]_{\mathfrak{m}} = 0. \quad (S_{\mathfrak{m}}^t)$$

Then the imaginary part of $\bar{\partial}^{\nabla^t} \partial f = 0$ gives us the ∇^t -harmonicity whereas the real part gives us

$$d\alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge \alpha_{\mathfrak{m}}] + t[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} = 0 \quad (\text{Re}(t))$$

which is nothing but the lift of the torsion free equation: $f^*T^t = 0$, where $T^t = T^{\nabla^t}$. Moreover the projection on \mathfrak{m} of the Maurer-Cartan equation (on α) gives us the structure equation

$$d\alpha_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge \alpha_{\mathfrak{m}}] + \frac{1}{2}[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} = 0 \quad [\text{MC}]_{\mathfrak{m}}$$

which is nothing but $(\text{Re}(\frac{1}{2}))$ (so that we recover that $T^{\frac{1}{2}} = 0$) but (since it can be written $((\text{Re}(0)) + \frac{1}{2}[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} = 0)$ it is also the lift of (the f -pullback of) the equation expressing the canonical torsion T^0 in term of the Lie bracket (see theorem 1.4 or equation (14)) :

$$T^0 + [,]_{[\mathfrak{m}]} = 0. \quad (132)$$

which combining with the fact that the left hand side of $(\text{Re}(t))$ is the lift of f^*T^t , gives us back $T^t = (2t - 1)[,]_{[\mathfrak{m}]}$ (see (14)).

Hence according to (131) and $[\text{MC}]_{\mathfrak{m}}$ the strongly harmonicity for one $t \neq \frac{1}{2}$ is equivalent to the harmonicity (imaginary part) and $[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} = 0$ (real part $(\text{Re}(t))$ combining with $[\text{MC}]_{\mathfrak{m}}$). We can also simply say that f is strongly harmonic if and only if f is harmonic and torsion free i.e. $[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} = 0$ according to (14). This proves the equivalence between (i), (ii), and (iii). Now, let us decompose the curvature of α_{λ} , with respect to powers of λ :

$$\begin{aligned} d\alpha_{\lambda} + \frac{1}{2}[\alpha_{\lambda} \wedge \alpha_{\lambda}] &= \lambda^{-1}(d\alpha'_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{m}}]) \\ &+ (d\alpha_{\mathfrak{k}} + \frac{1}{2}[\alpha_{\mathfrak{k}} \wedge \alpha_{\mathfrak{k}}] + \frac{1}{2}[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{k}}) + \frac{1}{2}[\alpha_{\mathfrak{m}} \wedge \alpha_{\mathfrak{m}}]_{\mathfrak{m}} \\ &+ \lambda(d\alpha''_{\mathfrak{m}} + [\alpha_{\mathfrak{k}} \wedge \alpha''_{\mathfrak{m}}]) \end{aligned}$$

hence using the fact that α_λ is real (i.e. \mathfrak{g} -valued)

$$d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0 \Leftrightarrow \begin{cases} d\alpha'_m + [\alpha_\mathfrak{k} \wedge \alpha'_m] = 0 & (S_m^0) \\ [\text{MC}]_\mathfrak{k} & \\ [\alpha_m \wedge \alpha_m]_m = 0 & \end{cases} \Leftrightarrow \begin{cases} (S_m^0) \\ [\text{MC}] \end{cases},$$

In the last equivalence, we use the fact that $(S_m^0) + \overline{(S_m^0)}$ is the equation $d\alpha_m + [\alpha_\mathfrak{k} \wedge \alpha_m] = 0$ which combined with $[\text{MC}]_m$ (above) gives us $[\alpha_m \wedge \alpha_m]_m = 0$. Thus the zero curvature equation on α_λ is equivalent to the strongly ∇^0 -harmonicity, i.e. the strongly ∇^t -harmonicity for all $t \in [0, 1] \setminus \{\frac{1}{2}\}$. Finally the last assertion is obvious. This completes the proof. \square

We are led naturally to the following definitions.

Definition 7.1 *We will say that $f: L \rightarrow G/K$ is torsion free if $f^*T^t = 0$ for $t \in [0, 1] \setminus \{\frac{1}{2}\}$ (this equation does not depend on t).*

Definition 7.2 *In the situation described by theorem 7.2-(iv), we will say that the \mathfrak{g} -valued 1-form on L , α , is solution of the the first elliptic system associated to the reductive homogeneous space G/K , and that the corresponding geometric map f is a geometric solution of this system.*

Remark 7.1 The independence of the ∇^t -tension field w.r.t. to t (and therefore the equivalence between the statements (i) and (ii) in theorem 7.1), are in fact a consequence of the geodesic equivalence of the connections ∇^t (see propositions 5.7 and 5.8). Moreover, the equivalence of the ∇^t -strongly harmonicities, for $t \neq \frac{1}{2}$, in theorem 7.2-(i, ii), is a particular case of remark 5.6.

Remark 7.2 Our so-called "strongly harmonic" maps in homogeneous space already appeared in the literature. It seems that they appeared for the first time in [11]. Moreover, in [15], a concrete example of these maps is given: Gauss maps of CMC surfaces in H^3 .

However, it seems to us very strange that nobody has remarked before that the torsion free condition $[\alpha_m \wedge \alpha_m]_m = 0$ is already contained in the zero curvature equation $d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0$, $\forall \lambda \in S^1$. Indeed everybody in the literature (in particular [11]) adds the torsion free condition to the zero curvature equation to characterise the torsion free harmonic maps - i.e. strongly harmonic maps -, whereas the strongly harmonicity is exactly and simply equivalent to the zero curvature equation.

Affine harmonic maps into symmetric spaces

Now, if we suppose in particular that N is (locally) symmetric, i.e. $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{k}$, then all the connections ∇^t , $0 \leq t \leq 1$, coincide. Moreover, if N is also Riemannian then these are equal to the Levi-Civita connection. Therefore we obtain:

Corollary 7.1 *The first elliptic integrable system associated to a (locally) symmetric space $N = G/K$ is the equation for ∇^0 -harmonic maps $f: L \rightarrow N$. If N is Riemannian this means that it is the equation for harmonic maps $f: L \rightarrow N$ (with respect to Levi-Civita in N).*

7.2 Affine/holomorphically harmonic maps into 3-symmetric spaces

Let us suppose now that $N = G/G_0$ is a (locally) 3-symmetric space. We use the notations of section 2. N is endowed with its canonical almost complex structure \underline{J} defined by (26). We continue here the study begun in 2.3.2 concerning the lowest order determined odd system.

Theorem 7.3 *Let (L, j) be a Riemann surface and $f: L \rightarrow N$ a smooth map. Let $F: L \rightarrow G$ be a (local) lift of f and $\alpha = F^{-1}.dF$. Then the following statement are equivalent*

- (i) $\bar{\partial}\alpha'_{-1} + [\alpha''_0 \wedge \alpha'_{-1}] + [\alpha''_1 \wedge \alpha'_1] = 0$ (S_1)
- (ii) $\bar{\partial}\alpha'_1 + [\alpha''_0 \wedge \alpha'_1] = 0$ (S_2)
- (iii) f is holomorphically ∇^1 -harmonic: $[\bar{\partial}^{\nabla^1} \partial f]^{1,0} = 0$.
- (iv) f is anti-holomorphically ∇^0 -harmonic: $[\bar{\partial}^{\nabla^0} \partial f]^{0,1} = 0$.
- (v) f is a geometric solution of the second elliptic integrable system associated to the (locally) 3-symmetric space G/G_0 :

$$d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0, \quad \forall \lambda \in S^1,$$

where $\alpha_\lambda = \lambda^{-2}\alpha'_1 + \lambda^{-1}\alpha'_{-1} + \alpha_0 + \lambda\alpha''_1 + \lambda^2\alpha''_{-1}$.

Proof. The equivalences (i) \Leftrightarrow (ii) \Leftrightarrow (v) have been proved in 2.3.2. To prove (i) \Leftrightarrow (iii): just take the (1, 0)-component in $TN^{\mathbb{C}}$ of (131) for $t = 1$. Idem for (ii) \Leftrightarrow (iv). This completes the proof. \square

Remark 7.3 In fact, the equivalences (iv) \Leftrightarrow (v) and (iii) \Leftrightarrow (iv) has also been already derived in 2.3.2, so that this theorem has already been completely proved there.

Now, additionning theorems 7.3, 7.2, 5.1 and proposition 5.1, we obtain

Corollary 7.2 *The following statements are equivalent*

- (i) f is strongly ∇^t -harmonic for one $t \in [0, 1] \setminus \{\frac{1}{2}\}$.
- (ii) f is ∇^t -harmonic for one $t \in [0, 1]$ and torsion free.
- (iii) f is holomorphically ∇^t -harmonic for one $t \in [0, 1]$ and torsion free.
- (iv) f is a geometric solution of the first elliptic system associated to the reductive homogeneous space G/G_0 .
- (v) f is a geometric solution of the second elliptic system associated to the 3-symmetric space G/G_0 , and moreover $[\alpha_1 \wedge \alpha_1] = 0$.
- (vi) f is in the same time a geometric solution of the determined odd elliptic systems (Syst(2, τ)) and (Syst(2, τ^{-1})).

Now, let us apply theorem 5.8 to the equivalence (iv) \Leftrightarrow (v) of theorem 7.3.

Theorem 7.4 *The second elliptic integrable system associated to a naturally reductive 3-symmetric space $N = G/G_0$ is the Euler-Lagrange equation for the sigma model in N with the Wess-Zumino term defined by the closed 3-form $H = -\frac{1}{3}d\Omega_{\underline{J}}$, where \underline{J} is the canonical almost complex structure.*

7.3 (Affine) vertically (holomorphically) harmonic maps

7.3.1 Affine vertically harmonic maps: general properties

Here we generalise the definition of vertical harmonicity for maps from a Riemannian surface into an affine manifold.

Definition 7.3 *Let (N, ∇) be an affine manifold. Let us suppose that we have a splitting $TN = \mathcal{V} \oplus \mathcal{H}$. In other words N is endowed with a Pfaffian system (the vertical subbundle \mathcal{V}) and with a connection on this Pfaffian system. Let $f: (L, b) \rightarrow N$ be a smooth map from a Riemannian manifold (L, b) into N . Then we set*

$$\tau^v(f) = \text{Tr}_b(\nabla^v d^v f) = *d^{\nabla^v} * d^v f,$$

where $\nabla^v d^v f$ is the vertical component of the covariant derivative of df with respect to the connection on $T^*L \otimes f^*TN$ induced by the Levi-Civita connection of L and the linear connection ∇ . We will say that f is affine vertically harmonic with respect to ∇ or ∇ -vertically harmonic if $\tau^v(f) = 0$.

Theorem 7.5 *Let (L, j) be a Riemann surface and $f: (L, j) \rightarrow (N, \nabla)$ a smooth map. Then we have*

$$2\bar{\partial}^{\nabla^v} \partial^v f = d^{\nabla^v} d^v f + id^{\nabla^v} * d^v f,$$

moreover $d^{\nabla^v} d^v f = f^*T^v$, where T^v is the vertical torsion (see 4.1.3) and $d^{\nabla^v} * d^v f = \tau^v(f)\text{vol}_b$ for any hermitian metric b in L . Therefore the following statements are equivalent:

- (i) $(\nabla'')^v \partial^v f = 0$.
- (ii) $\bar{\partial}^{\nabla^v} \partial^v f = 0$.
- (iii) $\nabla_{\frac{\partial}{\partial \bar{z}}}^v \left(\frac{\partial^v f}{\partial z} \right) = 0$, for any holomorphic local coordinate $z = x + iy$ (i.e. (x, y) are conformal coordinates for any hermitian metric in L).
- (iv) f is ∇^v -vertically harmonic with respect to any hermitian metric in L and vertically torsion free: $f^*T^v = 0$ (i.e. $T^v(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}) = 0$ for any x, y conformal coordinates).

We will say in this case that f is strongly ∇ -vertically harmonic.

7.3.2 Affine vertically holomorphically harmonic maps

Here we generalize the notion of holomorphic harmonicity by introducing a new notion of vertical holomorphic harmonicity (in the same way that the vertical harmonicity generalizes the harmonicity).

Definition 7.4 *Let (N, ∇) be an affine manifold with a splitting $TN = \mathcal{V} \oplus \mathcal{H}$ as in definition 7.3. Let us suppose that the subbundle \mathcal{V} admits a complex structure $J^{\mathcal{V}}$. Let us denote by $\mathcal{V}^{\mathbb{C}} = \mathcal{V}^{1,0} \oplus \mathcal{V}^{0,1}$ the splitting induced by the complex structure $J^{\mathcal{V}}$. Then we will say that a map $f: (L, j_L) \rightarrow N$ from a Riemann surface into N , is vertically holomorphically harmonic with respect to ∇ or ∇ -vert. hol. harmonic if*

$$\left[\bar{\partial}^{\nabla^v} \partial^v f \right]^{1,0} = 0.$$

Theorem 7.6 *Let (L, j_L) be a Riemann surface and (N, ∇) be an affine manifold with a splitting $TN = \mathcal{V} \oplus \mathcal{H}$ and a complex structure $J^\mathcal{V}$ on \mathcal{V} as in the previous definition. Then $f: L \rightarrow N$ is ∇ -vert. hol. harmonic if and only if*

$$f^*T^v + J^v d^{\nabla^v} * d^v f = 0.$$

Proof. The same as the one of proposition 5.1. □

7.4 Affine vertically harmonic maps into reductive homogeneous space

Let G be a Lie group, and $K \subset H \subset G$ subgroups of G such that $M = G/H$ and H/K are reductive. We use the notations of 4.3.1 (but we do not suppose a priori that the reductive homogeneous spaces are Riemannian).

Theorem 7.7 *Let (L, j) be a Riemann surface and $f: L \rightarrow N = G/K$ be a smooth map, $F: L \rightarrow G$ a (local) lift of f and $\alpha = F^{-1}.dF$. Then the following statements are equivalent:*

- (i) f is ∇^t -vertically harmonic for one $t \in [0, 1]$.
- (ii) f is ∇^t -vertically harmonic for all $t \in [0, 1]$.
- (iii) $d * \alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{p}}] = 0$.
- (iv) $\text{Im}(\bar{\partial} \alpha'_{\mathfrak{p}} + [\alpha''_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{p}}] + t[\alpha''_{\mathfrak{p}} \wedge \alpha'_{\mathfrak{p}}]_{\mathfrak{p}}) = 0, \forall t \in [0, 1]$.

The vertical tension field $\tau^{t,v}(f)$, with respect of ∇^t , is independent of $t \in [0, 1]$.

Proof. Setting $\nabla^{t,v} = (\nabla^t)^v$, we have

$$\begin{aligned} \tau^{t,v}(f) = *d^{\nabla^{t,v}} * d^v f &= * \text{Ad}F(d * \alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{p}}] + t[\alpha_{\mathfrak{n}} \wedge * \alpha_{\mathfrak{p}}]_{\mathfrak{p}}) \\ &= * \text{Ad}F(d * \alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{p}}] + t[\alpha_{\mathfrak{p}} \wedge * \alpha_{\mathfrak{p}}]_{\mathfrak{p}}) \\ &\quad \text{since } [\mathfrak{m}, \mathfrak{p}] \subset [\mathfrak{m}, \mathfrak{h}] \subset \mathfrak{m} \\ &= * \text{Ad}F(d * \alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{p}}]). \end{aligned}$$

This gives us the equivalences (i) \Leftrightarrow (ii) \Leftrightarrow (iii) as well as the last assertion of the theorem. Moreover let us compute the complex second derivative:

$$\bar{\partial}^{\nabla^{t,v}} \partial^v f = * \text{Ad}F(\bar{\partial} \alpha'_{\mathfrak{p}} + [\alpha''_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{p}}] + [\alpha''_{\mathfrak{p}} \wedge \alpha'_{\mathfrak{p}}]_{\mathfrak{p}}).$$

Then the equivalence (ii) \Leftrightarrow (iv) follows from theorem 7.5. For this equivalence, we could also remark that $[\alpha''_{\mathfrak{p}} \wedge \alpha'_{\mathfrak{p}}]_{\mathfrak{p}} = \frac{1}{2}[\alpha_{\mathfrak{p}} \wedge \alpha_{\mathfrak{p}}]_{\mathfrak{p}}$ is in the real subspace \mathfrak{p} and that $2 \text{Im}(\bar{\partial} \alpha'_{\mathfrak{p}} + [\alpha''_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{p}}]) = d * \alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge * \alpha_{\mathfrak{p}}]$. This completes the proof. □

Now, let $f: L \rightarrow N$ be an arbitrary map from a Riemann surface into N . Then the f -pullback of the vertical torsion with respect of ∇^t is

$$\begin{aligned} f^*T^{t,v} = d^{\nabla^{t,v}} d^v f &= \text{Ad}F(d\alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge \alpha_{\mathfrak{p}}] + t[\alpha_{\mathfrak{n}} \wedge \alpha_{\mathfrak{p}}]_{\mathfrak{p}}) \\ &= \text{Ad}F(d\alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge \alpha_{\mathfrak{p}}] + t[\alpha_{\mathfrak{p}} \wedge \alpha_{\mathfrak{p}}]_{\mathfrak{p}}) \\ &= f^*(T^{0,v} + t[\phi \wedge \phi]_{[\mathfrak{p}]}) \end{aligned}$$

where $\phi: TN \rightarrow [\mathfrak{p}]$ is the projection on the vertical subbundle along the horizontal subbundle $[\mathfrak{m}]$. Therefore

$$T^{t,v} = T^{0,v} + t[\phi \wedge \phi]_{[\mathfrak{p}]}. \quad (133)$$

Moreover, recall that, according to section 4.3.1, the projection on $[\mathfrak{p}]$ of the Maurer-Cartan equation gives us the homogeneous structure equation (see equations (71), (70) and footnote 32)

$$T^{0,v} = \Phi - \frac{1}{2}[\phi \wedge \phi]_{[\mathfrak{p}]}$$

where $\Phi = -\frac{1}{2}[\psi \wedge \psi]_{[\mathfrak{p}]}$ is the homogeneous curvature form and $\psi: TN \rightarrow [\mathfrak{m}]$ is the projection on $[\mathfrak{m}]$ along $[\mathfrak{p}]$. Then we have

$$T^{t,v} = \Phi + \left(t - \frac{1}{2}\right)[\phi \wedge \phi]_{[\mathfrak{p}]}. \quad (134)$$

Therefore

Theorem 7.8 *Let us consider the same situation as in theorem 7.7.*

• *If f is flat then the strongly ∇^t -vertical harmonicity and the freedom from torsion, for f , do not depend on t , if $t \in [0, 1] \setminus \{\frac{1}{2}\}$.*

Moreover $T^{\frac{1}{2},v} = \Phi$ so that (if f is flat) strongly vertical harmonicity and vertical harmonicity with respect to $\nabla^{\frac{1}{2}}$ are equivalent.

• *If H/K is locally symmetric, i.e. $[\mathfrak{p}, \mathfrak{p}] \subset \mathfrak{k}$, then $\forall t \in [0, 1]$, $T^{t,v} = \Phi$.*

In particular, the ∇^t -vertical torsion does not depend on $t \in [0, 1]$, and thus neither does strongly harmonicity.

Now, we can conclude

Corollary 7.3 *Let us suppose now that $N = G/K$ is a (locally) $2k$ -symmetric space and that $M = G/H$ is the corresponding (locally) k -symmetric space. Then the even minimal determined system $(\text{Syst}(k, \tau))$ associated to N means that the geometric map $f: L \rightarrow N$ is horizontally holomorphic and vertically harmonic with respect to any linear connection ∇^t , $0 \leq t \leq 1$. Moreover the horizontal holomorphicity implies the flatness of f and thus its freedom from vertical torsion (with respect to any connection ∇^t , $0 \leq t \leq 1$). More precisely the (last) equation (S_k) of the system means*

$$\bar{\partial}^{\nabla^{t,v}} \partial^v f = 0$$

i.e. that f is strongly ∇^t -vertically harmonic, so that its real part means that f is vertically torsion free and its imaginary part that f is vertically harmonic.

The Riemannian case

Now, let us suppose that the reductive homogeneous space $M = G/H$ is Riemannian, and then so is $N = G/K$. In other words, we are in the situation described by 4.3.1. Let us consider the metric connections in N :

$$\overset{\text{met}}{\nabla}^t = \nabla^0 + tB^N, \quad 0 \leq t \leq 1$$

with $B^N = [\cdot, \cdot]_{[\mathfrak{n}]} + U^N$ and U^N defined by equation (15).

For any $\text{Ad}K$ -invariant subspace $\mathfrak{l} \subset \mathfrak{n}$, we will denote by $U^{\mathfrak{l}}: \mathfrak{l} \times \mathfrak{l} \rightarrow \mathfrak{l}$ the bilinear symmetric map defined by

$$\langle U^{\mathfrak{l}}(X, Y), Z \rangle = \langle [Z, X]_{\mathfrak{l}}, Y \rangle + \langle X, [Z, Y]_{\mathfrak{l}} \rangle \quad \forall X, Y \in \mathfrak{l},$$

and by $U^{[l]}$ its extension to the subbundle $[l] \subset TN$. Then we have in particular

$$U^N = U^{[n]} \quad \text{and} \quad U = U^{[p]}$$

where, let us recall it, U is defined by (66).

Now, let us project the definition equation of $\overset{\text{met}}{\nabla}{}^t$ in the vertical subbundle: we obtain $\forall V \in \mathcal{C}(TN)$,

$$\phi(\overset{\text{met}}{\nabla}{}^t V) = \nabla^0 \phi V + t \phi \circ B^N(\cdot, V).$$

Moreover, according to 4.3.1, equation (74), we have $\phi \circ B^N = \phi^* B - \Phi$ so that

$$\phi(\overset{\text{met}}{\nabla}{}^t V) = \nabla^0 \phi V + t(\phi^* B - \Phi)(\cdot, V)$$

and in particular $\forall V \in \mathcal{C}(\mathcal{V}), \forall A \in TN$,

$$\phi(\overset{\text{met}}{\nabla}_A{}^t V) = \nabla_A^0 V + t \left([\phi A, V]_{[p]} + U^{[p]}(\phi A, V) \right). \quad (135)$$

Then according to theorem 4.4 and remark 4.4, and denoting by $\overset{\text{met}}{\nabla}{}^{t,v}$ the vertical component of $\overset{\text{met}}{\nabla}{}^t$, it follows:

Theorem 7.9 • *If H/K is naturally reductive, then the connections defined by the restriction to \mathcal{V} of $\overset{\text{met}}{\nabla}{}^{t,v}, 0 \leq t \leq 1$, are all ϕ -equivalent. Therefore the vertical harmonicity, with respect to $\overset{\text{met}}{\nabla}{}^t$, is the same for all $0 \leq t \leq 1$.*

• *If H/K is locally symmetric, then all the $\overset{\text{met}}{\nabla}{}^{t,v}, 0 \leq t \leq 1$, coincide in \mathcal{V} . In particular the strongly vertical harmonicity coincides for all the connections $\overset{\text{met}}{\nabla}{}^t, 0 \leq t \leq 1$.*

The vertical torsion of $\overset{\text{met}}{\nabla}{}^t$. We have seen in 1.6 that the torsion of $\overset{\text{met}}{\nabla}{}^t$ is the same as the one of ∇^t . Now let us see what happens for the vertical torsion. The vertical torsion with respect to $\overset{\text{met}}{\nabla}{}^{t,v}$ is given, according to (135), by

$$\begin{aligned} \overset{\text{met}}{T}{}^{t,v} &= d^{\overset{\text{met}}{\nabla}{}^{t,v}} \phi = d^{\nabla^0} \phi + t[\phi \wedge \phi]_{[p]} + U^{[p]}(\phi \wedge \phi) \\ &= T^{0,v} + t[\phi \wedge \phi]_{[p]} + U^{[p]}(\phi \wedge \phi). \end{aligned}$$

But $U^{[p]}(\phi \wedge \phi) = 0$ because $U^{[p]}$ is symmetric, so that, according to (133),

$$\overset{\text{met}}{T}{}^{t,v} = T^{t,v}. \quad (136)$$

Remark 7.4 We see that the value $t = \frac{1}{2}$, i.e. the Levi-Civita connection, plays a special role according to theorem 7.8 and equation (136). Indeed for the Levi-Civita connection, we always have $\overset{\text{met}}{T}{}^{\frac{1}{2},v} = \Phi$, so that if f is flat, the strongly harmonicity and the vertical harmonicity are equivalent.

However, if H/K is (locally) symmetric, then we have $\forall t \in [0, 1], \overset{\text{met}}{T}{}^{t,v} = T^{t,v} = \Phi$, and we have even more, since all the connections $\overset{\text{met}}{\nabla}{}^{t,v}$ coincides on \mathcal{V} . Therefore the special role played by the Levi-Civita connection is shared, in this case, with all the other connections $\overset{\text{met}}{\nabla}{}^t$.

Remark 7.5 As already mentioned in section 4.3.1, the equation (74), i.e. $\phi \circ B^N = \phi^* B - \Phi$, can be obtained directly by computation (without using the general theorem 4.2, as in section 4.3.1). Let us do this. Lifting this equation we then have to prove that $\forall X, Y \in \mathfrak{g}$,

$$[X_n, Y_n]_{\mathfrak{p}} + [U^n(X_n, Y_n)]_{\mathfrak{p}} = [X_{\mathfrak{p}}, Y_{\mathfrak{p}}]_{\mathfrak{p}} + U^{\mathfrak{p}}(X_{\mathfrak{p}}, Y_{\mathfrak{p}}) + [X_{\mathfrak{m}}, Y_{\mathfrak{m}}]_{\mathfrak{p}},$$

since $[\mathfrak{m}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$. It then suffices to prove that $[U^n(X_n, Y_n)]_{\mathfrak{p}} = 0$. This equality holds. Indeed, we have $\forall Z \in \mathfrak{g}$,

$$\langle U^n(X_n, Y_n), Z_{\mathfrak{p}} \rangle = \langle [Z_{\mathfrak{p}}, X_n]_{\mathfrak{n}}, Y_{\mathfrak{m}} \rangle + \langle X_n, [Z_{\mathfrak{p}}, Y_{\mathfrak{m}}]_{\mathfrak{n}} \rangle = \langle [Z_{\mathfrak{p}}, X_{\mathfrak{m}}], Y_{\mathfrak{m}} \rangle + \langle X_{\mathfrak{m}}, [Z_{\mathfrak{p}}, Y_{\mathfrak{m}}] \rangle = 0$$

since $\text{ad}_{\mathfrak{m}} \mathfrak{p} \subset \text{ad}_{\mathfrak{m}} \mathfrak{h} \subset \mathfrak{so}(\mathfrak{m})$.

The metric geometric interpretation of the even minimal determined system. Now, according to theorem 7.9, we can conclude by rewriting corollary 7.3 in terms of the metric connection $\overset{\text{met}}{\nabla}^t$ instead of the linear connection ∇^t .

Corollary 7.4 *Let us suppose now that $N = G/K$ is a (locally) $2k$ -symmetric space and that $M = G/H$ is the corresponding (locally) k -symmetric space. Then the even minimal determined system $(\text{Syst}(k, \tau))$ associated to N means that the geometric map $f: L \rightarrow N$ is horizontally holomorphic and vertically harmonic with respect to any metric connection $\overset{\text{met}}{\nabla}^t$, $0 \leq t \leq 1$. Moreover the horizontal holomorphicity implies the flatness of f and thus its freedom from vertical torsion (with respect to any connection $\overset{\text{met}}{\nabla}^t$, $0 \leq t \leq 1$). More precisely the (last) equation (S_k) of the system means*

$$\bar{\partial} \overset{\text{met}}{\nabla}^{t,v} \partial^v f = 0$$

i.e. that f is strongly $\overset{\text{met}}{\nabla}^t$ -vertically harmonic, so that its real part means that f is vertically torsion free and its imaginary part that f is vertically harmonic.

Remark 7.6 In particular for $t = \frac{1}{2}$, we recover theorem 4.10.

7.5 (Affine) vertically (holomorphically) harmonic maps into reductive homogeneous space with an invariant Pfaffian structure

Let $N = G/K$ be a reductive homogeneous space and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$ a reductive decomposition of \mathfrak{g} . Let us suppose that \mathfrak{m} admits an $\text{Ad}K$ -invariant decomposition

$$\mathfrak{m} = \mathfrak{m}' \oplus \mathfrak{p}.$$

Then \mathfrak{p} defines a vertical subbundle $\mathcal{V} = [\mathfrak{p}]$ and \mathfrak{m}' an horizontal subbundle $\mathcal{H} = [\mathfrak{m}']$ giving a splitting $TN = \mathcal{H} \oplus \mathcal{V}$.

The curvature of the horizontal distribution \mathcal{H} is given by

$$R^{\mathcal{H}} = -[\psi, \psi]_{[\mathfrak{p}]} = -\frac{1}{2}[\psi \wedge \psi]_{[\mathfrak{p}]}$$

where $\psi: TN \rightarrow [\mathfrak{m}']$ is the projection on $[\mathfrak{m}']$ along $[\mathfrak{p}]$. We will set

$$\Phi := R^{\mathcal{H}}.$$

The vertical torsion of the linear connection ∇^t is given by $T^{t,v} = d^{(\nabla^t)^v} \phi$ and lifts into

$$\widetilde{T}^{t,v} = d\theta_{\mathfrak{p}} + [\theta_{\mathfrak{k}} \wedge \theta_{\mathfrak{p}}] + t[\theta_{\mathfrak{m}} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}} \quad (137)$$

$$= d\theta_{\mathfrak{p}} + [\theta_{\mathfrak{k}} \wedge \theta_{\mathfrak{p}}] + t[\theta_{\mathfrak{m}'} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}} + t[\theta_{\mathfrak{p}} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}}. \quad (138)$$

On the other hand, the projection on \mathfrak{p} of the Maurer-Cartan equation gives

$$d\theta_{\mathfrak{p}} + [\theta_{\mathfrak{k}} \wedge \theta_{\mathfrak{p}}] + \frac{1}{2}[\theta_{\mathfrak{m}'} \wedge \theta_{\mathfrak{m}'}]_{\mathfrak{p}} + [\theta_{\mathfrak{m}'} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}} + \frac{1}{2}[\theta_{\mathfrak{p}} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}} = 0$$

so that (138) can be written

$$\widetilde{T}^{t,v} = -\frac{1}{2}[\theta_{\mathfrak{m}'} \wedge \theta_{\mathfrak{m}'}]_{\mathfrak{p}} + (t-1)[\theta_{\mathfrak{m}'} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}} + \left(t - \frac{1}{2}\right)[\theta_{\mathfrak{p}} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}}$$

which projected in N becomes

$$T^{t,v} = \Phi + (t-1)[\psi \wedge \phi]_{[\mathfrak{p}]} + \left(t - \frac{1}{2}\right)[\phi \wedge \phi]_{[\mathfrak{p}]}.$$

We remark that the values $t = \frac{1}{2}, 1$ play special roles. In particular:

- If $[\mathfrak{m}', \mathfrak{p}]_{\mathfrak{p}} = \{0\}$ then we have $T^{\frac{1}{2},v} = \Phi$. More generally we recover equation (134) and the results of theorem 7.8 (by taking the following values in the notations $\mathfrak{m} := \mathfrak{n}$ and $\mathfrak{m}' := \mathfrak{m}$).
- If $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$ then we have $T^{1,v} = \Phi$.

Now, if the two conditions are satisfied, $[\mathfrak{m}', \mathfrak{p}]_{\mathfrak{p}} = [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$, then we have

$$\forall t \in [0, 1], \quad T^{t,v} = \Phi.$$

Now let $f: (L, j_L) \rightarrow N$ be a map from a Riemann surface into N . Let us compute the vertical tension field $\tau^{t,v}(f)$ of f with respect to ∇^t (and some Hermitian metric b in L). In order to do that, let $F: L \rightarrow G$ be a lift of f and $\alpha = F^{-1}.dF$. Then we have

$$\begin{aligned} \tau^{t,v}(f) &= *d^{\nabla^t,v} *d^v f = *\text{Ad}F(d*\alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge *\alpha_{\mathfrak{p}}] + t[\alpha_{\mathfrak{m}} \wedge *\alpha_{\mathfrak{p}}]_{\mathfrak{p}}) \\ &= *\text{Ad}F(d*\alpha_{\mathfrak{p}} + [\alpha_{\mathfrak{k}} \wedge *\alpha_{\mathfrak{p}}] + t[\alpha_{\mathfrak{m}'} \wedge *\alpha_{\mathfrak{p}}]_{\mathfrak{p}} + t[\alpha_{\mathfrak{p}} \wedge *\alpha_{\mathfrak{p}}]_{\mathfrak{p}}) \\ &= \tau^{0,v}(f) + t*[f^*\psi \wedge *(f^*\phi)]_{[\mathfrak{p}]} \\ &= \tau^{0,v}(f) + t\text{Tr}_b([f^*\psi, f^*\phi]_{[\mathfrak{p}]}) \end{aligned}$$

Now, let us consider the $\text{Ad}K$ -invariant vector subspace

$$\mathfrak{m}_* = \{X \in \mathfrak{m}' \mid [X, \mathfrak{p}]_{\mathfrak{p}} = \{0\}\}$$

and let \mathfrak{m}_1 be an $\text{Ad}K$ -invariant complement⁷⁷ of \mathfrak{m}_* in \mathfrak{m}'

$$\mathfrak{m}' = \mathfrak{m}_* \oplus \mathfrak{m}_1.$$

Then we can rewrite the ∇^t -vertical torsion in the form

$$T^{t,v} = \Phi + (t-1)[\psi_1 \wedge \phi]_{[\mathfrak{p}]} + \left(t - \frac{1}{2}\right)[\phi \wedge \phi]_{[\mathfrak{p}]} \quad (139)$$

and the ∇^t -vertical tension field (of f) in the form:

$$\tau^{t,v}(f) = \tau^{0,v}(f) + t*[f^*\psi_1 \wedge *(f^*\phi)]_{[\mathfrak{p}]} \quad (140)$$

where $\psi_1: TN \rightarrow [\mathfrak{m}_1]$ is the projection on $[\mathfrak{m}_1]$ along $[\mathfrak{m}_*] \oplus [\mathfrak{p}]$ i.e. the $[\mathfrak{m}_1]$ -component of ψ .

⁷⁷Such an $\text{Ad}K$ -invariant complement always exists if \mathfrak{m} admits an $\text{Ad}K$ -invariant non degenerated inner product, which is always the case for example if \mathfrak{g} is semisimple (take the restriction to \mathfrak{m} of the Killing form).

Definition 7.5 Let us suppose that $N = G/K$ admits a G -invariant almost complex structure \underline{J} which leaves invariant the decomposition $TN = \mathcal{V} \oplus \mathcal{H}$, that is to say the vector space \mathfrak{m} admits an $\text{Ad}K$ -invariant almost complex structure \underline{J}_0 leaving invariant the decomposition $\mathfrak{m} = \mathfrak{m}' \oplus \mathfrak{p}$. Then we will say that \underline{J} anticommutes with the reductivity term $[\psi, \phi]_{[\mathfrak{p}]}$ if

$$\underline{J}[\psi, \phi]_{[\mathfrak{p}]} = -[\underline{J}\psi, \phi]_{[\mathfrak{p}]} = -[\psi, \underline{J}\phi]_{[\mathfrak{p}]}$$

If \underline{J} anticommutes with the reductivity term then \mathfrak{m}_* is \underline{J} -invariant so that it admits a \underline{J} -invariant complement \mathfrak{m}_1 in \mathfrak{m}' .

We obtain immediately the following characterisation of the anticommutation of \underline{J} with the reductivity term.

Proposition 7.1 Let us that $N = G/K$ is endowed with a G -invariant almost complex structure \underline{J} leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$. For any \underline{J}_0 -invariant $\text{Ad}K$ -invariant subspace $\mathfrak{l} \subset \mathfrak{m}$, let us denote by \mathfrak{l}^\pm respectively the $\pm i$ -eigenspace of $\underline{J}|_{\mathfrak{l}}$. Then \underline{J} anticommutes with the reductivity term $[\psi, \phi]_{[\mathfrak{p}]}$ if and only if

$$[\mathfrak{m}'^\pm, \mathfrak{p}^\pm]_{\mathfrak{p}} \subset \mathfrak{p}^\mp \quad \text{and} \quad [\mathfrak{m}'^\pm, \mathfrak{p}^\mp]_{\mathfrak{p}} \subset \{0\}.$$

In particular, if $\mathfrak{m}_* = \{X \in \mathfrak{m}' \mid [X, \mathfrak{p}]_{\mathfrak{p}} = \{0\}\}$ admits a \underline{J} -invariant $\text{Ad}K$ -invariant complement \mathfrak{m}_1 , then these conditions are equivalent to

$$[\mathfrak{m}_1^\pm, \mathfrak{p}^\pm]_{\mathfrak{p}} \subset \mathfrak{p}^\mp \quad \text{and} \quad [\mathfrak{m}_1^\pm, \mathfrak{p}^\mp]_{\mathfrak{p}} \subset \{0\}.$$

The following theorem presents some relations between vert. hol. harmonicity, vertical harmonicity, flatness and torsion freedom.

Theorem 7.10 Let us suppose that $N = G/K$ is endowed with a G -invariant almost complex structure \underline{J} leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$ and which anticommutes with the reductivity term $[\psi, \phi]_{[\mathfrak{p}]}$. Let \mathfrak{m}_1 be a \underline{J} -invariant $\text{Ad}K$ -invariant complement in \mathfrak{m}' of $\mathfrak{m}_* = \{X \in \mathfrak{m}' \mid [X, \mathfrak{p}]_{\mathfrak{p}} = \{0\}\}$. Let $f: (L, j_L) \rightarrow N$ be a map from a Riemann surface into N , $F: L \rightarrow G$ a (local) lift of f and $\alpha = F^{-1}.dF$.

• Then if f is flat, $f^*\Phi = 0$, and $[\mathfrak{m}_1]$ -holomorphic then the following statements are equivalent:

(i) f is vert. hol. harmonic w.r.t. ∇^1 and $\underline{J}: [\bar{\partial}^{\nabla^{1,v}} \partial^v f]^{1,0} = 0$.

(ii) f is vert. hol. harmonic w.r.t. ∇^0 and $-\underline{J}: [\bar{\partial}^{\nabla^{0,v}} \partial^v f]^{0,1} = 0$.

Moreover if $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$, then these are also equivalent to

(iii) f is vertically harmonic w.r.t. ∇^1 .

• Furthermore, if $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$ and f is flat, then f is ∇^1 -torsion free so that ∇^1 -vertical harmonicity is equivalent to strongly ∇^1 -vertical harmonicity.

Proof. The ∇^1 -vertical holomorphic harmonicity is written

$$f^*T^{1,v} + \underline{J}d^{\nabla^{1,v}} * d^v f = 0 \tag{141}$$

but $f^*T^{1,v} = \frac{1}{2}f^*[\phi \wedge \phi]_{[\mathfrak{p}]}$ and $f^*T^{0,v} = -[f^*\psi \wedge f^*\phi]_{[\mathfrak{p}]} - \frac{1}{2}f^*[\phi \wedge \phi]_{[\mathfrak{p}]}$ whereas

$$d^{\nabla^{1,v}} * d^v f = d^{\nabla^{0,v}} * d^v f + [f^*\psi \wedge (*f^*\phi)]_{[\mathfrak{p}]}$$

so that

$$f^*T^{1,v} + \underline{J}d^{\nabla^{1,v}} * d^v f = \frac{1}{2}[f^*\phi \wedge f^*\phi]_{[\mathfrak{p}]} + \underline{J}d^{\nabla^{0,v}} * d^v f + \underline{J}[f^*\psi \wedge (*f^*\phi)]_{[\mathfrak{p}]}.$$
 (142)

Now let us use the fact that \underline{J} anticommutes with $[\psi, \phi]_{[\mathfrak{p}]}$:

$$\begin{aligned} \underline{J}[f^*\psi \wedge *(f^*\phi)]_{[\mathfrak{p}]} &= -[\underline{J}(f^*\psi_1) \wedge *(f^*\phi)]_{[\mathfrak{p}]} \\ &= [*(f^*\psi_1) \wedge *(f^*\phi)] \quad \text{because } f \text{ is } [\mathfrak{m}_1]\text{-holomorphic,} \\ &= [f^*\psi_1 \wedge f^*\phi] = [f^*\psi \wedge f^*\phi]. \end{aligned}$$

Therefore, injecting this in (142), we obtain

$$f^*T^{1,v} + \underline{J}d^{\nabla^{1,v}} * d^v f = -\left(f^*T^{0,v} - \underline{J}d^{\nabla^{0,v}} * d^v f\right).$$

This proves the equivalence (i) \Leftrightarrow (ii). Now, if we suppose that $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$, then $f^*T^{1,v} = 0$. Therefore the ∇^1 -vertical holomorphic harmonicity (141) is equivalent to the ∇^1 -vertical harmonicity $d^{\nabla^{1,v}} * d^v f = 0$. This completes the proof. \square

Now, let us see how the vertical holomorphic harmonicity is written in terms of the Maurer-Cartan form α of a lift F of $f: L \rightarrow N$. We obtain immediately

Proposition 7.2 *Let us suppose that $N = G/K$ is endowed with a G -invariant almost complex structure \underline{J} leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$. Then $f: L \rightarrow N$ is vert. hol. harmonic w.r.t. ∇^0 and $-\underline{J}$ if and only if*

$$\bar{\partial}\alpha'_{\mathfrak{p}+} + [\alpha''_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{p}+}] = 0.$$

Moreover, if \underline{J} anticommutes with the reductivity term $[\psi, \phi]_{[\mathfrak{p}]}$ and \mathfrak{m}_* admits an $\text{Ad}K$ -invariant \underline{J} -invariant complement \mathfrak{m}_1 in \mathfrak{m}' , then a $[\mathfrak{m}_1]$ -holomorphic map $f: L \rightarrow N$ is vert. hol. harmonic w.r.t. ∇^1 and \underline{J} if and only if

$$\bar{\partial}\alpha'_{\mathfrak{p}+} + [\alpha''_{\mathfrak{k}} \wedge \alpha'_{\mathfrak{p}+}] + \left[\alpha''_{\mathfrak{m}_1^-} \wedge \alpha'_{\mathfrak{p}^-}\right]_{\mathfrak{p}} + [\alpha''_{\mathfrak{p}} \wedge \alpha'_{\mathfrak{p}}]_{\mathfrak{p}+} = 0.$$

Now, let us suppose that $N = G/K$ is a (locally) $(2k+1)$ -symmetric space, then the $\text{Ad}K$ -invariant decomposition $\mathfrak{m} = \mathfrak{m}' \oplus \mathfrak{p}$ is given by $\mathfrak{p} = \mathfrak{m}_k$ and $\mathfrak{m}' = \bigoplus_{j=1}^{k-1} \mathfrak{m}_j$ with the notations of 2.1.2. Moreover according to the commutation relations $[\mathfrak{g}_i^C, \mathfrak{g}_j^C] \subset \mathfrak{g}_{i+j}^C$, we have

$$\mathfrak{m}_* = \{X \in \mathfrak{m}', [X, \mathfrak{p}]_{\mathfrak{p}} = \{0\}\} = \bigoplus_{j=2}^{k-1} \mathfrak{m}_j$$

so that \mathfrak{m}_1 is an $\text{Ad}G_0$ -invariant supplement to \mathfrak{m}_* . Moreover N is endowed naturally with its canonical almost complex structure \underline{J} defined in 2.1.2, which leaves invariant all the \mathfrak{m}_j and thus the subspaces $\mathfrak{m}_1, \mathfrak{m}_*, \mathfrak{p}$. Furthermore, using once again the commutation relations, one can see that \underline{J} anticommutes with the reductivity term $[\psi, \phi]_{[\mathfrak{p}]}$. Finally, let us remark that $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = [\mathfrak{m}_k, \mathfrak{m}_k]_{\mathfrak{p}} = \{0\}$. Now, the theorem 7.10 can be applied.

Corollary 7.5 *Let us suppose that $N = G/K$ is a (locally) $(2k+1)$ -symmetric space endowed with its canonical almost complex structure \underline{J} , and with the \underline{J} -invariant splittings $TN = [\mathfrak{m}'] \oplus [\mathfrak{p}]$ and $[\mathfrak{m}'] = [\mathfrak{m}_1] \oplus [\mathfrak{m}_*]$. Let $f: L \rightarrow N$ be a map, $F: L \rightarrow G$ a lift of f and $\alpha = F^{-1}.dF$. Then if f is flat, $f^*\Phi = 0$, and $[\mathfrak{m}_1]$ -holomorphic then f is ∇^1 -vertically torsion free $f^*T^{1,v} = 0$, and the following statements are equivalent*

- (i) f is vert. hol. harmonic w.r.t. ∇^1 and $\underline{\mathbf{J}} : [\bar{\partial}^{\nabla^1, v} \partial^v f]^{1,0} = 0$.
- (ii) f is vert. hol. harmonic w.r.t. ∇^0 and $-\underline{\mathbf{J}} : [\bar{\partial}^{\nabla^0, v} \partial^v f]^{0,1} = 0$.
- (iii) f is vertically harmonic w.r.t. ∇^1 .
- (iv) f is strongly vertically harmonic w.r.t. ∇^1 .

Now, let us apply proposition 7.2.

Proposition 7.3 *Let us suppose that $N = G/K$ is a (locally) $(2k + 1)$ -symmetric space. Then $f: L \rightarrow N$ is vert. hol. harmonic w.r.t. ∇^0 and $-\underline{\mathbf{J}}$ if and only if*

$$\bar{\partial} \alpha'_k + [\alpha''_0 \wedge \alpha'_k] = 0.$$

Moreover, if $f: L \rightarrow N$ is flat and $[\mathfrak{m}_1]$ -holomorphic, then it is vert. hol. harmonic w.r.t. ∇^1 and $\underline{\mathbf{J}}$ if and only if

$$\bar{\partial} \alpha'_{-k} + [\alpha''_0 \wedge \alpha'_{-k}] + [\alpha''_1 \wedge \alpha'_k] = 0.$$

Furthermore, as in the even case (i.e. $N = G/K$ is (locally) $2k$ -symmetric), the horizontal holomorphicity implies the flatness.

Proposition 7.4 *Let us suppose that $N = G/K$ is a (locally) $(2k + 1)$ -symmetric space. Then if $f: L \rightarrow N$ is horizontally holomorphic (i.e. $[\mathfrak{m}']$ -holomorphic) then f is flat $f^* \Phi = 0$.*

Proof. It follows from the equation

$$\tilde{\Phi} = -[\theta_{\mathfrak{m}'}, \theta_{\mathfrak{m}'}]_{\mathfrak{p}} = - \sum_{\substack{i+j=k \\ 1 \leq |i|, |j| \leq k-1}} [\theta_i, \theta_j]_{\mathfrak{g}_k},$$

and the fact that if $i + j = k$ and $1 \leq |i|, |j| \leq k - 1$, then i and j have the same sign. This completes the proof. \square

Now let us conclude with the following geometric interpretation of the odd minimal determined system.

Corollary 7.6 *Let us suppose that $N = G/K$ is a (locally) $(2k + 1)$ -symmetric space. Then the odd minimal determined system $(\text{Syst}(k + 1, \tau))$ associated to N means that the geometric map $f: L \rightarrow N$ is horizontally holomorphic and vertically harmonic w.r.t. the linear connection ∇^1 . Moreover the horizontal holomorphicity implies the flatness of f and its freedom from ∇^1 -vertical torsion, $f^* T^{1, v} = 0$.*

More pecisely, the (last) equation (S_{k+1}) of the system (which lies in \mathfrak{g}_k) means the vert. hol. harmonicity of f w.r.t. ∇^0 and $-\underline{\mathbf{J}}$

$$[\bar{\partial}^{\nabla^0, v} \partial^v f]^{0,1} = 0$$

whereas the equation (S_k) (which lies in \mathfrak{g}_{-k}) means the vert. hol. harmonicity of f w.r.t. ∇^1 and $\underline{\mathbf{J}}$

$$[\bar{\partial}^{\nabla^1, v} \partial^v f]^{1,0} = 0$$

Moreover the sums $(S_k)+(S_{k+1})$ (which lies in \mathfrak{m}_k) means (taking account of the $[\mathfrak{m}_1]$ -holomorphicity $\alpha''_{-1} = 0$) the strongly vertical harmonicity of f w.r.t. ∇^1 :

$$\bar{\partial}^{\nabla^1, v} \partial^v f = 0,$$

so that its real part means that f is ∇^1 -vertically torsion free and its imaginary part that f is ∇^1 -vertically harmonic.

All the other equations of the system, (S_j) , $0 \leq j \leq k-1$ are (after having taken account of the horizontal holomorphicity $\alpha''_{-j} = 0$, $1 \leq j \leq k-1$) nothing but the projections on the subspace \mathfrak{g}_{-j} , $1 \leq j \leq k-1$ of the Maurer-Cartan equation (which means the existence of the geometric map f corresponding to α).

Proof. The first assertion has been proved in section 2.3.2, theorem 2.6. Moreover, the ∇^1 -vertical harmonicity will follow from the next assertions, according to corollary 7.5.

The second assertion follows from proposition 7.4 and theorem 7.10. The third assertion follows from propositions 2.7 and 7.3. The fourth assertion has been proved in subsection 2.3.2, paragraph *The strictly minimal determined case*. Finally, the fifth assertion follows from proposition 7.4. This completes the proof. \square

Strongly vertical harmonicity w.r.t. ∇^t . Let us see what the strongly harmonicity w.r.t. to ∇^t , with $t \in [0, 1] \setminus \{1\}$, means.

We have seen that the tension field of a map $f: L \rightarrow N = G/K$, with respect to ∇^t , does not depend on $t \in [0, 1]$ (see theorem 7.1). Let us set $\tau(f) := \tau^t(f)$.

Proposition 7.5 *Let $f: L \rightarrow N = G/K$ be a map.*

• *Then we have*

$$\tau^{0, v}(f) = [\tau(f)]^v = 2 \left[\bar{\partial}^{\nabla^{\frac{1}{2}}} \partial f \right]^v.$$

• *If $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$ and f is flat, then $f^*T^{1, v} = 0$ i.e. $\tau^{1, v}(f) = 2 \bar{\partial}^{\nabla^1, v} \partial f$.*

• *Moreover, under these hypothesis⁷⁸, let \mathfrak{m}_1 be an $\text{Ad}K$ -invariant complement of \mathfrak{m}_* , in \mathfrak{m}' . Then, the following statements are equivalent:*

(i) *$f^*T^{t, v} = 0$ for one $t \in [0, 1] \setminus \{1\}$,*

(ii) *$f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$,*

(iii) *$\tau^{t, v}(f) = \tau^{0, v}(f)$ for one $t \in [0, 1] \setminus \{0\}$.*

Proof. In the first point, the first equality comes from the fact that ∇^0 leaves invariant the splitting $TN = \mathcal{V} \oplus \mathcal{H}$, since then $(\nabla^0)^v d^v f = [\nabla^0 df]^v$. The second equality follows from theorem 5.1 and the fact that $T^{\frac{1}{2}} = 0$.

The second point follows from theorem 7.10. The third point follows from the equations (139) and (140). This completes the proof. \square

Corollary 7.7 *Let us suppose that $N = G/K$ is endowed with a G -invariant almost complex structure $\underline{\mathbf{J}}$ leaving invariant the decomposition $TN = \mathcal{H} \oplus \mathcal{V}$ and which anticommutes with the reductivity term $[\psi, \phi]_{[\mathfrak{p}]}$. We also suppose that there exists a $\underline{\mathbf{J}}$ -invariant $\text{Ad}K$ -invariant complement \mathfrak{m}_1 in \mathfrak{m}' of \mathfrak{m}_* . Lastly, we suppose that $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$.*

Let $f: L \rightarrow N = G/K$ be a map which is flat. Let $F: L \rightarrow N$ be a lift of f and $\alpha = F^{-1}.dF$. Then the following statements are equivalent

⁷⁸i.e. the hypothesis $[\mathfrak{p}, \mathfrak{p}] = \{0\}$ and f is flat.

- (i) $\bar{\partial}^{\nabla^{t,v}} \partial^v f = 0$ for one $t \in [0, 1] \setminus \{1\}$,
- (ii) $[\bar{\partial}^{\nabla^t} \partial f]^v = 0$ for one $t \in [0, 1] \setminus \{\frac{1}{2}\}$,
- (iii) $\tau^{1,v}(f) = 0$ and $f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$,
- (iv) $[\tau(f)]^v = 0$ and $f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$,
- (v) $[\bar{\partial}^{\nabla^{t,v}} \partial^v f]^{1,0} = 0$ for one $t \in [0, 1]$ and $f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$,
- (vi) $[\bar{\partial}^{\nabla^{1-t,v}} \partial^v f]^{0,1} = 0$ for one $t \in [0, 1]$ and $f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$,
- (vii) f is a geometric solution of the first elliptic integrable system associated to the $\text{ad}\mathfrak{k}$ -invariant decomposition $\mathfrak{g} = (\mathfrak{k} \oplus \mathfrak{m}') \oplus \mathfrak{p}$, i.e. the 1-form $\beta_\lambda = \lambda^{-1} \alpha'_\mathfrak{p} + (\alpha_\mathfrak{k} + \alpha_{\mathfrak{m}'}) + \lambda \alpha''_\mathfrak{p}$ satisfies the zero curvature equation

$$d\beta_\lambda + \frac{1}{2}[\beta_\lambda \wedge \beta_\lambda] = 0 \quad \forall \lambda \in \mathbb{C}^*.$$

Proof. The statement (i) is equivalent to the statement: " $f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$ and $\tau^{t,v}(f) = 0, \forall t \in [0, 1]$ ", according to proposition 7.5 and theorem 7.5. Moreover, since $\widetilde{[T^t]^v} = (2t-1)[\theta_{\mathfrak{m}}, \theta_{\mathfrak{m}}]_{\mathfrak{p}} = (2t-1)([\theta_{\mathfrak{m}'} \wedge \theta_{\mathfrak{m}'}]_{\mathfrak{p}} + \frac{1}{2}[\theta_{\mathfrak{m}_1} \wedge \theta_{\mathfrak{p}}]_{\mathfrak{p}})$, we have

$$[T^t]^v = (2t-1)(\Phi + [\psi_1 \wedge \phi]_{[\mathfrak{p}]}) .$$

Therefore, (ii) \Leftrightarrow ($[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$ and $[\tau(f)]^v = 0$), which is nothing but (iv). Moreover, according to theorem 7.6 and equations (139) and (140), we have (v) \Leftrightarrow (vi) \Leftrightarrow ($f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0$ and $\tau^{t,v}(f) = 0, \forall t \in [0, 1]$). Moreover, proposition 7.5 gives us the equivalences

$$(iii) \Leftrightarrow (iv) \Leftrightarrow (f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0 \text{ and } \tau^{t,v}(f) = 0, \forall t \in [0, 1]).$$

We have then proved the equivalence between the six first assertions. Finally, we have

$$(vii) \Leftrightarrow \begin{cases} \bar{\partial} \alpha'_\mathfrak{p} + [\alpha_\mathfrak{k} \wedge \alpha'_\mathfrak{p}] = 0 \\ [\alpha_{\mathfrak{m}'} \wedge \alpha'_\mathfrak{p}] = 0 \\ d(\alpha_\mathfrak{k} + \alpha_{\mathfrak{m}'}) + [\alpha'_\mathfrak{p} \wedge \alpha''_\mathfrak{p}] + \frac{1}{2}[(\alpha_\mathfrak{k} + \alpha_{\mathfrak{m}'}) \wedge (\alpha_\mathfrak{k} + \alpha_{\mathfrak{m}'})] = 0 \end{cases} \quad (143)$$

Moreover, we have $[\alpha'_\mathfrak{p} \wedge \alpha''_\mathfrak{p}]_{\mathfrak{p}} = 0$, since $[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = \{0\}$. Furthermore, $[(\alpha_\mathfrak{k} + \alpha_{\mathfrak{m}'}) \wedge (\alpha_\mathfrak{k} + \alpha_{\mathfrak{m}'})]_{\mathfrak{p}} = 0$ because $[\mathfrak{m}', \mathfrak{p}]_{\mathfrak{p}} = 0$ and the flatness of f means that $[\alpha_{\mathfrak{m}'} \wedge \alpha_{\mathfrak{m}'}]_{\mathfrak{p}} = 0$. Therefore, the last equation in (143) is nothing but $[\text{MC}]_{\mathfrak{k} \oplus \mathfrak{m}'}$, the projection of the Maurer-Cartan equation on $\mathfrak{k} \oplus \mathfrak{m}'$. Hence, we conclude that

$$(vii) \Leftrightarrow (\tau^{0,v}(f) = 0 \text{ and } f^*[\psi_1 \wedge \phi]_{[\mathfrak{p}]} = 0)$$

This completes the proof. □

Now, we come back to the case of a (locally) $(2k+1)$ -symmetric space.

Corollary 7.8 *Let us suppose that $N = G/K$ is a (locally) $(2k+1)$ -symmetric space. Let $f: L \rightarrow N = G/K$ be a map, $F: L \rightarrow N$ be a lift of f and $\alpha = F^{-1}.dF$. Then the following statements are equivalent:*

- (i) f is horizontally holomorphic and strongly ∇^t -vertically harmonic for one $t \in [0, 1] \setminus \{1\}$,
- (ii) f is a geometric solution of $(\text{Syst}(k+1, \tau))$ and $[\alpha_1 \wedge \alpha_k] = 0$,
- (iii) f is horizontally holomorphic and is a geometric solution of the first elliptic integrable system associated to the adg_0 -invariant decomposition $\mathfrak{g} = (\mathfrak{g}_0 \oplus \mathfrak{m}') \oplus \mathfrak{m}_k$.

Proof. According to corollary 7.7 and corollary 7.6, we have (i) \Leftrightarrow (iii). Moreover, according to corollaries 7.6 and 7.7, the equivalent statements (i) and (iii) are also equivalent to (ii). \square

7.6 The intermediate determined systems.

7.6.1 The odd case.

Here, we consider a (locally) $(2k+1)$ -symmetric space, $N = G/K$, endowed with its canonical almost complex structure \underline{J} and its canonical connection ∇^0 . Let $k+1 \leq m \leq 2k$ be an integer and set $\underline{m} = 2k - m$. Then we consider the G -invariant splitting $TN = \mathcal{H}^m \oplus \mathcal{V}^m$, where

$$\mathcal{H}^m = \bigoplus_{j=1}^{\underline{m}} [\mathfrak{m}_j] \quad \text{and} \quad \mathcal{V}^m = \bigoplus_{j=\underline{m}+1}^k [\mathfrak{m}_j].$$

Let us remark that this splitting is also \underline{J} -invariant and ∇^0 -parallel. According to proposition 2.3 and theorem 5.15, we have the following.

Proposition 7.6 *Let $N = G/K$ be a (locally) $(2k+1)$ -symmetric space endowed with its canonical almost complex structure \underline{J} and its canonical connection ∇^0 . Let m be an integer such that $k+1 \leq m \leq 2k$. Then the associated determined system, $\text{Syst}(m, \tau)$ means that the geometric map $f: L \rightarrow N$ is \star -stringy harmonic and \mathcal{H}^m -holomorphic.*

We will need the following definition.

Definition 7.6 *Let (N, J) be an almost complex manifold endowed with a linear connection ∇ . Let $E \subset TN$ be some subbundle. We will say that a map $f: L \rightarrow N$ from a Riemann surface into N is E - \star -stringy harmonic if*

$$[-\tau_g(f) + (J \star T)_g(f)]^E = 0.$$

where g is a Hermitian metric on L . If E inherits the name of vertical or horizontal subbundle then we will say that f is vertically, resp. horizontally \star -stringy harmonic.

Then the proposition 2.9 gives us:

Theorem 7.11 *Let $N = G/K$ be a (locally) $(2k+1)$ -symmetric space endowed with its canonical almost complex structure \underline{J} and its canonical connection ∇^0 . Let m be an integer such that $k+1 \leq m \leq 2k$. Then the associated determined system, $\text{Syst}(m, \tau)$ means that the geometric map $f: L \rightarrow N$ is \mathcal{H}^m -holomorphic and vertically \star -stringy harmonic.*

Proof. Let us come back to section 2.4, then the equation (E_m) , in theorem 2.7, means that f is \star -stringy harmonic, according to theorem 5.15 and its proof. Therefore, the projection $[(E)]_{\mathfrak{p}}$ in proposition 2.9 means that f is vertically \star -stringy harmonic. Then the proposition 2.9 allows us to conclude. This completes the proof. \square

In fact this theorem is a particular case of a more general result:

Proposition 7.7 *Let (N, J, ∇) be an almost complex affine manifold, with $TN = \overline{\mathcal{H}} \oplus \overline{\mathcal{V}}$ a J -invariant, ∇ -parallel splitting. Let T be the torsion of ∇ . Let us suppose that*

$$T^{++}(\overline{\mathcal{V}}, \overline{\mathcal{V}}) \subset \overline{\mathcal{V}} \quad \text{and} \quad T^{++}(\overline{\mathcal{H}}, \overline{\mathcal{V}}) \subset \overline{\mathcal{V}}. \quad (144)$$

Let (L, j) be a Riemann surface. Then any $\overline{\mathcal{H}}$ -holomorphic map $f: L \rightarrow N$ is horizontally \star -stringy harmonic. Therefore any horizontally holomorphic map $f: L \rightarrow N$ is \star -stringy harmonic if and only if it is vertically \star -stringy harmonic.

Proof. Using the fact that the splitting is ∇ -parallel and the horizontal holomorphicity, we have

$$[\tau^\nabla(f)]^{\overline{\mathcal{H}}} = d^\nabla * [df]^{\overline{\mathcal{H}}} = -Jd^\nabla [df]^{\overline{\mathcal{H}}} = -JT^{\overline{\mathcal{H}}}(df, df).$$

Moreover, recall that $J\star T = J(T^{++} - T^{--} - T^{1,1})$. We then have to prove that $f^*(T^{++})^{\overline{\mathcal{H}}} = 0$, which will imply the horizontal \star -stringy harmonic of f . Moreover, remark that we have

$$T^{\varepsilon, \varepsilon}(T^{1,0}N, T^{0,1}N) = 0, \quad \forall \varepsilon \in \mathbb{Z}_2.$$

Therefore, since f is horizontally holomorphic, we have $T^{\varepsilon, \varepsilon}([df]^{\overline{\mathcal{H}}}, [df]^{\overline{\mathcal{H}}}) = 0$. Furthermore, according to the hypothesis (144) on T^{++} , we conclude that $f^*(T^{++})^{\overline{\mathcal{H}}} = 0$. This completes the proof. \square

It is not difficult to see that a (locally) $(2k+1)$ -symmetric space endowed with its canonical almost complex structure \underline{J} , its canonical connection ∇^0 and the splitting $TN = \mathcal{H}^m \oplus \mathcal{V}^m$, satisfies the hypothesis of the previous proposition.

7.6.2 The even case.

Here, we consider a (locally) $2k$ -symmetric space, $N = G/K$, endowed with its canonical f -structure F and its canonical connection ∇^0 . Let $k \leq m \leq 2k-1$ be an integer and set $\underline{m} = 2k-1-m$. Then we consider the G -invariant splitting $TN = \mathcal{H}^m \oplus \mathcal{V}^m$, where

$$\mathcal{H}^m = \bigoplus_{j=1}^{\underline{m}} [\mathfrak{m}_j] \quad \text{and} \quad \mathcal{V}^m = \bigoplus_{j=\underline{m}+1}^k [\mathfrak{m}_j],$$

and $\mathfrak{m}_k = \mathfrak{g}_k$. This splitting is also \underline{J} -invariant and ∇^0 -parallel.

According to proposition 2.3 and theorem 6.11, we have the following.

Proposition 7.8 *Let $N = G/K$ be a (locally) $2k$ -symmetric space endowed with its canonical f -structure F and its canonical connection ∇^0 . Let m be an integer such that $k \leq m \leq 2k-1$. Then the associated determined system, $\text{Syst}(m, \tau)$ means that the geometric map $f: L \rightarrow N$ is \star -stringy harmonic and \mathcal{H}^m -holomorphic.*

We will need the following definition.

Definition 7.7 *Let (N, F) be a f -manifold endowed with a linear connection ∇ , and $E \subset TN$ some subbundle. We will say that a map $f: L \rightarrow N$ from a Riemann surface into N is E - \star -stringy harmonic if*

$$[-\tau_g(f) + (F \star T)_g(f)]^E = 0.$$

where g is a Hermitian metric on L . If E inherits the name of vertical or horizontal subbundle then we will say that f is vertically, resp. horizontally \star -stringy harmonic.

Then the proposition 2.9 gives us:

Theorem 7.12 *Let $N = G/K$ be a (locally) $2k$ -symmetric space endowed with its canonical f -structure F and its canonical connection ∇^0 . Let m be an integer such that $k \leq m \leq 2k - 1$. Then the associated determined system, $\text{Syst}(m, \tau)$ means that the geometric map $f: L \rightarrow N$ is \mathcal{H}^m -holomorphic and vertically \star -stringy harmonic.*

Proof. It is analogous to the one of theorem 7.11. \square

In fact this theorem is a particular case of a more general result:

Proposition 7.9 *Let (N, F, ∇) be a f -manifold endowed with a f -connection. As usual, we set $\mathcal{H} = \text{Im } F$ and $\mathcal{V} = \ker F$. Suppose that $R_{\mathcal{V}} = 0$. Moreover, let $TN = \overline{\mathcal{H}} \oplus \overline{\mathcal{V}}$ be a F -invariant, ∇ -parallel splitting, such that $\overline{\mathcal{H}} \subset \mathcal{H}$ and $\overline{\mathcal{V}} \supset \mathcal{V}$. Let us set $\overline{T} := T_{\overline{\mathcal{H}}^2}^{\mathcal{H}}$, where T is the torsion of ∇ . Let us suppose that*

$$\overline{T}^{++}(\overline{\mathcal{V}} \cap \mathcal{H}, \overline{\mathcal{V}} \cap \mathcal{H}) \subset \overline{\mathcal{V}} \cap \mathcal{H} \quad \text{and} \quad \overline{T}^{++}(\overline{\mathcal{H}}, \overline{\mathcal{V}} \cap \mathcal{H}) \subset \overline{\mathcal{V}} \cap \mathcal{H}.$$

Suppose also that $T_{\overline{\mathcal{H}} \times \mathcal{V}}^{\mathcal{H}}$ is pure⁷⁹.

Let (L, j) be a Riemann surface. Then any $\overline{\mathcal{H}}$ -holomorphic map $f: L \rightarrow N$ is horizontally \star -stringy harmonic. Therefore any horizontally holomorphic map $f: L \rightarrow N$ is \star -stringy harmonic if and only if it is vertically \star -stringy harmonic.

Proof. It is analogous to the one of theorem 7.11. \square

It is not difficult to see that a (locally) $2k$ -symmetric space endowed with its canonical f -structure F , its canonical connection ∇^0 and the splitting $TN = \mathcal{H}^m \oplus \mathcal{V}^m$, satisfies the hypothesis of the previous proposition.

7.6.3 Sigma model with a Wess-Zumino term.

We have seen that the maximal determined system has an interpretation in terms of a sigma models with a Wess-Zumino term defined by a 3-form H . In fact, more generally, let $m_{k'} \leq m \leq k' - 1$, and let us consider the splitting $TN = \mathcal{H}^m \oplus \mathcal{V}^m$ defined above. Then one can prove that any m -th determined system is the Euler-Lagrange equation w.r.t. vertical variations (i.e. in \mathcal{V}^m) of the following functional

$$E^{\overline{\mathcal{V}}}(f) = \frac{1}{2} \int_L |d^{\overline{\mathcal{V}}} f|^2 d\text{vol}_g + \int_B H^{\overline{\mathcal{V}}}$$

where $d^{\overline{\mathcal{V}}} f = [df]^{\mathcal{V}^m}$, $H^{\overline{\mathcal{V}}} = H - \overline{H} = H|_{\mathcal{S}(\mathcal{V}^m, \mathcal{H}^m)}$, $\overline{H} = H|_{(\mathcal{H}^m)^3}$, and B is a 3-submanifold of N with boundary $\partial B = f(L)$.

We will come back to this in [35].

7.7 Some remarks about the twistorial interpretation.

7.7.1 The even case.

We have seen that in the even case, each geometric property of the geometric map $f: L \rightarrow N = G/G_0$ like horizontal holomorphicity, vertical harmonicity, stringy harmonicity is naturally translated to the same geometric property of the twistor lift $J: L \rightarrow \mathcal{Z}_{2k,2}^{\alpha}(M, J_2)$ in the twistor space $\mathcal{Z}_{2k,2}^{\alpha}(M, J_2)$. Moreover, the very particular structure of homogeneous fibre f -bundle of $N + G/G_0$ can be realised as a subbundle of the universal homogeneous fibre f -bundle $\mathcal{Z}_{2k,2}^{\alpha}(M, J_2)$.

⁷⁹i.e. $T_{\overline{\mathcal{H}} \times \mathcal{V}}^{\mathcal{H}}$ anticommutes with \overline{J} , see definition 6.13.

Universal means that we can define it for any Riemannian manifold (M, g) endowed with a global k -structure J_2 (and a metric connection $\bar{\nabla}$). Remark that since $\mathcal{Z}_{2k,2}^\alpha(M, J_2)$ is a complete reduction of $\mathcal{Z}_{2k}(M)$ (w.r.t. to structures of the homogeneous fibre f -bundles defined by the canonical connection of M) then we can also as well embed N in $\mathcal{Z}_{2k}(M)$ which is more universal because defined for any Riemannian manifold (M, g) (endowed with a metric connection $\bar{\nabla}$). Then all the geometric properties below are also preserved under this embedding and hold in $\mathcal{Z}_{2k}(M)$. However, it is perhaps better to use the universal homogeneous fibre f -bundle containing N which looks like the most to N and which is also the smallest one (a kind of "universal homogeneous fibre f -bundle closure"). This exists and this is $\mathcal{Z}_{2k,2}^\alpha(M, J_2)$. For example, $\mathcal{Z}_{2k,2}^\alpha(M, J_2)$ has a symmetric fibre like N , whereas $\mathcal{Z}_{2k}(M)$ has a $2k$ -symmetric fibre.

7.7.2 The odd case.

In the odd case the use of the twistor space $\mathcal{Z}_{2k+1}(N)$ is less pertinent than in the even case. Indeed in the even case, we had some particular fibration that twistor space allows to realise more universally as some bundle of endomorphisms over M . Here we do not have this problem of fibration and therefore do not need a priori the twistor space. In the odd case, we have a canonical section⁸⁰ $J_1: G/G_0 \rightarrow \mathcal{Z}_{2k+1}(G/G_0)$, which allows to duplicate each geometric property satisfied by the geometric map $f: L \rightarrow N$ into 2 "identical" properties in each subbundle \mathcal{H} and \mathcal{V} of the tangent bundle of the twistor space.

However, the twistor space $\mathcal{Z}_{2k+1}(N)$ is still universal since it can be defined for any Riemannian manifold (N, h) (endowed with some metric connection $\bar{\nabla}$). Moreover, under the embedding defined by the canonical section $J_1: G/G_0 \rightarrow \mathcal{Z}_{2k+1}(G/G_0)$, any geometric property like \underline{J} -holomorphicity, $F^{[m]}$ -holomorphicity, vertical harmonicity, stringy harmonicity and so on is preserved and hold for the twistor lift $J = f^* J_1: L \rightarrow \mathcal{Z}_{2k+1}(N)$ in the twistor space.

We have to precise that the structure of homogeneous fibre bundle of $\mathcal{Z}_{2k+1}(N)$ is defined with respect to⁸¹ ∇^t . In particular, this connection defines a splitting $TN = \mathcal{H} \oplus \mathcal{V}$. Then, we can define a canonical almost complex structure $\check{J} := ((d\pi)^* \underline{J})|_{\mathcal{H}} \oplus \pi^*(\underline{\text{Ad}} J_1)$. That is to say on the horizontal subbundle we take complex structure defined by (the lift of) \underline{J} and moreover the fibre is $2k + 1$ -symmetric and is therefore endowed with a canonical almost complex structure, that we endow with the vertical subbundle.

Then, with respect to this almost complex structure, $J_1: (G/G_0, \underline{J}) \rightarrow (\mathcal{Z}_{2k+1}(G/G_0), \check{J})$ is holomorphic.

We will come back to that in [35].

⁸⁰Defined by lemma 3.3

⁸¹Recall that ∇^0 is metric w.r.t. any G -invariant metric, whereas ∇^t , for $t \neq 0$ is metric when N is endowed with a naturally reductive metric

8 Appendix

8.1 Vertical harmonicity

Theorem 8.1 *Let us consider the situation described by example 4.1 and suppose that $\pi: N \rightarrow M$ is a Riemannian submersion and $u: L \rightarrow M$ is an isometry. Then $f: L \rightarrow N$ is vertically harmonic if and only if the corresponding section $\tilde{f}: L \rightarrow u^*N$ is a harmonic section. Furthermore $f: L \rightarrow N$ is harmonic if and only if $\tilde{f}: L \rightarrow u^*N$ is harmonic and $[\tau(f)]_{u_*(TL)^\perp}^{\mathcal{H}} = 0$ i.e. the component of the tension field in the subspace of \mathcal{H} corresponding by the isometry $d\pi|_{\mathcal{H}}$ to the normal bundle $u_*(TL)^\perp$ in TM , vanishes, or equivalently $[d\pi(\tau(f))]_{u_*(TL)^\perp} = 0$.*

Proof. The Levi-Civita in u^*N is the orthogonal projection of the Levi-Civita connection in $L \times N$, on the tangent bundle $T(u^*N)$. Let us determine this orthonormal projection. First let us express clearly what is the tangent subbundle $T(u^*N)$ in $T(L \times N)$.

$$T_{(l,n)}(u^*N) = \{(\xi, \eta) \in T_{(l,n)}L \times N \mid d\pi(\xi) = d\pi(\eta)\}.$$

Let us do some identifications. First an usual one: consider that TL is a subbundle of $TM|_L$ (and forget the "u*" in $u_*(TL)$), secondly: we consider that $\pi^*TM = \mathcal{H}$, identifying these by the isometry $d\pi|_{\mathcal{H}}$, so that we will write $\mathcal{H}|_{\pi^{-1}(L)} = \pi^*TL \oplus \pi^*TL^\perp$, where TL^\perp is the normal bundle of L in M . Moreover, for any $\eta \in TN|_{\pi^{-1}(L)}$ let us write its decomposition following $TN_{\pi^{-1}(L)} = \pi^*TL \oplus \pi^*TL^\perp \oplus \mathcal{V}_{\pi^{-1}(L)}$ as

$$\eta = \eta_{TL}^{\mathcal{H}} + \eta_{TL^\perp}^{\mathcal{H}} + \eta^{\mathcal{V}}.$$

Then under the previous identifications, we have

$$\begin{aligned} T_{(l,n)}(u^*N) &= \{(\xi, \eta) \in T_lL \times T_nN \mid \eta_{TL}^{\mathcal{H}} = \xi, \eta_{TL^\perp}^{\mathcal{H}} = 0\} \\ &= \{(\xi, \xi + \eta^{\mathcal{V}}), \xi \in T_lL, \eta^{\mathcal{V}} \in \mathcal{V}_n\}. \end{aligned}$$

This gives us a splitting $T(u^*N) = \mathcal{V}^{u^*N} \oplus \mathcal{H}^{u^*N}$ where $\forall (l, n) \in u^*N$,

$$\mathcal{V}_{(l,n)}^{u^*N} = \{0\} \times \mathcal{V}_n \quad \text{and} \quad \mathcal{H}_{(l,n)}^{u^*N} = T_lL \times \mathcal{H}_n \cap T_{(l,n)}(u^*N) = \{(\xi, \xi) \in T_lL \times T_lL\}.$$

Let us determine the orthogonal of the tangent space $T(u^*N)$:

$$\begin{aligned} (\alpha, \beta) \in (T_{(l,n)}(u^*N))^\perp &\iff \\ \forall (\xi, \eta) \in T_{(l,n)}(u^*N), \quad 0 &= \langle (\xi, \eta), (\alpha, \beta) \rangle \\ &= \langle \xi, \alpha \rangle + \langle \eta, \beta \rangle \\ &= \langle \xi, \alpha \rangle + \langle \xi, \beta_{TL}^{\mathcal{H}} \rangle + \langle 0, \beta_{TL^\perp}^{\mathcal{H}} \rangle + \langle \eta^{\mathcal{V}}, \beta^{\mathcal{V}} \rangle \\ &= \langle \xi, \alpha + \beta_{TL}^{\mathcal{H}} \rangle + \langle \eta^{\mathcal{V}}, \beta^{\mathcal{V}} \rangle \\ &\iff (\alpha + \beta_{TL}^{\mathcal{H}}, \beta^{\mathcal{V}}) = 0. \end{aligned}$$

Therefore

$$(T_{(l,n)}(u^*N))^\perp = \{(-\beta_{TL}^{\mathcal{H}}, \beta), \beta \in \mathcal{H}_n\}.$$

Decomposing each $(a, b) \in T(L \times N)|_{u^*N}$ following the decomposition $T(L \times N)|_{u^*N} = T(u^*N) \oplus T(u^*N)^\perp$: $(a, b) = (\xi, \eta) + (\alpha, \beta)$, then we obtain

$$\begin{cases} a = \eta_{TL}^{\mathcal{H}} - \beta_{TL}^{\mathcal{H}} \\ b = (\eta_{TL}^{\mathcal{H}} + \beta_{TL}^{\mathcal{H}}) + \beta_{TL}^{\mathcal{H}} + \eta^{\mathcal{V}} \end{cases}$$

so that this decomposition is therefore given by

$$(a, b) = \left(\frac{a + b_{TL}^{\mathcal{H}}}{2}, a + b_{TL}^{\mathcal{H}} + b^{\mathcal{V}} \right) + \left(-\frac{(b_{TL}^{\mathcal{H}} - a)}{2}, \frac{(b_{TL}^{\mathcal{H}} - a)}{2} + b_{TL^{\perp}}^{\mathcal{H}} \right).$$

Now, let us come back to our function $f: L \rightarrow N$ and the corresponding section $\tilde{f}: (L, b) \rightarrow u^*N$. Then let us compute

$$\begin{aligned} \nabla^v d^v \tilde{f} &= \nabla^v (dl, df)^{\mathcal{V}^{u^*N}} = \nabla^v (0, d^v f) = \left([\nabla(0, d^v f)]_{T(u^*N)} \right)^{\mathcal{V}^{u^*N}} \\ &= \left([(0, \nabla d^v f)]_{T(u^*N)} \right)^{\mathcal{V}^{u^*N}} \\ &= \left(\frac{1}{2}(\nabla d^v f)_{TL}^{\mathcal{H}}, \frac{1}{2}(\nabla d^v f)_{TL}^{\mathcal{H}} + \nabla^v d^v f \right)^{\mathcal{V}^{u^*N}} \\ &= (0, \nabla^v d^v f) \end{aligned}$$

Finally, we have proved

$$\nabla^v d^v \tilde{f} = \nabla^v d^v f \quad (145)$$

and by taking the trace, we obtain the first assertion of the theorem.

Now, in the same way we obtain

$$\nabla d\tilde{f} = \left(\frac{1}{2}(\nabla df)_{TL}^{\mathcal{H}}, \frac{1}{2}(\nabla df)_{TL}^{\mathcal{H}} + \nabla^v df \right) \quad (146)$$

so that $\tilde{f}: N \rightarrow u^*N$ is harmonic if and only if $[\tau(f)]_{TL}^{\mathcal{H}} = 0$ and $[\tau(f)]^{\mathcal{V}} = 0$. Therefore $f: L \rightarrow N$ is harmonic if and only if $f: N \rightarrow u^*N$ is harmonic and $[\tau(f)]_{TL^{\perp}}^{\mathcal{H}} = 0$. This completes the proof. \square

From the proof of theorem 8.1 (more precisely from (145) and (146)), we obtain:

Theorem 8.2 *Let us consider the situation described by theorem 8.1. Then $f: L \rightarrow N$ is superflat if and only if the corresponding section $\tilde{f}: L \rightarrow u^*N$ is superflat. Furthermore $f: L \rightarrow N$ is totally geodesic if and only if $\tilde{f}: L \rightarrow u^*N$ is totally geodesic and $[\nabla df]_{TL^{\perp}}^{\mathcal{H}} = 0$ (i.e. $[d\pi(\nabla df)]_{TL^{\perp}} = 0$).*

Remark 8.1 The metric defined in example 4.1 in u^*N (and thus in theorems 8.1 and 8.2, i.e. the metric induced by the product metric, is given by

$$|(\xi, \eta)|^2 = 2|\xi|^2 + |\eta^{\mathcal{V}}|^2 \quad (147)$$

whereas, when $\pi: N \rightarrow M$ is a Homogeneous fibre bundle, the metric in u^*N , considered as an Homogeneous fibre bundle, is defined in 4.2 by equation (62) and is given by

$$|(\xi, \eta)|^2 = |\xi|^2 + |\eta^{\mathcal{V}}|^2. \quad (148)$$

However, theorems 8.1 and 8.2 hold, of course, also with the metric (148). Indeed, first remark that the theorems hold if we multiply the product metric in $L \times N$ by a constant factor. Then just apply these theorems with the same (M, g) (and thus the same (L, u^*g)), N endowed with the new metric $|\cdot|_{\mathcal{H}}^2 + 2|\cdot|_{\mathcal{V}}^2$ (the old one being $|\cdot|_{\mathcal{H}}^2 + |\cdot|_{\mathcal{V}}^2$) and endow $L \times N$ with $\frac{1}{2}$ times the product metric, then the induced metric on u^*N is (148): $\frac{1}{2}(|\xi|^2 + (|\xi|^2 + 2|\eta^{\mathcal{V}}|^2)) = |\xi|^2 + |\eta^{\mathcal{V}}|^2$.

8.2 G -invariant metrics

Here, we use the notations of section 2.1.

8.2.1 Existence of an $\text{Ad}H$ -invariant inner product on \mathfrak{m} for which $\tau|_{\mathfrak{m}}$ is an isometry.

Lemma 8.1 *Let G be a connected Lie group and $\sigma: G \rightarrow G$ an automorphism of order k . Then G^σ has finite many connected components.*

Proof. This is proved in [4, Lemma 2.7], for $k = 2$, i.e. σ is an involution. The proof in [4] can be generalized without any difficulty to the case of an order k automorphism. \square

Theorem 8.3 *Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of order $2k$, of a real Lie algebra \mathfrak{g} . Let G be a connected Lie group with Lie algebra \mathfrak{g} and H a subgroup such that $(G^\sigma)^0 \subset H \subset G^\sigma$, where $\sigma = \tau^2$. If $\text{Ad}_m H$ is compact, then the subgroup generated by $\text{Ad}_m H$ and $\tau|_{\mathfrak{m}}$ is compact.*

Proof. Follow the proof of [34, theorem 21] by using the lemma 8.1 above instead of [4, Lemma 2.7]. \square

8.2.2 Existence of a naturally reductive metric for which \underline{J} is an isometry, resp. F is metric.

The odd case Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of order $2k + 1$. We have

$$\tau_{\mathfrak{m}}[\text{ad}_{\mathfrak{m}}(X)]_{\mathfrak{m}}\tau_{\mathfrak{m}}^{-1} = [\text{ad}_{\mathfrak{m}}(\tau_{\mathfrak{m}}X)]_{\mathfrak{m}}, \quad \forall X \in \mathfrak{m}.$$

Therefore

Lemma 8.2 *Let $N = G/K$ be a (locally) $(2k + 1)$ -symmetric space endowed with its canonical almost complex structure \underline{J} . Suppose also that $N = G/K$ is naturally reductive. Denoting by $G(\mathfrak{m})$ the compact subgroup in $GL(\mathfrak{m})$ generated by $\Lambda_{\mathfrak{m}}(\mathfrak{m}) := \{[\text{ad}_{\mathfrak{m}}(X)]_{\mathfrak{m}}, X \in \mathfrak{m}\} \subset \mathfrak{gl}(\mathfrak{m})$, then $\langle G(\mathfrak{m}), \tau_{\mathfrak{m}} \rangle$, the subgroup generated by $G(\mathfrak{m})$ and $\tau_{\mathfrak{m}}$, is compact. Therefore for any metric h on \mathfrak{m} invariant by $\langle G(\mathfrak{m}), \tau_{\mathfrak{m}} \rangle$, then h is naturally reductive, and \underline{J}_0 is orthogonal.*

The even case Let $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ be an automorphism of order $2k$. We have

$$\tau_{\mathfrak{n}}[\text{ad}_{\mathfrak{m}}(X)]_{\mathfrak{n}}\tau_{\mathfrak{n}}^{-1} = [\text{ad}_{\mathfrak{n}}(\tau_{\mathfrak{n}}X)]_{\mathfrak{n}}, \quad \forall X \in \mathfrak{n}.$$

Therefore

Lemma 8.3 *Let $N = G/K$ be a (locally) $2k$ -symmetric space endowed with its canonical f -structure F . Suppose also that $N = G/K$ is naturally reductive. Denoting by $G(\mathfrak{n})$ the compact subgroup in $GL(\mathfrak{n})$ generated by $\Lambda_{\mathfrak{m}}(\mathfrak{n}) := \{[\text{ad}_{\mathfrak{n}}(X)]_{\mathfrak{n}}, X \in \mathfrak{n}\} \subset \mathfrak{gl}(\mathfrak{n})$, then $\langle G(\mathfrak{n}), \tau_{\mathfrak{n}} \rangle$, the subgroup generated by $G(\mathfrak{n})$ and $\tau_{\mathfrak{n}}$, is compact. Therefore for any metric h on \mathfrak{m} invariant by $\langle G(\mathfrak{n}), \tau_{\mathfrak{n}} \rangle$, then h is naturally reductive, and $I_0 = \underline{J}_0 \oplus -\text{Id}_{\mathfrak{g}_k}$ is orthogonal, i.e. F is metric.*

References

- [1] I. Agricola, *Connections on naturally reductive spaces, their Dirac operator and homogeneous models in string theory*, preprint, 2002.
- [2] I. Agricola, *The SRNI lectures on non-integrable geometries with torsion*, arXiv:math/0606705v1.
- [3] B. Alexandrov, Th. Friedrich, Nils Schoemann, *almost Hermitian 6-manifolds revisited*, J. Geom. Phys. 53 (2005), 1-30.
- [4] J. An, Z. Wang, *On the realization of Riemannian symmetric spaces in Lie groups II*, Topology and its Applications 153 (2006) 2943-2947.
- [5] A.L. Besse, *Einstein Manifolds*, Springer-Verlag, Berlin, Heidelberg, New York, 1987.
- [6] C. Bohle, *Constrained Willmore tori in the 4-sphere*, arxiv:0803.0633.
- [7] Bolton, J., Pedit, F., and Woodward, L., *Minimal surfaces and the affine field model*, J. reine angew. Math. 459 (1995), 119-150.
- [8] F.E. Burstall, *Harmonic tori in spheres and complex projective spaces*, J. reine angew. Math. 469 (1995), 149-177.
- [9] Burstall, F.E., Ferus, D., Pedit, F., Pinkall, U., *Harmonic tori in symmetric spaces and commuting Hamiltonian systems on loop algebras*, Ann. Math. 138, 173-212 (1993).
- [10] F.E. Burstall, I. Khemar, *Twistors, 4-symmetric spaces and integrable systems*, Mathematische Annalen (2009) 344: 451-461, (arXiv:0804.4235).
- [11] F.E. Burstall, F. Pedit, *Harmonic maps via Adler-Kostant-Symes Theory*, Harmonic maps and integrable systems, A.P. Fordy, J.C. Wood (Eds.), Vieweg (1994), 221-272.
- [12] F. Burstall, F. Pedit, U. Pinkall, *Schwarzian derivatives and flow of surfaces*, arxiv:math/011169.
- [13] F.E. Burstall and J.H. Rawnsley, *Twistor theory for Riemannian Symmetric Spaces with applications to harmonic maps of Riemann Surfaces* Lect. Notes in Math., vol. 1424, Springer, Berlin, Heidelberg, New York, 1990.
- [14] J. Dieudonné, *Éléments d'analyse, Tome 2*, Gauthiers-Villars.
- [15] J. Dorfmeister, Jun-Ichi Inoguchi, Shimpei Kobayashi, *Constant mean curvature surfaces in hyperbolic 3-space via loop groups*, preprint 2009.
- [16] J. Dorfmeister, F. Pedit and H.-Y. Wu, *Weierstrass type representation of harmonic maps into symmetric spaces*, Comm. in Analysis and Geometry, 6(4) (1998), p. 633-668.
- [17] T. Friedrich, S. Ivanov, *Parallel spinors and connections with skew-symmetric torsion in string theory*, Asian J. Math., Vol. 6, No. 2, pp. 303, June 2002.
- [18] *Hermitian connections and Dirac operators*, Boll. U.MI. (7) 11-B (1997), Suppl. fasc. 2, 257-288.
- [19] A. Gray and L. M. Hervella, *The sixteen classes of almost Hermitian manifolds and their linear invariants*, Ann. Mat. Pura e Appl. 123 (1980), 35-58.

- [20] R. Harvey, *Spinors and Calibrations*, Academic Press Inc., 1990.
- [21] R. Harvey and H. B. Lawson, *Calibrated geometries*, Acta Mathematica, 148 (1982), p. 47-157.
- [22] F. Hélein, *Applications harmoniques, lois de conservations et repères mobiles*, Diderot éditeur, Paris 1996; or *Harmonic maps, conservation laws and moving frames*, Cambridge University Press 2002.
- [23] F. Hélein, *Constant mean Curvature Surfaces, Harmonic maps and Integrable Systems*, Lecture in Mathematics, ETH Zürich, Birkhäuser 2001.
- [24] F. Hélein, *Willmore immersions and loop groups*, J. Diff. Geometry, 50(2) (1998), p.331-338.
- [25] F. Hélein and P. Romon, *Hamiltonian stationary Lagrangian surfaces in \mathbb{C}^2* , Comm. in Analysis and Geometry Vol. 10, N. 1, 2002, p. 79-126.
- [26] F. Hélein and P. Romon, *Weierstrass representation of lagrangian surfaces in four dimensional spaces using spinors and quaternions*, Comment. Math. Helv., 75 (2000), p. 668-680.
- [27] F. Hélein and P. Romon, *Hamiltonian stationnary Lagrangian surfaces in Hermitian symmetric spaces*, in *Differential Geometry and Integrable Systems*, Martin Guest, Reiko Miyaoka, and Yoshihiro Ohnita, Editors-AMS, 2002.
- [28] S. Helgason, *Differential geometry, Lie group and symmetric spaces*, Academic Press, Inc., 1978.
- [29] M. Higaki, *Actions of loop groups on the space of harmonic maps into reductive homogeneous spaces*, J. Math. Sci. Univ. Tokyo 5 (1998), 401-421.
- [30] J.A. Jimenez, *Riemannian 4-symmetric spaces*, Transactions of the American Mathematical Society, Vol.306, No.2. (Apr.,1988), pp. 715-734.
- [31] S. Ishihara, K. Yano, *On Integrability conditions of a structure satisfying $f^3 + f = 0$.*, Quart. J. math. Oxford (2), 15 (1964), 217-22.
- [32] I. Khemar, *Surfaces isotropes de \mathbb{O} et systèmes intégrables.*, Journal of Differential Geometry 79 (2008) 479-516.
- [33] I. Khemar, *Supersymmetric Harmonic Maps into Symmetric Spaces*, Journal of Geometry and Physics 57 (2007) 1601-1630.
- [34] I. Khemar, *Geometric interpretation of Second Elliptic Integrable Systems*, Differential Geometry and its Applications, 28 (2010) 40-64.
- [35] I. Khemar, *Some new examples of elliptic integrable systems*, in preparation.
- [36] I. Khemar, *Supersymmetric interpretation of even Elliptic Integrable Systems*, in preparation.
- [37] I. Khemar, *Elliptic integrable systems, WZW sigma models and string theory*, in preparation.
- [38] V.F. Kirichenko, *K-spaces of maximal rank*, Mat.Zam. 22 (1977), 465-476.

- [39] S. Kobayashi, *Differential geometry of complex vector bundles*, Publications of the mathematical society of Japan, Iwanami Shoten, Publishers and Princeton University Press, 1987.
- [40] S. Kobayashi, K. Nomizu, *Foundations of Differential Geometry*, Vols. 1,2, Wiley, New York, 1963, 1969.
- [41] Pinkall, U., Sterling, I. *On the classification of constant mean curvature tori*, Ann. of Math (2) 130 1989, no. 2 407-451.
- [42] O. Kowalski, *Generalized symmetric spaces*, Lect. Notes Math. 805. Springer 1980.
- [43] P. M. Quan, *Intoduction à la géométrie des variétés différentiables*, Monographies Universitaires de Mathématiques, Dunod Paris 1969.
- [44] A. Pressley and G. Segal, *Loop groups*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1986.
- [45] Áurea Casinhas Quintino, *Constrained Willmore Surfaces: Symmetries of a Moebius Invariant Integrable System*, arXiv:0912.5402.
- [46] J.H. Rawnsley, *f-structures, f-twistor spaces and harmonic maps*. In: Geometry seminar L.Bianci, Lect. Notes Math. 1164. Springer 1986.
- [47] C.L. Terng, *Geometries and Symmetries of Soliton Equations and Integrable Elliptic Equations*, preprint arXiv:math.DG/0212372.
- [48] K. Uhlenbeck, Harmonic maps into Lie group (classical solutions of the chiral model), J. Differential Geometry, **30** (1989), 1-50.
- [49] J.A. Wolf and A. Gray, *Homogeneous spaces defined by Lie group automorphisms*. I, II, J. Differential Geom. **2** (1968), 77-159.
- [50] C.M. Wood, *The Gauss Section of a Riemannian Immersion*, J. London Math. Soc. (2) 33 (1986) 157-168.
- [51] C.M. Wood, *Harmonic sections of homogeneous fibre bundles*, Differential Geometry and its applications 19 (2003) 193-210.