



HAL
open science

Fokker-Planck equation in bounded domain

Laurent Chupin

► **To cite this version:**

| Laurent Chupin. Fokker-Planck equation in bounded domain. 2009. hal-00375802

HAL Id: hal-00375802

<https://hal.science/hal-00375802>

Preprint submitted on 16 Apr 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FOKKER-PLANCK EQUATION IN BOUNDED DOMAIN

by Laurent CHUPIN

ABSTRACT. We study the existence and the uniqueness of a solution φ to the linear Fokker-Planck equation $-\Delta\varphi + \operatorname{div}(\varphi\mathbf{F}) = f$ in a bounded domain of \mathbb{R}^d when \mathbf{F} is a “confinement” vector field acting for instance like the inverse of the distance to the boundary. An illustration of the obtained results is given within the framework of fluid mechanics and polymer flows.

Equation de Fokker-Planck dans un domaine borné

RÉSUMÉ. On étudie l'existence et l'unicité de solution φ à l'équation de Fokker-Planck linéaire $-\Delta\varphi + \operatorname{div}(\varphi\mathbf{F}) = f$ sur un domaine borné de \mathbb{R}^d lorsque \mathbf{F} est un champ de vecteurs “confinant” par exemple comme l'inverse de la distance au bord. Une illustration des résultats obtenus est donnée dans le cadre de la mécanique des fluides et des écoulements de polymères.

1. Introduction

In this paper we are interested in the so called Fokker-Planck equation

$$(1.1) \quad -\Delta\varphi + \operatorname{div}(\varphi\mathbf{F}) = f.$$

- In the simplest case (that is $\mathbf{F} = \mathbf{0}$) this equation is known as the Laplace equation (when $f = 0$) or as the Poisson equation (when $f \neq 0$). The solutions of these equations are important in many fields of science, notably the fields of electromagnetism, astronomy and fluid dynamics, because they describe the behavior of electric, gravitational and fluid potentials.
- More generally, the main reason of the physical interest of equation (1.1) comes from the fact that it can be put in conservative form $\operatorname{div}(J) = f$ with $J = -\nabla\varphi + \varphi\mathbf{F}$. Thus it can be connected to a generalization to the Fick's law $J = -\nabla\varphi$ connecting diffusion flux J and concentration φ in inhomogeneous environments, see [9, 22].
- In the dynamical systems framework (see for instance [26]) the non-stationary Fokker-Planck equation $\partial_t\varphi = \varepsilon\Delta\varphi - \operatorname{div}(\varphi\mathbf{F})$ is usually introduced. In this case, the function φ represents the smooth probability density of a population driven by \mathbf{F} and subject to ε -small diffusion in the following sense. The term $\varphi\mathbf{F}$ is a vector field representing the population φ moving with the flow of \mathbf{F} , and so the divergence of this vector field represents a thinning out of the population due to \mathbf{F} , which therefore contributes negatively to the local growth rate of the population, $\partial_t\varphi$. This explains the drive term. Meanwhile the term $\varepsilon\Delta\varphi$ represents ε -small diffusion, and contributes positively to the growth rate. The study which is presented here concerns in particular the existence and the uniqueness of a steady-state solution. We note that the theory is closely related to applications, because the steady-state φ is an ε -smoothing of the measure on the attractors of the flow of \mathbf{F} (see [26]) and therefore in numerical and physical experiments φ can be used to model the data with ε -error.
- According to the contexts, the vector field \mathbf{F} can take various forms. In particular it may occur that physically realistic assumptions do not make it possible to conclude only with the already known results. We will give such a caricatural example in the last part.

Keywords: Fokker-Planck equation, Bounded domain, Stationary solution, Confinement, Fluid mechanics, Polymer flows.

Math. classification: 35J25, 35Q35, 35R60, 76A05, 82D60.

Besides the problems of the existence and the uniqueness, the question which interests us is to know which boundary conditions are needed to ensure the existence and the uniqueness of a solution of equation (1.1) in the bounded case.

We will see that this depends on \mathbf{F} . When \mathbf{F} is regular enough, i.e. does not diverge too quickly at the boundary, data on φ at the boundary of the domain enable to ensure the uniqueness of the solution. We remind of this result at the beginning (in particular because the proof resembles ours). We also say that when the domain does not have boundary, for instance if we are interested in the space \mathbb{R}^d or on a compact variety without boundary, uniqueness is ensured by imposing the average of φ .

We prove that, in the bounded case, when \mathbf{F} is not so regular, the "good" condition to ensure uniqueness is still to impose the average of φ , and that in that case, the unique solution vanishes on the boundary.

1.1. Some known results on equation (1.1)

Except for the case where $\mathbf{F} = \mathbf{0}$, a particularly simple case corresponds to $\mathbf{F} = \nabla V$ (V is assumed to be regular and differentiable) and $f = 0$. In this case $\varphi = \exp(V)$ is a solution of (1.1) in the bounded case as well as in the compact case or in the unbounded case. In the same way, we can easily prove that the case $\mathbf{F} = \nabla V + \mathbf{G}$ admits the solution $\varphi = \exp(V)$ if and only if $\operatorname{div}(\mathbf{G}) + \mathbf{G} \cdot \nabla V = 0$. Up to a renormalization, the average of such a solution will be equal to 1, as soon as $\exp(V) \in L^1(\Omega)$. The average value is consequently an essential ingredient to have uniqueness of the solution of equation (1.1) and we could be interested in the following problem

$$(1.2) \quad \begin{cases} -\Delta\varphi + \operatorname{div}(\varphi\mathbf{F}) = 0 & \text{in } \Omega, \\ \text{with } \int_{\Omega} \varphi = 1. \end{cases}$$

In the compact case without boundary E.C. Zeeman [26] proves the existence and uniqueness of a solution φ for an arbitrary smooth vector field \mathbf{F} (and without term source $f = 0$) on a compact manifold by the Perron-Fröbenius method:

THEOREM 1.1. — *Let Ω be a compact manifold without boundary. If $\mathbf{F} \in \mathcal{C}^\infty(\Omega)$ then there exists a unique non negative solution of (1.2).*

Without proof in the non-compact case (as writes E.C. Zeeman on p. 152, the extension of such results to non-compact case is an open question), E.C. Zeeman gives some example with $\Omega = \mathbb{R}^d$. These examples show that in the unbounded case uniqueness follows from some "boundary" conditions on \mathbf{F} , which are given by the behavior of \mathbf{F} outside large sphere. Many other works concern these equations of the Fokker-Planck type in \mathbb{R}^d . Most of these works describe specific assumptions for the potential \mathbf{F} at infinity. Let us quote as an example the beautiful recent series of works by Hérau, Nier and Helffer [13, 14] about the linear kinetic Fokker-Planck equation. See also the article [20] by Noarov in which the author gives some smallness conditions in some norm and rapid decay at infinity for \mathbf{F} to ensure the existence of a solution other than identical zero (in the case $f = 0$).

In this article, we are interested in a possible generalization in bounded domains. Usually, such a problem is coupled with boundary conditions (Dirichlet, Neumann or mixed boundary conditions). For instance, the natural weak formulation of the problem

$$\begin{cases} -\Delta\varphi + \operatorname{div}(\varphi\mathbf{F}) = f & \text{in } \Omega, \\ \text{with } \varphi = 0 & \text{on } \partial\Omega \end{cases}$$

is written

$$(1.3) \quad \begin{cases} \text{Find } \varphi \in H_0^1(\Omega) \text{ such that for all } \psi \in H_0^1(\Omega) \\ \int_{\Omega} \nabla \varphi \cdot \nabla \psi - \int_{\Omega} \varphi \mathbf{F} \cdot \nabla \psi = \langle f, \psi \rangle, \end{cases}$$

where $\langle f, \psi \rangle$ corresponds to the duality product $(H^{-1}(\Omega), H_0^1(\Omega))$. In order that all terms in (1.3) be defined, the minimal hypotheses on data are: $f \in H^{-1}(\Omega)$ and, thanks to the classical Sobolev injections, $\mathbf{F} \in L^{d_*}(\Omega)$ where $d_* = d$ for $d \geq 3$ and $d_* \in]2, +\infty[$ for $d = 2$. Within this framework, we have (see [7]):

THEOREM 1.2. — *Let Ω be a bounded domain of \mathbb{R}^d , $d \geq 2$. If $f \in H^{-1}(\Omega)$ and $\mathbf{F} \in L^{d_*}(\Omega)$ then there exists a unique solution of (1.3).*

The proof of the generalization which we propose is primarily based on the proof of this theorem. The main difficulty which appears for the study of problem (1.3) is the following: although the operator Δ is coercive, the operator $-\Delta + \operatorname{div}(\cdot \mathbf{F})$ is generally not coercive. The reason for which the result is still valid lies in the (conservative) form of the term $\operatorname{div}(\varphi \mathbf{F})$. Let us note that an equivalent theorem can be proved (see [7]) for equations of kind $-\Delta \varphi + \mathbf{G} \cdot \nabla \varphi = f$ but that it is not possible to obtain a similar general result for equations of the type $-\Delta \varphi + \operatorname{div}(\varphi \mathbf{F}) + \mathbf{G} \cdot \nabla \varphi = f$. In fact, the sum $\operatorname{div}(\varphi \mathbf{F}) + \mathbf{G} \cdot \nabla \varphi$ makes appear zeroth order terms and it is well known that the solutions of $-\Delta \varphi + \lambda \varphi = 0$, $\lambda \in \mathbb{R}$, with homogeneous Dirichlet boundary condition are not unique.

Moreover, J. Droniou proved that the same result is valid for other boundary conditions as non-homogeneous Dirichlet, Fourier or mixed boundary conditions (using more regularity for the domain Ω , say with Lipschitz continuous boundary). Concerning the Neumann boundary conditions, J. Droniou and J.-L. Vazquez recently showed that the same problem admits, for each fixed mean value, a unique solution with the said mean value (see [8]).

An other question is debated in the present paper: Which necessary and sufficient conditions must be placed on f , and what are the degrees of freedom on the solutions if \mathbf{F} is not regular enough ?

1.2. An (partial) answer

We show in this article that if the normal component of the vector $\mathbf{F}(\mathbf{x})$ behaves ⁽¹⁾ like $\frac{\alpha}{\operatorname{dist}(\mathbf{x}, \partial\Omega)}$ in a neighborhood of the boundary $\partial\Omega$ with $\alpha > 1$ then there exists a unique solution φ to the Fokker-Planck equation (1.1) as soon as the average of φ is given. Moreover, we can show that this unique solution automatically vanishes at the boundary $\partial\Omega$. More precisely we prove

THEOREM 1.3. — *Let Ω be a bounded domain of \mathbb{R}^d , $d \geq 2$. Let $f \in H_M^{-1}$ (this space will be precised later) and $\mathbf{F} = \boldsymbol{\kappa} + \nabla V$ where $\boldsymbol{\kappa} \in L^\infty(\Omega)$ and $V \in C^\infty(\Omega)$ satisfies $V = -\infty$ on the boundary of Ω . Under assumptions (\mathcal{H}_1) , (\mathcal{H}_2) and (\mathcal{H}_3) (see more details page 16) then there exists a unique solution of the Fokker-Planck equation (1.1) such that $\int_{\Omega} \varphi = 1$.*

Obviously, the additive assumptions (\mathcal{H}_1) , (\mathcal{H}_2) and (\mathcal{H}_3) enable to take into account the examples where the normal part of $\mathbf{F}(\mathbf{x})$ behaves like $\frac{\alpha}{\operatorname{dist}(\mathbf{x}, \partial\Omega)}$ in a neighborhood of the boundary $\partial\Omega$ with $\alpha > 1$.

They are not satisfied when $\alpha \leq 1$. Thus, there remain many cases without answers: for instance, when the normal part of \mathbf{F} behaves like $\frac{\alpha}{\operatorname{dist}(\mathbf{x}, \partial\Omega)}$ with $\alpha \leq 1$, and when the normal part of \mathbf{F} behaves like $\frac{\alpha}{\operatorname{dist}(\mathbf{x}, \partial\Omega)^\beta}$ with $\alpha > 0$ and $\beta > 1$.

The assumptions (\mathcal{H}_1) , (\mathcal{H}_2) and (\mathcal{H}_3) on the potential V are rather difficult to apprehend. The

(1) We can verify that we have $\mathbf{F} \notin L^{d_*}(\Omega)$, and that consequently the announced result is a generalization of the Theorem 1.2.

reason for which we have these assumptions is the following: they are used in this form in each steps of the proof (primarily in the various lemmas). Thus if one of the steps of the proof can be shown in another way that presented here, we can hope to be freed from certain corresponding assumptions.

1.3. Outline of the paper

The paper is organized as follows:

- In Section 2, we give mains tools adapted to the studied problem. First of all, some tools about differential geometry to understand “explosive” boundary conditions. Next we give all the lemmas which are used in the main proof.
- In Section 3 the precise statement of the main result is enunciated, see Theorem 3.3, page 16.
- The Section 4 is devoted to the proof of Theorem 3.3. It is composed of two parts: the existence proof and the uniqueness proof.
- The last section (Section 5) gives an application to fluid mechanics and some numerical results.

2. Main implements

In [5], the author gives an existence and a uniqueness result for a Fokker-Planck equation for a particular vector field \mathbf{F} and in a particular domain Ω which is a ball. For more complex domains, we must understand the effect of the geometry in the proof. We will present in this part some elements of differential geometry adapted for our calculus. Next, we will precise the functional framework adapted to the Fokker-Planck equation of this paper. Finally, we will give multiple fundamental lemmas which are use for the proof of the Theorem 3.3.

2.1. Elementary differential geometry

The results of this part are largely inspired on Subsection 2.1 of the paper [2] and on the annexe C of the book [3].

Let Ω be a smooth (say \mathcal{C}^2) bounded domain in \mathbb{R}^d , $d \geq 2$. We denote by Γ its boundary and by ν the outward unitary normal to Γ . The distance between any $\mathbf{x} \in \mathbb{R}^d$ and the boundary Γ is denoted by $\delta_\Gamma(\mathbf{x})$.

For any $\varepsilon \geq 0$, we introduce the open subset of Ω :

$$\mathcal{O}_\varepsilon = \{\mathbf{x} \in \Omega; \delta_\Gamma(\mathbf{x}) < \varepsilon\} \quad \text{and} \quad \Omega_\varepsilon = \{\mathbf{x} \in \Omega; \delta_\Gamma(\mathbf{x}) > \varepsilon\}.$$


It is classical that, for ε small enough, the two maps δ_Γ (called distance to Γ) and P_Γ (called projection on Γ) exist and are regular on $\overline{\mathcal{O}_\varepsilon}$. This allow to use tangential and normal variable near Γ , defining for any function $f : \overline{\mathcal{O}_\varepsilon} \rightarrow \mathbb{R}$ the corresponding function $\tilde{f} : [0, \varepsilon] \times \Gamma \rightarrow \mathbb{R}$ by⁽²⁾

$$\forall (r, \boldsymbol{\theta}) \in [0, \varepsilon] \times \Gamma, \quad \tilde{f}(r, \boldsymbol{\theta}) = f(\boldsymbol{\theta} - r \boldsymbol{\nu}(\boldsymbol{\theta})).$$

Moreover, for any $f \in L^1(\mathcal{O}_\varepsilon)$ we have the following change of variables formula:

$$(2.1) \quad \int_{\mathcal{O}_\varepsilon} f(\mathbf{x}) \, d\mathbf{x} = \int_\Gamma \int_0^\varepsilon \tilde{f}(r, \boldsymbol{\theta}) |J(r, \boldsymbol{\theta})| \, dr \, d\boldsymbol{\theta},$$

(2) For non-ambiguous cases, \tilde{f} and f will be together denote f .



where dr and $d\theta$ corresponds to the Lebesgue measure on $[0, \varepsilon]$ and Γ respectively, and where J is the jacobian determinant of the previous change of variables. Introduce $\kappa_M = \sup_{\theta \in \Gamma} \sup_i \kappa_i(\theta)$, where $\kappa_1(\theta), \dots, \kappa_{d-1}(\theta)$ are the principal curvatures of Γ at θ , we have for $0 \leq r \leq \varepsilon$ and for $\theta \in \Gamma$

$$(1 - \varepsilon \kappa_M)^{d-1} \leq J(r, \theta) \leq 1.$$

Using the change of variables (2.1) we deduce that for $f \in L^1(\mathcal{O}_\varepsilon)$ we have

$$(2.2) \quad (1 - \varepsilon \kappa_M)^{d-1} \int_{\Gamma} \int_0^\varepsilon \tilde{f}(r, \theta) dr d\theta \leq \int_{\mathcal{O}_\varepsilon} f(\mathbf{x}) d\mathbf{x} \leq \int_{\Gamma} \int_0^\varepsilon \tilde{f}(r, \theta) dr d\theta.$$

Roughly speaking these relations show us that for thin tubular neighborhood of Γ in Ω the jacobian $J(r, \theta)$ is equivalent to 1 for small r , uniformly with respect to $\theta \in \Gamma$. Consequently, the relations (2.2) give an approximation of $\int_{\mathcal{O}_\varepsilon} f$ by $\int_{\Gamma} \int_0^\varepsilon \tilde{f}$.

Notice too that it is possible to define δ_Γ as a regular function on $\bar{\Omega}$. This extension will be also noted δ_Γ .

2.2. Functional spaces

The Fokker-Planck equation (1.1) make appear a potential V , assume to be smooth on Ω , via the force $\mathbf{F} = \boldsymbol{\kappa} + \nabla V$. This potential is supposed to be confinant so that we assume that $e^V \in L^1(\Omega)$. We can always define a maxwellian function M by

$$M = \frac{e^V}{\int_{\Omega} e^V}.$$

Notice that $M \in C^\infty(\Omega : \mathbb{R}_+^*)$ with $\int_{\Omega} M = 1$. Notice too that the interesting cases for the present study correspond to a maxwellian M vanishing on the boundary $\Gamma = \partial\Omega$ (that is $V = -\infty$ on Γ). All the maxwellians considered in this paper will satisfy $M = 0$ on Γ . The Fokker-Planck equation (1.1) is written

$$(2.3) \quad \operatorname{div}(\varphi \boldsymbol{\kappa} - M \nabla \left(\frac{\varphi}{M} \right)) = f.$$

From the peculiar form of this Fokker-Planck equation (2.3), the adapted functional spaces use Sobolev weight spaces on the domain $\Omega \subset \mathbb{R}^d$. More precisely, we introduce

$$L_M^2 := ML^2(Mdx) \quad \text{and} \quad H_M^1 := MH^1(Mdx),$$

endowed with their natural norms respectively given by

$$\begin{aligned} \|\varphi\|_{L_M^2}^2 &= \int_{\Omega} M \left| \frac{\varphi}{M} \right|^2, \\ \|\varphi\|_{H_M^1}^2 &= \int_{\Omega} M \left| \frac{\varphi}{M} \right|^2 + M \left| \nabla \left(\frac{\varphi}{M} \right) \right|^2. \end{aligned}$$

Note that a large literature on weighted Sobolev spaces exists, see for instance [23, 25], the references given in the notes at the end of Chapter 1 of [12] or the classical book [24]. Among all the weights which are generally considered in the literature, only some are vanished on \mathbb{R}^d as it is the case of the weight M . We refer to [24] for references of spaces with weights which equal to a power of

the distance to the boundary. Nevertheless, the spaces $L^2(Mdx)$ and $H^1(Mdx)$ being spaces with "traditional" weights, the spaces L^2_M and H^1_M are not it. For this reasons, the results shown in the following section are not always "traditional" corollaries (except some whose proof is a direct consequence of the results in $L^2(Mdx)$ and $H^1(Mdx)$, see for example the proof of lemma 2.4). Note first that the two spaces L^2_M and H^1_M are Hilbert spaces (see [24, Th. 3.2.2a]). We introduce the following normalized subspace

$$H^1_{M,0} := \{\psi \in H^1_M ; \int_{\Omega} \psi = 0\}.$$

In the sequel, the space H^1_M will be equipped too with the semi-norm defined by

$$\|\psi\|_{H^1_{M,0}}^2 = \int_{\Omega} M \left| \nabla \left(\frac{\psi}{M} \right) \right|^2,$$

and we will see (Lemma 2.9 on page 13) that $\|\cdot\|_{H^1_{M,0}}$ is a norm on the space $H^1_{M,0}$. Finally, we denote H_M^{-1} the topological dual of $H^1_{M,0}$, that is the set of continuous linear forms on $H^1_{M,0}$. Each application $\chi \in H_M^{-1}$ will be defined by $\chi : \varphi \in H^1_{M,0} \mapsto \langle \chi, \varphi \rangle \in \mathbb{R}$. By its continuity, for each $\chi \in H_M^{-1}$ there exists $C \in \mathbb{R}$ such that

$$\forall \varphi \in H^1_{M,0} \quad |\langle \chi, \varphi \rangle| \leq C \|\varphi\|_{H^1_{M,0}}.$$

As it is usual, the smallest of these constants C is denoted $\|\chi\|_{H_M^{-1}}$: it is the norm of χ on H_M^{-1} .

2.3. Properties of the functional spaces

The proof of the main theorem (Theorem 3.3) follows the ideas of J. Droniou [7]. Nevertheless, the proof of J. Droniou, given in the case where $\mathbf{F} \in L^{d^*}(\Omega)$, does not use these "degenerated" spaces and use traditional results concerning usual Sobolev spaces. The essential contributions which are presented ties in the fact that these "classical" lemmas in the case where M does not vanished on $\overline{\Omega}$ are still true (sometimes in a weaker form) when M is a maxwellian as previously introduced, and in particular when $M = 0$ on Γ . So, the goal of this subsection is to give some essential properties of these functional spaces (Poincaré-type inequality, Sobolev injection, compactity result, Hardy-type inequality...).

Notice that in the estimates, the symbol \lesssim means "up to a harmless multiplicative constant", allowed to depend on the domain Ω only.

The first result that we present is a result allowing to controlled $\frac{1}{\delta_{\Gamma}} \frac{\varphi}{\sqrt{M}}$ in $L^2(\Omega)$ as soon as $\varphi \in H^1_M$. This result can be seen as an inequality of Hardy-type ⁽³⁾. We will prove this first lemma under the following assumption ⁽⁴⁾

$$(\mathcal{H}_1) \quad \begin{cases} \exists a < 1 \quad \left(\frac{\nabla_R M}{M} \right)^2 + 2 \nabla_R \left(\frac{\nabla_R M}{M} \right) \geq \frac{-a}{\delta_{\Gamma}^2}, \\ \nabla_R M(0) = 0, \\ \exists b > 0 \quad \nabla_R M \int_{\Omega} \frac{1}{M} < b. \end{cases}$$

⁽³⁾ Hardy inequality indicates that for a function f defined on a bounded domain Ω of \mathbb{R}^d and vanishing on the boundary Γ of Ω we get $|f/\delta_{\Gamma}|_{L^2(\Omega)} \lesssim |\nabla f|_{L^2(\Omega)}$ where δ_{Γ} corresponds to the distance to the boundary Γ .

⁽⁴⁾ Recall that, as it is specified just before, in this article the function M is a Maxwellian function which vanished on the boundary of the domain. The assumptions introduced here are consequently additional assumptions.

These hypotheses will be only used in the neighbourhood of the boundary Γ of Ω . In such neighbourhood, the notation ∇_R corresponds to the radial derivative (that is to say the derivative in the direction of the normal vector to the boundary).

The various assumptions introduced in this part will be discussed in Section 3. Nevertheless, it is important to notice that the three assumptions formulated in (\mathcal{H}_1) are independent. For example, in the radial case the function M defined by $M(r) = \sqrt{r}$ satisfy the last point of (\mathcal{H}_1) but does not satisfy the second point. Reciprocally, $M(r) = r/\ln(r)$ satisfy the second point but does not satisfy the last point.

LEMMA 2.1 (Hardy-type inequality). — *If M satisfies (\mathcal{H}_1) then for any $\varphi \in H_M^1$ we have*

$$\int_{\Omega} \frac{1}{\delta_{\Gamma}^2} M \left| \frac{\varphi}{M} \right|^2 \lesssim \|\varphi\|_{H_M^1}^2.$$

In fact, as can be seen from the proof, we shall not need entire norm of φ in H_M^1 since we will only use the radial part of the gradient.

Proof. — For any ε small enough (more precisely, $\varepsilon \leq \varepsilon_{\Omega}$, see Part 2.1) we have $\Omega = \overline{\Omega_{\varepsilon}} \cup \mathcal{O}_{\varepsilon}$ and

$$\int_{\Omega} \frac{M}{\delta_{\Gamma}^2} \left| \frac{\varphi}{M} \right|^2 = \int_{\Omega_{\varepsilon}} \frac{M}{\delta_{\Gamma}^2} \left| \frac{\varphi}{M} \right|^2 + \int_{\mathcal{O}_{\varepsilon}} \frac{M}{\delta_{\Gamma}^2} \left| \frac{\varphi}{M} \right|^2.$$

Since $1/\delta_{\Gamma}$ is bounded in Ω_{ε} we easily deduce that

$$\int_{\Omega_{\varepsilon}} \frac{M}{\delta_{\Gamma}^2} \left| \frac{\varphi}{M} \right|^2 \lesssim \int_{\Omega_{\varepsilon}} M \left| \frac{\varphi}{M} \right|^2 \leq \|\varphi\|_{L_M^2}^2$$


and the main difficulty to prove the Lemma 2.1 is concentrated in the control of the integral

$$I_0 := \int_{\mathcal{O}_{\varepsilon}} \frac{M}{\delta_{\Gamma}^2} \left| \frac{\varphi}{M} \right|^2.$$

For similar reasons, we can suppose that

$$(2.4) \quad \varphi|_{\overline{\Omega_{\varepsilon}} \cap \overline{\mathcal{O}_{\varepsilon}}} = 0.$$

In fact, introduce a regular function γ_{ε} (only depending on Ω and ε) which vanished outside the ε -neighborhood $\mathcal{O}_{\varepsilon}$ of Γ and is equal to 1 in the $\frac{\varepsilon}{2}$ -neighborhood. The estimate in Lemma 2.1 clearly holds for $(1 - \gamma_{\varepsilon})\varphi$ and the proof therefore has to be conducted only for $\gamma_{\varepsilon}\varphi$, which for sake of simplicity will be denote φ in the following.



The value of ε being able to be chosen as small as desired, we shall use copiously the change of variables (2.1) as well as approximation (2.2).

For each $\theta \in \Gamma$ let introduce the function

$$h_{\theta} : r \in]0, \varepsilon] \mapsto \frac{\varphi(r, \theta)}{\sqrt{M(r, \theta)}}.$$

Note that the relation (2.4) implies that $h_{\theta}(\varepsilon) = 0$.

To simplify the notations, when there are no ambiguities, we do not note the dependence with

respect to the variable $\boldsymbol{\theta}$ in $h_{\boldsymbol{\theta}}$ and in M . The integral I_0 writes using the approximation (2.2)

$$I_0 = \int_{\Gamma} \int_0^{\varepsilon} \frac{h(r)^2}{r^2} dr d\boldsymbol{\theta}$$

The goal is to control I_0 with $\|\varphi\|_{H_M^1}$. We will proceed in two steps:

- (1) We prove that $h(r)M'(r)/M(r) = 0$ on $r = 0$;
- (2) We control I_0 with $\|\varphi\|_{H_M^1}$.

• Step (1): We use the following change of variable adapted to the maxwellian M :

$$\Phi : (r, \boldsymbol{\theta}) \in]0, \varepsilon] \times \Gamma \longmapsto (s, \boldsymbol{\theta}) \in [0, +\infty[\times \Gamma,$$

$$\text{where } s = \int_r^{\varepsilon} \frac{dt}{\widetilde{M}(t, \boldsymbol{\theta})}.$$

Notice that the jacobian determinant of $\Phi(r, \boldsymbol{\theta})$ equals $-1/\widetilde{M}(r, \boldsymbol{\theta})$ and therefore, M being positive on Ω , it does not cancel on $]0, \varepsilon] \times \Gamma$. Moreover, for all $\boldsymbol{\theta} \in \Gamma$ we have $\lim_{r \rightarrow 0} \Phi(r, \boldsymbol{\theta}) = (+\infty, \boldsymbol{\theta})$ since using the assumption on M we get $\int_0^{\varepsilon} dr/\widetilde{M}(r, \boldsymbol{\theta}) = +\infty$. Consequently Φ is a local diffeomorphism from $]0, \varepsilon] \times \Gamma$ to $\Phi(]0, \varepsilon] \times \Gamma) = [0, +\infty[\times \Gamma$.

For any function $\widetilde{f} :]0, \varepsilon[\times \Gamma \mapsto \mathbb{R}$ we will define the function, noted $\widehat{f} :]0, +\infty[\times \Gamma$, by $\widetilde{f} = \widehat{f} \circ \Phi$. Using the change of variables introduce in Subsection 2.1, for any function $f : \mathcal{O}_{\varepsilon} \mapsto \mathbb{R}$ we have define the functions $\widetilde{f} :]0, \varepsilon[\times \Gamma \mapsto \mathbb{R}$ and $\widehat{f} :]0, +\infty[\times \Gamma$ such that, with the previous notations for the name of variables:

$$f(\mathbf{x}) = \widetilde{f}(r, \boldsymbol{\theta}) = \widehat{f}(s, \boldsymbol{\theta}).$$

In the nonambiguous cases, we will note f , all the functions f , \widetilde{f} or \widehat{f} .

As announced before the proof, we only need the radial part of the gradient in the desired estimate. In term of new coordinates, the radial part of the gradient of a function f defined on Ω_{ε} corresponds to the derivative with respect to the variable r in the new coordinates: $\nabla_R f(\mathbf{x}) = \partial_r \widetilde{f}(r, \boldsymbol{\theta})$.

For any $\boldsymbol{\theta} \in \Gamma$ let $g_{\boldsymbol{\theta}}$ be the function defined on $]0, +\infty[$ by

$$g_{\boldsymbol{\theta}}(s) := \frac{\varphi(\mathbf{x})}{M(\mathbf{x})} = \frac{h_{\boldsymbol{\theta}}(r)}{\sqrt{M(r, \boldsymbol{\theta})}}.$$

Derivating with respect to the variable s , we obtain (as previously the variable $\boldsymbol{\theta}$ will be understood as a parameter and we do not note its dependence):

$$M(\mathbf{x}) \left| \nabla_R \left(\frac{\varphi(\mathbf{x})}{M(\mathbf{x})} \right) \right|^2 d\mathbf{x} = -|g'(s)|^2 |J(r, \boldsymbol{\theta})| d\boldsymbol{\theta} ds.$$

We deduce that (using the approximation $J \sim 1$ valid for ε small enough)

$$I_1 := \int_{\Omega} M(\mathbf{x}) \left| \nabla_R \left(\frac{\varphi(\mathbf{x})}{M(\mathbf{x})} \right) \right|^2 d\mathbf{x} = \int_{\Gamma} \left(\int_0^{+\infty} |g'(s)|^2 ds \right) d\boldsymbol{\theta}.$$

Since $\varphi \in H_M^1$ the integral I_1 is finite. Consequently, for almost every $\boldsymbol{\theta} \in \Gamma$ we have

$$\int_0^{+\infty} |g'(s)|^2 ds < +\infty.$$

We deduce that

$$\forall \alpha > 0 \quad \exists s_{\alpha} > 0 \quad \int_{s_{\alpha}}^{+\infty} |g'(s)|^2 ds \leq \alpha^2.$$

Next, using the Cauchy-Schwarz inequality we obtain, for any $s \in]0, +\infty[$,

$$g(s) = g(s_{\alpha}) + \int_{s_{\alpha}}^s g'(t) dt \leq \begin{cases} g(s_{\alpha}) + \alpha \sqrt{s - s_{\alpha}} & \text{if } s > s_{\alpha}, \\ \sup_{[0, s_{\alpha}]} g & \text{if } 0 < s \leq s_{\alpha}. \end{cases}$$

We deduce that for almost every $\boldsymbol{\theta} \in \Gamma$, for all $\alpha > 0$ there exists a constant $C_{\boldsymbol{\theta},\alpha} > 0$ such that, for any $s \in]0, +\infty[$ we have $g(s) \leq \alpha\sqrt{s} + C_{\boldsymbol{\theta},\alpha}$. Using the r variable, this result is written: for any $r \in]0, \varepsilon]$

$$\frac{M'(r)}{M(r)}h(r)^2 \leq \alpha^2 M'(r) \int_r^\varepsilon \frac{dt}{M(t)} + M'(r)C_{\boldsymbol{\theta},\alpha}^2,$$

that enables, see assumption (\mathcal{H}_1) , to obtain for almost every $\boldsymbol{\theta} \in \Gamma$ the relation

$$(2.5) \quad \left. \frac{M'(r)}{M(r)}h(r)^2 \right|_{r=0} = 0.$$

• Step (2): Now, we prove the lemma. Since $\varphi \in H_M^1$, we know that

$$I_1 := \int_{\Omega} M(\mathbf{x}) \left| \nabla_R \left(\frac{\varphi(\mathbf{x})}{M(\mathbf{x})} \right) \right|^2 d\mathbf{x} \leq \|\varphi\|_{H_M^1}^2 < +\infty.$$

We express I_1 making appear the h function and using the change of variables $\mathbf{x} \mapsto (r, \boldsymbol{\theta})$ together the usual approximation for the jacobian determinant of this change of variable (see the Part 2.1 and the relations (2.2)). We obtain

$$I_1 = \int_{\Gamma} \int_0^\varepsilon M(r, \boldsymbol{\theta}) \left| \frac{\partial}{\partial r} \left(\frac{\varphi(r, \boldsymbol{\theta})}{M(r, \boldsymbol{\theta})} \right) \right|^2 dr d\boldsymbol{\theta} := \int_{\Gamma} I_1(\boldsymbol{\theta}) d\boldsymbol{\theta},$$

where, for $\boldsymbol{\theta} \in \Gamma$, the quantity $I_1(\boldsymbol{\theta})$ is defined by

$$\begin{aligned} I_1(\boldsymbol{\theta}) &= \int_0^\varepsilon M(r) \left(\frac{1}{\sqrt{M(r)}} h'(r) - \frac{M'(r)}{2M(r)\sqrt{M(r)}} h(r) \right)^2 dr \\ &= \int_0^\varepsilon \left(h'(r)^2 - \frac{M'(r)}{M(r)} h'(r)h(r) + \frac{M'(r)^2}{4M(r)^2} h(r)^2 \right) dr. \end{aligned}$$

Moreover, an integration by part gives

$$- \int_0^\varepsilon \frac{M'(r)}{M(r)} h'(r)h(r) dr = \frac{1}{2} \int_0^\varepsilon \left(\frac{M'(r)}{M(r)} \right)' h(r)^2 dr - \frac{1}{2} \left[\frac{M'(r)}{M(r)} h(r)^2 \right]_0^\varepsilon.$$

We deduce that

$$\begin{aligned} I_1(\boldsymbol{\theta}) &= \int_0^\varepsilon \left(h'(r)^2 + \lambda(r) \frac{h(r)^2}{4r^2} \right) dr - \frac{1}{2} \left[\frac{M'(r)}{M(r)} h(r)^2 \right]_0^\varepsilon \\ &\text{with } \lambda(r) = \left(\left(\frac{M'(r)}{M(r)} \right)^2 + 2 \left(\frac{M'(r)}{M(r)} \right)' \right) r^2. \end{aligned}$$

The assumption (\mathcal{H}_1) on M is written $\lambda \geq -a > -1$. Moreover using the equation (2.5) and the relation (2.4) the bracket term is vanished. We obtain

$$I_1(\boldsymbol{\theta}) \geq \int_0^\varepsilon \left(h'(r)^2 - a \frac{h(r)^2}{4r^2} \right) dr.$$

Moreover, thanks to the Hardy inequality (holds since h vanishes at 0, this is a direct consequence of equation (2.5)), we deduce

$$(2.6) \quad I_1(\boldsymbol{\theta}) \geq (1-a) \int_0^\varepsilon h'(r)^2 dr.$$

Since $a < 1$, this control allows us to estimate $\int_0^\varepsilon h'(r)^2 dr$. From the Hardy inequality again, we obtain the following estimate

$$\int_0^\varepsilon \frac{h(r)^2}{r^2} dr \leq 4 \int_0^\varepsilon h'(r)^2 dr \leq \frac{4}{1-a} I_1(\boldsymbol{\theta}).$$

Integrate with respect to the variable $\boldsymbol{\theta}$, we obtain

$$I_0 = \int_{\Gamma} \int_0^\varepsilon \frac{h(r)^2}{r^2} dr d\boldsymbol{\theta} \leq \frac{4}{1-a} I_1.$$

The function φ being in H_M^1 , right hand side terms are bounded by $\|\varphi\|_{H_M^1}^2$. Up to the change of variables we deduce

$$\int_{\mathcal{O}_\varepsilon} \frac{M}{\delta_\Gamma^2} \left| \frac{\varphi}{M} \right|^2 \lesssim \|\varphi\|_{H_M^1}^2,$$

which implies the announced result. \square

The additional hypothesis

$$(\mathcal{H}_2) \quad \exists c > 0 \quad |\nabla M| \leq \frac{1}{c} \frac{M}{\delta_\Gamma}$$

implies from Lemma 2.1 that if $\varphi \in H_M^1$ then $\frac{\nabla M}{M} \frac{\varphi}{\sqrt{M}} \in L^2(\Omega)$. It is in this form that the Lemma 2.1 will be generally used in this article.

Note that in term of \mathbf{x} variable, the inequality (2.6) show us that $\nabla_R(\frac{\varphi}{\sqrt{M}})$ belongs to $L^2(\Omega)$. This propertie is completed by the next lemma:

LEMMA 2.2 (Inclusion). — *If M satisfies (\mathcal{H}_1) and (\mathcal{H}_2) then we have the following inclusions*

$$L_M^2 \subset L^2(\Omega) \quad \text{and} \quad H_M^1 \subset H_0^1(\Omega).$$

More precisely, if $\varphi \in L_M^2$ then $\varphi/\sqrt{M} \in L^2(\Omega)$ and if $\varphi \in H_M^1$ then $\varphi/\sqrt{M} \in H^1(\Omega)$.

Proof. — Inclusion $L_M^2 \subset L^2(\Omega)$ is obvious since on the one hand, by definition of L_M^2 , we have $\varphi \in L_M^2$ if and only if $\varphi/\sqrt{M} \in L^2(\Omega)$, and on the other hand $\sqrt{M} \in L^\infty(\Omega)$.

To prove the inclusion $H_M^1 \subset H_0^1(\Omega)$ we use the Lemma 2.1 with additional assumption (\mathcal{H}_2) : if $\varphi \in H_M^1$ then we have

$$\nabla\left(\frac{\varphi}{\sqrt{M}}\right) = \nabla\left(\frac{\varphi}{M}\sqrt{M}\right) = \underbrace{\sqrt{M}\nabla\left(\frac{\varphi}{M}\right)}_{L^2(\Omega)} + \underbrace{\frac{1}{2}\frac{\nabla M}{M}\frac{\varphi}{\sqrt{M}}}_{L^2(\Omega)} \in L^2(\Omega).$$

Consequently $\varphi/\sqrt{M} \in H^1(\Omega)$. Hence this function φ/\sqrt{M} has a trace on the boundary Γ . Since M is a regular function on Ω vanishing on Γ , we deduce that $\varphi = \frac{\varphi}{\sqrt{M}}\sqrt{M} \in H_0^1(\Omega)$. \square

This next lemma is interesting in themselves for understanding the space H_M^1 better. Moreover, it will be used in the proof of the Lemma 2.7.

LEMMA 2.3 (Density). — *If M satisfies (\mathcal{H}_1) and (\mathcal{H}_2) then we have the following equality*

$$\overline{C_0^\infty}^{H_M^1} = H_M^1.$$

Proof. — Let $\varphi \in H_M^1$ and define, for $n \in \mathbb{N}^*$, the function φ_n by

$$\varphi_n = \varphi \chi\left(\frac{1}{nM}\right) \quad \text{where} \quad \chi(t) = \begin{cases} 1 & \text{if } 0 \leq t < 1, \\ 2-t & \text{if } 1 \leq t < 2, \\ 0 & \text{if } t \geq 2. \end{cases}$$

We successively prove that

- (1) the functions φ_n are in H_M^1 ,
- (2) we can approach these functions φ_n with C_0^∞ functions,
- (3) the sequence $\{\varphi_n\}_{n \in \mathbb{N}^*}$ converges to φ in H_M^1 sense.

These three points clearly implicate the lemma.

- (1) By definition of χ , we have for all $n \in \mathbb{N}^*$ the relation $|\varphi_n| \leq |\varphi|$ hence $\int_\Omega M \left| \frac{\varphi_n}{M} \right|^2 \leq \int_\Omega M \left| \frac{\varphi}{M} \right|^2$

which is bounded since $\varphi \in H_M^1 \subset L_M^2$. To control the gradient part of the H_M^1 -norm of φ_n , $n \in \mathbb{N}^*$, we write

$$\begin{aligned} \left| \nabla \left(\frac{\varphi_n}{M} \right) \right| &= \left| \nabla \left(\frac{\varphi}{M} \right) \chi \left(\frac{1}{nM} \right) + \frac{\varphi}{M} \nabla \left(\chi \left(\frac{1}{nM} \right) \right) \right| \\ &\leq \left| \nabla \left(\frac{\varphi}{M} \right) \right| + \left| \frac{\varphi}{M} \nabla \left(\chi \left(\frac{1}{nM} \right) \right) \right|. \end{aligned}$$

By definition of the truncature function χ , the last term is not equal to 0 if and only if $\frac{1}{2n} \leq M < \frac{1}{n}$. In this case it write $\left| \frac{\varphi}{n} \frac{\nabla M}{M^2} \right|$ and can be controlled by $2 \left| \frac{\varphi}{M} \frac{\nabla M}{M} \right|$. We have

$$\left| \nabla \left(\frac{\varphi_n}{M} \right) \right|^2 \leq \left| \nabla \left(\frac{\varphi}{M} \right) \right|^2 + 4 \left| \frac{\varphi}{M} \frac{\nabla M}{M} \right|^2.$$

Since $\varphi \in H_M^1$ and using the Lemma 2.1 together with the assumption (\mathcal{H}_2) , we deduce that $\int_{\Omega} M \left| \nabla \left(\frac{\varphi_n}{M} \right) \right|^2$ is bounded. Consequently, $\varphi_n \in H_M^1$.

• (2) For each $n \in \mathbb{N}^*$ the function $\varphi_n \in H_M^1 \subset H^1$ is equal to 0 in a neighborhood of the boundary Γ . Approaching φ_n by a sequence $\{\varphi_{n,m}\}_{m \in \mathbb{N}}$ (which is equal to 0 on a neighborhood of Γ) in H_0^1 allows to approach φ_n by the same sequence $\{\varphi_{n,m}\}_{m \in \mathbb{N}}$ in H_M^1 .

• (3) Note that $\varphi(\mathbf{x}) - \varphi_n(\mathbf{x})$ equals to 0 if $M(\mathbf{x}) > \frac{1}{n}$. In the other case, that is for all $\mathbf{x} \in \Omega$ such that $M(\mathbf{x}) \leq \frac{1}{n}$, we get $|\varphi_n(\mathbf{x}) - \varphi(\mathbf{x})| \leq |\varphi(\mathbf{x})|$ since $0 \leq \chi \leq 1$. Hence

$$\int_{\Omega} M \left| \frac{\varphi_n - \varphi}{M} \right|^2 \leq \int_{\mathbf{x} \in \Omega; M(\mathbf{x}) \leq \frac{1}{n}} M \left| \frac{\varphi}{M} \right|^2.$$

Since $\varphi \in H_M^1 \subset L_M^2$ and since the measure of the set $\{\mathbf{x} \in \Omega; M(\mathbf{x}) \leq \frac{1}{n}\}$ tends to 0 when n tends to $+\infty$, we have

$$(2.7) \quad \int_{\Omega} M \left| \frac{\varphi_n - \varphi}{M} \right|^2 \xrightarrow{n \rightarrow +\infty} 0.$$

Concerning the gradient part, we write

$$\nabla \left(\frac{\varphi_n - \varphi}{M} \right) = \nabla \left(\frac{\varphi}{M} \right) \left(\chi \left(\frac{1}{nM} \right) - 1 \right) + \frac{\varphi}{M} \nabla \left(\chi \left(\frac{1}{nM} \right) - 1 \right).$$

The first term is non zero if $M \leq \frac{1}{n}$ and is bounded by $\left| \nabla \left(\frac{\varphi}{M} \right) \right|$ in this case. The last term is non zero if $\frac{1}{2n} \leq M < \frac{1}{n}$ and is bounded by $2 \left| \frac{\nabla M}{M} \frac{\varphi}{M} \right|$. Hence we have

$$\int_{\Omega} M \left| \nabla \left(\frac{\varphi_n - \varphi}{M} \right) \right|^2 \leq \int_{\mathbf{x} \in \Omega; M(\mathbf{x}) \leq \frac{1}{n}} M \left| \nabla \left(\frac{\varphi}{M} \right) \right|^2 + 4 \left| \frac{\nabla M}{M} \frac{\varphi}{\sqrt{M}} \right|^2.$$

Since $\varphi \in H_M^1$, the first term is bounded, and using the result of the Lemma 2.1 again, we know that the last term is bounded too. More precisely, as previously, using the fact that the measure of the set $\{\mathbf{x} \in \Omega; M(\mathbf{x}) \leq \frac{1}{n}\}$ tends to 0 when n tends to $+\infty$, we have

$$(2.8) \quad \int_{\Omega} M \left| \nabla \left(\frac{\varphi_n - \varphi}{M} \right) \right|^2 \xrightarrow{n \rightarrow +\infty} 0.$$

The relations (2.7) and (2.8) ensure that the sequence $\{\varphi_n\}_{n \in \mathbb{N}^*}$ converges to φ in H_M^1 . \square

Now we introduce a compacity result for the spaces L_M^2 and H_M^1 which is comparable to the classical compact injection $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$.

LEMMA 2.4 (Compacity). — *The injection $H_M^1 \hookrightarrow L_M^2$ is compact.*

Proof. — To prove this lemma, we use the following result due to G. Metivier [19, Proposition 3.1 p. 221] affirming that the weight Sobolev space injection $H^1(Mdx) \subset L^2(Mdx)$ as soon as we have $M > 0$ on Ω , $M = 0$ on Γ and $\nabla M \neq 0$ out of Γ . Notice that this last point is a consequence of the two first points $M > 0$ on Ω and $M = 0$ on Γ , at least in a neighborhood of the boundary Γ , what is sufficient for our results. For sake of simplicity, we will nevertheless work with Ω .

Consider a sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ bounded in H_M^1 and show that a convergent sub-sequence can be extracted. By definition of H_M^1 , for all $n \in \mathbb{N}$, there exists $g_n \in H^1(Mdx)$ such that $\varphi_n = Mg_n$. The sequence $\{\varphi_n\}_{n \in \mathbb{N}}$ being bounded in H_M^1 , the sequence $\{g_n\}_{n \in \mathbb{N}}$ is bounded in L_M^2 . We use here the result of G. Métivier. We can extract from the sequence $\{g_n\}_{n \in \mathbb{N}}$ a sub-sequence, still noted $\{g_n\}_{n \in \mathbb{N}}$ and such that

$$g_n \rightharpoonup g \quad \text{in } H^1(Mdx) \quad \text{and} \quad g_n \rightarrow g \quad \text{in } L^2(Mdx).$$

By definition of the spaces H_M^1 and L_M^2 we conclude that

$$\varphi_n \rightharpoonup Mg \quad \text{in } H_M^1 \quad \text{and} \quad \varphi_n \rightarrow Mg \quad \text{in } L_M^2,$$

which proves that the injection $H_M^1 \hookrightarrow L_M^2$ is compact. \square

In the same way, we present the next lemma which proves that functions in H_M^1 are in certain L_M^p , $p > 2$, where the weighted-space L_M^p is defined by

$$L_M^p = \left\{ \varphi \in L_{loc}^1(\Omega) ; \left(\int_{\Omega} M \left| \frac{\varphi}{M} \right|^p \right)^{1/p} < +\infty \right\}$$

and endowed with its usual norm.

This kind of result is essential for the proof of the main theorem (Theorem 3.3); it is proved under the assumptions (\mathcal{H}_1) and (\mathcal{H}_2) . We will note that assumption (\mathcal{H}_2) is used in the following weak formulation (obtained by integration):

$$(\mathcal{H}_2) \quad \implies \quad \exists C' > 0 \quad M \geq C' \delta_{\Gamma}^{1/c}.$$

More exactly, we have

LEMMA 2.5 (Sobolev-type injection). — *If M satisfies (\mathcal{H}_1) and (\mathcal{H}_2) then there exists $p > 2$ such that the injection $H_M^1 \hookrightarrow L_M^p$ is continuous.*

Proof. — First, let us note that $\varphi \in L_M^p$ if and only if $\frac{\varphi}{M^{1-1/p}} \in L^p(\Omega)$ where the spaces $L^p(\Omega)$ are the classical Sobolev spaces on the set Ω . Let $\varphi \in H_M^1$. In the next three steps we will prove that there exists $p > 2$ such that $\frac{\varphi}{M^{1-1/p}} \in L^p(\Omega)$.

• Since $\varphi \in H_M^1$ we clearly have $\varphi \in L_M^2$ and so $\varphi/\sqrt{M} \in L^2$. Moreover, using assumptions Lemma 2.1 with assumption (\mathcal{H}_2) we obtain

$$\nabla \left(\frac{\varphi}{\sqrt{M}} \right) = \underbrace{\frac{\nabla \varphi}{\sqrt{M}}}_{L^2(\Omega)} + \underbrace{\frac{\nabla M}{M} \frac{\varphi}{\sqrt{M}}}_{L^2(\Omega)} \in L^2(\Omega).$$

Consequently, $\varphi/\sqrt{M} \in H^1(\Omega)$ and using the classical Sobolev injections, we deduce that $\varphi/\sqrt{M} \in L^q(\Omega)$ for all $q \leq 2d/(d-2)$ (and for $q < +\infty$ in the 2-dimensional case).

• Using the assumption (\mathcal{H}_2) and the Hardy inequality (Lemma 2.1) we obtain

$$\frac{\varphi}{\sqrt{M}M^c} = \underbrace{\frac{1}{\delta_{\Gamma}} \frac{\varphi}{\sqrt{M}}}_{L^2(\Omega)} \times \underbrace{\frac{\delta_{\Gamma}}{M^c}}_{L^{\infty}(\Omega)} \in L^2(\Omega).$$

• From the two previous steps, we can write that for any $\beta \in \mathbb{R}$ such that $0 \leq \beta \leq c$ we obtain

$$\frac{\varphi}{\sqrt{M}M^{\beta}} = \underbrace{\left(\frac{\varphi}{\sqrt{M}M^c} \right)^{\beta/c}}_{L^{2c/\beta}(\Omega)} \times \underbrace{\left(\frac{\varphi}{\sqrt{M}} \right)^{1-\beta/c}}_{L^{qc/(c-\beta)}(\Omega)} \in L^r(\Omega)$$

with $r = \frac{2cq}{q\beta+2(c-\beta)}$. Let us note p the real such that $1 - \frac{1}{p} = \beta + \frac{1}{2}$. The previous result is written: for any $p \in \mathbb{R}$ such that $2 \leq p \leq \frac{2}{1-2c}$ we obtain

$$\frac{\varphi}{M^{1-1/p}} \in L^r \quad \text{with} \quad r = \frac{4cpq}{qp - 2q + 4cp - 2p + 4}.$$

In particular, we have $\frac{\varphi}{M^{1-1/p}} \in L^p$ as soon as $r \geq p$. The inequality $r \geq p$ holds if and only if $p \leq 2 + \frac{4c(q-2)}{4c+q-2}$. It is thus possible to find $p > 2$ such that $\frac{\varphi}{M^{1-1/p}} \in L^p$. \square

Remark 2.6. — According to the previous proof, we obtain the inclusion $H_M^1 \subset L_M^p$ for all $p \leq 2 + \frac{4c}{c(d-2)+1}$. For instance, in the two dimensional case ($d = 2$) the inclusion $H_M^1 \subset L_M^p$ holds for all $p \leq 2 + 4c$.

To build functions in $H_{M,0}^1$ we use the following lemma which will be important to obtain a lot of test functions in the weak formulation later. The hypothesis (\mathcal{H}_1) allows to use the Lemma 2.3.

LEMMA 2.7. — Assume (\mathcal{H}_1) , (\mathcal{H}_2) hold. Let $\psi \in H_M^1$ and $\xi : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous application, piecewise- \mathcal{C}^1 such that ξ' is bounded on \mathbb{R} . Then we have

$$\varphi := M\xi\left(\frac{\psi}{M}\right) - M \int_{\Omega} M\xi\left(\frac{\psi}{M}\right) \in H_{M,0}^1,$$

with $\nabla\left(\frac{\varphi}{M}\right) = \xi'\left(\frac{\psi}{M}\right)\nabla\left(\frac{\psi}{M}\right)$ and $\|\varphi\|_{H_{M,0}^1} \leq \|\xi'\|_{L^\infty(\mathbb{R})}\|\psi\|_{H_{M,0}^1}$.

Proof. — The proof of this lemma uses the Stampacchia lemma which affirms that if $g \in H^1(\omega)$, ω being an open subset of \mathbb{R}^d , and $\xi : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, piecewise- \mathcal{C}^1 , such that ξ' is bounded on \mathbb{R} then we have $\xi(g) \in H^1(\omega)$ and $\nabla\xi(g) = \xi'(g)\nabla g$.

The Stampacchia lemma is a local result, hence applied to $g = \psi/M$ which is in $H_{\text{loc}}^1(\Omega)$ it shows that the formula $\nabla\xi\left(\frac{\psi}{M}\right) = \xi'\left(\frac{\psi}{M}\right)\nabla\left(\frac{\psi}{M}\right)$ holds in $\mathcal{D}'(\Omega)$. From this formula, it is obvious that $\varphi \in H_M^1$. The fact that φ is null average is then immediate since $\int_{\Omega} M = 1$. \square

One more important ingredient in our study is the following linear operator

$$\mathcal{L}\psi = -\text{div}\left(M\nabla\left(\frac{\psi}{M}\right)\right)$$

on the space L_M^2 and with domain, see [18, Remark 3.8, p. 9] given by

$$D(\mathcal{L}) = \left\{ \psi \in H_M^1 ; \int_{\Omega} \frac{1}{M} \left| \text{div}\left(M\nabla\left(\frac{\psi}{M}\right)\right) \right|^2 < +\infty \right\}.$$

We also find in [18, Proposition 3.6, p. 8] the following result and its proof which will be used to introduce the Galerkin approximation method later.

LEMMA 2.8. — The operator \mathcal{L} is self-adjoint and positive. Moreover, it has a discrete spectrum formed by a sequence $(\ell_n)_{n \in \mathbb{N}}$ such that ℓ_n tends to $+\infty$ when n tends to $+\infty$.

Concerning the uniqueness results for a linear operator, it is known that the eigenvalue 0, that is the kernel of the operator \mathcal{L} , is particularly important.

LEMMA 2.9. — The kernel of the operator \mathcal{L} is the set $\{\lambda M, \lambda \in \mathbb{R}\}$.

Proof. — This lemma is an immediate consequence of the following formulation of the operator \mathcal{L} :

$$(2.9) \quad \langle \mathcal{L}\psi, \varphi \rangle_{L_M^2} = \int_{\Omega} M\nabla\left(\frac{\psi}{M}\right) \cdot \nabla\left(\frac{\varphi}{M}\right)$$

where $\langle \cdot, \cdot \rangle_{L_M^2}$ corresponds to the scalar product subordinated to the norm $\|\cdot\|_{L_M^2}$ on L_M^2 . In fact, let ψ be a function such that $\mathcal{L}\psi = 0$. We obtain $\langle \mathcal{L}\psi, \psi \rangle_{L_M^2} = 0$ and the formulation (2.9) yields $\nabla(\psi/M) = 0$. Thus, thanks to the connexity of Ω , we deduce that $\psi = \lambda M$ with $\lambda \in \mathbb{R}$. \square

The last lemma is a generalized Poincaré inequality adapted to the weighted spaces introduced before. To obtain such a lemma, we use the fact that the potential $V = \ln(M)$ is concave. More precisely we will suppose that

$$(\mathcal{H}_3) \quad \exists \gamma > 0 \quad \nabla \left(\frac{\nabla M}{M} \right) \leq -\gamma \text{Id}.$$

LEMMA 2.10 (Poincaré-type inequality). — *If M satisfies (\mathcal{H}_3) then for all $\varphi \in H_M^1$ we get the following Poincaré-type inequality*

$$\frac{1}{\gamma} \int_{\Omega} M \left| \nabla \left(\frac{\varphi}{M} \right) \right|^2 + \left(\int_{\Omega} \varphi \right)^2 \geq \|\varphi\|_{L_M^2}^2.$$

For the free-average functions (that is for $\psi \in H_{M,0}^1$) this Lemma 2.10 show that the two norms $\|\cdot\|_{H_M^1}$ and $\|\cdot\|_{H_{M,0}^1}$ on this space are equivalents. This equivalence will be usually useful in the remainder of the paper.

Proof. — Let $\varphi \in H_M^1$ and introduce the non-stationary problem

$$\begin{cases} u_t(t, \mathbf{x}) + \mathcal{L}u(t, \mathbf{x}) = 0 & \text{for } (t, \mathbf{x}) \in \mathbb{R} \times \Omega, \\ u(0, \mathbf{x}) = \varphi(\mathbf{x}) & \text{for } \mathbf{x} \in \Omega. \end{cases}$$

The following time-dependant functions

$$D(t) = \int_{\Omega} M \left| \frac{u}{M} \right|^2 \quad \text{and} \quad H(t) = \int_{\Omega} M \left| \nabla \left(\frac{u}{M} \right) \right|^2$$

satisfy

$$D'(t) = 2 \int_{\Omega} \frac{u}{M} \cdot u_t = -2 \int_{\Omega} \frac{u}{M} \cdot \mathcal{L}u = -2 \int_{\Omega} M \left| \nabla \left(\frac{u}{M} \right) \right|^2 = -2H(t).$$

Moreover we have

$$H'(t) = -2 \int_{\Omega} M \nabla \left(\frac{u}{M} \right) \cdot \nabla \left(\frac{\mathcal{L}u}{M} \right).$$

For clarify the following computations, let us introduce the duality operator \mathcal{L}^* such that

$$\mathcal{L}^*v := -\frac{1}{M} \text{div}(M \nabla v) = -\Delta v - \frac{\nabla M}{M} \cdot \nabla v.$$

We have $\mathcal{L}u = M \mathcal{L}^* \left(\frac{u}{M} \right)$ and

$$H'(t) = -2 \int_{\Omega} M \nabla \left(\frac{u}{M} \right) \cdot \nabla \left(\mathcal{L}^* \left(\frac{u}{M} \right) \right).$$

Since $\nabla(\mathcal{L}^*v) = \mathcal{L}^*(\nabla v) - \nabla \left(\frac{\nabla M}{M} \right) \cdot \nabla v$ we deduce that

$$\begin{aligned} H'(t) &= -2 \int_{\Omega} M \nabla \left(\frac{u}{M} \right) \cdot \mathcal{L}^* \left(\nabla \left(\frac{u}{M} \right) \right) \\ &\quad + 2 \int_{\Omega} M \nabla \left(\frac{u}{M} \right) \cdot \left[\nabla \left(\frac{\nabla M}{M} \right) \cdot \nabla \left(\frac{u}{M} \right) \right]. \end{aligned}$$

The first term of the right hand side is written $-2 \int_{\Omega} M \left| \nabla^2 \left(\frac{u}{M} \right) \right|^2$ and is non-positive. Using the assumption (\mathcal{H}_3) , the last term of the right hand side is controled by $-2\gamma \int_{\Omega} M \left| \nabla \left(\frac{u}{M} \right) \right|^2$, that is by $-2\gamma H(t)$. We obtain

$$H'(t) \leq -2\gamma H(t).$$

Hence $H(t) \leq H(0)e^{-2\gamma t}$. Integrate in time, we obtain:

$$(2.10) \quad D(0) - D(+\infty) = 2 \int_0^{+\infty} H(t) dt \leq \frac{1}{\gamma} H(0).$$

To evaluate $D(+\infty)$, we consider a stationary solution u_{∞} . We note that due to the spectral properties of the operator \mathcal{L} (see Lemma 2.8), for any initial data, u tends to a stationary solution u_{∞} as $t \rightarrow +\infty$. By definition it is in the kernel of \mathcal{L} and following the Lemma 2.9 there exists a constant λ

such that $u_\infty = \lambda M$. But the evolution equation on u implies that the mean value $\int_\Omega u$ is conserved: $\int_\Omega u_\infty = \int_\Omega \varphi$, that allows to obtain the constant $\lambda = \int_\Omega \varphi$. We deduce that

$$D(+\infty) = \int_\Omega M \left| \frac{u_\infty}{M} \right|^2 = \left(\int_\Omega \varphi \right)^2.$$

Consequently, the inequality (2.10) corresponds to the following one

$$\int_\Omega M \left| \frac{\varphi}{M} \right|^2 - \left(\int_\Omega \varphi \right)^2 \leq \frac{1}{\gamma} \int_\Omega M \left| \nabla \left(\frac{\varphi}{M} \right) \right|^2$$

which exactly is the inequality announced by the Lemma 2.10. \square

Notice that it is possible to obtain a proof of this Poincaré-type inequality by contradiction, see for instance [18, p.7], or perhaps using the hole-space case (for example for $\Omega = \mathbb{R}^d$) proved in H.J. Brascamp [4] (see also Proposition 2.1 in [6]).

3. Statement of the main theorem

3.1. Definition of weak solution

When we consider the Fokker-Planck equation (1.1) with vector field \mathbf{F} decomposed as the sum $\mathbf{F} = \boldsymbol{\kappa} + \nabla V$ where $\boldsymbol{\kappa} \in L^\infty(\Omega)$ and $e^V \in L^1(\Omega)$, we can introduce the maxwellian function M by $M = \frac{e^V}{\int_\Omega e^V}$ and rewrite (1.1) as $\operatorname{div}(\varphi \boldsymbol{\kappa} - M \nabla(\frac{\varphi}{M})) = f$. If we look for a solution with given average, that is for instance a solution such that $\int_\Omega \varphi = 1$, then we can reduce to the case where φ is free-average exchanging φ into $\varphi - M$ and f into $f - \operatorname{div}(M \boldsymbol{\kappa})$. We obtain the following problem

$$\begin{cases} \operatorname{div}(\varphi \boldsymbol{\kappa} - M \nabla(\frac{\varphi}{M})) = f & \text{in } \Omega, \\ \text{with } \int_\Omega \varphi = 0. \end{cases}$$

Using the adapted spaces introduce in the previous part, the weak formulation of this equation is written: find $\varphi \in H_{M,0}^1$ such that for all $\psi \in H_{M,0}^1$

$$(3.1) \quad \int_\Omega M \nabla \left(\frac{\varphi}{M} \right) \cdot \nabla \left(\frac{\psi}{M} \right) - \int_\Omega \varphi \boldsymbol{\kappa} \cdot \nabla \left(\frac{\psi}{M} \right) = \langle f, \psi \rangle$$

where $\langle \cdot, \cdot \rangle$ denote the duality brackets between H_M^{-1} and $H_{M,0}^1$.

3.2. Assumptions on the potential

In this article we are interested in the case where the vector fields \mathbf{F} quickly explodes near to the boundary. The fact that \mathbf{F} is decomposed as a sum of two terms makes it possible to describe all the “explosive” behavior in the part ∇V . In addition to the fact that V equals $-\infty$ on Γ to ensure the explosion, the assumptions given on V (or on M , which is equivalent) can be checked only in a neighborhood of the boundary Γ . More precisely in order to use the lemmas proved we will use the following assumptions

$$(H_1) \quad \begin{cases} \exists a < 1 \quad \left(\frac{\nabla_R M}{M} \right)^2 + 2 \nabla_R \left(\frac{\nabla_R M}{M} \right) \geq \frac{-a}{\delta_\Gamma^2}, \\ \nabla_R M(0) = 0, \\ \exists b > 0 \quad \nabla_R M \int_\Omega \frac{1}{M} < b, \end{cases}$$

$$(\mathcal{H}_2) \quad \exists c > 0 \quad |\nabla M| \leq \frac{1}{c} \frac{M}{\delta_\Gamma},$$

$$(\mathcal{H}_3) \quad \exists \gamma > 0 \quad \nabla \left(\frac{\nabla M}{M} \right) \leq -\gamma \text{Id},$$

where we recall that ∇_R corresponds to the normal derivative and where δ_Γ represents the distance to Γ .

Notice that we can rewrite these assumptions in term of the potential V (wich is given with respect to the maxwellian M by $V = \ln M$), see for instance Theorem 3.3, page 16. It is important to note that these assumptions are satisfied for the radial functions M (i.e. functions depending only on the distance to the boundary) on the following form near to the boundary

$$M(r) = r^\alpha \quad \text{with } \alpha > 1.$$

In other words, the result is shown for vector fields \mathbf{F} whose the normal component explodes like $\frac{\alpha}{\delta_\Gamma}$ with $\alpha > 1$.

Remark 3.1. — As it was announced as introduction, an interesting case corresponds to the following Fokker-Planck equation

$$-\varepsilon \Delta \varphi + \text{div}(\varphi \tilde{\mathbf{F}}) = f,$$

making appear a small parameter ε . We can come back to the previous case using $\mathbf{F} = \frac{1}{\varepsilon} \tilde{\mathbf{F}}$. We note that if we define a Maxwellian M such that $\mathbf{F} = \nabla M / M$ then the Maxwellian \tilde{M} adapted to $\tilde{\mathbf{F}}$, i.e. such that $\tilde{\mathbf{F}} = \nabla \tilde{M} / \tilde{M}$, satisfies $M = C \tilde{M}^{1/\varepsilon}$. The assumptions on M can thus be interpreted on \tilde{M} and we show that they are less constraining in the following sense: they are checked when the normal component of $\tilde{\mathbf{F}}$ behave like $\frac{\alpha}{\delta_\Gamma}$ for all $\alpha > \varepsilon$.

Concerning the assumption on the “interior” part κ of $\mathbf{F} = \kappa + \nabla V$, that is about $\kappa \in L^\infty(\Omega)$, we can note that this assumption is stronger than that announced by J. Droniou in [7]. In fact, we will see during the proof that the regularity required on κ comes from a product lemma. Roughly speaking, if the product of a function $H^1(\Omega)$ by a function $L^p(\Omega)$ is a function $L^2(\Omega)$ then the theorem is true as soon as κ belongs to $L^p(\Omega)$. In the classical case the usual Sobolev injections $H^1(\Omega) \subset L^{2d/(d-2)}$ imply that $p = d_*$ is sufficient. In our case the injections of “Sobolev” type (see the Lemma 2.5) are not also “generous” and a product $H_M^1 \times L^p(\Omega)$ will not belong to $L^2(\Omega)$ for as many values of p . We can possibly improve the result of the theorem by taking $\kappa \in L^p(\Omega)$ with $p \geq d + 1/c$.

3.3. Main theorem

We prove in Part 4 the following theorem.

THEOREM 3.2. — *Let Ω be a bounded domain of \mathbb{R}^d , $d \geq 2$. We denote by Γ its boundary which is assumed to be of class \mathcal{C}^2 . Let $f \in H_M^{-1}$ and $\mathbf{F} = \kappa + \nabla V$ where $\kappa \in L^\infty(\Omega)$ and $V \in \mathcal{C}^\infty(\Omega)$ satisfies $V = -\infty$ on Γ .*

If the assumptions (\mathcal{H}_1) , (\mathcal{H}_2) and (\mathcal{H}_3) hold then the problem (3.1) admits a unique solution $\varphi \in H_{M,0}^1$.

We can deduce - see the link between a free-average solution and a solution with given average on Subsection 3.1 - the following theorem where we recall all the assumptions

THEOREM 3.3. — *Let Ω be a bounded domain of \mathbb{R}^d , $d \geq 2$. We denote by Γ its boundary which is assumed to be of class \mathcal{C}^2 . Let $f \in H_M^{-1}$ and $\mathbf{F} = \kappa + \nabla V$ where $\kappa \in L^\infty(\Omega)$ and $V \in \mathcal{C}^\infty(\Omega)$*

satisfies $V = -\infty$ on Γ .

If we assume that, in a neighborhood of the boundary Γ , we have

$$(\mathcal{H}_1) \quad \begin{cases} \exists a < 1 & (\nabla_R V)^2 + 2 \nabla_R^2 V \geq \frac{-a}{\delta_\Gamma^2}, \\ \nabla_R V e^V = 0 & \text{on } \Gamma, \\ \exists b > 0 & \nabla_R V e^V \int_\Omega e^{-V} < b, \end{cases}$$

$$(\mathcal{H}_2) \quad \exists c > 0 \quad |\nabla V| \leq \frac{c}{\delta_\Gamma},$$

$$(\mathcal{H}_3) \quad \exists \gamma > 0 \quad \nabla^2 V \leq -\gamma \text{ Id},$$

where ∇_R corresponds to the normal derivative and where δ_Γ represents the distance to Γ , then there exists a unique (weak) solution of the Fokker-Planck equation

$$-\Delta \varphi + \text{div}(\varphi \mathbf{F}) = f \quad \text{in } \Omega,$$

such that $\int_\Omega \varphi = 1$.

4. Proof of the Theorem 3.2

4.1. Existence proof in Theorem 3.2

Principle for the existence proof of Theorem 3.2 - The maxwellian M satisfying the assumptions (\mathcal{H}_1) , (\mathcal{H}_2) and (\mathcal{H}_3) , we use the different lemmas proved in Part 2. For instance, using the equivalence between the norms $\|\cdot\|_{H_M^1}$ and $\|\cdot\|_{H_{M,0}^1}$ on the space $H_{M,0}^1$, see Lemma 2.10, the operator $-\text{div}(M \nabla(\frac{\cdot}{M}))$ is coercive on $H_{M,0}^1$ thus we can (see for instance the Lax-Milgram theorem) prove that there exists a weak solution (that is belonging to $H_{M,0}^1$) to equations like

$$-\text{div}\left(M \nabla\left(\frac{\psi}{M}\right)\right) = f$$

as soon as the source term f belongs in H_M^{-1} . Moreover in this case we have $\|\psi\|_{H_M^1} \lesssim \|f\|_{H_M^{-1}}$.

Because of the non-coercivity of the operator $-\text{div}(M \nabla(\frac{\cdot}{M})) + \text{div}(\cdot \boldsymbol{\kappa})$, we start by studying an approach problem. For each $n \in \mathbb{N}$, let us consider the application $T_n : r \in \mathbb{R} \mapsto \max(\min(r, n), -n) \in \mathbb{R}$ and let us denote by F_n the following application: $F_n : \tilde{\psi} \in L_M^2 \mapsto \psi \in H_{M,0}^1 \subset L_M^2$ where ψ is the weak solution of

$$(4.1) \quad -\text{div}\left(M \nabla\left(\frac{\psi}{M}\right)\right) = f - \text{div}\left(M T_n\left(\frac{\tilde{\psi}}{M}\right) \boldsymbol{\kappa}\right).$$

For $\tilde{\psi} \in L_M^2$ we have $M T_n(\tilde{\psi}/M) \in L_M^2$ and since ⁽⁵⁾ $\boldsymbol{\kappa} \in L^\infty(\Omega)$ we get $\text{div}(M T_n(\frac{\tilde{\psi}}{M}) \boldsymbol{\kappa}) \in H_M^{-1}$. The function F_n is then well defined.

Let us prove that F_n is a compact application by showing that its image $F_n(L_M^2)$ is bounded in H_M^1 . Consider $\psi = F_n(\tilde{\psi}) \in F_n(L_M^2)$. Taking ψ as a test function in the weak formulation of the equation (4.1) we obtain

$$\int_\Omega M \left| \nabla\left(\frac{\psi}{M}\right) \right|^2 = \langle f, \psi \rangle + \int_\Omega M T_n\left(\frac{\tilde{\psi}}{M}\right) \boldsymbol{\kappa} \cdot \nabla\left(\frac{\psi}{M}\right).$$

⁽⁵⁾ Here the assumption on $\boldsymbol{\kappa}$ is essential. Following the proof of J. Droniou [7] it is possible to improve this assumption using the Sobolev injection 2.5 more finely. See discussion concerning the assumptions on $\boldsymbol{\kappa}$ page 16.

In other words, by using the duality definition and the Cauchy-Schwarz inequality, we have

$$\|\psi\|_{H_{M,0}^1}^2 \lesssim \|\psi\|_{H_{M,0}^1} + \|\kappa\|_{L^\infty(\Omega)} \sqrt{\int_{\Omega} M |T_n\left(\frac{\tilde{\psi}}{M}\right)|^2} \sqrt{\int_{\Omega} M |\nabla\left(\frac{\psi}{M}\right)|^2}.$$

Using the fact successively that for all $r \in \mathbb{R}$ we have $|T_n(r)| \leq n$ and that $\int_{\Omega} M = 1$ we deduce that

$$\|\psi\|_{H_{M,0}^1}^2 \lesssim \|\psi\|_{H_{M,0}^1} + n \|\kappa\|_{L^\infty(\Omega)} \|\psi\|_{H_{M,0}^1}.$$

Consequently we have

$$\|F_n(\tilde{\psi})\|_{H_M^1} = \|\psi\|_{H_{M,0}^1} \lesssim 1 + n \|\kappa\|_{L^\infty(\Omega)}.$$

Thus, the image of L_M^2 by the application F_n is contained in the ball of H_M^1 of radius $1 + n \|\kappa\|_{L^\infty(\Omega)}$ (up to a multiplicative constant depending on Ω , which appears in the symbol \lesssim). Moreover, the injection $H_M^1 \hookrightarrow L_M^2$ is compact (see Lemma 2.4) and the application F_n is clearly continuous. Applying the Schauder fixed point theorem, we conclude that the application F_n admits a fixed point, denoted by ψ_n , in L_M^2 . This fixed point is consequently a solution of

$$(4.2) \quad \int_{\Omega} M \nabla\left(\frac{\psi_n}{M}\right) \cdot \nabla\left(\frac{\varphi}{M}\right) - \int_{\Omega} M T_n\left(\frac{\psi_n}{M}\right) \kappa \cdot \nabla\left(\frac{\varphi}{M}\right) = \langle f, \varphi \rangle$$

for all test functions $\varphi \in H_{M,0}^1$.

The continuation of the proof consists of obtaining estimates on these functions ψ_n in order to be able to pass to the limit when n tends to $+\infty$.

Estimate of $M \ln(1 + |\psi_n/M|)$ in $H_{M,0}^1$ -norm - Let ξ be the application from \mathbb{R} to \mathbb{R} defined by $\xi(r) = \int_0^r \frac{ds}{(1+|s|)^2}$. This application is continuous, piecewise- \mathcal{C}^1 and with a bounded derivative. According to Lemma 2.7 we can choose $\varphi = M\xi(\psi_n/M) - M \int_{\Omega} M\xi(\psi_n/M)$ as a test function in formulation (4.2).

- The first of the three terms obtained is treated in the following way

$$(4.3) \quad \begin{aligned} \int_{\Omega} M \nabla\left(\frac{\psi_n}{M}\right) \cdot \nabla\left(\xi\left(\frac{\psi_n}{M}\right)\right) &= \int_{\Omega} M \frac{|\nabla\left(\frac{\psi_n}{M}\right)|^2}{\left(1 + \left|\frac{\psi_n}{M}\right|\right)^2} \\ &= \left\| M \ln\left(1 + \left|\frac{\psi_n}{M}\right|\right) \right\|_{H_{M,0}^1}^2. \end{aligned}$$

- For the second term we obtain

$$\begin{aligned} \left| \int_{\Omega} M T_n\left(\frac{\psi_n}{M}\right) \kappa \cdot \nabla\left(\xi\left(\frac{\psi_n}{M}\right)\right) \right| &= \left| \int_{\Omega} \frac{M T_n\left(\frac{\psi_n}{M}\right)}{1 + \left|\frac{\psi_n}{M}\right|} \kappa \cdot \frac{\nabla\left(\frac{\psi_n}{M}\right)}{1 + \left|\frac{\psi_n}{M}\right|} \right| \\ &\leq \|\kappa\|_{L^\infty(\Omega)} \int_{\Omega} \left| \frac{T_n\left(\frac{\psi_n}{M}\right)}{1 + \left|\frac{\psi_n}{M}\right|} \right| M \left| \nabla\left(\ln\left(1 + \left|\frac{\psi_n}{M}\right|\right)\right) \right|. \end{aligned}$$

Using the fact that for all $r \in \mathbb{R}$, we have $|T_n(r)| \leq |r|$, we deduce that⁽⁶⁾

$$(4.4) \quad \left| \int_{\Omega} M T_n\left(\frac{\psi_n}{M}\right) \kappa \cdot \nabla\left(\xi\left(\frac{\psi_n}{M}\right)\right) \right| \lesssim \left\| M \ln\left(1 + \left|\frac{\psi_n}{M}\right|\right) \right\|_{H_{M,0}^1}.$$

⁽⁶⁾ We also use the Cauchy-Schwarz inequality to show that

$$\int_{\Omega} M f \leq \sqrt{\int_{\Omega} M} \sqrt{\int_{\Omega} M f^2} = \sqrt{\int_{\Omega} M f^2}.$$

- For the last term, using $f \in H_M^{-1}$, we deduce

$$\begin{aligned}
 (4.5) \quad |\langle f, \varphi \rangle| &\lesssim \|\varphi\|_{H_{M,0}^1} = \sqrt{\int_{\Omega} M \xi' \left(\frac{\psi_n}{M} \right) \nabla \left(\frac{\psi_n}{M} \right)} \\
 &= \sqrt{\int_{\Omega} M \frac{|\nabla \left(\frac{\psi_n}{M} \right)|^2}{\left(1 + \left| \frac{\psi_n}{M} \right|\right)^2}} = \left\| M \ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right\|_{H_{M,0}^1}.
 \end{aligned}$$

The three estimates (4.3), (4.4) and (4.5) enable us to obtain for all $n \in \mathbb{N}$

$$(4.6) \quad \left\| M \ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right\|_{H_{M,0}^1} \lesssim 1.$$

Estimate of $\mu(\{\mathbf{Q} \in \Omega ; |\psi_n(\mathbf{Q})| \geq kM(\mathbf{Q})\})$ - In this paragraph, we control the size of the set where ψ_n has a large value, that is the set $\mathcal{E}_k = \{\mathbf{Q} \in \Omega ; |\psi_n(\mathbf{Q})| \geq kM(\mathbf{Q})\}$ for $k \in \mathbb{N}$. The natural measure in the present context is the measure $d\mu = M(\mathbf{Q})d\mathbf{Q}$ ($d\mathbf{Q}$ being the classical Lebesgue measure on $\Omega \subset \mathbb{R}^d$), which enables to take into account the weight of the Maxwellian M . Writing $\mathcal{E}_k = \{\mathbf{Q} \in \Omega ; (\ln(1 + |\psi_n(\mathbf{Q})/M(\mathbf{Q})|))^2 \geq (\ln(1 + k))^2\}$ we obtain

$$\int_{\Omega} M \left(\ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right)^2 = \int_{\mathcal{E}_k} M \left(\ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right)^2 + \int_{\Omega \setminus \mathcal{E}_k} M \left(\ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right)^2.$$

We easily deduce the following estimate

$$\int_{\Omega} M \left(\ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right)^2 \geq \int_{\mathcal{E}_k} M \left(\ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right)^2 \geq \int_{\mathcal{E}_k} M (\ln(1 + k))^2.$$

Introducing the measure $d\mu = M(\mathbf{Q})d\mathbf{Q}$ this inequality is also rewritten

$$\mu(\mathcal{E}_k) \leq \frac{1}{(\ln(1 + k))^2} \left\| M \ln \left(1 + \left| \frac{\psi_n}{M} \right| \right) \right\|_{L_M^2}^2.$$

Taking into account the estimate (4.6), the previous equation is written

$$(4.7) \quad \mu(\{\mathbf{Q} \in \Omega ; |\psi_n(\mathbf{Q})| \geq kM(\mathbf{Q})\}) \lesssim \frac{1}{(\ln(1 + k))^2}.$$

Estimate of $MS_k(\psi_n/M)$ in $H_{M,0}^1$ -norm - Recall that for $k \in \mathbb{N}$ the application T_k is given by $T_k : r \in \mathbb{R} \mapsto \max(\min(r, k), -k) \in \mathbb{R}$. We now define the application S_k such that $T_k + S_k = \text{id}$. To obtain an estimate on ψ_n we successively obtain an estimate on $MS_k(\psi_n/M)$ and then on $MT_k(\psi_n/M)$ for a sufficiently large $k \in \mathbb{N}$.

Let $k \in \mathbb{N}$. Taking $\varphi = MS_k(\psi_n/M) - M \int_{\Omega} MS_k(\psi_n/M)$ as a test function test in (4.2). According to Lemma 2.7, this choice is possible and we obtain

$$\underbrace{\int_{\Omega} M \nabla \left(\frac{\psi_n}{M} \right) \cdot \nabla \left(S_k \left(\frac{\psi_n}{M} \right) \right)}_A - \underbrace{\int_{\Omega} MT_n \left(\frac{\psi_n}{M} \right) \kappa \cdot \nabla \left(S_k \left(\frac{\psi_n}{M} \right) \right)}_B = \underbrace{\langle f, \varphi \rangle}_C.$$

- Since $S_k + T_k = \text{id}$ and for all $r \in \mathbb{R}$ we have $S'_k(r) = 0$ or $T'_k(r) = 0$ we deduce that the first term A is written

$$(4.8) \quad A = \int_{\Omega} M \left| \nabla \left(S_k \left(\frac{\psi_n}{M} \right) \right) \right|^2 = \left\| MS_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1}^2.$$

- Using the fact that for all $r \in \mathbb{R}$ we have $|T_n(r)| \leq |r|$ and using the Cauchy-Schwarz inequality, we estimate the second term B in the following way

$$|B| \leq \|\kappa\|_{L^\infty(\Omega)} \sqrt{\int_{\Omega} \frac{|\psi_n|^2}{M}} \sqrt{\int_{\Omega} M \left| \nabla \left(S_k \left(\frac{\psi_n}{M} \right) \right) \right|^2}.$$

However $|\psi_n/M| = |T_k(\psi_n/M) + S_k(\psi_n/M)| \leq k + |S_k(\psi_n/M)|$ thus $|\psi_n/\sqrt{M}| \leq k\sqrt{M} + \sqrt{M}|S_k(\psi_n/M)|$ and using the triangular inequality we obtain

$$\sqrt{\int_{\Omega} \frac{|\psi_n|^2}{M}} \leq \sqrt{\int_{\Omega} k^2 M} + \sqrt{\int_{\Omega} M |S_k\left(\frac{\psi_n}{M}\right)|^2} = k + \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{L_M^2}.$$

Since $S_k(r) = 0$ for $|r| < k$, we can estimate this last term as follows:

$$\left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{L_M^2}^2 = \int_{\Omega} M |S_k\left(\frac{\psi_n}{M}\right)|^2 = \int_{\mathcal{E}_k} M |S_k\left(\frac{\psi_n}{M}\right)|^2,$$

where we recall that $\mathcal{E}_k = \{\mathbf{Q} \in \Omega ; |\psi_n(\mathbf{Q})| \geq kM(\mathbf{Q})\}$. According to the Hölder inequality, for all $p > 1$, denoting by q the conjugate of p (i.e. such that $\frac{1}{p} + \frac{1}{q} = 1$) and using the estimate (4.7), we obtain

$$\begin{aligned} \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{L_M^2}^2 &\leq \left(\int_{\mathcal{E}_k} M \right)^{1/q} \left(\int_{\mathcal{E}_k} M |S_k\left(\frac{\psi_n}{M}\right)|^{2p} \right)^{1/p} \\ &\lesssim \frac{1}{(\ln(1+k))^{2/q}} \left(\int_{\Omega} M |S_k\left(\frac{\psi_n}{M}\right)|^{2p} \right)^{1/p}. \end{aligned}$$

We thus control the L_M^2 -norm of $MS_k(\psi_n/M)$ using his L_M^{2p} -norm. But this L_M^{2p} -norms can itself be controlled, for an adapted value of p by the H_M^1 -norm. In fact, using the weighted Sobolev embedding (see Lemma 2.5) there exists $p > 1$ for which we have the inequality

$$\begin{aligned} \left(\int_{\Omega} M |S_k\left(\frac{\psi_n}{M}\right)|^{2p} \right)^{1/p} &\lesssim \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_M^1}^2 \\ &\lesssim \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1}^2 + \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{L_M^2}^2. \end{aligned}$$

We deduce a control on the L_M^2 -norm of $MS_k(\psi_n/M)$ using his $H_{M,0}^1$ -norm:

$$\left(1 - \frac{1}{(\ln(1+k))^{2/q}} \right) \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{L_M^2}^2 \lesssim \frac{1}{(\ln(1+k))^{2/q}} \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1}^2,$$

that is a control of the form $\left\| MS_k(\psi_n/M) \right\|_{L_M^2} \leq \mathcal{A}(k) \left\| MS_k(\psi_n/M) \right\|_{H_{M,0}^1}$ where $\mathcal{A}(k)$ tends to 0 when k tends to $+\infty$. Hence, we obtain the following estimate for the term B of the left hand side of equation (4.1):

$$(4.9) \quad |B| \leq \|\kappa\|_{L^\infty(\Omega)} \left(k + \mathcal{A}(k) \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1} \right) \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1}.$$

• The last term of the equation (4.1) is controlled as follow

$$(4.10) \quad \begin{aligned} |C| = |\langle f, \varphi \rangle| &\lesssim \|\varphi\|_{H_{M,0}^1} = \sqrt{\int_{\Omega} M \left| \nabla \left(\frac{\varphi}{M} \right) \right|^2} \\ &\lesssim \sqrt{\int_{\Omega} M \left| \nabla \left(S_k \left(\frac{\varphi}{M} \right) \right) \right|^2} = \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1}. \end{aligned}$$

The previous estimates (4.8), (4.9) and (4.10) enable the deduction, from equation (4.1), for all $k \in \mathbb{N}$, of the following inequality

$$\left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1} \lesssim \|\kappa\|_{L^\infty(\Omega)} \left(k + \mathcal{A}(k) \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1} \right) + 1.$$

Since $\mathcal{A}(k)$ tends to 0 when k tends to $+\infty$, it possible to obtain for a sufficiently large k , the inequality

$$(4.11) \quad \left\| MS_k\left(\frac{\psi_n}{M}\right) \right\|_{H_{M,0}^1} \lesssim 1.$$

Estimate of $MT_k(\psi_n/M)$ in $H_{M,0}^1$ -norm - Choose now $\varphi = MT_k(\psi_n/M) - M \int_{\Omega} MT_k(\psi_n/M)$ as a test function in equation (4.2) (according to Lemma 2.7 we have $\varphi \in H_{M,0}^1$). As for the estimate of $MS_k(\psi_n/M)$, we study each of three terms, named A , B and C as previously, present in equation (4.2).

- The first is written

$$A = \int_{\Omega} M \left| \nabla \left(T_k \left(\frac{\psi_n}{M} \right) \right) \right|^2 = \left\| MT_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1}^2.$$

- For the second term, we proceed as follow:

$$\begin{aligned} |B| &\leq \|\kappa\|_{L^\infty(\Omega)} \int_{\Omega} M \left| T_n \left(\frac{\psi_n}{M} \right) \right| \left| \nabla \left(T_k \left(\frac{\psi_n}{M} \right) \right) \right| \\ &\leq \|\kappa\|_{L^\infty(\Omega)} \int_{\Omega} |\psi_n| \left| \nabla \left(T_k \left(\frac{\psi_n}{M} \right) \right) \right|. \end{aligned}$$

But for $|\psi_n/M| \geq k$ we have $\nabla(T_k(\psi_n/M)) = 0$ whereas for $|\psi_n/M| < k$ we clearly have $|\psi_n| < kM$ and consequently, according to the Cauchy-Schwarz inequality we obtain

$$\begin{aligned} |B| &\leq \|\kappa\|_{L^\infty(\Omega)} \int_{\Omega} kM \left| \nabla \left(T_k \left(\frac{\psi_n}{M} \right) \right) \right| \\ &\leq \|\kappa\|_{L^\infty(\Omega)} \sqrt{\int_{\Omega} k^2 M} \sqrt{\int_{\Omega} M \left| \nabla \left(T_k \left(\frac{\psi_n}{M} \right) \right) \right|^2} \\ &\leq k \|\kappa\|_{L^\infty(\Omega)} \left\| MT_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1}. \end{aligned}$$

- The last term is treated like those of the previous estimates:

$$|C| = |\langle f, \varphi \rangle| \lesssim \left\| MT_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1}.$$

These three estimates give (note that this estimate depends on k , but that k has been fixed)

$$(4.12) \quad \left\| MT_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1} \lesssim 1.$$

Estimate of ψ_n in $H_{M,0}^1$ - Since for all $k \in \mathbb{N}$ we have $S_k + T_k = \text{id}$ we obtain

$$\begin{aligned} \|\psi_n\|_{H_{M,0}^1} &= \left\| MS_k \left(\frac{\psi_n}{M} \right) + MT_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1} \\ &\leq \left\| MS_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1} + \left\| MT_k \left(\frac{\psi_n}{M} \right) \right\|_{H_{M,0}^1}. \end{aligned}$$

Using the estimates (4.11) and (4.12) we deduce that for all $n \in \mathbb{N}$ we have

$$(4.13) \quad \|\psi_n\|_{H_{M,0}^1} \lesssim 1.$$

Convergence of the sequence $\{\psi_n\}_{n \in \mathbb{N}}$ - According to the estimate (4.13) the sequence $\{\psi_n\}_{n \in \mathbb{N}}$ is bounded in $H_{M,0}^1$. According to the Lemma 2.4 a subsequence of the sequence $\{\psi_n\}_{n \in \mathbb{N}}$ (always denoted by $\{\psi_n\}_{n \in \mathbb{N}}$) admits a limit ψ , weak in $H_{M,0}^1$ and strong in L_M^2 . In order to perform the limit in equation (4.2), it is sufficient to prove that the sequence $\{MT_n(\psi_n/M)\}_{n \in \mathbb{N}}$ tends to ψ in L_M^2 . We obtain

$$\left\| MT_n \left(\frac{\psi_n}{M} \right) - \psi \right\|_{L_M^2}^2 \leq \left\| MT_n \left(\frac{\psi_n}{M} \right) - MT_n \left(\frac{\psi}{M} \right) \right\|_{L_M^2}^2 + \left\| MT_n \left(\frac{\psi}{M} \right) - \psi \right\|_{L_M^2}^2.$$

However the application $T : \mathbb{R} \rightarrow \mathbb{R}$ is 1-lipschitz and we have

$$\begin{aligned} \left\| MT_n\left(\frac{\psi_n}{M}\right) - MT_n\left(\frac{\psi}{M}\right) \right\|_{L_M^2}^2 &= \int_{\Omega} M \left| T_n\left(\frac{\psi_n}{M}\right) - T_n\left(\frac{\psi}{M}\right) \right|^2 \\ &\leq \int_{\Omega} M \left| \frac{\psi_n}{M} - \frac{\psi}{M} \right|^2 = \|\psi_n - \psi\|_{L_M^2}^2, \end{aligned}$$

which proves that $\|MT_n(\psi_n/M) - MT_n(\psi/M)\|_{L_M^2}$ tends to 0 when n tends to $+\infty$. As regards the other term, the Lebesgue convergence dominated theorem directly affirms that $\|MT_n(\psi/M) - \psi\|_{L_M^2}$ also tends to 0 when n tends to $+\infty$. Finally, it was shown that the sequence $\{MT_n(\psi_n/M)\}_{n \in \mathbb{N}}$ converges to ψ in L_M^2 and consequently that ψ is a solution of

$$\int_{\Omega} M \nabla\left(\frac{\psi}{M}\right) \cdot \nabla\left(\frac{\varphi}{M}\right) - \int_{\Omega} \psi \boldsymbol{\kappa} \cdot \nabla\left(\frac{\varphi}{M}\right) = \langle f, \varphi \rangle \quad \forall \varphi \in H_{M,0}^1.$$

4.2. Uniqueness proof in Theorem 3.2

Main steps for the uniqueness proof - To prove uniqueness, we proceed as follows: We start by introducing the dual problem. It is shown that this dual problem admits a solution by using the Schauder topological degree method. Then, by using the existence both problem and its dual, we deduce uniqueness from these two problems.

Introduction of the dual problem - For $g \in H_M^{-1}$ let us consider the elliptic partial differential equation

$$(4.14) \quad -\operatorname{div}\left(M \nabla\left(\frac{\phi}{M}\right)\right) - M \boldsymbol{\kappa} \cdot \nabla\left(\frac{\phi}{M}\right) = g \quad \text{on } \Omega$$

and we look for a solution $\phi \in H_{M,0}^1$ to this equation.

A compact application for the dual problem - For $\tilde{\phi} \in H_{M,0}^1$ we have $M \boldsymbol{\kappa} \cdot \nabla(\tilde{\phi}/M) \in L_M^2 \subset H_M^{-1}$ since $\|M \boldsymbol{\kappa} \cdot \nabla(\tilde{\phi}/M)\|_{L_M^2} \leq \|\boldsymbol{\kappa}\|_{L^\infty(\Omega)} \|\tilde{\phi}\|_{H_{M,0}^1}$. Since the operator $\varphi \mapsto -\operatorname{div}(M \nabla(\varphi/M))$ is coercive in $H_{M,0}^1$, there exists thus a unique solution $\phi = G(\tilde{\phi}) \in H_{M,0}^1$ such that for all $\varphi \in H_{M,0}^1$

$$(4.15) \quad \int_{\Omega} M \nabla\left(\frac{\phi}{M}\right) \cdot \nabla\left(\frac{\varphi}{M}\right) - \int_{\Omega} \varphi \boldsymbol{\kappa} \cdot \nabla\left(\frac{\tilde{\phi}}{M}\right) = \langle g, \varphi \rangle.$$

This defines an application $G : H_{M,0}^1 \rightarrow H_{M,0}^1$. It is quite easy to see that G is continuous; indeed, if $\tilde{\phi}_n$ tends to $\tilde{\phi}$ in $H_{M,0}^1$ then $M \boldsymbol{\kappa} \cdot \nabla(\tilde{\phi}_n/M)$ tends to $M \boldsymbol{\kappa} \cdot \nabla(\tilde{\phi}/M)$ in H_M^{-1} (more precisely in L_M^2). Thus $\operatorname{div}(M \nabla(\phi_n/M))$ tends to $\operatorname{div}(M \nabla(\phi/M))$ which implies that $\phi_n = G(\tilde{\phi}_n)$ tends to $\phi = G(\tilde{\phi})$ in $H_{M,0}^1$.

We will now prove that G is a compact operator. Suppose that the sequence $\{\tilde{\phi}_n\}_{n \in \mathbb{N}}$ is bounded in $H_{M,0}^1$; then $\{M \boldsymbol{\kappa} \cdot \nabla(\tilde{\phi}_n/M)\}_{n \in \mathbb{N}}$ is bounded in H_M^{-1} so that, using $\varphi = G(\tilde{\phi}_n) = \phi_n$ as a test function in the equation satisfied by ϕ_n , we get using the Lemma 2.10

$$\|\phi_n\|_{H_{M,0}^1}^2 \lesssim \left(1 + \left\| M \boldsymbol{\kappa} \cdot \nabla\left(\frac{\tilde{\phi}_n}{M}\right) \right\|_{H_M^{-1}}\right) \|\tilde{\phi}_n\|_{H_{M,0}^1},$$

which implies that the sequence $\{\phi_n\}_{n \in \mathbb{N}}$ is bounded in $H_{M,0}^1$. Using the Lemma 2.4, up to a subsequence, we can thus suppose that $\{\phi_n\}_{n \in \mathbb{N}}$ converges a.e. on Ω and is bounded in L_M^2 . Let $(n, m) \in \mathbb{N}^2$; subtract the equation satisfied by ϕ_m to the equation satisfied by ϕ_n and use $\varphi = \phi_n - \phi_m$ as a test function, this gives using the Lemma 2.10 again

$$\|\phi_n - \phi_m\|_{H_{M,0}^1}^2 \leq \left| \int_{\Omega} (\phi_n - \phi_m) \boldsymbol{\kappa} \cdot \nabla\left(\frac{\tilde{\phi}_n - \tilde{\phi}_m}{M}\right) \right| \lesssim \|\phi_n - \phi_m\|_{L_M^2}.$$

From the strong convergence of $\{\phi_n\}_{n \in \mathbb{N}}$ to ϕ in L^2_M we deduce that the sequence $\{\phi_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence in $H^1_{M,0}$ and converges in this space. We deduce that the application G is compact.

Existence result for the dual problem using the Leray-Schauder topological degree -

According to the Leray-Schauder topological theory (see the founder article of J. Leray and J. Schauder [16]) since the operator G introduced with equation (4.15) is a compact operator, to prove that it has a fixed point, we just have to find $R > 0$ such that for all $s \in [0, 1]$ there exists no solution of $\phi - sG(\phi) = 0$ satisfying $\|\phi\|_{H^1_{M,0}} = R$.

Let $s \in [0, 1]$ and suppose that $\phi \in H^1_{M,0}$ satisfies $\phi = sG(\phi)$. We have for all $\varphi \in H^1_{M,0}$

$$(4.16) \quad \int_{\Omega} M \nabla \left(\frac{\phi}{M} \right) \cdot \nabla \left(\frac{\varphi}{M} \right) - s \int_{\Omega} \varphi \kappa \cdot \nabla \left(\frac{\phi}{M} \right) = \langle sg, \varphi \rangle.$$

Using the ‘‘non-dual’’ problem (see the existence proof of Theorem 3.2 where we obtain an existence solution of equation (4.2)), we know that for all $f \in H^{-1}_M$ there exists at least one solution $\psi \in H^1_{M,0}$ such that for all $\varphi \in H^1_{M,0}$

$$(4.17) \quad \int_{\Omega} M \nabla \left(\frac{\psi}{M} \right) \cdot \nabla \left(\frac{\varphi}{M} \right) - s \int_{\Omega} \psi \kappa \cdot \nabla \left(\frac{\varphi}{M} \right) = \langle f, \varphi \rangle.$$

Moreover, according to estimate (4.13) there exists $C_1 \in \mathbb{R}^+$ such that for all $f \in H^{-1}_M$ with $\|f\|_{H^{-1}_M} \leq 1$ and for all $s \in [0, 1]$ we have $\|\psi\|_{H^1_{M,0}} \leq C_1$. We can verify that this constant C_1 depends only on $\|f\|_{H^{-1}_M}$ and can be selected independently on the function f when $\|f\|_{H^{-1}_M} \leq 1$. In addition according to the estimates obtained in the existence proof of Theorem 3.2 this constant C_1 depends on $\|s\kappa\|_{L^\infty(\Omega)} \leq \|\kappa\|_{L^\infty(\Omega)}$ and consequently the constant C_1 can also be selected independently of s .

By taking $\varphi = \phi$ in the equation (4.17) satisfied by ψ and $\varphi = \psi$ in the equation (4.16) satisfied by ϕ , we obtain

$$\langle f, \phi \rangle = \langle sg, \psi \rangle \leq s \|g\|_{H^{-1}_M} C_1 \leq \|g\|_{H^{-1}_M} C_1 := C_2.$$

Since this inequality is satisfied for all $f \in H^{-1}_M$ such that $\|f\|_{H^{-1}_M} \leq 1$, we deduce that $\|\phi\|_{H^1_{M,0}} \leq C_2$. Now take $R = C_2 + 1$. We have just proven that, for any $s \in [0, 1]$, any solution of $\phi - sG(\phi) = 0$ satisfies $\|\phi\|_{H^1_{M,0}} < R$; thus by the Leray-Schauder topological degree theory, the application G has a fixed point, that is to say a solution of (4.15).

Uniqueness - Since the equation (3.1) is linear, it is sufficient to prove that the only solution of (3.1) without source term, i.e. taking $f = 0$, is the null function. Let ψ be a solution of (3.1) with $f = 0$ and let ϕ be a solution of (4.14) with $g = \text{sign}(\psi) \in H^{-1}_M$. By putting $\varphi = \phi$ as a test function in the equation (3.1) satisfied by ψ and $\varphi = \psi$ as a test function in the weak formulation of the equation (4.14) satisfied by ϕ , we respectively get

$$\begin{aligned} \int_{\Omega} M \nabla \left(\frac{\psi}{M} \right) \cdot \nabla \left(\frac{\phi}{M} \right) - \int_{\Omega} \psi \kappa \cdot \nabla \left(\frac{\phi}{M} \right) &= 0 \quad \text{and} \\ \int_{\Omega} M \nabla \left(\frac{\phi}{M} \right) \cdot \nabla \left(\frac{\psi}{M} \right) - \int_{\Omega} \psi \kappa \cdot \nabla \left(\frac{\phi}{M} \right) &= \langle \text{sign}(\psi), \psi \rangle. \end{aligned}$$

We deduce that $\langle \text{sign}(\psi), \psi \rangle = 0$, that is to say $\int_{\Omega} |\psi| = 0$ and then $\psi = 0$.

Remark 4.1. — A similar reasoning gives the uniqueness of the solution of the dual problem (4.15).

5. Application to fluid mechanics

5.1. The FENE model for dilute polymers

A natural framework where vectors fields strongly explode at the boundaries of a domain is the framework of the modeling of the spring whose extension is finite (that is physically realist). In fluid mechanics, such an approach is used to develop polymer models in solution. It is this point of view which we have chooses to present in order to illustrate the preceding theoretical study.

The simplest micro-mechanical approach to model the polymer molecules in a dilute solution is the dumbbell model in which the polymers are represented by two beads connected by a spring. The configuration vector \mathbf{Q} describes the orientation and the elongation of such a dumbbell [15, 17]. The force of the spring is governed by some law that should be derived from physical arguments. We choose here the popular FENE model, in which the maximum extensibility of the dumbbell is fixed at some value determined by the dimensionless parameter ℓ and the spring force takes the simple form

$$\mathbf{E}(\mathbf{Q}) = \frac{\mathbf{Q}}{1 - |\mathbf{Q}|^2/\ell^2}.$$

The configuration vector \mathbf{Q} depends on time t and macroscopic position of the dumbbell \mathbf{x} in the flow. Moreover, it satisfies the following stochastic differential equation (see [21] for details):

$$(5.1) \quad d\mathbf{Q} = \left((\nabla \mathbf{u})^T \cdot \mathbf{Q} - \frac{1}{2\mathcal{D}e} \mathbf{E}(\mathbf{Q}) \right) dt + \sqrt{\frac{1}{2\mathcal{D}e}} d\mathbf{W},$$

where the 2-tensor $(\nabla \mathbf{u})^T$ is the transposed velocity gradient, $\mathcal{D}e$ is a dimensionless number called the Deborah number (linked to the relaxation time of the fluid) and \mathbf{W} is the Wiener random process that accounts for the Brownian forces acting on each bead. Equation (5.1) should be understood as the Itô ordinary stochastic differential equations along the particle paths since the dumbbells' centers of mass are supposed on average to follow the particules of the solvent fluid.

As is well known (see Section 3.3 of [21]), every Itô ordinary stochastic differential equation can be associated with a partial differential equation for the probability density function $\varphi(t, \mathbf{x}, \mathbf{Q})$ of the random process $\mathbf{Q}(t, \mathbf{x})$. In particular, equation (5.1) implies the following, also called Fokker-Planck, equation for $\varphi(t, \mathbf{x}, \mathbf{Q})$:

$$(5.2) \quad \frac{\partial \varphi}{\partial t} + \mathbf{u} \cdot \nabla_{\mathbf{x}} \varphi = \frac{1}{2\mathcal{D}e} \Delta_{\mathbf{Q}} \varphi - \operatorname{div}_{\mathbf{Q}} \left(\varphi \left((\nabla \mathbf{u})^T \cdot \mathbf{Q} - \frac{1}{2\mathcal{D}e} \mathbf{E} \right) \right).$$

In certain modes the dominating terms correspond to the terms of the right-hand side member of equation (5.2). It is the case, for instance, when the flow is supposed to be thin, see [5]. In these configurations, the distribution φ can be seen like depending only on \mathbf{Q} (to be rigorous, φ also depends on time t and on the macroscopic position \mathbf{x} , *via* the presence of the gradient $\nabla \mathbf{u}(t, \mathbf{x})$ but these dependences can be seen as parameters) and the equation (5.2) is approached by the following Fokker-Planck equation on $\varphi(\mathbf{Q})$:

$$(5.3) \quad -\Delta \varphi + \operatorname{div}(\varphi \mathbf{F}) = 0 \quad \text{in } B(0, \ell),$$

with $\mathbf{F} = 2\mathcal{D}e (\nabla \mathbf{u})^T \cdot \mathbf{Q} - \mathbf{E}$. It exactly corresponds to those studied in the first part of this paper (see equation (1.1)) in the case $\Omega = B(0, \ell)$ and without source term: $f = 0$.

Although it is wished that the solution $\varphi(\mathbf{Q})$ cancels for values \mathbf{Q} such that $|\mathbf{Q}| = \ell$ (i.e. we wish that the maximum length of the springs is ℓ and that there is no spring of this length), the classical framework of the Theorem 1.2 does not correspond to this equation provided with the homogeneous Dirichlet boundary conditions. In fact, the force \mathbf{F} is not sufficiently regular: we have $\mathbf{F} \notin L^d(B(0, \ell))$. Roughly speaking, the FENE model takes into account the finite extensibility of the polymer chain, through an important explosive force when $|\mathbf{Q}|$ tends to ℓ .




Figure 5.1. On the left, we have drawn the physical domain of a real flow for a diluted polymers solution. From the microscopic point of view, the polymer chains are identified to independent mass-springs systems (called dumbbells). The orientation and the length of each dumbbell is governed by a quantity (denoted by φ in this paper) distributed in a ball whose radius corresponds to the maximum extension of the spring. On the right, the colors correspond to the various probabilities that dumbbell be in the given position. For instance, the drawn dumbbell is the dumbbell which has the most chance to be present (with its “symmetrical” compared to the center of the ball).

On the other hand, this force \mathbf{F} perfectly corresponds to the principal result shown in this article (see Theorem 3.3). More precisely, the vector field $\mathbf{Q} \in B(0, \ell) \mapsto 2De (\nabla \mathbf{u})^T \cdot \mathbf{Q}$ is clearly bounded and we can write the “explosive” term \mathbf{E} as follow:

$$\mathbf{E} = -\nabla V \quad \text{with} \quad V(\mathbf{Q}) = \frac{\ell^2}{2} \ln \left(1 - \frac{|\mathbf{Q}|^2}{\ell^2} \right).$$

To make appear the maxwellian function M as it is used in this paper, we write

$$(5.4) \quad M(\mathbf{Q}) = \frac{e^{V(\mathbf{Q})}}{\int_{\Omega} e^{V(\mathbf{R})} d\mathbf{R}} = \frac{\left(1 - |\mathbf{Q}|^2/\ell^2 \right)^{\ell^2/2}}{\int_{B(0,\ell)} \left(1 - |\mathbf{R}|^2/\ell^2 \right)^{\ell^2/2} d\mathbf{R}}.$$

From Theorem 3.3 we deduce that if $\ell > \sqrt{2}$ then for any $\rho \in \mathbb{R}$ there exists a unique (weak) solution of the Fokker-Planck equation (5.3) such that $\int_{B(0,\ell)} \varphi = \rho$. According to H.C. Öttinger [21], the number ℓ roughly measures the number of monomer units represented by a bead and it is generally larger than 10. The assumption $\ell > \sqrt{2}$ is not constraining from the physical point of view. In fact, according to H.C. Öttinger [21], the number ℓ roughly measures the number of monomer units represented by a bead and it is generally larger than 10.

Moreover, impose the quantity $\int_{B(0,\ell)} \varphi$ physically corresponds to given the density of the polymer chains. Hence this condition is relevant for the studied problem.

Remark 5.1. — If the tensor $(\nabla_{\mathbf{x}} \mathbf{u})^T$ is replaced by its anti-symmetric part $\frac{1}{2}(\nabla_{\mathbf{x}} \mathbf{u} - (\nabla_{\mathbf{x}} \mathbf{u})^T)$ in the force term \mathbf{F} then we get the so-called co-rotational FENE model. This case corresponds to a particular cases presented page 2: $\varphi = M$ is a trivial solution of equation (5.3) (see [5, 18]).

5.2. Numerical results

In this subsection, we present numerical result for the Fokker-Planck equation (1.1) for a confinement vector field \mathbf{F} coupled with the normalization condition $\int_{\Omega} \varphi = \rho$, $\rho \in \mathbb{R}$, and then we apply the algorithm in the framework of fluid mechanics. The main difficulty to obtain a numerical scheme for the Fokker-Planck equation within the normalized condition is to treat this normalized condition

since the equation is not numerically difficult itself. Precisely, this condition is implemented by penalization. For simulation, we use the FreeFem++ program⁽⁷⁾ which is based on weak formulation of the problem and finite elements method.

In the fluid mechanics context, we want to observe the distribution of the orientation dumbbells in a dilute polymer under shear (for instance with a given stationary velocity flow given of the form $\mathbf{u}(x_1, x_2) = (\dot{\gamma} x_2, 0)$, $\dot{\gamma} \in \mathbb{R}$, in the 2-dimensional case). For simplicity, we make the presentation with the 2-dimensional model. According to the previous subsection, the searched distribution satisfies the following Fokker-Planck equation

$$(5.5) \quad -\Delta\varphi + \operatorname{div}(\varphi \mathbf{F}) = 0 \quad \text{on } B(0, \ell),$$

where the vector field \mathbf{F} is given by

$$\mathbf{F} : \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} \in B(0, \ell) \mapsto 2De\dot{\gamma} \begin{pmatrix} Q_2 \\ 0 \end{pmatrix} - \frac{1}{1 - |\mathbf{Q}|^2/\ell^2} \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix}.$$

Moreover, the solution must satisfy the relation $\int_{B(0, \ell)} \varphi = \rho$. Notice that if we have a solution such that $\int_{B(0, \ell)} \varphi = 1$ then, by linearity, the function $\bar{\varphi} = \rho\varphi$ is a solution such that $\int_{B(0, \ell)} \bar{\varphi} = \rho$. In the numerical test, we always take $\rho = 1$. The only two parameters which are interest are the product $De\dot{\gamma}$ and the coefficient ℓ which corresponds to the maximal elongation of the dumbbells. Without shear (that is for $\dot{\gamma} = 0$), a trivial solution of the Fokker-Planck equation (5.5) exists: it is the maxwellian M (see its expression (5.4)). For three characteristic maximal lengths of the dumbbells ($\ell = 2$, $\ell = 5$ and $\ell = 10$), we have been represented this maxwellian on the figure 5.2.




Figure 5.2. Solution without shear for different maximal lengths of dumbbells: $\ell = 2$, $\ell = 5$ and $\ell = 10$.

To observe the influence of the shear on the distribution, taking $De = 10$, $\ell = 5$ and different values of the shear coefficient $\dot{\gamma} \in \{0.1; 0.2; 0.5; 1\}$. The for results are descibe on figure 5.3.




Figure 5.3. Shear influence on the distribution of the length of dumbbells. The more the shear is raised, the more the dumbbells tend to lengthen in the direction of the flow. The four figures above correspond (from left to right) to the values $\dot{\gamma} = 0.1$, $\dot{\gamma} = 0.2$, $\dot{\gamma} = 0.5$ and $\dot{\gamma} = 1$.

⁽⁷⁾ see <http://www.freefem.org/ff++>

BIBLIOGRAPHY

- [1] A. ARNOLD, P. MARKOWICH, G. TOSCANI & A. UNTERREITER, “On convex Sobolev Inequalities and the rate of convergence to equilibrium for Fokker-Planck type equations”, *Communications in Partial Differential Equations* **26** (2001), no. 1, p. 43-100.
- [2] F. BOYER, “Trace theorems and spatial continuity properties for the solutions of the transport equation”, *Differential Integral Equations* **18** (2005), no. 8, p. 891-934.
- [3] F. BOYER & P. FABRIE, *Eléments d’analyse pour l’étude de quelques modèles d’écoulements de fluides visqueux incompressibles*, Mathématiques & Applications (Berlin) [Mathematics & Applications], vol. 52, Springer-Verlag, Berlin, 2006, xii+398 pages.
- [4] H. J. BRASCAMP & E. H. LIEB, “On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation”, *J. Functional Analysis* **22** (1976), no. 4, p. 366-389.
- [5] L. CHUPIN, “The FENE model for viscoelastic thin film flows: Justification of new models and applications.”, *Submitted* (2008).
- [6] P. DEGOND, M. LEMOU & M. PICASSO, “Constitutive relations for viscoelastic fluid models derived from kinetic theory”, in *Dispersive transport equations and multiscale models (Minneapolis, MN, 2000)*, IMA Vol. Math. Appl., vol. 136, Springer, New York, 2004, p. 77-89.
- [7] J. DRONIOU, “Non-coercive linear elliptic problems”, *Potential Anal.* **17** (2002), no. 2, p. 181-203.
- [8] J. DRONIOU & J.-L. VAZQUEZ, “Noncoercive convection-diffusion elliptic problems with Neumann boundary conditions”, *Calc. Var.* **34** (2009), no. 4, p. 413-434.
- [9] D. ESCANDE & F. SATTIN, “When Can the Fokker-Planck Equation Describe Anomalous or Chaotic Transport?”, *Physical Review Letters* **99** (2007), no. 18, p. 185005.
- [10] I. GHOSH, G. MCKINLEY, R. BROWN & R. ARMSTRONG, “Deficiencies of FENE dumbbell models in describing the rapid stretching of dilute polymer solutions”, *Journal of Rheology* **45** (2001), p. 721.
- [11] J. GUÍÑEZ & A. D. RUEDA, “Steady states for a Fokker-Planck equation on S_n ”, *Acta Math. Hungar.* **94** (2002), no. 3, p. 211-221.
- [12] J. HEINONEN, T. KILPELÄINEN & O. MARTIO, *Nonlinear potential theory of degenerate elliptic equations*, Dover Publications Inc., Mineola, NY, 2006, Unabridged republication of the 1993 original, xii+404 pages.
- [13] B. HELFFER & F. NIER, *Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians*, Lecture Notes in Mathematics, vol. 1862, Springer-Verlag, Berlin, 2005, x+209 pages.
- [14] F. HÉRAU & F. NIER, “Isotropic hypoellipticity and trend to equilibrium for the Fokker-Planck equation with a high-degree potential”, *Arch. Ration. Mech. Anal.* **171** (2004), no. 2, p. 151-218.
- [15] B. JOURDAIN, T. LELIÈVRE & C. LE BRIS, “Existence of solution for a micro-macro model of polymeric fluid: the FENE model”, *J. Funct. Anal.* **209** (2004), no. 1, p. 162-193.
- [16] J. LERAY & J. SCHAUDER, “Topologie et équations fonctionnelles”, *Ann. Sci. Éc. Norm. Supér., III. Ser.* **51** (1934), p. 45-78.
- [17] A. LOZINSKI & C. CHAUVIÈRE, “A fast solver for Fokker-Planck equation applied to viscoelastic flows calculations: 2D FENE model”, *J. Comput. Phys.* **189** (2003), no. 2, p. 607-625.
- [18] N. MASMOUDI, “Well posedness for the FENE dumbbell model of polymeric flows”, *Preprint* (2007).
- [19] G. MÉTIVIER, “Comportement asymptotique des valeurs propres d’opérateurs elliptiques dégénérés”, in *Journées: Équations aux Dérivées Partielles de Rennes (1975)*, Soc. Math. France, Paris, 1976, p. 215-249. Astérisque, No. 34-35.
- [20] A. I. NOAROV, “Generalized solvability of the stationary Fokker-Planck equation”, *Differ. Uravn.* **43** (2007), no. 6, p. 813-819, 863.
- [21] H. C. ÖTTINGER, *Stochastic processes in polymeric fluids*, Springer-Verlag, Berlin, 1996, Tools and examples for developing simulation algorithms, xxiv+362 pages.
- [22] F. SATTIN, “Fick’s law and Fokker-Planck equation in inhomogeneous environments”, *Physics Letters A* (2008).
- [23] E. W. STREDULINSKY, *Weighted inequalities and degenerate elliptic partial differential equations*, Lecture Notes in Mathematics, vol. 1074, Springer-Verlag, Berlin, 1984, iv+143 pages.
- [24] H. TRIEBEL, “Interpolation Theory, Function Spaces, Differential Operators. Joh”, *Ambrosius Barth Publ* (1995).
- [25] B. O. TURESSON, *Nonlinear potential theory and weighted Sobolev spaces*, Lecture Notes in Mathematics, vol. 1736, Springer-Verlag, Berlin, 2000, xiv+173 pages.
- [26] E. C. ZEEMAN, “Stability of dynamical systems”, *Nonlinearity* **1** (1988), no. 1, p. 115-155.

Laurent CHUPIN
 Université de Lyon
 INSA de Lyon - Pôle de Mathématiques
 CNRS, UMR5208, Institut Camille Jordan
 21 av. Jean Capelle, 69621 Villeurbanne Cedex, France
 laurent.chupin@insa-lyon.fr