



HAL
open science

Semiclassical resolvent estimates in chaotic scattering

Stéphane Nonnenmacher, Maciej Zworski

► **To cite this version:**

Stéphane Nonnenmacher, Maciej Zworski. Semiclassical resolvent estimates in chaotic scattering. 2009. hal-00376771v1

HAL Id: hal-00376771

<https://hal.science/hal-00376771v1>

Preprint submitted on 20 Apr 2009 (v1), last revised 11 Sep 2009 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SEMICLASSICAL RESOLVENT ESTIMATES IN CHAOTIC SCATTERING

STÉPHANE NONNENMACHER AND MACIEJ ZWORSKI

1. STATEMENT OF RESULTS

In this short note which is a companion to our recent paper [17] we prove a resolvent estimate in the pole free strip for certain Schrödinger-type operators whose classical Hamiltonian flows are assumed to be hyperbolic on the trapped set which is assumed to be sufficiently filamentary – see (1.4) for the precise condition. We will refer to [17, §3] for the preliminary material and assumptions on the operator.

The example to keep in mind is the operator

$$(1.1) \quad Pu = P(h)u = -h^2 \frac{1}{\sqrt{\bar{g}}} \sum_{i,j=1}^n \partial_{x_j} (\sqrt{\bar{g}} g^{ij} \partial_{x_i} u) + V(x), \quad x \in \mathbb{R}^n,$$

$G(x) \stackrel{\text{def}}{=} (g^{ij}(x))_{i,j}$ is a symmetric positive definite matrix, $\bar{g} \stackrel{\text{def}}{=} 1/\det G(x)$, and

$$g^{ij}(x) = \delta_{ij}, \quad V(x) = -1, \quad |x| > R.$$

We refer to [17, §3.2] for the complete set of assumptions which allow long range perturbations, at the expense of some analyticity assumptions standard for the definition of resonances – see [21] and references given there. We note that for $g^{ij} \equiv 1$ we obtain a class of semiclassical Schrödinger operators and for $V \equiv -1$, the Helmholtz equation for a Laplace-Beltrami operator, with $h = 1/\lambda$, playing the rôle of wavelength. The euclidean space can be without any changes replaced by a smooth manifold coinciding with $(\mathbb{R}^n \setminus B(0, R)) \sqcup \dots \sqcup (\mathbb{R}^n \setminus B(0, R))$ outside of a compact set.

The key assumption is that the flow is hyperbolic on the trapped set at energy 0, K :

$$(1.2) \quad K \stackrel{\text{def}}{=} \{(x, \xi) : p(x, \xi) = 0, \exp tH_p(x, \xi) \not\rightarrow \infty, t \rightarrow \pm\infty\},$$

$$p(x, \xi) \stackrel{\text{def}}{=} \sum_{i,j=1}^n g^{ij}(x) \xi_i \xi_j + V(x), \quad H_p(x, \xi) \stackrel{\text{def}}{=} \sum_{k=1}^n \partial_{\xi_k} p \partial_{x_k} - \partial_{x_k} p \partial_{\xi_k},$$

$\exp(tH_p)|_K$ is hyperbolic in the sense of [17, §3.3], $dp|_{p^{-1}(0)} \neq 0$.

The Hamiltonian flow defined by H_p corresponds to the geodesic flow on $S^*\mathbb{R}^n$ when $V = -1$, and the hyperbolicity assumption means that for $\rho \in K$, we have a flow invariant

decomposition,

$$T_\rho p^{-1}(0) = \mathbb{R}H_p(\rho) \oplus E_\rho^- \oplus E_\rho^+,$$

into neutral ($\mathbb{R}H_p(\rho)$), stable (E_ρ^-), and unstable (E_ρ^+) directions – see [17, (3.11)] for the full description.

The condition that the trapped set is “filamentary” is formulated in terms of a topological pressure. We refer to [17, §3.3] and texts on dynamical systems [13],[25] for precise definitions, recalling only the definition in the special, but often satisfied, case in which the periodic orbits are dense in K . Let $f \in C^0(K)$. Then the *pressure* of f with respect to the Hamiltonian flow on K is given by

$$(1.3) \quad \mathcal{P}(f) \stackrel{\text{def}}{=} \lim_{T \rightarrow \infty} \frac{1}{T} \log \sum_{\gamma: T_\gamma < T} \exp \int_0^{T_\gamma} (\exp tH_p)^* f dt,$$

where the sum runs over all periodic orbits γ of periods $T_\gamma \leq T$.

We can now formulate our result:

Theorem. *Suppose that $P(h)$ satisfies (1.1) or the more general assumptions of [17, §3.2]. Suppose also that the Hamiltonian flow (1.2) is hyperbolic on the trapped set K , and that*

$$(1.4) \quad \mathcal{P}(-\varphi_+/2) < 0, \quad \varphi_+(\rho) \stackrel{\text{def}}{=} \frac{d}{dt} (d \exp tH_p|_{E^+(\rho)})|_{t=0}, \quad \rho \in K.$$

Then for any $\chi \in C_c^\infty(\mathbb{R}^n)$ and $\epsilon > 0$, there exist $\delta(\epsilon) > 0$ and $h(\epsilon) > 0$ such that the cut-off resolvent $\chi(P(h) - z)^{-1}\chi$, $\text{Im } z > 0$, continues analytically to

$$\Omega_\epsilon(h) \stackrel{\text{def}}{=} \{z : \text{Im } z > h(\mathcal{P}(-\varphi_+/2) + \epsilon), \quad |\text{Re } z| < \delta(\epsilon)\}, \quad 0 < h < h(\epsilon).$$

For $z \in \Omega_\epsilon(h) \cap \{\text{Im } z \leq 0\}$, this resolvent is polynomially bounded in h :

$$(1.5) \quad \|\chi(P(h) - z)^{-1}\chi\|_{L^2 \rightarrow L^2} \leq C(\epsilon, \chi) h^{-1+c_E \text{Im } z/h} \log(1/h),$$

$$c_E \stackrel{\text{def}}{=} \frac{n}{2|P(-\varphi_+/2) + \epsilon/2|}.$$

We used the continuity of the pressure with respect to the energy in order to simplify the statement – see [17, Theorem 3] for a slightly more precise formulation. In dimension $n = 2$ the condition (1.4) is equivalent to the statement that the Hausdorff dimension of $K \subset p^{-1}(0)$ is less than 2. Since the energy surface $p^{-1}(0)$ has dimension 3 and the minimal dimension of a non-empty K is 1, the condition means that we are less than “half-way” and K is *filamentary*. Trapped sets with dimensions greater than 2 are referred to as *bulky*.

The first part of the theorem is the main result of [17], see Theorem 3 there. Here we use the techniques developed in that paper to prove (1.5). For $-\Delta$ outside several convex obstacles (satisfying a condition guaranteeing strict hyperbolicity of the flow) with Dirichlet or Neumann boundary condition, the theorem was proved by Ikawa [12], with the pressure

being only implicit in the statement which gave an explicit condition on distances and sizes of the obstacles. For more recent developments in that setting see [2],[18], and [15].

In particular, for z on the real axis the bound (1.5) gives

$$(1.6) \quad \|\chi(P(h) - z)^{-1}\chi\|_{L^2 \rightarrow L^2} \leq C \frac{\log(1/h)}{h}, \quad z \in [-\delta(\epsilon), \delta(\epsilon)], \quad 0 < h < h(\epsilon).$$

This result was already given in [17, Theorem 5] with a less direct proof. It has been generalized to a larger class of manifolds in [7] and (1.5) provides no new insight in that setting. One of the applications is local smoothing with a minimal loss [5] (see [3] for the original application in the setting of obstacle scattering) and a recent no loss Strichartz estimate [4] under the assumption (1.4). The advantage of having (1.5) lies in applications to the energy decay for the wave equation – see [3, 6, 10] and references given there. That is particularly important in view of the recent results of Schenck on the use of pressure estimates in the setting of the damped wave equation [20].

To prove (1.5) we show in §3 that the estimates obtained in [17, §7] can be used to obtain a good parametrix for the complex-scaled operator, which leads to an estimate for the inverse. As was pointed out to us by Burq the construction of the parametrix for the outgoing resolvent was the, somewhat implicit, key step in the work of Ikawa [12] on the resonance gap for several convex obstacle. That insight lead us to re-examine the consequences of [17].

We follow the notation of [17] with precise references given as we go along. For the needed aspects of semiclassical microlocal analysis [17, §3] and the references to [8] and [9] should be consulted.

ACKNOWLEDGEMENTS. The first author was partially supported by the Agence Nationale de la Recherche under the grant ANR-05-JCJC-0107-01, and the second author by the National Science Foundation under the grant DMS 0654436. In addition to Nicolas Burq, we would like to thank Nalini Anantharaman and Jared Wunsch for helpful discussions related to [17].

2. REVIEW OF THE HYPERBOLIC DISPERSION ESTIMATES

The dispersion estimate we need is proved in [17] for a modified operator which we will now describe.

The first modification of $P(h)$ comes from the method of complex scaling reviewed in [17, §3.4]. For any fixed sufficiently large $R_0 > 0$, it results in the operator $P_\theta(h)$, with the following properties. To formulate them put

$$(2.1) \quad \Omega_\theta \stackrel{\text{def}}{=} [-\delta, \delta] + i[-\theta/C, C], \quad \theta = M_1 h \log(1/h).$$

Then

$$(2.2) \quad P_\theta(h) - z : H_h^2(\mathbb{R}^n) \longrightarrow L^2(\mathbb{R}^n) \text{ is a Fredholm operator for } z \in \Omega_\theta,$$

$$(2.3) \quad \forall \chi \in \mathcal{C}_c^\infty(B(0, R_0)), \quad \chi R(z, h)\chi = \chi R_\theta(z, h)\chi.$$

Here and below we set

$$R_\bullet(z, h) \stackrel{\text{def}}{=} (P_\bullet(h) - z)^{-1}, \quad \text{Im } z > 0,$$

and (2.3) shows the meromorphic continuation of $\chi R(z, h)\chi$ to Ω_θ , the meromorphy being guaranteed by the Fredholm property of $P_\theta(h) - z$.

The operator $P_\theta(h)$ is further modified by an exponential weight, $G^w = G^w(x, hD)$,

$$G \in \mathcal{C}_c^\infty(T^*\mathbb{R}^n), \quad \text{supp } G \subset p^{-1}((-2\delta, 2\delta)), \quad \partial^\alpha G = \mathcal{O}(h \log(1/h)),$$

where $\delta > 0$ is a fixed number:

$$(2.4) \quad P_{\theta, \epsilon}(h) \stackrel{\text{def}}{=} e^{-\epsilon G^w/h} P_\theta(h) e^{\epsilon G^w/h}, \quad \epsilon = M_2 \theta, \quad \theta = M_1 h \log(1/h).$$

The new operator has the same spectrum as $P_\theta(h)$ and the following properties:

$$(2.5) \quad \text{Im } \psi_0^w(x, hD) P_{\theta, \epsilon}(h) \psi_0^w(x, hD) \leq C h$$

$$\text{if } \text{supp } \psi_0 \subset p^{-1}((-3\delta/2, 3\delta/2)), \quad \psi_0 \in S(1),$$

$$(2.6) \quad \|\exp(\pm \epsilon G^w/h)\|_{L^2 \rightarrow L^2} = \mathcal{O}(h^{-CM_2}).$$

The main reason for introducing the weight G is to ensure the bound (2.5). The specific choice of G is explained in [17, §6.1]. In particular G vanishes in some neighbourhood of the trapped set K , and

$$\exp(tG^w(x, hD)) = B_t^w(x, hD), \quad B_t \in h^{-N} S_\delta(T^*\mathbb{R}^n), \quad B_t|_{\mathfrak{C}_{\text{supp } G}} = 1 + \mathcal{O}_{S_\delta}(h^\infty).$$

Hence, if the spatial cutoff χ is supported away from $\pi \text{supp } G$, calculus of semiclassical pseudodifferential operators ensures that

$$(2.7) \quad \chi R(z, h)\chi = \chi R_{\theta, \epsilon}(z, h)\chi + \mathcal{O}_{L^2 \rightarrow L^2}(h^\infty) \|R_{\theta, \epsilon}(z, h)\|,$$

so our objective is to estimate $\|R_{\theta, \epsilon}(z, h)\|_{L^2 \rightarrow L^2}$.

We consider a final modification of $P(h)$ near the energy surface. Let $\psi_0 \in S(1)$ be supported in $p^{-1}((-3\delta/2, 3\delta/2))$ and equal to 1 in $p^{-1}(-\delta, \delta)$. Define

$$(2.8) \quad \tilde{P}_{\theta, \epsilon}(h) \stackrel{\text{def}}{=} \psi_0^w(x, hD) P_{\theta, \epsilon} \psi_0^w(x, hD),$$

and the modified propagator [17, (6.9)]:

$$(2.9) \quad U(t) \stackrel{\text{def}}{=} \exp\{-it\tilde{P}_{\theta, \epsilon}(h)/h\}.$$

The crucial ingredients in proving (2.13) are the good upper bounds for the norms

$$\|U(t)\psi^w(x, hD)\|_{L^2 \rightarrow L^2}, \quad 0 \leq t \leq M \log(1/h),$$

where $M > 0$ is fixed but large, and

$$(2.10) \quad \psi \in S(1), \quad \text{supp } \psi \subset p^{-1}((-\delta/2, \delta/2)), \quad \psi = 1 \text{ on } p^{-1}((-\delta/4, \delta/4)).$$

From the bound (2.5) on the imaginary part of the modified Hamiltonian, we get an exponential control on the propagator:

$$(2.11) \quad \|U(t)\|_{L^2 \rightarrow L^2} \leq \exp(Ct), \quad t \geq 0.$$

The reason to conjugate P_θ with the weight G^w was indeed to ensure this exponential bound. Together with the hyperbolic bound (2.13), this exponential bound would suffice to get a polynomial bound $\mathcal{O}(h^{-L})$ in (1.5), for some (unknown) $L > 0$. To obtain the explicit value,

$$-1 + \frac{c_E \operatorname{Im} z}{h},$$

for the exponent, we need to improve (2.11) into the following uniform bound:

Lemma 2.1. *Let ψ satisfy the conditions (2.10). Then, there exists a constant $C_0 > 0$ such that, for any small enough h and any time $0 \leq t \leq M \log(1/h)$,*

$$(2.12) \quad \|U(t)\psi^w(x, hD)\|_{L^2 \rightarrow L^2} \leq C_0.$$

Before proving this Lemma, we state the major consequence of the *hyperbolic dispersion estimate* [17, Proposition 6.3], in the spirit of [17, §6.4] (see also [17, Proposition 9.1] for a simpler case). As above, we take ψ as in (2.10). For any $\epsilon > 0$ and $0 < h < h(\epsilon)$, we then have

$$(2.13) \quad \begin{aligned} \|U(t)\psi^w(x, hD)\|_{L^2 \rightarrow L^2} &\leq C h^{-n/2} \exp(-\lambda t) + \mathcal{O}(h^{M_3}), \\ \lambda &\stackrel{\text{def}}{=} -\mathcal{P}(\varphi_+/2) + \epsilon/2, \end{aligned}$$

uniformly in the time range

$$0 < t < M \log(1/h).$$

We have used the notation of (1.4), M is arbitrarily large, and M_3 can be taken as large as we wish, provided we choose M_1 in (2.1) large enough depending on M .

If the pressure $\mathcal{P}(\varphi_+/2)$ is negative, one can take ϵ small enough to ensure $\lambda > \epsilon/2 > 0$. The above estimate is then sharper than (2.12) for times beyond the *Ehrenfest time*

$$(2.14) \quad t_E \stackrel{\text{def}}{=} c_E \log(1/h), \quad c_E \stackrel{\text{def}}{=} \frac{n}{2\lambda}.$$

The large constant M will always be chosen (much) larger than c_E .

Proof of Lemma 2.12. To motivate the proof we start with a heuristic argument for the bound (2.12). As mentioned above, the exponential bound (2.11) is due to the fact that the imaginary part of the operator $\tilde{P}_{\theta, \epsilon}(h)$ can take positive values of order $\mathcal{O}(h)$ (see (2.5)). However, the construction of the weight G shows that outside a bounded region of phase space of the form

$$V_{\text{pos}} = p^{-1}((-2\delta, 2\delta)) \cap T_{\{R_1 < |x| < R_2\}}^* \mathbb{R}^n,$$

the imaginary part of $\tilde{P}_{\theta, \epsilon}(h)$ is negative up to $\mathcal{O}(h^\infty)$ errors.

The radius R_1 above is large enough, so that V_{pos} lies *finite distance* from the trapped set. As a result, any trajectory crossing the region V_{pos} will only spend a bounded time in that region. For this reason, the propagator $U(t)$ on a large time $t \gg 1$ will “accumulate” exponential growth during a uniformly bounded time only.

We now provide a rigorous proof, using ideas and results from [17, §6.3]. The phase space $T^*\mathbb{R}^n$ is split using a smooth partition of unity:

$$1 = \sum_{b=0,1,2,\infty} \pi_b, \quad \pi_b \in C^\infty(T^*\mathbb{R}^n, [0, 1]).$$

These four functions have specific localization properties:

- $\text{supp } \pi_b \subset p^{-1}((-\delta, \delta))$ for $b = 0, 1, 2$
- π_∞ is localized outside $p^{-1}((-3\delta/4, 3\delta/4))$
- π_1 is supported near K , in particular, its support does not intersect V_{pos}
- π_2 is supported away from K but inside $\{|x| < R_2 + 1\}$
- π_0 is supported near spatial infinity, that is on $\{|x| > R_2 - 1\}$ where the operator $\tilde{P}_{\theta,\epsilon}(h)$ is *absorbing* (the imaginary part of its symbol is negative).

Employing a positive (Wick) quantization scheme (see for instance [14], and for the semiclassical setting [19, §3.3]), $\Pi_b = \text{Op}_h^+(\pi_b)$, we produce a quantum partition of unity

$$\text{Id} = \sum_{b=0,1,2,\infty} \Pi_b, \quad \|\Pi_b\| \leq 1.$$

It was the positive quantization that ensured $\|\Pi_b\| \leq 1$ above.

The evolution $U(t)$ is then split between time intervals of length t_0 , where $t_0 > 0$ is large but independent of h . Using the partition of unity, we decompose the propagator at time $t = Nt_0$ into

$$U(Nt_0) \psi^w(x, hD) = \left(\sum_{b=0,1,2,\infty} U_b \right)^N \psi^w(x, hD), \quad \text{where } U_b \stackrel{\text{def}}{=} U(t_0) \Pi_b.$$

Expanding the power, we obtain a sum of terms $U_{b_N} \cdots U_{b_1} \psi^w$ and we will see that many of these terms are negligible.

Taking into account the energy cutoff $\psi \in C^\infty(p^{-1}((-\delta/2, \delta/2)))$ and assuming $N \leq C \log(1/h)$, it is shown in [17, Lemma 6.5] that any term containing at least one factor U_∞ (localized outside the energy shell) is $\mathcal{O}_{L^2 \rightarrow L^2}(h^\infty)$.

Using the fact that any classical trajectory can travel in $\text{supp } \pi_2$ at most for a finite time $\leq N_0 t_0$, [17, Lemma 6.6] shows that the *relevant* sequences $b_1 \cdots b_N$ are of the form

$$b_i = 1 \quad \text{for } N_0 < i < N - N_0.$$

They correspond to trajectories spending most of the time near K . Then [17, Lemma 6.5] shows that

$$U(Nt_0) \psi^w(x, hD) = U(N_0 t_0) (U_1)^{N-2N_0} U(N_0 t_0) \psi^w(x, hD) + \mathcal{O}_{L^2 \rightarrow L^2}(h^{M_5}),$$

uniformly for any $2N_0 \leq N < M \log(1/h)$, where $M_5 > 0$.

Finally, [17, Lemma 6.3] shows that

$$U_1 = U(t_0)\Pi_1 = U_0(t_0)\Pi_1 + \mathcal{O}_{L^2 \rightarrow L^2}(h^\infty),$$

where $U_0(t_0) = \exp(-it_0 P(h)/h)$ is *unitary*. Hence, $\|U_1\| \leq 1 + \mathcal{O}(h^\infty)$, while $\|U(N_0 t_0)\|$ is estimated using (2.11). \square

3. RESOLVENT ESTIMATES

We can now prove the resolvent estimate (1.5) by constructing a parametrix for $P_{\theta,\epsilon}(h) - z$, $z \in \Omega_\epsilon(h)$ defined in the statement of the theorem. We will use the notation

$$\zeta \stackrel{\text{def}}{=} z/h$$

to shorten some of the formulæ. We want to find an approximate solution to

$$(P_{\theta,\epsilon}(h) - z)u = f, \quad f \in L^2(\mathbb{R}^n), \quad z \in \Omega_\epsilon(h).$$

First, the ellipticity away from the energy surface $p^{-1}(0)$ shows that, for ψ as in (2.10), there exists an operator, $T_0 = \mathcal{O}(1) : L^2(\mathbb{R}^n) \rightarrow H_h^2(\mathbb{R}^n)$, such that

$$(P_{\theta,\epsilon}(h) - z)T_0 f = (1 - \psi^w(x, hD))f + R_0 f, \quad R_0 = \mathcal{O}_{L^2 \rightarrow L^2}(h^\infty).$$

To treat the vicinity of $p^{-1}(0)$ we put

$$T_1 f = (i/h) \int_0^{t_M} dt e^{i\zeta t} U(t) \psi^w(x, hD) f, \quad t_M = M \log(1/h),$$

which satisfies

$$(3.1) \quad (\tilde{P}_{\theta,\epsilon}(h) - z) T_1 f = \psi^w(x, hD) f + R_1 f, \quad R_1 \stackrel{\text{def}}{=} -e^{i\zeta t_M} U(t_M) \psi^w(x, hD).$$

The estimate (2.13) shows that, if $\lambda + \text{Im } \zeta > \epsilon/2$, and for arbitrary $M_4 > 0$, one can choose M and M_3 large enough such that $R_1 = \mathcal{O}_{L^2 \rightarrow L^2}(h^{M_4})$. We can estimate the norm of T_1 by the triangle inequality,

$$(3.2) \quad \|T_1\|_{L^2 \rightarrow L^2} \leq h^{-1} \int_0^{t_M} e^{-\text{Im } \zeta t} \|U(t) \psi^w(x, hD)\|_{L^2 \rightarrow L^2} dt,$$

and then use the bounds (2.12) for times $0 \leq t \leq t_E$ and (2.13) for times $t_E < t \leq t_M$.

When $\text{Im } \zeta = 0$, the above integral can be estimated by the integral over the interval $t \in [0, t_E]$:

$$\text{Im } \zeta = 0 \implies \|T_1\|_{L^2 \rightarrow L^2} \leq h^{-1} \left(C_0 t_E + \frac{1}{\lambda} \right) \leq C h^{-1} \log h^{-1}.$$

In the case $0 > \text{Im } \zeta > -\lambda + \epsilon/2$, the dominant part of the integral comes from $t = t_E$:

$$0 > \text{Im } \zeta > -\lambda + \epsilon/2 \implies \|T_1\|_{L^2 \rightarrow L^2} \leq C_\epsilon h^{-1} e^{-\text{Im } \zeta t_E} = C_\epsilon h^{-1 + c_E \text{Im } \zeta}.$$

Now, (3.1) reads

$$\psi_0^w(x, hD)(P_{\theta, \epsilon}(h) - z)\psi_0^w(x, hD)T_1 f = \psi^w(x, hD)f + R_1 f.$$

From the inclusion $\psi_0|_{\text{supp } \psi} \equiv 1$, one can show (as in [17, Lemma 6.5]) that

$$\psi_0^w(x, hD)(P_{\theta, \epsilon}(h) - z)\psi_0^w(x, hD)T_1 = (P_{\theta, \epsilon}(h) - z)T_1 + R_2, \quad R_2 = \mathcal{O}_{L^2 \rightarrow L^2}(h^\infty),$$

and also that

$$\|T_1\|_{H_h^2} \leq C \|T_1\|_{L^2}.$$

Putting $T = T_0 + T_1$ and $R = R_0 + R_1 + R_2$, we obtain

$$(P_{\theta, \epsilon}(h) - z)T = \text{Id} + R, \quad R = \mathcal{O}_{L^2 \rightarrow L^2}(h^{M_4}).$$

This means that $(P_{\theta, \epsilon}(h) - z)$ can be inverted, with

$$\|(P_{\theta, \epsilon}(h) - z)^{-1}\|_{L^2 \rightarrow H_h^2} = (1 + \mathcal{O}(h^{M_4}))\|T\|_{L^2 \rightarrow H_h^2}.$$

The above estimates on the norms of T_0 and T_1 can be summarized by

$$(3.3) \quad 0 \geq \text{Im } \zeta \geq \epsilon + \mathcal{P}(-\varphi_+/2) \implies \|T\|_{L^2 \rightarrow H_h^2} \leq C_\epsilon h^{-1+c_E \text{Im } \zeta} \log h^{-1}.$$

Using (2.7), this proves the bound (1.5). \square

Remark. By using a sharper energy cutoff ψ_h belonging to an exotic symbol class (see [23, §4]) and supported in the energy layer $p^{-1}((-h^{1-\delta}, h^{1-\delta}))$ (as in [1]), the bound (2.13) is likely to be improved to

$$(3.4) \quad \|U(t)\psi_h^w(x, hD)\| \leq Ch^{-(n-1+\delta)/2} \exp(-\lambda t) + \mathcal{O}(h^{M_3}).$$

This bound becomes sharper than (2.12) around the time $t'_E = c'_E \log(1/h)$, where

$$c'_E \stackrel{\text{def}}{=} \frac{n-1+\delta}{2\lambda} < c_E.$$

As a result, the bounds on the norm of the corresponding operator T'_1 are modified accordingly. At the same time, as shown in [1, Prop. 5.4], the ellipticity away from the energy surface provides an operator T'_0 satisfying

$$(P_{\theta, \epsilon}(h) - z)T'_0 = (1 - \psi_h^w(x, hD)) + \mathcal{O}_{L^2 \rightarrow L^2}(h^\infty),$$

and of norm $\|T'_0\|_{L^2 \rightarrow H_h^2} = \mathcal{O}(h^{-1+\delta})$. The norm of $T' = T'_0 + T'_1$ is still dominated by that of T'_1 , so that we eventually get

$$\|\chi(P(h) - z)^{-1}\chi\|_{L^2 \rightarrow H_h^2} \leq C_\epsilon h^{-1+c'_E \text{Im } z/h} \log(1/h), \quad z \in \Omega_\epsilon(h) \cap \{\text{Im } z \leq 0\}.$$

Since it is not clear that even this bound is optimal, and that proving (3.4) would require some effort, we have limited ourselves to using the established bound (2.13).

One advantage of the approach presented in this note (compared with the method of [17, §9]) is that, to obtain the bound (1.6) we did not have to use the complex interpolation arguments of [3] and [24].

REFERENCES

- [1] N. Anantharaman and S. Nonnenmacher, *Half-delocalization of eigenfunctions of the laplacian*, Ann. Inst. Fourier. **57**, 2465–2523 (2007)
- [2] N. Burq, *Contrôle de l'équation des plaques en présence d'obstacle strictement convexes*. Mémoires de la Société Mathématique de France, Sér. 2, **55** (1993), 3-126.
- [3] N. Burq, *Smoothing effect for Schrödinger boundary value problems*, Duke Math. J. **123**(2004), 403–427.
- [4] N. Burq, C. Guillarmou, and A. Hassel, *Strichartz estimates without loss in hyperbolic trapping situations*. in preparation.
- [5] H. Christianson, *Cutoff resolvent estimates and the semilinear Schrödinger equation* Proc. Amer. Math. Soc. **136**(2008), 3513–3520.
- [6] H. Christianson, *Applications of cutoff Resolvent estimates to the wave equation*, preprint.
- [7] K. Datchev, *Local smoothing for scattering manifolds with hyperbolic trapped sets*, Comm. Math. Phys. **286**(2009), 837-850.
- [8] M. Dimassi and J. Sjöstrand, *Spectral Asymptotics in the semi-classical limit*, Cambridge University Press, 1999.
- [9] L.C. Evans and M. Zworski, *Lectures on Semiclassical Analysis*, <http://math.berkeley.edu/~zworski/seminclassical.pdf>
- [10] C. Guillarmou and F. Naud, *Wave decay on convex co-compact hyperbolic manifolds*, Commun. Math. Phys. **287**, 489–511 (2009)
- [11] L. Hörmander, *The Analysis of Linear Partial Differential Operators*, Vol. I, II, Springer-Verlag, Berlin, 1983.
- [12] M. Ikawa, *Decay of solutions of the wave equation in the exterior of several convex bodies*, Ann. Inst. Fourier, **38**(1988), 113-146.
- [13] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems*, Cambridge University Press, 1997.
- [14] N. Lerner, *The Wick calculus of pseudodifferential operators and some of its applications*, Cubo Mat. Edu. **5**(2003), 213–236.
- [15] S. Nonnenmacher, J. Sjöstrand, and M. Zworski, *Hyperbolic quantum monodromy operators*, in preparation.
- [16] S. Nonnenmacher and M. Zworski, *Distribution of resonances for open quantum maps*, Comm. Math. Phys. **269**(2007), 311–365, *Fractal Weyl laws in discrete models of chaotic scattering*, Journal of Physics A, **38** (2005), 10683-10702.
- [17] S. Nonnenmacher and M. Zworski, *Quantum decay rates in chaotic scattering*, Acta Math., to appear.
- [18] V. Petkov and L. Stoyanov, *Analytic continuation of the resolvent of the Laplacian and the dynamical zeta function*, preprint 2007, <http://www.math.u-bordeaux.fr/~petkov/publications/publi2.html>
- [19] K. Pravda-Starov, *Pseudo-spectrum for a class of semi-classical operators*, Bull. Soc. Math. France **136**(2008), 329–372.
- [20] E. Schenck, *Energy decay for the damped wave equation under a pressure condition*, in preparation.
- [21] J. Sjöstrand, *A trace formula and review of some estimates for resonances*, in *Microlocal analysis and spectral theory* (Luca, 1996), 377–437, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 490, Kluwer Acad. Publ., Dordrecht, 1997.
- [22] J. Sjöstrand and M. Zworski, *Complex scaling and the distribution of scattering poles*, Journal of AMS **4**(1991), 729–769
- [23] J. Sjöstrand and M. Zworski, *Asymptotic distribution of resonances for convex obstacles*, Acta Math. **183**(1999), 191-253.

- [24] S.H. Tang and M. Zworski, *From quasimodes to resonances*, Math. Res. Lett. **5**(1998), 261-272.
- [25] P. Walters, *An Introduction to Ergodic Theory*, Springer, Heidelberg, 1982.

INSTITUT DE PHYSIQUE THÉORIQUE, CEA/DSM/PhT, UNITÉ DE RECHERCHE ASSOCIÉ AU CNRS,
CEA/SACLAY,, 91191 GIF-SUR-YVETTE, FRANCE

E-mail address: `nonnen@spht.saclay.cea.fr`

MATHEMATICS DEPARTMENT, UNIVERSITY OF CALIFORNIA, EVANS HALL, BERKELEY, CA 94720,
USA

E-mail address: `zworski@math.berkeley.edu`