



HAL
open science

Pharmacological characterization of (4R)-alkyl glutamate analogues at the ionotropic glutamate receptors - Focus on subtypes iGluR5-7

Lennart Bunch, Thierry Gefflaut, Sébastien Alaux, Emmanuelle Sagot,
Brigitte Nielsen, Darryl S. Pickering

► To cite this version:

Lennart Bunch, Thierry Gefflaut, Sébastien Alaux, Emmanuelle Sagot, Brigitte Nielsen, et al.. Pharmacological characterization of (4R)-alkyl glutamate analogues at the ionotropic glutamate receptors - Focus on subtypes iGluR5-7. *European Journal of Pharmacology*, 2009, 609, pp.1-4. hal-00376953

HAL Id: hal-00376953

<https://hal.science/hal-00376953>

Submitted on 20 Apr 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Pharmacological characterization of (4*R*)-alkyl glutamate analogues at the ionotropic glutamate receptors — Focus on subtypes iGlu₅₋₇

Lennart Bunch^a, Thierry Gefflaut^c, Sebastien Alaux^c, Emanuelle Sagot^c, Birgitte Nielsen^a and Darryl S. Pickering^b

^aDepartment of Medicinal Chemistry, Faculty of Pharmaceutical Sciences, University of Copenhagen, Universitetsparken 2, DK-2100 Copenhagen, Denmark

^bDepartment of Pharmacology and Pharmacotherapy, Faculty of Pharmaceutical Sciences, University of Copenhagen, Universitetsparken 2, DK-2100 Copenhagen, Denmark

^cLaboratoire SEESIB, UMR 6504, Département de Chimie, Université Blaise Pascal, 63177 Aubière Cedex, France

Abstract

The kainic acid (kainate, KA) receptors belong to the class of ionotropic glutamate (iGlu) receptors in the central nervous system. Five subtypes have been identified, which have been termed KA_{1,2} and iGlu₅₋₇. In the search for subtype selective ligands, α -amino-5-*tert*-butyl-3-hydroxy-4-isoxazolyl)propionic acid (ATPA), (4*R*)-methyl Glu (**1a**), and *E*-4-neopentylidene Glu (**2f**) have all previously been reported as selective agonists for the iGlu₅ receptor subtype. In this paper, we present the pharmacological evaluation of a five-compound series of (4*R*)-alkyl Glu analogs (**1b–e,g**) which may be envisaged as conformationally released designs of ATPA and 4-alkylidenes **2a–h**. Most notable is the pharmacological profile for (4*R*)-isopentyl Glu (**1g**) which shows a 10-fold increase in binding affinity for the iGlu₅ receptor subtype ($K_i = 20.5$ nM) in comparison with its *E*-4-alkylidene structural isomer **2g**. Furthermore, **1g** displays high selectivity over other KA receptor subtypes (KA_{1,2} and iGlu_{6,7}), AMPA-, and NMDA receptors (2050 and > 5000 fold, respectively).

Keywords: Glutamate receptor; Kainate receptor; Subtype selectivity; iGlu₅; iGlu₆; iGlu₇

1. Introduction

(*S*)-Glutamic acid (glutamate, Glu) is the major excitatory neurotransmitter in the central nervous system (CNS) activating the plethora of ionotropic Glu (iGlu) receptors and metabotropic Glu (mGlu) receptors ([Bräuner-Osborne et al., 2000] and [Meldrum, 2000]). Whereas the iGlu receptors are ion channels and thus mediate a fast excitatory response (Na⁺, K⁺, Ca⁺⁺ flux), the mGlu receptors are classified as G-protein coupled receptors and generate a slower signal transduction through second messenger systems. On the basis of pharmacological studies, the iGlu receptors are further divided into: 2-amino-3-(3-hydroxy-5-methyl-4-isoxazolyl)propionic acid (AMPA) receptors (homo- or heteromeric receptors comprising the subunits iGlu₁₋₄), kainic acid (KA) receptors (homo- or heteromeric receptors comprising the subunits iGlu₅₋₇ and KA_{1,2}) (Wisden and Seeburg, 1993), and the *N*-methyl-d-aspartic acid (NMDA) receptors (heteromeric receptors comprising the subunits NR₁, NR_{2A-D}, NR_{3A-C}). The mGlu receptors comprise eight homodimeric subtypes, mGlu₁₋₈, which are grouped with respect to the second messenger system activated, pharmacology, and molecular biology (group I: mGlu_{1,5} receptors; group II: mGlu_{2,3} receptors; group III: mGlu_{4,6-8} receptors) (Ferraguti and Shigemoto, 2006). Termination of the excitatory signal is controlled by uptake of Glu from the synaptic cleft by the excitatory amino acid transporters (EAATs) (Beart and O'Shea, 2007). To date, five subtypes have been identified of which EAAT₁₋₃ are high capacity Glu transporting proteins, EAAT₄ functions predominantly as a chloride ion channel and EAAT₅ is present exclusively in the retina.

In order to study the function of a single receptor subtype, when found in its intrinsic biological environment, the application of subtype selective ligands (agonists, partial agonists or antagonists) is a useful pharmacological approach. In 1997 the AMPA-analog, α -amino-5-*tert*-butyl-3-hydroxy-4-isoxazolyl)propionic acid (ATPA), was shown to be a highly selective ligand for the iGlu₅ receptor subtype. Concurrently (4*R*)-methyl Glu (**1a**) was found to display preference for KA receptors over AMPA, NMDA, and the mGlu receptors ([BraunerOsborne et al., 1997], [Gu et al., 1995], [Sagot et al.,

2008] and [Zhou et al., 1997]), and in 2001 we reported that *E*-4-neopentylidene Glu (**2f**) (Bunch et al., 2001) is a highly selective iGlu₅ receptor ligand (Table 1). In this paper we present the pharmacological investigation of a five-compound series of (4*R*)-alkyl Glu analogs (**1b–e,g**) which may be viewed as conformationally *released* analogs of **2b–e,g**. Such a distinct – yet confined – change in physical chemical property is interesting as it allows for a detailed investigation of ligand flexibility as opposed to receptor subtype selectivity.

Table 1.

Binding affinities of (4*R*)-alkyl Glu analogs (**1a–e,g**), *E*-4-alkylidene Glu analogs (**2a–h**), AMPA, ATPA, and KA at native iGlu receptors (rat brain synaptosomes) and at cloned rat homomeric iGlu_{5–7} receptor subtypes.

Entry	[³ H]AMPA IC ₅₀ [nM]	[³ H]KA ^a IC ₅₀ [nM]	NMDA K _i [nM]	iGlu ₅ K _i [nM]	iGlu ₆ K _i [nM]	iGlu ₇ K _i [nM]
1a ^b	26,600	32	5900	0.7	17	6
1b	16,000 ± 1000	120 ± 10	28,000 ± 5000	2.08 ± 0.12	75.5 ± 5.7	27.5 ± 9.4
1c	78,000 ± 2000	700 ± 30	>100,000	11.31 ± 0.55	458 ± 29	141.8 ± 0.4
1d	>100,000	1400 ± 100	~75,000 (n = 1)	37.7 ± 0.73	2270 ± 376	383 ± 38
1e	>100,000	5600 ± 200	>100,000	63.5 ± 9.4	4690 ± 434	1596 ± 293
1g	42,000 ± 4000	>100,000	>100,000	20.5 ± 3.4	>100,000	6167 ± 216
2a ^b	150	230	1200	270	450	–
2b ^b	–	–	–	2200	2500	–
2c ^b	–	–	–	61	18,000	–
2d ^b	–	–	–	13,900	>400,000	–
2e ^b	–	–	–	54	83,000	520
2f	2000 ^b	>100,000 ^b	>100,000 ^b	10.4 ± 1.3	>1,000,000	2247 ± 155
2g ^b	–	–	–	224	634,000	–
2h	6900 ^b	>100,000 ^b	>100,000 ^b	122 ± 20	>1,000,000	27,270 ± 3690
AMPA	40 ^b	>100,000 ^b	>100,000 ^b	2000 ^b	>100,000 ^b	42,700 ± 12,500
ATPA	1800 ^b	23,000 ^b	>100,000 ^b	4 ^b	>100,000 ^b	2530 ± 250
KA ^b	4000	7	>100,000	76	13	33

^a KA_{1,2} are the predominant KA receptor subtypes expressed in native rat synaptosomes.

^b Data taken from original papers: **1a**, ([BraunerOsborne et al., 1997], [Gu et al., 1995], [Sagot et al., 2008] and [Zhou et al., 1997]); **2a**, ([Baker et al., 2000] and [BraunerOsborne et al., 1997]); **2b–e,g**, (Baker et al., 2000); **2f,h**, (Bunch et al., 2001); AMPA, ([Clarke et al., 1997] and [Vogensen et al., 2000]); ATPA, ([Clarke et al., 1997] and [Stensbol et al., 1999]); KA, ([Baker et al., 2000], [Conti et al., 1999] and [Sagot et al., 2008]).

2. Materials and methods

2.1. Binding affinities at native and homomeric ionotropic Glu receptors

Binding affinities for **1b–e,g** at native AMPA, KA, NMDA receptors (rat synaptosomes) were determined according to the previously published experimental procedure (Hermit et al., 2004) using radioligands [³H]AMPA, [³H]KA (representing predominantly subtypes KA_{1,2}), and [³H]CGP39653, respectively. Determination of binding affinities for **1b–e,g** and **2f,h** at cloned rat homomeric receptor subtypes iGlu_{5–7} were carried out following the procedure described earlier, using [³H]-SYM2081 as the radioligand (Sagot et al., 2008).

2.2. Molecular modelling

The modeling study was performed using the software package MOE (Molecular Operating Environment, v2004.03, Chemical Computing Group, 2004) using the build-in mmff94x forcefield and the GB/SA continuum solvent model. Compound **1g** was submitted to a stochastic conformational search and with respect to its global minimum returned (ΔG in kcal/mol), conformations above + 7 kcal/mol were discarded. The γ -carboxylate group was protonated prior to execution of the conformational search, as this gave a larger and thus more reliable number of output conformations. Superimpositions of ligands, were carried out using the built-in function in MOE, by fitting the ammonium group and the two carboxylate groups. The conformation of ATPA was adapted from X-ray structure when bound in the iGlu₂ subunit (PDB: 1 nnk).

3. Results

3.1. Binding affinities at native Glu receptor subtypes

In binding assays at native AMPA, KA and NMDA receptors, **1b** shows preference for the KA receptors with decreasing affinity as the 4-substituent is enlarged in size and bulk (**1a–e,g**). Eventually, compound **1g** displays very low affinity for any of the three iGlu receptor subgroups (42, > 100, > 100 μ M, respectively, Table 1).

3.2. Binding affinities at cloned homomeric Glu receptor subtypes

At cloned homomeric iGlu_{5–7} receptor subtypes, Glu analog **1b** displayed a 35- and 10-fold preference for iGlu₅ over subtypes iGlu_{6,7}, respectively. As the 4*R*-alkyl substituent is increased in length and bulkiness, this trend is strengthened (Table 1 and Fig. 1), and eventually Glu analog **1g** is highly selective for iGlu₅ receptor subtype ($K_i = 20.5$ nM), displaying more than 5000-fold and 300-fold selectivity over iGlu_{6,7} receptor subtype, respectively (Table 2).




Fig. 1. Binding affinities (pK_i [nM]) of (4*R*)-alkyl Glu analogs **1a–e,g** (top) and *E*-4-alkylidene Glu analogs **2a–h** (bottom) at cloned homomeric iGlu_{5–7} receptor subtypes.

Table 2.

Receptor subtype selectivity calculated and normalized from data presented in Table 1 (iGlu₅ receptor subtype set to 1).

	AMPA	KA _{1,2} ^a	NMDA	iGlu ₅	iGlu ₆	iGlu ₇
1g	2050	> 5000	> 5000	1	> 5000	300
2f	80	> 4200	> 4200	1	> 4200	90
2g	–	–	–	1	2830	–
ATPA	450	5750	> 25,000	1	> 25,000	2500

^a Predominant KA receptor subtypes expressed in native rat synaptosomes.

4. Discussion

Despite the fact that Glu is a highly flexible molecule (Fig. 2), it has been shown in several X-ray crystallographic and medicinal chemistry studies, that Glu agonizes iGlu receptors in a well-defined conformation termed the *folded* conformation ([Bunch and Krogsgaard-Larsen, 2009], [Bunch et al., 2003] and [Hogner et al., 2002]). On the other hand, when activating the mGlu receptors, Glu adopts an *extended* conformation ([Hayashi et al., 1992] and [Kozikowski et al., 1998]).




Fig. 2. Rotation of the C(2)–C(3) and C(3)–C(4) bonds allows Glu to adapt nine different staggered conformations. Of these, the two low-energy conformations are termed the Glu *folded* and *extended* conformations. These two Glu conformations are also observed when Glu is crystallized with iGlu receptor subunits (e.g. the iGlu₅ subunit: PDB code: 1TXF) and mGlu₁ subunit (PDB code: 1EWK).

Recently we reported the synthesis and pharmacological evaluation of a series of (4*R*)-alkyl Glu analogs at EAAT_{1–3} (Alaux et al., 2005). In extension from earlier findings, we showed that introduction of a large variety of longer and more bulky (4*R*)-alkyl substituents also endorses the *folded* conformation as the global low-energy conformation of Glu. In comparison, this low-energy conformational preference is also preferred for *E*-4-alkylidene Glu analogs **2a–h**, AMPA, and ATPA (Fig. 3). Furthermore, a general feature for all of these Glu analogs, is the observed increase in preference for the iGlu₅ receptor subtype, as the alkyl group is extended in length and bulk (**1a** → **1g**, **2a** → **2f** and AMPA → ATPA; Table 1). This observation may be explained by comparing the size of the iGlu receptor binding pockets: For subunit iGlu₅ it has been estimated to be approximately 20% larger as compared to subunit iGlu₆ and approximately 50% larger, when compared to subunit iGlu₂ (305 Å, 255 Å, and 218 Å, respectively) (Mayer, 2005).



Fig. 3. Superimposition of low-energy conformations of **1g** ($\Delta\Delta G = 0$ to + 1 kcal/mol) (type code/green) and ATPA (purple). The conformation of **1g** shown in green resembles the conformation of ATPA best.

(For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

(4*R*)-Alkyl Glu analogs **1a–e,g** are conformationally released structural analogs of 4-alkylidenes **2a–h**. This confined modification of physical chemical property may allow for a detailed investigation of ligand flexibility opposed to receptor subtype selectivity. Comparing the binding affinity profiles of series **1** and **2** at iGlu_{5–7} receptor subtypes reveals that both series display increasing selectivity as well as binding affinity for iGlu₅, as the 4-substituents is extended in length and bulk. However, from this resemblance two divergences are noted: Firstly, **1g** vs. **2g** displays a 10-fold difference in binding affinity at the iGlu₅ receptor subtype ($K_i = 20.5$ and 224 nM, respectively). This finding is intriguing and raises the question as to what origin this observation has. One explanation could be that the more flexible ligand, **1g**, may allow for a more tight domain closure, favoring both ligand–receptor and receptor–receptor interactions. A different explanation could be enhanced hydrophobic interactions with the receptor protein. Secondly, **2d** stands out from the trend, displaying significantly lower binding affinity to the iGlu₅ receptor subtype as compared to **2c** and **2e**, as well as **1c–e**. However, this is explained on the basis that **2d** is structurally quite distinct from preceding analogs **2a–c** and subsequent analogs **2e–f**, as it has a methyl group in the Z-1' position as opposed to hydrogen.

In conclusion, Glu analogs **1b–e,g** are a conformationally released structural design of **2a–h** and ATPA. Most notably, **1g** displays high affinity for the KA receptor subtype iGlu₅ with a high degree of selectivity over receptor subtypes iGlu_{6,7} (fold ratio: > 5000, 300, respectively), KA subtypes KA₁/KA₂ (fold ratio: > 5000, determined by KA binding at native rat synaptosomes), AMPA receptors (2050 fold, determined by AMPA binding at native rat synaptosomes), and NMDA receptors (> 5000 fold, determined by CGP39653 binding at native rat synaptosomes), making **1g** a valuable pharmacological tool. Furthermore, **1g** displays an intriguing 10-fold higher binding affinity at iGlu₅ receptor subtype than its corresponding *E*-4-alkylidene isomer **2g**. This observation encourages an expansion of the (4*R*)-alkyl Glu series by the synthesis of corresponding saturated analogs of **2f,h**, and a following investigation whether such difference in binding affinity at iGlu₅ receptor subtype is a generally observable fact and ultimately if it translates into functional differences.

Acknowledgements

We would like to thank the Carlsberg Foundation, the Lundbeck Foundation, and the Danish Research Council for financial support.

References

- Alaux et al., 2005 S. Alaux, M. Kusk, E. Sagot, J. Bolte, A.A. Jensen, H. Brauner-Osborne, T. Gefflaut and L. Bunch, Chemoenzymatic synthesis of a series of 4-substituted glutamate analogues and pharmacological characterization at human glutamate transporters subtypes 1–3, *J. Med. Chem.* **48** (2005), pp. 7980–7992.
- Baker et al., 2000 S.R. Baker, D. Bleakman, J. Ezquerra, B.A. Ballyk, M. Deverill, K. Ho, R.K. Kamboj, I. Collado, C. Dominguez, A. Escribano, A.I. Mateo, C. Pedregal and A. Rubio, 4-Alkylidenyl glutamic acids, potent and selective GluR5 agonists, *Bioorg. Med. Chem. Lett.* **10** (2000), pp. 1807–1810.
- Beart and O'Shea, 2007 P.M. Beart and R.D. O'Shea, Transporters for l-glutamate: an update on their molecular pharmacology and pathological involvement, *Br. J. Pharmacol.* **150** (2007), pp. 5–17.
- Brauner-Osborne et al., 1997 H. Brauner-Osborne, B. Nielsen, T.B. Stensbol, T.N. Johansen, N. Skjaerbaek and P. Krogsgaard-Larsen, Molecular pharmacology of 4-substituted glutamic acid analogues at ionotropic and metabotropic excitatory amino acid receptors, *Eur. J. Pharmacol.* **335** (1997), pp. R1–R3.
- Bräuner-Osborne et al., 2000 H. Bräuner-Osborne, J. Egebjerg, E.O. Nielsen, U. Madsen and P. Krogsgaard-Larsen, Ligands for glutamate receptors: design and therapeutic prospects, *J. Med. Chem.* **43** (2000), pp. 2609–2645.
- Bunch and Krogsgaard-Larsen, 2009 L. Bunch and P. Krogsgaard-Larsen, Subtype selective kainic acid receptor agonists: discovery and approaches to rational design, *Med. Res. Rev.* **29** (2009), pp. 3–28.
- Bunch et al., 2001 L. Bunch, T.H. Johansen, H. Bräuner-Osborne, T.B. Stensbol, T.N. Johansen, P. Krogsgaard-Larsen and U. Madsen, Synthesis and receptor binding affinity of new selective GluR5 ligands, *Bioorg. Med. Chem.* **9** (2001), pp. 875–879.
- Bunch et al., 2003 L. Bunch, T. Liljefors, J.R. Greenwood, K. Frydenvang, H. Bräuner-Osborne, P. Krogsgaard-Larsen and U. Madsen, Rational design, synthesis, and pharmacological evaluation of 2-azanorbornane-3-exo,5-endo-dicarboxylic acid: a novel conformationally restricted glutamic acid analogue, *J. Org. Chem.* **68** (2003), pp. 1489–1495.
- Clarke et al., 1997 V.R.J. Clarke, B.A. Ballyk, K.H. Hoo, A. Mandelzys, A. Pellizzari, C.P. Bath, J. Thomas, E.F. Sharpe, C.H. Davies, P.L. Ornstein, D.D. Schoepp, R.K. Kamboj, G.L. Collingridge, D. Lodge and D. Bleakman, A hippocampal GluR5 kainate receptor regulating inhibitory synaptic transmission, *Nature* **389** (1997), pp. 599–603.

- Conti et al., 1999 P. Conti, M. De Amici, G. De Sarro, M. Rizzo, T.B. Stensbol, H. Brauner-Osborne, U. Madsen, L. Toma and C. De Micheli, Synthesis and enantiopharmacology of new AMPA-kainate receptor agonists, *J. Med. Chem.* **42** (1999), pp. 4099–4107.
- Ferraguti and Shigemoto, 2006 F. Ferraguti and R. Shigemoto, Metabotropic glutamate receptors, *Cell Tissue Res.* **326** (2006), pp. 483–504.
- Gu et al., 1995 Z.Q. Gu, D.P. Hesson, J.C. Pelletier, M.L. Maccacchini, L.M. Zhou and P. Skolnick, Synthesis, resolution, and biological evaluation of the 4-stereoisomers of 4-methylglutamic acid — selective probes of kainate receptors, *J. Med. Chem.* **38** (1995), pp. 2518–2520.
- Hayashi et al., 1992 Y. Hayashi, Y. Tanabe, I. Aramori, M. Masu, K. Shimamoto, Y. Ohfuné and S. Nakanishi, Agonist analysis of 2-(carboxycyclopropyl)glycine isomers for cloned metabotropic glutamate receptor subtypes expressed in Chinese hamster ovary cells, *Br. J. Pharmacol.* **107** (1992), pp. 539–543.
- Hermit et al., 2004 M.B. Hermit, J.R. Greenwood, B. Nielsen, L. Bunch, C.G. Jorgensen, H.T. Vestergaard, T.B. Stensbol, C. Sanchez, P. Krosggaard-Larsen, U. Madsen and H. Bräuner-Osborne, Ibotenic acid and thioibotenic acid: a remarkable difference in activity at group III metabotropic glutamate receptors, *Eur. J. Pharmacol.* **486** (2004), pp. 241–250.
- Hogner et al., 2002 A. Hogner, J.S. Kastrup, R. Jin, T. Liljefors, M.L. Mayer, J. Egebjerg, I.K. Larsen and E. Gouaux, Structural basis for AMPA receptor activation and ligand selectivity: crystal structures of five agonist complexes with the GluR2 ligand-binding core, *J. Mol. Biol.* **322** (2002), pp. 93–109.
- Kozikowski et al., 1998 A.P. Kozikowski, D. Steensma, G.L. Araldi, W. Tuckmantel, S.M. Wang, S. Pshenichkin, E. Surina and J.T. Wroblewski, Synthesis and biology of the conformationally restricted ACPD analogue, 2-aminobicyclo[2.1.1]hexane-2,5-dicarboxylic acid-I, a potent mGluR agonist, *J. Med. Chem.* **41** (1998), pp. 1641–1650.
- Mayer, 2005 M.L. Mayer, Crystal structures of the GluR5 and GluR6 ligand binding cores: molecular mechanisms underlying kainate receptor selectivity, *Neuron* **45** (2005), pp. 539–552.
- Meldrum, 2000 B.S. Meldrum, Glutamate as a neurotransmitter in the brain: review of physiology and pathology, *J. Nutrition* **130** (2000), pp. 1007S–1015S.
- Sagot et al., 2008 E. Sagot, D.S. Pickering, X. Pu, M. Umberti, T.B. Stensbøl, B. Nielsen, M. Chapelet, J. Bolte, T. Gefflaut and L. Bunch, Chemo-enzymatic synthesis of a series of 2,4-syn-functionalized (S)-glutamate analogues: new insight into the structure–activity–relation of ionotropic glutamate receptor subtypes 5, 6 and 7, *J. Med. Chem.* **51** (2008), pp. 4093–4103.
- Stensbol et al., 1999 T.B. Stensbol, L. Borre, T.N. Johansen, J. Egebjerg, U. Madsen, B. Ebert and P. Krosggaard-Larsen, Resolution, absolute stereochemistry and molecular pharmacology of the enantiomers of ATPA, *Eur. J. Pharmacol.* **380** (1999), pp. 153–162.
- Vogensen et al., 2000 S.B. Vogensen, H.S. Jensen, T.B. Stensbol, K. Frydenvang, B. Bang-Andersen, T.N. Johansen, J. Egebjerg and P. Krosggaard-Larsen, Resolution, configurational assignment, and enantiopharmacology of 2-amino-3-[3-hydroxy-5-(2-methyl-2H-tetrazol-5-yl)isoxazol-4-yl]propionic acid, a potent GluR3- and GluR4-preferring AMPA receptor agonist, *Chirality* **12** (2000), pp. 705–713.
- Wisden and Seeburg, 1993 W. Wisden and P.H. Seeburg, A complex mosaic of high-affinity kainate receptors in rat-brain, *J. Neurosci.* **13** (1993), pp. 3582–3598.
- Zhou et al., 1997 L.M. Zhou, Z.Q. Gu, A.M. Costa, K.A. Yamada, P.E. Mansson, T. Giordano, P. Skolnick and K.A. Jones, (2S,4R)-4-methylglutamic acid (SYM 2081): a selective, high-affinity ligand for kainate receptors, *J. Pharmacol. Exp. Ther.* **280** (1997), pp. 422–427.