



**HAL**  
open science

# A note on uniqueness of entropy solutions to degenerate parabolic equations in $R^N$ .

Boris Andreianov, Mohamed Maliki

## ► To cite this version:

Boris Andreianov, Mohamed Maliki. A note on uniqueness of entropy solutions to degenerate parabolic equations in  $R^N$ . 2009. hal-00378475v1

**HAL Id: hal-00378475**

**<https://hal.science/hal-00378475v1>**

Preprint submitted on 24 Apr 2009 (v1), last revised 5 Sep 2009 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A NOTE ON UNIQUENESS OF ENTROPY SOLUTIONS TO DEGENERATE PARABOLIC EQUATIONS IN $\mathbb{R}^N$ .

BORIS ANDREIANOV AND MOHAMED MALIKI

ABSTRACT. We study the Cauchy problem in  $\mathbb{R}^N$  for the parabolic equation  $u_t + \operatorname{div} F(u) = \Delta\varphi(u)$ , which can degenerate into a hyperbolic equation for some intervals of values of  $u$ . In the context of conservation laws (the case  $\varphi \equiv 0$ ), it is known that an entropy solution can be non-unique when  $F'$  has strong singularities. We show the uniqueness of an entropy solution to the general parabolic problem for all  $L^\infty$  initial datum, under the isotropic condition on the flux  $F$  known for conservation laws. The only assumption on the diffusion term is that  $\varphi$  is a non-decreasing continuous function.

## 1. INTRODUCTION

Consider the convection-diffusion problem

$$(1) \quad u_t + \operatorname{div} F(u) - \Delta\varphi(u) = f \quad \text{in } (0, T) \times \mathbb{R}^N, \quad u|_{t=0} = u_0$$

with a source  $f$  measurable on  $(0, T) \times \mathbb{R}^N$ , with  $\int_0^T \|f(t, \cdot)\|_{L^\infty} dt < +\infty$  and an initial datum  $u_0 \in L^\infty(\mathbb{R}^N)$ . Here  $F = (F_1, \dots, F_N) : \mathbb{R} \mapsto \mathbb{R}^N$  is assumed merely continuous (a quantitative continuity assumption on  $F$  will be imposed in the sequel), and  $\varphi : \mathbb{R} \mapsto \mathbb{R}$  is a continuous non-decreasing function.

Degenerate and non-degenerate problems of the kind (1), often investigated under the local Lipschitz continuity assumption on  $\varphi$  and  $F$ , appear in various physical contexts, including e.g. flows and sedimentation in porous media. Mathematical treatment of these problems was boosted some ten years ago by the introduction of the adequate notion of entropy solution (see Carrillo [8]).

Indeed, the framework (1) includes conservation laws as a particular case. It is well known that in general, classical solutions of (1) may not exist; and that a weak solution in the sense of distributions may not be unique. The standard way to fix this problem is to work with the so-called entropy solutions (see Kruzhkov [9] for the case of conservation laws, and Carrillo [8] for the adaptation of this notion to the case of a degenerate elliptic-parabolic-hyperbolic equation, of which (1) is a particular case). The classical result of Kruzhkov establishes uniqueness of an entropy solution (together with the property of finite domain of dependence on initial data) for the case where  $\varphi \equiv 0$  and  $F$  is locally Lipschitz continuous. When the Lipschitz continuity assumption is dropped, uniqueness of entropy solutions for a conservation law can be lost (see the works [16, 13, 10, 14] of Kruzhkov and Panov). The effect of non-uniqueness is

---

*Date:* April 24, 2009.

*2000 Mathematics Subject Classification.* Primary 35K65; Secondary 35A05.

*Key words and phrases.* Degenerate hyperbolic-parabolic equation, conservation law, non-Lipschitz flux, entropy solution, Kato inequality, infinite speed of propagation, uniqueness.

closely connected to the property of “infinite speed of propagation”, which arises when  $F'$  is unbounded.

This is the goal of this note to investigate the question of uniqueness of entropy solutions of (1) for a general nonlinearity  $\varphi$  in the diffusion term. Technically, the problem amounts to the difficulty of exploiting the so-called Kato inequality

$$(2) \quad (u - \hat{u})_t^+ + \operatorname{div} (\operatorname{sign}^+(u - \hat{u})(F(u) - F(\hat{u}))) \\ \leq \Delta(\varphi(u) - \varphi(\hat{u}))^+ + (f - \hat{f})^+ \quad \text{in } \mathcal{D}'([0, T] \times \mathbb{R}^N);$$

here  $u, \hat{u}$  are two entropy solutions of (1) corresponding to source terms  $f, \hat{f}$ . Formally, if we let the test function in (2) converge to zero, we get the inequality

$$\frac{d}{dt} \int_{\mathbb{R}^N} (u - \hat{u})^+(t) \leq \int_{\mathbb{R}^N} (f - \hat{f})^+ \quad \text{in } \mathcal{D}'([0, T]),$$

which implies uniqueness of an entropy solution. Even if  $u - \hat{u} \in L^\infty(0, T; L^1(\mathbb{R}^N))$ , a too slow decrease of  $|F(u) - F(\hat{u})|$  as  $|x| \rightarrow \infty$  could be an obstacle to this formal reasoning. In fact, the issue of uniqueness of  $L^1 \cap L^\infty$  solutions for the conservation law  $u_t + \operatorname{div} F(u) = 0, u|_{t=0} = u_0$  is not yet clearly understood; see [2] for a discussion of the relevancy of known conditions.

For the case  $\varphi \equiv 0$  and  $L^\infty$  entropy solutions, sufficient uniqueness conditions for (1) in terms of the moduli of continuity of  $F$  were first formulated by Kruzhkov and Hil'debrand ([12]) (after that B enilan ([5] obtained the same condition for mild solutions in  $L^1$ ). Roughly speaking, the conditions of [5, 12] (see also [11]) amount to the H older continuity of  $F$  of order  $\alpha = 1 - \frac{1}{N}$ . This condition is “isotropic”: the possibility that some of the components  $F_1, \dots, F_N$  of  $F$  may have a “better” modulus of continuity than the vector-function  $F$  is not exploited. This contrasts with the following “anisotropic” assumption

$$(3) \quad \liminf_{r \downarrow 0} \frac{\prod_{i=1}^N \omega_i(r)}{r^{N-1}} < +\infty$$

due to Kruzhkov and Panov ([13, 16]); here for  $i = 1, \dots, N$ ,  $\omega_i : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a modulus of continuity of  $F_i$  on the interval  $[-M, M]$  containing the values of  $u$ . Assumption (3) restricts the behaviour of the moduli of continuity  $\omega_i(r)$  of  $F_i$  at  $r = 0$ . Assumption (3) is also sufficient for the uniqueness of entropy solutions of a scalar conservation law. A sharper condition is given in [14]. Notice that for the  $L^1 \cap L^\infty$  data, sufficient uniqueness conditions of a completely different nature were found by B enilan and Kruzhkov in [6] (see also [2]).

In [15], together with Tour e the second author of the present paper proved uniqueness of an entropy solution to the equation (1) upon replacing  $\omega_i(r)$  in (3) with  $(\omega_i(r) + (r\omega_\varphi(r))^{1/2})$ ; here  $\omega_\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a modulus of continuity of  $\varphi$ . Let us point out that, if the space dimension  $N$  equals 1, both the Kruzhkov-Panov and the Maliki-Tour e conditions are always satisfied. If the space dimension is equal to 2, then the Maliki-Tour e condition contains no restriction on the function  $\varphi$ .

In the work [3] we have investigated the exactness of the assumption of [15] for the case of a pure diffusion equation (i.e.,  $F \equiv 0$ ). It turns out that uniqueness of  $L^\infty$  solutions for this case holds for all  $\varphi$ . In the present paper, we show a similar result for the full convection-diffusion equation (1). More exactly, we show uniqueness of an entropy solution for (1) for all continuous non-decreasing  $\varphi$ , but under the assumption that  $F$  satisfies an isotropic condition

which essentially means that  $F$  is Hölder continuous of order  $\alpha = 1 - \frac{1}{N}$ . According to the previous remark, we can restrict our attention to the case  $N \geq 3$ .

Our technique does not allow to benefit from the anisotropy of the flux  $F$ ; in particular, unlike in [15], our result does not reduce (3) in the trivial case  $\varphi \equiv 0$ . The case of an anisotropic diffusion of the form  $\sum_{i=1}^N \partial_{x_i x_i} \varphi_i(u)$  cannot be covered neither; we refer to Maliki and Ouedraogo [1] for results on this non-isotropic diffusion case under suitable assumptions on the moduli of continuity of  $\varphi_i$ . The problem where the quasilinear diffusion operator  $\Delta\varphi(u)$  is replaced with a fully nonlinear diffusion operator such as the  $p$ -laplacian  $\Delta_p\varphi(u)$  is even more difficult; to our knowledge, no uniqueness result for entropy solutions in  $\mathbb{R}^N$  is available when  $p > 2$ .

## 2. DEFINITIONS AND RESULTS

Recall that  $\varphi$  is a continuous function on  $\mathbb{R}$ . For  $M > 0$ , we say that a sub-additive, continuous function  $\omega_\varphi : [0, 2M] \mapsto \mathbb{R}^+$  such that  $\omega_\varphi(0) = 0$  is a modulus of continuity of  $\varphi$ , if

$$(4) \quad \forall z, \hat{z} \in [-M, M] \quad |\varphi(z) - \varphi(\hat{z})| \leq \omega_\varphi(|z - \hat{z}|).$$

Recall that we only work with bounded solutions of (1). Without loss of generality, we can assume that  $M$  is fixed, moreover,  $\omega_\varphi$  is strictly convex (and thus, strictly increasing) and extended to the whole of  $\mathbb{R}^+$ ; we denote its inverse function by  $\Omega_\varphi$ . We use the Fenchel (Legendre) transform of  $\Omega_\varphi$ :

$$\Omega_\varphi^*(t) = \sup_{z \in \mathbb{R}} (zt - \Omega_\varphi(z)).$$

It easily follows that  $\Omega_\varphi^*$  is strictly increasing, thus it is invertible. We denote by  $\omega_\varphi^*$  the inverse of  $\Omega_\varphi$ . With the above notation, we have for all  $z, t \in \mathbb{R}^+$ ,

$$(5) \quad zt \leq \Omega_\varphi(z) + \Omega_\varphi^*(t), \quad (\Omega_\varphi \circ \omega_\varphi)(z) = z, \quad \text{and} \quad (\Omega_\varphi^* \circ \omega_\varphi^*)(t) = t.$$

For a vector-valued function  $F : \mathbb{R} \mapsto \mathbb{R}^N$ , we introduce the (isotropic) modulus of continuity  $\omega_F$  and the related functions  $\Omega_F, \Omega_F^*$  and  $\omega_F^*$  in the same way, replacing  $|\varphi(z) - \varphi(\hat{z})|$  in (4) with  $|F(z) - F(\hat{z})|$ . Here and in the sequel, we mean that  $\mathbb{R}^N$  is endowed with the euclidean norm, which we denote by  $|\cdot|$ .

**Definition 1.** *An entropy solution of (1) is a function  $u \in L^\infty((0, T) \times \mathbb{R}^N)$  such that  $\varphi(u) \in L^2(0, T; H_{loc}^1(\mathbb{R}^N))$  and the inequalities*

$$(6) \quad \begin{aligned} & - \int_0^T \int_{\mathbb{R}^N} \left( |u - k| \xi_t + \text{sign}(u - k) (F(u) - F(k)) \cdot \nabla \xi + \text{sign}(u - k) \nabla \varphi(u) \cdot \nabla \xi \right) \\ & \leq \int_0^T \int_{\mathbb{R}^N} \text{sign}(u - k) f \xi + \int_{\mathbb{R}^N} |u_0 - k| \xi(0, \cdot) \end{aligned}$$

hold for all  $k \in \mathbb{R}$  and all  $\xi \in \mathcal{D}([0, T) \times \mathbb{R}^N)$ .

Here and in the sequel, we mean that  $u_0 \in L^\infty(\mathbb{R}^N)$  and  $\|f(t, \cdot)\|_{L^\infty} \in L^1(0, T)$ . It is clear that an entropy solution is also a solution in the sense of distributions; it suffices to use (6) with  $k = \|u\|_{L^\infty}$ . It follows that in (6), one can replace  $\text{sign}(u - k)$  and  $|u - k|$  with  $\text{sign}^+(u - k)$  and  $(u - k)^+$ , respectively, where  $\text{sign}^+(r) := \max\{\text{sign } r, 0\}$  and  $r^+ := \max\{r, 0\}$ .

Existence of an entropy solution is standard; it can be obtained, e.g., by a regularization of  $F$  and  $\varphi$  (when  $\varphi \equiv 0$ , one takes  $\varphi_\varepsilon(z) = \varepsilon z$ , which is the so-called “vanishing viscosity” approximation), with the help of the translation invariance in  $x$  of the left-hand side of (1) and the  $L^1$  contraction result of Maliki and Touré [15].

Definition 1 implies the Kato inequality (2), with the help of the doubling-of-variables approach of Kruzhkov ([9]) and the techniques of Carrillo ([8]) and Blanchard and Porretta ([7]) (see e.g. [15, 4, 3]). Inequality (2) can also be obtained without doubling the variables, with the help of the so-called kinetic formulation (see Perthame [17] for the case of a conservation law). Uniqueness of an entropy solution of (1) (and, more generally, the  $L^1$  contraction and comparison principle) follows whenever we can pass to the limit in the Kato inequality (2) with test functions  $\xi \rightarrow 1$  on  $\mathbb{R}^N$ .

We assume that  $N \geq 3$  (for  $N \leq 2$ , the results of [15] apply with no restriction on  $\varphi$ ).

**Theorem 2.** *Let  $N \geq 3$ . Consider an arbitrary non-decreasing continuous function  $\varphi : \mathbb{R} \mapsto \mathbb{R}$ . Consider a continuous function  $F : \mathbb{R} \mapsto \mathbb{R}^N$  which possesses a modulus of continuity  $\omega_F$  such that, with the notation introduced above, there holds*

$$(7) \quad I_F(R, \varepsilon) := R^{N-2+\varepsilon} \int_R^{+\infty} \Omega_F^*\left(\frac{1}{r}\right) r^{1-\varepsilon} dr \leq \text{const} \quad \text{uniformly in } \varepsilon > 0, R > 1.$$

Assume in addition that the “ $\Delta_2$ -condition” on  $\Omega_F$  holds:

$$(8) \quad \Omega_F(2z) \leq \text{const} \Omega(z) \quad \forall z \in \mathbb{R}^+.$$

Then whenever  $(u_0 - \hat{u}_0)^+ \in L^1(\mathbb{R}^N)$ ,  $(f - \hat{f})^+ \in L^1((0, T) \times \mathbb{R}^N)$ ,

$$(9) \quad \int_{\mathbb{R}^N} (u - \hat{u})^+(t, \cdot) \leq \int_{\mathbb{R}^N} (u_0 - \hat{u}_0)^+ + \int_0^t \int_{\mathbb{R}^N} (f - \hat{f})^+ \quad \text{a.e. on } (0, T),$$

where  $u, \hat{u}$  are entropy solutions corresponding, respectively, to given initial data  $u_0, \hat{u}_0 \in L^\infty(\mathbb{R}^N)$  and sources  $f, \hat{f}$  with  $\|f(t, \cdot)\|_{L^\infty(\mathbb{R})}, \|\hat{f}(t, \cdot)\|_{L^\infty(\mathbb{R})} \in L^1(0, T)$ . In particular, under assumptions (7), (8) there exists a unique entropy solution  $u_{u_0, f}$  for given data  $u_0, f$ ; the map  $(u_0, f) \mapsto u_{u_0, f}$  is order-preserving, and the map  $u_0 \in L^\infty(\mathbb{R}^N) \cap L^1(\mathbb{R}^N) \mapsto u_{u_0, 0}$  is a  $T$ -contraction in  $L^1$ .

**Remark 3.**

- (i) If we replace (7) by the stronger requirement that  $\sup_{\varepsilon > 0} I_F(R, \varepsilon) \rightarrow 0$  as  $R \rightarrow +\infty$ , the assumption (8) can be dropped and the claim of the theorem continues to hold (see the proof of Theorem 2).
- (ii) A sufficient condition is that  $\omega_F(z) \leq \text{const} z^\alpha$  with  $\alpha = 1 - \frac{1}{N}$  (i.e., that  $F$  is Hölder continuous of order  $1 - \frac{1}{N}$ ); in this case, we can pick  $\Omega_F(z) = \text{const} z^{\frac{1}{\alpha}}$  and  $\Omega_F^*(t) = \text{const} t^{\frac{1}{1-\alpha}} \equiv \text{const} t^N$ , for which conditions (7), (8) are easy to check.

### 3. PROOF OF THEOREM 2

The technique of the proof of the theorem is closely related to the previous work [3]. Roughly speaking, we take in the Kato inequality (2) radial test functions  $\xi$  such that  $\Delta \xi \leq 0$ . This allows us to drop the second term in the right-hand side of (2); thus restrictions on  $\varphi$

are bypassed. The test functions are constructed by truncating the laplacian's fundamental solution profile  $\frac{1}{|x|^{N-2}}$ .

We are intended to use (2) with the test functions  $\rho_\varepsilon(x) = (\max\{\frac{|x|}{R}, 1\})^{2-N-\varepsilon}$ ,  $\varepsilon > 0$ ,  $R > 1$  (in order to lighten the notation, we do not indicate explicitly that  $\rho_\varepsilon$  also depends on  $R$ ). Then we let  $\varepsilon \rightarrow 0$ , and finally we let  $R \rightarrow +\infty$ .

To this end, let  $\rho_{\varepsilon,L}(x) = R^{N-2+\varepsilon} \left( (\max\{|x|, R\})^{2-N-\varepsilon} - (L)^{2-N-\varepsilon} \right)^+$ ,  $L > R$ . By a density argument, we take in (2) test functions under the form  $\xi(t, x) = \mu(t)\rho_{\varepsilon,L}(x)$ , where  $\mu \in \mathcal{D}([0, T])^+$ . Indeed, regularizing  $\rho_{\varepsilon,L}$  by convolution, dropping the negative measure part of the distribution  $\Delta\rho_{\varepsilon,L}$  concentrated of  $\{x \in \mathbb{R}^N \mid |x| = R\}$ , using the definition of  $\omega_\varphi, \omega_F$  we deduce

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^N} (u - \hat{u})^+ \rho_{\varepsilon,L}(-\mu_t) &\leq \int_0^T \int_{\mathbb{R}^N} \omega_F((u - \hat{u})^+) |\nabla \rho_{\varepsilon,L}| \mu + \int_0^T \int_{\mathbb{R}^N} \omega_\varphi((u - \hat{u})^+) \Delta \rho_{\varepsilon,L} \mu \\ &\quad + \int_{\{|x|=L\}} \omega_\varphi((u - \hat{u})^+) (2-N-\varepsilon) L^{1-N-\varepsilon} \mu \\ &\quad + \int_0^T \int_{\mathbb{R}^N} (f - \hat{f}) \text{sign}^+(u - \hat{u}) \rho_{\varepsilon,L} \mu + \int_{\mathbb{R}^N} (u_0 - \hat{u}_0)^+ \rho_{\varepsilon,L} \mu(0) \end{aligned}$$

for a.e.  $L > R'$ ; here  $\Delta\rho_{\varepsilon,L}$  is taken in the pointwise (a.e. on  $\mathbb{R}^N$ ) sense. Because  $u, \hat{u}$  are bounded, the integral on the sphere  $\{|x| = L\}$  is majorated by  $\text{const} L^{N-1} L^{1-N-\varepsilon} = \text{const} L^{-\varepsilon}$ . We calculate  $\Delta\rho_{\varepsilon,L} = \varepsilon(N-2+\varepsilon)|x|^{-2}\rho_\varepsilon \mathbb{1}_{R < |x| < L}$  and  $|\nabla \rho_{\varepsilon,L}| = (N-2+\varepsilon)|x|^{-1}\rho_\varepsilon \mathbb{1}_{R < |x| < L}$ . Notice that  $|\rho_{\varepsilon,L}| \leq 1$  for all  $\varepsilon, R, L$ . Now we let  $L \rightarrow \infty$ . Thanks to the boundedness of  $u, \hat{u}$  and to the integrability assumption on  $(u_0 - \hat{u}_0)^+$ ,  $(f - \hat{f})^+$  we deduce that

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^N} (u - \hat{u})^+ \rho_\varepsilon(-\mu_t) &\leq c \int_0^T \int_{\{|x|>R\}} \left( \omega_F((u - \hat{u})^+) |x|^{-1} + \varepsilon \omega_\varphi((u - \hat{u})^+) |x|^{-2} \right) \rho_\varepsilon \\ (10) \quad &\quad + \int_0^T \int_{\mathbb{R}^N} (f - \hat{f}) \text{sign}^+(u - \hat{u}) \mu + \int_{\mathbb{R}^N} (u_0 - \hat{u}_0)^+ \mu(0). \end{aligned}$$

Here and in the sequel, we fix  $c = (N-1)\|\mu\|_{L^\infty}$  for  $\varepsilon \in (0, 1)$ .

Our goal is to show that, letting  $\varepsilon \rightarrow 0$  and  $R \rightarrow \infty$ , we can drop the first term in the right-hand side of (10). The first step is to prove that  $(u - \hat{u})^+$  is integrable on  $(0, T) \times \mathbb{R}^N$ .

To this end, fix  $\delta > 0$ . Let us show that

$$(11) \quad c\varepsilon \int_0^T \int_{\{|x|>R\}} \omega_\varphi((u - \hat{u})^+) |x|^{-2} \rho_\varepsilon \leq \delta \int_0^T \int_{\{|x|>R\}} (u - \hat{u})^+ \rho_\varepsilon + r^{\delta,R}(\varepsilon),$$

where for all fixed  $\delta > 0$ ,  $R > 1$ , the quantity  $r^{\delta,R}(\varepsilon)$  tends to zero as  $\varepsilon$  decreases to zero. Indeed, using (5), we get

$$c\varepsilon \omega_\varphi((u - \hat{u})^+) |x|^{-2} = \delta \left( \omega_\varphi((u - \hat{u})^+) \frac{c\varepsilon}{\delta} |x|^{-2} \right) \leq \delta (u - \hat{u})^+ + \delta \Omega_\varphi^* \left( \frac{c\varepsilon}{\delta} |x|^{-2} \right);$$

thus (11) holds with

$$r^{\delta,R}(\varepsilon) := T\delta \int_{\{|x|>R\}} \Omega_\varphi^* \left( \frac{c\varepsilon}{\delta} |x|^{-2} \right) \rho_\varepsilon.$$

By the convexity of  $\Omega_\varphi^*$ , we have  $\Omega_\varphi^*(\frac{c\varepsilon}{\delta}|x|^{-2}) \leq (\Omega_\varphi^*)'(\frac{c\varepsilon}{\delta}R^{-2})\frac{c\varepsilon}{\delta}|x|^{-2}$  whenever  $|x| \geq R$ . Without loss of generality, we can assume that  $\Omega_\varphi$  is convex and strictly increasing; then it is not difficult to prove that  $\text{ess lim}_{t \downarrow 0} (\Omega_\varphi^*)'(t) = 0$  (see e.g. [3]). Hence we deduce that  $(\Omega_\varphi^*)'(\frac{c\varepsilon}{\delta}R^{-2}) =: \bar{\delta}_\varepsilon$  tends to zero as  $\varepsilon \rightarrow 0$ . Therefore using polar coordinates, we get

$$\begin{aligned} 0 \leq r^{\delta,R}(\varepsilon) &= \delta T \int_{\{|x|>R\}} \Omega_\varphi^*(\frac{c\varepsilon}{\delta}|x|^{-2}) \rho_\varepsilon \leq \bar{\delta}_\varepsilon cT \varepsilon \int_{\{|x|>R\}} |x|^{-2} \rho_\varepsilon \\ (12) \quad &= cT \varepsilon R^{N-2+\varepsilon} \bar{\delta}_\varepsilon \int_R^{+\infty} r^{-2+2-N-\varepsilon} r^{N-1} dr = cT \varepsilon R^{N-2+\varepsilon} \frac{R^{-\varepsilon}}{\varepsilon} \bar{\delta}_\varepsilon \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Similarly, from (5) we deduce

$$(13) \quad c\omega_F((u - \hat{u})^+) |x|^{-1} \rho_\varepsilon \leq \frac{1}{2}(u - \hat{u})^+ \rho_\varepsilon + \Omega_F^*(2c|x|^{-1}) \rho_\varepsilon;$$

moreover, using assumption (7) we get

$$(14) \quad \int_0^T \int_{\{|x|>R\}} \Omega_F^*(2c|x|^{-1}) \rho_\varepsilon = T R^{N-2+\varepsilon} \int_R^{+\infty} \Omega_F^*\left(\frac{2c}{r}\right) r^{1-\varepsilon} dr \leq \text{const } I_F(R/(2c), \varepsilon) \leq \text{const}.$$

By density, we can take  $\mu(\cdot) = (T - \cdot)^+$  in (10), so that  $-\mu_t(t) = \mathbb{1}_{(0,T)}(t)$ . Putting  $\delta = 1/2$  and letting  $\varepsilon \rightarrow 0$ , from (10) and (11)–(14) we deduce that

$$\int_0^s \int_{\{|x|<R\}} (u - \hat{u})^+ \leq T \int_{\mathbb{R}^N} (u_0 - \hat{u}_0)^+ + T \int_0^T \int_{\mathbb{R}^N} (f - \hat{f})^+ + \text{const} \leq \text{const}.$$

Letting  $R \rightarrow \infty$  and  $s \rightarrow T$ , by the monotone convergence theorem we deduce that

$$(15) \quad (u - \hat{u})^+ \in L^1((0, T) \times \mathbb{R}^N).$$

In order to conclude the proof, we return to (10). We now replace (13) with the estimate

$$(16) \quad c\omega_F((u - \hat{u})^+) |x|^{-1} \rho_\varepsilon \leq \delta c \Omega_F\left(\frac{1}{\delta}\omega_F((u - \hat{u})^+)\right) \rho_\varepsilon + \delta c \Omega_F^*(|x|^{-1}) \rho_\varepsilon.$$

Denote by  $C(\delta)$  a generic constant that does not depend on  $\varepsilon, R$ . Assumption (8) implies that for  $t \in \mathbb{R}^+$ ,  $\Omega_F(\frac{1}{\delta}\omega_F(t)) \leq C(\delta) \Omega_F(\omega_F(t)) = C(\delta)t$ . Now using (7), we infer

$$\begin{aligned} c \int_0^T \int_{\{|x|>R\}} \omega_F((u - \hat{u})^+) |x|^{-1} \rho_\varepsilon &\leq C(\delta) \int_0^T \int_{\{|x|>R\}} (u - \hat{u})^+ + \delta cT \int_{\{|x|>R\}} \Omega_F^*(|x|^{-1}) \rho_\varepsilon \\ (17) \quad &\leq C(\delta) \int_0^T \int_{\{|x|>R\}} (u - \hat{u})^+ + \text{const } \delta; \end{aligned}$$

The left-hand side of (17) converges to zero as  $R \rightarrow \infty$ , uniformly in  $\varepsilon$ ; indeed, (15) yields

$$(18) \quad \inf_{\delta>0} \left( C(\delta) \int_0^T \int_{\{|x|>R\}} (u - \hat{u})^+ + \text{const } \delta \right) \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

Using (11) and (12) once more, from (15) we get

$$\begin{aligned} c\varepsilon \int_0^T \int_{\{|x|>R\}} \omega_\varphi((u - \hat{u})^+) |x|^{-2} \rho_\varepsilon &\leq \inf_{\delta>0} \left( \delta \int_0^T \int_{\{|x|>R\}} (u - \hat{u})^+ + r^{\delta,R}(\varepsilon) \right) \\ (19) \quad &\rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \text{ and then } R \rightarrow \infty. \end{aligned}$$

Now we let  $\varepsilon \rightarrow 0$  and  $R \rightarrow +\infty$  in (10); notice that  $\rho_\varepsilon$  converge to 1 and remains bounded. With the help of (17),(18) and (19), we deduce

$$(20) \quad \int_0^T \int_{\mathbb{R}^N} (u - \hat{u})^+ (-\mu_t) \leq \int_0^T \int_{\mathbb{R}^N} (f - \hat{f}) \operatorname{sign}^+(u - \hat{u}) \mu + \int_{\mathbb{R}^N} (u_0 - \hat{u}_0)^+ \mu(0).$$

The claim (9) follows readily from (20). Because the existence of an entropy solution is already known, this ends the proof of Theorem 2.

**Acknowledgement** The first author acknowledges the hospitality of the FST Mohammédia, of the CMA Oslo, and of IZKS, Bonn University (supported by the DFG project 436 RUS 113/895/0-1).

## REFERENCES

- [1] M. Maliki and A. Ouedraogo. Renormalized solution for a nonlinear anisotropic degenerated parabolic equation with nonlipschitz convection and diffusion flux functions, *Int. J. of Evol. Eq.*, to appear
- [2] B. Andreianov, Ph. Bénilan and S.N. Kruzhkov.  $L^1$  theory of scalar conservation law with continuous flux function. *J. Funct. Anal.*, 171 (2000), no.1, pp.15–33.
- [3] B. Andreianov and M. Maliki. On well-posedness for nonlinear diffusion equations in  $\mathbb{R}^N$ . In preparation.
- [4] F. Andreu, N. Igbida, J.M. Mazón and J. Toledo. Renormalized solutions for degenerate elliptic-parabolic problems with nonlinear dynamical boundary conditions and L1-data. *J. Diff. Equ.*, to appear.
- [5] Ph. Bénilan. *Équations d'évolution dans un espace de Banach quelconque et applications*. Thèse d'état, 1972.
- [6] Ph. Bénilan and S.N. Kruzhkov. Conservation laws with continuous flux functions. *NoDEA* V.3(1996), pp.395–419.
- [7] D. Blanchard and A. Porretta. Stefan problems with nonlinear diffusion and convection. *J. Diff. Eq.*, 210(2005), no.2, pp. 383–428.
- [8] J. Carrillo. Entropy solutions for nonlinear degenerate problems. *Arch. Rational Mech. Anal.*, 147(1999), pp. 269–361.
- [9] S.N. Kruzhkov. Generalized solutions of the Cauchy problem in the large for first-order nonlinear equations. *Dokl.Akad.Nauk URSS* V.187(1969), no.1, pp.29–32 (Russian); English tr. in *Soviet Math.Dokl.* V.10(1969).
- [10] S.N. Kruzhkov. Conservation laws with the infinite domain of dependence on initial data. In *Differential equations and control theory*, ed. V. Barbu, Pitman Research Notes in Mathematics, V.250(1991), pp.149–162.
- [11] S.N. Kruzhkov and P.A. Andrejanov. On the nonlocal theory of the Cauchy problem for quasi-linear equations of first order in the class of locally summable functions. (English. Russian original) *Sov. Math. Dokl.*, 16(1975), pp.16–20 ; translation from *Dokl. Akad. Nauk SSSR* 220(1975), pp.23–26.
- [12] S.N. Kruzhkov and F. Hil'debrand. The Cauchy problem for first order quasilinear equations in the case when the domain of dependence on the initial data is infinite. (Russian). Collection of articles dedicated to the memory of I.G. Petrovskii. *Vestnik Moskov. Univ. Ser. I Mat. Meh.*, V.29(1974), no.1, pp.93–100.
- [13] S.N. Kruzhkov and E.Yu. Panov. First-order quasilinear conservation laws with infinite initial data dependence area. *Dokl.Akad.Nauk URSS* V.314(1990), no.1, pp.79-84 (Russian); English tr. in *Soviet Math.Dokl.* V.42(1991), no.2, pp.316–321.
- [14] S.N. Kruzhkov and E.Yu. Panov. Osgood's type conditions for uniqueness of entropy solutions to Cauchy problem for quasilinear conservation laws of the first order. *Ann. Univ. Ferrara*, V.15(1994), pp.31–53.
- [15] M. Maliki and H. Touré. Uniqueness of entropy solutions for nonlinear degenerate parabolic problem. *J. Evol. Equ.* V.3(2003), no.4, pp.603–622.
- [16] E.Yu. Panov. *Generalized solutions of the Cauchy problem for quasilinear conservation laws*. PhD thesis, Moscow State Univ., 1991.
- [17] B. Perthame. Uniqueness and error estimates in first order quasilinear conservation laws via the kinetic entropy defect measure, *J. Math. Pures Appl.*, (9) V.77(1998), no. 10, pp.1055–1064.



(Boris Andreianov)  
LABORATOIRE DE MATHÉMATIQUES  
UNIVERSITÉ DE FRANCHE-COMTÉ  
16, ROUTE DE GRAY  
25030 BESANÇON CEDEX, FRANCE

*E-mail address:* `boris.andreianov@univ-fcomte.fr`

(Mohamed Maliki)  
ÉQUIPE MODÉLISATION, EDP ET ANALYSE NUMÉRIQUE  
FST MOHAMMÉDIA  
B.P. 146 MOHAMMÉDIA, MAROC

*E-mail address:* `mohamedmaliki@yahoo.fr`