



HAL
open science

The semilinear wave equation on asymptotically euclidean manifolds

Jean Francois Bony, Dietrich Häfner

► **To cite this version:**

Jean Francois Bony, Dietrich Häfner. The semilinear wave equation on asymptotically euclidean manifolds. *Comm. Partial Differential Equations*, 2010, 35 (1), pp.23-67. hal-00384722

HAL Id: hal-00384722

<https://hal.science/hal-00384722>

Submitted on 15 May 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE SEMILINEAR WAVE EQUATION ON ASYMPTOTICALLY EUCLIDEAN MANIFOLDS

JEAN-FRANÇOIS BONY AND DIETRICH HÄFNER

ABSTRACT. We consider the quadratically semilinear wave equation on $(\mathbb{R}^d, \mathbf{g})$, $d \geq 3$. The metric \mathbf{g} is non-trapping and approaches the Euclidean metric like $\langle x \rangle^{-\rho}$. Using Mourre estimates and the Kato theory of smoothness, we obtain, for $\rho > 0$, a Keel–Smith–Sogge type inequality for the linear equation. Thanks to this estimate, we prove long time existence for the nonlinear problem with small initial data for $\rho \geq 1$. Long time existence means that, for all $n > 0$, the life time of the solution is at least δ^{-n} , where δ is the size of the initial data in some appropriate Sobolev space. Moreover, for $d \geq 4$ and $\rho > 1$, we obtain global existence for small data.

CONTENTS

1. Introduction	1
2. The general setting	4
3. The wave equation and the Mourre estimate	8
4. Proof of the linear estimates	20
5. Proof of the nonlinear result	29
Appendix A. Regularity	31
Appendix B. Resolvent estimates at low energies	32
References	39

1. INTRODUCTION

This paper is devoted to the study of the quadratically semilinear wave equation on asymptotically Euclidean non-trapping Riemannian manifolds. We show global existence in dimension $d \geq 4$ and long time existence in dimension $d = 3$ for small data solutions. In Minkowski space, the semilinear wave equation has been thoroughly studied. Global existence is known in dimension $d \geq 4$ for small initial data (see Klainerman and Ponce [23] and references therein). Almost global existence in dimension $d = 3$ for small data was shown by John and Klainerman in [18]. Almost global means that the life time of a solution is at least $e^{1/\delta}$, where δ is the size of the initial data in some appropriate Sobolev space. Note that, in dimension $d = 3$, Sideris [31] has proved that global existence does not hold in general (see also John [17]).

2000 *Mathematics Subject Classification.* 35L70, 35P25, 58J45.

Key words and phrases. Scattering estimates, Mourre theory, semilinear wave equation.

In [20], Keel, Smith and Sogge give a new proof of the almost global existence result in dimension 3 using estimates of the form

$$(1.1) \quad (\ln(2+T))^{-1/2} \|\langle x \rangle^{-1/2} u'\|_{L^2([0,T] \times \mathbb{R}^3)} \lesssim \|u'(0, \cdot)\|_{L^2(\mathbb{R}^3)} + \int_0^T \|G(s, \cdot)\|_{L^2(\mathbb{R}^3)} ds,$$

and a certain Sobolev type estimate due to Klainerman (see [22]). Here u solves the wave equation $\square u = G$ in $[0, +\infty[\times \mathbb{R}^3$ and $u' = (\partial_t u, \partial_x u)$. They also treat the non-trapping obstacle case. In [21], similar results are obtained for the corresponding quasilinear equation. The obstacle case in which the trapped trajectories are of hyperbolic type is treated by Metcalfe and Sogge [24].

Alinhac shows an estimate similar to (1.1) on a curved background. In his papers [1] and [2], the metric is depending on and decaying in time. The results of Metcalfe and Tataru [26] imply estimates analogous to (1.1) for a space-time variable coefficients wave equation outside a star shaped obstacle (see also [25]). Outside the obstacle, their wave operator is a small perturbation of the wave operator in Minkowski space.

The common point of the papers cited so far is that they all use vector field methods. We use in this paper a somewhat different approach. We will show how estimates of type (1.1) follow from a Mourre estimate [27]. This method will permit us to consider non-trapping Riemannian metrics which are asymptotically Euclidean without requiring that they are everywhere a small perturbation of the Euclidean metric. We will suppose for simplicity that the metric is C^∞ , but a C^k approach should in principle be possible. Spectral methods for proving dispersive estimates were previously used by Burq. In [5], he obtains global Strichartz estimates for compactly supported non-trapping perturbations of the Euclidean case. In more complicated geometries, conjugate operators are probably not vector fields and it is perhaps worth trying to mix the classical vector field approach with the Mourre theory.

Let us now state our precise results. We consider the asymptotically Euclidean manifold $(\mathbb{R}^d, \mathfrak{g})$ with $d \geq 3$ and

$$\mathfrak{g} = \sum_{i,j=1}^d g_{i,j}(x) dx^i dx^j.$$

We suppose $g_{i,j}(x) \in C^\infty(\mathbb{R}^d)$ and, for some $\rho > 0$,

$$(H1) \quad \forall \alpha \in \mathbb{N}^d \quad \partial_x^\alpha (g_{i,j} - \delta_{i,j}) = \mathcal{O}(\langle x \rangle^{-|\alpha| - \rho}).$$

We also assume that

$$(H2) \quad \mathfrak{g} \text{ is non-trapping.}$$

Let $g(x) = (\det(\mathfrak{g}))^{1/4}$. The Laplace–Beltrami operator associated to \mathfrak{g} is given by

$$\Delta_{\mathfrak{g}} = \sum_{i,j} \frac{1}{g^2} \partial_i g^{i,j} g^2 \partial_j,$$

where $g^{i,j}(x)$ denotes the inverse metric. Let us consider the following unitary transform

$$\mathcal{V} : \begin{cases} L^2(\mathbb{R}^d, g^2 dx) & \longrightarrow L^2(\mathbb{R}^d, dx) \\ v & \longmapsto gv. \end{cases}$$

The transformation \mathcal{V} sends $-\Delta_{\mathbf{g}}$ to

$$P = -\mathcal{V}\Delta_{\mathbf{g}}\mathcal{V}^* = -\sum_{i,j} \frac{1}{g} \partial_i g^{i,j} g^2 \partial_j \frac{1}{g},$$

which is the operator we are interested in. Let $\tilde{\partial}_j := \partial_j g^{-1}$ and $\Omega = \Omega^{k,\ell} := x_k \partial_\ell - x_\ell \partial_k$ be the rotational vector fields. We consider the following semilinear wave equation

$$(1.2) \quad \begin{cases} \square_{\mathbf{g}} u = Q(u'), \\ (u|_{t=0}, \partial_t u|_{t=0}) = (u_0, u_1). \end{cases}$$

Here $\square_{\mathbf{g}} = \partial_t^2 + P$ and $Q(u')$ is a quadratic form in $u' = (\partial_t u, \tilde{\partial}_x u)$. For $x \in \mathbb{R}$, $[x]$ (resp. $\lceil x \rceil$) denotes the largest (resp. smallest) integer such that $[x] \leq x \leq \lceil x \rceil$. Our main result is the following theorem.

Theorem 1.1. *Assume hypotheses (H1) and (H2). Suppose $u_0, u_1 \in C_0^\infty(\mathbb{R}^d)$ and that, for $M = 2(\lceil \frac{d-1}{2} \rceil + 1)$, we have*

$$(1.3) \quad \sum_{|\alpha|+j \leq M+1} \|\partial_x^j \Omega^\alpha u_0\| + \sum_{|\alpha|+j \leq M} \|\partial_x^j \Omega^\alpha u_1\| \leq \delta.$$

i) Assume $d \geq 3$ and $\rho \geq 1$. For all $n > 0$, there exists a constant $\delta_n > 0$ such that, for $\delta \leq \delta_n$, the problem (1.2) has a unique solution $u \in C^\infty([0, T] \times \mathbb{R}^d)$ with

$$T = \delta^{-n}.$$

ii) Assume $d \geq 4$ and $\rho > 1$. For δ small enough, the problem (1.2) has a unique global solution $u \in C^\infty([0, +\infty[\times \mathbb{R}^d)$.

Remark 1.2. *One may consider more general nonlinearities. For example, the previous result holds for quadratic nonlinearities of the form $Q(x)(\langle x \rangle^{-\mu} u, u')$ with $\mu > 1$ and $\|\partial_x^\alpha Q(x)\| = \mathcal{O}(\langle x \rangle^{-|\alpha|})$. In particular, one can replace $Q(u')$ by $Q(\partial_t u, \partial_x u)$ or work with the wave equation before the transformation by \mathcal{V} . To prove this remark, it is enough to combine the proof of Theorem 1.1 with Lemma 4.2.*

The main ingredient of the proof are estimates of type (1.1). Let us therefore consider the corresponding linear equation. Let u be solution of

$$(1.4) \quad \begin{cases} (\partial_t^2 + P)u = G(s), \\ (u|_{t=0}, \partial_t u|_{t=0}) = (u_0, u_1). \end{cases}$$

With the notation

$$F_\mu^\varepsilon(T) = \begin{cases} T^{1-2\mu+2\varepsilon} & \mu \leq 1/2, \\ 1 & \mu > 1/2, \end{cases}$$

we have the following estimate.

Theorem 1.3. *Assume that (H1) and (H2) hold with $\rho > 0$ and let $0 < \mu \leq 1$. For all $\varepsilon > 0$, the solution of (1.4) satisfies*

$$(1.5) \quad \|\langle x \rangle^{-\mu} u'\|_{L^2([0, T] \times \mathbb{R}^d)} \lesssim \langle F_\mu^\varepsilon(T) \rangle^{1/2} \left(\|u'(0, \cdot)\|_{L^2(\mathbb{R}^d)} + \int_0^T \|G(s, \cdot)\|_{L^2(\mathbb{R}^d)} ds \right).$$

To prove the nonlinear theorem, it will be useful to have higher order estimates. To this purpose, let us put $\tilde{\Omega}^{k,\ell} = x_k \tilde{\partial}_\ell - x_\ell \tilde{\partial}_k$, $Z = \{\partial_t, \tilde{\partial}_x, \tilde{\Omega}\}$, $Y = \{\tilde{\partial}_x, \tilde{\Omega}\}$, $X = \{\tilde{\partial}_x\}$, where $\{\tilde{\Omega}\}$ (resp. $\{\tilde{\partial}_x\}$) are the collections of rotational vector fields (resp. partial derivatives with respect to space variables). Then, we have

Theorem 1.4. *Assume that (H1) and (H2) hold with $\rho > 1$ and let $N > 0$ and $1/2 \leq \mu \leq 1$. For all $\varepsilon > 0$, the solution of (1.4) satisfies*

$$(1.6) \quad \sup_{0 \leq t \leq T} \sum_{1 \leq k+j \leq N+1} \|\partial_t^k P^{j/2} u(t, \cdot)\|_{L^2(\mathbb{R}^d)} + \sum_{|\alpha| \leq N} \langle F_\mu^\varepsilon(T) \rangle^{-1} \|\langle x \rangle^{-\mu} Z^\alpha u'\|_{L^2([0,T] \times \mathbb{R}^d)} \\ \lesssim \sum_{|\alpha| \leq N} \left(\|(Z^\alpha u)'(0, \cdot)\|_{L^2(\mathbb{R}^d)} + \int_0^T \|Z^\alpha G(s, \cdot)\|_{L^2(\mathbb{R}^d)} ds \right).$$

Moreover, for $\rho = 1$, the same inequality holds with $\langle F_\mu^\varepsilon(T) \rangle^{-1}$ replaced by $\langle T \rangle^{-\varepsilon}$.

Remark 1.5. *i) Note that, in Theorem 1.1 and Theorem 1.4, $\rho \geq 1$ is required whereas Theorem 1.3 is valid under a general long range condition $\rho > 0$.*

ii) Theorem 1.3 and Theorem 1.4 remain valid if we replace u' by $(\partial_t u, P^{1/2} u)$.

The paper is organized in the following way. In Section 2, we show scattering estimates in a general setting. Section 3 is devoted to the Mourre estimate for the wave equation on our asymptotically Euclidean manifold. Using these results, we prove the estimates for the linear wave equation (Theorem 1.3 and Theorem 1.4) in Section 4. From these estimates, we deduce the nonlinear result in Section 5. Appendix A collects some regularity properties of operators and Appendix B contains low frequency resolvent estimates.

2. THE GENERAL SETTING

In this section, we obtain some abstract estimates which will be used to prove Theorem 1.3 and Theorem 1.4. These estimates are not specific to the wave equation and could help to show analogous estimates for other equations. The key ingredients are the limiting absorption principle and the Kato theory of smoothness.

We begin this section with the notion of regularity with respect to an operator. A full presentation of this theory can be found in the book of Amrein, A. Boutet de Monvel and Georgescu [3]. In Appendix A, we recall the properties which will be used in this paper.

Definition 2.1. *Let $(A, D(A))$ and $(H, D(H))$ be self-adjoint operators on a separable Hilbert space \mathcal{H} . The operator H is of class $C^k(A)$ for $k > 0$, if there is $z \in \mathbb{C} \setminus \sigma(H)$ such that*

$$\mathbb{R} \ni t \longrightarrow e^{itA} (H - z)^{-1} e^{-itA},$$

is C^k for the strong topology of $\mathcal{L}(\mathcal{H})$.

Let $H \in C^1(A)$ and $I \subset \sigma(H)$ be an open interval. We assume that A and H satisfy a Mourre estimate on I :

$$(2.1) \quad \mathbb{1}_I(H) i[H, A] \mathbb{1}_I(H) \geq \delta \mathbb{1}_I(H),$$

for some $\delta > 0$. As usual, we define the multi-commutators $\text{ad}_A^j B$ inductively by $\text{ad}_A^0 B = B$ and $\text{ad}_A^{j+1} B = [A, \text{ad}_A^j B]$.

Theorem 2.2 (Limiting absorption principle). *Let $H \in C^2(A)$ be such that $\text{ad}_A^j H$, $j = 1, 2$, are bounded on \mathcal{H} . Assume furthermore (2.1). Then, for all closed intervals $J \subset I$ and $\mu > 1/2$, there exists $C_{J,\mu} > 0$ such that*

$$(2.2) \quad \sup_{\substack{\text{Re } z \in J \\ \text{Im } z \neq 0}} \|\langle A \rangle^{-\mu} (H - z)^{-1} \langle A \rangle^{-\mu}\| \leq C_{J,\mu}.$$

If A and H depend on a parameter, the constant in the limiting absorption principle can be specified according to this parameter. In fact, mimicking the proof of [28], we obtain the following estimate.

Remark 2.3. *Assume that (2.1) holds uniformly and that $[H, A]$ is uniformly bounded. Then, for all closed intervals $J \subset I$ and $\mu > 1/2$,*

$$\sup_{\substack{\text{Re } z \in J \\ \text{Im } z \neq 0}} \|\langle A \rangle^{-\mu} (H - z)^{-1} \langle A \rangle^{-\mu}\| \leq \tilde{C}_{J,\mu} \langle \|\text{ad}_A^2 H\| \rangle^{\tilde{C}_{J,\mu}},$$

for some $\tilde{C}_{J,\mu} > 0$.

We now state a result of Kato [19] which says that, under the conclusions of Theorem 2.2, $\langle A \rangle^{-\mu} \mathbb{1}_J(H)$ is H -smooth. For the proof and more details, we refer to Theorem XIII.25 and Theorem XIII.30 of [30].

Theorem 2.4 (H -smoothness). *Let A and H be two self-adjoint operators satisfying (2.2). Then, for all closed intervals $J \subset I$ and $\mu > 1/2$,*

$$\int_{\mathbb{R}} \|\langle A \rangle^{-\mu} e^{-itH} \mathbb{1}_J(H)u\|^2 dt \leq 8C_{J,\mu} \|u\|^2,$$

for all $u \in \mathcal{H}$.

In the previous theorem, $C_{J,\mu}$ is the constant appearing in (2.2). By interpolation, we get

Corollary 2.5. *Assume (2.2). Then, for all closed intervals $J \subset I$ and $0 < \mu \leq 1/2$,*

$$\int_0^T \|\langle A \rangle^{-\mu} e^{-itH} \mathbb{1}_J(H)u\|^2 dt \leq M_{J,\mu,\varepsilon} T^{1-2\mu+\varepsilon} \|u\|^2,$$

for all $0 < \varepsilon < 2\mu$. Here,

$$M_{J,\mu,\varepsilon} = (8C_{J,\mu/(2\mu-\varepsilon)})^{2\mu-\varepsilon}.$$

Proof. Since e^{-itH} is unitary,

$$\int_0^T \|e^{-itH} \mathbb{1}_J(H)u\|^2 dt \leq T \|u\|^2.$$

Combining Theorem 2.4, the previous estimate and an interpolation argument, we get

$$\int_0^T \|\langle A \rangle^{-(1-\theta)\nu} e^{-itH} \mathbb{1}_J(H)u\|^2 dt \leq (8C_{J,\nu})^{1-\theta} T^\theta \|u\|^2.$$

for all $0 \leq \theta \leq 1$ and $\nu > 1/2$. Taking $\theta = 1 - 2\mu + \varepsilon \in [0, 1]$ (since $\varepsilon < 2\mu$) and $\nu = \mu/(1 - \theta) = \mu/(2\mu - \varepsilon) > 1/2$, the corollary follows. \square

We now study the non-homogeneous equation using the Fourier transform. Let $G(t) \in L^1_{\text{loc}}(\mathbb{R}_t; \mathcal{H})$ be such that $\text{supp } G \subset [0, +\infty[$. We consider the solution u of

$$(2.3) \quad \begin{cases} (i\partial_t - H)u(t) = \varphi(H)G(t), \\ u|_{t=0} = 0, \end{cases}$$

with $\varphi \in L^\infty(\mathbb{R})$ and $\text{supp } \varphi \subset J$. This means that

$$(2.4) \quad u(t) = -i \int_0^t e^{-i(t-s)H} \varphi(H)G(s) ds,$$

and then $u \in C^0(\mathbb{R}_t; \mathcal{H}) \cap \mathcal{S}'(\mathbb{R}_t; \mathcal{H})$.

Lemma 2.6. *Let A and H be two self-adjoint operators satisfying (2.2). Then, for all $\mu > 1/2$ and $\varphi \in C^1(\mathbb{R})$ satisfying $\|\varphi\|_\infty \leq 1$, $\|\varphi'\|_\infty \leq C_1$ and $\text{supp } \varphi \subset J$, we have*

$$\|\langle A \rangle^{-\mu} \varphi(H) (H - z)^{-1} \langle A \rangle^{-\mu}\| \leq C_{J,\mu} + C_1,$$

for all $z \in \mathbb{C} \setminus \mathbb{R}$.

Proof. Using Taylor's expansion formula, we have

$$\varphi(x) = \varphi(y) + (x - y) \int_0^1 \varphi'(tx + (1-t)y) dt,$$

and then

$$\varphi(H)(H - z)^{-1} = \varphi(\text{Re } z)(H - z)^{-1} + \int_0^1 \varphi'(tH + (1-t)\text{Re } z) dt (H - \text{Re } z)(H - z)^{-1}.$$

Using the spectral theorem, we obtain the following estimates:

$$\begin{aligned} \left\| \int_0^1 \varphi'(tH + (1-t)\text{Re } z) dt \right\| &\leq \int_0^1 \|\varphi'(tH + (1-t)\text{Re } z)\| dt \leq C_1, \\ \|(H - \text{Re } z)(H - z)^{-1}\| &\leq \sup_{x \in \mathbb{R}} |x(x - i\text{Im } z)^{-1}| \leq 1. \end{aligned}$$

Therefore, for $\text{Re } z \in J$, we have

$$\begin{aligned} \|\langle A \rangle^{-\mu} \varphi(H)(H - z)^{-1} \langle A \rangle^{-\mu}\| &\leq |\varphi(\text{Re } z)| \|\langle A \rangle^{-\mu} (H - z)^{-1} \langle A \rangle^{-\mu}\| + C_1 \|\langle A \rangle^{-\mu}\|^2 \\ &\leq C_{J,\mu} + C_1. \end{aligned}$$

On the other hand, for $\text{Re } z \notin J$, $\varphi(\text{Re } z) = 0$ and then

$$\|\langle A \rangle^{-\mu} \varphi(H)(H - z)^{-1} \langle A \rangle^{-\mu}\| \leq C_1 \|\langle A \rangle^{-\mu}\|^2 \leq C_1.$$

The two last estimates give the lemma. \square

Proposition 2.7. *Let A and H be two self-adjoint operators satisfying (2.2) and $\varphi \in C^1(\mathbb{R})$ as in Lemma 2.6. Then, for all $\mu > 1/2$ and $G(t) \in L^2(\mathbb{R}_t; D(\langle A \rangle^\mu))$ with $\text{supp } G \subset [0, +\infty[$, the solution u of (2.3) satisfies*

$$\int_0^\infty \|\langle A \rangle^{-\mu} u(t)\|^2 dt \leq (C_{J,\mu} + C_1)^2 \int_0^\infty \|\langle A \rangle^\mu G(t)\|^2 dt.$$

Proof. Let $u_\varepsilon = (1 + i\varepsilon H)^{-1}u$. From (2.4), $u_\varepsilon \in C^1(\mathbb{R}_t; \mathcal{H}) \cap C^0(\mathbb{R}_t; D(H)) \cap \mathcal{S}'(\mathbb{R}_t; D(H))$ and u_ε is the solution of the problem

$$(2.5) \quad \begin{cases} (i\partial_t - H)u_\varepsilon(t) = (1 + i\varepsilon H)^{-1}\varphi(H)G(t), \\ u_\varepsilon|_{t=0} = 0. \end{cases}$$

Since the support of the temperate distributions u_ε and G is in $[0, +\infty[$, their Fourier transforms are analytic in $\text{Im } z < 0$. Then, (2.5) gives, for $\text{Im } z < 0$,

$$(z - H)\widehat{u}_\varepsilon(z) = (1 + i\varepsilon H)^{-1}\varphi(H)\widehat{G}(z).$$

Then

$$\langle A \rangle^{-\mu}\widehat{u}_\varepsilon(z) = \langle A \rangle^{-\mu}(1 + i\varepsilon H)^{-1}\varphi(H)(z - H)^{-1}\langle A \rangle^{-\mu}\langle A \rangle^\mu\widehat{G}(z).$$

Since $\|\varphi(x)(1 + i\varepsilon x)^{-1}\|_\infty \leq \|\varphi\|_\infty$ and $\|\partial_x(\varphi(x)(1 + i\varepsilon x)^{-1})\|_\infty \leq \varepsilon\|\varphi\|_\infty + \|\varphi'\|_\infty$, Lemma 2.6 implies

$$\|\langle A \rangle^{-\mu}\widehat{u}_\varepsilon(z)\| \leq (C_{J,\mu} + C_1 + \varepsilon)\|\langle A \rangle^\mu\widehat{G}(z)\|.$$

Thus, for all $\delta > 0$, Plancherel's theorem gives

$$\int_0^{+\infty} e^{-\delta t}\|\langle A \rangle^{-\mu}u_\varepsilon\|^2 dt \leq (C_{J,\mu} + C_1 + \varepsilon)^2 \int_0^{+\infty} e^{-\delta t}\|\langle A \rangle^\mu G\|^2 dt.$$

Letting δ and ε go to 0, we get the proposition. \square

By interpolation, we also have

Corollary 2.8. *Assume the hypotheses of Proposition 2.7. Then, for all $0 < \mu \leq 1/2$ and $G(t) \in L^2(\mathbb{R}_t; D(\langle A \rangle^\mu))$ with $\text{supp } G \subset [0, +\infty[$,*

$$\int_0^T \|\langle A \rangle^{-\mu}u(t)\|^2 dt \leq N_{J,\mu,\varepsilon} T^{2(1-2\mu+\varepsilon)} \int_0^T \|\langle A \rangle^\mu G(t)\|^2 dt,$$

for all $0 < \varepsilon < 2\mu$. Here,

$$N_{J,\mu,\varepsilon} = (C_{J,\mu/(2\mu-\varepsilon)} + C_1)^{4\mu-2\varepsilon}.$$

Proof. Let $P_T : L^2([0, T]; \mathcal{H}) \longrightarrow L^2([0, T]; \mathcal{H})$ be the operator defined by

$$(P_T G)(t) = -i \int_0^t e^{-i(t-s)H} \varphi(H)G(s) ds.$$

Proposition 2.7 gives

$$\|\langle A \rangle^{-\nu} P_T G\|_{L^2([0, T]; \mathcal{H})} \leq (C_{J,\nu} + C_1) \|\langle A \rangle^\nu G\|_{L^2([0, T]; \mathcal{H})},$$

for $\nu > 1/2$. Moreover,

$$\begin{aligned} \|P_T G\|_{L^2([0, T]; \mathcal{H})} &\leq \sqrt{T} \sup_{t \in [0, T]} \left\| \int_0^t e^{-i(t-s)H} \varphi(H)G(s) ds \right\| \\ &\leq \sqrt{T} \sup_{t \in [0, T]} \sqrt{t} \left(\int_0^t \left\| e^{-i(t-s)H} \varphi(H)G(s) \right\|^2 ds \right)^{1/2} \\ &\leq T \|G\|_{L^2([0, T]; \mathcal{H})}. \end{aligned}$$

With these two estimates in mind, one can prove the corollary by mimicking the proof of Corollary 2.5. \square

3. THE WAVE EQUATION AND THE MOURRE ESTIMATE

In this section we will show a Mourre estimate for the wave equation on our asymptotically Euclidean manifold:

$$(3.1) \quad \begin{cases} (\partial_t^2 + P)u = 0, \\ u|_{t=0} = u_0, \quad \partial_t u|_{t=0} = u_1. \end{cases}$$

Recall that

$$(3.2) \quad P = - \sum_{i,j} \frac{1}{g} \partial_i g^{i,j} g^2 \partial_j \frac{1}{g},$$

is self-adjoint on $L^2(\mathbb{R}^d, dx)$ with domain $D(P) = H^2(\mathbb{R}^d)$. We define $H_c^k(\mathbb{R}^d)$ as the closure of $H^k(\mathbb{R}^d)$ with respect to the norm

$$\|u\|_{H_c^k}^2 = \sum_{j=1}^k \|P^{j/2}u\|^2.$$

Let $\mathcal{E} := H_c^1(\mathbb{R}^d) \oplus L^2(\mathbb{R}^d)$ with

$$\|(u_0, u_1)\|_{\mathcal{E}}^2 = \langle Pu_0, u_0 \rangle + \|u_1\|^2,$$

be the energy space associated to (3.1). The energy of (3.1) is clearly conserved:

$$\|(u(t), \partial_t u(t))\|_{\mathcal{E}} = \|(u_0, u_1)\|_{\mathcal{E}}.$$

We will rewrite (3.1) as a first order system

$$(3.3) \quad \begin{cases} i\partial_t f = Rf, \\ f|_{t=0} = (u_0, u_1), \end{cases}$$

with

$$R = \begin{pmatrix} 0 & i \\ -iP & 0 \end{pmatrix}.$$

The operator R is self-adjoint on \mathcal{E} with domain $D(R) = H_c^2(\mathbb{R}^d) \oplus H^1(\mathbb{R}^d)$. Let $\mathcal{L} = L^2(\mathbb{R}^d) \oplus L^2(\mathbb{R}^d)$. It is useful to introduce the following unitary transform:

$$U : \mathcal{E} \longrightarrow \mathcal{L}, \quad U = \frac{1}{\sqrt{2}} \begin{pmatrix} P^{1/2} & i \\ P^{1/2} & -i \end{pmatrix},$$

which satisfies

$$U^* = U^{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} P^{-1/2} & P^{-1/2} \\ -i & i \end{pmatrix} \quad \text{and} \quad L = URU^* = \frac{1}{\sqrt{2}} \begin{pmatrix} P^{1/2} & 0 \\ 0 & -P^{1/2} \end{pmatrix}.$$

The operator $(L, D(L) = H^1(\mathbb{R}^d) \oplus H^1(\mathbb{R}^d))$ is self-adjoint. In order to establish a Mourre estimate for L , it is sufficient to establish a Mourre estimate for $P^{1/2}$. We divide this section into the study of the low, the intermediate and the high frequency part.

3.1. Low frequency Mourre estimate.

For low frequencies, we will make a dyadic decomposition and use a conjugate operator specific to each part of the decomposition. In this section, we will obtain a Mourre estimate for each part. For $\lambda \geq 1$, we set

$$(3.4) \quad \mathcal{A}_\lambda = \varphi(\lambda P)A_0\varphi(\lambda P),$$

where

$$A_0 = \frac{1}{2}(xD + Dx), \quad D(A_0) = \{u \in L^2(\mathbb{R}^d); A_0u \in L^2(\mathbb{R}^d)\},$$

is the generator of dilations and $\varphi \in C_0^\infty(]0, +\infty[;]0, +\infty[)$ satisfies $\varphi(x) > \delta > 0$ on some open bounded interval $I \subset]0, +\infty[$.

For the various estimates that we will establish in this section, the following formula for the square root of an operator will be useful. Making a change of contour and using the Cauchy formula, one can show that

$$(3.5) \quad \sigma^{-1/2} = \frac{1}{\pi} \int_0^{+\infty} s^{-1/2}(s + \sigma)^{-1} ds,$$

for $\sigma \neq 0$. Therefore, the functional calculus gives

$$(3.6) \quad \varphi(\lambda P)P^{1/2} = \frac{1}{\pi} \int_0^{+\infty} s^{-1/2}\varphi(\lambda P)P(s + P)^{-1} ds.$$

It is well known that $P \in C^1(A_0)$. In particular, $\varphi(\lambda P) : D(A_0) \rightarrow D(A_0)$ and \mathcal{A}_λ is well defined on $D(A_0)$. Its closure, again denoted \mathcal{A}_λ , is self-adjoint (see [3, Theorem 6.2.5, Lemma 7.2.15]).

Proposition 3.1. *i) We have $(\lambda P)^{1/2} \in C^2(\mathcal{A}_\lambda)$. The commutators $\text{ad}_{\mathcal{A}_\lambda}^j(\lambda P)^{1/2}$, $j = 1, 2$, can be extended to bounded operators and we have, uniformly in λ ,*

$$(3.7) \quad \|[\mathcal{A}_\lambda, (\lambda P)^{1/2}]\| \lesssim 1,$$

$$(3.8) \quad \|\text{ad}_{\mathcal{A}_\lambda}^2(\lambda P)^{1/2}\| \lesssim \begin{cases} 1 & \rho > 1, \\ \lambda^\varepsilon & \rho \leq 1, \end{cases}$$

where $\varepsilon > 0$ can be chosen arbitrary small.

ii) For λ large enough, we have the following Mourre estimate:

$$(3.9) \quad \mathbb{1}_I(\lambda P)[i(\lambda P)^{1/2}, \mathcal{A}_\lambda]\mathbb{1}_I(\lambda P) \geq \frac{\delta^2 \sqrt{\inf I}}{2} \mathbb{1}_I(\lambda P).$$

iii) For $0 \leq \mu \leq 1$ and $\psi \in C_0^\infty(]0, +\infty[)$, we have

$$(3.10) \quad \| |\mathcal{A}_\lambda|^\mu \langle x \rangle^{-\mu} \| \lesssim \lambda^{-\mu/2+\varepsilon},$$

$$(3.11) \quad \| \langle \mathcal{A}_\lambda \rangle^\mu \psi(\lambda P) \langle x \rangle^{-\mu} \| \lesssim \lambda^{-\mu/2+\varepsilon},$$

for all $\varepsilon > 0$.

The rest of this section will be devoted to the proof of the above proposition, which will be divided into several lemmas.

Lemma 3.2. *We have $(\lambda P)^{1/2} \in C^1(\mathcal{A}_\lambda)$.*

The proof of Lemma 3.2 is analogous to the proof of [13, Lemme 3.3]. In this lemma, it is shown that $((\lambda P)^{1/2}, \mathcal{A}_\lambda)$ fulfill the original conditions of Mourre which imply the C^1 regularity. We now have to estimate the commutators. First note that

$$[i(\lambda P)^{1/2}, \mathcal{A}_\lambda] = i\lambda^{1/2}\varphi(\lambda P)[P^{1/2}, A_0]\varphi(\lambda P).$$

Using formula (3.6), we find

$$(3.12) \quad \varphi(\lambda P)[P^{1/2}, A_0]\varphi(\lambda P) = \frac{1}{\pi} \int_0^{+\infty} s^{-1/2}\varphi(\lambda P)[P(s+P)^{-1}, A_0]\varphi(\lambda P) ds,$$

with

$$[P(s+P)^{-1}, A_0] = -s[(s+P)^{-1}, A_0] = s(s+P)^{-1}[P, A_0](s+P)^{-1}.$$

A direct calculation gives

$$(3.13) \quad \begin{aligned} [P, A_0] &= -i \sum_{j,k} g^{-1} D_j \left(2g^2 g^{j,k} - \sum_{\ell} x_{\ell} \partial_{\ell} (g^2 g^{j,k}) \right) D_k g^{-1} \\ &\quad - i \sum_{\ell} g^{-2} x_{\ell} (\partial_{\ell} g) \sum_{j,k} D_j g^2 g^{j,k} D_k g^{-1} - i \sum_{j,k} g^{-1} D_j g^2 g^{j,k} D_k g^{-2} \sum_{\ell} x_{\ell} (\partial_{\ell} g) \\ &= -2iP + i \sum_{j,k} g^{-1} D_j \left(\sum_{\ell} x_{\ell} \partial_{\ell} (g^2 g^{j,k}) - 2g^{j,k} g \sum_{\ell} x_{\ell} (\partial_{\ell} g) \right) D_k g^{-1} \\ &\quad + \sum_{j,k,\ell} g^{-1} \partial_j (g^{-1} x_{\ell} (\partial_{\ell} g)) g^2 g^{j,k} D_k g^{-1} - \sum_{j,k,\ell} g^{-1} D_j g^2 g^{j,k} \partial_k (g^{-1} x_{\ell} (\partial_{\ell} g)) g^{-1} \\ &= -2iP - 2i \sum_{j,k} g^{-1} D_j a_{j,k} D_k g^{-1} + 2i \sum_k b_k D_k g^{-1} - 2i \sum_j g^{-1} D_j b_j, \end{aligned}$$

where $\partial_x^\alpha a_{j,k} = \mathcal{O}(\langle x \rangle^{-\rho-|\alpha|})$ and $\partial_x^\alpha b = \mathcal{O}(\langle x \rangle^{-\rho-1-|\alpha|})$ by (H1). In the following, a term r_j , $j \in \mathbb{N}$, will denote a smooth function such that

$$(3.14) \quad \partial_x^\alpha r_j(x) = \mathcal{O}(\langle x \rangle^{-\rho-j-|\alpha|}).$$

Moreover, to clarify the statement, we will not write the sums over j, k and j, k and replace the remainder terms in (3.13) by $\tilde{\partial}^* r_0 \tilde{\partial}$, $\tilde{\partial}^* r_1$ and $r_1 \tilde{\partial}^*$. Then,

$$(3.15) \quad [P, A_0] = -2i(P + \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}^*).$$

and (3.12) becomes

$$\begin{aligned} \varphi(\lambda P)[P^{1/2}, A_0]\varphi(\lambda P) &= -\frac{2i}{\pi} \int_0^{+\infty} s^{1/2}\varphi(\lambda P)(s+P)^{-1} \\ &\quad (P + \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}^*)(s+P)^{-1}\varphi(\lambda P) ds. \end{aligned}$$

Proceeding as in (3.6), one can show that

$$\int_0^{+\infty} s^{1/2}\varphi^2(\lambda P)P(s+P)^{-2} ds = \frac{\pi}{2}\varphi^2(\lambda P)P^{1/2}.$$

Then, we finally obtain

$$(3.16) \quad [i(\lambda P)^{1/2}, \mathcal{A}_\lambda] = (\lambda P)^{1/2}\varphi^2(\lambda P) + R,$$

with

$$(3.17) \quad R = \frac{2}{\pi}\lambda^{1/2} \int_0^{+\infty} s^{1/2}\varphi(\lambda P)(s+P)^{-1}(\tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}^*)(s+P)^{-1}\varphi(\lambda P) ds.$$

The remainder term R can be estimated in the following way.

Lemma 3.3. *Assume $0 \leq \gamma \leq d/4$. Then, we have*

$$\|Ru\| \lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{-\min(\rho, d/2)+2\gamma+\varepsilon} u\|,$$

for all $\varepsilon > 0$.

Proof. First, we write

$$\varphi(\lambda P)(s+P)^{-1}\tilde{\partial}^* = \lambda^{-1/2}(s+P)^{-1}\varphi(\lambda P)(\lambda P)^{1/2}P^{-1/2}\tilde{\partial}^*.$$

Using Lemma B.13 and the functional calculus, this term can be estimated by

$$(3.18) \quad \|\varphi(\lambda P)(s+P)^{-1}\tilde{\partial}^*\| \lesssim \lambda^{-1/2} \|(s+P)^{-1}\mathbb{1}_{\text{supp } \varphi(\lambda P)}\| \lesssim \lambda^{-1/2}(s+\lambda^{-1})^{-1}.$$

Moreover, applying Lemma B.12 (with $\beta = 0$ and $\gamma = 1/2 \leq d/4$), we get

$$(3.19) \quad \|\varphi(\lambda P)(s+P)^{-1}\tilde{\partial}^*u\| \lesssim (s+\lambda^{-1})^{-1}\lambda^{-1+\varepsilon}\|\langle x \rangle u\|,$$

$$(3.20) \quad \|\varphi(\lambda P)(s+P)^{-1}u\| \lesssim (s+\lambda^{-1})^{-1}\lambda^{-1/2+\varepsilon}\|\langle x \rangle u\|.$$

On the other hand, we write, for $k \in \mathbb{N}$,

$$(s+P)^{-1}\varphi(\lambda P) = \lambda(\lambda s + \lambda P)^{-1}\psi(\lambda P)(\lambda P + 1)^{-k},$$

with $\psi(\sigma) = \varphi(\sigma)(\sigma + 1)^k \in C_0^\infty(]0, +\infty[)$. Using the spectral theorem, we have

$$(3.21) \quad (s+P)^{-1}\varphi(\lambda P) = \frac{\lambda}{\pi} \int (\lambda s + z)^{-1} \bar{\partial} \tilde{\psi}(z) (\lambda P - z)^{-1} (\lambda P + 1)^{-k} L(dz),$$

where $\tilde{\psi} \in C_0^\infty(\mathbb{C})$ is an almost analytic extension of ψ . From the form of φ , one can always assume that $\text{supp } \tilde{\psi} \subset \{z \in \mathbb{C}; \text{Re } z > \tilde{\varepsilon} > 0\}$. In particular,

$$(3.22) \quad |(\lambda s + z)^{-1}| \lesssim (\lambda s + 1)^{-1},$$

uniformly for $z \in \text{supp } \tilde{\psi}$.

Using Proposition B.9, Lemma B.10 and Remark B.11, we obtain

$$(3.23) \quad \begin{aligned} \|\langle x \rangle^{-\rho}(\lambda P - z)^{-1}u\| &\lesssim \|\langle x \rangle^{-\rho}(\lambda P - z)^{-1}\langle x \rangle^{\min(\rho, d/2)-\varepsilon}\langle x \rangle^{-\min(\rho, d/2)+\varepsilon}u\| \\ &\lesssim \frac{1}{|\text{Im } z|^C} \|\langle x \rangle^{-\min(\rho, d/2)+\varepsilon}u\|, \end{aligned}$$

and

$$(3.24) \quad \begin{aligned} \|\langle x \rangle^{-\rho}\tilde{\partial}(\lambda P - z)^{-1}u\| &\lesssim \|\langle x \rangle^{-\rho}\tilde{\partial}(\lambda P - z)^{-1}\langle x \rangle^{\min(\rho, d/2)-\varepsilon}\langle x \rangle^{-\min(\rho, d/2)+\varepsilon}u\| \\ &\lesssim \frac{\lambda^{-1/2}}{|\text{Im } z|^C} \|\langle x \rangle^{-\min(\rho, d/2)+\varepsilon}u\|, \end{aligned}$$

for all $\varepsilon > 0$.

Let $\gamma \leq d/4$ and fix $k > \gamma + 2$. Applying k times Proposition B.9, we get

$$(3.25) \quad \|\langle x \rangle^{-\min(\rho, d/2)+\varepsilon}(\lambda P + 1)^{-k}u\| \lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{-\min(\rho, d/2)+2\gamma+2\varepsilon}u\|,$$

for all $\varepsilon > 0$.

The formula (3.21) and the estimates (3.22), (3.23), (3.25) imply

$$\begin{aligned}
& \|\langle x \rangle^{-\rho} (s + P)^{-1} \varphi(\lambda P) u\| \\
& \lesssim \lambda (\lambda s + 1)^{-1} \int |\bar{\partial} \tilde{\psi}(z)| \|\langle x \rangle^{-\rho} (\lambda P - z)^{-1} (\lambda P + 1)^{-k} u\| L(dz) \\
& \lesssim \lambda (\lambda s + 1)^{-1} \left(\int |\operatorname{Im} z|^{-C} |\bar{\partial} \tilde{\psi}(z)| L(dz) \right) \|\langle x \rangle^{-\min(\rho, d/2) + \varepsilon} (\lambda P + 1)^{-k} u\| \\
(3.26) \quad & \lesssim \lambda^{1-\gamma+\varepsilon} (\lambda s + 1)^{-1} \|\langle x \rangle^{-\min(\rho, d/2) + 2\gamma + 2\varepsilon} u\|,
\end{aligned}$$

for all $\varepsilon > 0$. The same way, using (3.24) instead of (3.23), we obtain

$$(3.27) \quad \|\langle x \rangle^{-\rho} \tilde{\partial} (s + P)^{-1} \varphi(\lambda P) u\| \lesssim \lambda^{1/2-\gamma+\varepsilon} (\lambda s + 1)^{-1} \|\langle x \rangle^{-\min(\rho, d/2) + 2\gamma + 2\varepsilon} u\|,$$

for all $\varepsilon > 0$.

Let R_1 be the term of (3.17) with $\tilde{\partial}^* r_0 \tilde{\partial}$. Using $r_0 = \mathcal{O}(\langle x \rangle^{-\rho})$, (3.18) and (3.27), we get

$$\begin{aligned}
\|R_1 u\| & \lesssim \lambda^{1/2-\gamma+\varepsilon} \left(\int_0^{+\infty} s^{1/2} (s + \lambda^{-1})^{-1} (\lambda s + 1)^{-1} ds \right) \|\langle x \rangle^{-\min(\rho, d/2) + 2\gamma + 2\varepsilon} u\| \\
& \lesssim \lambda^{-\gamma+\varepsilon} \left(\int_0^{+\infty} s^{1/2} (s + 1)^{-1} (s + 1)^{-1} ds \right) \|\langle x \rangle^{-\min(\rho, d/2) + 2\gamma + 2\varepsilon} u\| \\
(3.28) \quad & \lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{-\min(\rho, d/2) + 2\gamma + 2\varepsilon} u\|,
\end{aligned}$$

for all $\varepsilon > 0$. The same estimate can be proved for the term of (3.17) with $\tilde{\partial}^* r_1$ (resp. $r_1 \tilde{\partial}$) from $r_1 = \mathcal{O}(\langle x \rangle^{-\rho-1})$, (3.19) and (3.26) (resp. (3.20) and (3.27)). \square

Lemma 3.4. *For all $\varepsilon > 0$,*

$$[[\langle \lambda P \rangle^{1/2}, \mathcal{A}_\lambda], \mathcal{A}_\lambda] = \mathcal{O}(\lambda^\varepsilon).$$

Remark 3.5. *If we assume $\rho > 1$, Lemma 3.4 can be proved more simply. In fact, Lemma 3.3 and Lemma 3.6 give $\|Ru\| \lesssim \|\langle x \rangle^{-1} u\|$ and $\|\mathcal{A}_\lambda u\| \lesssim \|\langle x \rangle u\|$. Using $R^* = R$, these estimates imply Lemma 3.4.*

Proof. • We start with the commutator between \mathcal{A}_λ and the first term on the right hand side of (3.16). Let $\psi(\sigma) = \sigma \varphi^2(\sigma^2)$ and $\tilde{\psi}$ be an almost analytic extension of ψ . Then, we have

$$\begin{aligned}
[(\lambda P)^{1/2} \varphi^2(\lambda P), \mathcal{A}_\lambda] & = [\psi((\lambda P)^{1/2}), \mathcal{A}_\lambda] \\
& = -\frac{1}{\pi} \int \bar{\partial} \tilde{\psi}(z) ((\lambda P)^{1/2} - z)^{-1} [(\lambda P)^{1/2}, \mathcal{A}_\lambda] ((\lambda P)^{1/2} - z)^{-1} L(dz).
\end{aligned}$$

From (3.16) and Lemma 3.3, $[(\lambda P)^{1/2}, \mathcal{A}_\lambda]$ is uniformly bounded. Therefore, the commutator $[(\lambda P)^{1/2} \varphi^2(\lambda P), \mathcal{A}_\lambda]$ is also uniformly bounded.

• We now study the commutator between \mathcal{A}_λ and R defined in (3.16). One can write

$$(3.29) \quad [\mathcal{A}_\lambda, R] = [\varphi(\lambda P), R] A_0 \varphi(\lambda P) + \varphi(\lambda P) [A_0, R] \varphi(\lambda P) + \varphi(\lambda P) A_0 [\varphi(\lambda P), R],$$

that we note $[\mathcal{A}_\lambda, R] = S_1 + S_2 + S_3$. Since $S_3 = -S_1^*$, we only study the two first terms.

★ With (3.17) in mind, the operator S_1 can be written

$$(3.30) \quad S_1 = \frac{2}{\pi} \lambda^{1/2} \int_0^{+\infty} s^{1/2} \varphi(\lambda P) (s + P)^{-1} [\varphi(\lambda P), \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] (s + P)^{-1} \varphi(\lambda P) A_0 \varphi(\lambda P) ds,$$

where

$$(3.31) \quad \begin{aligned} & [\varphi(\lambda P), \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] \\ &= -\frac{\lambda}{\pi} \int \bar{\partial} \tilde{\varphi}(z) (\lambda P - z)^{-1} [P, \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] (\lambda P - z)^{-1} L(dz), \end{aligned}$$

and $\tilde{\varphi}$ is an almost analytic extension of φ . A direct calculation gives

$$[P, \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] = \tilde{\partial}^* \tilde{\partial}^* r_1 \tilde{\partial} + \tilde{\partial}^* r_2 \tilde{\partial} + \tilde{\partial}^* r_3 + r_3 \tilde{\partial},$$

with the convention of (3.14). For the first term in this equality, we write

$$\begin{aligned} \tilde{\partial}^* \tilde{\partial}^* r_1 \tilde{\partial} &= \tilde{\partial}^* (\lambda P + 1) (\lambda P + 1)^{-1} \tilde{\partial}^* r_1 \tilde{\partial} \\ &= (\lambda P + 1) \tilde{\partial}^* (\lambda P + 1)^{-1} \tilde{\partial}^* r_1 \tilde{\partial} - \lambda [P, \tilde{\partial}^*] (\lambda P + 1)^{-1} \tilde{\partial}^* r_1 \tilde{\partial}. \end{aligned}$$

As before, a direct calculation gives

$$[P, \tilde{\partial}^*] = \tilde{\partial}^* r_1 \tilde{\partial} + \tilde{\partial}^* r_2,$$

with the usual decay on r_1 and r_2 . Summing up, we get

$$\begin{aligned} [P, \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] &= (\lambda P + 1) \tilde{\partial}^* (\lambda P + 1)^{-1} \tilde{\partial}^* r_1 \tilde{\partial} \\ &\quad - \lambda (\tilde{\partial}^* r_1 \tilde{\partial} + \tilde{\partial}^* r_2) (\lambda P + 1)^{-1} \tilde{\partial}^* r_1 \tilde{\partial} + \tilde{\partial}^* r_2 \tilde{\partial} + \tilde{\partial}^* r_3 + r_3 \tilde{\partial}^*. \end{aligned}$$

Applying Lemma B.10 (with $\beta = 1$ and $\gamma = 0$ satisfying $\gamma + \beta/2 \leq d/4$), Lemma B.10 (with $\beta = 0$ and $\gamma = 1/2$), Remark B.11 and Lemma B.13, one can show that all the terms (say \tilde{r}) of the last equation, with the exceptions of $\tilde{\partial}^* r_3$ and $r_3 \tilde{\partial}$, satisfy

$$(3.32) \quad \|(\lambda P + 1)^{-1} (\lambda P - z)^{-1} \tilde{r} (\lambda P - z)^{-1} u\| \lesssim \frac{\lambda^{-3/2+\varepsilon}}{|\operatorname{Im} z|^C} \|\langle x \rangle^{-1} u\|,$$

for all $\varepsilon > 0$. Writing

$$(\lambda P + 1)^{-1} (\lambda P - z)^{-1} \tilde{\partial}^* r_3 (\lambda P - z)^{-1} = (\lambda P + 1)^{-1} \langle x \rangle^{-1} \langle x \rangle (\lambda P - z)^{-1} \tilde{\partial}^* r_3 (\lambda P - z)^{-1},$$

and using Proposition B.9 (with $\beta = 0$ and $\gamma = 1/2$), Lemma B.10 (with $\beta = 1$ and $\gamma = 1/4$) and Proposition B.9 (with $\beta = 1$ and $\gamma = 1/4$), we get

$$(3.33) \quad \|(\lambda P + 1)^{-1} (\lambda P - z)^{-1} \tilde{\partial}^* r_3 (\lambda P - z)^{-1} u\| \lesssim \frac{\lambda^{-3/2+\varepsilon}}{|\operatorname{Im} z|^C} \|\langle x \rangle^{-1} u\|.$$

Note that, in the case $d = 3$, we have $\gamma + \beta/2 = d/4$. It is why we can not use the additional decay $\langle x \rangle^{-\rho}$ and loose λ^ε . In a similar manner, Proposition B.9 (with $\beta = 0$ and $\gamma = 3/4$) and Lemma B.10 (with $\beta = 1$ and $\gamma = 1/4$) imply

$$(3.34) \quad \|(\lambda P + 1)^{-1} (\lambda P - z)^{-1} r_3 \tilde{\partial} (\lambda P - z)^{-1} u\| \lesssim \frac{\lambda^{-3/2+\varepsilon}}{|\operatorname{Im} z|^C} \|\langle x \rangle^{-1} u\|.$$

Combining the estimates (3.32), (3.33) and (3.34) with the identity (3.31), we obtain

$$(3.35) \quad \|(\lambda P + 1)^{-1} [\varphi(\lambda P), \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] u\| \lesssim \lambda^{-1/2+\varepsilon} \|\langle x \rangle^{-1} u\|.$$

From the form of A_0 , we have

$$A_0 = -ig^{-1}\partial xg + a,$$

with $\partial_x^\alpha a(x) = \mathcal{O}(\langle x \rangle^{-|\alpha|})$. As in (3.21), we write

$$(3.36) \quad (s+P)^{-1}\varphi(\lambda P) = \frac{\lambda}{\pi} \int (\lambda s+z)^{-1} \bar{\partial} \tilde{\varphi}(z) (\lambda P-z)^{-1} L(dz).$$

The above expression for A_0 , together with Lemma B.10 (with $\beta = 1$ and $\gamma = 0$), Proposition B.9 (with $\beta = 0$ and $\gamma = 1/2$) and Remark B.11, gives

$$\|\langle x \rangle^{-1} (\lambda P - z)^{-1} A_0\| \lesssim \frac{\lambda^{-1/2+\varepsilon}}{|\operatorname{Im} z|^C}.$$

Then, (3.36) (see also (3.22)) implies

$$(3.37) \quad \|\langle x \rangle^{-1} (s+P)^{-1} \varphi(\lambda P) A_0\| \lesssim \lambda^{1/2+\varepsilon} (\lambda s+1)^{-1},$$

for all $\varepsilon > 0$.

Eventually, using the identity (3.30), the functional calculus and the estimates (3.35) and (3.37), we obtain

$$(3.38) \quad \begin{aligned} \|S_1\| &\lesssim \lambda^{1/2} \int_0^{+\infty} s^{1/2} (s+\lambda^{-1})^{-1} \lambda^{-1/2+\varepsilon} \lambda^{1/2+\varepsilon} (\lambda s+1)^{-1} ds \\ &\lesssim \lambda^{3/2+2\varepsilon} \int_0^{+\infty} s^{1/2} (\lambda s+1)^{-2} ds \lesssim \lambda^{2\varepsilon} \int_0^{+\infty} t^{1/2} (t+1)^{-2} dt \lesssim \lambda^{2\varepsilon}, \end{aligned}$$

for all $\varepsilon > 0$.

★ We now study $S_2 = \varphi(\lambda P)[A_0, R]\varphi(\lambda P)$. Using (3.36), S_2 can be decomposed as

$$(3.39) \quad S_2 = T_1 + T_2 + T_3,$$

with

$$\begin{aligned} T_1 &= \frac{2}{\pi^2} \lambda^{5/2} \iint_0^{+\infty} s^{1/2} \bar{\partial} \tilde{\varphi}(z) (\lambda s+z)^{-1} \varphi(\lambda P) (\lambda P-z)^{-1} [P, A_0] (\lambda P-z)^{-1} \\ &\quad (\tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}) (s+P)^{-1} \varphi^2(\lambda P) ds L(dz), \\ T_2 &= \frac{2}{\pi} \lambda^{1/2} \int_0^{+\infty} s^{1/2} \varphi^2(\lambda P) (s+P)^{-1} [A_0, \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] (s+P)^{-1} \varphi^2(\lambda P) ds, \\ T_3 &= \frac{2}{\pi^2} \lambda^{5/2} \iint_0^{+\infty} s^{1/2} \bar{\partial} \tilde{\varphi}(z) (\lambda s+z)^{-1} \varphi^2(\lambda P) (s+P)^{-1} (\tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}) \\ &\quad (\lambda P-z)^{-1} [P, A_0] (\lambda P-z)^{-1} \varphi(\lambda P) ds L(dz). \end{aligned}$$

Since $T_3^* = T_1$, we only treat T_1 and T_2 .

From (3.15), we know that

$$[P, A_0] = -2iP + \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}.$$

Let \tilde{r} be a term of the last equation and let \hat{r} be a term of the sum

$$\tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}.$$

Then the functional calculus, Proposition B.9 (with $\beta = 0$ and $\gamma = 1/2$), Lemma B.10 (with $\beta = 0$ and $\gamma = 1/2$), Remark B.11 and Lemma B.13 show that

$$\|(\lambda P - z)^{-1} \tilde{r} (\lambda P - z)^{-1} \tilde{r} (\lambda P + 1)^{-1}\| \lesssim \frac{\lambda^{-2+\varepsilon}}{|\operatorname{Im} z|^C},$$

for all $\varepsilon > 0$. Then, T_1 becomes

$$(3.40) \quad \|T_1\| \lesssim \lambda^{5/2} \int_0^{+\infty} s^{1/2} (1 + \lambda s)^{-1} \lambda^{-2+\varepsilon} (s + \lambda^{-1})^{-1} ds \lesssim \lambda^\varepsilon,$$

for all $\varepsilon > 0$.

A direct calculation shows that

$$[A_0, \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial}] = \tilde{\partial}^* r_0 \tilde{\partial} + \tilde{\partial}^* r_1 + r_1 \tilde{\partial} + r_2.$$

From Proposition B.9 (with $\beta = 0$ and $\gamma = 1/2$) and Lemma B.13, every term (say \tilde{r}) of the previous equation satisfies

$$\|(\lambda P + 1)^{-1} \tilde{r} (\lambda P + 1)^{-1}\| \lesssim \lambda^{-1+\varepsilon}.$$

Then, using the spectral theorem, T_2 fulfills

$$(3.41) \quad \|T_2\| \lesssim \lambda^{1/2} \int_0^{+\infty} s^{1/2} (s + \lambda^{-1})^{-1} \lambda^{-1+\varepsilon} (s + \lambda^{-1})^{-1} ds \lesssim \lambda^\varepsilon,$$

Combining (3.39) with the estimates (3.40), (3.41) and $T_3^* = T_1$, we obtain

$$(3.42) \quad \|S_2\| \lesssim \lambda^\varepsilon.$$

★ The lemma follows from (3.29), (3.38), (3.42) and $S_3 = -S_1^*$. □

Lemma 3.6. *Let $0 \leq \mu \leq 1$ and $\psi \in C_0^\infty([0, +\infty[)$. Then, we have*

$$(3.43) \quad \|\mathcal{A}_\lambda |^\mu \langle x \rangle^{-\mu}\| \lesssim \lambda^{-\mu/2+\varepsilon},$$

$$(3.44) \quad \|\langle \mathcal{A}_\lambda \rangle^\mu \psi(\lambda P) \langle x \rangle^{-\mu}\| \lesssim \lambda^{-\mu/2+\varepsilon},$$

for all $\varepsilon > 0$.

Proof. From (3.4), we have

$$(3.45) \quad \mathcal{A}_\lambda = \varphi(\lambda P) \left(g^{-1} D x g + i \left(\frac{d}{2} + g^{-1} x (\partial g) \right) \right) \varphi(\lambda P).$$

Lemma B.13 gives

$$\varphi(\lambda P) g^{-1} D = P^{1/2} \varphi(\lambda P) P^{-1/2} g^{-1} D = \mathcal{O}(\lambda^{-1/2}).$$

Moreover, Lemma B.12 (with $\beta = 1$ and $\gamma = 0$) implies

$$\|\langle x \rangle \varphi(\lambda P) u\| \lesssim \lambda^\varepsilon \|\langle x \rangle u\|,$$

for all $\varepsilon > 0$. Summing up the previous estimates, we get

$$(3.46) \quad \|\varphi(\lambda P) g^{-1} D x g \varphi(\lambda P) u\| \lesssim \lambda^{-1/2+\varepsilon} \|\langle x \rangle u\|,$$

for all $\varepsilon > 0$. Using Lemma B.12 (with $\beta = 0$ and $\gamma = 1/2$) and that $x(\partial g)$ is bounded by (H1), we obtain

$$(3.47) \quad \left\| \varphi(\lambda P) \left(\frac{d}{2} + g^{-1} x (\partial g) \right) \varphi(\lambda P) u \right\| \lesssim \|\varphi(\lambda P) u\| \lesssim \lambda^{-1/2+\varepsilon} \|\langle x \rangle u\|.$$

The inequality (3.43) follows from (3.45), (3.46) and (3.47) for $\mu = 1$ and for $0 \leq \mu \leq 1$ by interpolation. To prove (3.44), we write

$$\|\langle \mathcal{A}_\lambda \rangle^\mu \psi(\lambda P) \langle x \rangle^{-\mu}\| \lesssim \|\psi(\lambda P) \langle x \rangle^{-\mu}\| + \|\langle \mathcal{A}_\lambda \rangle^\mu \psi(\lambda P) \langle x \rangle^{-\mu}\| \lesssim \lambda^{-\mu/2+\varepsilon},$$

where we have again used Lemma B.12 with $\beta = 0$ and $\gamma = 1/2$. \square

Lemma 3.7. *For λ large enough, we have*

$$\mathbb{1}_I(\lambda P) [i(\lambda P)^{1/2}, \mathcal{A}_\lambda] \mathbb{1}_I(\lambda P) \geq \frac{\delta^2 \sqrt{\inf I}}{2} \mathbb{1}_I(\lambda P).$$

Proof. Recall that (3.16) gives

$$[i(\lambda P)^{1/2}, \mathcal{A}_\lambda] = (\lambda P)^{1/2} \varphi^2(\lambda P) + R.$$

On the other hand, we know by Lemma 3.3 that $\|Ru\| \lesssim \lambda^{-\tilde{\varepsilon}} \|u\|$ for some $\tilde{\varepsilon} > 0$. Using $\varphi(x) > \delta > 0$ on I and taking λ large enough, we get the lemma. \square

3.2. Intermediate frequency Mourre estimate.

Here, we obtain a Mourre estimate for frequencies inside the compact $[1/C, C]$. For that, we will use a standard argument in scattering theory. Mimicking Section 3.1, we set

$$(3.48) \quad \mathcal{A} = \varphi(P) A_0 \varphi(P),$$

where $\varphi \in C_0^\infty(]0, +\infty[;]0, +\infty[)$ with $\varphi = 1$ near $[1/C, C]$. As before, \mathcal{A} is essentially self-adjoint on $D(A_0)$ and we denote again \mathcal{A} its closure.

Proposition 3.8. *i) We have $P^{1/2} \in C^2(\mathcal{A})$. The commutators $\text{ad}_{\mathcal{A}}^j P^{1/2}$, $j = 1, 2$, can be extended to bounded operators.*

ii) For each $\sigma \in [1/C, C]$, there exists $\delta > 0$ such that

$$(3.49) \quad \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) [iP^{1/2}, \mathcal{A}] \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) \geq \frac{1}{2\sqrt{C}} \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P).$$

iii) For $0 \leq \mu \leq 1$, we have

$$(3.50) \quad \|\langle \mathcal{A} \rangle^\mu \langle x \rangle^{-\mu}\| \lesssim 1.$$

Proof. The points *i)* and *iii)* follow directly from Proposition 3.1 with $\lambda = 1$. Moreover, using (3.16) and Lemma 3.3, we get

$$[iP^{1/2}, \mathcal{A}] = P^{1/2} \varphi^2(P) + R,$$

where $\|Ru\| \lesssim \|\langle x \rangle^{-\nu} u\|$ for some $\nu > 0$. Then, $R\varphi(P)$ is a compact operator on $L^2(\mathbb{R}^d)$. Let $\sigma \in [1/C, C]$. Since σ is not an eigenvalue of P (see [9, Corollary 5.4]), we have

$$\text{s-}\lim_{\delta \rightarrow 0} \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) = 0.$$

Thus, we obtain

$$\lim_{\delta \rightarrow 0} \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) R\varphi(P) = 0,$$

in operator norm. Using

$$\begin{aligned} \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) [iP^{1/2}, \mathcal{A}] \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) &\geq \sqrt{\sigma - \delta} \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) \\ &\quad + \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P) R\varphi(P) \mathbb{1}_{[\sigma-\delta, \sigma+\delta]}(P), \end{aligned}$$

part *ii)* of the proposition follows. \square

3.3. High frequency Mourre estimate.

In this section, we construct a conjugate operator at high frequencies. We work with the simple σ -temperate metric

$$\gamma = \frac{dx^2}{1+x^2} + \frac{d\xi^2}{1+\xi^2}.$$

We refer to [15, Section XVIII] for the Weyl calculus of Hörmander. For $m(x, \xi)$ a weight function, let $S(m)$ be the set of functions $f \in C^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ such that

$$|\partial_x^\alpha \partial_\xi^\beta f(x, \xi)| \lesssim m(x, \xi) \langle x \rangle^{-|\alpha|} \langle \xi \rangle^{-|\beta|},$$

for all $\alpha, \beta \in \mathbb{N}^d$. In fact, $S(m)$ is the space of symbols of weight m for the metric γ . Let $\Psi(m)$ denote the set of pseudo-differential operators whose symbols are in $S(m)$.

Let $p(x, \xi) \in S(\langle \xi \rangle^2)$ be the symbol of P , and

$$p_0(x, \xi) = \sum_{j,k} g^{j,k}(x) \xi_j \xi_k \in S(\langle \xi \rangle^2),$$

be its principal part. We have $p - p_0 \in S(1)$. Let

$$H_{p_0} = \begin{pmatrix} \partial_\xi p_0 \\ -\partial_x p_0 \end{pmatrix},$$

be the Hamiltonian of p_0 . Since the metric \mathbf{g} is non-trapping by assumption, the energy $\{p_0 = 1\}$ is non-trapping for the Hamiltonian flow of p_0 . Then, using a result of C. Gérard and Martinez [12], one can construct a function $b(x, \xi) \in S(\langle x \rangle \langle \xi \rangle)$ such that $b = x \cdot \xi$ for x large enough, and

$$(3.51) \quad H_{p_0} b \geq \delta,$$

for some $\delta > 0$ and all $(x, \xi) \in p_0^{-1}([1 - \varepsilon, 1 + \varepsilon])$, $\varepsilon > 0$. We set $A = \text{Op}(a)$ with

$$a(x, \xi) = b(x, (p_0 + 1)^{-1/2} \xi) \in S(\langle x \rangle).$$

Let $f \in C^\infty(\mathbb{R}; \mathbb{R})$ be such that $f = 1$ on $[2, +\infty[$ and $f = 0$ on $] -\infty, 1]$. As conjugate operator at high frequency, we choose

$$(3.52) \quad A_\infty = f(P) A f(P).$$

Let $\varphi \in C_0^\infty(\mathbb{R})$ satisfy $\varphi + f = 1$ on $[-1, +\infty[$. Since $P \geq 0$, we have $f(P) = 1 - \varphi(P)$. On the other hand, from the functional calculus of pseudo-differential operators, $\varphi(P) \in \Psi(\langle \xi \rangle^{-\infty})$ and then $f(P) \in \Psi(1)$. To prove this assertion, one can, for instance, adapt Theorem 8.7 of [8] or Section D.11 of [7] to the metric γ . In particular, A_∞ is well defined as a pseudo-differential operator, and we have

$$(3.53) \quad A_\infty = A + \Psi(\langle x \rangle \langle \xi \rangle^{-\infty}) \in \Psi(\langle x \rangle).$$

The following proposition summarizes the useful properties of A_∞ .

Proposition 3.9. *i) The operator A_∞ is essentially self-adjoint on any core of $\langle x \rangle$ with $D(\langle x \rangle) = \{u \in L^2(\mathbb{R}^d); \langle x \rangle u \in L^2(\mathbb{R}^d)\}$. Moreover,*

$$\|A_\infty u\| \lesssim \|\langle x \rangle u\|,$$

for all $u \in D(\langle x \rangle)$.

ii) We have $P^{1/2} \in C^2(A_\infty)$. The commutators $[P^{1/2}, A_\infty]$ and $[[P^{1/2}, A_\infty], A_\infty]$ are in $\Psi(1)$ and can be extended as bounded operators on $L^2(\mathbb{R}^d)$.

iii) For $C > 0$ large enough,

$$\mathbb{1}_{[C, +\infty[}(P) i[P^{1/2}, A_\infty] \mathbb{1}_{[C, +\infty[}(P) \geq \frac{\delta}{8} \mathbb{1}_{[C, +\infty[}(P).$$

The rest of this subsection is devoted to the proof of this proposition. It is a direct consequence of the next lemmas. For the first part of the proposition, we will use the following extension of Nelson's theorem due to C. Gérard and Łaba [11, Lemma 1.2.5] (see also Reed and Simon [29, Theorem X.36]).

Theorem 3.10 (Nelson's theorem). *Let \mathcal{H} be a Hilbert space, $N \geq 1$ a self-adjoint operator on \mathcal{H} , H a symmetric operator such that $D(N) \subset D(H)$ and*

$$\begin{aligned} \|Hu\| &\lesssim \|Nu\|, \\ |(Hu, Nu) - (Nu, Hu)| &\lesssim \|N^{1/2}u\|^2, \end{aligned}$$

for all $u \in D(N)$. Then, H is essentially self-adjoint on any core of N .

Lemma 3.11. *The operator A_∞ is essentially self-adjoint on any core of $(\langle x \rangle, D(\langle x \rangle))$ (in particular, on the Schwartz space $\mathcal{S}(\mathbb{R}^d)$) and*

$$\|A_\infty u\| \lesssim \|\langle x \rangle u\|,$$

for all $u \in D(\langle x \rangle)$.

Proof. The operator $N = \langle x \rangle$ is self-adjoint on $D(N) = D(\langle x \rangle)$ and essentially self-adjoint on $\mathcal{S}(\mathbb{R}^d)$. Since $A_\infty \in \Psi(\langle x \rangle)$ and $N^{-1} \in \Psi(\langle x \rangle^{-1})$, the operator $A_\infty N^{-1} \in \Psi(1)$ is bounded on $L^2(\mathbb{R}^d)$ by Calderon and Vaillancourt's theorem. Then, A_∞ is defined on $D(N)$ and

$$\|A_\infty u\| \lesssim \|Nu\|,$$

for all $u \in D(N)$.

By pseudo-differential calculus, $\langle x \rangle^{-1/2} [A_\infty, \langle x \rangle] \langle x \rangle^{-1/2} \in \Psi(\langle \xi \rangle^{-1})$ is bounded on $L^2(\mathbb{R}^d)$. Then, working first on $\mathcal{S}(\mathbb{R}^d)$, this gives

$$|(Au, Nu) - (Nu, Au)| \lesssim \|N^{1/2}u\|^2,$$

for all $u \in D(N)$. Thus, Theorem 3.10 implies that A_∞ is essentially self-adjoint on any core of $D(\langle x \rangle)$. \square

Lemma 3.12. *Let $g \in C^\infty(\mathbb{R}; [0, +\infty[)$ be such that $g = 0$ on $] -\infty, a]$ and $g = 1$ on $[b, +\infty[$, for some $0 < a < b$. Then,*

$$g(P)P^{1/2} = \text{Op}((p_0 + 1)^{1/2}) + \Psi(1) \in \Psi(\langle \xi \rangle).$$

We omit the proof of this classical result. It follows from (3.5) and the Beals lemma [4]. We refer to Section 4.4 of [14] for similar arguments (see also [4, Theorem 4.9]).

Remark 3.13. *For the subsequent uses, a parametrix will be enough. In fact, since we work with the metric γ , the remainder terms will decay like $\langle (x, \xi) \rangle^{-\infty}$. Therefore, they can "absorb" the pseudo-differential operators of any weight. In particular, this allows to treat the commutators.*

Lemma 3.14. *We have $[P^{1/2}, A_\infty] \in \Psi(1)$ and $[[P^{1/2}, A_\infty], A_\infty] \in \Psi(\langle \xi \rangle^{-1})$. These commutators extend as bounded operators on $L^2(\mathbb{R}^d)$.*

Proof. Let $g \in C^\infty(\mathbb{R})$ as in Lemma 3.12 be such that $fg = f$. Then,

$$(3.54) \quad [P^{1/2}, A_\infty] = [g(P)P^{1/2}, A_\infty].$$

Since $g(P)P^{1/2} \in \Psi(\langle \xi \rangle)$ by Lemma 3.12 and $A_\infty \in \Psi(\langle x \rangle)$, the pseudo-differential calculus gives $[g(P)P^{1/2}, A_\infty] \in \Psi(1)$. The same way, $[[P^{1/2}, A_\infty], A_\infty] \in \Psi(\langle \xi \rangle^{-1})$. Using Calderon and Vaillancourt's theorem and working first on $\mathcal{S}(\mathbb{R}^d)$ which is dense in $D(\langle x \rangle) \cap H^1(\mathbb{R}^d)$, one can prove that these operators extend as bounded operator on $L^2(\mathbb{R}^d)$. \square

Lemma 3.15. *We have $P^{1/2} \in C^2(A_\infty)$.*

Proof. Let $H = \langle D \rangle = \text{Op}(\langle \xi \rangle) \in \Psi(\langle \xi \rangle)$ be the self-adjoint operator with domain $D(H) = D(P^{1/2}) = H^1(\mathbb{R}^d)$ (see Lemma B.13). We remark that $(H \pm z)^{-1} = \text{Op}(\langle \xi \rangle \pm z)^{-1} \in \Psi(1)$ is a Fourier multiplier. Thus, $D(\langle x \rangle)$, which is a core of A_∞ from Lemma 3.11, is stable by $(H \pm z)^{-1}$. On the other hand, $[H, A_\infty] \in \Psi(1)$ can be extend as a bounded operator on $L^2(\mathbb{R}^d)$. Then, Theorem A.1 implies $H \in C^1(A_\infty)$.

Since $H \in C^1(A_\infty)$ and $[H, A_\infty]$ is bounded on $L^2(\mathbb{R}^d)$, Lemma A.3 says that e^{itA_∞} leaves $D(H)$ invariant. Then, e^{itA_∞} leaves $D(P^{1/2}) = D(H)$ invariant and $[P^{1/2}, A_\infty]$ is bounded from Lemma 3.14. Then, Theorem A.2 implies that $P^{1/2} \in C^1(A_\infty)$.

The lemma follows from Theorem A.2, Remark A.4 and Lemma 3.14. \square

Lemma 3.16. *For $C > 0$ large enough,*

$$\mathbb{1}_{[C, +\infty[}(P) i[P^{1/2}, A_\infty] \mathbb{1}_{[C, +\infty[}(P) \geq \frac{\delta}{8} \mathbb{1}_{[C, +\infty[}(P).$$

Proof. Equation (3.53) and (3.54), Lemma 3.12 and the pseudo-differential calculus give

$$\begin{aligned} (3.55) \quad i[P^{1/2}, A_\infty] &= i[g(P)P^{1/2}, A_\infty] \\ &= i[\text{Op}((p_0 + 1)^{1/2}), \text{Op}(a)] + \Psi(\langle \xi \rangle^{-1}) \\ &= \frac{1}{2} \text{Op}((p_0 + 1)^{-1/2} \mathbb{H}_{p_0} a) + \Psi(\langle \xi \rangle^{-1}) \\ &= \frac{1}{2} \text{Op} \left((p_0 + 1)^{-1/2} (\partial_\xi p_0)(x, \xi) \cdot (\partial_x b)(x, (p_0 + 1)^{-1/2} \xi) \right. \\ &\quad \left. - (p_0 + 1)^{-1} (\partial_x p_0)(x, \xi) \cdot (\partial_\xi b)(x, (p_0 + 1)^{-1/2} \xi) \right) + \Psi(\langle \xi \rangle^{-1}) \\ &= \frac{1}{2} \text{Op}((\mathbb{H}_{p_0} b)(x, (p_0 + 1)^{-1/2} \xi)) + \Psi(\langle \xi \rangle^{-1}). \end{aligned}$$

For the last equality, we have used that p_0 is a homogeneous polynomial of order 2 in ξ .

Note that

$$p_0(x, (p_0 + 1)^{-1/2} \xi) = (p_0 + 1)^{-1} p_0 \in [1 - \varepsilon, 1 + \varepsilon],$$

for ξ large enough. Then, adding a cut-off function in ξ , (3.51) and (3.55) imply that

$$i[P^{1/2}, A_\infty] = \text{Op}(c) + \Psi(\langle \xi \rangle^{-1}),$$

with $c \in S(1)$ and $c(x, \xi) \geq \delta/2$. We write $c(x, \xi) = \delta/4 + d^2(x, \xi)$ with $d \in S(1)$ real valued. Thus, by the pseudo-differential calculus,

$$(3.56) \quad \begin{aligned} i[P^{1/2}, A_\infty] &\geq \delta/4 + \text{Op}(d)^* \text{Op}(d) + \Psi(\langle \xi \rangle^{-1}) \\ &\geq \delta/4 + \Psi(\langle \xi \rangle^{-1}), \end{aligned}$$

as self-adjoint operators (one can also apply the Gårding inequality).

Let $R \in \Psi(\langle \xi \rangle^{-1})$. Since $P \in \Psi(\langle \xi \rangle^2)$, the operator

$$R^*(P+1)R \in \Psi(1),$$

is a bounded operator on $L^2(\mathbb{R}^d)$. Then, $(P+1)^{1/2}R$ is also bounded on $L^2(\mathbb{R}^d)$. In particular, we have

$$(3.57) \quad \begin{aligned} \|\mathbb{1}_{[C, +\infty[}(P)R\| &= \|\mathbb{1}_{[C, +\infty[}(P)(P+1)^{-1/2}(P+1)^{1/2}R\| \\ &\lesssim \|\mathbb{1}_{[C, +\infty[}(P)(P+1)^{-1/2}\| \|(P+1)^{1/2}R\| \\ &\lesssim (C+1)^{-1/2}. \end{aligned}$$

The lemma follows from (3.56) and (3.57). \square

4. PROOF OF THE LINEAR ESTIMATES

In this section, we will show the main estimates for the linear wave equation (Theorem 1.3 and Theorem 1.4). To prove these results, we will make a dyadic decomposition of the low frequencies. We will often consider $\varphi \in C_0^\infty(]0, +\infty[; [0, +\infty[)$ such that

$$(4.1) \quad \sum_{\lambda=2^n, n \geq 0} \varphi(\lambda x) = 1,$$

for $x \in]0, 1]$. To φ , we will associate $\tilde{\varphi} \in C_0^\infty(]0, +\infty[; [0, +\infty[)$ satisfying $\tilde{\varphi}\varphi = \varphi$.

We begin with a technical lemma which proves Remark 1.5 *ii*).

Lemma 4.1. *For all $\tilde{\mu} < \mu \leq 3/2$, we have*

$$\begin{aligned} \|\langle x \rangle^{-\mu} \tilde{\partial}_\ell u\| &\lesssim \|\langle x \rangle^{-\tilde{\mu}} P^{1/2} u\|, \\ \|\langle x \rangle^{-\mu} P^{1/2} u\| &\lesssim \sum_{\ell=1}^d \|\langle x \rangle^{-\tilde{\mu}} \tilde{\partial}_\ell u\|. \end{aligned}$$

Proof. Since the two inequalities can be treated the same way, we only prove the first one. We write

$$\|\langle x \rangle^{-\mu} \tilde{\partial}_\ell u\| \leq \|\langle x \rangle^{-\mu} \tilde{\partial}_\ell \Psi(P \leq C)u\| + \|\langle x \rangle^{-\mu} \tilde{\partial}_\ell \Psi(P \geq C)u\| =: I_1 + I_2.$$

- We first estimate I_1 . Let φ be as in (4.1). For $\tilde{\mu} < \mu$, we have, using Lemma B.12,

$$\begin{aligned}
I_1 &\lesssim \sum_{\lambda \text{ dyadic}} \|\langle x \rangle^{-\mu} \tilde{\partial}_\ell \varphi(\lambda P) u\| \\
&= \sum_{\lambda \text{ dyadic}} \|\langle x \rangle^{-\mu} (\lambda^{1/2} \tilde{\partial}_\ell) \varphi(\lambda P) (\lambda P)^{-1/2} \langle x \rangle^{\tilde{\mu}} \langle x \rangle^{-\tilde{\mu}} P^{1/2} u\| \\
&\lesssim \sum_{\lambda \text{ dyadic}} \lambda^{-\frac{\mu-\tilde{\mu}}{2} + \varepsilon} \|\langle x \rangle^{-\tilde{\mu}} P^{1/2} u\| \\
&\lesssim \|\langle x \rangle^{-\tilde{\mu}} P^{1/2} u\|,
\end{aligned}$$

for ε small enough.

- We now estimate I_2 . By Lemma 3.12 and the pseudo-differential calculus, we know that the operator

$$\langle x \rangle^{-\mu} \tilde{\partial}_\ell \Psi(P \geq C) P^{-1/2} \langle x \rangle^\mu,$$

is bounded. Therefore,

$$I_2 = \|\langle x \rangle^{-\mu} \tilde{\partial}_\ell \Psi(P \geq C) P^{-1/2} \langle x \rangle^\mu \langle x \rangle^{-\mu} P^{1/2} u\| \lesssim \|\langle x \rangle^{-\mu} P^{1/2} u\|.$$

□

Using the same type of proof, one can show the following estimate.

Lemma 4.2. *For all $\mu > 1$, we have*

$$\|\langle x \rangle^{-\mu} u\| \lesssim \|P^{1/2} u\|.$$

Remark 4.3. *Let $\mu > 0$ be given. Then, for all $\varepsilon > 0$, there exist $0 < \tilde{\mu} < \mu$, $0 < \tilde{\varepsilon} < \varepsilon$ such that $F_{\tilde{\mu}}^{\tilde{\varepsilon}}(T) \leq F_{\mu}^{\varepsilon}(T)$. Then, it is sufficient to bound the different quantities we consider by $F_{\tilde{\mu}}^{\tilde{\varepsilon}}(T)$ rather than by $F_{\mu}^{\varepsilon}(T)$.*

Theorem 1.3 will follow from the corresponding result for the group $e^{-itP^{1/2}}$.

Proposition 4.4. *Let $0 < \mu \leq 1$. Then, for all $0 < \varepsilon < \mu$, we have*

$$\int_0^T \|\langle x \rangle^{-\mu} e^{-itP^{1/2}} v\|^2 dt \lesssim F_{\mu}^{\varepsilon}(T) \|v\|^2.$$

Proof. We write

$$\begin{aligned}
\int_0^T \|\langle x \rangle^{-\mu} e^{-itP^{1/2}} v\|^2 dt &\lesssim \int_0^T \|\langle x \rangle^{-\mu} e^{-itP^{1/2}} \Psi(P \leq 1/C) v\|^2 dt \\
&\quad + \int_0^T \|\langle x \rangle^{-\mu} e^{-itP^{1/2}} \Psi(1/C \leq P \leq C) v\|^2 dt \\
&\quad + \int_0^T \|\langle x \rangle^{-\mu} e^{-itP^{1/2}} \Psi(P \geq C) v\|^2 dt =: I_1 + I_2 + I_3.
\end{aligned}$$

- We first estimate I_1 . Let $\varphi, \tilde{\varphi}$ be as in (4.1). Proposition 3.1 gives

$$\|\langle x \rangle^{-\mu} \tilde{\varphi}(\lambda P) \langle \mathcal{A}_\lambda \rangle^\mu\|^2 \lesssim \lambda^{-\mu + \varepsilon_1},$$

for all $\varepsilon_1 > 0$. Then,

$$\begin{aligned} I_1 &\lesssim \sum_{\lambda \text{ dyadic}} \lambda^{-\mu+\varepsilon_1} \int_0^T \|\langle \mathcal{A}_\lambda \rangle^{-\mu} e^{-itP^{1/2}} \varphi(\lambda P)v\|^2 dt \\ &= \sum_{\lambda \text{ dyadic}} \lambda^{-\mu+\varepsilon_1+1/2} \int_0^{\lambda^{-1/2}T} \|\langle \mathcal{A}_\lambda \rangle^{-\mu} e^{-is(\lambda P)^{1/2}} \varphi(\lambda P)v\|^2 ds \\ &\lesssim \sum_{\lambda \text{ dyadic}} \lambda^{-\mu+\varepsilon_1+1/2+\varepsilon_3} F_\mu^{\varepsilon_2}(\lambda^{-1/2}T) \|v\|^2, \end{aligned}$$

for all $\varepsilon_2, \varepsilon_3 > 0$ with $\varepsilon_2 < \mu$. Here, we have used Proposition 3.1, Remark 2.3, Theorem 2.4 (for $\mu > 1/2$) and Corollary 2.5 (for $\mu \leq 1/2$) with $H = (\lambda P)^{1/2}$.

★ If $\mu > 1/2$, then, by choosing $\varepsilon_1, \varepsilon_3$ small enough, the sum is convergent and we find

$$I_1 \lesssim \|v\|^2.$$

★ If $\mu \leq 1/2$, we find

$$I_1 \lesssim \sum_{\lambda \text{ dyadic}} \lambda^{\varepsilon_1+\varepsilon_3-\varepsilon_2} T^{1-2\mu+2\varepsilon_2} \|v\|^2.$$

Fixing first ε_2 and then $\varepsilon_1, \varepsilon_3$ small enough makes the sum convergent.

• We now treat I_2 . Since $[1/C, C]$ is a compact interval, Proposition 3.8 gives us a finite number of open intervals I_k , $k = 1, \dots, K$, satisfying (3.49) and

$$[1/C, C] \subset \bigcup_{k=1}^K I_k.$$

Then, applying Theorem 2.4 (for $\mu > 1/2$) and Corollary 2.5 (for $\mu \leq 1/2$) on each I_k (slightly reduced), we obtain

$$I_2 \lesssim F_\mu^\varepsilon(T) \|v\|^2.$$

• Let us finally estimate I_3 . By Proposition 3.9 and an interpolation argument, we get

$$\|\langle x \rangle^{-\mu} \langle \mathcal{A}_\infty \rangle^\mu\| \lesssim 1.$$

Thus,

$$\begin{aligned} \int_0^T \|\langle x \rangle^{-\mu} e^{-itP^{1/2}} \Psi(P \geq C)v\|^2 dt &\lesssim \int_0^T \|\langle \mathcal{A}_\infty \rangle^{-\mu} e^{-itP^{1/2}} \Psi(P \geq C)v\|^2 dt \\ &\lesssim F_\mu^\varepsilon(T) \|v\|^2, \end{aligned}$$

where we have used Theorem 2.4 (for $\mu > 1/2$) and Corollary 2.5 (for $\mu \leq 1/2$). \square

For the proof of Theorem 1.3, we will need the following theorem of Christ and Kiselev [6] in a form available in the article of Burq [5].

Theorem 4.5 (Christ–Kiselev). *Consider a bounded operator $\mathcal{T} : L^p(\mathbb{R}; B_1) \longrightarrow L^q(\mathbb{R}; B_2)$ given by a locally integrable Kernel $K(t, s)$ with value operators from B_1 to B_2 , where B_1 and B_2 are Banach spaces. Suppose that $p < q$. Then, the operator*

$$\tilde{\mathcal{T}}f(t) = \int_{s < t} K(t, s)f(s) ds,$$

is bounded from $L^p(\mathbb{R}; B_1)$ to $L^q(\mathbb{R}; B_2)$ by

$$\|\tilde{\mathcal{T}}\|_{L^p(\mathbb{R}; B_1) \rightarrow L^q(\mathbb{R}; B_2)} \leq (1 - 2^{-(p^{-1}-q^{-1})}) \|\mathcal{T}\|_{L^p(\mathbb{R}; B_1) \rightarrow L^q(\mathbb{R}; B_2)}.$$

Proof of Theorem 1.3. By linearity and uniqueness it is sufficient to consider separately the cases $(u_0, u_1) = 0$, $G = 0$.

• $G = 0$. Thanks to the discussion at the beginning of Section 3, the solution of (1.4) is given by

$$\begin{pmatrix} u(t) \\ \partial_t u(t) \end{pmatrix} = e^{-itR} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} \quad \text{with} \quad R = \begin{pmatrix} 0 & i \\ -iP & 0 \end{pmatrix}, \quad R = U^* L U.$$

Using Lemma 4.1, we see that for $\tilde{\mu} < \mu$ we have

$$\|\langle x \rangle^{-\mu} u'\|_{L^2}^2 \lesssim \|\langle x \rangle^{-\tilde{\mu}} M \begin{pmatrix} u(t) \\ \partial_t u(t) \end{pmatrix}\|_{L^2 \times L^2}^2,$$

with

$$(4.2) \quad M := \begin{pmatrix} P^{1/2} & 0 \\ 0 & 1 \end{pmatrix}, \quad M U^* = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix}.$$

Using Proposition 4.4, we therefore have the following estimate

$$\begin{aligned} \int_0^T \|\langle x \rangle^{-\mu} u'\|_{L^2}^2 dt &\lesssim \int_0^T \|\langle x \rangle^{-\tilde{\mu}} M e^{-itR} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}\|_{L^2 \times L^2}^2 dt \\ &\lesssim \int_0^T \|\langle x \rangle^{-\tilde{\mu}} e^{-itL} U \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}\|_{L^2 \times L^2}^2 dt \\ &\lesssim F_{\tilde{\mu}}^\varepsilon(T) \left\| U \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} \right\|_{L^2 \times L^2}^2 = F_{\tilde{\mu}}^\varepsilon(T) \|(u_0, u_1)\|_{\mathcal{E}}^2. \end{aligned}$$

• $(u_0, u_1) = 0$. In this case, the solution of (1.4) is given by

$$\begin{pmatrix} u(t) \\ \partial_t u(t) \end{pmatrix} = \int_0^t e^{i(s-t)R} \begin{pmatrix} 0 \\ G(s) \end{pmatrix} ds.$$

Thus, for all $\tilde{\mu} < \mu$,

$$(4.3) \quad \int_0^T \|\langle x \rangle^{-\mu} u'\|_{L^2}^2 \lesssim \int_0^T \left\| \int_0^t \langle x \rangle^{-\tilde{\mu}} e^{i(s-t)L} U \begin{pmatrix} 0 \\ G(s) \end{pmatrix} ds \right\|_{L^2 \times L^2}^2 dt.$$

Let

$$\mathcal{T}f(t) = \int_{\mathbb{R}} \langle x \rangle^{-\tilde{\mu}} \mathbb{1}_{[0,T]}(s) \mathbb{1}_{[0,T]}(t) e^{i(s-t)L} f(s) ds.$$

We estimate

$$\begin{aligned} \|\mathcal{T}f\|_{L^2(\mathbb{R}; L^2 \times L^2)}^2 &= \int_0^T \left\| \langle x \rangle^{-\tilde{\mu}} e^{-itL} \int_0^T e^{isL} f(s) ds \right\|_{L^2 \times L^2}^2 dt \\ &\lesssim F_{\tilde{\mu}}^\varepsilon(T) \left\| \int_0^T e^{isL} f(s) ds \right\|_{L^2 \times L^2}^2 \\ &\lesssim F_{\tilde{\mu}}^\varepsilon(T) \left(\int_0^T \|f(s)\|_{L^2 \times L^2} ds \right)^2. \end{aligned}$$

It follows

$$\|\mathcal{T}\|_{L^1(\mathbb{R}; L^2 \times L^2) \rightarrow L^2(\mathbb{R}; L^2 \times L^2)}^2 \lesssim F_\mu^\varepsilon(T).$$

The expression on the right hand side of (4.3) is

$$\|\tilde{\mathcal{T}}U(0, G(s))\|_{L^2(\mathbb{R}; L^2 \times L^2)}^2,$$

with

$$\tilde{\mathcal{T}}f(t) = \int_{s < t} \langle x \rangle^{-\tilde{\mu}} \mathbb{1}_{[0, T]}(s) \mathbb{1}_{[0, T]}(t) e^{i(s-t)L} f(s) ds.$$

We can apply the theorem of Christ and Kiselev to conclude that

$$\begin{aligned} \|\tilde{\mathcal{T}}U(0, G(s))\|_{L^2(\mathbb{R}; L^2 \times L^2)}^2 &\lesssim F_\mu^\varepsilon(T) \left(\int_0^T \left\| U \begin{pmatrix} 0 \\ G(s) \end{pmatrix} \right\|_{L^2 \times L^2} ds \right)^2 \\ &= F_\mu^\varepsilon(T) \left(\int_0^T \|G(s)\|_{L^2} ds \right)^2, \end{aligned}$$

which finishes the proof. \square

Theorem 1.4 is now proved for $N = 0$ using in addition the usual energy estimate

$$(4.4) \quad \|u'\|_{L^2(\mathbb{R}^d)} \lesssim \|u'(0, \cdot)\|_{L^2(\mathbb{R}^d)} + \int_0^T \|G(s, \cdot)\|_{L^2(\mathbb{R}^d)} ds.$$

Note that in the usual energy estimate u' is replaced by $(\partial_t u, P^{1/2}u)$, but we have

$$\sum_k \|\tilde{\partial}_k u\| \lesssim \|P^{1/2}u\| \lesssim \sum_k \|\tilde{\partial}_k u\|,$$

by Lemma B.13. It will be useful to have similar estimates to the preceding containing a $L^2(\mathbb{R}^{d+1}, \langle x \rangle^\mu dt dx)$ norm of G on the right hand side rather than a $L_t^1 L_x^2$ norm.

Proposition 4.6. *Assume $0 < \mu \leq 1$.*

i) Let

$$(4.5) \quad \begin{cases} (i\partial_t - P^{1/2})v = G, \\ v|_{t=0} = 0. \end{cases}$$

Then we have, for all $0 < \varepsilon < \mu$,

$$(4.6) \quad \int_0^T \|\langle x \rangle^{-\mu} v\|^2 dt \lesssim (F_\mu^\varepsilon(T))^2 \int_0^T \|\langle x \rangle^\mu G\|^2 dt.$$

ii) Let

$$(4.7) \quad \begin{cases} (\partial_t^2 + P)u = G, \\ (u|_{t=0}, \partial_t u|_{t=0}) = 0. \end{cases}$$

Then we have, for all $0 < \varepsilon < \mu$,

$$(4.8) \quad \int_0^T \|\langle x \rangle^{-\mu} u'\|^2 dt \lesssim (F_\mu^\varepsilon(T))^2 \int_0^T \|\langle x \rangle^\mu G\|^2 dt.$$

Proof. *i)* We have

$$\begin{aligned} \int_0^T \|\langle x \rangle^{-\mu} v\|^2 dt &\lesssim \int_0^T \|\langle x \rangle^{-\mu} \Psi(P \leq 1/C)v\|^2 dt + \int_0^T \|\langle x \rangle^{-\mu} \Psi(1/C \leq P \leq C)v\|^2 dt \\ &\quad + \int_0^T \|\langle x \rangle^{-\mu} \Psi(P \geq C)v\|^2 dt =: I_1 + I_2 + I_3. \end{aligned}$$

• We first estimate I_1 . Let $\varphi, \tilde{\varphi}$ be as in (4.1). By Proposition 3.1, we know that

$$\|\langle x \rangle^{-\mu} \varphi(\lambda P)v\|^2 = \|\langle x \rangle^{-\mu} \tilde{\varphi}(\lambda P) \langle \mathcal{A}_\lambda \rangle^\mu \langle \mathcal{A}_\lambda \rangle^{-\mu} \varphi(\lambda P)v\|^2 \lesssim \lambda^{-\mu+\varepsilon_1} \|\langle \mathcal{A}_\lambda \rangle^{-\mu} \varphi(\lambda P)v\|^2.$$

Therefore, we have

$$\begin{aligned} I_1 &\lesssim \sum_{\lambda \text{ dyadic}} \lambda^{-\mu+\varepsilon_1} \int_0^T \|\langle \mathcal{A}_\lambda \rangle^{-\mu} \varphi(\lambda P)v(t)\|^2 dt \\ &= \sum_{\lambda \text{ dyadic}} \lambda^{-\mu+\varepsilon_1+1/2} \int_0^{\lambda^{-1/2}T} \|\langle \mathcal{A}_\lambda \rangle^{-\mu} \varphi(\lambda P)v(\lambda^{1/2}s)\|^2 ds. \end{aligned}$$

Now observe that $\tilde{v}(s) = v(\lambda^{1/2}s)$ is solution of the equation

$$\begin{cases} (i\partial_s - (\lambda P)^{1/2})\tilde{v} = \lambda^{1/2}G(\lambda^{1/2}s), \\ \tilde{v}|_{s=0} = 0. \end{cases}$$

We now apply Corollary 2.8 with $H = (\lambda P)^{1/2}$. Using also again Proposition 3.1, we obtain

$$\begin{aligned} &\int_0^{\lambda^{-1/2}T} \|\langle \mathcal{A}_\lambda \rangle^{-\mu} \varphi(\lambda P)v(\lambda^{1/2}s)\|^2 ds \\ &\lesssim (F_\mu^{\varepsilon_2}(\lambda^{-1/2}T))^2 \lambda \int_0^{\lambda^{-1/2}T} \|\langle \mathcal{A}_\lambda \rangle^\mu \varphi(\lambda P)G(\lambda^{1/2}s)\|^2 ds \\ &\lesssim (F_\mu^{\varepsilon_2}(\lambda^{-1/2}T))^2 \lambda^{1-\mu+\varepsilon_3} \int_0^{\lambda^{-1/2}T} \|\langle x \rangle^\mu G(\lambda^{1/2}s)\|^2 ds \\ &= (F_\mu^{\varepsilon_2}(\lambda^{-1/2}T))^2 \lambda^{1/2-\mu+\varepsilon_3} \int_0^T \|\langle x \rangle^\mu G(t)\|^2 dt. \end{aligned}$$

Thus,

$$I_1 \lesssim \sum_{\lambda \text{ dyadic}} \lambda^{1-2\mu+\varepsilon_1+\varepsilon_3} (F_\mu^{\varepsilon_2}(\lambda^{-1/2}T))^2 \int_0^T \|\langle x \rangle^\mu G(t)\|^2 dt.$$

If $\mu \leq 1/2$, then we see that

$$I_1 \lesssim \sum_{\lambda \text{ dyadic}} \lambda^{\varepsilon_1+\varepsilon_3-2\varepsilon_2} T^{2(1-2\mu+2\varepsilon_2)} \int_0^T \|\langle x \rangle^\mu G(t)\|^2 dt.$$

Once $0 < \varepsilon_2 < \mu$ fixed, it is therefore sufficient to choose $\varepsilon_1, \varepsilon_3$ small enough such that $\varepsilon_1 + \varepsilon_3 < 2\varepsilon_2$. If $\mu > 1/2$, we choose $\varepsilon_1, \varepsilon_3$ small enough such that $\varepsilon_1 + \varepsilon_3 < 2\mu - 1$. Then,

$$I_1 \lesssim \int_0^T \|\langle x \rangle^\mu G(t)\|^2 dt.$$

- We now study I_2 . Part *iii*) of Proposition 3.8 implies

$$I_2 \lesssim \int_0^T \|\langle \mathcal{A} \rangle^{-\mu} \Psi(1/C \leq P \leq C)v\|^2 dt.$$

As in the proof of Proposition 4.4, Proposition 3.8 gives us a finite number of open intervals I_k , $k = 1, \dots, K$, satisfying (3.49) and

$$[1/C, C] \subset \bigcup_{k=1}^K I_k.$$

Then, applying Corollary 2.8 on each I_k (slightly reduced) and using Proposition 3.8, we obtain

$$I_2 \lesssim (F_\mu^\varepsilon(T))^2 \int_0^T \|\langle x \rangle^\mu G\|^2 dt.$$

- We finally estimate I_3 . Proposition 3.9 and Corollary 2.8 yield

$$\begin{aligned} I_3 &\lesssim \int_0^T \|\langle A_\infty \rangle^{-\mu} \Psi(P \geq C)v\|^2 dt \lesssim (F_\mu^\varepsilon(T))^2 \int_0^T \|\langle A_\infty \rangle^\mu G\|^2 dt \\ &\lesssim (F_\mu^\varepsilon(T))^2 \int_0^T \|\langle x \rangle^\mu G\|^2 dt. \end{aligned}$$

ii) We first write (4.6) as a first order system

$$i\partial_t \begin{pmatrix} u \\ \partial_t u \end{pmatrix} = R \begin{pmatrix} u \\ \partial_t u \end{pmatrix} + i \begin{pmatrix} 0 \\ G \end{pmatrix}, \quad \begin{pmatrix} u \\ \partial_t u \end{pmatrix} \Big|_{t=0} = 0.$$

It is sufficient to estimate, for $\tilde{\mu} < \mu$,

$$\int_0^T \left\| \langle x \rangle^{-\tilde{\mu}} M \begin{pmatrix} u \\ \partial_t u \end{pmatrix} \right\|_{L^2 \times L^2}^2 dt = \int_0^T \left\| \langle x \rangle^{-\tilde{\mu}} M U^* U \begin{pmatrix} u \\ \partial_t u \end{pmatrix} \right\|_{L^2 \times L^2}^2 dt,$$

with M defined in (4.2). But $v = U \begin{pmatrix} u \\ \partial_t u \end{pmatrix}$ solves

$$(i\partial_t - L)v = iU \begin{pmatrix} 0 \\ G \end{pmatrix}, \quad v|_{t=0} = 0.$$

By (4.2) and part *i*) of the proposition, we find

$$\begin{aligned} \int_0^T \left\| \langle x \rangle^{-\tilde{\mu}} M \begin{pmatrix} u \\ \partial_t u \end{pmatrix} \right\|_{L^2 \times L^2}^2 dt &\lesssim \int_0^T \|\langle x \rangle^{-\tilde{\mu}} v\|_{L^2 \times L^2}^2 dt \\ &\lesssim (F_\mu^\varepsilon(T))^2 \int_0^T \left\| \langle x \rangle^{-\tilde{\mu}} U \begin{pmatrix} 0 \\ G \end{pmatrix} \right\|_{L^2 \times L^2}^2 dt \\ &= (F_\mu^\varepsilon(T))^2 \int_0^T \|\langle x \rangle^{-\tilde{\mu}} G\|_{L^2}^2 dt. \end{aligned}$$

which gives *ii*) thanks to Remark 4.3. \square

We now want to prove Theorem 1.4 for general N . In contrast to the Minkowski case, this does not follow directly from the case $N = 0$ because the vector fields $\tilde{\Omega}$, $\tilde{\partial}_x$ do not commute

with the equation. We will therefore need the form of certain commutators. As in (3.14), a term r_j or \tilde{r}_j , $j \in \mathbb{N}$, will denote a smooth function such that

$$\begin{aligned}\partial_x^\alpha r_j(x) &= \mathcal{O}(\langle x \rangle^{-\rho-j-|\alpha|}), \\ \partial_x^\alpha \tilde{r}_j(x) &= \mathcal{O}(\langle x \rangle^{-j-|\alpha|}).\end{aligned}$$

These functions can change from line to line. Direct computations give

Lemma 4.7. *We have*

$$\begin{aligned}[\tilde{\partial}_j, \tilde{\partial}_k] &= r_1 \tilde{\partial}, & [\tilde{\Omega}^{j,k}, P] &= r_0 \tilde{\partial} \tilde{\partial} + r_1 \tilde{\partial} \\ [\tilde{\partial}_j^*, \tilde{\partial}_k] &= \tilde{\partial}^* r_1 + r_2, & [\tilde{\Omega}^{j,k}, \tilde{\partial}_\ell] &= \tilde{r}_0 \tilde{\partial}, \\ [P, \tilde{\partial}_\ell] &= \tilde{\partial}^* r_1 \tilde{\partial} + r_2 \tilde{\partial}.\end{aligned}$$

As before, we have not written the sum over the indexes on the right hand sides.

We now observe that the vector fields $\tilde{\partial}_j$ can be replaced by powers of P .

Lemma 4.8. *For $0 < \mu \leq 3/2$ and $n \geq 2$, we have*

$$(4.9) \quad \|\langle x \rangle^{-\mu} \tilde{\partial}_{j_1} \cdots \tilde{\partial}_{j_n} u\| \lesssim \sum_{j=0}^{\lfloor \frac{n-1}{2} \rfloor} \sum_{q=1}^d \|\langle x \rangle^{-\mu} \tilde{\partial}_q P^j u\| + \sum_{j=1}^{\lfloor \frac{n}{2} \rfloor} \|\langle x \rangle^{-\mu} P^j u\|.$$

Proof. We first show

$$(4.10) \quad \|\langle x \rangle^{-\mu} \tilde{\partial}_k \tilde{\partial}_\ell u\| \lesssim \|\langle x \rangle^{-\mu} P u\| + \sum_{q=1}^d \|\langle x \rangle^{-\mu} \tilde{\partial}_q u\|.$$

Indeed, we have

$$\|\langle x \rangle^{-\mu} \tilde{\partial}_k \tilde{\partial}_\ell u\| \lesssim \|\langle x \rangle^{-\mu} \tilde{\partial}_k (P+1)^{-1} \tilde{\partial}_\ell (P+1) u\| + \|\langle x \rangle^{-\mu} \tilde{\partial}_k (P+1)^{-1} [P, \tilde{\partial}_\ell] u\| =: A + B.$$

We estimate A .

$$A \leq \|\langle x \rangle^{-\mu} \tilde{\partial}_k (P+1)^{-1} \tilde{\partial}_\ell P u\| + \|\langle x \rangle^{-\mu} \tilde{\partial}_k (P+1)^{-1} \tilde{\partial}_\ell u\|.$$

Noting that $\langle x \rangle^{-\mu} \tilde{\partial}_k (P+1)^{-1} \tilde{\partial}_\ell \langle x \rangle^\mu$ and $\langle x \rangle^{-\mu} \tilde{\partial}_k (P+1)^{-1} \langle x \rangle^\mu$ are bounded by Proposition B.9 and Lemma B.10, we obtain

$$(4.11) \quad A \lesssim \|\langle x \rangle^{-\mu} P u\| + \|\langle x \rangle^{-\mu} \tilde{\partial}_\ell u\|.$$

Now, recall from Lemma 4.7 that

$$[P, \tilde{\partial}_\ell] = \tilde{\partial}^* r_1 \tilde{\partial} + r_2 \tilde{\partial}.$$

Thus, as for (4.11), we see that

$$(4.12) \quad B \lesssim \sum_j \|\langle x \rangle^{-\mu} \tilde{\partial}_j u\|.$$

The inequalities (4.11), (4.12) give (4.10). We will show (4.9) by induction over $n \geq 2$. For $n = 2$ this is exactly (4.10). Assume $n \geq 3$. Using (4.10), we obtain

$$\|\langle x \rangle^{-\mu} \tilde{\partial}_{j_1} \tilde{\partial}_{j_2} \cdots \tilde{\partial}_{j_n} u\| \lesssim \|\langle x \rangle^{-\mu} P \tilde{\partial}_{j_3} \cdots \tilde{\partial}_{j_n} u\| + \sum_{k=1}^d \|\langle x \rangle^{-\mu} \tilde{\partial}_k \tilde{\partial}_{j_3} \cdots \tilde{\partial}_{j_n} u\|.$$

For the second term, we can use the induction hypothesis. For the first term we commute P through $\tilde{\partial}_{j_3} \cdots \tilde{\partial}_{j_n}$. The commutators give terms which can be estimated by terms of the form $\|\langle x \rangle^{-\mu} \tilde{\partial}_{k_m} \cdots \tilde{\partial}_{k_n} u\|$, with $2 \leq m \leq n$, which themselves can be estimated by the induction hypothesis. It remains to consider the term $\|\langle x \rangle^{-\mu} \tilde{\partial}_{j_3} \cdots \tilde{\partial}_{j_n} Pu\|$, which can either be kept ($n = 3$) or be estimated applying the induction hypothesis to Pu rather than to u . \square

In order to show (1.6), it is sufficient to use vector fields in X . This is shown in the next lemma.

Lemma 4.9. *Assume $\rho > 1$. Let $1/2 \leq \mu \leq 1$, $j \in \frac{1}{2}\mathbb{N}$, β be a multi-index and $N = 2j + |\beta|$. Then, for all $\varepsilon > 0$, there exists $\eta_\varepsilon > 1/2$ such that*

$$(4.13) \quad \langle F_\mu^\varepsilon(T) \rangle^{-1} \|\langle x \rangle^{-\mu} (P^j \tilde{\Omega}^\beta u)'\|_{L^2([0,T] \times \mathbb{R}^d)} \\ \lesssim \sum_{|\alpha| \leq N} \left(\|(Y^\alpha u)'(0, \cdot)\|_{L^2(\mathbb{R}^d)} + \int_0^T \|Y^\alpha G\|_{L^2(\mathbb{R}^d)} dt + \|\langle x \rangle^{-\eta_\varepsilon} (X^\alpha u)'\|_{L^2([0,T] \times \mathbb{R}^d)} \right).$$

Moreover, for $\rho = 1$ and $\varepsilon > 0$, the same inequality holds with $\langle F_\mu^\varepsilon(T) \rangle^{-1}$ replaced by $\langle T \rangle^{-\varepsilon}$.

Proof. The inequality will be proven by induction over $|\beta|$. Assume first $\rho > 1$. Since the wave equation commutes with P , the case $|\beta| = 0$ follows from Theorem 1.3 and Lemma B.13. Assume now $|\beta| \geq 1$ and let $v = P^j \tilde{\Omega}^\beta u$. The function v fulfills the following equation

$$(4.14) \quad \begin{cases} (\partial_t^2 + P)v = P^j \tilde{\Omega}^\beta G + P^j [P, \tilde{\Omega}^\beta]u, \\ (v|_{t=0}, \partial_t v|_{t=0}) = (P^j \tilde{\Omega}^\beta u_0, P^j \tilde{\Omega}^\beta u_1). \end{cases}$$

Let v_1, v_2 be the solutions of

$$(4.15) \quad \begin{cases} (\partial_t^2 + P)v_1 = P^j \tilde{\Omega}^\beta G, \\ (v_1|_{t=0}, \partial_t v_1|_{t=0}) = (P^j \tilde{\Omega}^\beta u_0, P^j \tilde{\Omega}^\beta u_1), \end{cases}$$

$$(4.16) \quad \begin{cases} (\partial_t^2 + P)v_2 = P^j [P, \tilde{\Omega}^\beta]u, \\ (v_2|_{t=0}, \partial_t v_2|_{t=0}) = 0. \end{cases}$$

Clearly $v = v_1 + v_2$. We have, for all $\tilde{\mu} < \mu$,

$$\|\langle x \rangle^{-\mu} v_1'\|_{L^2([0,T] \times \mathbb{R}^d)} \lesssim (F_\mu^\varepsilon(T))^{1/2} \left(\|(P^j \tilde{\Omega}^\beta u)'(0, \cdot)\|_{L^2(\mathbb{R}^d)} + \int_0^T \|P^j \tilde{\Omega}^\beta G\|_{L^2(\mathbb{R}^d)} dt \right),$$

where we have used Theorem 1.3. If $\mu > 1/2$, we choose $\tilde{\mu} > 1/2$. We further estimate, by Proposition 4.6,

$$(F_\mu^\varepsilon(T))^{-1} \|\langle x \rangle^{-\mu} v_2'\|_{L^2([0,T] \times \mathbb{R}^d)} \lesssim \|\langle x \rangle^{\tilde{\mu}} P^j [P, \tilde{\Omega}^\beta]u\|_{L^2([0,T] \times \mathbb{R}^d)}.$$

Using Lemma 4.7, we see that $\langle x \rangle^{\tilde{\mu}} P^j [P, \tilde{\Omega}^\beta]u$ is a sum of terms of the form

$$\langle x \rangle^{\tilde{\mu}-\rho} \tilde{\partial}_{k_1} \cdots \tilde{\partial}_{k_m} \tilde{\Omega}^\gamma u,$$

with $1 \leq m \leq 2j + 2$ and $|\gamma| \leq |\beta| - 1$. Using Lemma 4.8, we see that these terms can be estimated in norm by terms of the form

$$\|\langle x \rangle^{\tilde{\mu}-\rho} \tilde{\partial}_\ell (P^q \tilde{\Omega}^\gamma u)\| \quad \text{or} \quad \|\langle x \rangle^{\tilde{\mu}-\rho} P^r \tilde{\Omega}^\gamma u\|,$$

with $q, r \in \mathbb{N}$, $0 \leq q \leq (m-1)/2$ and $1 \leq r \leq m/2$. Applying Lemma 4.1, we see that we can replace $P^{1/2}$ in the second term by partial derivatives and apply the induction hypothesis with $\rho - \tilde{\mu} > 1/2$.

In the case $\rho = 1$, it is enough to choose $\tilde{\mu} = 1/2 - \delta$ with $\delta > 0$ small. \square

Proof of Theorem 1.4. The energy term is easily estimated by the observation that ∂_t and P commute with the equation. The same way, note that we can restrict our attention to vector fields in Y for the second term. Also, by Lemma 4.7, we can arrange for that the vector fields $\tilde{\partial}_x$ are always on the left of the vector fields $\tilde{\Omega}$. Using Lemma 4.7, we see that we can replace $Y^\alpha u'$ by $(Y^\alpha u)'$. Using Lemma 4.1, Lemma 4.8 and Lemma 4.9, we see that it is sufficient to estimate

$$\langle F_\mu^\varepsilon(T) \rangle^{-1} \|\langle x \rangle^{-\mu} P^j v\|_{L^2([0,T] \times \mathbb{R}^d)},$$

in the case $\rho > 1$ and

$$\langle T \rangle^{-\varepsilon} \|\langle x \rangle^{-\mu} P^j v\|_{L^2([0,T] \times \mathbb{R}^d)},$$

in the case $\rho = 1$. These terms can be estimated by Theorem 1.3, because P commutes with the equation. \square

5. PROOF OF THE NONLINEAR RESULT

In this section we will prove the main theorem, Theorem 1.1. The proof of the result will follow closely the arguments of Keel, Smith and Sogge in the Minkowski case (see [20]). We start with the now standard Sobolev estimate (see [22]).

Lemma 5.1. *Suppose that $h \in C^\infty(\mathbb{R}^d)$. Then, for $R > 1$,*

$$(5.1) \quad \|h\|_{L^\infty(R/2 \leq |x| \leq R)} \lesssim R^{\frac{1-d}{2}} \sum_{|\alpha| \leq \lceil \frac{d-1}{2} \rceil + 1} \|Y^\alpha h\|_{L^2(R/4 \leq |x| \leq 2R)}.$$

We now define the bilinear form \tilde{Q} by $\tilde{Q}(u', u') = Q(u')$. The following estimate for the nonlinear part will be crucial.

Lemma 5.2. *Let $\mu_d = \frac{d-1}{4}$. Then, for $L \geq \max(2(\lceil \frac{d-1}{2} \rceil + 1), |\beta|)$, we have*

$$\|Z^\beta \tilde{Q}(u', v')\|_{L^2(\mathbb{R}^d)}^2 \lesssim \left(\sum_{|\alpha| \leq L} \|\langle x \rangle^{-\mu_d} Z^\alpha u'\|_{L^2(\mathbb{R}^d)}^2 \right) \left(\sum_{|\alpha| \leq L} \|\langle x \rangle^{-\mu_d} Z^\alpha v'\|_{L^2(\mathbb{R}^d)}^2 \right).$$

Proof. We clearly have the pointwise bound:

$$\begin{aligned} |Z^\beta \tilde{Q}(u', v')(s, x)| &\lesssim \left(\sum_{|\alpha| \leq L} |Z^\alpha u'(s, x)| \right) \left(\sum_{|\alpha| \leq \lfloor \frac{L}{2} \rfloor} |Z^\alpha v'(s, x)| \right) \\ &\quad + \left(\sum_{|\alpha| \leq L} |Z^\alpha v'(s, x)| \right) \left(\sum_{|\alpha| \leq \lfloor \frac{L}{2} \rfloor} |Z^\alpha u'(s, x)| \right). \end{aligned}$$

We only estimate the first term. Using Lemma 5.1 for a given $R = 2^j$, $j \geq 0$, we get

$$\begin{aligned}
& \|Z^\beta \tilde{Q}(u', v')\|_{L^2(\{|x| \in [2^j, 2^{j+1}]\})}^2 \\
& \lesssim 2^{j(1-d)} \sum_{|\alpha| \leq L} \|Z^\alpha u'\|_{L^2(\{|x| \in [2^j, 2^{j+1}]\})}^2 \sum_{|\alpha| \leq \lfloor \frac{L}{2} \rfloor + \lceil \frac{d-1}{2} \rceil + 1} \|Z^\alpha v'\|_{L^2(\{|x| \in [2^{j-1}, 2^{j+2}]\})}^2 \\
& \lesssim \sum_{|\alpha| \leq L} \|\langle x \rangle^{-\mu_d} Z^\alpha u'\|_{L^2(\{|x| \in [2^j, 2^{j+1}]\})}^2 \sum_{|\alpha| \leq L} \|\langle x \rangle^{-\mu_d} Z^\alpha v'\|_{L^2(\{|x| \in [2^{j-1}, 2^{j+2}]\})}^2 \\
& \lesssim \sum_{|\alpha| \leq L} \|\langle x \rangle^{-\mu_d} Z^\alpha u'\|_{L^2(\{|x| \in [2^j, 2^{j+1}]\})}^2 \sum_{|\alpha| \leq L} \|\langle x \rangle^{-\mu_d} Z^\alpha v'\|_{L^2(\mathbb{R}^d)}^2.
\end{aligned}$$

We also have the bound

$$\|Z^\beta \tilde{Q}(u', v')\|_{L^2(\{|x| < 1\})}^2 \lesssim \sum_{|\alpha| \leq L} \|Z^\alpha u'\|_{L^2(\{|x| < 2\})}^2 \sum_{|\alpha| \leq L} \|Z^\alpha v'\|_{L^2(\{|x| < 2\})}^2.$$

Summing over j gives the lemma. \square

Proof of Theorem 1.1. We follow [20]. Let $u_{-1} = 0$. We define u_k , $k \in \mathbb{N}$ inductively by letting u_k solve

$$(5.2) \quad \begin{cases} \square_{\mathbf{g}} u_k = Q(u'_{k-1}), \\ (u_k|_{t=0}, \partial_t u_k|_{t=0}) = (u_0, u_1). \end{cases}$$

For $T > 0$, we denote

$$M_k(T) = \sup_{0 \leq t \leq T} \sum_{1 \leq i+j \leq M+1} \|\partial_t^i P^{j/2} u_k\|_{L^2(\mathbb{R}^d)} + \sum_{|\alpha| \leq M} K_n(T)^{-1} \|\langle x \rangle^{-\mu_d} Z^\alpha u'_k\|_{L^2([0, T] \times \mathbb{R}^d)},$$

with

$$K_n(T) = \begin{cases} T^{1/n} & d = 3 \text{ or } \rho = 1, \\ 1 & d \geq 4 \text{ and } \rho > 1. \end{cases}$$

Using Theorem 1.4, we see that there exists a constant C_0 such that

$$M_0(T) \leq C_0 \delta,$$

for any T . We claim that, for $k \geq 1$, we have

$$(5.3) \quad M_k(T_\delta) \leq 2C_0 \delta,$$

for δ sufficiently small and T_δ appropriately chosen. We will prove this inductively. Assume that the bound holds for $k-1$. By Theorem 1.4, we have, for δ small enough,

$$\begin{aligned}
M_k(T_\delta) & \leq C_0 \delta + C \sum_{|\alpha| \leq M} \int_0^{T_\delta} \|Z^\alpha Q(u'_{k-1})(s, \cdot)\|_{L^2(\mathbb{R}^d)} ds \\
& \leq C_0 \delta + C \sum_{|\alpha| \leq M} \int_0^{T_\delta} \|\langle x \rangle^{-\mu_d} Z^\alpha u'_{k-1}\|_{L^2(\mathbb{R}^d)}^2 ds \\
& \leq C_0 \delta + CK_n(T_\delta) M_{k-1}^2(T_\delta) \\
& \leq C_0 \delta + CK_n(T_\delta) (2C_0 \delta)^2,
\end{aligned}$$

where we have also used Lemma 5.2 and the induction hypothesis. Note that, to estimate the term $\|(Z^\alpha u_k)'(0, \cdot)\|_{L^2}$, we might have to use the equation and Lemma 5.2. We therefore need δ to be small enough. Then, to prove (5.3), it is enough to have

$$(5.4) \quad C_0\delta + CK_n(T_\delta)(2C_0\delta)^2 \leq 2C_0\delta \iff 4CC_0K_n(T_\delta)\delta \leq 1.$$

Therefore, we find:

- If $d = 3$ or $\rho = 1$, the estimate holds with $T_\delta = c_n\delta^{-n}$ and c_n small enough.
- If $d \geq 4$ and $\rho > 1$, (5.4) is fulfilled if δ is sufficiently small and we can take $T_\delta = \infty$.

To show that the sequence u_k converges, we estimate the quantity

$$A_k(T) = \sup_{0 \leq t \leq T} \sum_{1 \leq i+j \leq M+1} \|\partial_t^i P^{j/2}(u_k - u_{k-1})\|_{L^2(\mathbb{R}^d)} + \sum_{|\alpha| \leq M} K_n(T)^{-1} \|\langle x \rangle^{-\mu_d} Z^\alpha (u'_k - u'_{k-1})\|_{L^2([0, T] \times \mathbb{R}^d)}.$$

It is clearly sufficient to show

$$(5.5) \quad A_k(T) \leq \frac{1}{2} A_{k-1}(T).$$

Using Lemma 5.2 and repeating the above arguments, we obtain

$$A_k(T_\delta) \leq \tilde{C} \sum_{|\alpha| \leq M} \int_0^{T_\delta} \|\langle x \rangle^{-\mu_d} Z^\alpha (u'_{k-1} - u'_{k-2})\|_{L^2(\mathbb{R}^d)} \times \sum_{|\alpha| \leq M} \left(\|\langle x \rangle^{-\mu_d} Z^\alpha u'_{k-1}\|_{L^2(\mathbb{R}^d)} + \|\langle x \rangle^{-\mu_d} Z^\alpha u'_{k-2}\|_{L^2(\mathbb{R}^d)} \right) ds.$$

By the Cauchy–Schwarz inequality, we conclude that

$$A_k(T_\delta) \leq \tilde{C} K_n(T_\delta) (M_{k-1}(T_\delta) + M_{k-2}(T_\delta)) A_{k-1}(T_\delta).$$

Using (5.3), the above inequality leads to (5.5) if δ is small enough. Uniqueness and C^2 property of the solution follow from [16, Theorem 6.4.10, Theorem 6.4.11] using that the constructed solution is in $H_{loc}^{M+1}(\mathbb{R}^{d+1}) \subset C^2(\mathbb{R}^{d+1})$. Note also that the solution is bounded in C^2 on the interval $[0, T_\delta]$. \square

APPENDIX A. REGULARITY

Here, we give some results concerning the regularity with respect to an operator. More details can be found in the book of Amrein, A. Boutet de Monvel and Georgescu [3] and in the paper of C. Gérard and Georgescu [10]. We start with a useful characterization of the regularity $C^1(A)$.

Theorem A.1 ([3, Theorem 6.2.10]). *Let A and H be self-adjoint operators on a Hilbert space \mathcal{H} . Then H is of class $C^1(A)$ iff the following conditions are satisfied:*

i) there is a constant $c < \infty$ such that for all $u \in D(A) \cap D(H)$,

$$|(Au, Hu) - (Hu, Au)| \leq c (\|Hu\|^2 + \|u\|^2),$$

ii) for some $z \in \mathbb{C} \setminus \sigma(H)$, the set $\{u \in D(A); (H - z)^{-1}u \in D(A) \text{ and } (H - \bar{z})^{-1}u \in D(A)\}$ is a core for A .

If H is of class $C^1(A)$, then the following is true:

- i) The space $(H - z)^{-1}D(A)$ is independent of $z \in \mathbb{C} \setminus \sigma(H)$ and contained in $D(A)$. It is a core for H and a dense subspace of $D(A) \cap D(H)$ for the intersection topology (i.e. the topology associated to the norm $\|Hu\| + \|Au\| + \|u\|$).
- ii) The space $D(A) \cap D(H)$ is a core for H and the form $[A, H]$ has a unique extension to a continuous sesquilinear form on $D(H)$ (equipped with the graph topology). If this extension is denoted by $[A, H]$, the following identity holds on \mathcal{H} (in the form sense):

$$[A, (H - z)^{-1}] = -(H - z)^{-1}[A, H](H - z)^{-1},$$

for $z \in \mathbb{C} \setminus \sigma(H)$.

We also have the following theorem coming from [3, Theorem 6.3.4].

Theorem A.2. *Let A and H be self-adjoint operators in a Hilbert space \mathcal{H} . Assume that the unitary one-parameter group $\{\exp(iA\tau)\}_{\tau \in \mathbb{R}}$ leaves the domain $D(H)$ of H invariant. Then H is of class $C^1(A)$ iff $[H, A]$ is bounded from $D(H)$ to $D(H)^*$.*

A criterion for the above assumption to be satisfied is given by the following result of Georgescu and C. Gérard.

Lemma A.3 ([10, Lemma 2]). *Let A and H be self-adjoint operators in a Hilbert space \mathcal{H} . Let $H \in C^1(A)$ and suppose that the commutator $[iH, A]$ can be extended to a bounded operator from $D(H)$ to \mathcal{H} . Then e^{itA} preserves $D(H)$.*

In this paper, we will use the following characterization of the regularity $C^2(A)$.

Remark A.4. *From Section 6.2 of [3], it is known that H is of class $C^2(A)$ if the following conditions hold:*

- i) *For some $z \in \mathbb{C} \setminus \sigma(H)$, the set $\{u \in D(A); (H - z)^{-1}u \in D(A) \text{ and } (H - \bar{z})^{-1}u \in D(A)\}$ is a core for A .*
- ii) *$[H, A]$ and $[[H, A], A]$ extend as bounded operators on \mathcal{H} .*

APPENDIX B. RESOLVENT ESTIMATES AT LOW ENERGIES

B.1. Estimates for the free Laplacian.

We begin with some estimates for the free Laplacian $P_0 = -\Delta$.

Lemma B.1. *Let $\alpha > 0$. Then, for all $\varepsilon > 0$, we have*

$$\|(\lambda P_0 + 1)^{-\alpha} u\| \lesssim \lambda^{-\min(\alpha - \varepsilon, d/4)} \|\langle x \rangle^{\min(2\alpha, d/2 + \varepsilon)} u\|,$$

uniformly for $\lambda \geq 1$.

Proof. Here, we denote

$$\|u\|_p = \left(\int_{\mathbb{R}^d} |u(x)|^p dx \right)^{1/p},$$

the standard norm on $L^p(\mathbb{R}^d)$. Using the Hölder inequality, we get

$$\begin{aligned} \|(\lambda P_0 + 1)^{-\alpha} u\| &= \left(\int (\lambda \xi^2 + 1)^{-2\alpha} |\widehat{u}(\xi)|^2 d\xi \right)^{1/2} \\ &\leq \|(\lambda \xi^2 + 1)^{-\alpha}\|_{2p} \|\widehat{u}\|_{2q}, \end{aligned}$$

and we choose $p = \max(\frac{d}{4\alpha} + \mu, 1)$, $\mu > 0$, and $p^{-1} + q^{-1} = 1$. In particular, $2q \geq 2$ and $4\alpha p > d$. Then, by the Hausdorff–Young inequality, we obtain

$$(B.1) \quad \|(\lambda P_0 + 1)^{-\alpha} u\| \lesssim \lambda^{-d/4p} \|u\|_r,$$

with $r^{-1} = 1 - (2q)^{-1} = 2^{-1}(1 + p^{-1})$ satisfying $1 \leq r \leq 2$. Using one more time the Hölder inequality, we have

$$\|u\|_r \lesssim \left(\int |u|^{rs} \langle x \rangle^{\beta s} dx \right)^{1/rs} \left(\int \langle x \rangle^{-\beta t} dx \right)^{1/rt},$$

with $s^{-1} + t^{-1} = 1$ and $\beta > 0$. We choose $s = 2r^{-1}$ and $\beta = d/t + \nu$, $\nu > 0$. Thus,

$$(B.2) \quad \|u\|_r \lesssim \|\langle x \rangle^{\beta s/2} u\|_2.$$

The coefficient $\beta s/2$ satisfies

$$\begin{aligned} \frac{\beta s}{2} &= \frac{ds}{2t} + \frac{\nu s}{2} = \frac{ds}{2} - \frac{d}{2} + \frac{\nu s}{2} = \frac{d}{r} - \frac{d}{2} + \mathcal{O}(\nu) = \frac{d}{2p} + \mathcal{O}(\nu) \\ &= \frac{d}{2} \min\left(\left(\frac{d}{4\alpha} + \mu\right)^{-1}, 1\right) + \mathcal{O}(\nu) = \frac{d}{2} \min\left(\frac{4\alpha}{d} - \frac{16\alpha^2\mu}{d^2} + \mathcal{O}(\mu^2), 1\right) + \mathcal{O}(\nu) \\ &= \min\left(2\alpha - \frac{8\alpha^2\mu}{d} + \mathcal{O}(\mu^2), \frac{d}{2}\right) + \mathcal{O}(\nu). \end{aligned}$$

On the other hand,

$$\frac{d}{4p} = \frac{d}{4} \min\left(\left(\frac{d}{4\alpha} + \mu\right)^{-1}, 1\right) = \min\left(\alpha + \mathcal{O}(\mu), \frac{d}{4}\right).$$

Taking first μ and then ν small enough, the lemma follows from the estimates (B.1) and (B.2). \square

Lemma B.2. *Let $\beta \geq 0$, $0 \leq \gamma \leq \min(1, d/4)$ and $0 \leq \delta \leq d/4$. Then, for all $\varepsilon > 0$,*

$$\|\langle x \rangle^\beta (\lambda P_0 + 1)^{-1} u\| \lesssim \lambda^{\beta/2 - \delta + \varepsilon} \|\langle x \rangle^{2\delta} u\| + \lambda^{-\gamma + \varepsilon} \|\langle x \rangle^{\beta + 2\gamma} u\|,$$

uniformly for $\lambda \geq 1$.

Remark B.3. *In the previous lemma, assume $\gamma + \beta/2 \leq d/4$. Then, we can chose $\delta = \gamma + \beta/2$ and we have*

$$\|\langle x \rangle^\beta (\lambda P_0 + 1)^{-1} u\| \lesssim \lambda^{-\gamma + \varepsilon} \|\langle x \rangle^{\beta + 2\gamma} u\|,$$

uniformly for $\lambda \geq 1$.

Proof. Assume first that $\beta \in \mathbb{N}$. Using

$$\|\langle x \rangle u\|^2 = \langle (x^2 + 1)u, u \rangle = \sum_{j=1}^d \|x_j u\|^2 + \|u\|^2,$$

it is enough to estimate $\|x^a (\lambda P_0 + 1)^{-1} u\|$ where $a \in \mathbb{N}^d$ is a multi-index of length less or equal to β . Since

$$x_j (\lambda P_0 + 1)^{-1} = (\lambda P_0 + 1)^{-1} x_j - 2\lambda^{1/2} (\lambda^{1/2} \partial_j) (\lambda P_0 + 1)^{-2},$$

the operator $x^a (\lambda P_0 + 1)^{-1}$ can be written as a finite sum of terms of the form

$$T = \lambda^{\frac{|a|-|b|}{2}} (\lambda^{1/2} \partial)^c (\lambda P_0 + 1)^{-1 - \frac{|a+c-b|}{2}} x^b,$$

where b, c are non-negative multi-indexes such that $b + c \leq a$ and $|a + c - b| = |a| + |c| - |b|$ is even. Such a term can be written as

$$\begin{aligned} T &= \lambda^{\frac{|a|-|b|}{2}} (\lambda^{1/2} \partial)^c (\lambda P_0 + 1)^{-1 - \frac{|a+c-b|}{2}} (\lambda P_0 + 1)^{1 + \frac{|a|-|b|}{2}} (\lambda P_0 + 1)^{-1 - \frac{|a|-|b|}{2}} x^b \\ &= \lambda^{\frac{|a|-|b|}{2}} B (\lambda P_0 + 1)^{-1 - \frac{|a|-|b|}{2}} x^b, \end{aligned}$$

where B is a bounded operator on $L^2(\mathbb{R}^d)$ since it is a Fourier multiplier by a uniformly bounded function.

Using Lemma B.1 to estimate the powers of the resolvent, we get

$$(B.3) \quad T = B \lambda^{\frac{|a|-|b|}{2} - \min(\alpha, d/4) + \varepsilon} \langle x \rangle^{|b| + \min(2\alpha, d/2 + \varepsilon)}$$

where B is an other bounded operator, $0 < \varepsilon$ and $0 \leq \alpha \leq 1 + (|a| - |b|)/2$. We choose $\alpha = \min(\gamma + (|a| - |b|)/2, \delta) \leq d/4$ and note $b_0 = |a| + 2\gamma - 2\delta$.

If $|b| < b_0$, then $\alpha = \delta$ and (B.3) becomes

$$\begin{aligned} T &= B \lambda^{\frac{|a|-|b|}{2} - \delta + \varepsilon} \langle x \rangle^{|b| + 2\delta} \\ &= \mathcal{O}(\lambda^{\frac{|a|}{2} - \delta + \varepsilon} \langle x \rangle^{2\delta} + \lambda^{\frac{|a|-b_0}{2} - \delta + \varepsilon} \langle x \rangle^{b_0 + 2\delta}), \end{aligned}$$

since $y^{|b|} \leq y^{b_0} + y^0$ for $0 \leq b \leq b_0$ and $y \geq 0$. Using $|a| \leq \beta$, we get

$$(B.4) \quad T = \mathcal{O}(\lambda^{\beta/2 - \delta + \varepsilon} \langle x \rangle^{2\delta} + \lambda^{-\gamma + \varepsilon} \langle x \rangle^{\beta + 2\gamma}).$$

If $|b| \geq b_0$, then $\alpha = \gamma + (|a| - |b|)/2$ and (B.3) gives

$$(B.5) \quad T = \mathcal{O}(\lambda^{-\gamma + \varepsilon} \langle x \rangle^{\beta + 2\gamma}).$$

The estimates (B.4) and (B.5) imply the lemma for $\beta \in \mathbb{N}$. The case $\beta \in \mathbb{R}^+$ follows from an interpolation argument. \square

Mimicking the previous proofs, one can show the following results

Lemma B.4. *Let $j \in \{1, \dots, d\}$, $\beta \geq 0$ and $0 \leq \gamma \leq 1/2$ with $\gamma + \beta/2 \leq d/4$. Then, for all $\varepsilon > 0$, we have*

$$\|\langle x \rangle^\beta (\lambda^{1/2} \partial_j) (\lambda P_0 + 1)^{-1} u\| \lesssim \lambda^{-\gamma + \varepsilon} \|\langle x \rangle^{\beta + 2\gamma} u\|,$$

uniformly for $\lambda \geq 1$.

Lemma B.5. *Let $j, k \in \{1, \dots, d\}$ and $0 \leq \beta/2 \leq d/4$. Then, for all $\varepsilon > 0$, we have*

$$\|\langle x \rangle^\beta (\lambda^{1/2} \partial_j) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial_k) u\| \lesssim \lambda^\varepsilon \|\langle x \rangle^\beta u\|,$$

uniformly for $\lambda \geq 1$.

B.2. Estimates for an intermediate operator.

We now extend these results to the intermediate differential operator \tilde{P} defined by

$$(B.6) \quad \tilde{P} = - \sum_{j,k} \partial_j g^2 g^{j,k} \partial_k.$$

Recall from (H1) that $g^2 g^{j,k} - \delta_{j,k} = \mathcal{O}(\langle x \rangle^{-\rho})$. The square roots of \tilde{P} and P_0 are comparable. More precisely, we have

Lemma B.6. For $u \in D(\tilde{P}^{1/2}) = D(P_0^{1/2}) = H^1(\mathbb{R}^d)$,

$$\|\tilde{P}^{1/2}u\| \lesssim \|P_0^{1/2}u\| \lesssim \|\tilde{P}^{1/2}u\|.$$

Proof. For $u \in H^2(\mathbb{R}^d)$, we can write

$$(\tilde{P}u, u) = \sum_{j,k} (g^2 g^{j,k} \partial_j u, \partial_k u) \quad \text{and} \quad (P_0 u, u) = \sum_j (\partial_j u, \partial_j u).$$

Using the ellipticity of \tilde{P} and $g^2 g^{j,k} \in L^\infty(\mathbb{R}^d)$, we get

$$(P_0 u, u) \lesssim (\tilde{P}u, u) \lesssim (P_0 u, u).$$

In particular, we have, for $u \in H^2(\mathbb{R}^d)$,

$$\begin{aligned} \|\tilde{P}^{1/2}u\| &\lesssim \|P_0^{1/2}u\| \lesssim \|\tilde{P}^{1/2}u\| \\ \|(\tilde{P} + 1)^{1/2}u\| &\lesssim \|(P_0 + 1)^{1/2}u\| \lesssim \|(\tilde{P} + 1)^{1/2}u\|. \end{aligned}$$

Then, we obtain $D(\tilde{P}^{1/2}) = D(P_0^{1/2}) = H^1(\mathbb{R}^d)$ and the lemma follows. \square

Lemma B.7. Let $\beta \geq 0$ and $0 \leq \gamma \leq \min(1, d/4)$ with $\gamma + \beta/2 \leq d/4$. Then, for all $\varepsilon > 0$, we have

$$(B.7) \quad \|\langle x \rangle^\beta (\lambda \tilde{P} + 1)^{-1} u\| \lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{\beta+2\gamma} u\|,$$

uniformly for $\lambda \geq 1$.

Remark B.8. Mimicking the proof of Lemma B.7, one can show that Lemma B.4 (for the operators $(\lambda^{1/2} \partial_j)(\lambda \tilde{P} + 1)^{-1}$ and $(\lambda \tilde{P} + 1)^{-1}(\lambda^{1/2} \partial_j)$) and Lemma B.5 hold with P_0 replaced by \tilde{P} .

Proof. From (B.6), we have

$$P_0 - \tilde{P} = \sum_{j,k} \partial_j r_{j,k} \partial_k,$$

where $r_{j,k} = \delta_{j,k} - g^2 g^{j,k} = \mathcal{O}(\langle x \rangle^{-\rho})$. In the following, to clarify the statement, we will not write the sum over j, k and simply note $P_0 - \tilde{P} = \partial r \partial$. Iterating the resolvent identity, we have

$$\begin{aligned} (\lambda \tilde{P} + 1)^{-1} &= (\lambda P_0 + 1)^{-1} + (\lambda P_0 + 1)^{-1} \lambda^{1/2} \partial r \lambda^{1/2} \partial (\lambda P_0 + 1)^{-1} \\ &\quad + \sum_{j=1}^{2N} (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) \left(r (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) \right)^j r (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} \\ &\quad + (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) \left(r (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) \right)^N \\ &\quad \times r (\lambda^{1/2} \partial) (\lambda \tilde{P} + 1)^{-1} (\lambda^{1/2} \partial) r \\ (B.8) \quad &\quad \times \left((\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) r \right)^N (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1}. \end{aligned}$$

Thanks to Remark B.3, the first term of the previous equation satisfies (B.7). To treat the second term, we use two times Lemma B.4 with a gain equal to $\gamma/2 \leq \max(1/2, d/4)$.

The sum over j can be studied in a similar way: using Lemma B.4, each exterior term $(\lambda^{1/2}\partial_j)(\lambda P_0 + 1)^{-1}$ gives a factor $\lambda^{-\gamma/2+\tilde{\varepsilon}}$, and, using Lemma B.5, each interior factor $(\lambda^{1/2}\partial)(\lambda P_0 + 1)^{-1}(\lambda^{1/2}\partial)$ gives a factor $\lambda^{\tilde{\varepsilon}}$. Then, each term in the sum over j can be estimated by $\lambda^{-\gamma+(j+2)\tilde{\varepsilon}}$. Taking $\tilde{\varepsilon} = \varepsilon/(2N + 2)$, each term of the sum over j satisfies (B.7).

It remains to study the last term in (B.8). As usual, the first term can be estimated by Lemma B.4:

$$\|\langle x \rangle^\beta (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) u\| \lesssim \lambda^{-\gamma/2+\tilde{\varepsilon}} \|\langle x \rangle^{\beta+\gamma} u\|.$$

Now, using $r = \mathcal{O}(\langle x \rangle^{-\rho})$ together with Lemma B.5, we get

$$(B.9) \quad \|\langle x \rangle^\mu r (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) u\| \lesssim \lambda^{\tilde{\varepsilon}} \|\langle x \rangle^{\max(\mu-\rho, 0)} u\|,$$

for $\mu/2 \leq d/4 + \rho/2$. Using N times the last inequality, we obtain

$$(B.10) \quad \begin{aligned} & \left\| \langle x \rangle^\beta (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) \left(r (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) \right)^N u \right\| \\ & \lesssim \lambda^{-\gamma/2+(N+1)\tilde{\varepsilon}} \|\langle x \rangle^{\max(\beta+\gamma-\rho N, 0)} u\| \leq \lambda^{-\gamma/2+(N+1)\tilde{\varepsilon}} \|u\|, \end{aligned}$$

for N large enough. Using two times Lemma B.6 and the functional calculus,

$$(B.11) \quad \|(\lambda^{1/2} \partial) (\lambda \tilde{P} + 1)^{-1} (\lambda^{1/2} \partial) u\| \lesssim \|u\|.$$

Finally, applying N times (B.9), with N large enough, we get

$$(B.12) \quad \left\| \langle x \rangle^\gamma \left(r (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) \right)^N \right\| \lesssim \lambda^{N\tilde{\varepsilon}} \|u\|,$$

since $\gamma \leq d/4$. Moreover, using $\gamma/2 \leq 1/2$ and taking the adjoint in Lemma B.4, we have

$$\|\langle x \rangle^{-\gamma} (\lambda^{1/2} \partial_j) (\lambda P_0 + 1)^{-1} u\| \lesssim \lambda^{-\gamma/2+\tilde{\varepsilon}} \|u\|.$$

Combining the last estimate with the adjoint of (B.12), it follows

$$(B.13) \quad \left\| \left((\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} (\lambda^{1/2} \partial) r \right)^N (\lambda^{1/2} \partial) (\lambda P_0 + 1)^{-1} \right\| \lesssim \lambda^{-\gamma/2+(N+1)\tilde{\varepsilon}} \|u\|,$$

for N large enough. Summing up (B.10), (B.11), (B.13) and choosing $\tilde{\varepsilon}$ small enough with respect to ε , the last term in (B.8) satisfies (B.7). \square

B.3. Estimates for the perturbed Laplacian.

Here, we extend the previous results to the Laplacian P . From (B.6), we have $P = g^{-1} \tilde{P} g^{-1}$. In particular, the resolvent identity gives

$$(B.14) \quad \begin{aligned} (\lambda P + 1)^{-1} &= g (\lambda \tilde{P} + g^2)^{-1} g \\ &= g (\lambda \tilde{P} + 1)^{-1} g + g (\lambda \tilde{P} + 1)^{-1} (1 - g^2) g^{-1} (\lambda P + 1)^{-1} \end{aligned}$$

$$(B.15) \quad = g (\lambda \tilde{P} + 1)^{-1} g + (\lambda P + 1)^{-1} g^{-1} (1 - g^2) (\lambda \tilde{P} + 1)^{-1} g.$$

Note that, by (H1), $(1 - g^2) = \mathcal{O}(\langle x \rangle^{-\rho})$.

Proposition B.9. *Let $\beta \geq 0$ and $0 \leq \gamma \leq 1$ with $\gamma + \beta/2 \leq d/4$. Then, for all $\varepsilon > 0$, we have*

$$\|\langle x \rangle^\beta (\lambda P + 1)^{-1} u\| \lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{\beta+2\gamma} u\|,$$

uniformly for $\lambda \geq 1$.

Proof. As in the proof of Lemma B.7, we iterate the resolvent identity (B.14) and obtain

$$(B.16) \quad \begin{aligned} (\lambda P + 1)^{-1} &= g(\lambda \tilde{P} + 1)^{-1}g + g(\lambda \tilde{P} + 1)^{-1}(1 - g^2)(\lambda \tilde{P} + 1)^{-1}g \\ &\quad + \sum_{j=1}^N g(\lambda \tilde{P} + 1)^{-1} \left((1 - g^2)(\lambda \tilde{P} + 1)^{-1} \right)^j (1 - g^2)(\lambda \tilde{P} + 1)^{-1}g \\ &\quad + g(\lambda \tilde{P} + 1)^{-1} \left((1 - g^2)(\lambda \tilde{P} + 1)^{-1} \right)^{N+1} (1 - g^2)g^{-1}(\lambda P + 1)^{-1}. \end{aligned}$$

The two first terms and the sum over j can be directly estimated by Lemma B.7. For the last term in (B.16), we remark that Lemma B.7 gives

$$(B.17) \quad \left\| \langle x \rangle^\beta g(\lambda \tilde{P} + 1)^{-1}u \right\| \lesssim \lambda^{-\gamma + \tilde{\varepsilon}} \left\| \langle x \rangle^{\beta + 2\gamma}u \right\|,$$

and

$$(B.18) \quad \left\| \langle x \rangle^\mu (1 - g^2)(\lambda \tilde{P} + 1)^{-1}u \right\| \lesssim \lambda^{\tilde{\varepsilon}} \left\| \langle x \rangle^{\max(\mu - \rho, 0)}u \right\|,$$

for all $\mu/2 \leq d/4 + \rho/2$. Therefore, applying (B.17) and $N + 1$ times (B.18) (this can be made since $(\beta + 2\gamma)/2 \leq d/4$), we get

$$\begin{aligned} \left\| \langle x \rangle^\beta g(\lambda \tilde{P} + 1)^{-1} \left((1 - g^2)(\lambda \tilde{P} + 1)^{-1} \right)^{N+1} u \right\| &\lesssim \lambda^{-\gamma + (N+2)\tilde{\varepsilon}} \left\| \langle x \rangle^{\max(\mu - (N+1)\rho, 0)}u \right\| \\ &\lesssim \lambda^{-\gamma + (N+2)\tilde{\varepsilon}} \|u\|, \end{aligned}$$

for N large enough. Using $\|(\lambda P + 1)^{-1}\| \leq 1$ by the spectral theorem and taking $\tilde{\varepsilon} = \varepsilon/(N+2)$, this implies

$$\left\| \langle x \rangle^\beta g(\lambda \tilde{P} + 1)^{-1} \left((1 - g^2)(\lambda \tilde{P} + 1)^{-1} \right)^{N+1} (1 - g^2)g^{-1}(\lambda P + 1)^{-1}u \right\| \lesssim \lambda^{-\gamma + \varepsilon} \|u\|,$$

and the lemma follows. \square

Mimicking the proof of Proposition B.9 and using (B.15) and Remark B.8, one can prove, as for Lemma B.4, the following result.

Lemma B.10. *Let $j \in \{1, \dots, d\}$, $\beta \geq 0$ and $0 \leq \gamma \leq 1/2$ with $\gamma + \beta/2 \leq d/4$. Then, for all $\varepsilon > 0$, we have*

$$\begin{aligned} \left\| \langle x \rangle^\beta (\lambda P + 1)^{-1} (\lambda^{1/2} \tilde{\partial}_j^*) u \right\| &\lesssim \lambda^{-\gamma + \varepsilon} \left\| \langle x \rangle^{\beta + 2\gamma}u \right\| \\ \left\| \langle x \rangle^\beta (\lambda^{1/2} \tilde{\partial}_j) (\lambda P + 1)^{-1}u \right\| &\lesssim \lambda^{-\gamma + \varepsilon} \left\| \langle x \rangle^{\beta + 2\gamma}u \right\|. \end{aligned}$$

uniformly for $\lambda \geq 1$.

Let $j, k \in \{1, \dots, d\}$ and $0 \leq \beta/2 \leq d/4$. Then, for all $\varepsilon > 0$, we have

$$\left\| \langle x \rangle^\beta (\lambda^{1/2} \tilde{\partial}_j) (\lambda P + 1)^{-1} (\lambda^{1/2} \tilde{\partial}_k^*) u \right\| \lesssim \lambda^\varepsilon \left\| \langle x \rangle^\beta u \right\|,$$

uniformly for $\lambda \geq 1$.

Remark B.11. *The results of this section are given for $(\lambda P + 1)^{-1}$, but can be extended to $(\lambda P - z)^{-1}$, with $\text{Im } z \neq 0$. In fact, following the previous proofs, one can see that $(\lambda P - z)^{-1}$ satisfies the same results, if we accept a lose of the form $|\text{Im } z|^{-C}$, $C > 0$, in the estimates. This is due to $(\lambda P_0 + 1)(\lambda P_0 - z)^{-1} = \mathcal{O}(|\text{Im } z|^{-1})$ from the spectral theorem. Note that the constant C does not depend on $\varepsilon \in]0, 1]$, and is uniform with respect to $\alpha, \beta, \gamma, \delta$ in a compact subset.*

For example, Proposition B.9 gives the following estimate for $\beta \geq 0$, $\varepsilon > 0$ and $0 \leq \gamma \leq 1$ with $\gamma + \beta/2 \leq d/4$:

$$(B.19) \quad \|\langle x \rangle^\beta (\lambda P - z)^{-1} u\| \lesssim \frac{\lambda^{-\gamma+\varepsilon}}{|\operatorname{Im} z|^C} \|\langle x \rangle^{\beta+2\gamma} u\|,$$

uniformly for $\lambda \geq 1$ and z in a compact of \mathbb{C} .

Using the spectral theorem, this remark implies the following result.

Lemma B.12. *Let $\chi \in C_0^\infty(\mathbb{R})$, $j, k \in \{1, \dots, d\}$ and $\beta, \gamma \geq 0$ with $\gamma + \beta/2 \leq d/4$. Then, for all $\varepsilon > 0$, we have*

$$\begin{aligned} \|\langle x \rangle^\beta \chi(\lambda P) u\| &\lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{\beta+2\gamma} u\| \\ \|\langle x \rangle^\beta (\lambda^{1/2} \tilde{\partial}_j) \chi(\lambda P) u\| &\lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{\beta+2\gamma} u\| \\ \|\langle x \rangle^\beta \chi(\lambda P) (\lambda^{1/2} \tilde{\partial}_j^*) u\| &\lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{\beta+2\gamma} u\| \\ \|\langle x \rangle^\beta (\lambda^{1/2} \tilde{\partial}_j) \chi(\lambda P) (\lambda^{1/2} \tilde{\partial}_k^*) u\| &\lesssim \lambda^{-\gamma+\varepsilon} \|\langle x \rangle^{\beta+2\gamma} u\|, \end{aligned}$$

uniformly for $\lambda \geq 1$.

Proof. We only prove the first inequality since the others can be treated the same way. Let $k \in \mathbb{N}$ be such that $\gamma/k \leq 1$, $\varphi(\sigma) = \chi(\sigma)(\sigma + 1)^k \in C_0^\infty(\mathbb{R})$ and $\tilde{\varphi} \in C_0^\infty(\mathbb{C})$ be an almost analytic extension of φ . From the spectral theorem, we have

$$(B.20) \quad \langle x \rangle^\beta \chi(\lambda P) = \frac{1}{\pi} \int \bar{\partial} \tilde{\varphi}(z) \langle x \rangle^\beta (\lambda P - z)^{-1} (\lambda P + 1)^{-k} L(dz).$$

Estimate (B.19) with $\gamma = 0$ gives

$$(B.21) \quad \|\langle x \rangle^\beta (\lambda P - z)^{-1} u\| \lesssim \frac{\lambda^{\tilde{\varepsilon}}}{|\operatorname{Im} z|^C} \|\langle x \rangle^\beta u\|.$$

Proposition B.9 with $\gamma = \gamma/k \leq 1$ implies

$$\|\langle x \rangle^\mu (\lambda P + 1)^{-1} u\| \lesssim \lambda^{-\gamma/k+\tilde{\varepsilon}} \|\langle x \rangle^{\mu+2\gamma/k} u\|,$$

if $\gamma/k + \mu/2 \leq d/4$. By iteration, we obtain

$$\|\langle x \rangle^\beta (\lambda P + 1)^{-k} u\| \lesssim \lambda^{-\gamma+k\tilde{\varepsilon}} \|\langle x \rangle^{\beta+2\gamma} u\|,$$

since $\gamma + \beta/2 \leq d/4$. Combining this estimate with (B.21) and taking $\tilde{\varepsilon} = \varepsilon/(k+1)$, we get

$$\|\langle x \rangle^\beta (\lambda P - z)^{-1} (\lambda P + 1)^{-k} u\| \lesssim \frac{\lambda^{-\gamma+\varepsilon}}{|\operatorname{Im} z|^C} \|\langle x \rangle^{\beta+2\gamma} u\|,$$

and the lemma follows from (B.20). \square

We now state a result which will help us to estimate the square root of P . Since this lemma can be proved as Lemma B.6, we do not give the proof.

Lemma B.13. *We have, for $u \in D(P^{1/2}) = H^1(\mathbb{R}^d)$,*

$$\|P^{1/2} u\| \lesssim \|\nabla g^{-1} u\| \lesssim \|P^{1/2} u\|.$$

REFERENCES

1. S. Alinhac, *Free decay of solutions to wave equations on a curved background*, Bull. Soc. Math. France **133** (2005), no. 3, 419–458.
2. S. Alinhac, *On the Morawetz–Keel–Smith–Sogge inequality for the wave equation on a curved background*, Publ. Res. Inst. Math. Sci. **42** (2006), no. 3, 705–720.
3. W. Amrein, A. Boutet de Monvel, and V. Georgescu, *C_0 -groups, commutator methods and spectral theory of N -body Hamiltonians*, Progress in Mathematics, vol. 135, Birkhäuser Verlag, 1996.
4. R. Beals, *Weighted distribution spaces and pseudodifferential operators*, J. Analyse Math. **39** (1981), 131–187.
5. N. Burq, *Global Strichartz estimates for nontrapping geometries: about an article by H. Smith and C. Sogge*, Comm. Partial Differential Equations **28** (2003), no. 9–10, 1675–1683.
6. M. Christ and A. Kiselev, *WKB asymptotic behavior of almost all generalized eigenfunctions for one-dimensional Schrödinger operators with slowly decaying potentials*, J. Funct. Anal. **179** (2001), no. 2, 426–447.
7. J. Dereziński and C. Gérard, *Scattering theory of classical and quantum N -particle systems*, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997.
8. M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series, vol. 268, Cambridge University Press, Cambridge, 1999.
9. H. Donnelly, *Exhaustion functions and the spectrum of Riemannian manifolds*, Indiana Univ. Math. J. **46** (1997), no. 2, 505–527.
10. V. Georgescu and C. Gérard, *On the virial theorem in quantum mechanics*, Comm. Math. Phys. **208** (1999), no. 2, 275–281.
11. C. Gérard and I. Laba, *Multiparticle quantum scattering in constant magnetic fields*, Mathematical Surveys and Monographs, vol. 90, American Mathematical Society, Providence, RI, 2002.
12. C. Gérard and A. Martinez, *Principe d'absorption limite pour des opérateurs de Schrödinger à longue portée*, C. R. Acad. Sci. Paris Sér. I Math. **306** (1988), no. 3, 121–123.
13. D. Häfner, *Complétude asymptotique pour l'équation des ondes dans une classe d'espaces-temps stationnaires et asymptotiquement plats*, Ann. Inst. Fourier **51** (2001), no. 3, 779–833.
14. B. Helffer and F. Nier, *Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians*, Lecture Notes in Mathematics, vol. 1862, Springer-Verlag, Berlin, 2005.
15. L. Hörmander, *The analysis of linear partial differential operators. III*, Grundlehren der Mathematischen Wissenschaften, vol. 274, Springer-Verlag, Berlin, 1985, Pseudodifferential operators.
16. L. Hörmander, *Lectures on nonlinear hyperbolic differential equations*, Mathématiques & Applications, vol. 26, Springer-Verlag, Berlin, 1997.
17. F. John, *Blow-up for quasilinear wave equations in three space dimensions*, Comm. Pure Appl. Math. **34** (1981), no. 1, 29–51.
18. F. John and S. Klainerman, *Almost global existence to nonlinear wave equations in three space dimensions*, Comm. Pure Appl. Math. **37** (1984), no. 4, 443–455.
19. T. Kato, *Wave operators and similarity for some non-selfadjoint operators*, Math. Ann. **162** (1965/1966), 258–279.
20. M. Keel, H. Smith, and C. Sogge, *Almost global existence for some semilinear wave equations*, J. Anal. Math. **87** (2002), 265–279, Dedicated to the memory of Thomas H. Wolff.
21. M. Keel, H. Smith, and C. Sogge, *Almost global existence for quasilinear wave equations in three space dimensions*, J. Amer. Math. Soc. **17** (2004), no. 1, 109–153 (electronic).
22. S. Klainerman, *Uniform decay estimates and the Lorentz invariance of the classical wave equation*, Comm. Pure Appl. Math. **38** (1985), no. 3, 321–332.
23. S. Klainerman and G. Ponce, *Global, small amplitude solutions to nonlinear evolution equations*, Comm. Pure Appl. Math. **36** (1983), no. 1, 133–141.
24. J. Metcalfe and C. Sogge, *Hyperbolic trapped rays and global existence of quasilinear wave equations*, Invent. Math. **159** (2005), no. 1, 75–117.
25. J. Metcalfe and C. Sogge, *Long-time existence of quasilinear wave equations exterior to star-shaped obstacles via energy methods*, SIAM J. Math. Anal. **38** (2006), no. 1, 188–209 (electronic).
26. J. Metcalfe and D. Tataru, *Decay estimates for variable coefficient wave equations in exterior domains*, arXiv:0806.3409, (2008).

27. E. Mourre, *Absence of singular continuous spectrum for certain selfadjoint operators*, Comm. Math. Phys. **78** (1980/81), no. 3, 391–408.
28. P. Perry, I. M. Sigal, and B. Simon, *Spectral analysis of N -body Schrödinger operators*, Ann. of Math. (2) **114** (1981), no. 3, 519–567.
29. M. Reed and B. Simon, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*, Academic Press, 1975.
30. M. Reed and B. Simon, *Methods of modern mathematical physics. IV. Analysis of operators*, Academic Press, 1978.
31. T. Sideris, *Global behavior of solutions to nonlinear wave equations in three dimensions*, Comm. Partial Differential Equations **8** (1983), no. 12, 1291–1323.

E-mail address: `bony@math.u-bordeaux1.fr`

E-mail address: `hafner@math.u-bordeaux1.fr`

INSTITUT DE MATHÉMATIQUES DE BORDEAUX
UMR 5251 DU CNRS
UNIVERSITÉ DE BORDEAUX I
351 COURS DE LA LIBÉRATION
33 405 TALENCE CEDEX
FRANCE