



HAL
open science

A theorem of Poincaré-Hopf type

Stéphane Simon

► **To cite this version:**

| Stéphane Simon. A theorem of Poincaré-Hopf type. 2009. hal-00388853

HAL Id: hal-00388853

<https://hal.science/hal-00388853>

Preprint submitted on 27 May 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A theorem of Poincaré-Hopf type

Stéphane Simon

May 27, 2009

Abstract

We compute (algebraically) the Euler characteristic of a complex of sheaves with constructible cohomology. A stratified Poincaré-Hopf formula is then a consequence of the smooth Poincaré-Hopf theorem and of additivity of the Euler-Poincaré characteristic with compact supports, once we have a suitable definition of index.

AMS classification: 55N33 57R25

Keywords: intersection homology Poincaré-Hopf vector field radial

1 Introduction

M.-H. Schwartz has defined radial vector fields in [Sch65a] and extended the classical Poincaré-Hopf theorem to real analytic sets, equipped with a Whitney stratification for these vector fields [Sch86], [Sch91]. In their turn, H. King and D. Trotman have extended M.-H. Schwartz's result to more general singular spaces and generic vector fields [KT06].

Radial [Sch65a], [Sch65b] (and totally radial, see [KT06], [Sim95]) vector fields are important because of their relation with Chern-Schwartz-Mac Pherson classes. Chern-Mac Pherson classes are written as an integral combination of Mather classes of algebraic varieties with coefficients determined by local Euler obstructions [Mac74]. A transcendental definition (and the original one) of local Euler obstruction is the obstruction to extend a lift of a radial vector field, prescribed on the link of a point in the base, inside a whole neighborhood of Nash transform. Chern-Schwartz classes [Sch65a], [Sch65b] (which lie in cohomology of the complex analytic variety) are defined as the obstruction to extend a radial frame field given on a sub-skeleton of a fixed triangulation. These two points of view coincide: Chern-Mac Pherson classes are identified with Chern-Schwartz classes by Alexander duality [BS81]. In [BBF⁺95], it is shown that these Chern-MacPherson-Schwartz classes can be realised (in general not uniquely) in intersection homology with middle perversity.

This paper concerns a Poincaré-Hopf theorem in intersection homology for a stratified pseudo-manifold A ([GM83]) and a vector field v which does not necessarily admit a globally continuous flow. Our main result is that we still have a Poincaré-Hopf formula when the vector field is semi-radial [KT06] :

$$I\chi_c^{\bar{p}}(A) = \sum_{v(x)=0} Ind^{\bar{p}}(v, x).$$

More precisely, we compute (algebraically) the Euler characteristic of a complex of sheaves with constructible cohomology. A stratified Poincaré-Hopf formula is then a consequence of the smooth Poincaré-Hopf theorem and of additivity of the Euler-Poincaré characteristic with compact supports, once we have a suitable definition of index.

Given a vector field with isolated singularities on a singular space, which admits a globally continuous flow, one can already deduce a Poincaré-Hopf theorem from a Lefschetz formula in intersection homology with middle perversity [GM85], [GM93], [Mac84].

A. Dubson announced in [Dub84] a formula similar to ours for a constructible complex in a complex analytic framework. In [BDK81], J.-L. Brylinski, A. Dubson and M. Kashiwara expressed the “local characteristic” of a holonomic module as a function of multiplicities of polar varieties and local Euler obstructions.

M. Goresky and R. MacPherson have proved a Lefschetz fixed point theorem for a sub-analytic morphism and constructible complex of sheaves [GM93]. They show that a weakly hyperbolic morphism (*i.e.* whose fixed points are *weakly hyperbolic*) can be lifted to a morphism (not necessarily unique) at the level of sheaves. The Lefschetz number can be written as a sum of contributions of the various connected components of fixed points, a component being itself possibly stratified; every contribution is a sum of multiplicities (relative to the morphism), weighted by Euler characteristics in compactly supported cohomology of the strata of the connected component.

In Section 2 we give a formula to calculate the characteristic of a constructible complex of sheaves. Then, in section 3, we apply the preceding results to the intersection chain complex. A brief recall of definitions and results on stratified vector fields is given in section 4. A theorem of Poincaré-Hopf type appears in section 5, where the vector field considered is totally (or only semi-) radial. Sections 6 and 7 are devoted to illustrate the theorems of section 4.

I am very grateful to D. Trotman for numerous valuable conversations, and for his constant encouragement. I thank equally J.-P. Brasselet for the discussions we have had related to this work. I am also greatly indebted to E. Leichtnam for his active interest and for many helpful suggestions during the preparation of the paper.

2 A formula to calculate the Euler-Poincaré characteristic of a complex of sheaves with constructible cohomology

First we recall some definitions. Let R be a principal ideal domain. We shall consider sheaves of R -modules.

Definition 2.1 *A stratified set A is a topological space which is a union of a locally finite family of disjoint, connected subsets (strata) which are smooth manifolds, satisfying the frontier condition. We shall denote by \mathcal{A} the set of strata and suppose that this stratification is fixed once and for all.*

Definition 2.2 *Let A be a stratified set. We say that A is compactifiable if there exists a compact abstract stratified set (B, \mathcal{B}) ([Mat70], [Mat73], [Tho69], [Ver84]), such that $A \subseteq B$ is a locally closed subset of B which is a union of elements of \mathcal{B} . We then say that (B, \mathcal{B}) is a compactification of A .*

Definition 2.3 *Let A be a stratified set and \mathcal{F} a sheaf on A . We say that \mathcal{F} is A -constructible on A if for every stratum X of \mathcal{A} , the sheaf $\mathcal{F}|_X$ is locally constant of finite rank on R .*

Recall that $H_c(A; \mathcal{F}) \cong \mathbb{H}_c(A; \mathcal{F})$ where \mathbb{H}_c denotes hypercohomology with compact supports. As usual, suppose that $H_c^p(A; \mathcal{F})$ has finite rank for $p \geq 0$ and is null for large enough p . Then we call *Euler characteristic of A with compact supports and coefficients in \mathcal{F}* , the alternating sum of the ranks of the modules $H_c^p(A; \mathcal{F})$ and denote it by $\chi^c(A; \mathcal{F})$. When the sheaf \mathcal{F} is the constant sheaf R , we simply write $\chi^c(A)$. We shall see that the Euler characteristic is always defined in our situation.

Proposition 2.1 *Let X be a locally compact topological space, \mathcal{G} a locally constant sheaf on X of finite rank g and suppose that X admits a finite partition \mathcal{T} into open simplexes, *i.e.**

there exists a finite simplicial complex (resp. subcomplex, possibly empty) K (resp. L) and a homeomorphism $\varphi : K \setminus L \rightarrow X$. Then

$$\chi^c(X; \mathcal{G}) = \chi^c(X).g.$$

Proof. As simplexes of X are contractible, the restriction of \mathcal{G} is isomorphic to the constant sheaf over any one of them. Consider the finite union U of open simplexes of maximal dimension m . By induction on m and using the long cohomological exact sequence (with compact supports) of (U, X) , we are reduced to showing the result for U . But, applying Mayer-Vietoris to the partition of U , this shows that $\chi^c(U; \mathcal{G}|_U)$ is well defined and establishes the formula.

Proposition 2.2 *Let A be a compactifiable stratified set and (B, \mathcal{B}) a compactification of A . Let $\mathcal{A} = (X_i)_{i \in \{1, \dots, N\}}$ be the strata of A and \mathcal{F} an \mathcal{A} -constructible sheaf. Then we have :*

$$\chi^c(A; \mathcal{F}) = \sum_{i=1}^N \chi^c(X_i).rk \mathcal{F}|_{X_i}.$$

Proof. Write \overline{A} for the closure of A in B . Thanks to the triangulation theorem for abstract stratified sets of M. Goresky [Gor78], there exists a triangulation \mathcal{T} of \overline{A} adapted to the stratification \overline{A} . As \overline{A} is compact, this triangulation is finite. Moreover, it is also adapted to A .

We are going to do induction on the number of strata of A and apply the method of proof of proposition 2.1. Let X be a stratum of maximal depth ([Ver84]) in A . Remark that X is closed in A . If $A = X$ we apply proposition 2.1 with X and \mathcal{F} .

Suppose the cardinal of \mathcal{A} is strictly greater than 1.

We have then a long exact sequence in cohomology :

$$\cdots \rightarrow H_c^p(A \setminus X; \mathcal{F}|_{A \setminus X}) \rightarrow H_c^p(A; \mathcal{F}) \rightarrow H_c^p(X; \mathcal{F}|_X) \rightarrow \cdots$$

As the number of strata of $A \setminus X$ is strictly smaller than that in A , we can apply the induction hypothesis to $A \setminus X$ and $\mathcal{F}|_{A \setminus X}$. This shows that $rk H_c^p(A; \mathcal{F})$ is finite, so $\chi^c(A; \mathcal{F})$ is defined. On the other hand, we have :

$$\chi^c(A; \mathcal{F}) = \chi^c(A \setminus X; \mathcal{F}|_{A \setminus X}) + \chi^c(X; \mathcal{F}|_X).$$

We conclude by using the induction hypothesis and proposition 2.1.

Let \mathcal{F}^\bullet be a complex of sheaves. Let $\mathcal{H}^\bullet(\mathcal{F}^\bullet)$ be the complex of derived sheaves.

Definition 2.4 *Let A be a compactifiable stratified set and \mathcal{F}^\bullet a complex of sheaves on A . We say that \mathcal{F}^\bullet has \mathcal{A} -constructible cohomology if :*

(i) \mathcal{F}^\bullet is bounded

(ii) $\mathcal{H}^\bullet(\mathcal{F}^\bullet)$ is \mathcal{A} -constructible.

Theorem 2.1 *Let A be a compactifiable stratified set, $\mathcal{A} = (X_i)_{i \in \{1, \dots, N\}}$ its stratification and \mathcal{F}^\bullet a complex of c -acyclic sheaves with \mathcal{A} -constructible cohomology. Then we have :*

$$\chi^c(A; \mathcal{F}^\bullet) = \sum_{q=-N_1}^{N_2} (-1)^q rk \mathbb{H}_c^q(A; \mathcal{F}^\bullet) = \sum_{i=1}^N \chi^c(X_i) \chi((\mathcal{H}^\bullet(\mathcal{F}^\bullet))|_{X_i})_{x_i}$$

where $\mathcal{F}^p = 0$ except for $-N_1 \leq p \leq N_2$ and x_i is any point of X_i , $1 \leq i \leq N$.

Proof. As \mathcal{F}^p is c -acyclic for all $p \in \mathbb{Z}$, we have $H^p(H_c^q(A; \mathcal{F}^\bullet)) = 0$ for all $p \in \mathbb{Z}$ and $q \geq 1$. So the second spectral sequence, of second term ' $E_2^{p,q} = H^p(H_c^q(A; \mathcal{F}^\bullet))$ ', degenerates. As \mathcal{F}^\bullet is bounded, the filtration of the associated double complex is regular, so the first spectral sequence is convergent and we have according to theorem 4.6.1 of [God73] p. 178 :

$$E_2^{p,q} = H_c^p(A; \mathcal{H}^q(\mathcal{F}^\bullet)) \Rightarrow \mathbb{H}_c^{p+q}(A; \mathcal{F}^\bullet).$$

As \mathcal{F}^\bullet has \mathcal{A} -constructible cohomology and A is compactifiable, we can define :

$$\begin{aligned} \chi(E_2) &= \sum_{p \in \mathbb{N}, q \in \mathbb{Z}} (-1)^{p+q} rg H_c^p(A; \mathcal{H}^q(\mathcal{F}^\bullet)) \\ &= \sum_{q \in \mathbb{Z}} (-1)^q \sum_{p \in \mathbb{N}} (-1)^p rg H_c^p(A; \mathcal{H}^q(\mathcal{F}^\bullet)) \\ &= \sum_{q=-N_1}^{N_2} (-1)^q \chi^c(A; \mathcal{H}^q(\mathcal{F}^\bullet)) \end{aligned}$$

for \mathcal{F}^\bullet is bounded. Remark that, since A is triangulable, every point of A (which is paracompact) admits a neighborhood homeomorphic to a subspace of some \mathbb{R}^p , so that A is of cohomological dimension lower or equal to p ($< \infty$ because A is compactifiable), according to theorem 5.13.1 of [God73] p. 237.

Apply then proposition 2.2 to A and $\mathcal{H}^q(\mathcal{F}^\bullet)$:

$$\begin{aligned} \chi(E_2) &= \sum_{q=-N_1}^{N_2} (-1)^q \sum_{i=1}^N \chi^c(X_i) rg (\mathcal{H}^q(\mathcal{F}^\bullet)|_{X_i})_{x_i} \\ &= \sum_{i=1}^N \chi^c(X_i) \sum_{q=-N_1}^{N_2} (-1)^q rg (\mathcal{H}^q(\mathcal{F}^\bullet)|_{X_i})_{x_i} \\ &= \sum_{i=1}^N \chi^c(X_i) \sum_{q=-N_1}^{N_2} (-1)^q rg \mathcal{H}^q(\mathcal{F}^\bullet)_{x_i} \\ &= \sum_{i=1}^N \chi^c(X_i) \chi(\mathcal{H}^\bullet(\mathcal{F}^\bullet)_{x_i}) \end{aligned}$$

with $x_i \in X_i$ for $i \in \{1, \dots, N\}$. As $E_{r+1} = H(E_r)$, we have $\chi(E_{r+1}) = \chi(E_r)$ for all $r \geq 2$. So $\chi(E_r) = \chi(E_2)$ for all $r \geq 2$.

As \mathcal{F}^\bullet is bounded, $E_2^{p,q} = 0$ for q big enough or small enough and $p \in \mathbb{N}$. Thus the spectral sequence degenerates and so

$$E_r^{p,q} = E_\infty^{p,q}$$

for r big enough.

Hence

$$\chi(E_\infty) = \chi(E_r) = \chi(E_2).$$

But $(E_\infty^{p,q})_{p+q=s}$ is the associated graded module to $\mathbb{H}_c^s(A; \mathcal{F}^\bullet)$. We have thus :

$$rg \mathbb{H}_c^s(A; \mathcal{F}^\bullet) = \sum_{p+q=s} rg E_\infty^{p,q}.$$

Finally

$$\begin{aligned} \chi^c(A; \mathcal{F}^\bullet) &= \sum_{s \in \mathbb{Z}} (-1)^s rg \mathbb{H}_c^s(A; \mathcal{F}^\bullet) \\ &= \chi(E_\infty) \\ &= \chi(E_2) \\ &= \sum_{i=1}^N \chi^c(X_i) \chi(\mathcal{H}^\bullet(\mathcal{F}^\bullet)_{x_i}). \end{aligned}$$

Remark. Theorem 2.1 works also with the weaker hypothesis of (finite) triangulability.

3 Application to intersection homology

Suppose now that A is a pseudo-manifold, and let \mathcal{A} be its stratification. Here the strata of A will no longer be necessarily connected, but we shall work with connected components of strata. We denote by L_x the link of the point x in A .

Proposition 3.1 ([Ba84]) *Let A be an n -pseudo-manifold and \bar{p} a perversity. Let $IC_{\bullet}^{\bar{p}}$ be the intersection chain complex for perversity \bar{p} with coefficients in R [GM93] and set $\mathcal{IC}_{\bar{p}}^{\bullet}$ =sheaf associated to the presheaf $\{U \mapsto IC_{n-\bullet}^{\bar{p}}(U)\}$. Then $\mathcal{IC}_{\bar{p}}^{\bullet}$ is a complex of c -soft sheaves (so c -acyclic). Moreover we have :*

$$\mathbb{H}_c^{\bullet}(A; \mathcal{IC}_{\bar{p}}^{\bullet}) = IH_{n-\bullet}^{\bar{p}}(A; R).$$

Proposition 3.2 (Proposition 2.4 of [GM83]) *Let A be an n -pseudo-manifold, x any point in a stratum X^k of A of dimension k and L_x the link of X^k at x in A . The fibre of the complex of derived sheaves $\mathcal{H}^{\bullet}(\mathcal{IC}_{\bar{p}}^{\bullet})$ is given by :*

$$\mathcal{H}^i(\mathcal{IC}_{\bar{p}}^{\bullet})_x = \begin{cases} \begin{cases} IH_{n-i-k-1}^{\bar{p}}(L_x) & \text{if } i \leq p_{n-k} \\ 0 & \text{otherwise} \end{cases} & \text{if } x \in X^k \subset A \setminus A_{reg} \\ \begin{cases} R & \text{if } i = 0 \\ 0 & \text{otherwise} \end{cases} & \text{if } x \in X^n \subset A_{reg}. \end{cases}$$

As usual the *Euler-Poincaré characteristic in intersection homology* $I\chi_c^{\bar{p}}(A)$ of an n -pseudo-manifold A is the Euler-Poincaré characteristic with compact supports of the complex of sheaves $\mathcal{IC}_{\bar{p}}^{\bullet}$ multiplied by $(-1)^n$, i.e. $I\chi_c^{\bar{p}}(A) = (-1)^n \chi^c(A; \mathcal{IC}_{\bar{p}}^{\bullet})$.

Theorem 3.1 *Let A be an n -pseudo-manifold such that (A, \mathcal{A}) is compactifiable, N the number of connected components of strata of A and \bar{p} a perversity.*

We have :

$$I\chi_c^{\bar{p}}(A) = \sum_{i=1}^N (-1)^n \chi^c(X_i) \sum_{j=0}^{p_{n-\dim X_i}} (-1)^j rg IH_{n-j-\dim X_i-1}^{\bar{p}}(L_{x_i}; R)$$

where x_i is an arbitrary point of X_i for $1 \leq i \leq N$ and we make the convention that $rg IH_{-1}^{\bar{p}}(L_{x_i}; R) = 1$ if $\dim X_i = n$.

Proof. Application of theorem 2.1 and proposition 3.2.

Remark. As in theorem 2.1 we can weaken the hypothesis by only assuming the existence of a (finite) triangulation compatible with the stratification.

Proposition 3.3 *Let A be a $2n$ -pseudo-manifold such that (A, \mathcal{A}) is compactifiable, the dimension of strata being even and let \bar{m} be the middle perversity. We have :*

$$I\chi_c^{\bar{m}}(A) = \sum_{i=0}^n \sum_{k=1}^{N_i} \chi_c(X_k^{2i}) \sum_{j=0}^{n-i-1} (-1)^j rg IH_{2n-j-2i-1}^{\bar{m}}(L_{x_k^i}; R)$$

where we have written X_k^{2i} (resp. N_i) for the k -th connected component of the stratum (resp. number of connected components of the stratum) of dimension $2i$, x_k^i an arbitrary point of X_k^{2i}

and $\chi_c(X) = \sum_{i=0}^{\dim X} (-1)^i rg H_i^c(X; R)$.

Proof. We apply theorem 2.1 with $\bar{p} = \bar{m}$ and we remark that $\chi_c(X) = \chi^c(X)$ for a manifold X of even dimension.

4 Totally radial and semi-radial vector fields on abstract stratified sets

M.-H. Schwartz constructed certain frame fields to define (by obstruction) her Chern-Schwartz classes in the cohomology of a singular complex analytic variety equipped with a Whitney stratification [Sch65a], [Sch65b]. These were called radial fields. When one is concerned with 1-frame fields (i.e. vector fields), they are called radial vector fields. She showed that they verified a Poincaré-Hopf formula [Sch86], [Sch91].

This section is an easy transcription to abstract stratified sets of some notions and results of [KT06] which were given in the more general setting of “mapping cylinder stratified space with boundary”. In their paper, H. King and D. Trotman extend M.-H. Schwartz’s work on Poincaré-Hopf formulas, to more general spaces, and to generic vector fields. Notice that abstract stratified sets are not (necessarily) embedded nor are vector fields (necessarily) continuous.

Definition 4.1 ([KT06]) *Let (A, \mathcal{A}) be an abstract stratified set and v a stratified vector field on A ([Mat70],[Mat73], [Tho69], [Ver84]). We say that v is a totally radial vector field if for all strata $X \in \mathcal{A}$ there exists a neighborhood U_X of X in the control tube T_X such that $d\rho_X(v) > 0$ on $U_X \setminus X$ (i.e. v is pointing outwards with respect to the level hypersurfaces of the control function ρ_X).*

In [KT06] such a vector field was called *radial*. To avoid confusion with the radial vector fields of M.-H. Schwartz, we have adopted the terminology *totally radial*, which also expresses the fact that one imposes that $d\rho_X(v) > 0$ on a whole neighborhood U_X of X in T_X . The analogous condition is only imposed on a neighbourhood of some closed subset of X by M.H. Schwartz. See [Sim95] for a detailed discussion of the differences between the radial fields of [Sch86], [Sch91] and the radial fields of [KT06], called totally radial here.

Proposition 4.1 *Let (A, \mathcal{A}) be an abstract stratified set and Y a stratum of A . Then there exists a vector field ξ_Y on $T_Y \setminus Y$ such that :*

$$\text{for all } y \text{ in } T_Y \setminus Y \text{ we have } \begin{cases} \rho_{Y_*y}(\xi_Y) = 1 \\ \rho_{X_*y}(\xi_Y) = 0 \quad \text{if } X < Y. \end{cases}$$

Proof. It suffices to consider the stratified submersion $(\pi_Y, \rho_Y) : T_Y \setminus Y \rightarrow Y \times \mathbb{R}_+^*$ and to lift the constant field $(0, \partial_t)$ to a field ξ_Y on $T_Y \setminus Y$. Thanks to the compatibility conditions, we see that $\rho_{X_*}(\xi_Y) = 0$ for $X < Y$.

Definition 4.2 ([KT06]) *Let (A, \mathcal{A}) be an abstract stratified set, v a stratified vector field on A and Y a stratum of A . Let $(Y_i)_{1 \leq i \leq m}$ be the strata such that $Y < Y_i$. Set $B_Y(v) = \{x \in T_Y \setminus Y \mid (\exists c_i \in \mathbb{R}_- \mid 0 \leq i \leq m) : v(y) = c_0 \xi_Y(y) + \sum_{j=1}^m c_j \xi_{Y_j}(y) \text{ with } c_0 < 0\}$. A point $x \in \overline{B_Y(v)} \cap Y$ is called a virtual zero of v .*

Definition 4.3 ([KT06]) *Let (A, \mathcal{A}) be an abstract stratified set and v a stratified vector field on A . Then v is called semi-radial if v has no virtual zero.*

Examples. Totally radial vector fields, and controlled vector fields, are semi-radial.

Definition 4.4 *Let (A, \mathcal{A}) be a compactifiable stratified set, (B, \mathcal{B}) a compactification of A such that $\mathcal{A} \subseteq \mathcal{B}$ and v a stratified vector field on A . We say that v is strongly totally radial (resp. strongly semi-radial) if and only if there exists a totally radial (resp. semi-radial) extension u of v to (B, \mathcal{B}) .*

Lemma 4.1 ([KT06]) *Let (A, \mathcal{A}) be an abstract stratified set (resp. compactifiable stratified set) and v a semi-radial (resp. strongly semi-radial) vector field with isolated singularities on A . Then there exists a (resp. strongly) totally radial vector field v' having the same singularities as v and the same indices at these points.*

5 Towards a Poincaré-Hopf theorem

Definition 5.1 *Let A be an n -pseudo-manifold, \bar{p} a perversity and x a point of a stratum X . We call multiplicity of A at x for perversity \bar{p} the following integer :*

$$m_x^{\bar{p}}(A) = \begin{cases} \sum_{i=n-p_n-\dim X}^n (-1)^i \text{rg } IH_{i-\dim X-1}^{\bar{p}}(L_x; R) & \text{if } x \in A \setminus A_{\text{reg}} \\ (-1)^n & \text{if } x \in A_{\text{reg}}. \end{cases}$$

Remark. The multiplicity is nothing else than $I\chi_c^{\bar{p}}(A, A - \{x\})$ (which equals $(-1)^n$ if $x \in A_{\text{reg}}$).

Definition 5.2 *Let A be an n -pseudo-manifold such that (A, \mathcal{A}) is a compactifiable abstract stratified set, \bar{p} a perversity and v a stratified vector field having an isolated singularity at $x \in X$. We call singular index of v at x , and we denote by $\text{Ind}^{\bar{p}}(v, x)$ the integer:*

$$\text{Ind}^{\bar{p}}(v, x) = m_x^{\bar{p}}(A) \cdot \text{Ind}(v, x).$$

Recall that if the stratum X is reduced to a point, then $\text{Ind}(v, x) = 1$.

Theorem 5.1 *Let A be an n -pseudo-manifold such that (A, \mathcal{A}) is a compactifiable abstract stratified set, \bar{p} a perversity and v a strongly semi-radial vector field admitting a finite number of singularities on A . We have :*

$$I\chi_c^{\bar{p}}(A) = \sum_{v(x)=0} \text{Ind}^{\bar{p}}(v, x).$$

Proof. As in [Bek92], for all strata X of A , let f_X be a carpeting function, i.e. let $U_{b(X)}$ be a neighborhood of $b(X) = \bar{X} \setminus X$ in \bar{X} , and let $f_X : U_{b(X)} \rightarrow \mathbb{R}_+$ be a continuous function (constructed using the control functions $\{\rho_X\}_{X \subseteq A}$ induced by the compactification of A), smooth on the stratum X such that $f_X^{-1}(0) = b(X)$ and $f_X|_{U_{b(X)} \cap X}$ is submersive. Now, apply lemma 4.1 to v ; this gives a totally radial vector field v' . Then we remark that if v' is a totally radial vector field, for all strata X , v' is entering $X_{\geq \epsilon} = X \setminus \{f_X < \epsilon\}$ along ∂X_ϵ for ϵ small enough, where the symbol ∂X_ϵ denotes the level hypersurface $\{f_X = \epsilon\}$. This is because $\text{grad}(f_X) = \sum_{Y < X} a_Y \cdot \text{grad}(\rho_Y)$, where the a_Y are non-negative smooth functions, at every point of X . So we have $\chi_c(X_{\geq \epsilon}) - \chi_c(\partial X_\epsilon) = \sum_{v'(x)=0} \text{Ind}(v', x)$ thanks to the classical Poincaré-Hopf theorem. Finally, we have $\chi^c(M) = \chi_c(M) - \chi_c(\partial M)$ for every compactifiable manifold M by adding a boundary ∂M . Use the “additivity” formula of theorem 3.1 and the definition of the singular index to complete the proof.

6 A few examples

In the following computations, as we are only interested in the rank of intersection homology groups, we shall take $R = \mathbb{Q}$ and work with the dimension of \mathbb{Q} -vector spaces. Moreover, this will permit us to apply Poincaré duality to calculate some associated groups. In the remainder of the text, T^2 will denote the torus $S^1 \times S^1$. The stratifications of spaces will be the evident ones and we shall not go into details. See [Ba84] for classical tools to compute IH_\bullet of the following spaces.

6.1 An inevitable example : the pinched torus T_p^2

We have a unique perversity $\bar{p} = \bar{0}$ and we have evidently a totally radial vector field u on T_p^2 with a unique singularity at the isolated singular point x_0 of T_p^2 , of indice 1. The link at this point is $L_{x_0} = S^1 \sqcup S^1$. We have :

$$IH_i^{\bar{0}}(T_p^2) = \begin{cases} \mathbb{Q} & \text{si } i = 2 \\ 0 & \text{si } i = 1 \\ \mathbb{Q} & \text{si } i = 0 \end{cases}$$

so that

$$I\chi^{\bar{0}}(T_p^2) = 2.$$

On the other hand :

$$\begin{aligned} m_{x_0}^{\bar{0}}(T_p^2) &= \dim IH_1^{\bar{0}}(L_{x_0}) \\ &= \dim H_1(S^1 \sqcup S^1) \\ &= 2. \end{aligned}$$

Finally we have $I\chi^{\bar{0}}(T_p^2) = 2 = 2.1 = Ind^{\bar{0}}(u, x_0)$.

6.2 A well-known example : the suspension of the torus ΣT^2 (H. Poincaré, 1895)

This time, we have two different perversities $\bar{0}$ and \bar{t} and two isolated singularities (which are the two vertices of suspension). The link at these points is $L_{x_0} = L_{x_1} = T^2$. We still have a totally radial vector field v with two singular points of indice 1 at singularities of ΣT^2 . Remark that this pseudo-manifold is normal so we have $IH_*^{\bar{t}}(\Sigma T^2) = H_*(\Sigma T^2)$, i.e.

$$IH_i^{\bar{t}}(\Sigma T^2) = \begin{cases} \mathbb{Q} & \text{if } i = 3 \\ \mathbb{Q}^2 & \text{if } i = 2 \\ 0 & \text{if } i = 1 \\ \mathbb{Q} & \text{if } i = 0. \end{cases}$$

Hence

$$I\chi^{\bar{t}}(\Sigma T^2) = 2$$

and by duality we find

$$I\chi^{\bar{0}}(\Sigma T^2) = -2.$$

On the other hand :

$$\begin{aligned} m_{x_0}^{\bar{0}}(\Sigma T^2) &= -\dim IH_2^{\bar{0}}(L_{x_0}) \\ &= -\dim H_2(T^2) \\ &= -1 \end{aligned}$$

and

$$\begin{aligned} m_{x_0}^{\bar{t}}(\Sigma T^2) &= \dim IH_1^{\bar{t}}(L_{x_0}) - \dim IH_2^{\bar{t}}(L_{x_0}) \\ &= \dim H_1(T^2) - \dim H_2(T^2) \\ &= 1. \end{aligned}$$

Finally we have :

$$I\chi^{\bar{0}}(\Sigma T^2) = -2 = -1 - 1 = 2.Ind^{\bar{0}}(v, x_0)$$

and

$$I\chi^{\bar{t}}(\Sigma T^2) = 2 = 1 + 1 = 2.Ind^{\bar{t}}(v, x_0).$$

6.3 A hybrid example : the suspension of the torus of dimension 3, twice pinched, ΣT_{2p}^3

We have $\Sigma T_{2p}^3 = \Sigma(\Sigma(T^2 \sqcup T^2))$. Here we have four perversities $\bar{0}, \bar{m}, \bar{n}, \bar{t}$. Calculate to begin with the homology of T_{2p}^3 :

$$H_i(T_{2p}^3) = \begin{cases} \mathbb{Q}^2 & \text{si } i = 3 \\ \mathbb{Q}^4 & \text{if } i = 2 \\ \mathbb{Q} & \text{if } i = 1 \\ \mathbb{Q} & \text{if } i = 0 \end{cases}.$$

Then its intersection homology is :

$$\begin{aligned} IH_i^{\bar{0}}(T_{2p}^3) &= \begin{cases} H_i(T_{2p}^3) & \text{if } i > 2 \\ Im(H_i(T^2 \sqcup T^2) \rightarrow H_i(T_{2p}^3)) & \text{if } i = 2 \\ H_i(T^2 \sqcup T^2) & \text{if } i < 2 \end{cases} \\ &= \begin{cases} \mathbb{Q}^2 & \text{if } i = 3 \\ 0 & \text{if } i = 2 \\ \mathbb{Q}^4 & \text{if } i = 1 \\ \mathbb{Q}^2 & \text{if } i = 0 \end{cases} \end{aligned}$$

where we deduce

$$IH_i^{\bar{t}}(T_{2p}^3) = \begin{cases} \mathbb{Q}^2 & \text{if } i = 3 \\ \mathbb{Q}^4 & \text{if } i = 2 \\ 0 & \text{if } i = 1 \\ \mathbb{Q}^2 & \text{if } i = 0 \end{cases}.$$

And at last the intersection homology of the suspension ΣT_{2p}^3 is :

$$\begin{aligned} IH_i^{\bar{p}}(\Sigma T_{2p}^3) &= \begin{cases} IH_{i-1}^{\bar{p}}(T_{2p}^3) & \text{if } i > 3 - p_4 \\ 0 & \text{if } i = 3 - p_4 \\ IH_i^{\bar{p}}(T_{2p}^3) & \text{if } i < 3 - p_4 \end{cases} \\ &= \begin{cases} \begin{cases} \mathbb{Q}^2 & \text{if } i = 4 \\ 0 & \text{if } i = 3 \\ 0 & \text{if } i = 2 \end{cases} & \text{if } \bar{p} = \bar{m} \\ \begin{cases} \mathbb{Q}^4 & \text{if } i = 1 \\ \mathbb{Q}^2 & \text{if } i = 0 \end{cases} \\ \begin{cases} \mathbb{Q}^2 & \text{if } i = 4 \\ 0 & \text{if } i = 3 \\ 0 & \text{if } i = 2 \end{cases} & \text{if } \bar{p} = \bar{0}. \\ \begin{cases} \mathbb{Q}^4 & \text{if } i = 1 \\ \mathbb{Q}^2 & \text{if } i = 0 \end{cases} \end{cases} \end{aligned}$$

It is easy to construct a totally radial vector field w with four singularities : two at the vertices of suspension, say x_0, x_1 , of indice 1 and two others on strata of codimension 3, say x_2, x_3 , of indice -1 . Links are $L_{x_0} = L_{x_1} = T_{2p}^3$ and $L_{x_2} = L_{x_3} = T^2 \sqcup T^2$. Calculations of multiplicities give :

$$m_{x_0}^{\bar{p}}(\Sigma T_{2p}^3) = \begin{cases} \dim IH_1^{\bar{t}}(T_{2p}^3) - \dim IH_2^{\bar{t}}(T_{2p}^3) + \dim IH_3^{\bar{t}}(T_{2p}^3) = -2 & \text{if } \bar{p} = \bar{t} \\ -\dim IH_2^{\bar{n}}(T_{2p}^3) + \dim IH_3^{\bar{n}}(T_{2p}^3) = -2 & \text{if } \bar{p} = \bar{n} \\ -\dim IH_2^{\bar{m}}(T_{2p}^3) + \dim IH_3^{\bar{m}}(T_{2p}^3) = 2 & \text{if } \bar{p} = \bar{m} \\ \dim IH_3^{\bar{0}}(T_{2p}^3) = 2 & \text{if } \bar{p} = \bar{0} \end{cases}$$

and

$$m_{x_2}^{\bar{p}}(\Sigma T_{2p}^3) = \begin{cases} -\dim H_1(T^2 \sqcup T^2) + \dim H_2(T^2 \sqcup T^2) = -2 & \text{if } \bar{p} = \bar{t} \\ -\dim H_1(T^2 \sqcup T^2) + \dim H_2(T^2 \sqcup T^2) = -2 & \text{if } \bar{p} = \bar{n} \\ \dim H_2(T^2 \sqcup T^2) = 2 & \text{if } \bar{p} = \bar{m} \\ \dim H_2(T^2 \sqcup T^2) = 2 & \text{if } \bar{p} = \bar{0} \end{cases}.$$

Finally we have

$$\begin{aligned} I\chi^{\bar{0}}(\Sigma T_{2p}^3) &= 0 = 2 + 2 + 2 \cdot (-1) + 2 \cdot (-1) \\ I\chi^{\bar{m}}(\Sigma T_{2p}^3) &= 0 = 2 + 2 - 2 - 2 \\ I\chi^{\bar{n}}(\Sigma T_{2p}^3) &= 0 = -2 - 2 + (-2) \cdot (-1) + (-2) \cdot (-1) \\ I\chi^{\bar{t}}(\Sigma T_{2p}^3) &= 0 = -2 - 2 + 2 + 2. \end{aligned}$$

7 A partial converse

We present here a partial converse to theorem 5.1 in the sense that we study when a stratified set admits a strongly totally radial vector field without singularity. This result is in the line of [Sul71], [Ver72] or [Sch91], [Sch92]. See also [Mat73], theorem 8.5. The result is partial because of the example below. Indeed, it shows that we cannot expect the condition $I\chi_c^{\bar{p}}(A) = 0$ to imply the existence of a non singular totally radial vector field.

Theorem 7.1 *Let A be a compactifiable n -pseudo-manifold. There exists a strongly totally radial vector field (relatively to A) on A without singularity if and only if $\chi^c(X) = 0$ for all strata X of A .*

Proof. To show sufficiency, we use the carpeting functions of the proof of theorem 5.1. Let v be a strongly totally radial vector field on A with isolated singularities ; the vector field v_X is entering on the boundary $\partial X_{\geq \epsilon}$ (defined by a level hypersurface of a carpeting function). Remark that, as $\chi^c(X) = 0$, we can deform v_X on $X_{\geq \epsilon}$ (without modifying it near $\partial X_{\geq \epsilon}$) so as to have no singularities ([Hir88]). We have evidently $\rho_{X_*}(v) > 0$ on $T_X \setminus X$ for all strata X . Necessity is proved in an analogous manner.

Corollary 7.1 *Let A be a compactifiable n -pseudo-manifold, stratified with strata of odd dimension. Then there exists a strongly totally radial vector field without singularity on A .*

Remark. Existence of a totally radial vector field without singularity, on an abstract stratified set, is equivalent to the existence of a controlled vector field without singularity.

Example. Finally, here is an example of a compact pseudo-manifold without strata of dimension 0 for which $I\chi_c^{\bar{p}}(A) = 0$ for every perversity \bar{p} and admitting no totally radial vector field without a singularity. Consider $A = \Sigma(T_{2p}^3) \times S^2$; it is clear that $I\chi^{\bar{p}}(A; R) = 0$ for all \bar{p} . Nevertheless, there does not exist a totally radial vector field without a singularity (look at strata $\{*\} \times S^2$ or $\{**\} \times S^2$). This is also evident as a consequence of theorem 7.1.

References

- [Ba84] A. Borel and al. *Intersection cohomology*, volume 50 of *Progress in mathematics*. Birkhäuser Boston • Basel • Stuttgart, 1984.
- [BBF⁺95] G. Barthel, J.-P. Brasselet, K.-H. Fieseler, O. Gabber, and L. Kaup. Relèvement de cycles algébriques et homomorphismes associés en homologie d'intersection. *Annals of Mathematics*, 141:147–179, 1995.
- [BDK81] J.-L. Brylinski, A. Dubson, and M. Kashiwara. Formule d'indice pour les modules holonomes et obstruction d'Euler locale. *Comptes Rendus de l'Académie des Sciences*, 293:573–576, 1981.
- [Bek92] K. Bekka. Continuous vector fields and Thom's isotopy theorems, 1992.

- [BS81] J.-P. Brasselet and M.-H. Schwartz. Sur les classes de Chern d'un ensemble analytique complexe. In S.M.F., editor, *Séminaire E.N.S. 1978-1979 : Caractéristique d'Euler-Poincaré*, volume 82-83 of *Astérisque*, pages 93–147, 1981.
- [Dub84] A. Dubson. Formule pour l'indice des complexes constructibles et des modules holonomes. *Comptes Rendus de l'Académie des Sciences*, 298(6):113–116, 1984.
- [GM83] M. Goresky and R. MacPherson. Intersection homology theory II. *Inventiones Mathematicae*, 72:77–129, 1983.
- [GM85] M. Goresky and R. MacPherson. Lefschetz fixed point theorem for intersection homology. *Commentarii Mathematici Helvetici*, 60(3):386–391, 1985.
- [GM93] M. Goresky and R. MacPherson. Local contribution to the Lefschetz fixed point formula. *Inventiones Mathematicae*, 111:1–33, 1993.
- [God73] R. Godement. *Topologie algébrique et théorie des faisceaux*. Actualités scientifiques et industrielles de l'Université de Strasbourg. Hermann Paris, 1973.
- [Gor78] M. Goresky. Triangulation of stratified objects. *Proceedings of the American Mathematical Society*, 72:193–200, 1978.
- [Hir88] M. Hirsch. *Differential topology*, volume 33. Springer-Verlag New York, 1988.
- [KT06] H. King and D. Trotman. Poincaré-Hopf theorems on stratified spaces, 2006. Université de Provence.
- [Mac74] R. MacPherson. Chern classes for singular algebraic varieties. *Annals of Mathematics*, 100:423–432, 1974.
- [Mac84] R. MacPherson. Global questions in the topology of singular spaces. In *Proceedings of the International Congress of Mathematicians*, pages 213–235, Warszawa, 1984. August 16-24 1983.
- [Mat70] J. Mather. Notes on topological stability. Harvard University, 1970.
- [Mat73] J. Mather. Stratifications and mappings. *Dynamical Systems*, pages 195–232, 1973.
- [Sch65a] M.-H. Schwartz. Classes caractéristiques définies par une stratification d'une variété complexe. *Comptes Rendus de l'Académie des Sciences*, 260:3262–3264 and 3535–3537, 1965.
- [Sch65b] M.-H. Schwartz. Classes obstructrices d'un sous-ensemble analytique complexe d'une variété analytique complexe. Publication de l'Université de Lille, 1965.
- [Sch86] M.-H. Schwartz. Une généralisation du théorème de Hopf sur les champs sortants. *Comptes Rendus de l'Académie des Sciences*, 303(7):307–309, 1986.
- [Sch91] M.-H. Schwartz. *Champs radiaux sur une stratification analytique*. Travaux en cours. Hermann, 1991.
- [Sch92] M.-H. Schwartz. Un problème d'homologie dans l'espace des vecteurs tangents à une stratification. *Comptes Rendus de l'Académie des Sciences*, 314:833–836, 1992.
- [Sim95] S. Simon. Champs de vecteurs totalement radiaux sur une structure de Thom-Mather. *Annales de l'Institut Fourier*, 45(5):1423–1447, 1995.
- [Sul71] D. Sullivan. Combinatorial invariants of analytic spaces. In *Liverpool Singularities Symposium*, volume 192 of *Lecture notes in mathematics*, pages 165–168. Springer-Verlag, 1971.
- [Tho69] R. Thom. Ensembles et morphismes stratifiés. *Bulletin of the American Mathematical Society*, 75:240–284, 1969.
- [Ver72] A. Verona. Homological properties of abstract prestratifications. *Revue Roumaine de Mathématiques pures et appliquées*, XVII(7):1109–1121, 1972.
- [Ver84] A. Verona. *Stratified mappings—Structure and triangulability*, volume 1102 of *Lecture notes in mathematics*. Springer-Verlag Berlin, 1984.