



HAL
open science

New permutation coding and equidistribution of set-valued statistics

Dominique Foata, Guo-Niu Han

► **To cite this version:**

Dominique Foata, Guo-Niu Han. New permutation coding and equidistribution of set-valued statistics. Theoretical Computer Science, 2009, 13 p. hal-00395705

HAL Id: hal-00395705

<https://hal.science/hal-00395705>

Submitted on 16 Jun 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

April 9, 2009

NEW PERMUTATION CODING AND EQUIDISTRIBUTION OF SET-VALUED STATISTICS

Dominique Foata and Guo-Niu Han

ABSTRACT. A new coding for permutations is explicitly constructed and its association with the classical Lehmer coding provides a bijection of the symmetric group onto itself serving to show that six bivariable set-valued statistics are equidistributed on that group. This extends a recent result due to Cori valid for integer-valued statistics.

1. Introduction

In a recent paper Cori [Cor08] updates a classical algorithm constructed by Ossona de Mendez and Rosenstiehl [OR04] that provides a one-to-one correspondence between rooted hypermaps and indecomposable permutations. He further constructs a bijection of the symmetric group \mathfrak{S}_n onto itself that maps each permutation having p cycles and q left-to-right maxima onto another one having q cycles and p left-to-right maxima. Moreover, by using an encoding of permutations by Dyck paths due to Roblet and Viennot [RV96] he also shows that three bivariable *integer-valued* statistics, introduced in the next paragraph, are equidistributed on \mathfrak{S}_n . The purpose of this paper is to show that all those results can be extended to *set-valued* statistics and that the construction of the underlying bijection is very simple; it involves two permutation codings called the *A-code* and the *B-code*.

The first one, classically referred to as the *Lehmer code* [Le60] or the *inversion table*, goes back, in fact, to more ancient authors (Rothe, Rodrigues, Netto), as knowledgeably stated by Knuth ([Kn98], Ex. 5.1.1-7, p. 14). The second one is a *new* coding that takes the cycle decomposition of permutations into account. Although the motivation of the paper was to prove the equidistribution of several set-valued statistics, its novelty is to fully describe that B-code and exploit its basic properties.

The set-valued statistics in question are introduced as follows. Let $w = x_1x_2 \cdots x_n$ be a word of length n , whose letters are positive integers. The **L**eft to right **m**aximum **p**lace set, $\text{Lmap } w$, of w is defined to be the set of all *places* i such that $x_j < x_i$ for all $j < i$, while the **R**ight to left **m**inimum **l**etter set, $\text{Rmil } w$, of w is the set of all *letters* x_i such that $x_j > x_i$ for all $j > i$.

When the word w is a permutation of $12 \cdots n$ that we shall preferably denote by $\sigma = \sigma(1)\sigma(2) \cdots \sigma(n)$ and the bijection $i \mapsto \sigma(i)$ ($1 \leq i \leq n$) has

r disjoint cycles, whose *minimum* elements are c_1, c_2, \dots, c_r , respectively, define $\text{Cyc } \sigma$ to be the set

$$\text{Cyc } \sigma := \{c_1, c_2, \dots, c_r\}.$$

When σ is a permutation, the *cardinalities* of $\text{Lmap } \sigma$, $\text{Rmil } \sigma$ and $\text{Cyc } \sigma$ are denoted by $\text{lmap } \sigma$, $\text{rmil } \sigma$ and $\text{cyc } \sigma$, respectively, and classically referred to as the *number of left-to-right maxima*, *number of right-to-left minima*, *number of cycles*.

In Fig. 1 the graphs of the permutation $\sigma = 5, 7, 1, 4, 9, 2, 6, 3, 8$ and its inverse $\sigma^{-1} = 3, 6, 8, 4, 1, 7, 2, 9, 5$ have been drawn. The set $\text{Lmap } \sigma$ (resp. $\text{Lmap } \sigma^{-1}$) is the set of the *abscissas* of the “bullets,” while $\text{Rmil } \sigma$ (resp. $\text{Rmil } \sigma^{-1}$) is the set of the *ordinates* of the “crosses.” The set-valued statistics “Leh,” “RmilLeh” and “MaxLeh” will be further introduced. Notice that $\text{lmap } \sigma = \text{rmil } \sigma^{-1} = 3$, $\text{rmil } \sigma = \text{lmap } \sigma^{-1} = 4$. As σ is the product of the disjoint cycles $(15983)(4)(276)$, we have $\text{Cyc } \sigma = \text{Cyc } \sigma^{-1} = \{1, 2, 4\}$ and $\text{cyc } \sigma = \text{cyc } \sigma^{-1} = 3$.




Fig. 1. Graphs of σ and of its inverse σ^{-1}

First, recall Cori’s result [Cor08].

The three pairs of integer-valued statistics $(\text{rmil}, \text{cyc})$, $(\text{cyc}, \text{rmil})$ and $(\text{lmap}, \text{rmil})$ are equidistributed on \mathfrak{S}_n .

The equidistribution of the first two pairs (resp. of the last two ones) is proved by updating the Ossona-de-Mendez-Rosenstiehl algorithm [OR04]

on hypermaps (resp. by using the Roblet-Viennot Dyck path encoding [RV96]). Second, the set-valued statistics “Cyc” and “Rmil” (or “Lmap”) are known to be equidistributed on \mathfrak{S}_n . This is one of the properties of the first fundamental transformation [Lo83, chap. 10]. Our main result is the following theorem.

Theorem 1. *The six bivariable set-valued statistics (Cyc, Rmil), (Cyc, Lmap), (Rmil, Lmap), (Rmil, Cyc), (Lmap, Rmil), (Lmap, Cyc) are all equidistributed on \mathfrak{S}_n .*

Based on two permutation codings, the A-code and B-code, introduced in Sections 2 and 3, respectively, we construct a bijection ϕ of \mathfrak{S}_n onto itself (see (4.1)) having the following property:

$$(1.1) \quad (\text{Lmap, Rmil}) \sigma = (\text{Lmap, Cyc}) \phi(\sigma) \quad (\sigma \in \mathfrak{S}_n).$$

Let $\mathbf{i} : \sigma \mapsto \sigma^{-1}$. As

$$(1.2) \quad \text{Cyc } \mathbf{i} \sigma = \text{Cyc } \sigma;$$

$$(1.3) \quad \text{Rmil } \mathbf{i} \sigma = \text{Lmap } \sigma;$$

(see Fig. 1 for the second relation), it follows from (1.1) that the chain

$$(1.4) \quad \begin{array}{ccccccccc} \mathfrak{S}_n & \xrightarrow{\mathbf{i}} & \mathfrak{S}_n & \xrightarrow{\phi^{-1}} & \mathfrak{S}_n & \xrightarrow{\mathbf{i}} & \mathfrak{S}_n & \xrightarrow{\phi} & \mathfrak{S}_n & \xrightarrow{\mathbf{i}} & \mathfrak{S}_n \\ (\text{Cyc}) & & (\text{Cyc}) & & (\text{Rmil}) & & (\text{Lmap}) & & (\text{Lmap}) & & (\text{Rmil}) \\ (\text{Rmil}) & & (\text{Lmap}) & & (\text{Lmap}) & & (\text{Rmil}) & & (\text{Cyc}) & & (\text{Cyc}) \end{array}$$

provides all the bijections needed to prove Theorem 1. Note that (1.1), on the one hand, and (1.2)–(1.3), on the other hand, are reproduced as

$$\begin{array}{ccc} \mathfrak{S}_n & \xrightarrow{\phi} & \mathfrak{S}_n & \text{and} & \mathfrak{S}_n & \xrightarrow{\mathbf{i}} & \mathfrak{S}_n \\ (\text{Lmap}) & & (\text{Lmap}) & & (\text{Cyc}) & & (\text{Cyc}) \\ (\text{Rmil}) & & (\text{Cyc}) & & (\text{Rmil}) & & (\text{Lmap}) \end{array}$$

Let $A = (I_1, I_2, \dots, I_h)$ be an ordered partition of the set $[n] := \{1, 2, \dots, n\}$ into disjoint non-empty *intervals*, such that $\max I_j + 1 = \min I_{j+1}$ for $j = 1, 2, \dots, h - 1$. A permutation σ from \mathfrak{S}_n is said to be *A-decomposable*, if each I_j is the *smallest interval* such that $\sigma(I_j) = I_j$ (see [Com74], p. 261, exercise 14). For instance, $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 1 & 5 & 4 & 3 \end{pmatrix}$ is *A-decomposable*, with $A = (\{1, 2\}, \{3, 4, 5\})$. It is convenient to write $\text{Decomp } \sigma = A$, if σ is *A-decomposable*. A permutation is said to be *indecomposable*, if it is *A-decomposable*, with $A = ([n])$. The bijection ϕ defined in (4.1) has the further property

$$(1.5) \quad \text{Decomp } \phi(\sigma) = \text{Decomp } \sigma \quad (\sigma \in \mathfrak{S}_n).$$

As we evidently have

$$(1.6) \quad \text{Decomp } \mathbf{i} \sigma = \text{Decomp } \sigma,$$

the following result holds.

Theorem 2. *Let A be an ordered partition of the set $[n]$ into disjoint consecutive non-empty intervals. Then, $(\text{Cyc}, \text{Rmil})$, $(\text{Cyc}, \text{Lmap})$, $(\text{Rmil}, \text{Lmap})$, $(\text{Rmil}, \text{Cyc})$, $(\text{Lmap}, \text{Rmil})$, $(\text{Lmap}, \text{Cyc})$ are equidistributed on the set of all A -decomposable permutations from \mathfrak{S}_n .*

The next corollary is relevant to the study of hypermaps, as the set of rooted hypermaps with darts $1, 2, \dots, n$ is in one-to-one correspondence with the subset of indecomposable permutations from \mathfrak{S}_{n+1} (see [Cor08, CM92]).

Corollary 3. *The statistics $(\text{Cyc}, \text{Rmil})$, $(\text{Cyc}, \text{Lmap})$, $(\text{Rmil}, \text{Lmap})$, $(\text{Rmil}, \text{Cyc})$, $(\text{Lmap}, \text{Rmil})$, $(\text{Lmap}, \text{Cyc})$ are equidistributed on the set of all indecomposable permutations from \mathfrak{S}_n .*

The construction of the bijection ϕ together with the proofs of Theorem 2, and Corollary 3 are given in Section 4. It is followed by the algorithmic definitions of both A-code and B-code in Section 5. Tables and concluding remarks are reproduced in Section 6.

2. The A-code

The *Lehmer code* [Le60] of a permutation $\sigma = \sigma(1)\sigma(2)\cdots\sigma(n)$ of $12\cdots n$ is defined to be the sequence $\text{Leh } w = (a_1, a_2, \dots, a_n)$, where for each $i = 1, 2, \dots, n$

$$a_i := \#\{j : 1 \leq j \leq i, \sigma(j) \leq \sigma(i)\}.$$

The sequence $\text{Leh } w$ belongs to SE_n of all sequences $a = (a_1, a_2, \dots, a_n)$, called *subexcedant*, such that $1 \leq a_i \leq i$ for each $i = 1, 2, \dots, n$. For such a sequence it makes sense to define the set, denoted by $\text{Max } a$, of all letters (or places!) a_i such that $a_i = i$.

Under the graphs drawn in Fig. 1 the Lehmer codes $\text{Leh } \sigma$ and $\text{Leh } \sigma^{-1}$ have been calculated, as well as the four sets $\text{Rmil } \text{Leh } \sigma$, $\text{Rmil } \text{Leh } \sigma^{-1}$, $\text{Max } \text{Leh } \sigma$ and $\text{Max } \text{Leh } \sigma^{-1}$. The next Proposition is geometrically evident and given without proof. It shows that the set-valued statistics “Lmap” and “Rmip” can be directly read from the Lehmer code.

Proposition 4. *For each permutation σ we have:*

$$(2.1) \quad \text{Rmil } \text{Leh } \sigma = \text{Rmil } \sigma;$$

$$(2.2) \quad \text{Max } \text{Leh } \sigma = \text{Lmap } \sigma.$$

We then define the *A-code* of a permutation σ to be

$$(2.3) \quad \text{A-code } \sigma := \text{Leh } \mathbf{i} \sigma.$$

Hence, $\text{Max } \text{A-code } \sigma = \text{Max } \text{Leh } \mathbf{i} \sigma = \text{Lmap } \mathbf{i} \sigma = \text{Rmil } \sigma$. Furthermore, $\text{Rmil } \text{A-code } \sigma = \text{Rmil } \text{Leh } \mathbf{i} \sigma = \text{Rmil } \mathbf{i} \sigma = \text{Lmap } \sigma$. As Leh is a bijection of the symmetric group \mathfrak{S}_n onto SE_n , we obtain the following result.

Theorem 5. *The A-code is a bijection of \mathfrak{S}_n onto SE_n having the property:*

$$(2.4) \quad (\text{Rmil}, \text{Lmap}) \sigma = (\text{Max}, \text{Rmil}) \text{ A-code } \sigma \quad (\sigma \in \mathfrak{S}_n).$$

An algorithmic definition of the A-code will be given in Section 5.

3. The B-code

The B-code is based on the decomposition of each permutation as product of disjoint cycles. For a permutation $\sigma = \sigma(1)\sigma(2)\cdots\sigma(n)$ and each $i = 1, 2, \dots, n$ let $k := k(i)$ be the *smallest* integer $k \geq 1$ such that $\sigma^{-k}(i) \leq i$. Then, define

$$\text{B-code } \sigma = (b_1, b_2, \dots, b_n) \quad \text{with} \quad b_i := \sigma^{-k(i)}(i) \quad (1 \leq i \leq n).$$

For example, with the permutation $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 4 & 5 & 2 & 6 & 1 \end{pmatrix}$ we have:
 $\sigma^{-1}(1) = 6, \sigma^{-2}(1) = 5, \sigma^{-3}(1) = 3, \sigma^{-4}(1) = 1$, so that $b_1 = 1$;
 $\sigma^{-1}(2) = 4, \sigma^{-2}(2) = 2$, so that $b_2 = 2$; $\sigma^{-1}(3) = 1$, so that $b_3 = 1$;
 $\sigma^{-1}(4) = 2$, so that $b_4 = 2$; $\sigma^{-1}(5) = 3$, so that $b_5 = 3$;
 $\sigma^{-1}(6) = 5$, so that $b_6 = 5$. Thus, B-code $\sigma = (1, 2, 1, 2, 3, 5)$.

An alternate definition is the following. First, the B-code of the unique permutation from \mathfrak{S}_1 is defined to be the sequence $(1) \in SE_1$. Let $n \geq 2$. When writing each permutation $\sigma \in \mathfrak{S}_n$ of order $n \geq 2$ as a product of its disjoint cycles, the removal of n yields a permutation σ' of order $(n-1)$. Let $b' = (b'_1, b'_2, \dots, b'_{n-1})$ be the B-code of σ' . We define the B-code of σ to be $b := (b'_1, b'_2, \dots, b'_{n-1}, \sigma^{-1}(n))$. By induction on n , we immediately see that the B-code is a bijection of \mathfrak{S}_n onto SE_n .

The following Theorem shows that the set-valued statistics ‘‘Lmap’’ and ‘‘Cyc’’ can be directly read from the B-code.

Theorem 6. *The B-code is a bijection of \mathfrak{S}_n onto SE_n having the property:*

$$(3.1) \quad (\text{Cyc}, \text{Lmap}) \sigma = (\text{Max}, \text{Rmil}) \text{ B-code } \sigma \quad (\sigma \in \mathfrak{S}_n).$$

Proof. By induction, suppose that $\text{Lmap } \sigma' = \text{Rmil } b'$ and $\text{Cyc } \sigma' = \text{Max } b'$. If n is a fixed point of σ , so that $\sigma^{-1}(n) = n$ and $b = (b'_1, \dots, b'_{n-1}, n)$, then $\text{Lmap } \sigma = \text{Lmap } \sigma' \cup \{n\} = \text{Rmil } b' \cup \{n\} = \text{Rmil } \sigma$. Also, $\text{Cyc } \sigma = \text{Cyc } \sigma' \cup \{n\} = \text{Max } b' \cup \{n\} = \text{Max } b$.

When n is not a fixed point of σ , then σ is a product of the form

$$\sigma = \cdots (\cdots \sigma^{-1}(n)n\sigma(n)\cdots) \cdots$$

while σ' may be expressed as

$$\sigma' = \cdots (\cdots \sigma^{-1}(n)\sigma(n)\cdots) \cdots$$

In particular, $\sigma^{-1}(n) < n$, $\sigma(n) < n$ and $\sigma'(\sigma^{-1}(n)) = \sigma(n)$. We have $\text{Cyc } \sigma = \text{Cyc } \sigma' = \text{Max } b' = \text{Max } b$ since $\sigma^{-1}(n) < n$.

To prove $\text{Lmap } \sigma = \text{Rmil } b$, three cases are to be considered, (i) $\sigma(n) = n-1$; (ii) $\sigma(n) \neq n-1$ and $\sigma^{-1}(n-1) < \sigma^{-1}(n)$; (iii) $\sigma(n) \neq n-1$ and $\sigma^{-1}(n-1) > \sigma^{-1}(n)$, each of them materialized by the following three tableaux:

$$\begin{array}{l}
 (i) \quad \begin{array}{ccccccc}
 \text{Id} = & 1 & \cdots & \sigma^{-1}(n) & \cdots & n-1 & n \\
 \sigma = & \sigma(1) & \cdots & n & \cdots & \sigma(n-1) & \sigma(n) = n-1 \\
 \sigma' = & \sigma(1) & \cdots & \sigma(n) = n-1 & \cdots & \sigma(n-1) & *
 \end{array} \\
 \\
 (ii) \quad \begin{array}{ccccccc}
 \text{Id} = & 1 & \cdots & \sigma^{-1}(n-1) & \cdots & \sigma^{-1}(n) & \cdots & n-1 & n \\
 \sigma = & \sigma(1) & \cdots & n-1 & \cdots & n & \cdots & \sigma(n-1) & \sigma(n) \\
 \sigma' = & \sigma(1) & \cdots & n-1 & \cdots & \sigma(n) & \cdots & \sigma(n-1) & *
 \end{array} \\
 \\
 (iii) \quad \begin{array}{ccccccc}
 \text{Id} = & 1 & \cdots & \sigma^{-1}(n) & \cdots & \sigma^{-1}(n-1) & \cdots & n-1 & n \\
 \sigma = & \sigma(1) & \cdots & n & \cdots & n-1 & \cdots & \sigma(n-1) & \sigma(n) \\
 \sigma' = & \sigma(1) & \cdots & \sigma(n) & \cdots & n-1 & \cdots & \sigma(n-1) & *
 \end{array}
 \end{array}$$

In case (i) we get $\text{Lmap } \sigma = \text{Lmap } \sigma'$, $b' = (\dots, \sigma^{-1}(n))$ and $b = (\dots, \sigma^{-1}(n), \sigma^{-1}(n))$, then $\text{Rmil } b = \text{Rmil } b'$.

In case (ii) we clearly have: $\text{Lmap } \sigma = \text{Lmap } \sigma' \cup \{\sigma^{-1}(n)\}$. Also, $b' = (\dots, \sigma^{-1}(n-1))$ and $b = (\dots, \sigma^{-1}(n-1), \sigma^{-1}(n))$. Hence, $\text{Lmap } \sigma = \text{Lmap } \sigma' \cup \{\sigma^{-1}(n)\} = \text{Rmil } b' \cup \{\sigma^{-1}(n)\} = \text{Rmil } b$.

Finally, comes case (iii), which is the hardest one. We have $\text{Lmap } \sigma = (\text{Lmap } \sigma' \cap [1, \sigma^{-1}(n) - 1]) \cup \{\sigma^{-1}(n)\}$, also $b' = (\dots, b'_{n-2}, \sigma^{-1}(n-1))$, $b = (\dots, b'_{n-2}, \sigma^{-1}(n-1), \sigma^{-1}(n))$. But as $\sigma^{-1}(n) < \sigma^{-1}(n-1)$, we have $\text{Rmil } b = (\text{Rmil } b' \cap [1, \sigma^{-1}(n) - 1]) \cup \{\sigma^{-1}(n)\} = (\text{Lmap } \sigma' \cap [1, \sigma^{-1}(n) - 1]) \cup \{\sigma^{-1}(n)\} = \text{Lmap } \sigma$. \square

4. The bijection ϕ

The bijection ϕ , which is the main ingredient in the chain displayed in (1.4), is simply defined as

$$(4.1) \quad \phi := (\text{B-code})^{-1} \circ \text{A-code}.$$

It follows from Theorems 6 and 5 that

$$\begin{aligned}
 (\text{Cyc}, \text{Lmap}) \phi(\sigma) &= (\text{Max}, \text{Rmil}) \text{B-code } \phi(\sigma) \\
 &= (\text{Max}, \text{Rmil}) \text{A-code } \sigma = (\text{Rmil}, \text{Lmap}) \sigma.
 \end{aligned}$$

This proves relation (1.1) and consequently Theorem 1. It also follows from Theorem 5 and/or 6 that the distribution of each pair of statistics stated in Theorem 1 is also equal to the distribution of $(\text{Max}, \text{Rmil})$ on SE_n .

It remains to prove identity (1.5) to achieve the proofs of Theorem 2 and its Corollary. Let $A = ([p_1, q_1], [p_2, q_2], \dots, [p_h, q_h])$ be an ordered partition of $[n]$ into disjoint non-empty intervals, such that $p_j + 1 = q_{j+1}$ for $j = 1, 2, \dots, h-1$ and $p_1 = 1, q_h = n$. Let $G(\sigma) = \{(i, \sigma(i)) : 1 \leq i \leq n\}$ be the graph of a permutation σ from \mathfrak{S}_n . Referring to Fig. 2, where the square $[p_j, q_j] \times [p_j, q_j]$ has been materialized by the four points B, B'', D'', D , we see that σ is A -indecomposable, if for every $j = 1, 2, \dots, h$

- (i) the square $[BB''D''D]$ contains the subgraph $\{(i, \sigma(i)) : p_j \leq i \leq q_j\}$;
- (ii) for every l such that $p_j + 1 \leq l \leq q_j$ the rectangle $[B'B''C''C']$ contains at least one element from $G(\sigma)$.




Fig. 2. Graphs of σ and c

We are then led to the following definition.

Definition. Each subexcedant sequence $c = (c_1, c_2, \dots, c_n)$ from SE_n is said to be A -decomposable, if for every $j = 1, 2, \dots, h$

- (i) the triangle $[BB''D'']$ contains the subgraph $\{(i, c_i) : p_j \leq i \leq q_j\}$;
- (ii) for every l such that $p_j + 1 \leq l \leq q_j$ the rectangle $[B'B''C''C']$ contains at least one element (i, c_i) ($l \leq i \leq q_j$).

Proposition 6. A permutation σ from \mathfrak{S}_n is A -decomposable, if and only if its A -code (resp. B -code) is A -decomposable.

Proof. Let $a = (a_1, a_2, \dots, a_n)$ be the A -code of a permutation σ . If σ is A -decomposable, then for every $j = 1, 2, \dots, h$ and $l = p_j, p_j + 1, \dots, q_j$ the point $(\sigma^{-1}(l), l)$ belongs to the square $[BB''D''D]$. As a_l is equal to 1 plus the number of points $(i, \sigma(i))$ such that $1 \leq i < \sigma^{-1}(l)$ and $\sigma(i) < l$, we have $a_l \geq p_j$, so that the point (l, a_l) belongs to the triangle $[BB''D'']$. Conversely, if $(l, a_l) \in [BB''D'']$, then $(\sigma^{-1}(l), l) \in [BB''D''D]$.

Now, the rectangle $[B'B''C''C']$ contains no element from $G(\sigma)$ if and only if all the points $(\sigma^{-1}(l), l), \dots, (\sigma^{-1}(q_j), q_j)$ are in the square $[C'C''D''D']$. This is equivalent to saying that all the quantities $\sigma^{-1}(l), l, \dots, \sigma^{-1}(q_j), q_j$ lie between l and q_j , which is also equivalent to the fact

that a_l, \dots, a_{q_j} lie between l and q_j , that is, the rectangle $[B'B''C''C']$ has no element (i, a_i) ($l \leq i \leq q_j$).

Next, let $b = (b_1, b_2, \dots, b_n)$ be the B-code of σ . If σ is A -decomposable, the restriction of σ to the interval $[p_j, q_j]$ is a product of cycles all elements of which lie between p_j and q_j . By definition of the B-code all the terms b_{p_j}, \dots, b_{q_j} also lie between p_j and q_j and conversely, if it is the case, all the points $(p_j, \sigma(p_j)), \dots, (q_j, \sigma(q_j))$ belong to the square $[BB''D''D]$. The same argument can be applied when all the points $(l, \sigma(l)), \dots, (q_j, \sigma(q_j))$ belong to the square $[C'C''D''D']$. All terms b_l, \dots, b_{q_j} are greater than or equal to l and the rectangle $[B'B''C''C']$ contains no element of the form (i, b_i) with $l \leq i \leq q_j$. \square

Thus, if σ is A -decomposable, so are A-code σ and the composition product $(\text{B-code})^{-1} \text{A-code}(\sigma) = \phi(\sigma)$. This proves identity (1.5) and then Theorem 2 and its corollary.

5. Algorithmic definitions and examples

Although the A-code has been greatly described in various forms (see, e.g., [Kn98], p. 14), we give a full algorithmic definition, which is to be compared with the analogous definition for the B-code.

Algorithmic definition of A-code. Let $\sigma = \sigma(1)\sigma(2)\cdots\sigma(n)$ be a permutation of $12\cdots n$. By definition the A-code of σ is the sequence $a = (a_1, a_2, \dots, a_n)$ where for each $i = 1, 2, \dots, n$

$$a_i := \#\{j : 1 \leq j \leq i, \sigma^{-1}(j) \leq \sigma^{-1}(i)\},$$

or still

$$(5.1) \quad a_i := \#\{\sigma(k) : 1 \leq \sigma(k) \leq i, k \leq \sigma^{-1}(i)\}.$$

Thus, a_i is equal to 1 plus the number of letters less than i , to the left of i , in the word $\sigma = \sigma(1)\sigma(2)\cdots\sigma(n)$.

For instance, with $\sigma = 4, 6, 1, 2, 3, 5$ the A-code of σ is equal to $a = (1, 2, 3, 1, 5, 2)$: $a_1 = 1$, $a_2 = 2$ as 1 is to the left of 2, $a_3 = 3$ as 1 and 2 are to the left of 3, $a_4 = 1$, as 4 is the leftmost letter of σ , etc. Thus,

$$(5.2) \quad \text{A-code}(4, 6, 1, 2, 3, 5) = (1, 2, 3, 1, 5, 2).$$

Algorithmic definition of A-code⁻¹. Given $a = (a_1, a_2, \dots, a_n) \in \text{SE}_n$ write a word with n empty places numbered 1 to n from left to right. First, move the letter n to the a_n -th leftmost place; let σ_n be the resulting word (having one non-empty letter!). Next, move $(n-1)$ to the place having $a_{n-1} - 1$ empty letters to its left. Let σ_{n-2} be the resulting word (having

two non-empty letters). Move $(n - 2)$ to the place having $a_{n-2} - 1$ empty letters to its left, etc. Thus, $\text{A-code}^{-1}(a)$ is the final permutation σ_1 .

For instance, start with $a = (1, 2, 1, 2, 3, 5)$. We successively get:

$$\begin{array}{rcl}
 & * * * * * & \\
 \sigma_6 = & * * * * 6 * & a_6 = 5 \\
 \sigma_5 = & * * 5 * 6 * & a_5 = 3 \\
 \sigma_4 = & * 4 5 * 6 * & a_4 = 2 \\
 \sigma_3 = & 3 4 5 * 6 * & a_3 = 1 \\
 \sigma_2 = & 3 4 5 * 6 2 & a_2 = 2 \\
 \sigma_1 = & 3 4 5 1 6 2 & a_1 = 1
 \end{array}$$

Thus

$$(5.3) \quad \text{A-code}^{-1}(1, 2, 1, 2, 3, 5) = 3, 4, 5, 1, 6, 2.$$

Algorithmic definition of B-code. Let $\sigma = \sigma(1)\sigma(2) \cdots \sigma(n) \in \mathfrak{S}_n$. Its B-code $b = (b_1, b_2, \dots, b_n)$ is calculated as follows. First, b_n is the place occupied by n in $\sigma_n := \sigma$. Permute the two letters n and $\sigma(n)$ in the word σ . Let σ_{n-1} be the resulting word. Then, b_{n-1} is the place occupied by $(n - 1)$ in σ_{n-1} . Next, permute the two letters $(n - 2)$ and $\sigma(n - 2)$ in σ_{n-1} and let σ_{n-2} be the resulting word. Let b_{n-2} is the place occupied by $(n - 2)$ in σ_{n-2} . Permute $(n - 3)$ and $\sigma(n - 3)$ in σ_{n-2} , etc. The B-code of σ is (b_1, b_2, \dots, b_n) .

Start with $\sigma = 3, 4, 5, 2, 6, 1$. We successively get:

$$\begin{array}{rcl}
 \text{Id} = & 1 2 3 4 5 6 & \\
 \sigma_6 = & 3 4 5 2 6 1 & b_6 = 5 \\
 \sigma_5 = & 3 4 5 2 1 \mathbf{6} & b_5 = 3 \\
 \sigma_4 = & 3 4 1 2 \mathbf{5 6} & b_4 = 2 \\
 \sigma_3 = & 3 2 1 \mathbf{4 5 6} & b_3 = 1 \\
 \sigma_2 = & 1 2 \mathbf{3 4 5 6} & b_2 = 2 \\
 \sigma_1 = & 1 \mathbf{2 3 4 5 6} & b_1 = 1
 \end{array}$$

Thus

$$(5.4) \quad \text{B-code}(3, 4, 5, 2, 6, 1) = (1, 2, 1, 2, 3, 5).$$

Algorithmic definition of B-code⁻¹. Let $b = (b_1, b_2, \dots, b_n) \in \text{SE}_n$. Start with the identity permutation $\sigma_1 = 1, 2, \dots, n$. In σ_1 exchange 2 and the letter at the b_2 -th place. Let σ_2 be the resulting word. In σ_2 permute 3 and the letter at the b_3 -th place. Let σ_3 be the resulting word. In σ_3 permute 4 and the letter at the b_4 -th place, etc. The permutation $\sigma = \text{B-code}^{-1} b$ is the permutation σ_n .

For example, starting with $b = (1, 2, 3, 1, 5, 2)$. We successively form:

$$\begin{array}{ll} \sigma_1 = 1 \mathbf{2} \mathbf{3} \mathbf{4} \mathbf{5} \mathbf{6} & b_1 = 1 \\ \sigma_2 = 1 \ 2 \ \mathbf{3} \ \mathbf{4} \ \mathbf{5} \ \mathbf{6} & b_2 = 2 \\ \sigma_3 = 1 \ 2 \ 3 \ \mathbf{4} \ \mathbf{5} \ \mathbf{6} & b_3 = 3 \\ \sigma_4 = 4 \ 2 \ 3 \ 1 \ \mathbf{5} \ \mathbf{6} & b_4 = 1 \\ \sigma_5 = 4 \ 2 \ 3 \ 1 \ 5 \ \mathbf{6} & b_5 = 5 \\ \sigma_6 = 4 \ 6 \ 3 \ 1 \ 5 \ 2 & b_6 = 2 \end{array}$$

Thus,

$$(5.5) \quad \text{B-code}^{-1}(1, 2, 3, 1, 5, 2) = 4, 6, 3, 1, 5, 2.$$

Let $\Phi := \mathbf{i}\phi\mathbf{i}\phi^{-1}\mathbf{i}$ be the product of the bijections occurring in (1.4). With $\sigma = 6, 4, 1, 2, 3, 5$ the computation of $\Phi(\sigma)$ can be made as follows.

$$\begin{array}{ll} \text{Id} = 1 \ 2 \ 3 \ 4 \ 5 \ 6 & \\ \sigma = 6 \ 4 \ 1 \ 2 \ 3 \ 5 & \\ \mathbf{i}\sigma = 3 \ 4 \ 5 \ 2 \ 6 \ 1 & \\ \text{B-code } \mathbf{i}\sigma = 1 \ 2 \ 1 \ 2 \ 3 \ 5 & \text{(by (5.4))} \\ \text{A-code}^{-1} \text{B-code } \mathbf{i}\sigma = \phi^{-1} \mathbf{i}\sigma = 3 \ 4 \ 5 \ 1 \ 6 \ 2 & \text{(by (5.3))} \\ \mathbf{i}\phi^{-1} \mathbf{i}\sigma = 4 \ 6 \ 1 \ 2 \ 3 \ 5 & \\ \text{A-code } \mathbf{i}\phi^{-1} \mathbf{i}\sigma = 1 \ 2 \ 3 \ 1 \ 5 \ 2 & \text{(by (5.2))} \\ \text{B-code}^{-1} \text{A-code } \mathbf{i}\phi^{-1} \mathbf{i}\sigma = \phi \mathbf{i}\phi^{-1} \mathbf{i}\sigma = 4 \ 6 \ 3 \ 1 \ 5 \ 2 & \text{(by (5.5))} \\ \Phi(\sigma) = \mathbf{i}\phi\mathbf{i}\phi^{-1}\mathbf{i}\sigma = 4 \ 6 \ 3 \ 1 \ 5 \ 2. & \end{array}$$

We verify that

$$(\text{Cyc}, \text{Rmil}) \sigma = (\text{Rmil}, \text{Cyc}) \Phi(\sigma) = (\{1, 2\}, \{1, 2, 3, 5\}).$$

6. Concluding remarks and Tables

The bijection constructed by Cori [Cor08] only preserves the *cardinalities* “cyc” and “lmap”, but not the sets “Cyc” and “Lmap.” With the example used in his paper, the permutation

$$\theta = 6, 5, 7, 4, 2, 10, 3, 8, 9 = (1, 6, 10)(2, 5)(3, 7)(4)(8)(9)$$

is mapped onto

$$\theta' = 4, 6, 5, 7, 3, 8, 1, 9, 10, 2 = (1, 4, 7)(2, 6, 8, 9, 10)(3, 5),$$

so that $(\text{Lmap}, \text{Cyc}) \theta' = (\{1, 2, 4, 6, 8, 9\}, \{1, 2, 3\}) \neq (\{1, 2, 3, 4, 8, 9\}, \{1, 3, 6\}) = (\text{Cyc}, \text{Lmap}) \theta$. However, $(\text{cyc}, \text{lmap}) \theta = (\text{lmap}, \text{cyc}) \theta' = (6, 3)$.

B=	1, 2, 3, 4	1, 2, 4	1, 3, 4	2, 3, 4	1, 4	2, 4	3, 4	4	Σ
A=1,2,3,4	1								1
1,2,4		1							6
1,3,4			1		1				
2,3,4				1			1	1	
1,4			1		1				11
2,4						1	1	1	
3,4				1		1	2	2	
4				1		1	2	2	6
Σ	1	6			11			6	

$$n = 4$$

Fig. 3. Distribution of (Cyc, Rmil) over \mathfrak{S}_n .

There exist other bijections $\sigma \mapsto a$ such that the sum $\sum_i (a_i - 1)$ is equal to a statistic different from the inversion number “inv,” but having interesting properties. Let us quote the *Tompkins-Paige method* ([To56, Le60, We61]) for generating permutations on a computer. That method was further used in [Ha92, Ha94] to show that the corresponding sum $\sum_i (a_i - 1)$ is equal to the *major index* “maj”. Let us also mention the *Denert coding* [FZ90, Ha94], whose sum $\sum_i (a_i - 1)$ is equal to the *Denert statistic* “den”. Those codings serve to prove that the statistics “inv,” “maj” and “den” are equidistributed on \mathfrak{S}_n , their common distribution being called *Mahonian*.

Let $b = (b_1, b_2, \dots, b_n)$ be the B-code of a permutation $\sigma \in \mathfrak{S}_n$. In its turn the sum $\text{env } \sigma := \sum_i (b_i - 1)$ becomes a new Mahonian statistic. Moreover, it follows from the properties of the bijection ϕ defined in (4.1) that the two three-variable statistics (env, Cyc, Lmap) and (inv, Rmil, Lmap) are equidistributed on \mathfrak{S}_n .

References

- [Com74] Comtet, Louis, *Advanced Combinatorics*. D. Reidel, Boston, 1974.
- [Cor08] Cori, Robert, Indecomposable permutations, hypermaps and labeled Dyck paths, preprint, arXiv:0812.0440, 2008, 30 p., to appear in *J. Combin. Theory, ser. A*.
- [CM92] Cori, Robert; Machí, A., Maps, hypermaps and their automorphisms, a survey, *Expo. Math.*, vol. **10**, 1992, p. 403–467.
- [FZ90] Foata, Dominique; Zeilberger, Doron, Denert’s Permutation Statistic Is Indeed Euler-Mahonian, *Studies in Appl. Math.*, vol. **83**, 1990, p. 31-59.
- [Ha92] Han, Guo-Niu, Une courte démonstration d’un résultat sur la Z -statistique, *C. R. Acad. Sci. Paris*, vol. **314**, **Série I**, 1992, p. 969–971.
- [Ha94] Han, Guo-Niu, Une transformation fondamentale sur les réarrangements de mots, *Adv. in Math.*, vol. **105(1)**, 1994, p. 26–41.
- [Kn98] Knuth, Donald E., *The Art of Computer Programming*, vol. 3, *Sorting and Searching* (second edition). Addison-Wesley, Boston, 1998.
- [Le60] Lehmer, D. H., Teaching combinatorial tricks to a computer, Proc. Sympos. Appl. Math., vol. **10**, 1960, p. 179–193. Amer. Math. Soc., Providence, R.I.
- [Lo83] M. Lothaire, *Combinatorics on Words*. Addison-Wesley, Encyclopedia of Math. and its Appl., **17**, London, 1983.
- [OR04] Ossona de Mendez, P.; Rosenstiehl, P., Transitivity and connectivity of permutations, *Combinatorica*, vol. **24**, 2004, p. 487–502.
- [RV96] Roblet, E.; Viennot, X.G., Théorie combinatoire des T-fractions et approximants de Padé en deux points, *Discrete Math.*, vol. **153**, 1996, p. 271–288.
- [Ri58] Riordan, John, *An introduction to Combinatorial Analysis*. John Wiley, New York, 1958.
- [To56] Tompkins, C. B., Machine attacks on problems whose variables are permutations. *Proceedings of Symposia in Applied Mathematics*, vol. **VI**, Numerical Analysis, McGraw-Hill, New York, 1956, 195-211.
- [We61] Wells, Mark B., Generation of permutations by transposition, *Math. Computation*, vol. **15**, 1961, p. 192-195.

Dominique Foata
 Institut Lothaire
 1, rue Murner
 F-67000 Strasbourg, France
 foata@math.u-strasbg.fr

Guo-Niu Han
 I.R.M.A. UMR 7501
 Université de Strasbourg et CNRS
 7, rue René-Descartes
 F-67084 Strasbourg, France
 guoniu@math.u-strasbg.fr