



HAL
open science

On the frequency of partial quotients of regular continued fractions

Ai-Hua Fan, Lingmin Liao, Ji-Hua Ma

► **To cite this version:**

Ai-Hua Fan, Lingmin Liao, Ji-Hua Ma. On the frequency of partial quotients of regular continued fractions. *Mathematical Proceedings of the Cambridge Philosophical Society*, 2010, 148, pp.179–192. hal-00396379

HAL Id: hal-00396379

<https://hal.science/hal-00396379>

Submitted on 17 Jun 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE FREQUENCY OF PARTIAL QUOTIENTS OF REGULAR CONTINUED FRACTIONS

AI-HUA FAN, LING-MIN LIAO, AND JI-HUA MA

ABSTRACT. We consider sets of real numbers in $[0, 1)$ with prescribed frequencies of partial quotients in their regular continued fraction expansions. It is shown that the Hausdorff dimensions of these sets, always bounded from below by $1/2$, are given by a modified variational principle.

1. INTRODUCTION

Let \mathbb{Q}^c denote the set of irrational number. It is well-known that each $x \in [0, 1) \cap \mathbb{Q}^c$ possesses a unique continued fraction expansion of the form

$$x = \frac{1}{a_1(x) + \frac{1}{a_2(x) + \frac{1}{a_3(x) + \ddots}}}, \quad (1.1)$$

where $a_k(x) \in \mathbb{N} := \{1, 2, 3, \dots\}$ is the k -th *partial quotient* of x . This expansion is usually denoted by $x = [a_1(x), a_2(x), \dots]$. For each $j \in \mathbb{N}$, define the *frequency* of the digit j in the continued fraction expansion of x by

$$\tau_j(x) := \lim_{n \rightarrow \infty} \frac{\tau_j(x, n)}{n},$$

whenever the limit exists, where

$$\tau_j(x, n) := \text{Card}\{1 \leq k \leq n : a_k(x) = j\}.$$

This paper is concerned with sets of real numbers with prescribed digit frequencies in their continued fraction expansions. To be precise, let $\vec{p} = (p_1, p_2, \dots)$ be a probability vector with $p_j \geq 0$ for all $j \in \mathbb{N}$ and $\sum_{j=1}^{\infty} p_j = 1$, which will be called a *frequency vector* in the sequel. Our purpose is to determine the Hausdorff dimension of the set

$$\mathcal{E}_{\vec{p}} := \{x \in [0, 1) \cap \mathbb{Q}^c : \tau_j(x) = p_j \ \forall j \geq 1\}.$$

Let us first recall some notation. For any $a_1, a_2, \dots, a_n \in \mathbb{N}$, we call

$$I(a_1, a_2, \dots, a_n) := \{x \in [0, 1) : a_1(x) = a_1, a_2(x) = a_2, \dots, a_n(x) = a_n\}$$

a *rank- n basic interval*. Let $T : [0, 1) \rightarrow [0, 1)$ be the Gauss transformation defined by

$$T(0) = 0, \quad T(x) = 1/x \pmod{1} \quad \text{for } x \in (0, 1).$$

For a given frequency vector $\vec{p} = (p_1, p_2, \dots)$, we denote by $\mathcal{N}(\vec{p})$ the set of T -invariant ergodic probability measures μ such that

$$\int |\log x| d\mu < \infty \quad \text{and} \quad \mu(I(j)) = p_j \quad \text{for all } j \geq 1. \quad (1.2)$$

Let h_μ stand for the measure-theoretical entropy of μ , and $\dim_H A$ for the Hausdorff dimension of a set A . The main result of this paper can be stated as follows.

Theorem 1.1. *For any frequency vector \vec{p} , one has*

$$\dim_H(\mathcal{E}_{\vec{p}}) = \max \left\{ \frac{1}{2}, \sup_{\mu \in \mathcal{N}(\vec{p})} \frac{h_\mu}{2 \int |\log x| d\mu} \right\},$$

where the ‘‘sup’’ is set to be zero if $\mathcal{N}(\vec{p}) = \emptyset$.

By virtue of $\log |T'(x)| = 2|\log x|$, we see that $2 \int |\log x| d\mu$ is the Liapunov exponent of the measure μ . Therefore, the ‘‘sup’’ in the above is a variational formula which relates the Hausdorff dimension to the entropy and Liapunov exponent of measures.

Theorem 1.1 provides a complete solution to the long standing problem that requests an exact formula for $\dim_H(\mathcal{E}_{\vec{p}})$. Let us recall some partial results in the literature. In 1966, under the condition that $\sum_{j=1}^{\infty} p_j \log j < \infty$, Kinney and Pitcher [9] showed that

$$\dim_H(\mathcal{E}_{\vec{p}}) \geq \frac{-\sum_{j=1}^{\infty} p_j \log p_j}{2 \int |\log x| d\mu_{\vec{p}}},$$

where $\mu_{\vec{p}}$ is the Bernoulli measure on $[0, 1]$ defined by

$$\mu_{\vec{p}}(I(a_1, a_2, \dots, a_n)) = \prod_{j=1}^n p_{a_j}.$$

The above lower bound is just the Hausdorff dimension of the Bernoulli measure $\mu_{\vec{p}}$. However, by a result of Kifer, Peres and Weiss in 2001, this is not an optimal lower bound. Indeed, it is shown in [8] that, for any Bernoulli measure $\mu_{\vec{p}}$,

$$\dim_H \mu_{\vec{p}} \leq 1 - 10^{-7}$$

This surprising fact indicates that the collection of Bernoulli measures are insufficient for providing the correct lower bound for $\dim_H(\mathcal{E}_{\vec{p}})$.

In 1975, under the same condition $\sum_{j=1}^{\infty} p_j \log j < \infty$, Billingsley and Henningsen [2] obtained an improved lower bound

$$\dim_H(\mathcal{E}_{\vec{p}}) \geq \sup_{\mu \in \mathcal{N}(\vec{p})} \frac{h_\mu}{2 \int |\log x| d\mu}.$$

Moreover, they proved that, for any fixed $N \in \mathbb{N}$, this lower bound is the exact Hausdorff dimension of the set

$$\{x \in \mathcal{E}_{\vec{p}} : a_n(x) \leq N \text{ for all } n \geq 1\}$$

provided that $p_j = 0$ for all $j > N$. It is therefore quite natural to guess that this lower bound is the right value for $\dim_H \mathcal{E}_{\vec{p}}$ in general. However, as will be shown in Theorem 1.1, this is not the case.

Actually, the lower bound due to Billingsley and Henningsen is only a half of the correct lower bound. The other half of the lower bound, namely, $\dim_H(\mathcal{E}_{\vec{p}}) \geq 1/2$, can be proved basing on Lemma 2.4 in [11]. However, we will give a direct proof in this paper.

The upper bound estimate is more difficult. In its proof, we will use techniques from [11] and [2] to estimate the lengths of basic intervals. Not incidentally, an entropy-involved combinatorial lemma borrowed from [5] (see Lemma 2.7) will play an important role.

The paper is organized as follows. In Section 2, we give some preliminaries on the basic intervals and on the entropy of finite words. In Section 3, we establish the upper bound in Theorem 1.1. In Section 4, we prove that $\dim_H(\mathcal{E}_{\bar{p}}) \geq 1/2$ and show that we can drop the condition $\sum_{j=1}^{\infty} p_j \log j < \infty$ in Billingsley and Henningsen's theorem and then obtain the lower bound in Theorem 1.1. The last section serves as a remark.

2. PRELIMINARY

Let $x = [a_1(x), a_2(x), \dots] \in [0, 1) \cap \mathbb{Q}^c$. The n -th convergent in the continued fraction expansion of x is defined by

$$\frac{p_n}{q_n} := \frac{p_n(a_1(x), \dots, a_n(x))}{q_n(a_1(x), \dots, a_n(x))} = \frac{1}{a_1(x) + \frac{1}{a_2(x) + \dots + \frac{1}{a_n(x)}}}.$$

For ease of notation, we shall drop the argument x in what follows. It is known (see [6] p.9) that p_n, q_n can be obtained by the recursive relations:

$$\begin{aligned} p_{-1} &= 1, \quad p_0 = 0, \quad p_n = a_n p_{n-1} + p_{n-2} & (n \geq 2), \\ q_{-1} &= 0, \quad q_0 = 1, \quad q_n = a_n q_{n-1} + q_{n-2} & (n \geq 2). \end{aligned}$$

By the above recursion relations, we have the following results.

Lemma 2.1 ([6]). *Let $q_n = q_n(a_1, \dots, a_n)$ and $p_n = p_n(a_1, \dots, a_n)$, we have*

- (i) $p_{n-1}q_n - p_nq_{n-1} = (-1)^n$;
- (ii) $q_n \geq 2^{\frac{n-1}{2}}$, $\prod_{k=1}^n a_k \leq q_n \leq \prod_{k=1}^n (a_k + 1)$.

Lemma 2.2 ([14]). *For any $a_1, a_2, \dots, a_n, b \in \mathbb{N}$,*

$$\frac{b+1}{2} \leq \frac{q_{n+1}(a_1, \dots, a_j, b, a_{j+1}, \dots, a_n)}{q_n(a_1, \dots, a_j, a_{j+1}, \dots, a_n)} \leq b+1 \quad (\forall 1 \leq j < n).$$

Recall that for any $a_1, a_2, \dots, a_n \in \mathbb{N}$, the set

$$I(a_1, a_2, \dots, a_n) = \{x \in [0, 1) : a_1(x) = a_1, a_2(x) = a_2, \dots, a_n(x) = a_n\}$$

is a rank- n basic interval. We write $|I|$ for the length of an interval I .

Lemma 2.3 ([10] p.18). *The basic interval $I(a_1, a_2, \dots, a_n)$ is an interval with endpoints p_n/q_n and $(p_n + p_{n-1})/(q_n + q_{n-1})$. Consequently, one has*

$$\left| I(a_1, \dots, a_n) \right| = \frac{1}{q_n(q_n + q_{n-1})}, \quad (2.1)$$

and

$$\frac{1}{2q_n^2} \leq \left| I(a_1, \dots, a_n) \right| \leq \frac{1}{q_n^2}. \quad (2.2)$$

Lemma 2.4. *We have*

$$\left| I(x_1, \dots, j, \dots, x_n) \right| \leq \frac{8}{(j+1)^2} \left| I(x_1, \dots, \hat{j}, \dots, x_n) \right|,$$

where the notation \hat{j} means “deleting the digit j ”.

Proof. By Lemma 2.2 and (2.2), we have

$$\begin{aligned} |I(x_1, \dots, j, \dots, x_n)| &\leq q_n^{-2}(x_1, \dots, j, \dots, x_n) \\ &\leq \left(\frac{2}{j+1}\right)^2 q_{n-1}^{-2}(x_1, \dots, \hat{j}, \dots, x_n) \\ &\leq \frac{8}{(j+1)^2} |I(x_1, \dots, \hat{j}, \dots, x_n)|. \end{aligned}$$

□

We will simply denote by $I_n(x)$ the rank n basic interval containing x . Suppose that $a_n := a_n(x) \geq 2$ and consider

$$I'_n(x) = I(a_1, \dots, a_{n-1}, a_n - 1) \quad \text{and} \quad I''_n(x) = I(a_1, \dots, a_{n-1}, a_n + 1)$$

which are two rank n basic intervals adjacent to $I_n(x)$. By the recursive equation of q_n and (2.1), one has the following lemma.

Lemma 2.5. *Suppose that $a_n := a_n(x) \geq 2$. Then the lengths of the adjacent intervals $I'_n(x)$ and $I''_n(x)$ are bounded by $\frac{1}{3}|I_n(x)|$ from below and by $3|I_n(x)|$ from above.*

For any $x \in [0, 1] \setminus \mathbb{Q}$ and any word $i_1 \cdots i_k \in \mathbb{N}^k$, ($k \geq 1$), denote by $\tau_{i_1 \cdots i_k}(x, n)$ the number of j , $1 \leq j \leq n$, for which

$$a_j(x) \cdots a_{j+k-1}(x) = i_1 \cdots i_k.$$

For $N \in \mathbb{N}$, define $\Sigma_N := \{1, \dots, N\}$. We shall use the following estimate in [2].

Lemma 2.6 ([2]). *Let $N \geq 1$ and $n \geq 1$. For any $x = [x_1, x_2, \dots] \in [0, 1] \cap \mathbb{Q}^c$ with $x_j \in \Sigma_N$ for $1 \leq j \leq n$. Then for any $k \geq 1$, we have*

$$\log |I_n(x)| \leq 2 \sum_{i_1 \cdots i_k \in \Sigma_N^k} \tau_{i_1 \cdots i_k}(x, n) \log \frac{p_k(i_1, \dots, i_k)}{q_k(i_1, \dots, i_k)} + 8 + \frac{8n}{2^k}. \quad (2.3)$$

Now we turn to the key combinatorial lemma which will be used in the upper bound estimation. Let $\phi : [0, 1] \rightarrow \mathbb{R}$ denote the function

$$\phi(0) = 0, \quad \text{and} \quad \phi(t) = -t \log t \quad \text{for} \quad 0 < t \leq 1.$$

For every word $\omega \in \Sigma_N^n$ of length n and every word $u \in \Sigma_N^k$ of length k , denote by $p(u|\omega)$ the frequency of appearances of u in ω , i.e.,

$$p(u|\omega) = \frac{\tau_u(\omega)}{n - k + 1},$$

where $\tau_u(\omega)$ denote the number of j , $1 \leq j \leq n - k + 1$, for which

$$\omega_j \cdots \omega_{j+k-1} = u.$$

Define

$$H_k(\omega) := \sum_{u \in \Sigma_N^k} \phi(p(u|\omega)).$$

We have the following counting lemma.

Lemma 2.7 ([5]). *For any $h > 0, \epsilon > 0, k \in \mathbb{N}$, and for any $n \in \mathbb{N}$ large enough, we have*

$$\text{Card}\{\omega \in \Sigma_N^n : H_k(\omega) \leq kh\} \leq \exp(n(h + \epsilon)).$$

3. UPPER BOUND

3.1. Some Lemmas. Let $(p(i_1, \dots, i_k))_{i_1 \dots i_k \in \mathbb{N}^k}$ be a probability vector indexed by \mathbb{N}^k . As usual, we denote by $q_k(a_1, \dots, a_k)$ the denominator of the k -th convergent of a real number with leading continued fraction digits a_1, \dots, a_k .

Lemma 3.1. *For each $k \in \mathbb{N}$ and each probability vector $(p(i_1, \dots, i_k))_{i_1 \dots i_k \in \mathbb{N}^k}$,*

$$\sum_{i_1 \dots i_k \in \mathbb{N}^k} -p(i_1, \dots, i_k) \log p(i_1, \dots, i_k) \leq \sum_{i_1 \dots i_k \in \mathbb{N}^k} -p(i_1, \dots, i_k) \log |I(i_1, \dots, i_k)|.$$

Proof. Applying Jensen's inequality to the concave function \log , we have

$$\sum_{i_1 \dots i_k \in \mathbb{N}^k} p(i_1, \dots, i_k) \log \frac{|I(i_1, \dots, i_k)|}{p(i_1, \dots, i_k)} \leq \log \sum_{i_1 \dots i_k \in \mathbb{N}^k} |I(i_1, \dots, i_k)| = 0.$$

□

Lemma 3.2. *Let $\vec{p} = (p_1, p_2, \dots)$ be a probability vector and $\vec{q} = (q_1, q_2, \dots)$ a positive vector. Suppose $-\sum_{j=1}^{\infty} p_j \log q_j = \infty$ and $\sum_{j=1}^{\infty} q_j^s < \infty$ for some positive number s . Then*

$$\limsup_{n \rightarrow \infty} \frac{-\sum_{j=1}^n p_j \log p_j}{-\sum_{j=1}^n p_j \log q_j} \leq s.$$

Proof. This is a consequence of the following inequality (see [13], p.217): for non-negative numbers s_j ($1 \leq j \leq m$) such that $\sum_{j=1}^m s_j = 1$ and any real numbers t_j ($1 \leq j \leq m$), we have

$$\sum_{j=1}^m s_j (t_j - \log s_j) \leq \log \left(\sum_{j=1}^m e^{t_j} \right). \quad (3.1)$$

Fix $n \geq 1$. Let $s_j = p_j$ for $1 \leq j \leq n$ and $s_{n+1} = \sum_{j=n+1}^{\infty} p_j$. Let $t_j = s \log q_j$ for $1 \leq j \leq n$ and $t_{n+1} = 0$. Applying the above inequality (3.1) with $m = n + 1$, we get

$$s \sum_{j=1}^n p_j \log q_j - \sum_{j=1}^n p_j \log p_j - \left(\sum_{j=n+1}^{\infty} p_j \right) \log \left(\sum_{j=n+1}^{\infty} p_j \right) \leq \log \left(1 + \sum_{j=1}^n q_j^s \right).$$

Consequently,

$$\frac{-\sum_{j=1}^n p_j \log p_j}{-\sum_{j=1}^n p_j \log q_j} \leq s + \frac{(\sum_{j=n+1}^{\infty} p_j) \log(\sum_{j=n+1}^{\infty} p_j)}{-\sum_{j=1}^n p_j \log q_j} + \frac{\log(1 + \sum_{j=1}^n q_j^s)}{-\sum_{j=1}^n p_j \log q_j}.$$

Using the facts $-\sum_{j=1}^{\infty} p_j \log q_j = \infty$ and $\sum_{j=1}^{\infty} q_j^s < \infty$, we finish the proof by letting $n \rightarrow \infty$. □

Lemma 3.2 implies the following lemma. Recall that $\Sigma_N^k = \{1, \dots, N\}^k$.

Lemma 3.3. *Let $k \geq 1$. If $(p(i_1, \dots, i_k))_{i_1 \dots i_k \in \mathbb{N}^k}$ is a probability vector such that*

$$\sum_{i_1 \dots i_k \in \mathbb{N}^k} p(i_1, \dots, i_k) \log q_k(i_1, \dots, i_k) = \infty,$$

then we have

$$\limsup_{N \rightarrow \infty} \frac{-\sum_{i_1 \dots i_k \in \Sigma_N^k} p(i_1, \dots, i_k) \log p(i_1, \dots, i_k)}{2 \sum_{i_1 \dots i_k \in \Sigma_N^k} p(i_1, \dots, i_k) \log q_k(i_1, \dots, i_k)} \leq \frac{1}{2}.$$

Proof. By Lemma 2.1, for any $k \in \mathbb{N}$ and any $s > 1/2$, we have

$$\sum_{i_1, \dots, i_k} q_k(i_1, \dots, i_k)^{-2s} \leq \sum_{i_1, \dots, i_k} (i_1 \cdots i_k)^{-2s} = \left(\sum_{j=1}^{\infty} j^{-2s} \right)^k < \infty.$$

Thus we get the result by Lemma 3.2. \square

3.2. Proof of the upper bound. To prove the upper bound, we shall make use of multi-step Markov measures. Let $k \geq 1$, by a $(k-1)$ -step Markov measure, we mean a T -invariant probability measure P on $[0, 1)$ satisfying the Markov property

$$\frac{P(I(a_1, \dots, a_n))}{P(I(a_1, \dots, a_{n-1}))} = \frac{P(I(a_{n-k}, \dots, a_n))}{P(I(a_{n-k}, \dots, a_{n-1}))} \quad (3.2)$$

for all $n \geq 1$ and $a_1, \dots, a_n \in \mathbb{N}$ (see [3], p.9). We may regard a Bernoulli measure as a 0-step Markov measure.

For each $N \geq 2$, we denote by $\mathbb{P}_N^k = \mathbb{P}_N^k(\vec{p})$ the collection of $(k-1)$ -step Markov measures satisfying the condition

$$P(I(j)) = p_j \text{ for } 1 \leq j \leq N-1 \text{ and } P(I(N)) = 1 - \sum_{j=1}^{N-1} p_j. \quad (3.3)$$

These Markov measures are supported by the set of continued fractions for which the partial quotients are bounded from above by N .

For each $i_1 \cdots i_k \in \{1, 2, \dots, N\}^k$, write $p(i_1, \dots, i_k) = P(I(i_1, \dots, i_k))$. Put

$$\alpha_{N,k} := \sup_{P \in \mathbb{P}_N^k} \frac{-\frac{1}{k} \sum p(i_1, \dots, i_k) \log p(i_1, \dots, i_k)}{-2 \sum p(i_1, \dots, i_k) \log(p_k(i_1, \dots, i_k)/q_k(i_1, \dots, i_k))}. \quad (3.4)$$

The argument in [2] (pp.171-172) shows that the following limit

$$\alpha_N := \lim_{k \rightarrow \infty} \alpha_{N,k},$$

exists and coincides with each of the following three limits:

$$\alpha'_N := \lim_{k \rightarrow \infty} \sup_{P \in \mathbb{P}_N^k} \frac{-\sum p(i_1, \dots, i_k) \log p(i_1, \dots, i_k)}{2 \sum p(i_1, \dots, i_k) \log q_k(i_1, \dots, i_k)},$$

$$\alpha''_N := \lim_{k \rightarrow \infty} \sup_{P \in \mathbb{P}_N^k} \frac{-\sum p(i_1, \dots, i_k) \log p(i_1, \dots, i_k)}{-\sum p(i_1, \dots, i_k) \log |I(i_1, \dots, i_k)|},$$

and

$$\alpha'''_N := \lim_{k \rightarrow \infty} \sup_{P \in \mathbb{P}_N^k} \frac{h_P}{2 \int |\log x| dP}.$$

Let

$$\alpha := \limsup_{N \rightarrow \infty} \alpha_N = \limsup_{N \rightarrow \infty} \alpha'_N = \limsup_{N \rightarrow \infty} \alpha''_N = \limsup_{N \rightarrow \infty} \alpha'''_N. \quad (3.5)$$

To prove the upper bound, we need only to prove the following two propositions.

Proposition 3.4. *For any $N \in \mathbb{N}$ large enough, we have*

$$\dim_H(\mathcal{E}_{\vec{p}}) \leq \max \left\{ \frac{1}{2}, \alpha_N \right\}.$$

Proposition 3.5. *We have*

$$\alpha \leq \max \left\{ \frac{1}{2}, \sup_{\mu \in \mathcal{N}(\vec{p})} \frac{h_\mu}{2 \int |\log x| d\mu} \right\}.$$

Remark that we will finally establish the formula in Theorem 1.1, thus by Proposition 3.4 and Proposition 3.5 we have

$$\max \left\{ \frac{1}{2}, \liminf_{N \rightarrow \infty} \alpha_N \right\} = \max \left\{ \frac{1}{2}, \limsup_{N \rightarrow \infty} \alpha_N \right\} = \max \left\{ \frac{1}{2}, \sup_{\mu \in \mathcal{N}(\vec{p})} \frac{h_\mu}{2 \int |\log x| d\mu} \right\},$$

and if $\liminf_{N \rightarrow \infty} \alpha_N \geq 1/2$, then the limit of α_N exists and equals to

$$\sup_{\mu \in \mathcal{N}(\vec{p})} \frac{h_\mu}{2 \int |\log x| d\mu}.$$

Proof of Proposition 3.5. By virtue of (3.5), we set

$$\alpha = \limsup_{N \rightarrow \infty} \lim_{k \rightarrow \infty} \sup_{P \in \mathbb{P}_N^k} \frac{-\sum p(i_1, \dots, i_k) \log p(i_1, \dots, i_k)}{2 \sum p(i_1, \dots, i_k) \log q_k(i_1, \dots, i_k)}.$$

Denote

$$D := \limsup_{N \rightarrow \infty} \lim_{k \rightarrow \infty} \sup_{P \in \mathbb{P}_N^k} 2 \sum p(i_1, \dots, i_k) \log q_k(i_1, \dots, i_k).$$

If $D = \infty$, then by Lemma 3.3, we have $\alpha \leq 1/2$.

Now suppose that $D < \infty$, which is equivalent to

$$\limsup_{N \rightarrow \infty} \lim_{k \rightarrow \infty} \sup_{P \in \mathbb{P}_N^k} \int |\log x| dP < \infty.$$

By (3.5),

$$\alpha = \limsup_{N \rightarrow \infty} \lim_{k \rightarrow \infty} \sup_{P \in \mathbb{P}_N^k} \frac{h_P}{2 \int |\log x| dP}.$$

Without loss of generality, we may suppose that there is a sequence of Markov measures $P_{N,k} \in \mathbb{P}_N^k$ converging to a measure $\mu \in \mathcal{N}(\vec{p})$ in the *weak**-topology such that

$$\limsup_{N \rightarrow \infty} \lim_{k \rightarrow \infty} \frac{h_{P_{N,k}}}{2 \int |\log x| dP_{N,k}} = \alpha.$$

Then by the upper semi-continuity of the entropy function and the weak convergence, we have

$$\alpha \leq \sup_{\mu \in \mathcal{N}(\vec{p})} \frac{h_\mu}{2 \int |\log x| d\mu}.$$

□

Proof of Proposition 3.4. For any fixed integer N which is large enough, and any $\epsilon > 0$, we have

$$E_{\vec{p}} \subset \bigcup_{\ell=1}^{\infty} \bigcap_{n=\ell}^{\infty} H_n(\epsilon, N),$$

where

$$H_n(\epsilon, N) := \left\{ x \in [0, 1) \setminus \mathbb{Q} : \left| \frac{\tau_j(x, n)}{n} - p_j \right| < \epsilon, 1 \leq j \leq N \right\}.$$

For any $\gamma > \max\{1/2, \alpha_N\}$ and for any integer $k \in \mathbb{N}$, we have the γ -Hausdorff measure (see [4], for the definition of \mathcal{H}^γ)

$$\begin{aligned} & \mathcal{H}^\gamma \left(\bigcap_{n=\ell}^{\infty} H_n(\epsilon, N) \right) \\ & \leq \sum_{\substack{|\frac{\tau_j(x, n)}{n} - p_j| < \epsilon, 1 \leq j \leq N \\ n(p_j - \epsilon) < m_j < n(p_j + \epsilon), 1 \leq j \leq N}} |I_n(x)|^\gamma \quad (\forall n \geq \ell) \\ & = \sum_{n(p_j - \epsilon) < m_j < n(p_j + \epsilon), 1 \leq j \leq N} \sum_{x_1 \cdots x_n \in A} |I(x_1, \dots, x_n)|^\gamma, \end{aligned}$$

where

$$A := \{x_1 \cdots x_n \in \Sigma_N^n : \tau_j(x_1 \cdots x_n) = m_j, 1 \leq j \leq N\}.$$

(We recall that $\tau_j(x_1 \cdots x_n)$ denotes the times of appearances of j in $x_1 \cdots x_n$.)

Let $\tilde{n} := \sum_{j=1}^N m_j$. By Lemma 2.4, we have the following estimate by deleting the digits $j > N$ in the first n partial quotients x_1, \dots, x_n of $x \in I_n(x)$:

$$\sum_{x_1 \cdots x_n \in A} |I(x_1, \dots, x_n)|^\gamma \leq \left(\sum_{j=N+1}^{\infty} \frac{8}{(j+1)^{2\gamma}} \right)^{n-\tilde{n}} \sum_{x_1 \cdots x_{\tilde{n}} \in \tilde{A}} |I(x_1, \dots, x_{\tilde{n}})|^\gamma,$$

where

$$\tilde{A} := \{x_1 \cdots x_{\tilde{n}} \in \Sigma_N^{\tilde{n}} : \tau_j(x_1 \cdots x_{\tilde{n}}) = m_j, 1 \leq j \leq N\}.$$

Since $\gamma > 1/2$, for N large enough

$$\sum_{j=N+1}^{\infty} \frac{8}{(j+1)^{2\gamma}} < 1. \quad (3.6)$$

By applying Lemma 2.6, and noticing that $\tau_{i_1 \cdots i_k}(x, \tilde{n}) \leq \tau_{i_1 \cdots i_k}(x_1 \cdots x_{\tilde{n}}) + k$, we have

$$\begin{aligned} & |I(x_1, \dots, x_{\tilde{n}})| = \exp\{\log |I(x_1, \dots, x_{\tilde{n}})|\} \\ & \leq \exp \left\{ 2 \sum_{i_1 \cdots i_k \in \Sigma_N^k} (\tau_{i_1 \cdots i_k}(x_1 \cdots x_{\tilde{n}}) + k) \log \frac{p_k(i_1, \dots, i_k)}{q_k(i_1, \dots, i_k)} + 8 + \frac{8\tilde{n}}{2^k} \right\}. \end{aligned}$$

Thus

$$\begin{aligned} & \sum_{x_1 \cdots x_{\tilde{n}} \in \tilde{A}} |I(x_1, \dots, x_{\tilde{n}})|^\gamma \\ & \leq \sum_{m_{i_1 \cdots i_k}} \sum_{x_1 \cdots x_{\tilde{n}} \in B} \exp \left\{ 2\gamma \sum_{i_1 \cdots i_k \in \Sigma_N^k} (m_{i_1 \cdots i_k} + k) \log \frac{p_k}{q_k} + 8\gamma + \frac{8\tilde{n}\gamma}{2^k} \right\}, \end{aligned}$$

where

$$B := \{x_1 \cdots x_{\tilde{n}} \in \tilde{A} : \tau_{i_1 \cdots i_k}(x_1 \cdots x_{\tilde{n}}) = m_{i_1 \cdots i_k} \quad \forall i_1 \cdots i_k \in \Sigma_N^k\}.$$

Take

$$h = \frac{1}{k} \sum_{i_1 \cdots i_k \in \Sigma_N^k} \phi \left(\frac{m_{i_1 \cdots i_k}}{\tilde{n} - k + 1} \right) \quad (3.7)$$

in Lemma 2.7. We have for any $\delta > 0$ and for \tilde{n} large enough

$$\begin{aligned} & \sum_{x_1 \cdots x_{\tilde{n}} \in B} \exp \left\{ 2\gamma \sum_{i_1 \cdots i_k \in \Sigma_N^k} (m_{i_1 \cdots i_k} + k) \log \frac{p_k}{q_k} + 8\gamma + \frac{8\tilde{n}\gamma}{2^k} \right\} \\ & \leq \exp \left\{ \tilde{n}(h + \delta) + 2\gamma \sum_{i_1 \cdots i_k \in \Sigma_N^k} (m_{i_1 \cdots i_k} + k) \log \frac{p_k}{q_k} + 8\gamma + \frac{8\tilde{n}\gamma}{2^k} \right\}. \end{aligned}$$

Rewrite the right side of the above inequality as

$$\exp \{ \tilde{n} (L(\gamma, k, m_{i_1 \cdots i_k})) \},$$

where

$$L(\gamma, k, m_{i_1 \cdots i_k}) := h + 2\gamma \sum_{i_1 \cdots i_k \in \Sigma_N^k} \frac{m_{i_1 \cdots i_k} + k}{\tilde{n}} \log \frac{p_k}{q_k} + \frac{8\gamma}{\tilde{n}} + \frac{8\gamma}{2^k} + \delta.$$

Since there are at most $(\tilde{n} - k + 1)^{N^k}$ possible words of $i_1 \cdots i_k$ in $\Sigma_N^{\tilde{n}}$, we have

$$\begin{aligned} & \sum_{x_1 \cdots x_{\tilde{n}} \in \tilde{A}} |I(x_1, \cdots, x_{\tilde{n}})|^\gamma \\ & \leq (\tilde{n} - k + 1)^{N^k} \exp \left\{ \tilde{n} \left(\sup_{m_{i_1 \cdots i_k}} L(\gamma, k, m_{i_1 \cdots i_k}) \right) \right\}. \end{aligned}$$

Notice that by the definition of \tilde{A} and B , the possible values of $m_{i_1 \cdots i_k}$ are restricted to satisfy the condition that the frequency of digit j in $x_1 \cdots x_{\tilde{n}}$ is about p_j . Then when $\tilde{n} \rightarrow \infty$, for $i_1, \cdots, i_k \in \Sigma_N^k$

$$\frac{m_{i_1 \cdots i_k}}{\tilde{n} - k + 1} \rightarrow p(i_1, \cdots, i_k), \quad (3.8)$$

and $\{p(i_1, \cdots, i_k) : i_1, \cdots, i_k \in \Sigma_N^k\}$ defines a probability measure P in \mathbb{P}_N^k .

Now take $\delta > 0$ small enough and k large enough such that

$$\frac{8\gamma}{2^k} < \delta, \quad (3.9)$$

and

$$\gamma > \frac{-\frac{1}{k} \sum p(i_1, \cdots, i_k) \log p(i_1, \cdots, i_k) + 5\delta}{-2 \sum p(i_1, \cdots, i_k) \log(p_k(i_1, \cdots, i_k)/q_k(i_1, \cdots, i_k))}. \quad (3.10)$$

The last inequality comes from the definition of α_N , (3.4) and the assumption $\gamma > \alpha_N$.

By (3.8), for sufficiently large \tilde{n} , we have $\frac{8\gamma}{\tilde{n}} < \delta$ and

$$\begin{aligned} & \left| \frac{1}{k} \sum_{i_1 \cdots i_k \in \Sigma_N^k} \phi \left(\frac{m_{i_1 \cdots i_k}}{\tilde{n} - k - 1} \right) - \frac{1}{k} \sum_{i_1 \cdots i_k \in \Sigma_N^k} \phi(p(i_1 \cdots i_k)) \right| < \delta, \\ & \left| \sum_{i_1 \cdots i_k \in \Sigma_N^k} \frac{m_{i_1 \cdots i_k} + k}{\tilde{n}} \log \frac{p_k}{q_k} - \sum_{i_1 \cdots i_k \in \Sigma_N^k} p(i_1 \cdots i_k) \log \frac{p_k}{q_k} \right| < \delta. \end{aligned} \quad (3.11)$$

By (3.7) and (3.11),

$$L(\gamma, k, m_{i_1 \dots i_k}) < \frac{1}{k} \sum_{i_1 \dots i_k \in \Sigma_N^k} \phi(p(i_1 \dots i_k)) + 2\gamma \sum_{i_1 \dots i_k \in \Sigma_N^k} p(i_1 \dots i_k) \log \frac{p_k}{q_k} + 5\delta.$$

Thus (3.10) implies

$$L(\gamma, k, m_{i_1 \dots i_k}) < 0.$$

Hence finally, by (3.6), we can obtain for any $\gamma > \max\{1/2, \alpha_N\}$,

$$\mathcal{H}^\gamma \left(\bigcap_{n=k}^{\infty} H_n(\epsilon, N) \right) < \infty.$$

This implies that $\dim_H(\mathcal{E}_{\bar{p}}) \leq \max\{1/2, \alpha_N\}$ as desired. \square

4. LOWER BOUND

In this section, we first prove $\dim_H(\mathcal{E}_{\bar{p}}) \geq 1/2$. Then we examine what happens if the condition $\sum_{j=1}^{\infty} p_j \log j < \infty$ in Billingsley and Henningsen's theorem is violated. We will see that if the condition is not satisfied, then $\dim_H(\mathcal{E}_{\bar{p}}) = 1/2$ and $\mathcal{N}(p) = \emptyset$.

The following is the key lemma for proving that $\dim_H(\mathcal{E}_{\bar{p}}) \geq 1/2$.

Lemma 4.1. *For any given sequence of positive integers $\{c_n\}_{n \geq 1}$ tending to the infinity, there exists a point $z = (z_1, z_2, \dots) \in \mathcal{E}_{\bar{p}}$ such that $z_n \leq c_n$ for all $n \geq 1$.*

Proof. For any $n \geq 1$, we construct a probability vector $(p_1^{(n)}, p_2^{(n)}, \dots, p_k^{(n)}, \dots)$ such that $p_k^{(n)} > 0$ for all $1 \leq k \leq c_n$ and $\sum_{k=1}^{c_n} p_k^{(n)} = 1$, and that for any $k \geq 1$,

$$\lim_{n \rightarrow \infty} p_k^{(n)} = p_k. \quad (4.1)$$

Consider a product Bernoulli probability \mathbb{P} supported by $\prod_{n=1}^{\infty} \{1, \dots, c_n\}$. For each digit $k \geq 1$, consider the random variables of $x \in \mathbb{N}^{\mathbb{N}}$, $X_i(x) = 1_{\{k\}}(x_i)$, ($i \geq 1$). By Kolmogorov's strong law of large numbers (see [12] p.388), we have for each digit k ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left(\sum_{i=1}^n 1_{\{k\}}(x_i) - \sum_{i=1}^n \mathbb{E}(1_{\{k\}}(x_i)) \right) = 0 \quad \mathbb{P} - a.s.,$$

which implies

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n 1_{\{k\}}(x_i) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n p_k^{(i)} = p_k \quad \mathbb{P} - a.s.. \quad (4.2)$$

That is to say, for \mathbb{P} almost every point in the space $\prod_{n=1}^{\infty} \{1, \dots, c_n\}$, the digit k has the frequency p_k . Considering each point in $\mathbb{N}^{\mathbb{N}}$ as a continued fraction expansion of a number in $[0, 1]$, we complete the proof. \square

Proof of $\dim_H(\mathcal{E}_{\bar{p}}) \geq 1/2$. Take $c_n = n$ in Lemma 4.1, we find a point $z \in \mathcal{E}_{\bar{p}}$, such that

$$z_n = a_n(z) \leq n \quad (\forall n \geq 1). \quad (4.3)$$

For a positive number $b > 1$, set

$$\mathcal{F}_z(b) := \{x \in [0, 1) : a_{k^2}(x) \in (b^{k^2}, 2b^{k^2}]; \ a_k(x) = a_k(z) \text{ if } k \text{ is nonsquare}\}.$$

It is clear that $\mathcal{F}_z(b) \subset \mathcal{E}_{\bar{p}}$ for all $b > 1$. We define a measure μ on $\mathcal{F}_z(b)$. For $n^2 \leq m < (n+1)^2$, set

$$\mu(I_m(x)) = \prod_{k=1}^n \frac{1}{b^{k^2}}. \quad (4.4)$$

Denote by $B(x, r)$ the ball centered at x with radius r . We will show that for any $\theta > 0$, there exists $b > 1$, such that for all $x \in \mathcal{F}_z(b)$,

$$\liminf_{r \rightarrow 0} \frac{\log \mu(B(x, r))}{\log r} \geq \frac{1}{2} - \theta. \quad (4.5)$$

In fact, for any positive number r , there exist integers m and n such that

$$|I_{m+1}(x)| < 3r \leq |I_m(x)| \quad \text{and} \quad n^2 \leq m < (n+1)^2. \quad (4.6)$$

By the construction of $\mathcal{F}_z(b)$, $a_{n^2}(x) > b^{n^2} > 1$. Let $x = [x_1, x_2, \dots]$. By Lemma 2.5, $B(x, r)$ is covered by the union of three adjacent rank n^2 basic intervals, i.e.,

$$B(x, r) \subset I(x_1, x_2, \dots, x_{n^2} - 1) \cup I(x_1, x_2, \dots, x_{n^2}) \cup I(x_1, x_2, \dots, x_{n^2} + 1).$$

By the definition of μ , the above three intervals admit the same measure. Hence by (4.6), we have

$$\frac{\log \mu(B(x, r))}{\log r} \geq \frac{\log 3\mu(I(x_1, x_2, \dots, x_{n^2}))}{\log \frac{1}{3}|I(x_1, x_2, \dots, x_{m+1})|}. \quad (4.7)$$

However, on the one hand, by (4.4)

$$-\log \mu(I(x_1, x_2, \dots, x_{n^2})) = -\log \prod_{k=1}^n \frac{1}{b^{k^2}} = \sum_{k=1}^n k^2 \log b. \quad (4.8)$$

On the other hand, by (2.2) and Lemma 2.1, we have

$$-\log |I(x_1, x_2, \dots, x_{m+1})| \leq \log 2 + \sum_{k=1}^{m+1} 2 \log(x_k + 1).$$

Let us estimate the second term of the sum. First we have

$$\sum_{k=1}^{m+1} 2 \log(x_k + 1) \leq 2 \sum_{k=1}^{n+1} \log(x_{k^2} + 1) + 2 \sum_{k=1}^{m+1} \log(z_k + 1).$$

Since $x_{k^2} \leq 2b^{k^2}$ for all $k \geq 1$, we deduce

$$\begin{aligned} \sum_{k=1}^{m+1} \log(x_{k^2} + 1) &\leq \sum_{k=1}^{n+1} \log(2b^{k^2} + 1) \leq \sum_{k=1}^{n+1} \log(3b^{k^2}) \\ &= (n+1) \log 3 + \sum_{k=1}^{n+1} k^2 \log b. \end{aligned}$$

By (4.3), since $z_n \leq n$, for all $n \geq 1$, we know

$$\sum_{k=1}^{m+1} \log(z_k + 1) \leq \sum_{k=1}^{(n+1)^2} \log(k + 1).$$

Thus

$$\begin{aligned} & -\log |I(x_1, x_2, \dots, x_{m+1})| \\ & \leq \log 2 + 2(n+1) \log 3 + 2 \sum_{k=1}^{n+1} k^2 \log b + 2 \sum_{k=1}^{(n+1)^2} \log(k+1). \end{aligned} \quad (4.9)$$

Combining (4.8) and (4.9), for any $\theta > 0$, take $b > 1$ to be large enough, we have

$$\liminf_{n \rightarrow \infty} \frac{\log \mu(I(x_1, x_2, \dots, x_{n^2}))}{\log |I(x_1, x_2, \dots, x_{m+1})|} \geq \frac{1}{2} - \theta, \quad \forall x \in \mathcal{F}_z(b).$$

Hence by (4.7), we obtain (4.5).

Since θ can be arbitrary small, by Billingsley Theorem ([1]), we have

$$\dim_H(\mathcal{E}_{\vec{p}}) \geq \frac{1}{2}.$$

□

Thus Theorem 1.1 is already proved under the condition $\sum_{j=1}^{\infty} p_j \log j < \infty$.

Now assume $\sum_{j=1}^{\infty} p_j \log j = \infty$. Then for any invariant and ergodic measure μ such that $\mu(I(j)) = p_j$ for all $j \geq 1$, we have

$$\int |\log x| d\mu(x) \geq \sum_{j=1}^{\infty} \mu(I(j)) \log j = \sum_{j=1}^{\infty} p_j \log j = \infty. \quad (4.10)$$

This implies $D = \infty$ (see the proof of Proposition 3.5 for the definition of D). Then by Proposition 3.4, we have $\dim_H(\mathcal{E}_{\vec{p}}) \leq \frac{1}{2}$. Since we have already proved $\dim_H(\mathcal{E}_{\vec{p}}) \geq \frac{1}{2}$, we get

$$\dim_H(\mathcal{E}_{\vec{p}}) = \frac{1}{2} \quad \text{if} \quad \sum_{j=1}^{\infty} p_j \log j = \infty.$$

This is in accordance with the formula of Theorem 1.1 under the convention that $\sup \emptyset = 0$, because (4.10) implies $\mathcal{N}(\vec{p}) = \emptyset$.

Finally, we remark that $\mathcal{N}(\vec{p}) = \emptyset$ if and only if $\sum_{j=1}^{\infty} p_j \log j = \infty$. We have seen the “if” part. For the other part, assume $\sum_{j=1}^{\infty} p_j \log j < \infty$. Then the Bernoulli measure μ such that $\mu(I(j)) = p_j$ satisfies

$$\int |\log x| d\mu(x) \leq \sum_{j=1}^{\infty} \mu(I(j)) \log(j+1) = \sum_{j=1}^{\infty} p_j \log(j+1) < \infty.$$

which implies that $\mathcal{N}(\vec{p}) \neq \emptyset$.

5. A REMARK

As suggested by the referee, we add a remark on a problematic argument appearing in the literature. To obtain an upper bound of the Hausdorff dimension of a set, one usually applies the Billingsley’s theorem by constructing a finite measure P on the set such that

$$|U|^s \leq P(U)$$

(see [4], p.67). For the set $\mathcal{E}_{\vec{p}}$, where $p_j = 0$ for some j , the Markov measure P satisfying (3.3) does not match because the cylinders starting with j do not

charge the measure and the above inequality is obviously not true. This appeared unnoticed for long (see the remarks of Kifer [7], p. 2012).

This problem did exist in the proof of Theorem 2 in [2]. Let us briefly indicate how to get around the problem in the proof of Theorem 2 of [2] when $p_j = 0$ for some j 's. The basic idea is similar to that of Cajar (see [3], p.67) and that of Kifer [7]. Recall that P is a $(k - 1)$ -step Markov measure supported on the set

$$\{x \in [0, 1) \cap \mathbb{Q}^c : a_n(x) \leq N \text{ for all } n \geq 1\}.$$

It is uniquely determined by its values on the k -cylinders, namely,

$$p(i_1, \dots, i_k) = P([i_1, \dots, i_k]), \quad (i_1 \cdots i_k) \in \{1, 2, \dots, N\}^k.$$

Let $0 < \epsilon < 1$ and P_ϵ be the perturbed $(k - 1)$ -Markov measure determined by

$$p_\epsilon(i_1, \dots, i_k) = (1 - \epsilon)P([i_1, \dots, i_k]) + \frac{\epsilon}{N^k}, \quad (i_1 \cdots i_k) \in \{1, 2, \dots, N\}^k.$$

Now, we can apply the Billingsley's theorem with P_ϵ to find an upper bound, and then get the desired result by letting $\epsilon \rightarrow 0$.

In the present paper, we have intentionally avoided using the Billingsley's theorem. The proof of upper bound consists of Propositions 3.4 and 3.5. Proposition 3.5 concerns some calculations for which the zero frequency of some digits will not cause any trouble. In the proof of Proposition 3.4, we have used a "covering argument" depending on the estimate (2.3) instead of using the Billingsley's theorem. This enables us to get the upper bound of the Hausdorff dimension.

Acknowledgments : The authors are grateful to the referee for a number of valuable comments. This work was partially supported by NSFC10728104 (A. H. Fan) and NSFC10771164 (J. H. Ma).

REFERENCES

1. P. Billingsley. *Ergodic theory and information*. (John Wiley and Sons, Inc., New York-London-Sydney 1965).
2. P. Billingsley and I. Henningsen. Hausdorff dimension of some continued-fraction sets. *Z. Wahrscheinlichkeitstheorie verw. Geb.* **31** (1975) 163-173.
3. H. Cajar. *Billingsley dimension in probability spaces*. Lecture Notes in Mathematics, **892** (Springer-Verlag, Berlin-New York, 1981).
4. K. J. Falconer. *Fractal Geometry, Mathematical Foundations and Application*. (John Wiley & Sons, Ltd., Chichester, 1990).
5. E. Glasner and B. Weiss. On the interplay between measurable and topological dynamics. *Handbook of dynamical systems. Vol. 1B* Elsevier B. V., Amsterdam. (2006) 597-648.
6. A. Ya. Khintchine. *Continued Fractions*. (University of Chicago Press, Ill.-London, 1964).
7. Y. Kifer. Fractal dimensions and random transformations. *Trans. A.M.S.* **348** (1996) 2003-2008.
8. Y. Kifer, Y. Peres and B. Weiss. A dimension gap for continued fractions with independent digits. *Israel J. Math.* **124**(1), (2001) 61-76.
9. J. R. Kinney and T. S. Pitcher. The dimension of some sets defined in terms of f -expansions. *Z. Wahrscheinlichkeitstheorie verw. Geb.* **4** (1966) 293-315.
10. M. Iosifescu and C. Kraaikamp. *The Metrical Theory on Continued Fractions*. Mathematics and its Applications, **547** (Kluwer Academic Publishers, Dordrecht, 2002).
11. L. M. Liao, J. H. Ma and B. W. Wang. Dimension of some non-normal continued fraction sets. *Math. Proc. Cambridge Philos. Soc.* **145**(1) (2008) 215-225.
12. A. N. Shiryaev. *Probability, Second Edition*. GTM **95** (Springer-Verlag, New York, 1996).
13. P. Walters. *An introduction to ergodic theory*. (Springer-Verlag, New York-Berlin, 2001).
14. J. Wu. A remark on the growth of the denominators of convergents. *Monatsh. Math.* **147**(3) (2006) 259-264.

AI-HUA FAN: DEPARTMENT OF MATHEMATICS, WUHAN UNIVERSITY, WUHAN, 430072, P.R. CHINA & CNRS UMR 6140-LAMFA, UNIVERSITÉ DE PICARDIE 80039 AMIENS, FRANCE
E-mail address: `ai-hua.fan@u-picardie.fr`

LING-MIN LIAO: DEPARTMENT OF MATHEMATICS, WUHAN UNIVERSITY, WUHAN, 430072, P.R. CHINA & CNRS UMR 6140-LAMFA, UNIVERSITÉ DE PICARDIE 80039 AMIENS, FRANCE
E-mail address: `lingmin.liao@u-picardie.fr`

JI-HUA MA: DEPARTMENT OF MATHEMATICS, WUHAN UNIVERSITY, WUHAN, 430072, P.R. CHINA
E-mail address: `jhma@whu.edu.cn`