



Assessment of metal, metalloid and radionuclide bioaccessibility from mussels to human consumers, using centrifugation and simulated digestion methods coupled with radiotracer techniques

Marc Metian, Laetitia Charbonnier, François Oberhänsli, Paco Bustamante, Ross Jeffree, Jean-Claude Amiard, Michel Warnau

► To cite this version:

Marc Metian, Laetitia Charbonnier, François Oberhänsli, Paco Bustamante, Ross Jeffree, et al.. Assessment of metal, metalloid and radionuclide bioaccessibility from mussels to human consumers, using centrifugation and simulated digestion methods coupled with radiotracer techniques. *Ecotoxicology and Environmental Safety*, 2009, 72 (5), pp.1499-1502. 10.1016/j.ecoenv.2008.10.009 . hal-00397345

HAL Id: hal-00397345

<https://hal.science/hal-00397345>

Submitted on 21 Jun 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Assessment of metal, metalloid and radionuclide bioaccessibility from mussels to human consumers, using centrifugation and simulated digestion methods coupled with radiotracer techniques

Marc Metian^{a,b}, Laetitia Charbonnier^c, François Oberhaënsli^a, Paco Bustamante^b, Ross Jeffree^a,
Jean-Claude Amiard^c & Michel Warnau^{a*}

a. International Atomic Energy Agency – Marine Environment Laboratories (IAEA-MEL),
Principality of Monaco

b. Littoral, Environnement et Sociétés (LIENSs), UMR 6250, CNRS-Université de La Rochelle,
France

c. Nantes Atlantique Universités, Molécules Mer et Santé, EA2160, Université de Nantes, France

*Corresponding Author : Michel Warnau

Present address: LIENSs, UMR 6250 CNRS-Université de La Rochelle

2 rue Olympe de Gouges

F-17042 La Rochelle Cedex 01 (France)

Phone : +33 6 62 53 83 27

E-mail : warnaumichel@yahoo.com

ABSTRACT: The dietary bioaccessibility of 7 elements (^{241}Am , Cd, Co, Cs, Mn, Se and Zn) in the Mediterranean mussels *Mytilus galloprovincialis* (Lamarck, 1819) was assessed for human consumers. In this respect we assessed and compared the proportion of elements associated with the cellular cytosolic (“soluble”) fraction vs. the bioaccessible fraction derived, respectively, from (1) differential centrifugation method and (2) simulated digestion method. Comparisons were carried out on both raw and cooked mussels. Results showed that (1) the centrifugation method systematically underestimated (up to a factor 4) element bioaccessibility in raw mussels compared to the *in vitro* digestion method (e.g., 10 vs. 42% for ^{241}Am), and (2) the cooking process (5 min at 200°C) lead to concentrating the elements in mussel tissues (e.g., by a factor 2 for Zn) and reducing their bioaccessibility. Overall, the simulated *in vitro* digestion method appears as a powerful tool for seafood safety assessment and cooking could contribute in reducing substantially the global trace element intake from mussel tissues (up to 65% for Cd and Cs).

Keywords: Bioavailability; seafood; digestion simulation; differential centrifugation; seafood safety

43 **Statement (according to requirements indicated in “EES Instructions to Authors”).**

44

45 Funding sources that supported the work described in the manuscript were from IAEA-MEL
46 regular budget.

47

48 The present work was mainly carried out using *in vitro* techniques; however it also involved the
49 use of marine bivalves. All experimental studies were conducted in accordance with national and
50 institutional guidelines for the protection of animal welfare.

51

1. Introduction

Among seafood, bivalves generally display a high capacity in bioaccumulating metals, which suggests that the risk for their human consumers may not be negligible (e.g., Chouvelon et al., 2008; Metian et al., 2008). This risk depends primarily on the dietary bioavailable fraction of metal (viz., the fraction that is actually assimilated from the food and can reach the systemic circulation of an organism), that depends itself on the metal speciation within the seafood soft tissues. Before the 1990's, it was generally assumed that the bioavailable fraction was reliably assessed by the metal content of the cytosolic ("soluble") fraction in the cells. Later on, studies on the bioavailability of metals focusing on the "insoluble" fraction of the cells showed that the use of the cytosolic fraction alone underestimated the fraction of the metals that was bioavailable to the higher trophic levels. Nowadays, it is considered that the metals contained in both cytosolic and organelles fractions better reflect the bioavailable fraction (Wallace and Lopez, 1996; Wallace and Luoma, 2003). More recently, simulated digestion methodologies were developed to provide a more realistic assessment of the dietary bioavailable fraction of contaminants (Oomen et al., 2003; Versantvoort et al., 2005; Amiard et al., 2008).

The objective of the present study was to assess the dietary bioaccessible fraction (viz., the fraction resulting from the digestive process that can potentially be assimilated by the organism) of selected elements from mussels to human consumers, using two different methods, i.e., differential centrifugation and *in vitro* simulated digestion. Both methods were coupled with highly sensitive radiotracer techniques by using the corresponding γ -emitting radiotracers of the selected elements, i.e., four metals (Cd, Co, Cs, Mn, Zn), one metalloid (Se) and one artificial radionuclide (^{241}Am). It is noteworthy that γ -emitting radiotracers of Co and Cs are also radionuclides commonly associated with nuclear industry wastes and fallout from nuclear

weapon testing. The present work also investigated the effect of cooking (raw vs. cooked mussels) on the metal content in the mussel soft tissues and on the resulting change in dietary bioaccessibility.

2. Material and Methods

Individuals of the Mediterranean mussel, *Mytilus galloprovincialis* (n = 32; 4-6 cm length), were collected along the coast of the Principality of Monaco and exposed for 3 weeks to the 7 selected radiotracers (^{54}Mn , ^{60}Co , ^{65}Zn , ^{75}Se , ^{109}Cd , ^{134}Cs and ^{241}Am) dissolved in natural 0.45- μm filtered seawater (closed-circuit 70-l aquarium; salinity: 36 p.s.u.; temperature: $21 \pm 0.5^\circ\text{C}$; pH: 8.0 ± 0.1 ; light/dark cycle: 12h/12h). Activity of the radiotracers measured in seawater over the exposure duration was: 0.23 kBq $^{54}\text{Mn l}^{-1}$, 0.48 kBq $^{60}\text{Co l}^{-1}$, 0.58 kBq $^{65}\text{Zn l}^{-1}$, 0.31 kBq $^{75}\text{Se l}^{-1}$, 1 kBq $^{109}\text{Cd l}^{-1}$, 0.58 kBq $^{134}\text{Cs l}^{-1}$ and 0.14 kBq $^{241}\text{Am l}^{-1}$. These radiotracer additions corresponded to stable metal concentrations of 1.1 pmol Mn l^{-1} , 0.6 pmol Co l^{-1} , 30 pmol Zn l^{-1} , 7 pmol Se l^{-1} , 0.3 pmol Cd l^{-1} , 11 pmol Cs l^{-1} , i.e., concentrations more than two orders of magnitude lower than the background concentrations in open seas (Ward, 2000). During this period, scallops were fed twice daily a phytoplankton diet (*Isochrysis galbana*; $5 \cdot 10^4$ cells ml^{-1}) and the seawater and spike were renewed every day for one week then every two days (Hédouin et al., 2006). After the exposure, mussels were placed in clean flowing seawater for 2 days in order to remove the radiotracers loosely bound to the mussels and to clear the gut contents (Metian et al., 2007).

A subsample of the mussels (n = 16) was then collected and kept raw whereas the remaining individuals (n = 16) were placed individually in 50-ml glass beakers, covered and heated for 5 min at 200°C on a hot plate. This temperature of cooking was selected in order to reflect a

99 intermediate temperature between relative low cooking temperatures (~100°C in boiling water)
100 and high ones (250 to 350°C with oven or barbecue).

101 In order to determine the possible effect of cooking on the radiotracer contents in the mussels,
102 tissues and fluids of 8 raw and 8 cooked mussels were radioanalyzed. Gamma-counting was
103 carried out according to the method described in Rodriguez y Baena et al. (2006), using a high-
104 resolution γ -spectrometry system consisting of four coaxial Germanium (N- or P-type) detectors
105 (EGNC 33-195-R, Canberra[®] and Eurysis[®]) connected to a multi-channel analyzer and a
106 computer loaded with a spectra analysis software (Interwinner[®] 6).

107 For the remaining mussels (n = 8 for both cooked and raw individuals), the whole soft tissues
108 were treated (1) by differential centrifugation according to the method described in Bustamante
109 and Miramand (2005) to isolate the cellular cytosolic fraction by ultracentrifugation (28,000 G
110 for 1 h, using a Sorvall RC28S ultracentrifuge) or (2) according to the *in vitro* simulated
111 digestion method as described by Versantvoort et al. (2005). Briefly, this latter method consists
112 of a three-step procedure which simulates quite closely the human digestive processes occurring
113 in the mouth, stomach and small intestine (Versantvoort et al., 2005; Brandon et al., 2006;
114 Amiard et al., 2008). The food matrix was first minced at 4°C and exposed to artificial saliva at
115 pH 6.8 for 5 min. Artificial gastric juice at pH 1.3 was then added for 2 h and finally a mixture of
116 artificial duodenal juice, bile and HCO₃ at pH 8.1–8.2 was added for a further 2 h. The incubation
117 temperature was 37.2 ± 0.2°C. Chemicals and enzymes used are the same as described in
118 Versantvoort et al. (2005) and Amiard et al. (2008) and were purchased from Sigma[®]. After the
119 *in vitro* digestion, the resulting chyme was centrifuged at 1,572 G for 15 min at 37.2 ± 0.2°C.
120 According to Versantvoort et al. (2005), the elements in the supernatant are representative of
121 those occurring in the food (i.e. the mussels) and that are bioaccessible to humans (i.e. the

fraction that can be absorbed by the human gut enterocytes). Pellets and supernatants from both treatments (centrifugation and simulated digestion) were radioanalyzed in order to determine the activity of the 7 radiotracers.

3. Results

The overall results of the subcellular fractioning and simulated *in vitro* digestion methods showed that the bioaccessible fraction of the metals in mussels varied from 10% (^{241}Am in raw mussels) to 92% (^{134}Cs in raw mussels) (Table 1). However, for all the studied elements, the bioaccessible fraction determined using the *in vitro* digestion method was always higher (by a factor of up to 4) than the one assessed using the differential centrifugation method (Table 1).

Examination of Table 2 indicates that the cooking process resulted in concentrating systematically all elements but ^{134}Cs in the mussel soft tissues (by 20 to 70%) and in releasing a significant amount of all elements but ^{241}Am in the cooking juice.

It was also observed that although cooking increased the element concentration in the mussel flesh, the elements remaining in the cooked flesh were less bioaccessible than when occurring in raw tissues, in particular for Cd, Se, and Zn (Table 1).

4. Discussion

The results of our study show clearly that the bioaccessible fraction determined using the simulated *in vitro* digestion method was systematically higher (by a factor of up to 4) than when assessed using the differential centrifugation method (see Table 1). In particular, the latter method considerably underestimated the bioaccessibility of ^{241}Am , Co, Mn, Se and Zn. This

observation was not that surprising as, from the theory, it was expected that the differential centrifugation method would provide a lower bioaccessibility estimate since it is only related to the cytosolic subcellular fraction (Wallace and Luoma, 2003). In contrast, by definition, the simulated *in vitro* digestion method would allow assessing the global metal fraction which is bioaccessible, regardless of the subcellular partitioning of the elements. By using a simplified empirical digestion simulation (acetic acid solution at pH 4), Bragigand et al. (2004) had already shown that some metals (Ag, Cd, Cu and Zn) could be bioavailable from the insoluble subcellular fraction of oyster cells. The *in vitro* digestion previously developed by Oomen et al. (2003) and Versantvoort et al. (2005) and used in the present study is a step forward to evaluate bioaccessibility of metals from seafood products by humans as it mimics quite closely the conditions occurring all along the human digestive tract (constant temperature, succession of enzymatic activities and pH, corresponding to each digestive step occurring from the mouth to the intestine).

The analysis of radiotracer content in raw and cooked mussels demonstrated that the cooking process resulted in concentrating by 20 to 70% most of the studied elements due to the loss of moisture as well as in releasing metals into the cooking juice. In particular, the cooking juice displayed metal concentrations higher by up to one order of magnitude than in the inter-valve fluid of raw mussels (see Table 2), corresponding to similar differences in terms of metal load (both cooking juice and inter-valve fluid were of similar volume).

Nevertheless, the comparison of our data with those previously published showed that a trend can hardly be generalized regarding the cooking effect on the metal content in seafood. Indeed, while some elements were found to concentrate in seafood after cooking (e.g. total and inorganic As in fish and molluscs; Devesa et al., 2001), other elements were not (e.g. Cd, Cu, Pb and Zn in the fish *Tilapia nilotica*; Atta et al., 1997). Furthermore, the element concentration in tissues after

cooking appears to depend on (1) the species considered (e.g. ^{134}Cs is not concentrated in cooked mussel tissues whereas it is in fish tissues; present study and Burger et al., 2004, respectively), (2) the element considered and its chemical speciation (Devesa et al., 2001), and (3) the type of cooking (Ersoy et al., 2006). Although the aforementioned factors are generally reported as influencing the cooking effect, it has to be noted that the species-dependence factor could be partly due to the difference in tissue consideration when different species are investigated. Indeed, studies dealing with bivalves as ours generally consider the whole soft tissues as edible target, whereas fish related studies generally consider only the muscle tissues (fillet). Therefore, differences in the nature of the tissues (e.g., protein composition and moisture content) and in metal interactions with the cellular components in different tissues could lead to contrasting results. This is particularly true when storage organs such as liver / hepatopancreas or kidneys are considered. Indeed these organs are the main sites where detoxification processes take place, which usually result in an increase of excretion capacity or, more generally, in an increase in sequestration capacity and thus in different binding strength of the metals with cellular components (e.g., Metian et al., 2005).

Finally, although cooking resulted in an increase in element concentration in whole mussel flesh, it also appeared that the elements remaining in the cooked flesh were significantly less bioaccessible than those occurring in the raw tissues. This was particularly obvious for Cd, Se and Zn (see Table 1).

Considering the simulated *in vitro* digestion method and taking into account all the parameters (i.e., change in weight and in metal concentration of the cooked flesh, change in bioaccessible fraction), we have assessed that the metal intake for a consumer eating mussels would be reduced by 25% for Mn, 35% for Zn, 40% for Co, 50% for Se and 65% for Cd and ^{134}Cs if the mussels are previously cooked and the cooking juice discarded before consumption. This decrease in

metal intake would be even more important if the bioaccessibility assessed via the subcellular fractioning method was considered (as the decrease in bioaccessibility between raw and cooked mussels was more marked). However, the information provided by the latter method is probably not directly comparable between raw and cooked mussels. Indeed, cooking of the flesh results in changes/damages of the cellular structure (e.g., protein agglutination), which will affect substantially the cytosol (both in its nature and occurrence) as well as the results of a centrifugation approach and their meaning. Hence the use of the centrifugation fractioning method is most probably relevant and informative only when the raw product is considered.

5. Conclusion

The simulated *in vitro* digestion method represents a powerful tool for the safety assessment of commercially important seafood products. Our study indicates that the bioaccessible fraction of metals in mussel soft tissues for human consumers depends on the metal chemical properties (e.g. distribution among cellular components and binding properties) and on the preparation of this seafood (in the present case, raw vs. cooked at 200°C for 5 min). The present work also showed that the cooking process generally concentrated the elements in mussel soft tissues, decreased their bioaccessibility for the consumers and released a significant part of the elements into the cooking juice. Results indicated that, providing the cooking juice is discarded, consumption of cooked mussels can contribute in reducing significantly the global intake of trace elements from seafood in humans (by as much as 65% for Cd and ¹³⁴Cs). Further studies are required on seafood containing norm-exceeding levels of trace elements in order to assess whether the risk for consumers could be decreased in cooked products down to levels allowing safe marketing. This

could be of particular interest in the case of Cd for which the maximum concentration allowed in marketed bivalves is low ($1 \mu\text{g g}^{-1}$ wet wt; EC, 2006).

Acknowledgments

Michel Warnau is an Honorary Senior Research Associate of the National Fund for Scientific Research (NFSR, Belgium) and holds a 2008 Invited Expert position at LIENSs (CNRS-Université de La Rochelle), supported by the Conseil Régional de Poitou-Charente. The work described in the manuscript was supported by IAEA-MEL regular budget and a GIP Seine-Aval PhD grant awarded to Marc Metian. The IAEA is grateful for the support provided to its Marine Environment Laboratories by the Government of the Principality of Monaco. Authors are grateful to three anonymous reviewers for constructive comments on the manuscript.

References

- Amiard, J.C., Amiard-Triquet, C., Charbonnier, L., Mesnil, A., Rainbow, P.S., Wang, W.X. 2008. Bioaccessibility of essential and non-essential metals in commercial shellfish from Western Europe and Asia. *Food Chem. Toxicol.* 46, 2010-2022.
- Atta, M.B., El-Sebaie, L.A., Noaman, M.A., Kassab, H.E. 1997. The effect of cooking on the content of heavy metals in fish (*Tilapia nilotica*). *Food Chem.* 58, 1-4.
- Bragigand, V., Berthet, B., Amiard, J.C., Rainbow, P.S. 2004. Estimates of trace metal bioavailability to humans ingesting contaminated oysters. *Food Chem. Toxicol.* 42, 1893-1902.

240 Brandon, E.F.A., Oomen, A.G., Rompelberg, C.J.M., Versantvoort, C.H.M., van Engelen,
 241 J.G.M., Sips, A.J.A.M. 2006. Consumer product in vitro digestion model: Bioaccessibility of
 242 contaminants and its application in risk assessment. Regul. Toxicol. Pharm. 44, 161-171.

243 Burger, J., Gaines, K.F., Boring, C.S., Snodgrass, J., Stephens Jr., W.L., Gochfeld, M. 2004.
 244 Effects of cooking on radiocesium in fish from the Savannah River: Exposure differences for
 245 the public. Arch. Environ. Contam. Toxicol. 46, 231-235.

246 Bustamante, P., Miramand, P. 2005. Subcellular and body distributions of 17 trace elements in
 247 the variegated scallop *Chlamys varia* from the French coast of the Bay of Biscay. Sci. Total
 248 Environ. 337, 59–73.

249 Chouvelon, T., Warnau, M., Churlaud, C, Bustamante, P. 2008. Hg concentrations and related
 250 risk assessment in coral reef crustaceans, molluscs and fish from New Caledonia. Environ.
 251 Pollut. doi:[10.1016/j.envpol.2008.06.027](https://doi.org/10.1016/j.envpol.2008.06.027)

252 Devesa, V., Macho, M.L., Jalón, M., Urieta, I., Muñoz, O., Súnier, M.A., López, F., Vélez, D.,
 253 Montoro, R. 2001. Arsenic in cooked seafood products: study on the effect of cooking on
 254 total and inorganic arsenic contents. J. Agric. Food Chem. 49, 4132-4140.

255 EC. 2006. Commission Regulation No 1881/2006 of 19 December 2006 setting maximum levels
 256 for certain contaminants in foodstuffs (Text with EEA relevance). Official J. Europ.
 257 Communities L364, 5-24.

258 Ersoy, B., Yanar, Y., Kucukgulmez, A., Celik, M. 2006. Effects of four cooking methods on the
 259 heavy metal concentrations of sea bass fillets (*Dicentrarchus labrax* Linne, 1785). Food
 260 Chem. 99, 748-751.

261 Hédouin, L., Metian, M., Cotret, O., Teyssié, J.-L., Fowler, S.W., Fichez, R., Warnau, M. 2006.
 262 Allometry of heavy metal bioconcentration in the edible tropical clam *Gafrarium tumidum*.
 263 Sci. Total Environ. 66, 154-163.

264 Metian, M., Bustamante, P., Hédouin, L., Warnau, M. 2008. Accumulation of trace elements in
 265 the tropical scallop *Comptopallium radula* from coral reefs in New Caledonia. Environ.
 266 Pollut. 152, 543-552.

267 Metian, M., Bustamante, P., Oberhänsli, F., Teyssié, J.-L., Warnau, M. 2007. Interspecific
 268 comparison of Cd bioaccumulation in European Pectinidae (*Chlamys varia* and *Pecten*
 269 *maximus*). J. Exp. Mar. Biol. Ecol. 353, 58-67.

270 Metian, M., Hédouin, L., Barbot, C., Teyssié, J.L., Fowler, S.W., Goudard, F., Bustamante, P.,
 271 Durand, J.P., Piéri, J., Warnau, M. 2005. Use of radiotracer techniques to study subcellular
 272 distribution of metals and radionuclides in bivalves from the Noumea lagoon, New
 273 Caledonia. Bull. Environ. Contam. Toxicol. 75, 89-93.

274 Oomen, A.G., Rompelberg, C.J.M., Bruil, M.A., Dobbe, C.J.G., Pereboom, D.P.K.H., Sips,
 275 A.J.A.M. 2003. Development of an *in vitro* digestion model for estimating the
 276 bioaccessibility of soil contaminants. Arch. Environ. Contam. Toxicol. 44, 281-287.

277 Rodriguez y Baena, A.M., Metian, M., Teyssié, J.L., De Broyer, C., Warnau, M. 2006.
 278 Experimental evidence for ²³⁴Th bioaccumulation in three Antarctic crustaceans: Potential
 279 implications for particle flux studies. Mar. Chem. 100, 354-365.

280 Versantvoort, C.H.M., Oomen, E., van de Kamp, E., Rompelberg, C.J.M., Sips, A.J. 2005.
 281 Applicability of an *in vitro* digestion model in assessing the bioaccessibility of mycotoxins
 282 from food. Food Chem. Toxicol. 43, 31-40.

283 Wallace, W.G., Lopez, G.R. 1996. Relationship between subcellular cadmium distribution in
 284 prey and cadmium trophic transfer to a predator. Estuaries 19, 923-930.

285 Wallace, W.G., Luoma, S.N. 2003. Subcellular compartmentalization of Cd and Zn in two
 286 bivalves. II. Significance of trophically available metal (TAM). Mar. Ecol. Prog. Ser. 257,
 287 125-137.

288 Ward, N.I. 2000. Trace elements. In: Fifield, F.W., Haines, P.J. (Eds), Environmental Analytical
289 Chemistry, 2nd edition. Blackwell Science, Oxford, pp. 360-392.




Figure 1. Repartition of radiotracer activities (Bq g⁻¹) between tissues and liquid of the mussel *Mytilus galloprovincialis* before and after a cooking process.




Figure 2. Radiotracer availabilities (% of initial activity in tissues) for mussel consumers following either subcellular or digestion methods (with or without the cooking process).

Table 1. Radiotracer bioaccessibility (%; mean \pm SD, n = 8) from mussel soft tissues (raw or cooked) to human consumers, assessed using either differential centrifugation fractioning method or simulated *in vitro* digestion method.

Mussel preparation	Method for assessing bioaccessibility	⁵⁴ Mn	⁶⁰ Co	⁶⁵ Zn	⁷⁵ Se	¹⁰⁹ Cd	¹³⁴ Cs	²⁴¹ Am
Raw	fractioning	36 \pm 6	37 \pm 4	38 \pm 4	65 \pm 3	77 \pm 3	82 \pm 6	10 \pm 3
	digestion	79 \pm 13	79 \pm 10	70 \pm 5	85 \pm 3	72 \pm 4	92 \pm 1	42 \pm 13
Cooked	fractioning	17 \pm 5	17 \pm 3	3 \pm 1	39 \pm 2	20 \pm 3	59 \pm 21	2 \pm 1
	digestion	82 \pm 2	68 \pm 7	47 \pm 12	63 \pm 5	34 \pm 9	80 \pm 8	52 \pm 8

Table 2. Radiotracer activity (Bq g⁻¹ wet wt; mean \pm SD, n = 8) in mussel soft tissues and fluid before and after cooking. Fluid: inter-valve fluid (in raw mussels) or cooking juice (in cooked mussels).

Mussel preparation	Compartment	Weight (g wet wt)	⁵⁴ Mn	⁶⁰ Co	⁶⁵ Zn	⁷⁵ Se	¹⁰⁹ Cd	¹³⁴ Cs	²⁴¹ Am
Raw	soft tissues	2.4 \pm 0.7	41 \pm 10	34 \pm 9	379 \pm 74	73 \pm 27	366 \pm 64	4 \pm 0.6	11 \pm 2
	fluid	5.4 \pm 1.9	13 \pm 7	7 \pm 4	60 \pm 26	23 \pm 5	30 \pm 13	3 \pm 0.7	3 \pm 2
Cooked	soft tissues	1.3 \pm 0.5	54 \pm 21	42 \pm 17	648 \pm 152	86 \pm 34	450 \pm 133	3 \pm 0.6	15 \pm 5
	fluid	4.1 \pm 1.3	75 \pm 49	32 \pm 13	74 \pm 17	74 \pm 22	322 \pm 87	6 \pm 1	2 \pm 2