



## Formal geometric quantization II

Paul-Emile Paradan

### ► To cite this version:

Paul-Emile Paradan. Formal geometric quantization II. Pacific Journal of Mathematics, 2011, 1 (253), pp.169-211. 10.2140/pjm.2011.253.169 . hal-00398077

**HAL Id: hal-00398077**

**<https://hal.science/hal-00398077>**

Submitted on 24 Jun 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# FORMAL GEOMETRIC QUANTIZATION II

PAUL-EMILE PARADAN

ABSTRACT. In this paper we pursue the study of formal geometric quantization of non-compact Hamiltonian manifolds. Our main result is the proof that two quantization process coincide. This fact was obtained by Ma and Zhang in the preprint Arxiv:0812.3989 by completely different means.

## CONTENTS

1. Introduction and statement of results	1
2. Quantizations of non-compact manifolds	4
3. Proof of Theorem 1.4	19
4. Other properties of $\mathcal{Q}^\Phi$	25
5. Example: the cotangent bundle of an orbit	28
References	32

In the previous article [21], we have studied some functorial properties of the “formal geometric quantization” process  $\mathcal{Q}^{-\infty}$ , which is defined on *proper Hamiltonian manifolds*, e.g. *non-compact* Hamiltonian manifolds with *proper* moment map.

There is another way, denoted  $\mathcal{Q}^\Phi$ , of quantizing proper Hamiltonian manifolds by localizing the index of the Dolbeault Dirac operator on the critical points of the square of the moment map [15, 19, 20].

The main purpose of this paper is to provide a geometric proof that the quantization process  $\mathcal{Q}^{-\infty}$  and  $\mathcal{Q}^\Phi$  coincide. This fact was proved by Ma and Zhang in the recent preprint [15] by completely different means.

*Keywords:* moment map ; symplectic reduction ; geometric quantization ; transversally elliptic symbol.

## 1. INTRODUCTION AND STATEMENT OF RESULTS

Let us first recall the definition of the geometric quantization of a smooth and compact Hamiltonian manifold. Then we show two way of extending the notion of geometric quantization to the case of a *non-compact* Hamiltonian manifold.

Let  $K$  be a compact connected Lie group, with Lie algebra  $\mathfrak{k}$ . In the Kostant-Souriau framework, a Hamiltonian  $K$ -manifold  $(M, \Omega, \Phi)$  is pre-quantized if there

---

*Date:* June 2009.

*Keywords :* moment map, reduction, geometric quantization, transversally elliptic symbol.

*1991 Mathematics Subject Classification :* 58F06, 57S15, 19L47.

is an equivariant Hermitian line bundle  $L$  with an invariant Hermitian connection  $\nabla$  such that

$$(1.1) \quad \mathcal{L}(X) - \nabla_{X_M} = i\langle \Phi, X \rangle \quad \text{and} \quad \nabla^2 = -i\Omega,$$

for every  $X \in \mathfrak{k}$ . Here  $X_M$  is the vector field on  $M$  defined by  $X_M(m) = \frac{d}{dt}e^{-tX}m|_0$ .

The data  $(L, \nabla)$  is also called a Kostant-Souriau line bundle, and  $\Phi : M \rightarrow \mathfrak{k}^*$  is the moment map. Remark that conditions (1.1) imply via the equivariant Bianchi formula the relation

$$(1.2) \quad \iota(X_M)\Omega = -d\langle \Phi, X \rangle, \quad X \in \mathfrak{k}.$$

Let us recall the notion of geometric quantization when  $M$  is **compact**. Choose a  $K$ -invariant almost complex structure  $J$  on  $M$  which is compatible with  $\Omega$  in the sense that the symmetric bilinear form  $\Omega(\cdot, J\cdot)$  is a Riemannian metric. Let  $\bar{\partial}_L$  be the Dolbeault operator with coefficients in  $L$ , and let  $\bar{\partial}_L^*$  be its (formal) adjoint. The *Dolbeault-Dirac operator* on  $M$  with coefficients in  $L$  is  $D_L = \bar{\partial}_L + \bar{\partial}_L^*$ , considered as an elliptic operator from  $\mathcal{A}^{0,\text{even}}(M, L)$  to  $\mathcal{A}^{0,\text{odd}}(M, L)$ . Let  $R(K)$  be the representation ring of  $K$ .

**Definition 1.1.** *The geometric quantization of a compact Hamiltonian  $K$ -manifold  $(M, \Omega, \Phi)$  is the element  $\mathcal{Q}_K(M) \in R(K)$  defined as the equivariant index of the Dolbeault-Dirac operator  $D_L$ .*

Let us consider the case of a **proper** Hamiltonian  $K$ -manifold  $M$ : the manifold is (perhaps) **non-compact** but the moment map  $\Phi : M \rightarrow \mathfrak{k}^*$  is supposed to be proper. Under this properness assumption, one defines the *formal geometric quantization* of  $M$  as an element  $\mathcal{Q}_K^-(M)$  that belongs to  $R^{-\infty}(K)$  [30, 21]. Let us recall the definition.

Let  $T$  be a maximal torus of  $K$ . Let  $\mathfrak{t}^*$  be the dual of the Lie algebra of  $T$  containing the weight lattice  $\Lambda^* : \alpha \in \Lambda^*$  if  $i\alpha : \mathfrak{t} \rightarrow i\mathbb{R}$  is the differential of a character of  $T$ . Let  $C_K \subset \mathfrak{t}^*$  be a Weyl chamber, and let  $\widehat{K} := \Lambda^* \cap C_K$  be the set of dominant weights. The ring of characters  $R(K)$  has a  $\mathbb{Z}$ -basis  $V_\mu^K, \mu \in \widehat{K} : V_\mu^K$  is the irreducible representation of  $K$  with highest weight  $\mu$ .

A representation  $E$  of  $K$  is *admissible* if it has finite  $K$ -multiplicities :  $\dim(\text{hom}_K(V_\mu^K, E)) < \infty$  for every  $\mu \in \widehat{K}$ . Let  $R^{-\infty}(K)$  be the Grothendieck group associated to the  $K$ -admissible representations. We have an inclusion map  $R(K) \hookrightarrow R^{-\infty}(K)$  and  $R^{-\infty}(K)$  is canonically identified with  $\text{hom}_{\mathbb{Z}}(R(K), \mathbb{Z})$ .

For any  $\mu \in \widehat{K}$  which is a regular value of moment map  $\Phi$ , the reduced space (or symplectic quotient)  $M_\mu := \Phi^{-1}(K \cdot \mu)/K$  is a *compact* orbifold equipped with a symplectic structure  $\Omega_\mu$ . Moreover  $L_\mu := (L|_{\Phi^{-1}(\mu)} \otimes \mathbb{C}_{-\mu})/K_\mu$  is a Kostant-Souriau line orbibundle over  $(M_\mu, \Omega_\mu)$ . The definition of the index of the Dolbeault-Dirac operator carries over to the orbifold case, hence  $\mathcal{Q}(M_\mu) \in \mathbb{Z}$  is defined. In Section 2.3, we explain how this notion of geometric quantization extends further to the case of singular symplectic quotients. So the integer  $\mathcal{Q}(M_\mu) \in \mathbb{Z}$  is well defined for every  $\mu \in \widehat{K}$ : in particular  $\mathcal{Q}(M_\mu) = 0$  if  $\mu \notin \Phi(M)$ .

**Definition 1.2.** *Let  $(M, \Omega, \Phi)$  be a proper Hamiltonian  $K$ -manifold which is pre-quantized by a Kostant-Souriau line bundle  $L$ . The formal quantization of  $(M, \Omega, \Phi)$*

is the element of  $R^{-\infty}(K)$  defined by

$$\mathcal{Q}_K^{-\infty}(M) = \sum_{\mu \in \hat{K}} \mathcal{Q}(M_\mu) V_\mu^K.$$

When  $M$  is compact, the fact that

$$(1.3) \quad \mathcal{Q}_K(M) = \mathcal{Q}_K^{-\infty}(M)$$

is known as the “quantization commutes with reduction Theorem”. This was conjectured by Guillemin-Sternberg in [9] and was first proved by Meinrenken [17] and Meinrenken-Sjamaar [18]. Other proofs of (1.3) were also given by Tian-Zhang [26] and the author [19]. For complete references on the subject the reader should consult [25, 28].

One of the main feature of the formal geometric quantization  $\mathcal{Q}^{-\infty}$  is its stability relatively to the restriction to subgroups.

**Theorem 1.3** ([21]). *Let  $M$  be a pre-quantized Hamiltonian  $K$ -manifold which is proper. Let  $H \subset K$  be a closed connected Lie subgroup such that  $M$  is still proper as a Hamiltonian  $H$ -manifold. Then  $\mathcal{Q}_K^{-\infty}(M)$  is  $H$ -admissible and we have  $\mathcal{Q}_K^{-\infty}(M)|_H = \mathcal{Q}_H^{-\infty}(M)$  in  $R^{-\infty}(H)$ .*

When  $M$  is a proper Hamiltonian  $K$ -manifold, we can also define another “formal geometric quantization”, denoted

$$(1.4) \quad \mathcal{Q}_K^\Phi(M) \in R^{-\infty}(K),$$

by localizing the index of the Dolbeault-Dirac operator  $D_L$  on the set  $\text{Cr}(\|\Phi\|^2)$  of critical points of the square of the moment map (see Section 2.2 for the precise definition). We proved in previous papers [20, 21, 23] that

$$(1.5) \quad \mathcal{Q}_K^{-\infty}(M) = \mathcal{Q}_K^\Phi(M).$$

in some situations:

- $M$  is a coadjoint orbit of a semi-simple Lie group  $S$  that parametrizes a representation of the discrete series of  $S$ ,
- $M$  is a Hermitian vector space.

In her ICM 2006 plenary lecture [29], Vergne conjectured that (1.5) holds when  $\text{Cr}(\|\Phi\|^2)$  is compact. Recently, Ma and Zhang [15] prove the following generalisation of this conjecture.

**Theorem 1.4.** *The equality (1.5) holds for **any** proper Hamiltonian  $K$ -manifold.*

This article is dedicated to the study of the quantization map  $\mathcal{Q}^\Phi$ :

- In Section 2.2, we give the precise definition of the quantization process  $\mathcal{Q}^\Phi$ . In particular, we refine the constant  $c_\gamma$  appearing in [15][Theorem 0.1].
- In Section 2.4, we explain how to compute the quantization of a point.
- We give in Section 3 another proof of Theorem 1.4 by using the technique of symplectic cutting developed in [21].
- In Section 4, we consider the case where  $K = K_1 \times K_2$  acts on  $M$  in a way that the symplectic reduction  $M//_0 K_1$  is a *smooth* proper  $K_2$ -Hamiltonian manifold. We show then that the  $K_1$ -invariant part of  $\mathcal{Q}_{K_1 \times K_2}^\Phi(M)$  is equal to  $\mathcal{Q}_{K_2}^{\Phi_2}(M//_0 K_1)$ .

In Section 5, we study the example where  $M$  is the cotangent bundle of a homogeneous space:  $M = T^*(K/H)$  where  $H$  is a closed subgroup of  $K$ . We see that  $T^*(K/H)$  is a proper Hamiltonian  $K$ -manifold prequantized by the trivial line bundle. A direct computation gives

$$(1.6) \quad \mathcal{Q}_K^\Phi(T^*(K/H)) = L^2(K/H) \quad \text{in} \quad R^{-\infty}(K).$$

Let us denote  $[T^*(K/H)]_{\mu, K}$  the symplectic reduction at  $\mu \in \widehat{K}$  of the  $K$ -Hamiltonian manifold  $T^*(K/H)$ . Theorem 1.4 together with (1.6) give

$$\mathcal{Q}([T^*(K/H)]_{\mu, K}) = \dim[V_\mu^K]^H,$$

for any  $\mu \in \widehat{K}$ . Here  $[V_\mu^K]^H \subset V_\mu^K$  is the subspace of  $H$ -invariant vectors.

Then we consider the action of a closed connected subgroup  $G \subset K$  on  $T^*(K/H)$ . We first check that  $T^*(K/H)$  is a *proper* Hamiltonian  $G$ -manifold if and only if the restriction  $L^2(K/H)|_G$  is an admissible  $G$ -representation. Then, using Theorem 1.3, we get that

$$(1.7) \quad \mathcal{Q}_G^{-\infty}(T^*(K/H)) = L^2(K/H)|_G \quad \text{in} \quad R^{-\infty}(G).$$

In other words, the multiplicity of  $V_\lambda^G$  in  $L^2(K/H)$  is equal to the quantization of the reduced space  $[T^*(K/H)]_{\lambda, G}$ .

## 2. QUANTIZATIONS OF NON-COMPACT MANIFOLDS

In this section we define the quantization process  $\mathcal{Q}^\Phi$ , and we give another definition of the quantization process  $\mathcal{Q}^{-\infty}$  that uses the notion of symplectic cutting [21].

**2.1. Transversally elliptic symbols.** Here we give the basic definitions from the theory of transversally elliptic symbols (or operators) defined by Atiyah-Singer in [1]. For an axiomatic treatment of the index morphism see Berline-Vergne [6, 7] and Paradan-Vergne [22]. For a short introduction see [19].

Let  $\mathcal{X}$  be a *compact*  $K$ -manifold. Let  $p : T\mathcal{X} \rightarrow \mathcal{X}$  be the projection, and let  $(-, -)_\mathcal{X}$  be a  $K$ -invariant Riemannian metric. If  $E^0, E^1$  are  $K$ -equivariant complex vector bundles over  $\mathcal{X}$ , a  $K$ -equivariant morphism  $\sigma \in \Gamma(T\mathcal{X}, \text{hom}(p^*E^0, p^*E^1))$  is called a *symbol* on  $\mathcal{X}$ . The subset of all  $(x, v) \in T\mathcal{X}$  where<sup>1</sup>  $\sigma(x, v) : E_x^0 \rightarrow E_x^1$  is not invertible is called the *characteristic set* of  $\sigma$ , and is denoted by  $\text{Char}(\sigma)$ .

In the following, the product of a symbol  $\sigma$  by a complex vector bundle  $F \rightarrow M$ , is the symbol

$$\sigma \otimes F$$

defined by  $\sigma \otimes F(x, v) = \sigma(x, v) \otimes \text{Id}_{F_x}$  from  $E_x^0 \otimes F_x$  to  $E_x^1 \otimes F_x$ . Note that  $\text{Char}(\sigma \otimes F) = \text{Char}(\sigma)$ .

Let  $T_K\mathcal{X}$  be the following subset of  $T\mathcal{X}$  :

$$T_K\mathcal{X} = \{(x, v) \in T\mathcal{X}, (v, X_\mathcal{X}(x))_\mathcal{X} = 0 \quad \text{for all } X \in \mathfrak{k}\}.$$

A symbol  $\sigma$  is *elliptic* if  $\sigma$  is invertible outside a compact subset of  $T\mathcal{X}$  (i.e.  $\text{Char}(\sigma)$  is compact), and is  *$K$ -transversally elliptic* if the restriction of  $\sigma$  to  $T_K\mathcal{X}$  is invertible outside a compact subset of  $T_K\mathcal{X}$  (i.e.  $\text{Char}(\sigma) \cap T_{K_2}\mathcal{X}$  is compact). An elliptic symbol  $\sigma$  defines an element in the equivariant  $\mathbf{K}$ -theory of  $T\mathcal{X}$  with compact support, which is denoted by  $\mathbf{K}_K(T\mathcal{X})$ , and the index of  $\sigma$  is a virtual finite dimensional representation of  $K$ , that we denote  $\text{Index}_\mathcal{X}^K(\sigma) \in R(K)$  [2, 3, 4, 5].

---

<sup>1</sup>The map  $\sigma(x, v)$  will be also denote  $\sigma|_x(v)$

Let

$$R_{tc}^{-\infty}(K) \subset R^{-\infty}(K)$$

be the  $R(K)$ -submodule formed by all the infinite sum  $\sum_{\mu \in \widehat{K}} m_\mu V_\mu^K$  where the map  $\mu \in \widehat{K} \mapsto m_\mu \in \mathbb{Z}$  has at most a *polynomial* growth. The  $R(K)$ -module  $R_{tc}^{-\infty}(K)$  is the Grothendieck group associated to the *trace class* virtual  $K$ -representations: we can associate to any  $V \in R_{tc}^{-\infty}(K)$ , its trace  $k \rightarrow \text{Tr}(k, V)$  which is a generalized function on  $K$  invariant by conjugation. Then the trace defines a morphism of  $R(K)$ -module

$$(2.8) \quad R_{tc}^{-\infty}(K) \hookrightarrow \mathcal{C}^{-\infty}(K)^K.$$

A  $K$ -transversally elliptic symbol  $\sigma$  defines an element of  $\mathbf{K}_K(\text{T}_K \mathcal{X})$ , and the index of  $\sigma$  is defined as a trace class virtual representation of  $K$ , that we still denote  $\text{Index}_{\mathcal{X}}^K(\sigma) \in R_{tc}^{-\infty}(K)$ .

Remark that any elliptic symbol of  $\text{T}\mathcal{X}$  is  $K$ -transversally elliptic, hence we have a restriction map  $\mathbf{K}_K(\text{T}\mathcal{X}) \rightarrow \mathbf{K}_K(\text{T}_K \mathcal{X})$ , and a commutative diagram

$$(2.9) \quad \begin{array}{ccc} \mathbf{K}_K(\text{T}\mathcal{X}) & \longrightarrow & \mathbf{K}_K(\text{T}_K \mathcal{X}) \\ \text{Index}_{\mathcal{X}}^K \downarrow & & \downarrow \text{Index}_{\mathcal{X}}^K \\ R(K) & \longrightarrow & R_{tc}^{-\infty}(K) . \end{array}$$

Using the *excision property*, one can easily show that the index map  $\text{Index}_{\mathcal{U}}^K : \mathbf{K}_K(\text{T}_K \mathcal{U}) \rightarrow R_{tc}^{-\infty}(K)$  is still defined when  $\mathcal{U}$  is a  $K$ -invariant relatively compact open subset of a  $K$ -manifold (see [19][section 3.1]).

Suppose now that the group  $K$  is equal to the product  $K_1 \times K_2$ . When a symbol  $\sigma$  is  $K_1 \times K_2$ -transversally elliptic we will be interested in the  $K_1$ -invariant part of its index, that we denote

$$\left[ \text{Index}_{\mathcal{X}}^{K_1 \times K_2}(\sigma) \right]^{K_1} \in R_{tc}^{-\infty}(K_2).$$

An intermediate notion between the “ellipticity” and “ $K_1 \times K_2$ -transversal ellipticity” is the “ $K_1$ -transversal ellipticity”. When a  $K_1 \times K_2$ -equivariant morphism  $\sigma$  is  $K_1$ -transversally elliptic, its index  $\text{Index}_{\mathcal{X}}^{K_1 \times K_2}(\sigma) \in R_{tc}^{-\infty}(K_1 \times K_2)$ , viewed as a generalized function on  $K_1 \times K_2$ , is *smooth* relatively to the variable in  $K_2$ . It implies that  $\text{Index}_{\mathcal{X}}^{K_1 \times K_2}(\sigma) = \sum_{\lambda} \theta(\lambda) \otimes V_{\lambda}^{K_1}$  with

$$\theta(\lambda) \in R(K_2), \quad \forall \lambda \in \widehat{K_1}.$$

In particular, we know that

$$\left[ \text{Index}_{\mathcal{X}}^{K_1 \times K_2}(\sigma) \right]^{K_1} = \theta(0)$$

belongs to  $R(K_2)$ .

Let us recall the multiplicative property of the index map for the product of manifolds that was proved by Atiyah-Singer in [1]. Consider a compact Lie group  $K_2$  acting on two manifolds  $\mathcal{X}_1$  and  $\mathcal{X}_2$ , and assume that another compact Lie group  $K_1$  acts on  $\mathcal{X}_1$  commuting with the action of  $K_2$ .

The external product of complexes on  $\text{T}\mathcal{X}_1$  and  $\text{T}\mathcal{X}_2$  induces a multiplication (see [1, 22]):

$$\odot : K_{K_1 \times K_2}(\text{T}_{K_1} \mathcal{X}_1) \times K_{K_2}(\text{T}_{K_2} \mathcal{X}_2) \longrightarrow K_{K_1 \times K_2}(\text{T}_{K_1 \times K_2}(\mathcal{X}_1 \times \mathcal{X}_2)).$$

The following property will be used frequently in the paper.

**Theorem 2.1** (Multiplicative property). *For any  $[\sigma_1] \in K_{K_1 \times K_2}(\mathrm{T}_{K_1} \mathcal{X}_1)$  and any  $[\sigma_2] \in K_{K_2}(\mathrm{T}_{K_2} \mathcal{X}_2)$  we have*

$$\mathrm{Index}_{\mathcal{X}_1 \times \mathcal{X}_2}^{K_1 \times K_2}([\sigma_1] \odot [\sigma_2]) = \mathrm{Index}_{\mathcal{X}_1}^{K_1 \times K_2}([\sigma_1]) \otimes \mathrm{Index}_{\mathcal{X}_2}^{K_2}([\sigma_2]).$$

We will use in this article the notion of *support* of a generalized character.

**Definition 2.2.** *The support of  $\chi := \sum_{\mu \in \widehat{K}} a_\mu V_\mu^K \in R^{-\infty}(K)$  is the set of  $\mu \in \widehat{K}$  such that  $a_\mu \neq 0$ .*

We will say that  $\chi \in R^{-\infty}(K)$  is supported outside  $B \subset \mathfrak{t}^*$  if the support of  $\chi$  does not intersect  $B$ . Note that an infinite sum  $\sum_{i \in I} \chi_i$  converges in  $R^{-\infty}(K)$  if for each ball

$$B_r = \{\xi \in \mathfrak{t}^* \mid \|\xi\| < r\}$$

the set  $\{i \in I \mid \mathrm{support}(\chi_i) \cap B_r \neq \emptyset\}$  is finite.

**Definition 2.3.** *We denote by  $O(r)$  any character of  $R^{-\infty}(K)$  which is supported outside the ball  $B_r$ .*

**2.2. Definition and first properties of  $\mathcal{Q}^\Phi$ .** Let  $(M, \Omega, \Phi)$  be a proper Hamiltonian  $K$ -manifold prequantized by an equivariant line bundle  $L$ . Let  $J$  be an invariant almost complex structure compatible with  $\Omega$ . Let  $p : \mathrm{T}M \rightarrow M$  be the projection.

Let us first describe the principal symbol of the Dolbeault-Dirac operator  $\overline{\partial}_L + \overline{\partial}_L^*$ . The complex vector bundle  $(\mathrm{T}^*M)^{0,1}$  is  $K$ -equivariantly identified with the tangent bundle  $\mathrm{T}M$  equipped with the complex structure  $J$ . Let  $h$  be the Hermitian structure on  $(\mathrm{T}M, J)$  defined by  $h(v, w) = \Omega(v, Jw) - i\Omega(v, w)$  for  $v, w \in \mathrm{T}M$ . The symbol

$$\mathrm{Thom}(M, J) \in \Gamma(M, \mathrm{hom}(p^*(\wedge_{\mathbb{C}}^{\mathrm{even}} \mathrm{T}M), p^*(\wedge_{\mathbb{C}}^{\mathrm{odd}} \mathrm{T}M)))$$

at  $(m, v) \in \mathrm{T}M$  is equal to the Clifford map

$$(2.10) \quad \mathbf{c}_m(v) : \wedge_{\mathbb{C}}^{\mathrm{even}} \mathrm{T}_m M \longrightarrow \wedge_{\mathbb{C}}^{\mathrm{odd}} \mathrm{T}_m M,$$

where  $\mathbf{c}_m(v).w = v \wedge w - \iota(v)w$  for  $w \in \wedge_{\mathbb{C}}^{\bullet} \mathrm{T}_m M$ . Here  $\iota(v) : \wedge_{\mathbb{C}}^{\bullet} \mathrm{T}_m M \rightarrow \wedge_{\mathbb{C}}^{\bullet-1} \mathrm{T}_m M$  denotes the contraction map relative to  $h$ . Since  $\mathbf{c}_m(v)^2 = -\|v\|^2 \mathrm{Id}$ , the map  $\mathbf{c}_m(v)$  is invertible for all  $v \neq 0$ . Hence the characteristic set of  $\mathrm{Thom}(M, J)$  corresponds to the 0-section of  $\mathrm{T}M$ .

It is a classical fact that the principal symbol of the Dolbeault-Dirac operator  $\overline{\partial}_L + \overline{\partial}_L^*$  is equal to<sup>2</sup>

$$(2.11) \quad \mathrm{Thom}(M, J) \otimes L,$$

see [8]. Here also we have  $\mathrm{Char}(\mathrm{Thom}(M, J) \otimes L) = 0$  – section of  $\mathrm{T}M$ .

**Remark 2.4.** *When the manifold  $M$  is a product  $M_1 \times M_2$  the symbol  $\mathrm{Thom}(M, J) \otimes L$  is equal to the product  $\sigma_1 \odot \sigma_2$  where  $\sigma_k = \mathrm{Thom}(M_k, J_k) \otimes L_k$ .*

---

<sup>2</sup>Here we use an identification  $\mathrm{T}^*M \simeq \mathrm{T}M$  given by an invariant Riemannian metric.

When  $M$  is compact, the symbol  $\text{Thom}(M, J) \otimes L$  is elliptic and then defines an element of the equivariant  $\mathbf{K}$ -group of  $TM$ . The topological index of  $\text{Thom}(M, J) \otimes L \in \mathbf{K}_K(TM)$  is equal to the analytical index of the Dolbeault-Dirac operator  $\bar{\partial}_L + \bar{\partial}_L^*$ :

$$(2.12) \quad \mathcal{Q}_K(M) = \text{Index}_M^K(\text{Thom}(M, J) \otimes L) \quad \text{in} \quad R(K).$$

When  $M$  is not compact the topological index of  $\text{Thom}(M, J) \otimes L$  is not defined. In order to extend the notion of geometric quantization to this setting we deform the symbol  $\text{Thom}(M, J) \otimes L$  in the ‘‘Witten’’ way [19, 20]. Consider the identification  $\xi \mapsto \hat{\xi}, \mathfrak{k}^* \rightarrow \mathfrak{k}$  defined by a  $K$ -invariant scalar product on  $\mathfrak{k}^*$ . We define the *Kirwan vector field* on  $M$ :

$$(2.13) \quad \kappa_m = \left( \widetilde{\Phi(m)} \right)_M(m), \quad m \in M.$$

**Definition 2.5.** *The symbol  $\text{Thom}(M, J) \otimes L$  pushed by the vector field  $\kappa$  is the symbol  $\mathbf{c}^\kappa$  defined by the relation*

$$\mathbf{c}^\kappa|_m(v) = \text{Thom}(M, J) \otimes L|_m(v - \kappa_m)$$

for any  $(m, v) \in TM$ .

Note that  $\mathbf{c}^\kappa|_m(v)$  is invertible except if  $v = \kappa_m$ . If furthermore  $v$  belongs to the subset  $T_K M$  of tangent vectors orthogonal to the  $K$ -orbits, then  $v = 0$  and  $\kappa_m = 0$ . Indeed  $\kappa_m$  is tangent to  $K \cdot m$  while  $v$  is orthogonal.

Since  $\kappa$  is the Hamiltonian vector field of the function  $\frac{1}{2}\|\Phi\|^2$ , the set of zeros of  $\kappa$  coincides with the set  $\text{Cr}(\|\Phi\|^2)$  of critical points of  $\|\Phi\|^2$ . Finally we have

$$\text{Char}(\mathbf{c}^\kappa) \cap T_K M \simeq \text{Cr}(\|\Phi\|^2).$$

In general  $\text{Cr}(\|\Phi\|^2)$  is not compact, so  $\mathbf{c}^\kappa$  does not define a transversally elliptic symbol on  $M$ . In order to define a kind of index of  $\mathbf{c}^\kappa$ , we proceed as follows. For any invariant open relatively compact subset  $U \subset M$  the set  $\text{Char}(\mathbf{c}^\kappa|_U) \cap T_K U \simeq \text{Cr}(\|\Phi\|^2) \cap U$  is compact when

$$(2.14) \quad \partial U \cap \text{Cr}(\|\Phi\|^2) = \emptyset.$$

When (2.14) holds we denote

$$(2.15) \quad \mathcal{Q}_K^\Phi(U) := \text{Index}_U^K(\mathbf{c}^\kappa|_U) \quad \in \quad R_{tc}^{-\infty}(K)$$

the equivariant index of the transversally elliptic symbol  $\mathbf{c}^\kappa|_U$ .

It will be useful to understand the dependance of the generalized character  $\mathcal{Q}_K^\Phi(U)$  relatively to the data  $(U, \Omega, L)$ . So we consider two proper Hamiltonian  $K$ -manifolds  $(M, \Omega, \Phi)$  and  $(M', \Omega', \Phi')$  respectively prequantized by the line bundles  $L$  and  $L'$ . Let  $V \subset M$  and  $V' \subset M'$  two invariant open subsets.

**Proposition 2.6.** • *The generalized character  $\mathcal{Q}_K^\Phi(U)$  does not depend of the choice of an invariant almost complex structure on  $U$  which is compatible with  $\Omega|_U$ .*

• *Suppose that there exists an equivariant diffeomorphism  $\Psi : V \rightarrow V'$  such that*

- (1)  $\Psi^*(\Phi') = \Phi$ ,
- (2)  $\Psi^*(L') = L$ ,
- (3) *there exists an homotopy of symplectic forms taking  $\Psi^*(\Omega'|_{V'})$  to  $\Omega|_V$ .*

Let  $U' \subset \overline{U'} \subset V'$  be an invariant open relatively compact subset such that  $\partial U'$  satisfies (2.14). Take  $U = \Psi^{-1}(U')$ . Then  $\partial U$  satisfies (2.14) and

$$\mathcal{Q}_K^{\Phi'}(U') = \mathcal{Q}_K^{\Phi}(U) \in R^{-\infty}(K).$$

*Proof.* Let us prove the first point. Let  $\mathbf{c}_i^{\kappa}|_U, i = 0, 1$  be the transversally elliptic symbols defined with the compatible almost complex structure  $J_i, i = 0, 1$ . Since the space of compatible almost complex structure is contractible, there exist an homotopy  $J_t, t \in [0, 1]$  of almost complex structures linking  $J_0$  and  $J_1$ . If we use Lemma 2.2 in [19], we know that there exists an invertible bundle map  $A \in \Gamma(U, \text{End}(TU))$ , homotopic to the identity, such that  $A \circ J_0 = J_1 \circ A$ . With the help of  $A$  we prove then that the symbols  $\mathbf{c}_0^{\kappa}|_U$  and  $\mathbf{c}_1^{\kappa}|_U$  define the same class in  $\mathbf{K}_K(T_K U)$  (see [19][Lemma 2.2]). Hence their equivariant index coincide.

Let us prove the second point. The characters  $\mathcal{Q}_K^{\Phi}(U)$  and  $\mathcal{Q}_K^{\Phi'}(U')$  are computed as the equivariant index of the symbols  $\mathbf{c}^{\kappa}|_U$  and  $\mathbf{c}^{\kappa'}|_{U'}$ . Let  $\tilde{\mathbf{c}}^{\kappa}|_U$  the pull back of  $\mathbf{c}^{\kappa'}|_{U'}$  by  $\Psi$ . Thanks to the point (1) and (2), the only thing which differs in the definitions of the symbols  $\mathbf{c}^{\kappa}|_U$  and  $\tilde{\mathbf{c}}^{\kappa}|_U$  are the almost complex structures  $J$  and  $\tilde{J} = \Psi^*(J') : \text{the first one is comptible with } \Omega \text{ and the second one with } \Psi^*(\Omega'|_{V'})$ . Since these two symplectic structure are homotopic, one sees that the almost complex structures  $J$  and  $\tilde{J}$  are also homotopic. So we can conclude like in the first point.  $\square$

Let us recall the basic fact concerning the singular values of  $\|\Phi\|^2$ .

**Lemma 2.7.** *The set of singular values of  $\|\Phi\|^2 : M \rightarrow \mathbb{R}$  forms a sequence  $0 \leq r_1 < r_2 < \dots < r_k < \dots$  which is finite iff  $\text{Cr}(\|\Phi\|^2)$  is compact. In the other case  $\lim_{k \rightarrow \infty} r_k = \infty$ .*

At each regular value  $R$  of  $\text{Cr}(\|\Phi\|^2)$ , we associate the invariant open subset  $M_{<R} := \{\|\Phi\|^2 < R\}$  which satisfies (2.14). The restriction  $\mathbf{c}^{\kappa}|_{M_{<R}}$  defines then a transversally elliptic symbol on  $M_{<R}$ : let  $\mathcal{Q}_K^{\Phi}(M_{<R})$  be its equivariant index.

Let us show that  $\mathcal{Q}_K^{\Phi}(M_{<R})$  has a limit when  $R \rightarrow \infty$ . The set  $\text{Cr}(\|\Phi\|^2)$  has the following decomposition

$$(2.16) \quad \text{Cr}(\|\Phi\|^2) = \bigcup_{\beta \in \mathcal{B}} \underbrace{K \cdot (M^{\tilde{\beta}} \cap \Phi^{-1}(\beta))}_{Z_{\beta}}$$

where the  $\mathcal{B}$  is a subset of the Weyl chamber  $\mathfrak{t}_+^*$ . Note that each part  $Z_{\beta}$  is compact, hence  $\mathcal{B}$  is finite only if  $\text{Cr}(\|\Phi\|^2)$  is compact. When  $\text{Cr}(\|\Phi\|^2)$  is non-compact, the set  $\mathcal{B}$  is infinite, but it is easy to see that  $\mathcal{B} \cap B_r$  is finite for any  $r \geq 0$ . For any  $\beta \in \mathcal{B}$ , we consider a relatively compact open invariant neighborhood  $\mathcal{U}_{\beta}$  of  $Z_{\beta}$  such that  $\text{Cr}(\|\Phi\|^2) \cap \overline{\mathcal{U}_{\beta}} = Z_{\beta}$ .

**Definition 2.8.** *We denote*

$$\mathcal{Q}_K^{\beta}(M) \in R_{tc}^{-\infty}(K)$$

*the index<sup>3</sup> of the transversally elliptic symbol  $\mathbf{c}^{\kappa}|_{\mathcal{U}_{\beta}}$ .*

A simple application of the excision property [19] gives that

$$(2.17) \quad \mathcal{Q}_K^{\Phi}(M_{<R}) = \sum_{\|\beta\|^2 < R} \mathcal{Q}_K^{\beta}(M).$$

---

<sup>3</sup>The index of  $\mathbf{c}^{\kappa}|_{\mathcal{U}_{\beta}}$  was denoted  $RR_{\beta}^K(M, L)$  in [19].

We have now the key fact

**Theorem 2.9.** *The generalized character  $\mathcal{Q}_K^\beta(M)$  is supported outside the open ball  $B_{\|\beta\|}$ .*

*Proof.* Proposition 2.9 follows directly from the computations done in [19]. First consider the case where  $\beta \neq 0$  is a  $K$ -invariant element of  $\mathcal{B}$ . Let  $i : \mathbb{T}_\beta \hookrightarrow T$  be the compact torus generated by  $\beta$ . If  $F$  is  $\mathbb{Z}$ -module we denote  $F \hat{\otimes} R^{-\infty}(\mathbb{T}_\beta)$  the  $\mathbb{Z}$ -module formed by the infinite formal sums  $\sum_a E_a h^a$  taken over the set of weights of  $\mathbb{T}_\beta$ , where  $E_a \in F$  for every  $a$ .

Since  $\mathbb{T}_\beta$  lies in the center of  $K$ , the morphism  $\pi : (k, t) \in K \times \mathbb{T}_\beta \mapsto kt \in K$  induces a map  $\pi^* : R^{-\infty}(K) \rightarrow R^{-\infty}(K) \hat{\otimes} R^{-\infty}(\mathbb{T}_\beta)$ .

The normal bundle  $\mathcal{N}$  of  $M^{\tilde{\beta}}$  in  $M$  inherits a canonical complex structure  $J_{\mathcal{N}}$  on the fibers. We denote by  $\overline{\mathcal{N}} \rightarrow M^{\tilde{\beta}}$  the complex vector bundle with the opposite complex structure. The torus  $\mathbb{T}_\beta$  is included in the center of  $K$ , so the bundle  $\overline{\mathcal{N}}$  and the virtual bundle  $\wedge_{\mathbb{C}}^{\bullet} \overline{\mathcal{N}} := \wedge_{\mathbb{C}}^{\text{even}} \overline{\mathcal{N}} \xrightarrow{0} \wedge_{\mathbb{C}}^{\text{odd}} \overline{\mathcal{N}}$  carry a  $K \times \mathbb{T}_\beta$ -action: they can be considered as elements of  $K_{K \times \mathbb{T}_\beta}(M^{\tilde{\beta}}) = K_K(M^{\tilde{\beta}}) \otimes R(\mathbb{T}_\beta)$ .

In [19], we have defined an inverse of  $\wedge_{\mathbb{C}}^{\bullet} \overline{\mathcal{N}}$ ,  $[\wedge_{\mathbb{C}}^{\bullet} \overline{\mathcal{N}}]_\beta^{-1} \in K_K(M^{\tilde{\beta}}) \hat{\otimes} R^{-\infty}(\mathbb{T}_\beta)$ , which is polarized by  $\beta$ . It means that  $[\wedge_{\mathbb{C}}^{\bullet} \overline{\mathcal{N}}]_\beta^{-1} = \sum_a N_a h^a$  with  $N_a \neq 0$  only if  $\langle a, \beta \rangle \geq 0$ .

We prove in [19] the following localization formula :

$$(2.18) \quad \pi^* \left[ \mathcal{Q}_K^\beta(M) \right] = R R_\beta^{K \times \mathbb{T}_\beta} \left( M^{\tilde{\beta}}, L|_{M^{\tilde{\beta}}} \otimes [\wedge_{\mathbb{C}}^{\bullet} \overline{\mathcal{N}}]_\beta^{-1} \right),$$

as an equality in  $R^{-\infty}(K) \hat{\otimes} R^{-\infty}(\mathbb{T}_\beta)$ . With (2.18) in hand, it is easy to see that  $V_\mu^K$  occurs in the character  $\mathcal{Q}_K^\beta(M)$  only if  $(\mu, \beta) \geq \|\beta\|^2$  (See Lemma 9.4 in [19]).

Now we consider the case where  $\beta \in \mathcal{B}$  is not a  $K$ -invariant element. Let  $\sigma$  be the unique open face of the Weyl chamber  $\mathfrak{t}_+^*$  which contains  $\beta$ . Let  $K_\sigma$  be the corresponding stabilizer subgroup. We consider the symplectic slice  $\mathcal{Y}_\sigma \subset M$ : it is a  $K_\sigma$  invariant Hamiltonian submanifold of  $M$  which is prequantized by the line bundle  $L|_{\mathcal{Y}_\sigma}$ . The restriction of  $\Phi$  to  $\mathcal{Y}_\sigma$  is a moment map  $\Phi_\sigma : \mathcal{Y}_\sigma \rightarrow \mathfrak{k}_\sigma^*$  which is proper in a neighborhood of  $\beta \in \mathfrak{k}_\sigma^*$ . The set

$$K_\sigma \cdot (\mathcal{Y}_\sigma^{\tilde{\beta}} \cap \Phi_\sigma^{-1}(\beta)) = M^{\tilde{\beta}} \cap \Phi^{-1}(\beta)$$

is a component of  $\text{Cr}(\|\Phi_\sigma\|^2)$ . Let  $\mathcal{Q}_{K_\sigma}^\beta(\mathcal{Y}_\sigma) \in R_{tc}^{-\infty}(K_\sigma)$  be the corresponding character (see Definition 2.8).

We prove in [19][Section 7], the following induction formula:

$$(2.19) \quad \mathcal{Q}_K^\beta(M) = \text{Hol}_{K_\sigma}^K \left( \mathcal{Q}_{K_\sigma}^\beta(\mathcal{Y}_\sigma) \right)$$

where  $\text{Hol}_{K_\sigma}^K : R^{-\infty}(K_\sigma) \rightarrow R^{-\infty}(K)$  is the holomorphic induction map. See the Appendix in [19] for the definition and properties of these induction maps.

We know from the previous case that

$$\mathcal{Q}_{K_\sigma}^\beta(\mathcal{Y}_\sigma) = \sum_{\mu \in \widehat{K_\sigma}} m_\mu V_\mu^{K_\sigma}$$

where  $m_\mu \neq 0 \implies (\mu, \beta) \geq \|\beta\|^2$ . Then, with (2.19), we get

$$\begin{aligned} \mathcal{Q}_{K_\sigma}^\beta(\mathcal{Y}_\sigma) &= \sum_{(\mu, \beta) \geq \|\beta\|^2} m_\mu \operatorname{Hol}_{K_\sigma}^K(V_\mu^{K_\sigma}) \\ &= \sum_{(\mu, \beta) \geq \|\beta\|^2} m_\mu \operatorname{Hol}_T^K(t^\mu), \end{aligned}$$

where  $\operatorname{Hol}_T^K : R^{-\infty}(T) \rightarrow R^{-\infty}(K)$  is the holomorphic induction map.

Let  $\rho$  be half the sum of the positive roots. The term  $\operatorname{Hol}_T^K(t^\mu)$  is equal to 0 when  $\mu + \rho$  is not a regular element of  $\mathfrak{t}^*$ . When  $\mu + \rho$  is a regular element of  $\mathfrak{t}^*$ , we have  $\operatorname{Hol}_T^K(t^\mu) = (-1)^{|\omega|} V_{\mu_\omega}^K$  where

$$\mu_\omega = \omega(\mu + \rho) - \rho$$

is dominant for a unique  $\omega \in W$ .

Finally, a representation  $V_\lambda^K$  appears in the character  $\mathcal{Q}_K^\beta(M)$  only if  $\lambda = \mu_\omega$  for a weight  $\mu$  satisfying  $(\mu, \beta) \geq \|\beta\|^2$ . Hence, for such  $\lambda$ , we have

$$\begin{aligned} \|\lambda\| &= \|\mu + \rho - \omega^{-1}\rho\| \\ &\geq (\mu + \rho - \omega^{-1}\rho, \frac{\beta}{\|\beta\|}) \\ &\geq \|\beta\|. \end{aligned}$$

In the last inequality we use that  $(\rho - \omega^{-1}\rho, \beta) \geq 0$  since  $\rho - \omega^{-1}\rho$  is a sum of positive roots, and  $\beta \in \mathfrak{t}_+^*$ . □

With the help of Theorem 2.9 and decomposition (2.17), we see that the multiplicity of  $V_\gamma^K$  in  $\mathcal{Q}_K^\Phi(M_{<R})$  does not depend on the regular value  $R > \|\gamma\|^2$ . We can refine the constant  $c_\gamma$  appearing in [15][Theorem 0.1]: take  $c_\gamma$  equal to  $\|\gamma\|^2$  instead of<sup>4</sup>  $\|\gamma + \rho\|^2 - \|\rho\|^2 \geq \|\gamma\|^2$ .

**Definition 2.10.** *The generalized character  $\mathcal{Q}_K^\Phi(M)$  is defined as the limit in  $R^{-\infty}(K)$  of  $\mathcal{Q}_K^\Phi(M_{<R})$  when  $R$  goes to infinity. In other words*

$$(2.20) \quad \mathcal{Q}_K^\Phi(M) = \sum_{\beta \in \mathcal{B}} \mathcal{Q}_K^\beta(M).$$

Note that for any regular value  $R$  of  $\|\Phi\|^2$  we have the useful relation

$$(2.21) \quad \mathcal{Q}_K^\Phi(M) = \mathcal{Q}_K^\Phi(M_{<R}) + O(\sqrt{R}).$$

**2.3. Quantization of a symplectic quotient.** We will now explain how we define the geometric quantization of *singular* compact Hamiltonian manifolds : here “singular” means that the manifold is obtained by symplectic reduction.

Let  $(N, \Omega)$  be a smooth symplectic manifold equipped with a Hamiltonian action of  $K_1 \times K_2$  : we denote  $(\Phi_1, \Phi_2) : N \rightarrow \mathfrak{k}_1^* \times \mathfrak{k}_2^*$  the corresponding moment map. We assume that  $N$  is pre-quantized by a  $K_1 \times K_2$ -equivariant line bundle  $L$  and we

---

<sup>4</sup>Here  $\rho$  is half the sum of the positive roots. Hence  $\|\gamma + \rho\|^2 - \|\rho\|^2 - \|\gamma\|^2 = 2(\rho, \gamma) \geq 0$  and  $(\rho, \gamma) = 0$  only if the weight  $\gamma$  belongs to the center of  $\mathfrak{k} \simeq \mathfrak{k}^*$ .

suppose that the map  $\Phi_1$  is **proper**. One wants to define the geometric quantization of the (compact) symplectic quotient

$$N//_0 K_1 := \Phi_1^{-1}(0)/K_1.$$

Let  $\kappa_1$  be the Kirwan vector field attached to the moment map  $\Phi_1$ . We denote by  $\mathbf{c}^{\kappa_1}$  the symbol  $\text{Thom}(N, J) \otimes L$  pushed by the vector field  $\kappa_1$ . For any regular value  $R_1$  of  $\|\Phi_1\|^2$ , we consider the restriction  $\mathbf{c}^{\kappa_1}|_{N_{<R_1}}$  to the invariant, open subset  $N_{<R_1} := \{\|\Phi_1\|^2 < R_1\}$ . The symbol  $\mathbf{c}^{\kappa_1}|_{N_{<R_1}}$  is  $K_1 \times K_2$ -equivariant and  $K_1$ -transversally elliptic, hence we can consider its index

$$\text{Index}_{N_{<R_1}}^{K_1 \times K_2}(\mathbf{c}^{\kappa_1}|_{N_{<R_1}}) \in R^{-\infty}(K_1 \times K_2).$$

which is smooth relatively to the parameter in  $K_2$ . We consider the following extension of Definition 2.10.

**Definition 2.11.** *The generalized character  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(N)$  is defined as the limit in  $R^{-\infty}(K_1 \times K_2)$  of  $\text{Index}_{N_{<R_1}}^{K_1 \times K_2}(\mathbf{c}^{\kappa_1}|_{N_{<R_1}})$  when  $R_1$  goes to infinity.*

Here  $\text{Cr}(\|\Phi_1\|^2)$  is equal to the disjoint union of the compact  $K_1 \times K_2$ -invariant subsets  $Z_{\beta_1} := K_1 \cdot (M^{\beta_1} \cap \Phi_1^{-1}(\beta_1))$ ,  $\beta_1 \in \mathcal{B}_1$ . For  $\beta_1 \in \mathcal{B}_1$ , we consider an invariant relatively compact open subset  $\mathcal{U}_{\beta_1}$  such that:  $Z_{\beta_1} \subset \mathcal{U}_{\beta_1}$  and  $Z_{\beta_1} = \text{Cr}(\|\Phi_1\|^2) \cap \overline{\mathcal{U}_{\beta_1}}$ . Let  $\mathcal{Q}_{K_1 \times K_2}^{\beta_1}(N) \in R^{-\infty}(K_1 \times K_2)$  be the equivariant index of the  $K_1$ -transversally elliptic symbol  $\mathbf{c}_{\kappa_1}^L|_{\mathcal{U}_{\beta_1}}$ . The  $K_1$ -transversality condition imposes that  $\mathcal{Q}_{K_1 \times K_2}^{\beta_1}(N) = \sum_{\lambda} \theta^{\beta_1}(\lambda) \otimes V_{\lambda}^{K_1}$  with

$$\theta^{\beta_1}(\lambda) \in R(K_2), \quad \forall \lambda \in \widehat{K_1}.$$

We have the following extension of Theorem 2.9

**Theorem 2.12.** *We have  $\mathcal{Q}_{K_1 \times K_2}^{\beta_1}(N) = \sum_{\lambda \in \widehat{K_1}} \theta^{\beta_1}(\lambda) \otimes V_{\lambda}^{K_1}$  where  $\theta^{\beta_1}(\lambda) \neq 0$  only if  $\|\lambda\| \geq \|\beta_1\|$ .*

*Proof.* The proof works exactly like the one of Theorem 2.9.  $\square$

Let us explain the “quantization commutes with reduction theorem”, or why we can consider the geometric quantization of

$$N//_0 K_1 := \Phi_1^{-1}(0)/K_1$$

as the  $K_1$ -invariant part of  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(N)$ .

Let us first suppose that 0 is a regular value of  $\Phi_1$ . Then  $N//_0 K_1$  is a compact symplectic *orbifold* equipped with a Hamiltonian action of  $K_2$ : the corresponding moment map is induced by the restriction of  $\Phi_2$  to  $\Phi_1^{-1}(0)$ . The symplectic quotient  $N//_0 K_1$  is pre-quantized by the line orbibundle

$$L_0 := (L|_{\Phi_1^{-1}(0)})/K_1.$$

Definition 1.1 extends to the orbifold case. We can still define the geometric quantization of  $N//_0 K_1$  as the index of an elliptic operator: we denote it by  $\mathcal{Q}_{K_2}(N//_0 K_1) \in R(K_2)$ . We have

**Theorem 2.13.** *If 0 is a regular value of  $\Phi_1$ , the  $K_1$ -invariant part of  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(N)$  is equal to  $\mathcal{Q}_{K_2}(N//_0 K_1) \in R(K_2)$ .*

Suppose now that 0 is not a regular value of  $\Phi_1$ . Let  $T_1$  be a maximal torus of  $K_1$ , and let  $C_1 \subset \mathfrak{t}_1^*$  be a Weyl chamber. Since  $\Phi_1$  is proper, the convexity Theorem says that the image of  $\Phi_1$  intersects  $C_1$  in a closed locally polyhedral convex set, that we denote  $\Delta_{K_1}(N)$  [14].

We consider an element  $a \in \Delta_{K_1}(N)$  which is generic and sufficiently close to  $0 \in \Delta_{K_1}(N)$  : we denote  $(K_1)_a$  the subgroup of  $K_1$  which stabilizes  $a$ . When  $a \in \Delta_{K_1}(N)$  is generic, one can show (see [18]) that

$$N//_a K_1 := \Phi_{K_1}^{-1}(a)/(K_1)_a$$

is a compact Hamiltonian  $K_2$ -orbifold, and that

$$L_a := \left( L|_{\Phi_{K_1}^{-1}(a)} \right) / (K_1)_a.$$

is a  $K_2$ -equivariant line orbibundle over  $N//_a K_1$  : we can then define, like in Definition 1.1, the element  $\mathcal{Q}_{K_2}(N//_a K_1) \in R(K_2)$  as the equivariant index of the Dolbeault-Dirac operator on  $N//_a K_1$  (with coefficients in  $L_a$ ).

**Theorem 2.14.** *The  $K_1$ -invariant part of  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M)$  is equal to  $\mathcal{Q}_{K_2}(N//_a K_1) \in R(K_2)$ . In particular, the elements  $\mathcal{Q}_{K_2}(N//_a K_1)$  do not depend on the choice of the generic element  $a \in \Delta_H(N)$ , when  $a$  is sufficiently close to 0.*

PROOFS OF THEOREM 2.13 AND THEOREM 2.14 . When  $N$  is compact and  $K_2 = \{e\}$ , the proofs can be found in [18] and in [19]. Let us explain briefly how the  $\mathbf{K}$ -theoretic proof of [19] extends naturally to our case. Like in Definition 2.10, we have the following decomposition

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(N) = \sum_{\beta \in \mathcal{B}_1} \mathcal{Q}_{K_1 \times K_2}^{\beta_1}(N),$$

And Theorem 2.12 tells us that  $\left[ \mathcal{Q}_{K_1 \times K_2}^{\beta_1}(N) \right]^{K_1} = 0$  if  $\beta_1 \neq 0$ . We have proved the first step:

$$\left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(N) \right]^{K_1} = \left[ \mathcal{Q}_{K_1 \times K_2}^0(N) \right]^{K_1}.$$

The analysis of the term  $\left[ \mathcal{Q}_{K_1 \times K_2}^0(N) \right]^{K_1}$  is undertaken in [19] when  $K_2 = \{e\}$ : we explain that this term is equal either to  $\mathcal{Q}(N//_0 K_1)$  when 0 is a regular value, or to  $\mathcal{Q}(N//_a K_1)$  with  $a$  generic. It work similarly with an action of a compact Lie group  $K_2$ .  $\square$

**Definition 2.15.** *The geometric quantization of  $N//_0 K_1 := \Phi_1^{-1}(0)/K_1$  is taken as the  $K_1$ -invariant part of  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(N)$ . We denote it  $\mathcal{Q}_{K_2}(N//_0 K_1)$ .*

**2.4. Quantization of points.** Let  $(M, \Omega, \Phi)$  be a proper Hamiltonian  $K$ -manifold prequantized by a Kostant-Souriau line bundle  $L$ . Let  $\mu \in \hat{K}$  be dominant weight such that  $\Phi^{-1}(K \cdot \mu)$  is a  $K$ -orbit in  $M$ . Let  $m^o \in \Phi^{-1}(\mu)$  so that

$$\Phi^{-1}(K \cdot \mu) = K \cdot m^o$$

Then the reduced space  $M_\mu := \Phi^{-1}(K \cdot \mu)/K$  is a point. The aim of this section is to compute the quantization of  $M_\mu$ :  $\mathcal{Q}(M_\mu) \in \mathbb{Z}$ .

Let  $H$  be the stabilizer subgroup of  $m^o$ . We have a linear action of  $H$  on the 1-dimensional vector space  $L_{m^o} \subset L$ . We have  $H \subset K_\mu$  where  $K_\mu$  is the connected

subgroup of  $K$  that fixes  $\mu \in \mathfrak{t}^*$ . Let  $\mathbb{C}_{-\mu}$  be the 1-dimensional representation of  $K_\mu$  associated to the infinitesimal character  $-i\mu$ .

Let us denote  $\chi$  be the character of  $H$  defined by the 1-dimensional representation  $\mathbb{C}_\chi := L_{m^o} \otimes \mathbb{C}_{-\mu}$ . We know from the Kostant formula (1.1) that  $\chi = 1$  on the identity component  $H^o \subset H$ .

**Theorem 2.16.** *We have*

$$(2.22) \quad \mathcal{Q}(M_\mu) = \begin{cases} 1 & \text{if } \chi = 1 \text{ on } H \\ 0 & \text{in the other case.} \end{cases}$$

This Theorem tells us in particular that  $\mathcal{Q}(M_\mu) = 1$  when the stabiliser subgroup  $H \subset K$  of a point  $m^o \in \Phi^{-1}(\mu)$  is *connected*.

*Proof.* Let  $N = M \times \overline{K \cdot \mu}$  be the proper Hamiltonian  $K$ -manifold which is prequantized by the line bundle  $L_N := L \otimes [\mathbb{C}_{-\mu}]$ . Let us denote  $\Phi_N$  the moment map on  $N$ . Since  $\Phi^{-1}(K \cdot \mu)$  is a  $K$ -orbit in  $M$ , we see that  $\Phi_N^{-1}(0)$  is the  $K$ -orbit through  $n^o := (m^o, \mu)$  where  $m^o \in \Phi^{-1}(\mu)$ . Note that  $H$  is the stabilizer subgroups of  $n^o$ .

Let  $\mathcal{Q}_K^{\Phi_N}(N) \in R^{-\infty}(K)$  be the formal quantization of  $N$  through the proper map  $\Phi_N$ . By definition

$$\begin{aligned} \mathcal{Q}(M_\mu) &= [\mathcal{Q}_K^{\Phi_N}(N)]^K \\ &= [\mathcal{Q}_K^0(N)]^K. \end{aligned}$$

where  $\mathcal{Q}_K^0(N)$  depends only of a neighborhood of  $\Phi_N^{-1}(0)$ .

The orbit  $K \cdot n^o \hookrightarrow N$  is an isotropic embedding since it is the 0-level of the moment map  $\Phi_N$ . Then to describe a  $K$ -invariant neighborhood of  $K \cdot n^o$  in  $N$  we can use the normal-form recipe of Marle, Guillemin and Sternberg.

First we consider, following Weinstein (see [11, 31]), the symplectic normal bundle

$$(2.23) \quad \mathcal{V} := T(K \cdot n^o)^{\perp, \Omega} / T(K \cdot n^o),$$

where the orthogonal  $(^{\perp, \Omega})$  is taken relatively to the symplectic 2-form. We have

$$\mathcal{V} = K \times_H V$$

where the vector space  $V := T_{n^o}(K \cdot n^o)^{\perp, \Omega} / T_{n^o}(K \cdot n^o)$  inherits a symplectic structure and an Hamiltonian action of the group  $H$ : we denote  $\Phi_H : V \rightarrow \mathfrak{h}^*$  the corresponding moment map.

Consider now the following symplectic manifold

$$(2.24) \quad \tilde{N} := \mathcal{V} \oplus T^*(K/H) = K \times_H \left( (\mathfrak{k}/\mathfrak{h})^* \oplus V \right).$$

The action of  $H$  on  $\tilde{N}$  is Hamiltonian and the moment map  $\Phi_{\tilde{N}} : \tilde{N} \rightarrow \mathfrak{k}^*$  is given by the equation

$$(2.25) \quad \Phi_{\tilde{N}}([k; \xi, v]) = k \cdot (\xi + \Phi_H(v)) \quad k \in K, \xi \in (\mathfrak{k}/\mathfrak{h})^*, v \in V.$$

The Hamiltonian  $K$ -manifold  $\tilde{N}$  is prequantized by the line bundle  $L_{\tilde{N}} := K \times_H \mathbb{C}_\chi$ .

The *local normal form* Theorem (see [10], [24] Proposition 2.5 ) tells us that there exists a  $K$ -Hamiltonian isomorphism  $\Upsilon : \mathcal{U}_1 \xrightarrow{\sim} \mathcal{U}_2$  between a  $K$ -invariant neighborhood  $\mathcal{U}_1$  of  $K \cdot n^o$  in  $N$ , and a  $K$ -invariant neighborhood  $\mathcal{U}_2$  of  $K/H$  in  $\tilde{N}$ . This isomorphism  $\Upsilon$ , when restricted to  $K \cdot n^o$ , corresponds to the natural isomorphism  $K \cdot n^o \xrightarrow{\sim} K/H$ .

Thanks to  $\Upsilon$ , we know that the fiber  $\Phi_H^{-1}(0) \subset V$  is reduced to  $\{0\}$ . This last point is equivalent to the fact that  $\Phi_H$  (and then  $\Phi_{\tilde{N}}$ ) is proper map (see [21]). We check easily that the set of critical points of  $\|\Phi_{\tilde{N}}\|^2$  is reduced to  $\Phi_{\tilde{N}}^{-1}(0) = K/H$ . Then, thank to the isomorphism  $\Upsilon$ , we have that

$$(2.26) \quad \mathcal{Q}_K^0(N) = \mathcal{Q}_K^0(\tilde{N}) = \mathcal{Q}_K^{\Phi_{\tilde{N}}}(\tilde{N}).$$

Let  $\text{Ind}_H^K : R^{-\infty}(H) \rightarrow R^{-\infty}(K)$  be the induction map that is defined by the relation  $\langle \text{Ind}_H^K(\phi), E \rangle = \langle \phi, E|_H \rangle$  for any  $\phi \in R^{-\infty}(H)$  and  $E \in R(K)$ . Note that

$$[\text{Ind}_H^K(\phi)]^K = \langle \text{Ind}_H^K(\phi), \mathbb{C} \rangle = \langle \phi, \mathbb{C} \rangle = [\phi]^H.$$

Since  $\Phi_H : V \rightarrow \mathfrak{h}^*$  is proper one can consider the quantization of the vector space  $V$  through the map  $\Phi_H : \mathcal{Q}_H^{\Phi_H}(V) \in R^{-\infty}(H)$ .

**Proposition 2.17.** • *We have*

$$(2.27) \quad \mathcal{Q}_K^{\Phi_{\tilde{N}}}(\tilde{N}) = \text{Ind}_H^K \left( \mathcal{Q}_H^{\Phi_H}(V) \otimes \mathbb{C}_\chi \right)$$

- *The formal quantization  $\mathcal{Q}_H^{\Phi_H}(V)$  coincides, as a generalized  $H$ -module, to the  $H$ -module  $S(V^*)$  of polynomial function on  $V$ .*
- *The set  $[S(V^*)]^{H^o}$  of polynomials invariant by the connected component  $H^o$  is reduced to the scalars.*

With the last Proposition we can finish the proof of Theorem 2.16 as follows. We have

$$\begin{aligned} \mathcal{Q}(M_\mu) &= [\mathcal{Q}_K^{\Phi_{\tilde{N}}}(\tilde{N})]^K \\ &= [\mathcal{Q}_K^{\Phi_{\tilde{N}}}(\tilde{N})]^K \\ &= [\mathcal{Q}_H^{\Phi_H}(V) \otimes \mathbb{C}_\chi]^H \\ &= [S(V^*) \otimes \mathbb{C}_\chi]^H = [\mathbb{C}_\chi]^H. \end{aligned}$$

*Proof.* The first point of Proposition 2.17 follows from the property of induction defined by Atiyah (see Section 3.4 in [19]). Let us explain the arguments. We work with the  $H$ -manifold  $\mathcal{Y} = (\mathfrak{k}/\mathfrak{h})^* \oplus V$  and the  $H$ -equivariant map  $j : \mathcal{Y} \hookrightarrow \tilde{N} := K \times_H \mathcal{Y}, y \mapsto [e, y]$ .

We notice<sup>5</sup> that  $T\tilde{N} \simeq K \times_H (\mathfrak{k}/\mathfrak{h} \oplus T\mathcal{Y})$ , and that  $T_K \tilde{N} \simeq K \times_H (T_H \mathcal{Y})$ . Hence the map  $j$  induces an isomorphism  $j_* : K_H(T_H \mathcal{Y}) \rightarrow K_K(T_K \tilde{N})$ . Theorem 4.1 of

---

<sup>5</sup> These identities come from the following  $K \times H$ -equivariant isomorphism of vector bundles over  $K \times \mathcal{Y}$ :  $T_H(\tilde{N}) \rightarrow K \times (\mathfrak{k}/\mathfrak{h} \oplus T\mathcal{Y}), (k, m; \frac{d}{dt}|_{t=0}(ke^{tX}) + v_m) \mapsto (k, m; pr_{\mathfrak{k}/\mathfrak{h}}(X) + v_m)$ . Here  $pr_{\mathfrak{k}/\mathfrak{h}} : \mathfrak{k} \rightarrow \mathfrak{k}/\mathfrak{h}$  is the orthogonal projection.

Atiyah [1] tells us that the following diagram

$$(2.28) \quad \begin{array}{ccc} K_H(\mathrm{T}_H \mathcal{Y}) & \xrightarrow{j_*} & K_K(\mathrm{T}_K \tilde{N}) \\ \mathrm{Index}_{\mathcal{Y}}^H \downarrow & & \downarrow \mathrm{Index}_{\tilde{N}}^K \\ R^{-\infty}(H) & \xrightarrow{\mathrm{Ind}_H^K} & R^{-\infty}(K) . \end{array}$$

is commutative.

The tangent bundle  $\mathrm{T}\tilde{N}$  is equivariantly diffeomorphic to

$$K \times_H [\mathfrak{k}/\mathfrak{h} \oplus (\mathfrak{k}/\mathfrak{h})^* \oplus \mathrm{TV}] \simeq K \times_H [(\mathfrak{k}/\mathfrak{h})_{\mathbb{C}} \oplus \mathrm{TV}]$$

where  $(\mathfrak{k}/\mathfrak{h})_{\mathbb{C}}$  is the complexification of the real vector space  $\mathfrak{k}/\mathfrak{h}$ . We consider on  $\tilde{N}$  the almost complex structure  $J_{\tilde{N}} = (i, J_V)$  where  $i$  is the complex structure on  $(\mathfrak{k}/\mathfrak{h})_{\mathbb{C}}$  and  $J_V$  is a compatible (constant) complex structure on the symplectic vector space  $V$ . Note that  $J_{\tilde{N}}$  is compatible with the symplectic structure on a neighborhood  $U$  of the 0-section of the bundle  $\tilde{N} \rightarrow K/H$ .

Let  $\kappa_{\tilde{N}}$  be the Kirwan vector field on  $\tilde{N}$ :

$$\kappa_{\tilde{N}}([k; \xi, v]) = -\xi + i[\xi, \Phi_H(v)] \oplus \kappa_V(v) \in (\mathfrak{k}/\mathfrak{h})_{\mathbb{C}} \oplus V.$$

Here  $\kappa_V$  is the Kirwan vector field relative to the Hamiltonian action of  $H$  on the symplectic vector space  $V$ . Note that  $\kappa_{\tilde{N}}$  vanishes exactly on the 0-section of the bundle  $\tilde{N} \rightarrow K/H$ .

Let  $\mathbf{c}^{\kappa_{\tilde{N}}}$  be the symbol  $\mathrm{Thom}(\tilde{N}, J_{\tilde{N}}) \otimes L_{\tilde{N}}$  pushed by the vector field  $\kappa_{\tilde{N}}$ . The generalized character  $\mathcal{Q}_K^{\Phi_{\tilde{N}}}(\tilde{N})$  is either computed as the equivariant index of the symbols  $\mathbf{c}^{\kappa_{\tilde{N}}}$  or  $\mathbf{c}^{\kappa_{\tilde{N}}}|_U$ .

**Remark 2.18.** *The fact that  $J_{\tilde{N}}$  is not compatible on the entire manifold  $\tilde{N}$  is not problematic, since  $J_{\tilde{N}}$  is compatible in a neighborhood  $U$  of the set where  $\kappa_{\tilde{N}}$  vanishes. See the first point of Lemma 2.6.*

For  $X + i\eta \oplus w \in \mathrm{T}_{[k; \xi, v]} \tilde{N} \simeq (\mathfrak{k}/\mathfrak{h})_{\mathbb{C}} \oplus V$ , the map

$$(2.29) \quad \mathbf{c}^{\kappa_{\tilde{N}}}(X + i\eta \oplus w) = \mathbf{c}\left(X + \xi + i(\eta - [\xi, \Phi_H(v)])\right) \odot \mathbf{c}\left(w - \kappa_V(v)\right)$$

acts on the vector space  $\wedge_{\mathbb{C}}(\mathfrak{k}/\mathfrak{h})_{\mathbb{C}} \otimes \wedge_{J_V} V \otimes \mathbb{C}_{\chi}$ .

Let  $\mathrm{Bott}(\mathfrak{k}/\mathfrak{h})$  be the Bott morphism of the vector space  $\mathfrak{k}/\mathfrak{h}$ . It is an elliptic morphism defined by

$$\mathrm{Bott}(\mathfrak{k}/\mathfrak{h})|_{\xi}(\eta) = \mathbf{c}(\xi + i\eta) \text{ acting on } \wedge_{\mathbb{C}}(\mathfrak{k}/\mathfrak{h})_{\mathbb{C}},$$

for  $\eta \in \mathrm{T}_{\xi}(\mathfrak{k}/\mathfrak{h})$ . Let  $\mathbf{c}^{\kappa_V}$  be the symbol  $\mathrm{Thom}(V, J_V)$  pushed by the vector field  $\kappa_V$ .

**Lemma 2.19.** *We have*

$$\mathbf{c}^{\kappa_{\tilde{N}}} = j_* \left( \mathrm{Bott}(\mathfrak{k}/\mathfrak{h}) \odot \mathbf{c}^{\kappa_V} \otimes \mathbb{C}_{\chi} \right).$$

*Proof.* We work with the symbol

$$\sigma^T|_{(\xi, v)}(\eta) = \mathbf{c}(\xi + i\eta - iT[\xi, \Phi_H(v)])$$

acting on  $\wedge_{\mathbb{C}}(\mathfrak{k}/\mathfrak{h})_{\mathbb{C}}$ . Note that  $\mathrm{Bott}(\mathfrak{k}/\mathfrak{h}) = \sigma^0$ . From (2.29), we see that  $\mathbf{c}^{\kappa_{\tilde{N}}} = j_* \left( \sigma^1 \odot \mathbf{c}^{\kappa_V} \otimes \mathbb{C}_{\chi} \right)$ . It is now easy to check that  $\sigma^T \odot \mathbf{c}^{\kappa_V} \otimes \mathbb{C}_{\chi}$ ,  $T \in [0, 1]$  is an homotopy of transversally elliptic symbols on  $\mathfrak{k}/\mathfrak{h} \times V$ .  $\square$

The commutative diagram (2.28) and the last Lemma gives

$$\begin{aligned}
\mathcal{Q}_K^{\Phi_{\tilde{N}}}(\tilde{N}) &= \text{Index}_{\tilde{N}}^K(\mathbf{c}^{\kappa_{\tilde{N}}}) \\
&= \text{Ind}_H^K \left( \text{Index}_{\mathfrak{k}/\mathfrak{h} \times V}^K \left( \text{Bott}(\mathfrak{k}/\mathfrak{h}) \odot \mathbf{c}^{\kappa_V} \right) \otimes \mathbb{C}_\chi \right) \\
&= \text{Ind}_H^K \left( \text{Index}_{\mathfrak{k}/\mathfrak{h}}^K(\text{Bott}(\mathfrak{k}/\mathfrak{h})) \otimes \text{Index}_V^K(\mathbf{c}^{\kappa_V}) \otimes \mathbb{C}_\chi \right) \\
&= \text{Ind}_H^K \left( \mathcal{Q}_H^{\Phi_H}(V) \otimes \mathbb{C}_\chi \right).
\end{aligned}$$

We have used here that the equivariant index of  $\text{Bott}(\mathfrak{k}/\mathfrak{h})$  is equal to 1 (e.g. the trivial representation).

Let us prove now the second point of Proposition 2.17. The Kirwan vector field  $\kappa^V$  satisfies the simple rule:

$$(2.30) \quad (\kappa^V(v), J_V v) = -\Omega(\kappa^V(v), v) = \frac{1}{2} \|\Phi_H(v)\|^2, \quad v \in V.$$

It shows in particular that  $\kappa^V(v) = 0 \Leftrightarrow \Phi_H(v) = 0$ . Since the moment map  $\Phi_H : V \rightarrow \mathfrak{h}^*$  is quadratic, the fact that  $\Phi_H$  is proper is equivalent to the fact that  $\Phi_H^{-1}(0) = 0$ .

We consider on  $V$  the family of symbol  $\sigma^s$  :

$$\sigma^s|_v(w) = \mathbf{c}(w - s\kappa^V(v) - (1-s)J_V v)$$

viewed as a map from  $\wedge_{\mathbb{C}}^{\text{even}} V$  to  $\wedge_{\mathbb{C}}^{\text{odd}} V$ . Thanks to (2.30), one sees that  $\sigma^s$  is a family of  $K$ -transversally elliptic symbol on  $V$ . Hence  $\sigma^1 = \mathbf{c}^{\kappa_V}$  and  $\sigma^0 = \mathbf{c}(w - J_V v)$  defines the same class in the group  $K_K(\text{T}_K V)$ . The symbol  $\sigma^0$  was first studied by Atiyah [1] when  $\dim_{\mathbb{C}} V = 1$ . The author considered the general case in [19]. We have

$$\text{Index}_V^K(\sigma^0) = S(V^*) \quad \text{in } R^{-\infty}(K).$$

The last point of Proposition 2.17 is a consequence of the properness of the moment map  $\Phi_H$  (see Section 5 of [21]).

□

□

**Example 2.20** ([21]). *We consider the action of the unitary group  $U_n$  on  $\mathbb{C}^n$ . The symplectic form on  $\mathbb{C}^n$  is defined by  $\Omega(v, w) = \frac{i}{2} \sum_k v_k \overline{w_k} - \overline{v_k} w_k$ . Let us identify the Lie algebra  $\mathfrak{u}_n$  with its dual through the trace map. The moment map  $\Phi : \mathbb{C}^n \rightarrow \mathfrak{u}_n$  is defined by  $\Phi(v) = \frac{1}{2i} v \otimes v^*$  where  $v \otimes v^* : \mathbb{C}^n \rightarrow \mathbb{C}^n$  is the linear map  $w \mapsto (\sum_k \overline{v_k} w_k) v$ . One checks easily that the pull-back by  $\Phi$  of a  $U_n$ -orbit in  $\mathfrak{u}_n$  is either empty or a  $U_n$ -orbit in  $\mathbb{C}^n$ . We know also that the stabiliser subgroup of a non-zero vector of  $\mathbb{C}^n$  is connected since it is diffeomorphic to  $U_{n-1}$ . Finally we have*

$$(2.31) \quad \mathcal{Q}((\mathbb{C}^n)_\mu) = \begin{cases} 1 & \text{if } \mu \in \widehat{U_n} \text{ belongs to the image of } \Phi \\ 0 & \text{if } \mu \in \widehat{U_n} \text{ does not belong to the image of } \Phi. \end{cases}$$

*Then one checks that  $\mathcal{Q}_{U_n}^{-\infty}(\mathbb{C}^n)$  coincides in  $R^{-\infty}(U_n)$  with the algebra  $S((\mathbb{C}^n)^*)$  of polynomial function on  $\mathbb{C}^n$ .*

**Example 2.21** ([23]). We consider the Lie group  $\mathrm{SL}_2(\mathbb{R})$  and its compact torus of dimension 1 denoted by  $T$ . The Lie algebra  $\mathfrak{sl}_2(\mathbb{R})$  is identified with its dual through the trace map, and the Lie algebra  $\mathfrak{t}$  is naturally identified with  $\mathfrak{sl}_2(\mathbb{R})^T$ . For  $l \in \mathbb{Z} \setminus \{0\}$ , we consider the character  $\chi_l$  of  $T$  defined by

$$\chi_l \left( \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \right) = e^{il\theta}.$$

Its differential  $\frac{1}{i}d\chi_l \in \mathfrak{t}^*$  correspond (through the trace map) to the matrix

$$X_l = \begin{pmatrix} 0 & l/2 \\ -l/2 & 0 \end{pmatrix}.$$

Let  $\mathcal{O}_l$  be the coadjoint orbit of the group  $\mathrm{SL}_2(\mathbb{R})$  through the matrix  $X_l$ . It is a Hamiltonian  $\mathrm{SL}_2(\mathbb{R})$ -manifold prequantized by the  $\mathrm{SL}_2(\mathbb{R})$ -equivariant line bundle  $L_l \simeq \mathrm{SL}_2(\mathbb{R}) \times_T \mathbb{C}_l$ , where  $\mathbb{C}_l$  is the  $T$ -module associated to the character  $\chi_l$ . We look at the Hamiltonian action of  $T$  on  $\mathcal{O}_l$ . Let  $\Phi_T : \mathcal{O}_l \rightarrow \mathfrak{t}^*$  be the corresponding moment map. One checks that the moment map  $\Phi_T$  is proper and that its image is equal to the half-line  $\{aX_l, a \geq 1\} \subset \mathfrak{t}^*$ .

We check that for each  $\xi \in \{aX_l, a \geq 1\}$  the fiber  $\Phi_T^{-1}(\xi)$  is equal to a  $T$ -orbit in  $\mathcal{O}_l$ . For  $k \in \mathbb{Z}$ , let us denote  $(\mathcal{O}_l)_k$  the symplectic reduction of  $\mathcal{O}_l$  at the level  $X_k$ . We know that  $(\mathcal{O}_l)_k = \emptyset$  if  $k \notin \{al, a \geq 1\}$ , and that  $(\mathcal{O}_l)_k$  is a point if  $k \in \{al, a \geq 1\}$ .

In order to compute  $\mathcal{Q}((\mathcal{O}_l)_k)$  we look at the stabilizer subgroup  $T_m := \{t \in T \mid t \cdot m = m\}$  for each point  $m \in \mathcal{O}_l$ . One sees that  $T_m = T$  if  $m = X_l$  and  $T_m$  is equal to the center  $\{\pm Id\}$  of  $\mathrm{SL}_2(\mathbb{R})$ , when  $m \neq X_l$ .

Theorem 2.16 gives in this setting that, for  $k \in \{al, a \geq 1\}$ ,

$$(2.32) \quad \mathcal{Q}((\mathcal{O}_l)_k) = \begin{cases} 1 & \text{if } l - k \text{ is even} \\ 0 & \text{if } l - k \text{ is odd.} \end{cases}$$

Hence the formal geometric quantization of the proper  $T$ -manifold  $\mathcal{O}_l$  is

$$(2.33) \quad \mathcal{Q}_T^{-\infty}(\mathcal{O}_l) = \begin{cases} \mathbb{C}_l \cdot \sum_{p \geq 0} \mathbb{C}_{2p} & \text{if } l > 0 \\ \mathbb{C}_l \cdot \sum_{p \geq 0} \mathbb{C}_{-2p} & \text{if } l < 0. \end{cases}$$

Here we recognize that  $\mathcal{Q}_T^{-\infty}(\mathcal{O}_l)$  coincides with the restriction of the holomorphic (resp. anti-holomorphic) discrete series representation  $\Theta_l$  to the group  $T$  when  $l > 0$  (resp.  $l < 0$ ).

**2.5. Wonderful compactifications and symplectic cuts.** Another equivalent definition of the quantization  $\mathcal{Q}^{-\infty}$  uses a generalisation of the technique of symplectic cutting (originally due to Lerman [13]) that was introduced in [21] and was motivated by the wonderful compactifications of De Concini and Procesi. Let us recall the method.

We recall that  $T$  is a maximal torus in the compact connected Lie group  $K$ , and  $W$  is the Weyl group. We define a  $K$ -adapted polytope in  $\mathfrak{t}^*$  to be a  $W$ -invariant Delzant polytope  $P$  in  $\mathfrak{t}^*$  whose vertices are regular elements of the weight lattice  $\Lambda^*$ . If  $\{\lambda_1, \dots, \lambda_N\}$  are the dominant weights lying in the union of all the closed one-dimensional faces of  $P$ , then there is a  $G \times G$ -equivariant embedding of  $G = K_{\mathbb{C}}$  into

$$\mathbb{P} \left( \bigoplus_{i=1}^N V_{\lambda_i}^* \otimes V_{\lambda_i} \right)$$

associating to  $g \in G$  its representation on  $\bigoplus_{i=1}^N V_{\lambda_i}$ . The closure  $\mathcal{X}_P$  of the image of  $G$  in this projective space is smooth and is equipped with a  $K \times K$  that we denote:

$$(k_1, k_2) \cdot x = k_2 \cdot x \cdot k_1^{-1}.$$

Let  $\Omega_{\mathcal{X}_P}$  be the symplectic 2-form on  $\mathcal{X}_P$  which given by the Kahler structure. We recall briefly the different properties of  $(\mathcal{X}_P, \Omega_{\mathcal{X}_P})$  : all the details can be found in [21].

- (1)  $\mathcal{X}_P$  is equipped with an Hamiltonian action of  $K \times K$ . Let  $\Phi = (\Phi_l, \Phi_r) : M \rightarrow \mathfrak{k}^* \times \mathfrak{k}^*$  be the corresponding moment map.
- (2) The image of  $\Phi := (\Phi_l, \Phi_r)$  is equal to  $\{(k \cdot \xi, -k' \cdot \xi) \mid \xi \in P \text{ and } k, k' \in K\}$ .
- (3) The Hamiltonian manifold  $(\mathcal{X}_P, K \times K)$  has no multiplicities: the pull-back by  $\Phi$  of a  $K \times K$ -orbit in the image is a  $K \times K$ -orbit in  $\mathcal{X}_P$ .

Let  $\mathcal{U}_P := K \cdot P^\circ$  where  $P^\circ$  is the interior of  $P$ . We define

$$\mathcal{X}_P^\circ := \Phi_l^{-1}(\mathcal{U}_P)$$

which is an invariant, open and dense subset of  $\mathcal{X}_P$ . We have the following important property concerning  $\mathcal{X}_P^\circ$ .

- (4) There exists an equivariant diffeomorphism  $\Upsilon : K \times \mathcal{U}_P \rightarrow \mathcal{X}_P^\circ$  such that  $\Upsilon^*(\Phi_l)(k, \xi) = k \cdot \xi$  and  $\Upsilon^*(\Phi_r)(k, \xi) = -\xi$ .
- (5) This diffeomorphism  $\Upsilon$  is a quasi-symplectomorphism in the sense that there is a homotopy of symplectic forms taking the symplectic form on the open subset  $K \times \mathcal{U}_P$  of the cotangent bundle  $T^*K$  to the pullback of the symplectic form  $\Omega_{\mathcal{X}_P}$  on  $\mathcal{X}_P^\circ$ .
- (6) The symplectic manifold  $(\mathcal{X}_P, \Omega_{\mathcal{X}_P})$  is prequantized by the restriction of the hyperplane line bundle  $\mathcal{O}(1) \rightarrow \mathbb{P}(\bigoplus_{i=1}^N V_{\lambda_i}^* \otimes V_{\lambda_i})$  to  $\mathcal{X}_P$ : let us denoted  $L_P$  the corresponding  $K \times K$ -equivariant line bundle.
- (7) The pull-back of the line bundle  $L_P$  by the map  $\Upsilon : K \times \mathcal{U}_P \hookrightarrow \mathcal{X}_P$  is trivial.

Let  $(M, \Omega_M, \Phi_M)$  be a proper Hamiltonian  $K$ -manifold. We also consider the Hamiltonian  $K \times K$ -manifold  $\mathcal{X}_P$  associated to a  $K$ -adapted polytope  $P$ . We consider now the product  $M \times \mathcal{X}_P$  with the following  $K \times K$  action:

- the action  $k \cdot_1 (m, x) = (k \cdot m, x \cdot k^{-1})$  : the corresponding moment map is  $\Phi_1(m, x) = \Phi_M(m) + \Phi_r(x)$ ,
- the action  $k \cdot_2 (m, x) = (m, k \cdot x)$  : the corresponding moment map is  $\Phi_2(m, x) = \Phi_l(x)$ .

**Definition 2.22.** We denote  $M_P$  the symplectic reduction at 0 of  $M \times \mathcal{X}_P$  for the action  $\cdot_1 : M_P := (\Phi_1)^{-1}(0)/(K, \cdot_1)$ .

Then  $M_P$  inherits a Hamiltonian  $K$ -action with moment map  $\Phi_{M_P} : M_P \rightarrow \mathfrak{k}^*$  whose image is  $\Phi(M) \cap K \cdot P$ .

One checks that  $M_P$  contains an open and dense subset of smooth points which quasi-symplectomorphic to the open subset  $(\Phi_M)^{-1}(\mathcal{U}_P)$ . If the polytope  $P$  is fixed, we can work with the dilated polytopes  $nP$  for  $n \geq 1$ . We have then the family of compact, perhaps singular,  $K$ -hamiltonian manifolds  $M_{nP}$ ,  $n \geq 1$ : in Section 2.3, we have explained how was defined their geometric quantization  $\mathcal{Q}_K(M_{nP}) \in R(K)$ .

We have a convenient definition for  $\mathcal{Q}^{-\infty}$ .

**Proposition 2.23** ([21]). *We have the following equality in  $R^{-\infty}(K)$ :*

$$(2.34) \quad \mathcal{Q}_K^{-\infty}(M) = \lim_{n \rightarrow \infty} \mathcal{Q}_K(M_{nP}).$$

### 3. PROOF OF THEOREM 1.4

The main result of this section is

**Theorem 3.1.** *Let  $r_P := \inf_{\xi \in \partial P} \|\xi\|$ . The generalized character*

$$\mathcal{Q}_K^\Phi(M) - \mathcal{Q}_K(M_P) \in R^{-\infty}(K)$$

*is supported outside the ball  $B_{r_P}$ .*

Then, for the dilated polytope  $nP, n \geq 1$ , the character  $\mathcal{Q}_K^\Phi(M) - \mathcal{Q}_K(M_{nP})$  is supported outside the ball  $B_{nr_P}$ . Taking the limit when  $n$  goes to infinity gives

$$(3.35) \quad \mathcal{Q}_K^\Phi(M) = \lim_{n \rightarrow \infty} \mathcal{Q}_K(M_{nP}).$$

Finally, the identity of Theorem 1.4,

$$\mathcal{Q}_K^\Phi(M) = \mathcal{Q}_K^{-\infty}(M),$$

is a direct consequence of (2.34) and (3.35).

Recall that  $O(r) \in R^{-\infty}(K)$  denoted any generalized character supported outside the ball  $B_r$ .

Theorem 3.1 follows from the comparison of three different geometrical situation. All of them concern Hamiltonian actions of  $K_1 \times K_2$ , where  $K_1$  and  $K_2$  are two copies of  $K$ .

**First setting.** We work with the Hamiltonian  $K_1 \times K_2$ -manifold  $M \times \mathcal{X}_P$ : here  $K_1$  acts both on  $M$  and on  $\mathcal{X}_P$ . Since the moment map  $\Phi_1$  (relative to the  $K_1$ -action) is proper we may “quantize”  $M \times \mathcal{X}_P$  via the map  $\|\Phi_1\|^2$ : let

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M \times \mathcal{X}_P) \in R^{-\infty}(K_1 \times K_2)$$

be the corresponding generalized character. Recall that  $\mathcal{Q}_{K_2}(M_P)$  is equal to  $[\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M \times \mathcal{X}_P)]^{K_1}$ .

**Second setting.** We consider the same setting than before : the Hamiltonian action of  $K_1 \times K_2$  on  $M \times \mathcal{X}_P$ . But we “quantize”  $M \times \mathcal{X}_P$  through the global moment map  $\Phi = (\Phi_1, \Phi_2)$ . Here we have some liberty in the choice of the scalar product on  $\mathfrak{k}_1^* \times \mathfrak{k}_2^*$ . If  $\|\xi\|^2$  is an invariant Euclidean norm on  $\mathfrak{k}^*$ , we take on  $\mathfrak{k}_1^* \times \mathfrak{k}_2^*$  the Euclidean norm

$$(3.36) \quad \|(\xi_1, \xi_2)\|_\rho^2 = \|\xi_1\|^2 + \rho \|\xi_2\|^2$$

depending on a parameter  $\rho > 0$ . Let us consider the quantization of  $M \times \mathcal{X}_P$  via the map  $\|\Phi\|_\rho^2$ :

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) \in R^{-\infty}(K_1 \times K_2).$$

**Third setting.** We consider the cotangent bundle  $T^*K$  with the Hamiltonian action of  $K_1 \times K_2$ :  $K_1$  acts by *right* translations, and  $K_2$  by *left* translations. We consider the Hamiltonian action of  $K_1 \times K_2$  on  $M \times T^*K$ : here  $K_1$  acts both on

$M$  and on  $T^*K$ . Let  $\Phi = (\Phi_1, \Phi_2)$  be the global moment map on  $M \times T^*K$ . Since the moment map  $\Phi$  is proper we can “quantize”  $M \times T^*K$  via the map  $\|\Phi\|_\rho^2$ : let

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K) \in R^{-\infty}(K_1 \times K_2)$$

be the corresponding generalized character.

Theorem 3.1 is a consequence of the following propositions.

First we compare  $\mathcal{Q}_{K_2}^\Phi(M)$  with the  $K_1$ -invariant part of  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K)$ .

**Proposition 3.2.** *For any  $\rho \in ]0, 1]$ , we have*

$$(3.37) \quad \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K) \right]^{K_1} = \mathcal{Q}_{K_2}^\Phi(M) \quad \text{in } R^{-\infty}(K_2).$$

Then we compare the  $K_1$ -invariant part of the generalized characters  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K)$  and  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P)$ .

**Proposition 3.3.** *For any  $\rho \in ]0, 1]$ , we have the following relation in  $R^{-\infty}(K_2)$*

$$(3.38) \quad \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) \right]^{K_1} - \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K) \right]^{K_1} = O(r_P)$$

Finally we compare the  $K_1$ -invariant part of the generalized characters  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P)$  and  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M \times \mathcal{X}_P)$ .

**Proposition 3.4.** *There exists  $\epsilon > 0$  such that*

$$(3.39) \quad \mathcal{Q}_{K_2}(M_P) - \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) \right]^{K_1} = O((\epsilon/\rho)^{1/2}) \quad \text{in } R^{-\infty}(K_2)$$

if  $\rho > 0$  is small enough.

If we sum the relations (3.37), (3.38) and (3.39) we get

$$\mathcal{Q}_{K_2}^\Phi(M) = \mathcal{Q}_{K_2}(M_P) + O(r_P) + O((\epsilon/\rho)^{1/2})$$

if  $\rho$  is small enough. So Theorem 3.1 follows by taking  $(\epsilon/\rho)^{1/2} \geq r_P$ .

**3.1. Proof of Proposition 3.2.** The cotangent bundle  $T^*K$  is identified with  $K \times \mathfrak{k}^*$ . The data is then (see Section 5.1):

- the Liouville 1-form  $\lambda = \sum_j \omega_j \otimes E_j$ . Here  $(E_j)$  is a basis of  $\mathfrak{k}$  with dual basis  $(E_j^*)$ , and  $\omega_j$  is the left invariant 1-form on  $K$  defined by  $\omega_j(\frac{d}{dt}a e^{tX}|_0) = \langle E_j^*, X \rangle$ .
- the symplectic form  $\Omega := -d\lambda$ ,
- the action of  $K_1 \times K_2$  on  $K \times \mathfrak{k}^*$  is  $(k_1, k_2) \cdot (a, \xi) = (k_2 a k_1^{-1}, k_1 \cdot \xi)$ ,
- the moment map relative to the  $K_1$ -action is  $\Phi_r(a, \xi) = -\xi$ ,
- the moment map relative to the  $K_2$ -action is  $\Phi_l(a, \xi) = a \cdot \xi$ .

We work now with the Hamiltonian action of  $K_1 \times K_2$  on  $M \times T^*K$  given by

$$(k_1, k_2) \cdot (m, a, \xi) = (k_1 \cdot m, k_2 a k_1^{-1}, k_1 \cdot \xi).$$

The corresponding moment map is  $\Phi = (\Phi_1, \Phi_2)$ :  $\Phi_1(m, a, \xi) = \Phi_M(m) - \xi$  and  $\Phi_2(m, a, \xi) = a \cdot \xi$ .

Let  $\mathbf{c}_1$  be a symbol  $\text{Thom}(M, J_1) \otimes L$  attached to the prequantized Hamiltonian  $K_1$ -manifold  $(M, \Omega)$ . The cotangent bundle  $T^*K$  is prequantized by the trivial line bundle: let  $\mathbf{c}_2$  be the symbol  $\text{Thom}(T^*K, J_2)$  attached to the prequantized

Hamiltonian  $K_1 \times K_2$ -manifold  $T^*K$ . The product  $\mathbf{c} = \mathbf{c}_1 \odot \mathbf{c}_2$  corresponds to the symbol  $\text{Thom}(N, J) \otimes L$  on  $N = M \times T^*K$ .

Let  $\kappa_\rho$  be the Kirwan vector field associated to the map  $\|\Phi\|_\rho^2 : M \times T^*K \rightarrow \mathbb{R}$ . We check that  $\|\Phi\|_\rho^2(m, k, \xi) = \|\Phi_M(m) - \xi\|^2 + \rho\|\xi\|^2$ , and

$$\kappa_\rho(m, k, \xi) = \left( \underbrace{(\Phi_M(m) - \xi) \cdot m}_{\kappa_I}; \underbrace{\tilde{\Phi}_M(m) - (1 + \rho)\xi}_{\kappa_{II, \rho}}; \underbrace{-[\tilde{\Phi}_M(m), \tilde{\xi}]}_{\kappa_{III}} \right).$$

Here  $T_{(m, k, \xi)}(M \times T^*K) \simeq T_m M \times \mathfrak{k} \times \mathfrak{k}$ . We have

$$\begin{aligned} \text{Cr}(\|\Phi\|_\rho^2) &= \{\kappa_\rho = 0\} \\ &= \bigcup_{\beta \in \mathcal{B}} K_1 \times K_2 \cdot \left[ M^{\tilde{\beta}} \cap \Phi_M^{-1}(\beta) \times \{1\} \times \left\{ \frac{\beta}{\rho + 1} \right\} \right] \end{aligned}$$

where  $\mathcal{B}$  parametrizes  $\text{Cr}(\|\Phi_M\|^2)$ . Hence one checks that the critical values of  $\|\Phi\|_\rho^2$  are  $\frac{\rho}{\rho+1}\|\beta\|^2, \beta \in \mathcal{B}$ .

Let  $\mathbf{c}^{\kappa_\rho}$  be the symbol  $\mathbf{c}$  pushed by the vector field  $\kappa_\rho$ : we have

$$\mathbf{c}^{\kappa_\rho}(v; X; Y) = \mathbf{c}_1(v - \kappa_I) \odot \mathbf{c}_2(X - \kappa_{II, \rho}; Y - \kappa_{III})$$

for  $(v; X; Y) \in T_{(m, k, \xi)}(M \times T^*K) \simeq T_m M \times \mathfrak{k} \times \mathfrak{k}$ .

For a real  $R > 0$  we define the open invariant subsets of  $M \times T^*K$

$$\begin{aligned} U_R &:= \{\|\Phi\|_\rho^2 < R\} \\ V_R &:= \{\|\Phi_M\|^2 < R\} \times T^*K. \end{aligned}$$

By definition the generalized index  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K)$  is defined as the limit of the equivariant index

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(U_R) := \text{Index}_{\mathcal{U}_R}^{K_1 \times K_2}(\mathbf{c}^{\kappa_\rho}|_{U_R}),$$

when  $R$  goes to infinity (and stays outside the critical values of  $\|\Phi\|_\rho^2$ ).

In the other hand, when  $R'$  is a regular value of  $\|\Phi_M\|^2$ , we see that the symbol  $\mathbf{c}_\rho|_{V_{R'}}$  is  $K_1 \times K_2$ -transversally elliptic. Let

$$(3.40) \quad \text{Index}_{V_{R'}}^{K_1 \times K_2}(\mathbf{c}^{\kappa_\rho}|_{V_{R'}})$$

be its equivariant index. Notice that the index map is well-defined on  $V_R = \{\|\Phi_M\|^2 < R\} \times T^*K$  since  $T^*K$  can be seen as a open subset of a compact manifold.

It is easy to check that for any  $R > 0$  there exists  $R' > R$  such that  $U_R \subset V_{R'}$ . It implies that  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K)$  is also defined as the limit of (3.40) when  $R'$  goes to infinity.

We look now to the deformation  $\kappa_\rho(s) = (\kappa_I^s; \kappa_{II, \rho}^s; s\kappa_{III})$ ,  $s \in [0, 1]$  where

$$\kappa_I^s(m, \xi) = (\Phi_M(m) - s\xi) \cdot m \quad \text{and} \quad \kappa_{II, \rho}^s(m, \xi) = s\tilde{\Phi}_M(m) - (1 + s\rho)\tilde{\xi}.$$

Let  $\mathbf{c}^{\kappa_\rho(s)}$  be the symbol  $\mathbf{c}$  pushed by the vector field  $\kappa_\rho(s)$ .

**Lemma 3.5.** *Let  $R'$  be a regular value of  $\|\Phi_M\|^2$ .*

- *The family  $\mathbf{c}^{\kappa_\rho(s)}|_{V_{R'}}$ ,  $s \in [0, 1]$  defines an homotopy of  $K_1 \times K_2$ -transversally elliptic symbols on  $V_{R'}$ .*
- *The  $K_1$ -invariant part of  $\text{Index}_{V_{R'}}^{K_1 \times K_2}(\mathbf{c}^{\kappa_\rho(0)}|_{V_{R'}})$  is equal to  $\mathcal{Q}_{K_2}^\Phi(M_{< R'})$ .*

*Proof.* The first point follows from the fact that  $\text{Char}(\mathbf{c}^{\kappa_\rho(s)}|_{V_{R'}}) \cap T_{K_1 \times K_2}(V_{R'})$ , which is equal to

$$\left\{ (m, k, \frac{s}{1+s\rho} \Phi_M(m)), k \in K \text{ and } m \in \text{Cr}(\|\Phi_M\|^2) \cap \{\|\Phi_M\|^2 < R'\} \right\},$$

stays in a compact set when  $s \in [0, 1]$ .

The symbol  $\mathbf{c}^{\kappa_\rho(0)}|_{V_{R'}}$  is equal to the product of the symbol  $\mathbf{c}_1^\kappa|_{M < R'}$ , which is  $K_1$ -transversally elliptic, with the symbol

$$\mathbf{c}_2^\kappa(X; Y) = \mathbf{c}_2(X + \xi; Y)$$

which is a  $K_2$ -transversally elliptic on  $T^*K$ . A basic computation done in section 5.1.2 gives that

$$\begin{aligned} \text{Index}_{T^*K}^{K_1 \times K_2}(\mathbf{c}_2^\kappa) &= L^2(K) \\ &= \sum_{\mu \in \hat{K}} (V_\mu^{K_1})^* \otimes V_\mu^{K_2} \end{aligned}$$

in  $R^{-\infty}(K_1 \times K_2)$ . Finally the “multiplicative property” (see Theorem 2.1) gives

$$\begin{aligned} \text{Index}_{V_{R'}}^{K_1 \times K_2}(\mathbf{c}^{\kappa_\rho(0)}|_{V_{R'}}) &= \text{Index}_{M < R'}^{K_1}(\mathbf{c}_1^\kappa|_{M < R'}) \otimes \text{Index}_{T^*K}^{K_1 \times K_2}(\mathbf{c}_2^\kappa) \\ &= \sum_{\mu \in \hat{K}} \mathcal{Q}_{K_1}^\Phi(M < R') \otimes (V_\mu^{K_1})^* \otimes V_\mu^{K_2} \end{aligned}$$

Taking the  $K_1$ -invariant completes the proof of the second point.  $\square$

Finally we have proved that the generalized character  $[\text{Index}_{V_{R'}}^{K_1 \times K_2}(\mathbf{c}^{\kappa_\rho}|_{V_{R'}})]^{K_1}$  is equal to  $\mathcal{Q}_{K_2}^\Phi(M < R')$ . Taking the limit  $R' \rightarrow \infty$  gives

$$\begin{aligned} \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K) \right]^{K_1} &= \lim_{R' \rightarrow \infty} \left[ \text{Index}_{V_{R'}}^{K_1 \times K_2}(\mathbf{c}^{\kappa_\rho}|_{V_{R'}}) \right]^{K_1} \\ &= \lim_{R' \rightarrow \infty} \mathcal{Q}_{K_2}^\Phi(M < R') = \mathcal{Q}_{K_2}^\Phi(M). \end{aligned}$$

**3.2. Proof of Proposition 3.3.** We work here with the Hamiltonian action of  $K_1 \times K_2$  on  $M \times \mathcal{X}_P$ . The action is  $(k_1, k_2) \cdot (m, x) = (k_1 \cdot m, k_2 \cdot x \cdot k_1^{-1})$  and the corresponding moment map is  $\Phi = (\Phi_1, \Phi_2)$  with  $\Phi_1(m, x) = \Phi_M(m) + \Phi_r(x)$  and  $\Phi_2(m, x) = \Phi_l(x)$ . Let  $\|(\xi_1, \xi_2)\|_\rho^2 = \|\xi_1\|^2 + \rho\|\xi_2\|^2$  be the Euclidean norm  $\mathfrak{k}_1^* \times \mathfrak{k}_2^*$  attached to  $\rho > 0$ .

Let us consider the quantization of  $M \times \mathcal{X}_P$  via the map  $\|\Phi\|_\rho^2$ :

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) \in R^{-\infty}(K_1 \times K_2)$$

The critical set  $\text{Cr}(\|\Phi\|_\rho^2)$  admits the decomposition

$$(3.41) \quad \text{Cr}(\|\Phi\|_\rho^2) = \bigcup_{\gamma \in \mathcal{B}_\rho} K_1 \times K_2 \cdot \mathcal{C}_\gamma$$

where  $(m, x) \in \mathcal{C}_\gamma$  if and only if  $\gamma = (\gamma_1, \gamma_2)$  with

$$(3.42) \quad \begin{cases} \Phi_M(m) + \Phi_r(x) = \gamma_1 \\ \Phi_l(x) = \gamma_2 \\ \tilde{\gamma}_1 \cdot m = 0 \\ \tilde{\gamma}_1 \cdot_r x + \rho \tilde{\gamma}_2 \cdot_l x = 0. \end{cases}$$

We have

$$(3.43) \quad \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) = \sum_{\gamma \in \mathcal{B}_\rho} \mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)$$

where the generalized character  $\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)$  is computed as an index of a transversally elliptic symbol in a neighborhood of

$$K_1 \times K_2 \cdot \mathcal{C}_\gamma \subset M \times \Phi_l^{-1}(K_2 \cdot \gamma_2).$$

Thanks to Theorem 2.9 we know that the support of the generalized character  $\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)$  is contained in  $\{(a, b) \in \widehat{K_1} \times \widehat{K_2} \mid \|a\|^2 + \rho\|b\|^2 \geq \|\gamma\|_\rho^2\}$ . Hence

$$\text{support} \left( [\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)]^{K_1} \right) \subset \left\{ b \in \widehat{K_2} \mid \rho\|b\|^2 \geq \|\gamma\|_\rho^2 \right\}$$

Let  $r_P = \inf_{\xi \in \partial P} \|\xi\|$ . We know then that

$$\left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) \right]^{K_1} = \sum_{\substack{\gamma \in \mathcal{B}_\rho \\ \|\gamma\|_\rho^2 < \rho r_P^2}} [\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)]^{K_1} + O(r_P).$$

Let  $R_P < \rho r_P^2$  be a regular value of  $\|\Phi\|_\rho^2 : M \times \mathcal{X}_P \rightarrow \mathbb{R}$  such that for all  $\gamma \in \mathcal{B}_\rho$  we have  $\|\gamma\|_\rho^2 < \rho r_P^2 \iff \|\gamma\|_\rho^2 < R_P$ . Then

$$(3.44) \quad \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) \right]^{K_1} = \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}((M \times \mathcal{X}_P)_{< R_P}) \right]^{K_1} + O(r_P).$$

For the generalized index  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K)$  we have also a decomposition

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K) = \sum_{\gamma \in \mathcal{B}'_\rho} \mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times T^*K)$$

where  $\mathcal{B}'_\rho$  parametrizes the critical set of  $\|\Phi\|_\rho^2 : M \times T^*K \rightarrow \mathbb{R}$ . Like before we get

$$(3.45) \quad \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K) \right]^{K_1} = \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}((M \times T^*K)_{< R'_P}) \right]^{K_1} + O(r_P).$$

Here  $R'_P < \rho r_P^2$  is a regular value of  $\|\Phi\|_\rho^2 : M \times T^*K \rightarrow \mathbb{R}$  such that for all  $\gamma \in \mathcal{B}'_\rho$  we have  $\|\gamma\|_\rho^2 < \rho r_P^2 \iff \|\gamma\|_\rho^2 < R'_P$ .

**Lemma 3.6.** *We have*

$$(3.46) \quad \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}((M \times \mathcal{X}_P)_{< R_P}) = \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}((M \times T^*K)_{< R'_P}).$$

*Proof.* The Lemma will follow from Proposition 2.6. We take here  $V' = M \times \mathcal{X}_P^o$ ,  $V = M \times K \times \mathcal{U}_P \subset M \times T^*K$  and the equivariant diffeomorphism  $\Psi : V \rightarrow V'$  is equal to  $\text{Id} \times \Upsilon$  where  $\Upsilon$  was introduced in Section 2.5. Note that  $\Psi$  satisfies points (1) – (3) of Proposition 2.6.

Note that  $\|\Phi(m, x)\|_\rho^2 < \rho r_P^2$  implies that  $\|\Phi_l(x)\| < r_P$  and then  $x \in \mathcal{X}_P^o$ . Hence the open subset  $U' := (M \times \mathcal{X}_P)_{< R_P}$  is contained in  $V' = M \times \mathcal{X}_P^o$ . In the same way the open subset  $U := (M \times T^*K)_{< R'_P}$  is contained in  $V$ . We have  $\Psi(U) = U'$  if  $R_P = R'_P$ .

We have proved that (3.46) is a consequence of Proposition 2.6.  $\square$

Finally, if we take the difference between (3.44) and (3.45), we get

$$\left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) \right]^{K_1} - \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times T^*K) \right]^{K_1} = O(r_P).$$

which is the relation of Proposition 3.3.

**3.3. Proof of Proposition 3.4.** Here we want to compare the  $K_1$ -invariant part of the characters  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P)$  and  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M \times \mathcal{X}_P)$ .

We know after Theorem 2.14 that

$$\begin{aligned} \mathcal{Q}_{K_2}(M_P) &= \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M \times \mathcal{X}_P) \right]^{K_1} \\ &= \left[ \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(U_\epsilon) \right]^{K_1} \end{aligned}$$

when  $\epsilon > 0$  is any regular value of  $\|\Phi_1\|^2$ , and  $U_\epsilon := \{\|\Phi_1\|^2 < \epsilon\} \subset M \times \mathcal{X}_P$ .

In this section we fix once for all  $\epsilon > 0$  small enough so that

$$(3.47) \quad \text{Cr}(\|\Phi_1\|^2) \cap \{\|\Phi_1\|^2 \leq \epsilon\} = \{\Phi_1 = 0\}.$$

Let  $\mathbf{c}_1$  be the symbol  $\text{Thom}(M, J_1) \otimes L$  attached to the prequantized Hamiltonian  $K_1$ -manifold  $(M, \Omega)$ . Let  $\mathbf{c}_3$  be the symbol  $\text{Thom}(\mathcal{X}_P, J_3) \otimes L_P$  attached to the prequantized Hamiltonian  $K_1 \times K_2$ -manifold  $\mathcal{X}_P$ . The product  $\mathbf{c} = \mathbf{c}_1 \odot \mathbf{c}_3$  corresponds to the symbol  $\text{Thom}(N, J) \otimes L$  on  $N = M \times \mathcal{X}_P$ .

Let  $\kappa_0$  and  $\kappa_\rho$  be the Kirwan vector fields associated to the functions  $\|\Phi_1\|^2$  and  $\|\Phi\|_\rho^2$  on  $M \times \mathcal{X}_P$ :

$$\kappa_0(m, x) = \left( \underbrace{\Phi_1(m, x) \cdot m}_{\kappa_I}; \underbrace{\Phi_1(m, x) \cdot_\tau x}_{\kappa_{II}} \right), \quad \kappa_\rho(m, x) = \kappa^0(m, x) + \rho(0, \underbrace{\Phi_l(x) \cdot_l x}_{\kappa_{III}}).$$

Let  $\mathbf{c}^{\kappa_\rho}$  be the symbol  $\mathbf{c}$  pushed by the vector field  $\kappa_\rho$ : we have

$$\mathbf{c}^{\kappa_\rho}(v; \eta) = \mathbf{c}_1(v - \kappa_I) \odot \mathbf{c}_3(\eta - \kappa_{II} - \rho \kappa_{III})$$

for  $(v; \eta) \in T_{(m, x)}(M \times \mathcal{X}_P)$ .

The character  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(U_\epsilon)$  is given by the index of the  $K_1$ -transversally elliptic symbol  $\mathbf{c}^{\kappa_0}|_{U_\epsilon}$ . The character  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P)$  is given by the index of the  $K_1 \times K_2$ -transversally elliptic symbol  $\mathbf{c}^{\kappa_\rho}$ .

**Lemma 3.7.** • *There exists  $\rho(\epsilon) > 0$  such that*

$$\text{Cr}(\|\Phi\|_\rho^2) \cap \{\|\Phi_1\|^2 \leq \epsilon\} \subset \left\{ \|\Phi_1\|^2 \leq \frac{\epsilon}{2} \right\}$$

for any  $0 \leq \rho \leq \rho(\epsilon)$ .

*Proof.* With the help of Riemannian metrics on  $M$  and  $\mathcal{X}_P$  we define

$$\begin{aligned} a(\epsilon) &:= \inf_{\epsilon/2 \leq \|\Phi_1(m, x)\| \leq \epsilon} \|\kappa^0(m, x)\| \\ b &:= \sup_{x \in \mathcal{X}_P} \|\Phi_l(x) \cdot_l x\|. \end{aligned}$$

We have  $a(\epsilon) > 0$  thanks to (3.47), and  $b < \infty$  since  $\mathcal{X}_P$  is compact. It is now easy to check that  $\{\kappa_\rho = 0\} \cap \{\epsilon/2 \leq \|\Phi_1\|^2 \leq \epsilon\} = \emptyset$  if  $0 \leq \rho < \frac{a(\epsilon)}{b}$ .  $\square$

The symbols  $\mathbf{c}^{\kappa_\rho}|_{U_\epsilon}$ ,  $\rho \in [0, \rho(c)]$  are  $K_1 \times K_2$ -transversally elliptic, and they define the same class in  $\mathbf{K}_{K_1 \times K_2}(T_{K_1 \times K_2} U_\epsilon)$ . Hence  $\mathcal{Q}_{K_2}(M_P)$  can be computed as the  $K_1$ -invariant part of

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(U_\epsilon) := \text{Index}_{K_1 \times K_2}^{U_\epsilon}(\mathbf{c}^{\kappa_\rho}|_{U_\epsilon}) \in R^{-\infty}(K_1 \times K_2)$$

for  $\rho \in [0, \rho(\epsilon)]$ .

A component  $K_1 \times K_2 \cdot \mathcal{C}_\gamma$  of  $\text{Cr}(\|\Phi\|_\rho^2)$  is contained in  $U_\epsilon$  if and only  $\|\gamma_1\| < \epsilon$  : hence the decomposition (3.43) for the character  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P)$  gives

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P) = \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(\mathcal{U}_\epsilon) + \sum_{\substack{\gamma \in \mathcal{B}_\rho \\ \|\gamma_1\|^2 \geq \epsilon}} \mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P).$$

where

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(\mathcal{U}_\epsilon) = \sum_{\substack{\gamma \in \mathcal{B}_\rho \\ \|\gamma_1\|^2 < \epsilon}} \mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P).$$

Taking the  $K_1$ -invariant gives

$$(3.48) \quad [\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P)]^{K_1} = \mathcal{Q}_{K_2}(M_P) + \sum_{\substack{\gamma \in \mathcal{B}_\rho \\ \|\gamma_1\|^2 \geq \epsilon}} [\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)]^{K_1}.$$

In general we know that the support of the generalized character  $[\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)]^{K_1}$  is included in  $\{b \in \widehat{K_2} \mid \rho\|b\|^2 \geq \|\gamma_1\|^2 + \rho\|\gamma_2\|^2\}$ . When  $\|\gamma_1\|^2 \geq \epsilon$  we have then that the support of  $[\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M \times \mathcal{X}_P)]^{K_1}$  is contained in  $\{b \in \widehat{K_2} \mid \rho\|b\|^2 \geq \epsilon\}$ .

Finally (3.48) imposes that

$$[\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M \times \mathcal{X}_P)]^{K_1} = \mathcal{Q}_{K_2}(M_P) + O((\epsilon/\rho)^{1/2}).$$

when  $0 < \rho \leq \rho(\epsilon)$ , which is the precise content of Proposition 3.4.

#### 4. OTHER PROPERTIES OF $\mathcal{Q}^\Phi$

Let  $(M, \omega, \Phi)$  be a proper Hamiltonian  $K$ -manifold which is prequantized by a line bundle  $L$ . The character  $\mathcal{Q}_K^\Phi(M)$  is computed by means of a scalar product on  $\mathfrak{k}^*$ . The fact that  $\mathcal{Q}_K^\Phi(M) = \mathcal{Q}_K^{-\infty}(M)$  gives the following

**Proposition 4.1.** *The character  $\mathcal{Q}_K^\Phi(M)$  does not depend of the choice of a scalar product on  $\mathfrak{k}^*$*

In this section we work in the setting where  $K = K_1 \times K_2$ . Let  $\Phi_1$  be the moment map relative to the  $K_1$ -action.

**4.1.  $\Phi_1$  is proper.** In this subsection we suppose that the moment map  $\Phi_1$  relative to the  $K_1$ -action is *proper*. We fix an invariant Euclidean norm  $\|\bullet\|^2$  on  $\mathfrak{k}$  in such a way that  $\mathfrak{k}_1 = \mathfrak{k}_2^\perp$ .

Let us “quantize”  $(M, \Omega)$  via the invariant proper function  $\|\Phi_1\|^2$ : let

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M) \in R^{-\infty}(K_1 \times K_2)$$

be the corresponding generalized character.

**Theorem 4.2.** *We have*

$$(4.49) \quad \mathcal{Q}_{K_1 \times K_2}^\Phi(M) = \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M) \quad \text{in} \quad R^{-\infty}(K_1 \times K_2).$$

*Proof.* On  $\mathfrak{k} = \mathfrak{k}_1 \oplus \mathfrak{k}_2$  we may consider the family of invariant Euclidean norms:  $\|X_1 \oplus X_2\|_\rho^2 = \|X_1\|^2 + \rho\|X_2\|^2$  for  $X_j \in \mathfrak{k}_j$ . Let

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M) \in R^{-\infty}(K_1 \times K_2)$$

be the quantization of  $M$  computed via the map  $\|\Phi\|_\rho^2 = \|\Phi_1\|^2 + \rho\|\Phi_2\|^2$ . By definition,  $\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M)$  is equal to  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, 0}(M)$ , and we know after Proposition 4.1 that  $\mathcal{Q}_{K_1 \times K_2}^{\Phi}(M)$  coincides with the generalized character  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M) \in R^{-\infty}(K)$  for any  $\rho > 0$ .

Let us prove that  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M) = \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M)$ . We denote  $O(r) \in R^{-\infty}(K_1 \times K_2)$  any generalized character supported outside the ball

$$\{\xi \in \mathfrak{k}_1^* \times \mathfrak{k}_2^* \mid \|\xi_1\|^2 + \|\xi_2\|^2 < r^2\}.$$

And we denote  $O_1(r) \in R^{-\infty}(K_1 \times K_2)$  any generalized character supported outside the

$$\{\xi \in \mathfrak{k}_1^* \times \mathfrak{k}_2^* \mid \|\xi_1\| < r\}.$$

Let  $R_1 > 0$  be a regular value of  $\|\Phi_1\|^2$ : the open subset  $\{\|\Phi_1\|^2 < R_1\}$  is denoted  $M_{<R_1}$ . We know that

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M) = \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M_{<R_1}) + O_1(\sqrt{R_1}).$$

Like in the Lemma 3.7, we know that

$$(4.50) \quad \text{Cr}(\|\Phi\|_\rho^2) \cap \{\|\Phi_1\|^2 = R_1\} = \emptyset.$$

for  $\rho \geq 0$  small enough. The identity (4.50) first implies that

$$\begin{aligned} \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M) &= \sum_{\substack{\gamma \in \mathcal{B}_\rho \\ \|\gamma_1\|^2 < R_1}} \mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M) + \sum_{\substack{\gamma \in \mathcal{B}_\rho \\ \|\gamma_1\|^2 > R_1}} \mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M) \\ &= \mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M_{<R_1}) + O(\sqrt{R_1}). \end{aligned}$$

In the second equality we have used that  $\mathcal{Q}_{K_1 \times K_2}^{\gamma, \rho}(M) = O(\sqrt{R_1})$  when  $\|\gamma_1\|^2 > R_1$  since the ball  $\{(\xi_1, \xi_2) \in \mathfrak{k}_1^* \times \mathfrak{k}_2^* \mid \|\xi_1\|^2 + \|\xi_2\|^2 < R_1\}$  is contained in

$$\{(\xi_1, \xi_2) \in \mathfrak{k}_1^* \times \mathfrak{k}_2^* \mid \|(\xi_1, \xi_2)\|_\rho^2 < \|(\gamma_1, \gamma_2)\|_\rho^2\}.$$

The identity (4.50) shows also that the symbol  $\mathbf{c}^{\kappa_\rho}|_{M_{<R_1}}$  are homotopic for  $\rho \geq 0$  small enough. Hence

$$\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M_{<R_1}) = \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M_{<R_1})$$

We get finally that  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M) - \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M) = O(\sqrt{R_1}) + O_1(\sqrt{R_1})$  for any regular value  $R_1$  of  $\|\Phi_1\|^2$ . We have proved that  $\mathcal{Q}_{K_1 \times K_2}^{\Phi, \rho}(M) - \mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M) = 0$ .  $\square$

Let us explain how Theorem 4.2 contains the identity that we called “*quantization commutes with reduction in the singular setting*” in [21]. By definition the  $K_1$ -invariant part of the right hand side of (4.49) is equal to the geometric quantization of the (possibly singular) compact Hamiltonian  $K_2$ -manifold

$$M//_0 K_1 := \Phi_1^{-1}(0)/K_1.$$

Using now the fact that the left hand side of (4.49) is equal to  $\mathcal{Q}_{K_1 \times K_2}^{-\infty}(M)$ , we see that the multiplicity of  $V_\mu^{K_2}$  in  $\mathcal{Q}_{K_2}(M//_0 K_1)$  is equal to the geometric quantization of the (possibly singular) compact manifold

$$M \times \overline{K_2 \cdot \mu} //_{(0, \mu)} K_1 \times K_2.$$

**4.2. The symplectic reduction  $M//_0 K_1$  is smooth.** Let  $(M, \Omega)$  be an Hamiltonian  $K_1 \times K_2$ -manifold with a proper moment map  $\Phi = (\Phi_1, \Phi_2)$ . In this section we suppose that 0 is a regular value of  $\Phi_1$  and that  $K_1$  acts freely on  $\Phi_1^{-1}(0)$ . We work then with the (smooth) Hamiltonian  $K_2$ -manifold

$$N := \Phi_1^{-1}(0)/K_1.$$

We still denote by  $\Phi_2 : N \rightarrow \mathfrak{k}_2^*$  the moment map relative to the  $K_2$ -action: note that this map is proper. Hence we can quantize the  $K_2$ -action on  $N$  via the map  $\Phi_2$ . Let  $\mathcal{Q}_{K_2}^{\Phi_2}(N) \in R^{-\infty}(K_2)$  be the corresponding character.

**Proposition 4.3.** *We have*

$$(4.51) \quad [\mathcal{Q}_{K_1 \times K_2}^{\Phi}(M)]^{K_1} = \mathcal{Q}_{K_2}^{\Phi_2}(N) \quad \text{in } R^{-\infty}(K_2).$$

*Proof.* When  $\Phi_1$  is proper, the manifold  $N$  is compact. Then the right hand side of (4.51) is equal to  $\mathcal{Q}_{K_2}(N)$ , and we know from Theorem 4.2 that the left hand side of (4.51) is equal to  $[\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M)]^{K_1}$ . In this case (4.51) becomes  $[\mathcal{Q}_{K_1 \times K_2}^{\Phi_1}(M)]^{K_1} = \mathcal{Q}_{K_2}(M//_0 K_1)$  which is the content of Theorem 2.13.

Let us consider the general case where  $\Phi_1$  is not proper. Thanks to Theorem 1.4 one knows that the multiplicities of  $V_\mu^{K_2}$  in  $[\mathcal{Q}_{K_1 \times K_2}^{\Phi}(M)]^{K_1}$  and  $\mathcal{Q}_{K_2}^{\Phi_2}(N)$  are respectively equal to the quantization of the (possibly singular) symplectic reductions

$$\mathcal{M}_\mu := M \times \overline{K_2 \cdot \mu} //_{(0, \mu)} K_1 \times K_2.$$

and

$$\mathcal{M}'_\mu := N \times \overline{K_2 \cdot \mu} //_0 K_2, \quad \text{with } N = M//_0 K_1.$$

Note that  $\mathcal{M}_\mu$  and  $\mathcal{M}'_\mu$  coincide as symplectic reduced space. Let us prove that their geometric quantization are identical also. The proof will be done for  $\mu = 0$ : the other case follows from the shifting trick.

Let  $\mathbf{c}$  be the  $K_1 \times K_2$ -equivariant symbol  $\text{Thom}(M, J) \otimes L_M$ . Let  $\kappa$  be the Kirwan vector field attached to the moment map  $\Phi = (\Phi_1, \Phi_2)$ . Let  $\mathbf{c}^\kappa$  be the symbol  $\mathbf{c}$  pushed by  $\kappa$ . Let us denote  $M_{<\epsilon}$  the open subset  $\{\|\Phi\|^2 < \epsilon\}$ . For  $\epsilon > 0$  small enough, the symbol  $\mathbf{c}^\kappa|_{M_{<\epsilon}}$  is  $K_1 \times K_2$ -transversally elliptic, and  $\mathcal{Q}(\mathcal{M}_0)$  is the  $K_1 \times K_2$ -invariant part of  $\text{Index}_{M_{<\epsilon}}^{K_1 \times K_2}(\mathbf{c}^\kappa|_{M_{<\epsilon}})$ .

Let  $\mathbf{c}_2$  be the  $K_2$ -equivariant symbol  $\text{Thom}(N, J) \otimes L_N$ . Let  $\kappa_2$  be the Kirwan vector field attached to the moment map  $\Phi_2$ . Let  $\mathbf{c}_2^{\kappa_2}$  be the symbol  $\mathbf{c}_2$  pushed by  $\kappa_2$ . Let us denote  $N_{<\epsilon}$  the open subset  $\{\|\Phi_2\|^2 < \epsilon\}$ . For  $\epsilon > 0$  small enough, the symbol  $\mathbf{c}_2^{\kappa_2}|_{N_{<\epsilon}}$  is  $K_2$ -transversally elliptic, and  $\mathcal{Q}(\mathcal{M}'_0)$  is the  $K_2$ -invariant part of  $\text{Index}_{N_{<\epsilon}}^{K_2}(\mathbf{c}_2^{\kappa_2}|_{N_{<\epsilon}})$ .

Our proof follows from the comparison of the classes

$$[\mathbf{c}^\kappa|_{M_{<\epsilon}}] \in \mathbf{K}_{K_1 \times K_2}(\text{T}_{K_1 \times K_2} M_{<\epsilon})$$

and

$$[\mathbf{c}_2^{\kappa_2}|_{N_{<\epsilon}}] \in \mathbf{K}_{K_2}(\text{T}_{K_2} N_{<\epsilon})$$

A neighborhood of the smooth submanifold  $Z := \Phi_1^{-1}(0)$  in  $M$  is diffeomorphic to a neighborhood of the 0-section of the bundle  $Z \times \mathfrak{k}_1^* \rightarrow Z$ . Let  $Z_{<\epsilon} = Z \cap M_{<\epsilon}$  so that  $N_{<\epsilon} = Z_{<\epsilon}/K_1$ . Hence  $[\mathbf{c}^\kappa|_{M_{<\epsilon}}]$  can be seen naturally a class in the K-group  $\mathbf{K}_{K_1 \times K_2}(\mathbf{T}_{K_1 \times K_2}(Z_{<\epsilon} \times \mathfrak{k}_1^*))$ .

Following Atiyah [1][Theorem 4.3], the inclusion map  $j : Z_{<\epsilon} \hookrightarrow Z_{<\epsilon} \times \mathfrak{k}_1^*$  induces the Thom isomorphism

$$j! : \mathbf{K}_{K_1 \times K_2}(\mathbf{T}_{K_1 \times K_2} Z_{<\epsilon}) \longrightarrow \mathbf{K}_{K_1 \times K_2}(\mathbf{T}_{K_1 \times K_2}(Z_{<\epsilon} \times \mathfrak{k}_1^*)),$$

with the commutative diagram

$$(4.52) \quad \begin{array}{ccc} \mathbf{K}_{K_1 \times K_2}(\mathbf{T}_{K_1 \times K_2} Z_{<\epsilon}) & \xrightarrow{j!} & \mathbf{K}_{K_1 \times K_2}(\mathbf{T}_{K_1 \times K_2}(Z_{<\epsilon} \times \mathfrak{k}_1^*)) \\ & \searrow \text{Index}_{Z_{<\epsilon}}^{K_1 \times K_2} & \downarrow \text{Index}_{Z_{<\epsilon} \times \mathfrak{k}_1^*}^{K_1 \times K_2} \\ & & R^{-\infty}(K_1 \times K_2) \end{array}$$

Let  $\pi_1 : Z_{<\epsilon} \rightarrow N_{<\epsilon}$  be the quotient relative to the free action of  $K_1$ . The corresponding isomorphism

$$\pi_1^* : \mathbf{K}_{K_2}(\mathbf{T}_{K_2} N_{<\epsilon}) \longrightarrow \mathbf{K}_{K_1 \times K_2}(\mathbf{T}_{K_1 \times K_2} Z_{<\epsilon})$$

satisfies the following rule :

$$(4.53) \quad \left[ \text{Index}_{K_1 \times K_2}^{Z_{<\epsilon}}(\pi_1^* \theta) \right]^{K_1} = \text{Index}_{K_2}^{N_{<\epsilon}}(\theta)$$

for any  $\theta \in \mathbf{K}_{K_2}(\mathbf{T}_{K_2} N_{<\epsilon})$ .

**Lemma 4.4** ([19]). *We have*

$$j! \circ \pi_1^* \left( [\mathbf{c}_2^{\kappa_2}|_{N_{<\epsilon'}}] \right) = [\mathbf{c}^\kappa|_{M_{<\epsilon}}]$$

in  $\mathbf{K}_{K_1 \times K_2}(\mathbf{T}_{K_1 \times K_2}(Z_{<\epsilon} \times \mathfrak{k}_1^*))$ .

*Proof.* This Lemma is proven in [19][Section 6.2] when the group  $K_2$  is trivial. It is easy to check that the proof extends naturally to our setting.  $\square$

If one uses Lemma 4.4 together with (4.52) and (4.53), we get that

$$\begin{aligned} \mathcal{Q}(\mathcal{M}_0) &= \left[ \text{Index}_{Z_{<\epsilon} \times \mathfrak{k}_1^*}^{K_1 \times K_2}(\mathbf{c}^\kappa|_{M_{<\epsilon}}) \right]^{K_1 \times K_2} \\ &= \left[ \text{Index}_{N_{<\epsilon}}^{K_2}(\mathbf{c}_2^{\kappa_2}|_{N_{<\epsilon}}) \right]^{K_2} = \mathcal{Q}(\mathcal{M}'_0). \end{aligned}$$

$\square$

## 5. EXAMPLE: THE COTANGENT BUNDLE OF AN ORBIT

**5.1. The formal quantization of  $\mathbf{T}^*K$ .** Let  $K$  be a compact connected Lie group equipped with the action of two copies of  $K$ :  $(k_1, k_2) \cdot a = k_2 a k_1^{-1}$ . Then we have a Hamiltonian action of  $K_1 \times K_2$  on the cotangent bundle  $\mathbf{T}^*K$ . In this section, we check that each formal geometric quantization of  $\mathbf{T}^*K$ ,  $\mathcal{Q}_{K_1 \times K_2}^{-\infty}(\mathbf{T}^*K)$  and  $\mathcal{Q}_{K_1 \times K_2}^\Phi(\mathbf{T}^*K)$ , are both equal to the  $K_1 \times K_2$ -module  $L^2(K)$ .

The tangent bundle  $TK$  is identified with  $K \times \mathfrak{k}$  through the right translations: to  $(a, X) \in K \times \mathfrak{k}$  we associate  $\frac{d}{dt}ae^{tX}|_0$ . The action of  $K_1 \times K_2$  on the cotangent bundle  $T^*K \simeq K \times \mathfrak{k}^*$  is then

$$(k_1, k_2) \cdot (a, \xi) = (k_2 a k_1^{-1}, k_1 \cdot \xi).$$

The symplectic form on  $T^*K$  is  $\Omega := -d\lambda$ , where  $\lambda$  is the Liouville 1-form. Let us compute these two form in coordinates. The tangent bundle of  $T^*K \simeq K \times \mathfrak{k}^*$  is identified with  $T^*K \times \mathfrak{k} \times \mathfrak{k}^*$ : for each  $(a, \xi) \in T^*K$ , we have a two form  $\Omega_{(a, \xi)}$  on  $\mathfrak{k} \times \mathfrak{k}^*$ . A direct computation gives

$$\Omega_{(a, \xi)}(X, Y) = \langle \xi, [X, Y] \rangle, \quad \Omega_{(a, \xi)}(\eta, \eta') = 0, \quad \Omega_{(a, \xi)}(X, \eta) = \langle \eta, X \rangle$$

for  $X, Y \in \mathfrak{k}$  and  $\eta, \eta' \in \mathfrak{k}^*$ . So  $\Omega_{(a, \xi)} = \Omega_0 + \pi_\xi$  where  $\Omega_0$  is the canonical (constant) symplectic form on  $\mathfrak{k} \times \mathfrak{k}^*$  and  $\pi_\xi$  is the closed two form on  $\mathfrak{k}$  defined by  $\pi_\xi(X, Y) = \langle \xi, [X, Y] \rangle$ .

If we identify  $\mathfrak{k} \simeq \mathfrak{k}^*$  through an invariant Euclidean norm, the symplectic structure on  $T_{(a, \xi)}(T^*K) \simeq \mathfrak{k} \times \mathfrak{k}^*$  is given by a skew-symmetric matrix

$$A_\xi := \begin{pmatrix} \text{ad}(\xi) & I_n \\ -I_n & 0 \end{pmatrix}.$$

We will work with the following compatible almost complex structure on the tangent bundle of  $T^*K$ :  $J_\xi = -A_\xi(-A_\xi^2)^{-1/2}$ . When  $\xi = 0$ , the complex structure  $J_0$  on  $\mathfrak{k} \times \mathfrak{k}^*$  is defined by the matrix

$$J_0 := \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}.$$

Hence the complex  $K$ -module  $(\mathfrak{k} \times \mathfrak{k}^*, J_0)$  is naturally identified with the complexification  $\mathfrak{k}_\mathbb{C}$  of  $\mathfrak{k}$ .

One checks easily that the moment map relative to the  $K_1 \times K_2$ -action is the *proper* map  $\Phi : T^*K \rightarrow \mathfrak{k}_1^* \times \mathfrak{k}_2^*$  defined by  $\Phi(a, \xi) = (-\xi, a \cdot \xi)$ .

Here the symplectic manifold  $T^*K$  is prequantized by the trivial line bundle.

5.1.1. *Computation of  $\mathcal{Q}_{K_1 \times K_2}^{-\infty}(T^*K)$ .* Let  $\mathcal{O}_1 \times \mathcal{O}_2$  be a coadjoint orbit of  $K_1 \times K_2$  in  $\mathfrak{k}_1^* \times \mathfrak{k}_2^*$ . One checks that

$$(5.54) \quad \Phi^{-1}(\mathcal{O}_1 \times \mathcal{O}_2) = \begin{cases} \emptyset & \text{if } \mathcal{O}_1 \neq -\mathcal{O}_2 \\ \text{a } K_1 \times K_2 \text{-orbit} & \text{if } \mathcal{O}_1 = -\mathcal{O}_2. \end{cases}$$

We know that the stabiliser subgroup  $K_\xi$  of an element  $\xi \in \mathfrak{k}^*$  is connected. Then the stabilizer subgroup  $(K_1 \times K_2)_{(a, \xi)} = \{(k_1, a k_1 a^{-1}), k_1 \in K_\xi\}$  is also connected.

Let  $(T^*K)_{(\mu, \lambda)}$  be the symplectic reduction of  $T^*K$  at the level  $(\mu, \lambda) \in \widehat{K}^2$ . For any  $\mu \in \widehat{K}$ , we define  $\mu^* \in \widehat{K}$  by the relation  $-K \cdot \mu = K \cdot \mu^*$ : note that  $V_{\mu^*}^K \simeq (V_\mu^K)^*$ . If one uses Theorem 2.16, one has

$$(5.55) \quad \mathcal{Q}((T^*K)_{(\mu, \lambda)}) = \begin{cases} 0 & \text{if } \lambda \neq \mu^* \\ 1 & \text{if } \lambda = \mu^*. \end{cases}$$

Finally

$$\begin{aligned}\mathcal{Q}_{K_1 \times K_2}^{-\infty}(\mathrm{T}^*K) &= \sum_{(\mu, \lambda) \in \widehat{K} \times \widehat{K}} \mathcal{Q}((\mathrm{T}^*K)_{(\mu, \lambda)}) V_{\mu}^{K_1} \otimes V_{\lambda}^{K_2} \\ &= \sum_{\mu \in \widehat{K}} V_{\mu}^{K_1} \otimes (V_{\mu}^{K_2})^* = L^2(K).\end{aligned}$$

5.1.2. *Computation of  $\mathcal{Q}_{K_1 \times K_2}^{\Phi}(\mathrm{T}^*K)$ .* The Kirwan vector field on  $\mathrm{T}^*K$  is

$$\kappa(a, \xi) = -2\xi \in \mathfrak{k}_{\mathbb{C}}.$$

Let  $\mathbf{c}^{\kappa}$  be the symbol  $\mathrm{Thom}(\mathrm{T}^*K, J)$  pushed by the vector field  $\frac{1}{2}\kappa$ . At each  $(a, \xi) \in \mathrm{T}^*K$ , the map  $\mathbf{c}_{(a, \xi)}^{\kappa}(X \oplus \eta)$  from  $\wedge_{J_{\xi}}^{\mathrm{even}}(\mathfrak{k} \times \mathfrak{k}^*)$  to  $\wedge_{J_{\xi}}^{\mathrm{odd}}(\mathfrak{k} \times \mathfrak{k}^*)$  is equal to the clifford map  $\mathbf{c}(X + \xi \oplus \eta)$ . Note that  $\mathbf{c}^{\kappa}$  is a  $K_2$ -transversally elliptic symbol on  $\mathrm{T}^*K$ : we have  $\mathrm{Char}(\mathbf{c}^{\kappa}) \cap \mathrm{T}_{K_2}(\mathrm{T}^*K) = \{(1, 0)\}$ . We will now compute the equivariant index of  $\mathbf{c}^{\kappa}$ .

First we consider the homotopy  $t \in [0, 1] \rightarrow J_{t\xi}$  of symplectic structure on  $\mathrm{T}^*K$ . Let  $\tilde{\mathbf{c}}^{\kappa}$  be the symbol acting on  $\wedge_{J_0}^{\bullet}(\mathfrak{k} \times \mathfrak{k}^*) = \wedge_{\mathbb{C}}^{\bullet} \mathfrak{k}_{\mathbb{C}}$ . Proposition 2.6 shows that the symbols  $\mathbf{c}^{\kappa}$  and  $\tilde{\mathbf{c}}^{\kappa}$  define the same class in  $K_{K_1 \times K_2}(\mathrm{T}_{K_2}(\mathrm{T}^*K))$ .

The projection  $\pi : \mathrm{T}^*K \rightarrow \mathfrak{k}^*$  corresponds to the quotient map relative to the free action of  $K_2$ . At the level of  $K$ -groups we get an isomorphism

$$\pi_* : K_{K_1 \times K_2}(\mathrm{T}_{K_2}(\mathrm{T}^*K)) \rightarrow K_{K_1}(\mathrm{T}\mathfrak{k}^*).$$

Atiyah [1] proves that

$$\mathrm{Index}_{K_1 \times K_2}^{\mathrm{T}^*K}(\sigma) = \sum_{\mu \in \widehat{K}} \mathrm{Index}_{K_1}^{\mathfrak{k}^*}(\pi_*(\sigma \otimes V_{\mu}^{K_2})) \otimes (V_{\mu}^{K_2})^*$$

for any class  $\sigma \in K_{K_1 \times K_2}(\mathrm{T}_{K_2}(\mathrm{T}^*K))$ . In our case the symbol  $\pi^*(\tilde{\mathbf{c}}^{\kappa})$  is equal to the Bott symbol  $\mathrm{Bott}(\mathfrak{k}^*)$ , and for any  $K_2$ -module  $E_2$  we have

$$\pi_*(\tilde{\mathbf{c}}^{\kappa} \otimes E_2) = \mathrm{Bott}(\mathfrak{k}^*) \otimes E_1$$

where  $E_1$  is the module  $E_2$  with the action of  $K_1$ . Then

$$\begin{aligned}\mathcal{Q}_{K_1 \times K_2}^{\Phi}(\mathrm{T}^*K) &= \mathrm{Index}_{K_1 \times K_2}^{\mathrm{T}^*K}(\tilde{\mathbf{c}}^{\kappa}) \\ &= \sum_{\mu \in \widehat{K}} \mathrm{Index}_{K_1}^{\mathfrak{k}^*}(\mathrm{Bott}(\mathfrak{k}^*) \otimes V_{\mu}^{K_1}) \otimes (V_{\mu}^{K_2})^* \\ &= \sum_{\mu \in \widehat{K}} V_{\mu}^{K_1} \otimes (V_{\mu}^{K_2})^* = L^2(K),\end{aligned}$$

since  $\mathrm{Index}_{K_1}^{\mathfrak{k}^*}(\mathrm{Bott}(\mathfrak{k}^*)) = 1$ .

5.2. **The formal quantization of  $\mathrm{T}^*(K/H)$ .** Let  $H$  be a closed connected subgroup of  $K$ . Look at  $\mathrm{T}^*K$  as a Hamiltonian manifold relatively to the action of  $H \times K \subset K_1 \times K_2$ . The moment map  $\Phi = (\Phi_H, \Phi_K)$  is defined by :  $\Phi_H(a, \xi) = -\mathrm{pr}(\xi)$  and  $\Phi_K(a, \xi) = a \cdot \xi$ , where  $\mathrm{pr} : \mathfrak{k}^* \rightarrow \mathfrak{h}^*$  is the projection. Note that  $\Phi$  is a proper map.

The cotangent bundle  $\mathrm{T}^*(K/H)$ , viewed as  $K$ -manifold, is equal to the symplectic reduction of  $\mathrm{T}^*K$  relatively to the  $H$ -action: if the kernel of the projection  $\mathrm{pr}$  is denoted  $\mathfrak{h}^{\perp}$ , we have

$$\Phi_H^{-1}(0)/H = K \times_H \mathfrak{h}^{\perp} = \mathrm{T}^*(K/H).$$

We are here in the setting of section 4.2. The reduction of the  $H \times K$  proper Hamiltonian manifold  $T^*K$  relatively to the  $H$ -action is smooth, then its formal quantization is computed as follows

$$\begin{aligned}
 \mathcal{Q}_K^\Phi(T^*(K/H)) &= [\mathcal{Q}_{H \times K}^\Phi(T^*K)]^H = [\mathcal{Q}_{K_1 \times K_2}^\Phi(T^*K)|_{H \times K}]^H \\
 (5.56) \qquad \qquad \qquad &= [L^2(K)]^H \\
 &= L^2(K/H).
 \end{aligned}$$

Here the fact that  $\mathcal{Q}_{H \times K}^\Phi(T^*K)$  is equal to the restriction of  $\mathcal{Q}_{K_1 \times K_2}^\Phi(T^*K) = L^2(K)$  to  $H \times K$  is a consequence of Theorem 1.3.

Let us denote  $\left[T^*(K/H)\right]_\mu$  the symplectic reduction at  $\mu \in \widehat{K}$  of the  $K$ -Hamiltonian manifold  $T^*(K/H)$ . Theorem 1.4 together with (1.6) gives

**Corollary 5.1.** *For any  $\mu \in \widehat{K}$ , we have*

$$\mathcal{Q}\left(\left[T^*(K/H)\right]_\mu\right) = \dim[V_\mu^K]^H,$$

where  $[V_\mu^K]^H$  is the subspace of  $H$ -invariant vector.

**5.3. The formal quantization of  $T^*(K/H)$  relatively to the action of  $G$ .** Let  $G$  be a closed connected subgroup of  $K$ . We look at the hamiltonian action of  $G$  on  $T^*(K/H)$ . Let  $\Phi_G : T^*(K/H) \rightarrow \mathfrak{g}^*$  be the moment map. We consider also the restriction of the  $K$ -module  $L^2(K/H)$  to  $G$ .

We have

**Proposition 5.2.** *The following statements are equivalent*

- (1) *The moment map  $\Phi_G : T^*(K/H) \rightarrow \mathfrak{g}^*$  is proper.*
- (2)  *$\Phi_G^{-1}(0)$  is equal to the zero section.*
- (3)  *$k \cdot \mathfrak{g} + \mathfrak{h} = \mathfrak{k}$ , for any  $k \in K$ .*
- (4)  *$\mathfrak{g} + \mathfrak{h} = \mathfrak{k}$*
- (5)  *$G$  acts transitively on  $K/H$ .*
- (6)  *$L^2(K/H)]^G \simeq \mathbb{C}$*
- (7)  *$L^2(K/H)|_G$  is an admissible  $G$ -representation.*

*Proof.* (1)  $\implies$  (7) is a consequence of Theorem 1.3. Let us prove that (7)  $\implies$  (6). Suppose now that

$$L^2(K/H)|_G = \sum_{\mu \in \widehat{K}} [V_\mu^K]^H \otimes (V_\mu^K)^*|_G$$

is an admissible  $G$ -representation. It means that for any  $\lambda \in \widehat{G}$  the set

$$A_\lambda := \left\{ \mu \in \widehat{K} \mid [V_\mu^K]^H \neq \{0\} \text{ and } [(V_\lambda^G)^* \otimes (V_\mu^K)^*|_G]^G \neq \{0\} \right\}$$

is finite. Then the vector space  $L^2(K/H)]^G$  is equal to the finite dimensional vector space  $\sum_{\mu \in A_0} [V_\mu^K]^H \otimes [(V_\mu^K)^*]^G$ . It is not difficult to check that if  $\mu \in A_0$ , then  $k\mu \in A_0$  for  $k \gg 1$ . Finally the fact that  $A_0$  is finite implies that  $A_0$  is reduced to  $\mu = 0$ . Hence the only  $G$ -invariant functions on  $K/H$  are the scalars.

(6)  $\iff$  (5)  $\iff$  (4)  $\iff$  (3) is a general fact concerning smooth actions of a compact connected Lie group  $G$  on a compact connected manifold  $M$ . The manifold  $M$  does not have  $G$ -invariant functions which are not scalar if and only if the action of  $G$  on  $M$  is transitive. And given a point  $m \in M$ , the orbit  $G \cdot m$  is all of  $M$  if

and only if tangent spaces  $T_m(G \cdot m)$  and  $T_m M$  are equal. If we take  $m = \overline{k^{-1}}$  in  $M = K/H$ , the condition  $T_m(G \cdot m) = T_m M$  is equivalent to  $k \cdot \mathfrak{g} + \mathfrak{h} = \mathfrak{k}$ .

Let us check (3)  $\implies$  (2). Let  $[k, \xi] \in K \times_H \mathfrak{h}^\perp = T^*(K/H)$ . We have  $\Phi_G([k, \xi]) = 0$  if and only if  $k \cdot \xi \in \mathfrak{g}^\perp$ . Hence the vector  $\xi$  belongs to

$$k^{-1} \cdot \mathfrak{g}^\perp \bigcap \mathfrak{h}^\perp = (k^{-1} \cdot \mathfrak{g} + \mathfrak{h})^\perp.$$

Hence condition (3) imposes that  $\xi = 0$ .

(2)  $\iff$  (1) comes from the fact that  $\Phi_G$  is a homogeneous map of degree one between the vector bundle  $T^*(K/H)$  and the vector space  $\mathfrak{g}^*$ .  $\square$

Suppose now that the cotangent bundle  $T^*(K/H)$  is a *proper* Hamiltonian  $G$ -manifold. Let us denote  $[T^*(K/H)]_{\mu, G}$  the (compact) symplectic reduction at  $\mu \in \widehat{G}$  of the  $G$ -Hamiltonian manifold  $T^*(K/H)$ . Then,

**Corollary 5.3.** *The multiplicity of  $V_\mu^G$  in  $L^2(K/H)$  is equal to the quantization of the reduced space  $[T^*(K/H)]_{\mu, G}$ .*

*Proof.* Using Theorem 1.3, equality (5.56) gives then

$$\begin{aligned} \mathcal{Q}_G^{-\infty}(T^*(K/H)) &= \mathcal{Q}_K^{-\infty}(T^*(K/H))|_G \\ &= L^2(K/H)|_G. \end{aligned}$$

In other words, the multiplicity of  $V_\mu^G$  in  $L^2(K/H)$  is equal to the quantization of the reduced space  $[T^*(K/H)]_{\mu, G}$ .  $\square$

#### REFERENCES

- [1] M.F. ATIYAH, Elliptic operators and compact groups, Springer, 1974. Lecture notes in Mathematics, **401**.
- [2] M.F. ATIYAH, G.B. SEGAL, *The index of elliptic operators II*, Ann. Math. **87**, 1968, p. 531-545.
- [3] M.F. ATIYAH, I.M. SINGER, *The index of elliptic operators I*, Ann. Math. **87**, 1968, p. 484-530.
- [4] M.F. ATIYAH, I.M. SINGER, *The index of elliptic operators III*, Ann. Math. **87**, 1968, p. 546-604.
- [5] M.F. ATIYAH, I.M. SINGER, *The index of elliptic operators IV*, Ann. Math. **93**, 1971, p. 139-141.
- [6] N. BERLINE and M. VERGNE, The Chern character of a transversally elliptic symbol and the equivariant index, *Invent. Math.*, **124**, 1996, p. 11-49.
- [7] N. BERLINE and M. VERGNE, L'indice équivariant des opérateurs transversalement elliptiques, *Invent. Math.*, **124**, 1996, p. 51-101.
- [8] J. J. DUISTERMAAT, *The heat equation and the Lefschetz fixed point formula for the Spin<sup>c</sup>-Dirac operator*, Progress in Nonlinear Differential Equation and Their Applications, vol. 18, Birkhauser, Boston, 1996.
- [9] V. GUILLEMIN and S. STERNBERG, Geometric quantization and multiplicities of group representations, *Invent. Math.*, **67**, 1982, p. 515-538.
- [10] V. GUILLEMIN and S. STERNBERG, A normal form for the moment map, in *Differential Geometric Methods in Mathematical Physics*(S. Sternberg, ed.), Reidel Publishing Company, Dordrecht, 1984.
- [11] V. GUILLEMIN and S. STERNBERG, Symplectic techniques in physics, Cambridge University Press, Cambridge, 1990.
- [12] B. KOSTANT, Quantization and unitary representations, in *Modern Analysis and Applications*, Lecture Notes in Math., Vol. 170, Springer-Verlag, 1970, p. 87-207.
- [13] E. LERMAN, Symplectic cut, *Math Res. Lett.* **2**, 1995, p. 247-258.
- [14] E. LERMAN, E. MEINRENKEN, S. TOLMAN and C. WOODWARD, Non-Abelian convexity by symplectic cuts, *Topology*, **37**, 1998, p. 245-259.
- [15] X. MA, W. ZHANG, Geometric quantization for proper moment map, Arxiv:0812.3989

- [16] E. MEINRENKEN, On Riemann-Roch formulas for multiplicities, *J. Amer. Math. Soc.*, **9**, 1996, p. 373-389.
- [17] E. MEINRENKEN, Symplectic surgery and the  $\text{Spin}^c$ -Dirac operator, *Advances in Math.*, **134**, 1998, p. 240-277.
- [18] E. MEINRENKEN, R. SJAMAAR, Singular reduction and quantization, *Topology*, **38**, 1999, p. 699-762.
- [19] P-E. PARADAN, Localization of the Riemann-Roch character, *J. Funct. Anal.* **187**, 2001, p. 442-509.
- [20] P-E. PARADAN,  $\text{Spin}^c$  quantization and the  $K$ -multiplicities of the discrete series, *Annales Scientifiques de l'E. N. S.*, **36**, 2003, p. 805-845.
- [21] P-E. PARADAN, Formal geometric quantization, *Ann. Inst. Fourier* **59**, 2009, p. 199-238.
- [22] P-E. PARADAN M. VERGNE, Index of transversally elliptic operators, 40 pages, to appear in *Astérisque, Soc. Math. Fr.*. Arxiv math/08041225.
- [23] P-E. PARADAN, Multiplicities of the discrete series, 38 pages. ArXiv:0812.0059.
- [24] R. SJAMAAR, E. LERMAN, Stratified symplectic spaces and reduction, *Annals of Math.*, **134**, 1991, p. 375-422.
- [25] R. SJAMAAR, Symplectic reduction and Riemann-Roch formulas for multiplicities, *Bull. Amer. Math. Soc.* **33**, 1996, p. 327-338.
- [26] Y. TIAN, W. ZHANG, An analytic proof of the geometric quantization conjecture of Guillemin-Sternberg, *Invent. Math.* **132**, 1998, p. 229-259.
- [27] M. VERGNE, Multiplicity formula for geometric quantization, Part I, Part II, and Part III, *Duke Math. Journal*, **82**, 1996, p. 143-179, p 181-194, p 637-652.
- [28] M. VERGNE, Quantification géométrique et réduction symplectique, *Séminaire Bourbaki* **888**, 2001.
- [29] M. VERGNE, Applications of Equivariant Cohomology, International Congress of Mathematicians 2006, Vol. I, Eur. Math. Soc., Zürich, 2007, p. 635-664. Arxiv: math/0607389.
- [30] J. WEITSMAN, Non-abelian symplectic cuts and the geometric quantization of noncompact manifolds. EuroConférence Moshé Flato 2000, Part I (Dijon). *Lett. Math. Phys.*, **56**, 2001, no. 1, p. 31-40.
- [31] A. WEINSTEIN, Lecture on symplectic manifold, CBMS Regional Conf. Series in Math., **29**, 1983.

INSTITUT DE MATHÉMATIQUES ET DE MODÉLISATION DE MONTPELLIER (I3M), UNIVERSITÉ MONTPELLIER 2

*E-mail address:* Paul-Emile.Paradan@math.univ-montp2.fr