



Perturbation Method for Particle-like Solutions of the Einstein-Dirac Equations

Simona Rota Nodari

► To cite this version:

Simona Rota Nodari. Perturbation Method for Particle-like Solutions of the Einstein-Dirac Equations. Annales Henri Poincaré, 2009, 10 (7), p. 1377-1393. 10.1007/s00023-009-0015-x . hal-00399486v2

HAL Id: hal-00399486

<https://hal.science/hal-00399486v2>

Submitted on 5 Oct 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PERTURBATION METHOD FOR PARTICLELIKE SOLUTIONS OF EINSTEIN-DIRAC EQUATIONS

SIMONA ROTA NODARI

ABSTRACT. The aim of this work is to prove by a perturbation method the existence of solutions of the coupled Einstein-Dirac equations for a static, spherically symmetric system of two fermions in a singlet spinor state. We relate the solutions of our equations to those of the nonlinear Choquard equation and we show that the nondegenerate solution of Choquard's equation generates solutions for Einstein-Dirac equations.

1. INTRODUCTION

In this paper, we study the coupled Einstein-Dirac equations for a static, spherically symmetric system of two fermions in a singlet spinor state. Using numerical methods, F. Finster, J. Smoller and ST. Yau found, in [1], particlelike solutions; our goal is to give a rigorous proof of their existence by a perturbation method ¹.

The Einstein-Dirac equations take the form

$$(1.1) \quad (D - m)\psi = 0$$

$$(1.2) \quad R_j^i - \frac{1}{2}R\delta_j^i = -8\pi T_j^i$$

where D denotes the Dirac operator, ψ is the wave function of a fermion of mass m , R_j^i is the Ricci curvature tensor, R indicates the scalar curvature and, finally, T_j^i is the energy-momentum tensor of the Dirac particle.

In [1], Finster, Smoller and Yau work with the Dirac operator into a static, spherically symmetric space-time where the metric, in polar coordinates $(t, r, \vartheta, \varphi)$, is given by

$$(1.3) \quad g_{ij} = \text{diag} \left(\frac{1}{T^2}, -\frac{1}{A}, -r^2, -r^2 \sin^2 \vartheta \right)$$

$$(1.4) \quad g^{ij} = \text{diag} \left(T^2, -A, -\frac{1}{r^2}, -\frac{1}{r^2 \sin^2 \vartheta} \right)$$

with $A = A(r)$, $T = T(r)$ positive functions; so, the Dirac operator can be written as

$$(1.5) \quad D = i\gamma^t \partial_t + \gamma^r \left(i\partial_r + \frac{i}{r} \left(1 - A^{-1/2} \right) - \frac{i}{2} \frac{T'}{T} \right) + i\gamma^\vartheta \partial_\vartheta + i\gamma^\varphi \partial_\varphi$$

Date: October 2, 2009.

¹After completing this work, we learned from professor Joel Smoller that Erik J. Bird had proved the existence of small solutions of the Einstein-Dirac equations in his doctoral thesis in 2005 [2]. His method is quite different from ours: he uses Schauder's fixed point theorem.

with

$$(1.6) \quad \gamma^t = T\bar{\gamma}^0$$

$$(1.7) \quad \gamma^r = \sqrt{A} (\bar{\gamma}^1 \cos \vartheta + \bar{\gamma}^2 \sin \vartheta \cos \varphi + \bar{\gamma}^3 \sin \vartheta \sin \varphi)$$

$$(1.8) \quad \gamma^\vartheta = \frac{1}{r} (-\bar{\gamma}^1 \sin \vartheta + \bar{\gamma}^2 \cos \vartheta \cos \varphi + \bar{\gamma}^3 \cos \vartheta \sin \varphi)$$

$$(1.9) \quad \gamma^\varphi = \frac{1}{r \sin \vartheta} (-\bar{\gamma}^2 \sin \varphi + \bar{\gamma}^3 \cos \varphi)$$

where $\bar{\gamma}^i$ are the Dirac matrices in Minkowski space (see [1]).

Moreover, Finster, Smoller and Yau are looking for solutions taking the form

$$(1.10) \quad \psi = e^{-i\omega t} r^{-1} T^{1/2} \begin{pmatrix} \Phi_1 \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ i\Phi_2 \sigma^r \begin{pmatrix} 1 \\ 0 \end{pmatrix} \end{pmatrix},$$

where $\sigma^r = (\bar{\sigma}^1 \cos \vartheta + \bar{\sigma}^2 \sin \vartheta \cos \varphi + \bar{\sigma}^3 \sin \vartheta \sin \varphi)$ is a linear combination of the Pauli matrices $\bar{\sigma}^i$ and $\Phi_1(r)$, $\Phi_2(r)$ are radial real functions.

We remind also that the energy-momentum tensor is obtained as the variation of the classical Dirac action

$$S = \int \bar{\psi} (D - m) \psi \sqrt{|g|} d^4x$$

and takes the form

$$T_j^i = \frac{1}{r^2} \text{diag} \left(2\omega T^2 |\Phi|^2, -2\omega T^2 |\Phi|^2 + 4T \frac{1}{r} \Phi_1 \Phi_2 + 2mT (\Phi_1^2 - \Phi_2^2), \right. \\ \left. -2T \frac{1}{r} \Phi_1 \Phi_2, -2T \frac{1}{r} \Phi_1 \Phi_2 \right)$$

(see [1] for more details).

In this case, the coupled Einstein-Dirac equations can be written as

$$(1.11) \quad \sqrt{A} \Phi_1' = \frac{1}{r} \Phi_1 - (\omega T + m) \Phi_2$$

$$(1.12) \quad \sqrt{A} \Phi_2' = (\omega T - m) \Phi_1 - \frac{1}{r} \Phi_2$$

$$(1.13) \quad rA' = 1 - A - 16\pi\omega T^2 (\Phi_1^2 + \Phi_2^2)$$

$$(1.14) \quad 2rA \frac{T'}{T} = A - 1 - 16\pi\omega T^2 (\Phi_1^2 + \Phi_2^2) + 32\pi \frac{1}{r} T \Phi_1 \Phi_2 + \\ + 16\pi mT (\Phi_1^2 - \Phi_2^2)$$

with the normalization condition

$$(1.15) \quad \int_0^\infty |\Phi|^2 \frac{T}{\sqrt{A}} dr = \frac{1}{4\pi}.$$

In order that the metric be asymptotically Minkowskian, Finster, Smoller and Yau assume that

$$\lim_{r \rightarrow \infty} T(r) = 1.$$

Finally, they also require that the solutions have finite (ADM) mass; namely

$$\lim_{r \rightarrow \infty} \frac{r}{2} (1 - A(r)) < \infty.$$

In this paper, we will prove the existence of solutions of (1.1-1.2) in the form (1.10) by a perturbation method.

In particular, we follow the idea described by Ounaies in [6] (see also [9] for a rigorous existence proof of nonlinear Dirac solitons based on Ounaies' approach). Ounaies, by a perturbation parameter, relates the solutions of a nonlinear Dirac equation to those of nonlinear Schrödinger equation. Imitating the idea of Ounaies, we relate the solutions of our equations to those of nonlinear Choquard's equation (see [4], [5] for more details on Choquard's equation) and we obtain the following result.

Theorem 1.1. *Given $0 < \omega < m$ such that $m - \omega$ is sufficiently small, there exists a non trivial solution of (1.11-1.14).*

In Section 2, we solve the Einstein-Dirac equations by means of the perturbation method suggested by Ounaies; in particular in the first subsection we describe a useful rescaling and some properties of the operators involved, whereas in the second subsection we prove the existence of solutions generated by the solution of the Choquard equation.

2. PERTURBATION METHOD FOR EINSTEIN-DIRAC EQUATIONS

First of all, we observe that writing $T(r) = 1 + t(r)$ and using equation (1.13), the coupled Einstein-Dirac equations become

$$(2.1) \quad \sqrt{A}\Phi_1' = \frac{1}{r}\Phi_1 - (\omega + m)\Phi_2 - \omega t\Phi_2$$

$$(2.2) \quad \sqrt{A}\Phi_2' = (\omega - m)\Phi_1 + \omega t\Phi_1 - \frac{1}{r}\Phi_2$$

$$(2.3) \quad \begin{aligned} 2rAt' &= (A - 1)(1 + t) - 16\pi\omega(1 + t)^3 (\Phi_1^2 + \Phi_2^2) + \\ &\quad + 32\pi\frac{1}{r}(1 + t)^2\Phi_1\Phi_2 + 16\pi m(1 + t)^2 (\Phi_1^2 - \Phi_2^2) \end{aligned}$$

where

$$(2.4) \quad A(r) = 1 - \frac{16\pi\omega}{r} \int_0^r (1 + t(s))^2 (\Phi_1(s)^2 + \Phi_2(s)^2) ds := 1 - \frac{16\pi\omega}{r} Q(r).$$

Furthermore, because we want $A(r) > 0$, we have that the following condition must be satisfied

$$(2.5) \quad 0 \leq \frac{Q(r)}{r} < \frac{1}{16\pi\omega}$$

for all $r \in (0, \infty)$.

Now, to find a solution of the equations (2.1-2.3), we exploit the idea used by Ounaies in [6]. In particular, we proceed as follow: in a first step we use a rescaling argument to transform (2.1-2.3) in a perturbed system of the form

$$(2.6) \quad \begin{cases} \sqrt{A(\varepsilon, \varphi, \chi, \tau)} \frac{d}{dr} \varphi - \frac{1}{r} \varphi + 2m\chi + K_1(\varepsilon, \varphi, \chi, \tau) = 0 \\ \sqrt{A(\varepsilon, \varphi, \chi, \tau)} \frac{d}{dr} \chi + \frac{1}{r} \chi + \varphi - m\varphi\tau + K_2(\varepsilon, \varphi, \chi, \tau) = 0 \\ A(\varepsilon, \varphi, \chi, \tau) \frac{d}{dr} \tau + \frac{8\pi m}{r^2} \int_0^r \varphi^2 ds + K_3(\varepsilon, \varphi, \chi, \tau) = 0 \end{cases}$$

where $\varphi, \chi, \tau : (0, \infty) \rightarrow \mathbb{R}$.

Second, we relate the solutions of (2.6) to those of the nonlinear system

$$(2.7) \quad \begin{cases} -\frac{d^2}{dr^2}\varphi + 2m\varphi - 16\pi m^3 \left(\int_0^\infty \frac{\varphi^2}{\max(r,s)} ds \right) \varphi = 0 \\ \chi(r) = \frac{1}{2m} \left(\frac{1}{r}\varphi - \frac{d}{dr}\varphi \right) \\ \tau(r) = 8\pi m \int_0^\infty \frac{\varphi^2}{\max(r,s)} ds. \end{cases}$$

We remark that φ is a solution of (2.7) if and only if $u(x) = \frac{\varphi(|x|)}{|x|}$ solves the nonlinear Choquard equation

$$(2.8) \quad -\Delta u + 2mu - 4m^3 \left(\int_{\mathbb{R}^3} \frac{|u(y)|^2}{|x-y|} dy \right) u = 0 \quad \text{in } H^1(\mathbb{R}^3).$$

To prove this fact it's enough to remind that for a radial function ρ ,

$$\Delta \rho = \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \rho \right)$$

and

$$(| \cdot | \star \rho)(x) = 4\pi \left(\int_0^\infty \frac{s^2 \rho(s)}{\max(r,s)} ds \right)$$

with $r = |x|$.

We observe also that if we write

$$\begin{pmatrix} u(x) \\ v(x) \end{pmatrix} = r^{-1} \begin{pmatrix} \varphi(r) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ i\chi(r)\sigma^r \begin{pmatrix} 1 \\ 0 \end{pmatrix} \end{pmatrix}$$

with $r = |x|$, (φ, χ) is a solution of (2.7) if and only if $(u(x), v(x))$ solve

$$(2.9) \quad -\Delta u + 2mu - 4m^3 \left(\int_{\mathbb{R}^3} \frac{|u(y)|^2}{|x-y|} dy \right) u = 0 \quad v = \frac{-i\bar{\sigma}\nabla u}{2m}$$

in \mathbb{R}^3 where $\bar{\sigma}\nabla = \sum_{i=1}^3 \bar{\sigma}^i \partial_i$.

It's well known that Choquard's equation (2.8) has a unique radial, positive solution u_0 with $\int |u_0|^2 = N$ for some $N > 0$ given. Furthermore, u_0 is infinitely differentiable and goes to zero at infinity; more precisely there exist some positive constants $C_{\delta,\eta}$ such that $|D^\eta(u_0)| \leq C_{\delta,\eta} \exp(-\delta|x|)$ for $x \in \mathbb{R}^3$. At last, $u_0 \in H^1(\mathbb{R}^3)$ is a radial nondegenerate solution; by this we mean that the linearization of (2.8) around u_0 has a trivial nullspace in $L_r^2(\mathbb{R}^3)$. In particular, the linear operator \mathcal{L} given by

$$\mathcal{L}\xi = -\Delta\xi + 2m\xi - 4m^3 \left(\int_{\mathbb{R}^3} \frac{|u_0(y)|^2}{|x-y|} dy \right) \xi - 8m^3 \left(\int_{\mathbb{R}^3} \frac{\xi(y)u_0(y)}{|x-y|} dy \right) u_0$$

satisfies $\ker \mathcal{L} = \{0\}$ when \mathcal{L} is restricted to $L_r^2(\mathbb{R}^3)$ (see [4], [5], [3] for more details).

The main idea is that the solutions of (2.6) are the zeros of a \mathcal{C}^1 operator $D : \mathbb{R} \times X_\varphi \times X_\chi \times X_\tau \rightarrow Y_\varphi \times Y_\chi \times Y_\tau$. If we denote by $D_{\varphi,\chi,\tau}(\varepsilon, \varphi, \chi, \tau)$ the derivative of $D(\varepsilon, \cdot, \cdot, \cdot)$, by $(\varphi_0, \chi_0, \tau_0)$ the ground state solution of (2.7) and we observe that $D_{\varphi,\chi,\tau}(\varepsilon, \varphi_0, \chi_0, \tau_0)$ is an isomorphism, the application of the implicit function theorem (see [7]) yields the following result, which is equivalent to theorem 1.1.

Theorem 2.1. *Let $(\varphi_0, \chi_0, \tau_0)$ be the ground state solution of (2.7), then there exists $\delta > 0$ and a function $\eta \in \mathcal{C}((0, \delta), X_\varphi \times X_\chi \times X_\tau)$ such that $\eta(0) = (\varphi_0, \chi_0, \tau_0)$ and $(\varepsilon, \eta(\varepsilon))$ is a solution of (2.6), for $0 \leq \varepsilon < \delta$.*

2.1. Rescaling. In this subsection we are going to introduce the new variable (φ, χ, τ) such that $\Phi_1(r) = \alpha\varphi(\lambda r)$, $\Phi_2(r) = \beta\chi(\lambda r)$ and $t(r) = \gamma\tau(\lambda r)$, where Φ_1, Φ_2, t satisfy (2.1-2.3) and $\alpha, \beta, \gamma, \lambda > 0$ are constants to be chosen later.

Using the explicit expressions of A , given in (2.4), we have

$$\begin{aligned} A(\Phi_1, \Phi_2, t) &= 1 - \frac{16\pi\omega\alpha^2}{r} \int_0^r (1 + \gamma\tau)^2 \left(\varphi^2 + \left(\frac{\beta}{\alpha}\chi \right)^2 \right) ds \\ (2.10) \quad &:= A_{\alpha, \beta, \gamma}(\varphi, \chi, \tau) \end{aligned}$$

It's now clear that if Φ_1, Φ_2, t satisfy (2.1-2.3), then φ, χ, τ satisfy the system

$$(2.11) \quad \begin{cases} \sqrt{A_{\alpha, \beta, \gamma}} \frac{\alpha\lambda}{\beta} \frac{d}{dr} \varphi - \frac{\alpha\lambda}{\beta} \frac{1}{r} \varphi + (m + \omega)\chi + \omega\gamma\tau\chi = 0 \\ \sqrt{A_{\alpha, \beta, \gamma}} \frac{d}{dr} \chi + \frac{1}{r} \chi + \frac{\alpha}{\beta\lambda} (m - \omega)\varphi - \frac{\alpha\gamma}{\beta\lambda} \omega\tau\varphi = 0 \\ 2A_{\alpha, \beta, \gamma} r \frac{d}{dr} \tau + K_{\alpha, \beta, \gamma} = 0 \end{cases}$$

with

$$\begin{aligned} K_{\alpha, \beta, \gamma}(\varphi, \chi, \tau) &= \frac{16\pi\omega}{r} \frac{\alpha^2}{\gamma} \left(\int_0^r (1 + \gamma\tau)^2 \left(\varphi^2 + \left(\frac{\beta}{\alpha}\chi \right)^2 \right) ds \right) (1 + \gamma\tau) + \\ &+ 16\pi\omega \frac{\alpha^2}{\gamma} (1 + \gamma\tau)^3 \left(\varphi^2 + \left(\frac{\beta}{\alpha}\chi \right)^2 \right) - 32\pi \frac{\lambda\alpha\beta}{\gamma} \frac{1}{r} (1 + \gamma\tau)^2 \varphi\chi + \\ &- 16\pi m \frac{\alpha^2}{\gamma} (1 + \gamma\tau)^2 \left(\varphi^2 - \left(\frac{\beta}{\alpha}\chi \right)^2 \right). \end{aligned}$$

By adding the conditions $\frac{\alpha}{\beta\lambda}(m - \omega) = 1$, $\frac{\alpha\lambda}{\beta} = 1$, $\frac{\alpha\gamma}{\beta\lambda} = 1$, $\frac{\alpha^2}{\gamma} = 1$ and $m - \omega \geq 0$, we obtain $\alpha = (m - \omega)^{1/2}$, $\lambda = (m - \omega)^{1/2}$, $\beta = m - \omega$ and $\gamma = m - \omega$.

Denoting $\varepsilon = m - \omega$, (2.11) is equivalent to

$$(2.12) \quad \begin{cases} \sqrt{A(\varepsilon, \varphi, \chi, \tau)} \frac{d}{dr} \varphi - \frac{1}{r} \varphi + 2m\chi + K_1(\varepsilon, \varphi, \chi, \tau) = 0 \\ \sqrt{A(\varepsilon, \varphi, \chi, \tau)} \frac{d}{dr} \chi + \frac{1}{r} \chi + \varphi - m\varphi\tau + K_2(\varepsilon, \varphi, \chi, \tau) = 0 \\ A(\varepsilon, \varphi, \chi, \tau) \frac{d}{dr} \tau + \frac{8\pi m}{r^2} \int_0^r \varphi^2 ds + K_3(\varepsilon, \varphi, \chi, \tau) = 0 \end{cases}$$

where $A(\varepsilon, \varphi, \chi, \tau)$, $K_1(\varepsilon, \varphi, \chi, \tau)$, $K_2(\varepsilon, \varphi, \chi, \tau)$ and $K_3(\varepsilon, \varphi, \chi, \tau)$ are defined by

$$(2.13) \quad A(\varepsilon, \varphi, \chi, \tau) = 1 - \frac{16\pi(m - \varepsilon)\varepsilon}{r} \int_0^r (1 + \varepsilon\tau)^2 (\varphi^2 + \varepsilon\chi^2) ds;$$

$$(2.14) \quad K_1(\varepsilon, \varphi, \chi, \tau) = -\varepsilon\chi + \varepsilon(m - \varepsilon)\tau\chi;$$

$$(2.15) \quad K_2(\varepsilon, \varphi, \chi, \tau) = \varepsilon\tau\varphi;$$

and

$$\begin{aligned}
K_3(\varepsilon, \varphi, \chi, \tau) = & \frac{8\pi m \varepsilon}{r^2} \int_0^r \chi^2 ds + \frac{16\pi m \varepsilon}{r^2} \int_0^r \tau (\varphi^2 + \varepsilon \chi^2) ds + \\
& + \frac{8\pi m \varepsilon^2}{r^2} \int_0^r \tau^2 (\varphi^2 + \varepsilon \chi^2) ds + \\
& - \frac{8\pi \varepsilon}{r^2} \left(\int_0^r (1 + \varepsilon \tau)^2 (\varphi^2 + \varepsilon \chi^2) ds \right) \tau + \\
& + \frac{8\pi(m - \varepsilon)\varepsilon}{r^2} \left(\int_0^r (1 + \varepsilon \tau)^2 (\varphi^2 + \varepsilon \chi^2) ds \right) \tau + \\
& + 16\pi m \varepsilon \frac{\chi^2}{r} + 8\pi m \varepsilon (3\tau + 3\varepsilon \tau^2 + \varepsilon^2 \tau^3) \frac{(\varphi^2 + \varepsilon \chi^2)}{r} + \\
& - 8\pi \varepsilon (1 + \varepsilon \tau)^3 \frac{(\varphi^2 + \varepsilon \chi^2)}{r} - 16\pi \varepsilon (1 + \varepsilon \tau)^2 \frac{\varphi \chi}{r^2} + \\
(2.16) \quad & - 8\pi m \varepsilon (2\tau + \varepsilon \tau^2) \frac{(\varphi^2 - \varepsilon \chi^2)}{r}.
\end{aligned}$$

For $\varepsilon = 0$, (2.12) becomes

$$(2.17) \quad \begin{cases} \frac{d}{dr} \varphi - \frac{1}{r} \varphi + 2m\chi = 0 \\ \frac{d}{dr} \chi + \frac{1}{r} \chi + \varphi - m\varphi\tau = 0 \\ \frac{d}{dr} \tau + \frac{8\pi m}{r^2} \int_0^r \varphi^2 ds = 0 \end{cases}$$

that is equivalent to

$$(2.18) \quad \begin{cases} -\frac{d^2}{dr^2} \varphi + 2m\varphi - 16\pi m^3 \left(\int_0^\infty \frac{\varphi^2}{\max(r,s)} ds \right) \varphi = 0 \\ \chi(r) = \frac{1}{2m} \left(\frac{1}{r} \varphi - \frac{d}{dr} \varphi \right) \\ \tau(r) = 8\pi m \int_0^\infty \frac{\varphi^2}{\max(r,s)} ds. \end{cases}$$

Then, we denote by $(\varphi_0, \chi_0, \tau_0)$ a solution of (2.18); in particular

$$\begin{aligned}
\chi_0(r) &= -\frac{r}{2m} \frac{d}{dr} \left(\frac{\varphi_0}{r} \right) \\
\tau_0(r) &= 8\pi m \int_0^\infty \frac{\varphi_0^2}{\max(r,s)} ds.
\end{aligned}$$

Now, to obtain a solution of (2.12) from $(\varphi_0, \chi_0, \tau_0)$, we define the operators $L_1 : \mathbb{R} \times X_\varphi \times X_\chi \times X_\tau \rightarrow Y_\varphi$, $L_2 : \mathbb{R} \times X_\varphi \times X_\chi \times X_\tau \rightarrow Y_\chi$, $L_3 : \mathbb{R} \times X_\varphi \times X_\chi \times X_\tau \rightarrow Y_\tau$ and $D : \mathbb{R} \times X_\varphi \times X_\chi \times X_\tau \rightarrow Y_\varphi \times Y_\chi \times Y_\tau$ by

$$\begin{aligned}
L_1(\varepsilon, \varphi, \chi, \tau) &= \sqrt{A(\varepsilon, \varphi, \chi, \tau)} \frac{1}{r} \frac{d}{dr} \varphi - \frac{\varphi}{r^2} + 2m \frac{\chi}{r} + \frac{1}{r} K_1(\varepsilon, \varphi, \chi, \tau) \\
L_2(\varepsilon, \varphi, \chi, \tau) &= \sqrt{A(\varepsilon, \varphi, \chi, \tau)} \frac{1}{r} \frac{d}{dr} \chi + \frac{\chi}{r^2} + \frac{\varphi}{r} - m \frac{\varphi}{r} \tau + \frac{1}{r} K_2(\varepsilon, \varphi, \chi, \tau) \\
L_3(\varepsilon, \varphi, \chi, \tau) &= A(\varepsilon, \varphi, \chi, \tau) \frac{d}{dr} \tau + \frac{8\pi m}{r^2} \int_0^r \varphi^2 ds + K_3(\varepsilon, \varphi, \chi, \tau)
\end{aligned}$$

and

$$D(\varepsilon, \varphi, \chi, \tau) = (L_1(\varepsilon, \varphi, \chi, \tau), L_2(\varepsilon, \varphi, \chi, \tau), L_3(\varepsilon, \varphi, \chi, \tau))$$

where

$$\begin{aligned}
X_\varphi &= \left\{ \varphi : (0, \infty) \rightarrow \mathbb{R} \left| \frac{\varphi(|x|)}{|x|} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \in H^1(\mathbb{R}^3, \mathbb{R}^2) \right. \right\} \\
X_\chi &= \left\{ \chi : (0, \infty) \rightarrow \mathbb{R} \left| \frac{\chi(|x|)}{|x|} \sigma^r \begin{pmatrix} 1 \\ 0 \end{pmatrix} \in H^1(\mathbb{R}^3, \mathbb{C}^2) \right. \right\} \\
X_\tau &= \left\{ \tau : (0, \infty) \rightarrow \mathbb{R} \left| \lim_{r \rightarrow \infty} \tau(r) \rightarrow 0, \frac{d}{dr} \tau \in L^1((0, \infty), dr) \right. \right\} \\
Y_\varphi &= Y_\chi = L^2(\mathbb{R}^3) \\
Y_\tau &= L^1((0, \infty), dr).
\end{aligned}$$

Furthermore we define the following norms:

$$\begin{aligned}
\|\varphi\|_{X_\varphi} &= \left\| \frac{\varphi(|x|)}{|x|} \right\|_{H^1(\mathbb{R}^3)}, \\
\|\chi\|_{X_\chi} &= \left\| \frac{\chi(|x|)}{|x|} \sigma^r \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\|_{H^1(\mathbb{R}^3)}, \\
\|\tau\|_{X_\tau} &= \left\| \frac{d}{dr} \tau \right\|_{L^1((0, \infty), dr)}.
\end{aligned}$$

It's well known that

$$\begin{aligned}
H^1(\mathbb{R}^3) &\hookrightarrow L^q(\mathbb{R}^3) \quad 2 \leq q \leq 6 \\
X_\tau &\hookrightarrow L^\infty((0, \infty), dr).
\end{aligned}$$

Moreover, using Hardy's inequality

$$\int_{\mathbb{R}^3} \frac{|f|^2}{|x|^2} dx \leq 4 \int_{\mathbb{R}^3} |\nabla f|^2 dx,$$

we get the following properties:

$$\begin{aligned}
(2.19) \quad \rho &\in H^1((0, \infty), dr) \hookrightarrow L^\infty((0, \infty), dr) \\
\frac{\rho}{r} &\in L^2((0, \infty), dr).
\end{aligned}$$

$\forall \rho \in X_\varphi, \forall \rho \in X_\chi$.

Since the operator $A(\varepsilon, \varphi, \chi, \tau)$ must be strictly positive, we consider $B_\varphi, B_\chi, B_\tau$, defined as the balls of the spaces $X_\varphi, X_\chi, X_\tau$, and $\varepsilon_1, \varepsilon_2$, depending on m and on the radius of $B_\varphi, B_\chi, B_\tau$, such that

$$1 - \frac{16\pi(m-\varepsilon)\varepsilon}{r} \int_0^r (1+\varepsilon\tau)^2 (\varphi^2 + \varepsilon\chi^2) ds \geq \delta > 0$$

for all $(\varepsilon, \varphi, \chi, \tau) \in (-\varepsilon_1, \varepsilon_2) \times B_\varphi \times B_\chi \times B_\tau$. The existence of $\varepsilon_1, \varepsilon_2$ is assured by the fact that φ, χ, τ are bounded; in particular, if $\varepsilon \geq 0$,

$$\begin{aligned}
&1 - \frac{16\pi(m-\varepsilon)\varepsilon}{r} \int_0^r (1+\varepsilon\tau)^2 (\varphi^2 + \varepsilon\chi^2) ds \geq \\
&\geq 1 - 20m\varepsilon \|\varphi\|_{X_\varphi}^2 - 8m\varepsilon^2 \left(5 \|\tau\|_{X_\tau} \|\varphi\|_{X_\varphi}^2 + \|\chi\|_{X_\chi}^2 \right) + \\
&- 4m\varepsilon^3 \|\tau\|_{X_\tau} \left(5 \|\tau\|_{X_\tau} \|\varphi\|_{X_\varphi}^2 + 4 \|\chi\|_{X_\chi}^2 \right) - 8m\varepsilon^4 \|\tau\|_{X_\tau}^2 \|\chi\|_{X_\chi}^2,
\end{aligned}$$

then there exists $\varepsilon_2 > 0$ such that $A(\varepsilon, \varphi, \chi, \tau) > 0$ for all $\varepsilon \in [0, \varepsilon_2]$. In the same way, if $\varepsilon < 0$,

$$\begin{aligned} 1 - \frac{16\pi(m-\varepsilon)\varepsilon}{r} \int_0^r (1+\varepsilon\tau)^2 (\varphi^2 + \varepsilon\chi^2) ds &\geq \\ &\geq 1 - 8m\varepsilon^2 \|\chi\|_{X_\chi}^2 - 8m|\varepsilon|^3 \|\chi\|_{X_\chi}^2 (1 + 2\|\tau\|_{X_\tau}) + \\ &\quad - 8m\varepsilon^4 \|\chi\|_{X_\chi}^2 \|\tau\|_{X_\tau} (2 + 1\|\tau\|_{X_\tau}) - 8m|\varepsilon|^5 \|\tau\|_{X_\tau}^2 \|\chi\|_{X_\chi}^2, \end{aligned}$$

then there exists $\varepsilon_1 > 0$ such that $A(\varepsilon, \varphi, \chi, \tau) > 0$ for all $\varepsilon \in (-\varepsilon_1, 0)$.

Lemma 2.2. *The operators $L_1, L_2 \in \mathcal{C}^1((-\varepsilon_1, \varepsilon_2) \times B_\varphi \times B_\chi \times B_\tau, Y_\varphi)$ and $L_3 \in \mathcal{C}^1((-\varepsilon_1, \varepsilon_2) \times B_\varphi \times B_\chi \times B_\tau, Y_\tau)$.*

Before starting the proof of the lemma we observe that for a radial function ρ such that $\frac{\rho}{r} \in H_r^1(\mathbb{R}^3)$ we have

$$(2.20) \quad |\rho(r)| \leq r^{1/2} \left\| \frac{d}{dr} \left(\frac{\rho(r)}{r} \right) \right\|_{L_{rad}^2}.$$

We remind that $H_r^1(\mathbb{R}^3) = \{u \in H^1(\mathbb{R}^3) \mid u \text{ is radial}\}$.

Proof. We begin with L_3 ; first, we have to prove that it is well defined in $Y_\tau = L^1((0, \infty), dr)$. We remark that

$$\begin{aligned} |L_3(\varepsilon, \varphi, \chi, \tau)| &\leq C_1 \left| \frac{d}{dr} \tau \right| + \frac{C_2}{r^2} \int_0^r |\varphi^2 + \varepsilon\chi^2| ds + \frac{C_3}{r} |\varphi^2 + \varepsilon\chi^2| + \\ &\quad + \frac{C_4}{r^2} |\varphi\chi| + \frac{C_5}{r} |\varphi^2 - \varepsilon\chi^2| \end{aligned}$$

where C_1, C_2, C_3, C_4, C_5 are positive constants and, by definition, we have that $\frac{d}{dr} \tau \in L^1((0, \infty), dr)$.

Next, we have

$$\int_0^\infty \frac{1}{r^2} \int_0^r |\varphi^2 + \varepsilon\chi^2| ds dr = \int_0^\infty \frac{|\varphi^2 + \varepsilon\chi^2|}{s} ds < +\infty,$$

using Hölder's inequality, then $\frac{1}{r^2} \int_0^r |\varphi^2 + \varepsilon\chi^2| ds \in Y_\tau$. In the same way, we can conclude that $\frac{1}{r} (\varphi^2 + \varepsilon\chi^2), \frac{1}{r} (\varphi^2 - \varepsilon\chi^2) \in Y_\tau$.

Finally,

$$\int_0^\infty \frac{|\varphi|}{r} \frac{|\chi|}{r} dr \leq C \left\| \frac{\varphi}{r} \right\|_{L^2((0, \infty))} \left\| \frac{\chi}{r} \right\|_{L^2((0, \infty))} < +\infty$$

thanks to (2.19), then $\frac{1}{r^2} \varphi\chi \in Y_\tau$.

Now, we have to prove that $L_3(\varepsilon, \varphi, \chi, \tau)$ is \mathcal{C}^1 ; by classical arguments, it's enough to show that for $(h_1, h_2, h_3) \in B_\varphi \times B_\chi \times B_\tau$

$$\begin{aligned} \frac{\partial}{\partial \varphi} (L_3(\varepsilon, \varphi, \chi, \tau)) h_1 &\in Y_\tau, \\ \frac{\partial}{\partial \chi} (L_3(\varepsilon, \varphi, \chi, \tau)) h_2 &\in Y_\tau, \\ \frac{\partial}{\partial \tau} (L_3(\varepsilon, \varphi, \chi, \tau)) h_3 &\in Y_\tau. \end{aligned}$$

We begin with $\frac{\partial}{\partial \varphi} (L_3 (\varepsilon, \varphi, \chi, \tau))$,

$$\begin{aligned} \frac{\partial}{\partial \varphi} (L_3 (\varepsilon, \varphi, \chi, \tau)) h_1 &= \left(\frac{\partial}{\partial \varphi} (A (\varepsilon, \varphi, \chi, \tau)) h_1 \right) \frac{d}{dr} \tau + \\ &+ \frac{16\pi(m-\varepsilon)}{r^2} \left(\int_0^r (1+\varepsilon\tau)^2 \varphi h_1 ds \right) (1+\varepsilon\tau) + \\ &+ 16\pi(m-\varepsilon)(1+\varepsilon\tau)^3 \frac{\varphi h_1}{r} - 16\pi\varepsilon(1+\varepsilon\tau)^2 \frac{h_1 \chi}{r^2} + \\ &- 16\pi m(1+\varepsilon\tau)^2 \frac{\varphi h_1}{r}; \end{aligned}$$

for $\frac{\partial}{\partial \chi} (L_3 (\varepsilon, \varphi, \chi, \tau))$,

$$\begin{aligned} \frac{\partial}{\partial \chi} (L_3 (\varepsilon, \varphi, \chi, \tau)) h_2 &= \left(\frac{\partial}{\partial \chi} (A (\varepsilon, \varphi, \chi, \tau)) h_2 \right) \frac{d}{dr} \tau + \\ &+ \frac{16\pi(m-\varepsilon)\varepsilon}{r^2} \left(\int_0^r (1+\varepsilon\tau)^2 \chi h_2 ds \right) (1+\varepsilon\tau) + \\ &+ 16\pi(m-\varepsilon)\varepsilon(1+\varepsilon\tau)^3 \frac{\chi h_2}{r} - 16\pi\varepsilon(1+\varepsilon\tau)^2 \frac{\varphi h_2}{r^2} + \\ &- 16\pi m\varepsilon(1+\varepsilon\tau)^2 \frac{\chi h_2}{r} \end{aligned}$$

and, finally,

$$\begin{aligned} \frac{\partial}{\partial \tau} (L_3 (\varepsilon, \varphi, \chi, \tau)) h_3 &= \left(\frac{\partial}{\partial \tau} (A (\varepsilon, \varphi, \chi, \tau)) h_3 \right) \frac{d}{dr} \tau + \\ &+ A (\varepsilon, \varphi, \chi, \tau) \frac{d}{dr} h_3 + \\ &+ \frac{16\pi(m-\varepsilon)\varepsilon}{r^2} \left(\int_0^r (1+\varepsilon\tau) h_3 (\varphi^2 + \varepsilon\chi^2) ds \right) (1+\varepsilon\tau) + \\ &+ \frac{8\pi(m-\varepsilon)\varepsilon}{r^2} \left(\int_0^r (1+\varepsilon\tau)^2 (\varphi^2 + \varepsilon\chi^2) ds \right) h_3 + \\ &+ 24\pi(m-\varepsilon)\varepsilon(1+\varepsilon\tau)^2 \frac{(\varphi^2 + \varepsilon\chi^2)}{r} h_3 - 32\pi\varepsilon^2(1+\varepsilon\tau) \frac{\varphi\chi}{r^2} h_3 + \\ &- 16\pi m\varepsilon(1+\varepsilon\tau) \frac{(\varphi^2 - \varepsilon\chi^2)}{r} h_3. \end{aligned}$$

First of all, we remark that if $\varphi, h_1 \in B_\varphi$, $\chi, h_2 \in B_\chi$ and $\tau, h_3 \in B_\tau$, then $\frac{\partial}{\partial \varphi} (A (\varepsilon, \varphi, \chi, \tau)) h_1$, $\frac{\partial}{\partial \chi} (A (\varepsilon, \varphi, \chi, \tau)) h_2$ and $\frac{\partial}{\partial \tau} (A (\varepsilon, \varphi, \chi, \tau)) h_3$ are bounded. So, we have that

$$\begin{aligned} \left| \frac{\partial L_3}{\partial \varphi} h_1 \right| &\leq C_1 \left| \frac{d}{dr} \tau \right| + \frac{C_2}{r^2} \left(\int_0^r |\varphi h_1| ds \right) + C_3 \frac{|\varphi h_1|}{r} + C_4 \frac{|h_1 \chi|}{r^2} \\ \left| \frac{\partial L_3}{\partial \chi} h_2 \right| &\leq C_5 \left| \frac{d}{dr} \tau \right| + \frac{C_6}{r^2} \left(\int_0^r |\chi h_2| ds \right) + C_7 \frac{|\chi h_2|}{r} + C_8 \frac{|\varphi h_2|}{r^2} \\ \left| \frac{\partial L_3}{\partial \tau} h_3 \right| &\leq C_9 \left| \frac{d}{dr} \tau \right| + C_{10} \left| \frac{d}{dr} h_3 \right| + \frac{C_{11}}{r^2} \int_0^r |\varphi^2 + \varepsilon\chi^2| ds + C_{12} \frac{|\varphi^2 + \varepsilon\chi^2|}{r} + \\ &+ C_{13} \frac{|\varphi\chi|}{r^2} + C_{14} \frac{|\varphi^2 - \varepsilon\chi^2|}{r} \end{aligned}$$

with C_i positive constants. With exactly the same arguments used above, we conclude that

$$\begin{aligned} \int_0^\infty \left| \frac{\partial}{\partial \varphi} (L_3(\varepsilon, \varphi, \chi, \tau)) h_1 \right| dr &< +\infty \\ \int_0^\infty \left| \frac{\partial}{\partial \chi} (L_3(\varepsilon, \varphi, \chi, \tau)) h_2 \right| dr &< +\infty \\ \int_0^\infty \left| \frac{\partial}{\partial \tau} (L_3(\varepsilon, \varphi, \chi, \tau)) h_3 \right| dr &< +\infty \end{aligned}$$

if $(\varepsilon, \varphi, \chi, \tau) \in (-\varepsilon_1, \varepsilon_2) \times B_\varphi \times B_\chi \times B_\tau$, and $(h_1, h_2, h_3) \in B_\varphi \times B_\chi \times B_\tau$. Furthermore $\frac{\partial L_3}{\partial \varphi}$, $\frac{\partial L_3}{\partial \chi}$ and $\frac{\partial L_3}{\partial \tau}$ are continuous; thus the proof for L_3 .

We consider now L_1 ; first, we have to prove that it is well defined in Y_φ . We observe that

$$|L_1(\varepsilon, \varphi, \chi, \tau)| \leq C_1 \left| \frac{1}{r} \frac{d}{dr} \varphi \right| + \left| \frac{\varphi}{r^2} \right| + C_2 \left| \frac{\chi}{r} \right|$$

with C_1, C_2 positive constants then, $L_1(\varepsilon, \varphi, \chi, \tau) \in L^2(\mathbb{R}^3)$, thanks to conditions (2.19).

Now, we have to prove that $L_1(\varepsilon, \varphi, \chi, \tau)$ is \mathcal{C}^1 ; by classical arguments, it's enough to show that for $(h_1, h_2, h_3) \in B_\varphi \times B_\chi \times B_\tau$

$$\begin{aligned} \frac{\partial}{\partial \varphi} (L_1(\varepsilon, \varphi, \chi, \tau)) h_1 &\in Y_\varphi, \\ \frac{\partial}{\partial \chi} (L_1(\varepsilon, \varphi, \chi, \tau)) h_2 &\in Y_\varphi, \\ \frac{\partial}{\partial \tau} (L_1(\varepsilon, \varphi, \chi, \tau)) h_3 &\in Y_\varphi. \end{aligned}$$

By a straightforward computation, we find out

$$\begin{aligned} \frac{\partial L_1}{\partial \varphi} h_1 &= \frac{1}{2} A^{-1/2} \left(\frac{\partial A}{\partial \varphi} h_1 \right) \frac{1}{r} \frac{d}{dr} \varphi + A^{1/2} \frac{1}{r} \frac{d}{dr} h_1 - \frac{h_1}{r^2}, \\ \frac{\partial L_1}{\partial \chi} h_2 &= \frac{1}{2} A^{-1/2} \left(\frac{\partial A}{\partial \chi} h_2 \right) \frac{1}{r} \frac{d}{dr} \varphi + (2m - \varepsilon) \frac{h_2}{r} + \varepsilon(m - \varepsilon) \tau \frac{h_2}{r}, \\ \frac{\partial L_1}{\partial \tau} h_3 &= \frac{1}{2} A^{-1/2} \left(\frac{\partial A}{\partial \tau} h_3 \right) \frac{1}{r} \frac{d}{dr} \varphi + \varepsilon(m - \varepsilon) h_3 \frac{\chi}{r}; \end{aligned}$$

and, using the positivity of A ,

$$\begin{aligned} \left| \frac{\partial L_1}{\partial \varphi} h_1 \right| &\leq C_1 \left| \frac{1}{r} \frac{d}{dr} \varphi \right| + C_2 \left| \frac{1}{r} \frac{d}{dr} h_1 \right| + \left| \frac{h_1}{r^2} \right| \\ \left| \frac{\partial L_1}{\partial \chi} h_2 \right| &\leq C_3 \left| \frac{1}{r} \frac{d}{dr} \varphi \right| + C_4 \left| \frac{h_2}{r} \right| \\ \left| \frac{\partial L_1}{\partial \tau} h_3 \right| &\leq C_5 \left| \frac{1}{r} \frac{d}{dr} \varphi \right| + C_6 \left| \frac{\chi}{r} \right|. \end{aligned}$$

with C_i positive constants. Then, we can conclude that

$$\begin{aligned} \int_{\mathbb{R}^3} \left| \frac{\partial}{\partial \varphi} (L_3(\varepsilon, \varphi, \chi, \tau)) h_1 \right|^2 dx &< +\infty \\ \int_{\mathbb{R}^3} \left| \frac{\partial}{\partial \chi} (L_3(\varepsilon, \varphi, \chi, \tau)) h_2 \right|^2 dx &< +\infty \\ \int_{\mathbb{R}^3} \left| \frac{\partial}{\partial \tau} (L_3(\varepsilon, \varphi, \chi, \tau)) h_3 \right|^2 dx &< +\infty \end{aligned}$$

if $(\varepsilon, \varphi, \chi, \tau) \in (-\varepsilon_1, \varepsilon_2) \times B_\varphi \times B_\chi \times B_\tau$, and $(h_1, h_2, h_3) \in B_\varphi \times B_\chi \times B_\tau$. Furthermore $\frac{\partial L_1}{\partial \varphi}$, $\frac{\partial L_1}{\partial \chi}$ and $\frac{\partial L_1}{\partial \tau}$ are continuous; thus the proof for L_1 and with the same arguments for L_2 . \square

2.2. Branches generated by solutions of Choquard equation. In this subsection, we show that a solution $\phi_0 = (\varphi_0, \chi_0, \tau_0)$ of (2.7) can generate a local branch of solutions of (2.6).

First, we linearize the operator D on (φ, χ, τ) around $(0, \phi_0)$

$$D_{\varphi, \chi, \tau}(0, \phi_0)(h, k, l) = \begin{pmatrix} \frac{1}{r} \frac{d}{dr} h - \frac{h}{r^2} + 2m \frac{k}{r} \\ \frac{1}{r} \frac{d}{dr} k + \frac{k}{r^2} + \frac{h}{r} - m \frac{h}{r} \tau_0 - m \frac{\varphi_0}{r} l \\ \frac{d}{dr} l + \frac{16\pi m}{r^2} \int_0^r \varphi_0 h ds \end{pmatrix}.$$

Now, if we prove that $D_{\varphi, \chi, \tau}(0, \phi_0)$ is an isomorphism, the implicit function theorem can be applied and we can find solutions of (2.6) near the ground state ϕ_0 .

Lemma 2.3. *We define the operator $V : X_\varphi \times X_\chi \rightarrow Y_\varphi \times Y_\chi$, by*

$$V(\varphi, \chi) = \begin{pmatrix} \frac{1}{r} \frac{d}{dr} \varphi - \frac{\varphi}{r^2} + 2m \frac{\chi}{r} \\ \frac{1}{r} \frac{d}{dr} \chi + \frac{\chi}{r^2} + \frac{\varphi}{r} \end{pmatrix},$$

then V is an isomorphism of $X_\varphi \times X_\chi$ onto $Y_\varphi \times Y_\chi$.

This lemma is obvious if we remind that $L^2(\mathbb{R}^3, \mathbb{C}^4)$ can be written as the direct sum of partial wave subspaces and that the Dirac operator leaves invariant all these subspaces (see [8]). So, thanks to lemma 2.3 of [6], we know that $\overline{V} : H^1(\mathbb{R}^3, \mathbb{C}^2) \times H^1(\mathbb{R}^3, \mathbb{C}^2) \rightarrow L^2(\mathbb{R}^3, \mathbb{C}^2) \times L^2(\mathbb{R}^3, \mathbb{C}^2)$ defined by

$$\overline{V}(u, v) = \begin{pmatrix} i\bar{\sigma} \nabla u + 2mv \\ -i\bar{\sigma} \nabla v + u \end{pmatrix}$$

is an isomorphism of $H^1(\mathbb{R}^3, \mathbb{C}^2) \times H^1(\mathbb{R}^3, \mathbb{C}^2)$ onto $L^2(\mathbb{R}^3, \mathbb{C}^2) \times L^2(\mathbb{R}^3, \mathbb{C}^2)$ and then \overline{V} is an isomorphism of each partial wave subspace. In particular, V coincide with \overline{V} on the partial wave subspace $X_\varphi \times X_\chi$.

Lemma 2.4. *We define the operator $W : X_\varphi \times X_\chi \times X_\tau \rightarrow Y_\varphi \times Y_\chi \times Y_\tau$, by*

$$W(h, k, l) = \begin{pmatrix} \frac{1}{r} \frac{d}{dr} h - \frac{h}{r^2} + 2m \frac{k}{r} \\ \frac{1}{r} \frac{d}{dr} k + \frac{k}{r^2} + \frac{h}{r} - m \frac{\varphi_0}{r} l \\ \frac{d}{dr} l \end{pmatrix},$$

then W is an isomorphism of $X_\varphi \times X_\chi \times X_\tau$ onto $Y_\varphi \times Y_\chi \times Y_\tau$.

Proof. First we prove that W is one to one. We observe that $W(h, k, l) = 0$ if and only if (h, k, l) satisfies

$$\begin{cases} \frac{1}{r} \frac{d}{dr} h - \frac{h}{r^2} + 2m \frac{k}{r} = 0 \\ \frac{1}{r} \frac{d}{dr} k + \frac{k}{r^2} + \frac{h}{r} - m \frac{\varphi_0}{r} l = 0 \\ \frac{d}{dr} l = 0 \end{cases}$$

in $Y_\varphi \times Y_\chi \times Y_\tau$. In particular, we must have $l \equiv 0$ and (h, k) solution of

$$\begin{cases} \frac{1}{r} \frac{d}{dr} h - \frac{h}{r^2} + 2m \frac{k}{r} = 0 \\ \frac{1}{r} \frac{d}{dr} k + \frac{k}{r^2} + \frac{h}{r} = 0 \end{cases}$$

that is equivalent to $V(h, k) = 0$. So, thank to lemma 2.3, $h \equiv k \equiv 0$ and W is one to one in $Y_\varphi \times Y_\chi \times Y_\tau$.

Secondly, we have to prove that for $f = (f_1, f_2, f_3) \in Y_\varphi \times Y_\chi \times Y_\tau$, there exists $(h, k, l) \in X_\varphi \times X_\chi \times X_\tau$ such that $W(h, k, l) = f$. This means that the system

$$\begin{cases} \frac{1}{r} \frac{d}{dr} h - \frac{h}{r^2} + 2m \frac{k}{r} = f_1 \\ \frac{1}{r} \frac{d}{dr} k + \frac{k}{r^2} + \frac{h}{r} - m \frac{\varphi_0}{r} l = f_2 \\ \frac{d}{dr} l = f_3 \end{cases}$$

has a solution in $X_\varphi \times X_\chi \times X_\tau$ for all $(f_1, f_2, f_3) \in Y_\varphi \times Y_\chi \times Y_\tau$. We observe that $\forall f_3 \in L^1((0, \infty), dr)$ there exist $l^*(r) = -\int_r^\infty f_3 ds$ such that $\frac{d}{dr} l^* = f_3$; furthermore $l^* \in X_\tau$. So, we have to show that

$$(2.21) \quad \begin{cases} \frac{1}{r} \frac{d}{dr} h - \frac{h}{r^2} + 2m \frac{k}{r} = f_1 \\ \frac{1}{r} \frac{d}{dr} k + \frac{k}{r^2} + \frac{h}{r} = f_2 + m \frac{\varphi_0}{r} l^* \end{cases}$$

has a solution in $X_\varphi \times X_\chi$ for all $(f_1, f_2) \in Y_\varphi \times Y_\chi$.

Now, we remark that $\frac{\varphi_0}{r} l^* \in L^2(\mathbb{R}^3)$ and then, thanks to lemma 2.3, (2.21) has a solution in $X_\varphi \times X_\chi$ for all $(f_1, f_2) \in Y_\varphi \times Y_\chi$.

In conclusion W is an isomorphism of $X_\varphi \times X_\chi \times X_\tau$ onto $Y_\varphi \times Y_\chi \times Y_\tau$. \square

Finally, we observe that $D_{\varphi, \chi, \tau}(0, \phi_0)(h, k, l)$ can be written as

$$(2.22) \quad D_{\varphi, \chi, \tau}(0, \phi_0)(h, k, l) = W(h, k, l) + S(h)$$

with

$$(2.23) \quad S(h) = \begin{pmatrix} 0 \\ -m \frac{h}{r} \tau_0 \\ \frac{16\pi m}{r^2} \int_0^r \varphi_0 h ds \end{pmatrix}.$$

Theorem 2.5. *Let ϕ_0 be the ground state solution of (2.7), then there exists $\delta > 0$ and a function $\eta \in \mathcal{C}((0, \delta), X_\varphi \times X_\chi \times X_\tau)$ such that $\eta(0) = \phi_0$ and $D(\varepsilon, \eta(\varepsilon)) = 0$ for $0 \leq \varepsilon < \delta$.*

Proof. Since $D(0, \phi_0) = 0$ and D is continuously differentiable in a neighborhood of $(0, \phi_0)$, to apply the implicit function theorem we have to prove that $D_{\varphi, \chi, \tau}(0, \phi_0)$

is an isomorphism of $X_\varphi \times X_\chi \times X_\tau$ onto $Y_\varphi \times Y_\chi \times Y_\tau$.

We observe that $D_{\varphi,\chi,\tau}(0, \phi_0)(h, k, l) = 0$ if and only if (h, k, l) satisfies

$$(2.24) \quad \begin{cases} \frac{d}{dr}h - \frac{h}{r} + 2mk = 0 \\ \frac{d}{dr}k + \frac{k}{r} + h - mh\tau_0 - m\varphi_0l = 0 \\ \frac{d}{dr}l + \frac{16\pi m}{r^2} \int_0^r \varphi_0 h ds = 0 \end{cases}$$

that means

$$(2.25) \quad \begin{cases} -\frac{d^2}{dr^2}h + 2mh - 16\pi m^3 \left(\int_0^\infty \frac{\varphi_0^2}{\max(r,s)} ds \right) h - 32\pi m^3 \left(\int_0^\infty \frac{\varphi_0 h}{\max(r,s)} ds \right) \varphi_0 = 0 \\ \frac{d}{dr}h - \frac{h}{r} + 2mk = 0 \\ l = 16\pi m \int_0^\infty \frac{\varphi_0 h}{\max(r,s)} ds \end{cases}$$

Now, if we write $\xi(x) = \frac{h(|x|)}{|x|}$ and we remind that $\varphi_0(|x|) = |x|u_0(x)$ with u_0 solution of (2.8), we have that (h, k, l) is a solution of (2.25) if

$$\begin{pmatrix} \xi(x) \\ \zeta(x) \end{pmatrix} = r^{-1} \begin{pmatrix} h(r) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ ik(r)\sigma^r \begin{pmatrix} 1 \\ 0 \end{pmatrix} \end{pmatrix}$$

satisfies

$$(2.26) \quad \begin{cases} -\Delta\xi + 2m\xi - 4m^3 \left(\int_{\mathbb{R}^3} \frac{|u_0(y)|^2}{|x-y|} dy \right) \xi - 8m^3 \left(\int_{\mathbb{R}^3} \frac{\xi(y)u_0(y)}{|x-y|} dy \right) u_0 = 0 \\ \zeta = \frac{-i\sigma \nabla \xi}{2m} \end{cases}$$

and

$$(2.27) \quad l(x) = 4m \int_{\mathbb{R}^3} \frac{\xi(y)u_0(y)}{|x-y|} dy.$$

It's well known that the unique solution of the first equation of (2.26) in $H_r^1(\mathbb{R}^3)$ is $\xi \equiv 0$ (see [3] for more details) and that implies $\zeta \equiv l \equiv 0$. So the unique solution of (2.24) is $h \equiv k \equiv l \equiv 0$ and $D_{\varphi,\chi,\tau}(0, \phi_0)$ is one to one in $X_\varphi \times X_\chi \times X_\tau$.

Next, if we show that $S(h)$ is a compact operator, we have that $D_{\varphi,\chi,\tau}(0, \phi_0)$ is a one to one operator that can be written as a sum of an isomorphism and a compact operator and then it's an isomorphism.

First, we can easily see that $T(h) = \frac{1}{r^2} \left(\int_0^r \varphi_0 h ds \right)$ is a compact operator from X_φ on Y_τ ; in particular, we use the fact that $H_r^1(\mathbb{R}^3)$ is compactly embedded in $L^q(\mathbb{R}^3)$, for $2 < q < 6$, to prove that for any bounded sequence $\{h_n\} \subset X_\varphi$, the sequence $\{T(h_n)\} \subset Y_\tau$ contains a Cauchy subsequence.

Second, we have to show that the operator $\frac{h}{r}\tau_0$ from X_φ to $L^2(\mathbb{R}^3)$ is compact. If $\{\frac{h_n}{r}\}$ is a bounded sequence in $H^1(\mathbb{R}^3)$ then $\{\frac{h_n}{r}\tau_0\}$ is precompact on $L_{loc}^2(\mathbb{R}^3)$, thanks to compact Sobolev embedding and, since $\tau_0(r) \rightarrow 0$ when $r \rightarrow +\infty$, we can conclude that $\{\frac{h_n}{r}\tau_0\}$ is precompact on $L^2(\mathbb{R}^3)$.

So $S(h)$ is a compact operator from X_φ on $Y_\varphi \times Y_\chi \times Y_\tau$ and $D_{\varphi,\chi,\tau}(0, \phi_0)$ is an isomorphism of $X_\varphi \times X_\chi \times X_\tau$ onto $Y_\varphi \times Y_\chi \times Y_\tau$.

In conclusion, we can apply the implicit function theorem to find that there exists $\delta > 0$ and a function $\eta \in \mathcal{C}((0, \delta), X_\varphi \times X_\chi \times X_\tau)$ such that $\eta(0) = \phi_0$ and $D(\varepsilon, \eta(\varepsilon)) = 0$ for $0 \leq \varepsilon < \delta$.

□

Acknowledgment. The author would like to thank professor Eric Séré for helpful discussions and useful comments.

REFERENCES

- [1] F. Finster, J. Smoller, S.T. Yau, *Particlelike solutions of the Einstein-Dirac equations*. Physical Review. D. Particles and Fields. Third Series **59** (1999).
- [2] E. J. Bird, *A proof of existence of particle-like solutions of Einstein Dirac Equations*. Ph.D. thesis, University of Michigan, 2005.
- [3] E. Lenzmann, *Uniqueness of ground states for pseudo-relativistic Hartree equations*. Preprint (2008).
- [4] E. H. Lieb, *Existence and Uniqueness of the Minimizing Solution of Choquard's Nonlinear Equation*. Studies in Applied Mathematics **57** (1977), 93–105.
- [5] P.L. Lions, *The Choquard equation and related questions*. Nonlinear Analysis. Theory, Methods and Applications **4** (1980), no. 6, 1063–1073.
- [6] H. Ounaies, *Perturbation method for a class of non linear Dirac equations*. Differential Integral Equations **13** (2000), no. 4-6, 707–720.
- [7] M. Renardy, R. C. Rogers, *An Introduction to Partial Differential Equations*, 337–338. Springer, 1993.
- [8] B. Thaller, *The Dirac Equation*. Springer, 1992.
- [9] M. Guan, *Solitary Wave Solutions for the Nonlinear Dirac Equations*. Preprint (2008).

CEREMADE (UMR CNRS 7534) UNIVERSITÉ PARIS-DAUPHINE
 PLACE MARÉCHAL LATTRE DE TASSIGNY
 75775 PARIS CEDEX 16 FRANCE
E-mail address: `rotanodari@ceremade.dauphine.fr`