

On $sl(3)$ KZ equations and \mathcal{W}_3 null-vector equations

Sylvain Ribault (LPTA, Montpellier)

Based mostly on the article [arxiv:0811.4587](https://arxiv.org/abs/0811.4587) [hep-th]

Families of non-rational 2d CFTs

Algebra	Target Dim.	Theory
$\widehat{sl(2)}$	3d	$SL(2, \mathbb{R})$ WZNW \sim Strings in AdS_3
$\widehat{sl(2)}$	3d	H_3^+ model
$\widehat{sl(2)/u(1)}$	2d	Strings in 2d black hole
$Vir = \mathcal{W}_2$	1d	Liouville theory $L = \partial\phi\bar{\partial}\phi + e^{b\phi}$

Algebra	Target Dim.	Theory
$\widehat{sl(3)}$	8d	$SL(3, \mathbb{R})$ WZNW
...	$2 \leq d \leq 8$...
\mathcal{W}_3	2d	Conformal $sl(3)$ Toda theory $L = (\partial\phi, \bar{\partial}\phi) + e^{b(e_1, \phi)} + e^{b(e_2, \phi)}$

Plan of the talk

- Conjectured relation between correlation functions in the $SL(N)$ WZNW model and $sl(N)$ conformal Toda theory
- Case $N = 2$ and an application
- Case $N = 3$

Correlation functions

$$\Omega_m = \left\langle \prod_{i=1}^m \Phi^{j_i}(x_i|z_i) \right\rangle^{SL(N)_k \text{ WZNW}} \quad (1)$$

- $z \in \mathbb{C}$: position on Riemann sphere
- $j \in \mathbb{C}^{N-1} = \widehat{s\ell(N)}$ spin; $x \in \mathbb{C}^{\frac{N(N-1)}{2}}$ isospin

→ order- N Knizhnik–Zamolodchikov equations

$$\tilde{\Omega}_m = \left\langle \prod_{i=1}^m V_{\alpha(j_i)}(z_i) \prod_{a=1}^{\frac{N(N-1)}{2}(m-2)} V_{\alpha_d}(y_a) \right\rangle^{s\ell(N) \text{ Toda } b=\frac{1}{\sqrt{k-N}}} \quad (2)$$

- $y \in \mathbb{C}$: position on Riemann sphere (Sklyanin variable)
- $\alpha(j) \in \mathbb{C}^{N-1} = \mathcal{W}_N$ momentum; V_{α_d} = degenerate field

The relation

Sklyanin's separation of variables = integral transformation

$$\Psi(\{x_i\}) = S \cdot \tilde{\Psi}(\{y_a\}) = \int \prod_a dy_a \ S(\{x_i\}|\{y_a\}) \tilde{\Psi}(\{y_a\}) \quad (3)$$

Twist $\Theta_m = \prod_{a < b} (y_a - y_b)^\lambda \prod_{i,a} (y_a - z_i)^\mu \prod_{i < j} (z_i - z_j)^\nu$

Conjectured relation:

$$\Omega_m = S \cdot \Theta_m \tilde{\Omega}_m$$

Status of relation	$sl(2)$ case	$sl(3)$ case
KZ-Compatible	Yes [Stoyanovsky +Feigin+Frenkel]	Only if $k = 3$
Proved in a model	Yes [SR+Teschner]	

KZ equations and the Gaudin model

Knizhnik–Zamolodchikov equations

$$\left[(k - N) \frac{\delta}{\delta z_i} + H_i \right] \Omega_m^{\text{WZNW}} = 0 \quad , \quad H_i = \sum_{\ell \neq i} \frac{\vec{D}_{(i)} \cdot \vec{D}_{(\ell)}}{z_i - z_\ell} \quad (4)$$

H_i – Gaudin Hamiltonians

$s\ell(2)$ differential operators

$$\left\{ \begin{array}{l} D_{(i)}^- = \frac{\partial}{\partial x_i} \\ D_{(i)}^3 = x_i \frac{\partial}{\partial x_i} - j_i \\ D_{(i)}^+ = x_i^2 \frac{\partial}{\partial x_i} - 2j_i x_i \end{array} \right.$$

Lax matrix

$$L(u) = \sum_{i=1}^m \frac{\vec{t} \cdot \vec{D}_{(i)}}{u - z_i}$$
$$\left\{ \begin{array}{l} u = \text{spectral parameter} \\ L(u) \in Mat_{N \times N} \end{array} \right.$$

Sklyanin's Separation of Variables

$s\ell(2)$ Lax matrix

$$L(u) = \begin{bmatrix} \sum_{i=1}^m \frac{1}{u-z_i} \left(x_i \frac{\partial}{\partial x_i} - j_i \right) & \sum_{i=1}^m \frac{1}{u-z_i} \frac{\partial}{\partial x_i} \\ \sum_{i=1}^m \frac{1}{u-z_i} \left(x_i^2 \frac{\partial}{\partial x_i} - 2j_i x_i \right) & - \sum_{i=1}^m \frac{1}{u-z_i} \left(x_i \frac{\partial}{\partial x_i} - j_i \right) \end{bmatrix}$$

Building separated variables y_a from $L(u)$:

- Define y_a as the zeroes of $B(u) = L_1^2(u)$
- Find associated momenta $p_a = A(y_a)$ where $A(u) = L_1^1(u)$
- Compute the kernel $S(\{x_i\} | \{y_a\}) = \prod_{i=1}^m e^{x_i \frac{\prod_a (z_i - y_a)}{\prod_{\ell \neq i} (z_i - z_\ell)}}$ of the integral transformation

$sl(2)$ KZ equations in Sklyanin variables

Characteristic equation [Sklyanin]

$$p_a^2 = \frac{1}{2} (L_\alpha^\beta L_\beta^\alpha)(y_a) = \sum_i \frac{1}{y_a - z_i} \left(H_i - \frac{1}{2} \frac{j_i(j_i + 1)}{y_a - z_i} \right) \quad (5)$$

Apply to $S^{-1}\Omega_m^{\text{WZNW}}$ and inject KZ equations:

$$\left[\frac{\partial^2}{\partial y_a^2} + \sum_{i=1}^m \frac{k-2}{y_a - z_i} \left(S^{-1} \frac{\delta}{\delta z_i} S + \frac{\Delta_{j_i}}{y_a - z_i} \right) \right] S^{-1}\Omega_m = 0 \quad (6)$$

$$\sum_i \frac{1}{y_a - z_i} S^{-1} \frac{\delta}{\delta z_i} S = \sum_i \frac{1}{y_a - z_i} \left[\frac{\partial}{\partial z_i} + \frac{\partial}{\partial y_a} \right] + \sum_b \frac{1}{y_a - y_b} \left[\frac{\partial}{\partial y_b} - \frac{\partial}{\partial y_a} \right]$$

→ Second order BPZ equations for $\Theta_m \tilde{\Omega}_m^{\text{Liouville}}$
with $V_{\alpha_d} = V_{-\frac{1}{2b}}$ [Stoyanovsky+Feigin+Frenkel]

A family of solvable non-rational CFTs

H_3^+ -Liouville $S \cdot \Theta_m \tilde{\Omega}_m^{\text{Liouville}} = \Omega_m^{H_3^+}$

Now if $\tilde{\Omega}_m^{(r)} \equiv \left\langle \prod_{i=1}^m V_{\alpha_i}(z_i) \prod_{a=1}^{m-2} V_{-\frac{r}{2b}}(y_a) \right\rangle \quad (r \neq 1)$

then $S \cdot \Theta_m^{(r)} \tilde{\Omega}_m^{(r)} = ?$ [SR, 2008]

Answer: Lagrangian $L^{(r)} = \partial\phi\bar{\partial}\phi + \beta\bar{\partial}\gamma + \bar{\beta}\partial\bar{\gamma} + (-\beta\bar{\beta})^r e^{2b\phi}$
with ϕ, β, γ bosons, $\Delta(\phi) = \Delta(\beta) = 0$, $\Delta(\gamma) = 1$

- Correlation functions of descendants?
- Differential equation if $r \in \mathbb{N} + b^2\mathbb{N}$?

→ extend $sl(2)$ family by a subfamily with 2 parameters (r, b)
and $c(r, b) = 3 + 6(b + b^{-1}(1 - r))^2$

$sl(3)$ Separation of Variables

$L(u) = \sum_{i=1}^m \frac{\vec{t} \cdot \vec{D}_{(i)}}{u - z_i}$: the 3×3 Lax matrix

- $B(u) = L_2^1 L_3^2 L_3^2 - L_3^2 L_3^1 L_2^2 + L_3^1 L_3^2 L_1^1 - L_1^2 L_3^1 L_3^1$
- $A(u) = -L_1^1 + \frac{L_3^1 L_1^2}{L_3^2}$
- Kernel $S(\{x_i\}, \{y_a\})$: no explicit expression

As before $B(y_a) = 0$ and $p_a = A(y_a)$

Characteristic equation:

$$\begin{aligned} p_a^3 - p_a \cdot \frac{1}{2} (L_\alpha^\beta L_\beta^\alpha)(y_a) \\ + \frac{1}{4} (L_\alpha^\beta L_\beta^\alpha)'(y_a) + \frac{1}{6} \left(L_\alpha^\beta L_\beta^\gamma L_\gamma^\alpha + L_\beta^\alpha L_\gamma^\beta L_\alpha^\gamma \right) (y_a) = 0 \quad (7) \end{aligned}$$

$sl(3)$ KZ equations in Sklyanin variables

$$\left[\frac{\partial^3}{\partial y_a^3} + \frac{\partial}{\partial y_a} \cdot \sum_{i=1}^m \frac{k-3}{y_a - z_i} S^{-1} \frac{\delta}{\delta z_i} S + \sum_{i=1}^m \frac{(k-3)\Delta_{j_i}}{(y_a - z_i)^2} \frac{\partial}{\partial y_a} \right. \\ \left. + \sum_{i=1}^m \frac{(k-3)^{\frac{3}{2}} q_{j_i} - (k-3)\Delta_{j_i}}{(y_a - z_i)^3} \right] S^{-1} \Omega_m^{\text{WZNW}} \sim 0 \quad (8)$$

Null-vector equations in $sl(3)$ Toda theory with $\alpha_d = -b\omega_1$:

$$\left[\frac{\partial^3}{\partial y_a^3} + \frac{1}{b^2} D_2 + \frac{1}{b^4} D_1 + \frac{1}{b^2} \sum_{i=1}^m \frac{\Delta_{j_i}}{(y_a - z_i)^2} \frac{\partial}{\partial y_a} \right. \\ \left. + \sum_{i=1}^m \frac{\frac{1}{b^3} q_{j_i} - \frac{1}{b^2} \Delta_{j_i}}{(y_a - z_i)^3} \right] \Theta_m \tilde{\Omega}_m^{\text{Toda}} \sim 0 \quad (9)$$

$$\frac{1}{b^2} = k - 3 , \quad \alpha(j) = -bj + b^{-1}(e_1 + e_2) , \quad j \in \mathbb{C}^2 \Leftrightarrow (\Delta_j, q_j)$$

Comparison of differential equations

- $D_2 \stackrel{?}{\sim} \frac{\partial}{\partial y_a} \cdot \sum_{i=1}^m \frac{1}{y_a - z_i} S^{-1} \frac{\delta}{\delta z_i} S$
- Problem with D_1 term where

$$D_1 = - \sum_i \frac{1}{(y_a - z_i)^2} \frac{\partial}{\partial y_a} + 3 \sum_i \frac{1}{y_a - z_i} \sum_b \frac{1}{y_a - y_b} \left(\frac{\partial}{\partial y_b} - \frac{\partial}{\partial y_a} \right) \\ + 2 \left(\sum_i \frac{1}{y_a - z_i} \right)^2 \frac{\partial}{\partial y_a} - 2 \sum_{b \neq c} \frac{1}{y_a - y_b} \frac{1}{y_b - y_c} \left(\frac{\partial}{\partial y_b} - \frac{\partial}{\partial y_a} \right) \quad (10)$$

A solution: $k \rightarrow 3 \Leftrightarrow b \rightarrow \infty$ (critical level)

An explanation: Langlands correspondence?

A speculation: $S \cdot \Theta_m \tilde{\Omega}_m^{\text{Toda}} \neq \Omega_m^{WZNW}$
but $S \cdot \Theta_m \tilde{\Omega}_m^{\text{Toda}} = \Omega_m^{\text{Some new CFT}}$