



**HAL**  
open science

## Fractal tiles associated with shift radix systems

Valerie Berthe, Anne Siegel, Wolfgang Steiner, Paul Surer, Jörg Thuswaldner

► **To cite this version:**

Valerie Berthe, Anne Siegel, Wolfgang Steiner, Paul Surer, Jörg Thuswaldner. Fractal tiles associated with shift radix systems. 2009. hal-00407999v1

**HAL Id: hal-00407999**

**<https://hal.science/hal-00407999v1>**

Preprint submitted on 28 Jul 2009 (v1), last revised 31 Aug 2010 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# FRACTAL TILES ASSOCIATED WITH SHIFT RADIX SYSTEMS

VALÉRIE BERTHÉ, ANNE SIEGEL, WOLFGANG STEINER, PAUL SURER,  
AND JÖRG M. THUSWALDNER

ABSTRACT. Shift radix systems form a collection of dynamical systems depending on a parameter  $\mathbf{r}$  which varies in the  $d$ -dimensional real vector space. They generalize well-known numeration systems such as beta-expansions, expansions with respect to rational bases, and canonical number systems. Beta-numeration and canonical number systems are known to be intimately related to fractal shapes, such as the classical Rauzy fractal and the twin dragon. These fractals turned out to be important for studying properties of expansions in several settings.

In the present paper we associate a collection of fractal tiles with shift radix systems. We show that for certain classes of parameters  $\mathbf{r}$  these tiles coincide with affine copies of the well-known tiles associated with beta-expansions and canonical number systems. On the other hand, these tiles provide natural families of tiles for beta-expansions with (non-unit) Pisot numbers as well as canonical number systems with (non-monic) expanding polynomials.

We also prove basic properties for tiles associated with shift radix systems. Indeed, we prove that under some algebraic conditions on the parameter  $\mathbf{r}$  of the shift radix system, these tiles provide multiple tilings and even tilings of the  $d$ -dimensional real vector space. These tilings turn out to have a more complicated structure than the tilings arising from the known number systems mentioned above. Such a tiling may consist of tiles having infinitely many different shapes. Moreover, the tiles need not be self-affine (or graph directed self-affine).

## 1. INTRODUCTION

**Number systems, dynamics and fractal geometry.** In the last decades, dynamical systems and fractal geometry have been proved to be deeply related to the study of number systems (see *e.g.* the survey [13]). Famous examples of fractal tiles that stem from number systems are given by the twin dragon fractal (upper left part of Figure 1 below) which is related to expansions of Gaussian integers in base  $-1 + i$  (see [26, p. 206]), or by the classical Rauzy fractal (upper right part of Figure 1) which is related to beta-expansions with respect to the Tribonacci number  $\beta$  (satisfying  $\beta^3 = \beta^2 + \beta + 1$ ; *cf.* [34, 40]). Moreover, we mention the Hokkaido tile (lower left part of Figure 1) which is related to the smallest Pisot number (see [3]) and has been studied frequently in the literature.

For several notions of number system geometric and dynamical considerations on fractals imply various non-trivial number theoretical properties. The boundary of these fractals is intimately related to the addition of 1 in the underlying number system. Moreover, the fact that the origin is an inner point of such a fractal has several implications. For instance, in beta-numeration as well as for the case of canonical number systems it implies that the underlying number system admits finite expansions (all these relations are discussed in [13]). In the case of the classical Rauzy fractal, this allows the computation of best rational simultaneous approximations of the vector  $(1, 1/\beta, 1/\beta^2)$ , where  $\beta$  is the Tribonacci number (see [17, 21]). Another example providing a relation between fractals and numeration is given by the local spiral shape of the boundary of the Hokkaido tile: by constructing a realization of the natural extension for the so-called beta-transformation, one proves that unexpected non-uniformity phenomena appear in the beta-numeration associated with

---

*Date:* July 28, 2009.

*2000 Mathematics Subject Classification.* 11A63, 28A80, 52C22.

*Key words and phrases.* Beta expansion, canonical number system, shift radix system, tiling.

This research was supported by the Austrian Science Foundation (FWF), project S9610, which is part of the national research network FWF-S96 “Analytic combinatorics and probabilistic number theory”, by the Agence Nationale de la Recherche, grant ANR-JCJC06-134288 “DyCoNum”, by the “Amadée” grant FR-13-2008 and the “PHC Amadeus” grant 17111UB.

the smallest Pisot number. Indeed, the smallest positive real number that can be approximated by rational numbers with non-purely periodic beta-expansion is an irrational number slightly smaller than  $2/3$  (see [1, 10]).

All the fractals mentioned so far are examples of the new class of tiles associated with shift radix systems which forms the main object of the present paper.



FIGURE 1. Examples of SRS tiles: The upper left tile is the so-called twin dragon fractal (cf. [26]), right beside it is the well-known Rauzy fractal associated with the Tribonacci number (cf. [34]). The lower left tile is known as “Hokkaido fractal” and corresponds to the smallest Pisot number which has minimal polynomial  $x^3 - x - 1$  (cf. [3]). The lower right one seems to be new, and is an SRS tile associated with the parameter  $\mathbf{r} = (9/10, -11/20)$ .

**Shift radix systems: a common dynamical formalism.** Shift radix systems have been proposed in [5] to unify various notions of radix expansion such as beta-expansions (see [19, 32, 35]), canonical number systems (CNS for short, see [27, 33]) and number systems with respect to rational bases (in the sense of [9]) under a common roof. Instead of starting with a base number (or base polynomial), one considers a vector  $\mathbf{r} \in \mathbb{R}^d$  and defines the mapping  $\tau_{\mathbf{r}} : \mathbb{Z}^d \rightarrow \mathbb{Z}^d$  by

$$\tau_{\mathbf{r}}(\mathbf{z}) = (z_1, \dots, z_{d-1}, -\lfloor \mathbf{r}\mathbf{z} \rfloor)^t \quad (\mathbf{z} = (z_0, \dots, z_{d-1})^t).$$

Here,  $\mathbf{r}\mathbf{z}$  denotes the scalar product of the vectors  $\mathbf{r}$  and  $\mathbf{z}$ ; moreover, for  $x$  being a real number,  $\lfloor x \rfloor$  denotes the largest integer less than or equal to  $x$ . We call  $(\mathbb{Z}^d, \tau_{\mathbf{r}})$  a *shift radix system* (SRS, for short). A vector  $\mathbf{r}$  gives rise to an SRS with *finiteness property* if each integer vector  $\mathbf{z} \in \mathbb{Z}^d$  can be finitely expanded with respect to the vector  $\mathbf{r}$ , that is, if for each  $\mathbf{z} \in \mathbb{Z}^d$  there is an  $n \in \mathbb{N}$  such that the  $n$ -th iterate of  $\tau_{\mathbf{r}}$  satisfies  $\tau_{\mathbf{r}}^n(\mathbf{z}) = \mathbf{0}$ .

In the papers written on SRS so far (see *e.g.* [5, 6, 7, 8, 39]), relations between SRS and well-known notions of number system such as beta-expansions with respect to unit Pisot numbers and canonical number systems with a monic expanding polynomial have been established (we will give a short account on these relations in Section 5 and 6, respectively). In particular, SRS turned out to be a fruitful tool in order to deal with the problem of finite representations in these number systems.

**SRS tiles: an extension of arithmetically meaningful known constructions.** In the present paper, we study geometric properties of SRS: we introduce a family of tiles for each  $\mathbf{r} \in \mathbb{R}^d$  with contractive and regular companion matrix, and prove geometric properties of these tiles. Figure 1 shows four examples of such tiles, called *SRS tiles*.

We prove that our definition unifies the notions of self-affine tiles known for CNS with respect to monic polynomial (see [13, 25]) and beta-numeration related to a unit Pisot number (see [3]).

**Result 1.** *The following relations between SRS tiles and classical tiles related to number systems hold:*

- *If the parameter  $\mathbf{r}$  of an SRS is related to a monic expanding polynomial over  $\mathbb{Z}$ , the SRS tile is a linear image of the self-affine tile associated with this polynomial.*
- *If the parameter  $\mathbf{r}$  of an SRS comes from a unit Pisot number, the SRS tile is a linear image of the central tile associated with the corresponding beta-numeration.*

It is well-known that these classical tiles are self-affine and associated with (multiple) tilings that are highly structured (see *e.g.* the surveys [12, 16]). Indeed, these tiles satisfy a set equation expressed as a graph-directed iterated function system (GIFS) in the sense of Mauldin and Williams [31] which means that each tile can be decomposed with respect to the action of an affine mapping into several copies of a finite number of tiles, with the decomposition being produced by a finite graph (see [13]).

In the present SRS situation, we prove that our construction gives rise to new classes of tiles, in particular we want to emphasize on tiles related to a non-monic expanding polynomial or to a non-unit Pisot number. In both cases, we are actually able to extend the usual definition of tiles used for unit Pisot numbers and monic polynomials. The tiles defined in the usual way are not satisfactory in this more general setting, since they always produce overlaps in their self-affine decomposition. A first strategy to remove overlaps consists in enlarging the space of representation by adding arithmetic components ( $p$ -adic factors) as proposed in [4]. However, such tiles are of limited topological importance since they have totally disconnected factors.

We prove that our construction, however, allows to insert arithmetic criteria in the construction of the tiles: roughly speaking, the SRS mapping  $\tau_{\mathbf{r}}$  naturally selects points in appropriate submodules. This arithmetic selection process removes the overlaps.

**Result 2.** *SRS provide a natural collection of tiles for number systems related to non-monic expanding polynomials as well as to non-unit Pisot numbers.*

Nonetheless, the geometrical structure of these tiles is much more complicated than the structure of the classical ones. As we shall illustrate for  $\mathbf{r} = (-2/3)$ , there may be infinitely many shapes of tiles associated with certain parameters  $\mathbf{r}$ . Actually, the description of SRS tiles requires a set equation that cannot be captured by a finite graph. Equation (3.3) suggests that an infinite hierarchy of set equations is needed to describe an SRS tile. Therefore, SRS tiles in general cannot be regarded as GIFS attractors. Furthermore, an SRS tile is not always equal to the closure of its interior (Example 3.12 exhibits a case of an SRS tile that is equal to a single point). Also, due to the lack of a GIFS structure, we have no information on the measure of the boundary of tiles.

**Tiling properties.** Despite of their complicated geometrical structure, we are able to prove tiling properties for SRS tiles. We first prove that, for each fixed parameter  $\mathbf{r}$ , the associated tiles form a covering, extending the results known for unit Pisot numbers and monic expanding polynomials. In the classical cases, exhibiting a multiple tiling (that is, a covering with an almost everywhere constant degree) from this covering is usually done by exploiting specific features of the tiling together with the GIFS structure of the tiles: this allows to transfer local information to the whole space in order to obtain global information. The finiteness property is then used to prove that  $\mathbf{0}$  belongs to only one tile, leading to a tiling. In the present SRS situation, however, this strategy does not work any more. The finiteness property is still equivalent to the fact that  $\mathbf{0}$  belongs to exactly one SRS tile, but the fact that SRS tiles are no longer GIFS attractors prevents us from spreading this local information to the whole space. In order to get global information, we impose additional algebraic conditions on the parameter  $\mathbf{r}$ , and we use these conditions to

exhibit a dense set of points that are proved to belong to a fixed number of tiles. This leads to the multiple tiling property.

Our main result can thus be stated as follows (for the corresponding definitions and for a more precise statement, see Section 4, in particular Theorem 4.6 and Corollary 4.7):

**Result 3.** *Let  $\mathbf{r}$  be an SRS parameter with contractive and regular companion matrix, and assume that either  $\mathbf{r} \in \mathbb{Q}^d$  or  $\mathbf{r}$  is related to a Pisot number or  $\mathbf{r}$  has algebraically independent coordinates. Then the following assertions hold.*

- *The collection of SRS tiles associated with the parameter  $\mathbf{r}$  forms a weak multiple tiling.*
- *If the finiteness property is satisfied, then the collection of SRS tiles forms a weak tiling.*

By weak tiling, we mean here that the tiles cover the whole space and have disjoint interiors. Stating a “strong” tiling property would require to have information on the boundaries of the tiles, which is deeply intricate if  $d \geq 2$  since the tiles are no longer GIFS attractors, and deserves a specific study. For  $d = 1$  the situation becomes easier. Indeed, we will prove in Theorem 4.9 that in this case SRS tiles are (possibly degenerate) intervals which form a tiling of  $\mathbb{R}$ .

**Structure of the paper.** In Section 2, we introduce a way of representing integer vectors by using the shift radix transformation  $\tau_{\mathbf{r}}$ . In Section 3, SRS tiles are defined and fundamental geometric properties of them are studied. Section 4 is devoted to tiling properties of SRS tiles. We show that SRS tiles form tilings or multiple tilings for large classes of parameters  $\mathbf{r} \in \mathbb{R}^d$ . As the tilings are no longer self-affine we have to use new methods in our proofs. In Section 5, we analyse the relation between tiles associated with expanding polynomials and SRS tiles more closely. We prove that the SRS tiles given by a monic CNS parameter  $\mathbf{r}$  coincide up to a linear transformation with the self-affine CNS tile. For non-monic expanding polynomials, SRS tiles give rise to a new class of tiles. At the end of this section, we prove for the parameter  $\mathbf{r} = (-2/3)$  that the associated tiling has infinitely many shapes of tiles. In Section 6, after proving that the shift radix transformation is conjugate to the beta-transformation restricted to  $\mathbb{Z}[\beta]$ , we investigate the relation between beta-tiles and SRS tiles. It turns out that beta-tiles associated with a unit Pisot number are linear images of SRS tiles related to a parameter associated with this Pisot number. Moreover, we define a new class of tiles associated with non-unit Pisot numbers.

## 2. THE SRS REPRESENTATION

**2.1. Definition of shift radix systems and their parameter domains.** Shift radix systems are dynamical systems defined on  $\mathbb{Z}^d$  as follows (see [5]).

**Definition 2.1** (Shift radix system, finiteness property). For  $\mathbf{r} = (r_0, \dots, r_{d-1}) \in \mathbb{R}^d$ ,  $d \geq 1$ , set

$$\begin{aligned} \tau_{\mathbf{r}} : \mathbb{Z}^d &\rightarrow \mathbb{Z}^d, \\ \mathbf{z} = (z_0, z_1, \dots, z_{d-1})^t &\mapsto (z_1, \dots, z_{d-1}, -\lfloor \mathbf{r}\mathbf{z} \rfloor)^t, \end{aligned}$$

where  $\mathbf{r}\mathbf{z}$  denotes the scalar product of  $\mathbf{r}$  and  $\mathbf{z}$ . We call the dynamical system  $(\mathbb{Z}^d, \tau_{\mathbf{r}})$  a *shift radix system* (SRS, for short). The SRS parameter  $\mathbf{r}$  is said to be *reduced* if  $r_0 \neq 0$ .

We say that  $(\mathbb{Z}^d, \tau_{\mathbf{r}})$  satisfies the *finiteness property* if for every  $\mathbf{z} \in \mathbb{Z}^d$  there exists some  $n \in \mathbb{N}$  such that  $\tau_{\mathbf{r}}^n(\mathbf{z}) = \mathbf{0}$ .

*Remark 2.2.* If  $r_0 = 0$ , then every vector  $\tau_{\mathbf{r}}^n(\mathbf{z})$ ,  $\mathbf{z} \in \mathbb{Z}^d$ ,  $n \geq 1$ , can be easily obtained from the SRS  $(\mathbb{Z}^{d-1}, \tau_{\mathbf{r}'})$  with  $\mathbf{r}' = (r_1, \dots, r_{d-1})$  since  $\tau_{\mathbf{r}'} \circ \pi = \pi \circ \tau_{\mathbf{r}}$ , where  $\pi : \mathbb{Z}^d \rightarrow \mathbb{Z}^{d-1}$  denotes the projection defined by  $\pi(z_0, z_1, \dots, z_{d-1}) = (z_1, \dots, z_{d-1})$ .

The companion matrix of  $\mathbf{r} = (r_0, \dots, r_{d-1})$  is denoted by

$$M_{\mathbf{r}} := \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ \vdots & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ 0 & 0 & \cdots & 0 & 1 \\ -r_0 & -r_1 & \cdots & -r_{d-2} & -r_{d-1} \end{pmatrix} \in \mathbb{R}^{d \times d}.$$

It is regular if  $r_0 \neq 0$ . We will work with  $M_{\mathbf{r}}^{-1}$  (see *e.g.* Proposition 2.6), hence we will assume  $\mathbf{r}$  reduced in all that follows. For  $\mathbf{z} = (z_0, \dots, z_{d-1})^t$ , note that

$$M_{\mathbf{r}}\mathbf{z} = (z_1, \dots, z_{d-1}, -\mathbf{r}\mathbf{z})^t$$

and thus

$$(2.1) \quad \tau_{\mathbf{r}}(\mathbf{z}) = M_{\mathbf{r}}\mathbf{z} + (0, \dots, 0, \{\mathbf{r}\mathbf{z}\})^t,$$

where  $\{x\} = x - \lfloor x \rfloor$  denotes the fractional part of  $x$ .

The sets

$$\begin{aligned} \mathcal{D}_d &:= \{\mathbf{r} \in \mathbb{R}^d \mid (\tau_{\mathbf{r}}^n(\mathbf{z}))_{n \in \mathbb{N}} \text{ is eventually periodic for all } \mathbf{z} \in \mathbb{Z}^d\} \quad \text{and} \\ \mathcal{D}_d^{(0)} &:= \{\mathbf{r} \in \mathbb{R}^d \mid \tau_{\mathbf{r}} \text{ is an SRS with finiteness property}\} \end{aligned}$$

are intimately related to the notion of SRS. Obviously we have  $\mathcal{D}_d^{(0)} \subset \mathcal{D}_d$ . Apart from its boundary, the set  $\mathcal{D}_d$  is easy to describe.

**Lemma 2.3** (see [5]). *A point  $\mathbf{r} \in \mathbb{R}^d$  is contained in the interior of  $\mathcal{D}_d$  if and only if the spectral radius of the companion matrix  $M_{\mathbf{r}}$  generated by  $\mathbf{r}$  is strictly less than 1.*

The shape of  $\mathcal{D}_d^{(0)}$  is of a more complicated nature. While  $\mathcal{D}_1^{(0)} = [0, 1)$  is quite easy to describe (*cf.* [5, Proposition 4.4]), already for  $d = 2$  no complete description of  $\mathcal{D}_2^{(0)}$  is known. The available partial results (see [6, 39]) indicate that  $\mathcal{D}_d^{(0)}$  is of a quite irregular structure for  $d \geq 2$ . Nonetheless, an algorithm is given in [5, Proposition 4.4] which decides whether a given  $\mathbf{r}$  belongs to  $\mathcal{D}_d^{(0)}$ . This algorithm was used to draw the approximation of  $\mathcal{D}_2^{(0)}$  depicted in Figure 2.




FIGURE 2. An approximation of  $\mathcal{D}_2^{(0)}$ .

Apart from the easier case  $d = 1$ , in this paper, we are going to consider three classes of points  $\mathbf{r}$  assumed to be reduced which belong to  $\mathcal{D}_d$ . The first two classes are dense in  $\text{int}(\mathcal{D}_d)$ , while the third one has full measure in  $\text{int}(\mathcal{D}_d)$ .

- (1)  $\mathbf{r} \in \mathbb{Q}^d \cap \text{int}(\mathcal{D}_d)$ . This class includes the parameters  $\mathbf{r} = \left(\frac{1}{a_0}, \frac{a_{d-1}}{a_0}, \dots, \frac{a_1}{a_0}\right) \in \text{int}(\mathcal{D}_d)$  with  $a_0, \dots, a_{d-1} \in \mathbb{Z}$ , which correspond to expansions with respect to monic expanding polynomials, including CNS. If the first coordinate of  $\mathbf{r}$  has numerator greater than one, this extends to non-monic polynomials in a natural way (see Section 5 for details).
- (2)  $\mathbf{r} = (r_0, \dots, r_{d-1})$  is obtained by decomposing the minimal polynomial of a Pisot number  $\beta$  as  $(x - \beta)(x^d + r_{d-1}x^{d-1} + \dots + r_0)$ . In view of Lemma 2.3, this implies that  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$ , and by [8] this set of parameters is dense in  $\text{int}(\mathcal{D}_d)$ . These parameters correspond to beta-numeration with respect to Pisot numbers (see Section 6). Note that even non-unit Pisot numbers are covered here.
- (3)  $\mathbf{r} = (r_0, \dots, r_{d-1}) \in \text{int}(\mathcal{D}_d)$  with algebraically independent coordinates  $r_0, \dots, r_{d-1}$ .

**2.2. SRS representation of  $d$ -dimensional integer vectors.** For  $\mathbf{r} \in \mathbb{R}^d$  we can use the SRS transformation  $\tau_{\mathbf{r}}$  to define an expansion for  $d$ -dimensional integer vectors.

**Definition 2.4** (SRS representation). Let  $\mathbf{r} \in \mathbb{R}^d$ . For  $\mathbf{z} \in \mathbb{Z}^d$ , the *SRS representation* of  $\mathbf{z}$  with respect to  $\mathbf{r}$  is defined to be the sequence  $(v_1, v_2, v_3, \dots)$ , with  $v_n = \{\mathbf{r}\tau_{\mathbf{r}}^{n-1}(\mathbf{z})\}$  for all  $n \geq 1$ .

The representation is said to be *finite* if there is an  $n_0$  such that  $v_n = 0$  for all  $n \geq n_0$ . It is said to be *eventually periodic* if there are  $n_0, p$  such that  $v_n = v_{n+p}$  for all  $n \geq n_0$ .

By definition, every SRS representation is finite if  $\mathbf{r} \in \mathcal{D}_d^{(0)}$ , and every SRS representation is eventually periodic if  $\mathbf{r} \in \mathcal{D}_d$ . The following simple properties of SRS representations show that integer vectors can be expanded according to  $\tau_{\mathbf{r}}$ . The matrix  $M_{\mathbf{r}}$  acts as a base and the vectors  $(0, \dots, 0, v_j)$  are the digits.

**Lemma 2.5.** *Let  $\mathbf{r} \in \mathbb{R}^d$  and  $(v_1, v_2, \dots)$  be the SRS representation of  $\mathbf{z} \in \mathbb{Z}^d$  with respect to  $\mathbf{r}$ . Then the following properties hold for all  $n \in \mathbb{N}$ :*

- (1)  $0 \leq v_n < 1$ ,
- (2)  $\tau_{\mathbf{r}}^n(\mathbf{z})$  has the SRS representation  $(v_{n+1}, v_{n+2}, v_{n+3}, \dots)$ ,
- (3) we have

$$(2.2) \quad M_{\mathbf{r}}^n \mathbf{z} = \tau_{\mathbf{r}}^n(\mathbf{z}) - \sum_{j=1}^n M_{\mathbf{r}}^{n-j} (0, \dots, 0, v_j)^t.$$

*Proof.* Assertions (1) and (2) follow immediately from the definition of the SRS representation. By iterating (2.1), we obtain

$$\tau_{\mathbf{r}}^n(\mathbf{z}) = M_{\mathbf{r}}^n \mathbf{z} + \sum_{j=1}^n M_{\mathbf{r}}^{n-j} (0, \dots, 0, \{\mathbf{r}\tau_{\mathbf{r}}^{j-1}(\mathbf{z})\})^t,$$

which yields (2.2). □

Note that the set of possible SRS digits  $(0, \dots, 0, v)$  is infinite unless  $\mathbf{r} \in \mathbb{Q}^d$ .

We prove that the SRS representation is unique in the following sense.

**Proposition 2.6.** *Let  $\mathbf{r} \in \mathbb{R}^d$  be reduced and suppose that the SRS-representation of an element  $\mathbf{z}_0 \in \mathbb{Z}^d$  is  $(v_1, v_2, v_3, \dots)$ . Assume that for some reals  $v_0, v_{-1}, \dots, v_{-n+1} \in [0, 1)$ ,  $n \in \mathbb{N}$ , we have*

$$\mathbf{z}_{-k} := M_{\mathbf{r}}^{-k} \left( \mathbf{z}_0 - \sum_{j=0}^{k-1} M_{\mathbf{r}}^j (0, \dots, 0, v_{-j})^t \right) \in \mathbb{Z}^d \quad \text{for all } 1 \leq k \leq n.$$

*Then  $\tau_{\mathbf{r}}^n(\mathbf{z}_{-n}) = \mathbf{z}_0$  and  $\mathbf{z}_{-n}$  has the SRS representation  $(v_{-n+1}, v_{-n+2}, v_{-n+3}, \dots)$ .*

*Proof.* The assertion is obviously true for  $n = 0$ . Now continue by induction on  $n$ . We have

$$\tau_{\mathbf{r}}(\mathbf{z}_{-n}) = \tau_{\mathbf{r}}(M_{\mathbf{r}}^{-1}(\mathbf{z}_{-n+1} - (0, \dots, 0, v_{-n+1})^t)) = \mathbf{z}_{-n+1} - (0, \dots, 0, v_{-n+1})^t + (0, \dots, 0, \{\mathbf{r}\mathbf{z}_{-n}\})^t.$$

Since  $\tau_{\mathbf{r}}(\mathbf{z}_{-n}) \in \mathbb{Z}^d$ , we obtain  $\tau_{\mathbf{r}}(\mathbf{z}_{-n}) = \mathbf{z}_{-n+1}$  and  $v_{-n+1} = \{\mathbf{r}\mathbf{z}_{-n}\}$ . Therefore the first SRS digit of  $\mathbf{z}_{-n}$  is  $v_{-n+1}$ . By induction, we have  $\tau_{\mathbf{r}}^n(\mathbf{z}_{-n}) = \mathbf{z}_0$ , hence, the SRS representation of  $\mathbf{z}_{-n}$  is  $(v_{-n+1}, v_{-n+2}, v_{-n+3}, \dots)$ . □

## 3. DEFINITION AND FIRST PROPERTIES OF SRS TILES

We now define a new type of tiles based on the mapping  $\tau_{\mathbf{r}}$  when the matrix  $M_{\mathbf{r}}$  is contractive and  $\mathbf{r}$  is reduced. By analogy with the definition of tiles for other dynamical systems (see *e.g.* [2, 16, 25, 34, 37, 40]), we consider elements of  $\mathbb{Z}^d$  which are mapped to a given  $\mathbf{x} \in \mathbb{Z}^d$  by  $\tau_{\mathbf{r}}^n$ , renormalize by a multiplication with  $M_{\mathbf{r}}^n$ , and let  $n$  tend to  $\infty$ . To build this set, we thus consider vectors whose SRS expansion coincides with the expansion of  $\mathbf{x}$  up to an added finite prefix and we then renormalize this expansion. We will see in Sections 5 and 6 that some of these tiles are related to well-known types of tiles, namely CNS tiles and beta-tiles. We recall that four examples of central SRS tiles are depicted in Figure 1.

**3.1. Definition of SRS tiles.** An SRS tile will turn out to be the limit of the sequence of compact sets  $(M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x}))_{n \geq 0}$  with respect to the Hausdorff metric. As it is *a priori* not clear that this limit exists, we first define the tiles as lower Hausdorff limits of these sets and then show that the Hausdorff limit exists. Recall that the lower Hausdorff limit  $\text{Li}_{n \rightarrow \infty} A_n$  of a sequence  $(A_n)$  of subsets of  $\mathbb{R}^d$  is the (closed) set of all  $\mathbf{t} \in \mathbb{R}^d$  having the property that each neighborhood of  $\mathbf{t}$  intersects  $A_n$  provided that  $n$  is sufficiently large. If the sets  $A_n$  are compact and  $(A_n)$  is a Cauchy sequence w.r.t. the Hausdorff metric  $\delta$ , then the Hausdorff limit  $\text{Lim}_{n \rightarrow \infty} A_n$  exists and

$$\text{Lim}_{n \rightarrow \infty} A_n = \text{Li}_{n \rightarrow \infty} A_n$$

(see [28, Chapter II, §29] for details on Hausdorff limits).

**Definition 3.1** (SRS tile). Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced and  $\mathbf{x} \in \mathbb{Z}^d$ . The *SRS tile* associated with  $\mathbf{r}$  is defined as

$$\mathcal{T}_{\mathbf{r}}(\mathbf{x}) := \text{Li}_{n \rightarrow \infty} M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x}).$$

*Remark 3.2.* Note that this means that  $\mathbf{t} \in \mathbb{R}^d$  is an element of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  if and only if there exist vectors  $\mathbf{z}_{-n} \in \mathbb{Z}^d$ ,  $n \in \mathbb{N}$ , such that  $\tau_{\mathbf{r}}^n(\mathbf{z}_{-n}) = \mathbf{x}$  for all  $n \in \mathbb{N}$  and  $\lim_{n \rightarrow \infty} M_{\mathbf{r}}^n \mathbf{z}_{-n} = \mathbf{t}$ .

We will see in Theorem 3.5 that the lower limit in Definition 3.1 is equal to the limit with respect to the Hausdorff metric.

**3.2. Compactness, a set equation, and a covering property of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$ .** In this subsection we will show that each SRS tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is a compact set that can be decomposed into subtiles which are obtained by multiplying other tiles by  $M_{\mathbf{r}}$ . Moreover, we prove that for each reduced  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  covers the real vector space  $\mathbb{R}^d$ .

If  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$ , then the matrix  $M_{\mathbf{r}}$  is contractive by Lemma 2.3. Let  $\rho < 1$  be the spectral radius of  $M_{\mathbf{r}}$ ,  $\rho < \tilde{\rho} < 1$  and  $\|\cdot\|$  be a norm satisfying

$$\|M_{\mathbf{r}} \mathbf{x}\| \leq \tilde{\rho} \|\mathbf{x}\| \quad \text{for all } \mathbf{x} \in \mathbb{R}^d$$

(for the construction of such a norm see for instance [29, Equation (3.2)]). Then we have

$$(3.1) \quad R := \sum_{n=0}^{\infty} \|M_{\mathbf{r}}^n(0, \dots, 0, 1)^t\| \leq \frac{\|(0, \dots, 0, 1)^t\|}{1 - \tilde{\rho}}.$$

**Lemma 3.3.** *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced. Then every  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$ ,  $\mathbf{x} \in \mathbb{Z}^d$ , is contained in the closed ball of radius  $R$  with center  $\mathbf{x}$ . Therefore the cardinality of the sets  $\{\mathbf{x} \in \mathbb{Z}^d \mid \mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})\}$  is uniformly bounded in  $\mathbf{t} \in \mathbb{R}^d$ . Furthermore, the family of SRS tiles  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is locally finite, that is, any open ball meets only a finite number of tiles of the family.*

*Proof.* Let  $\mathbf{x} \in \mathbb{Z}^d$ ,  $\mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$  and  $\mathbf{z}_{-n}$  as in Remark 3.2. Let the SRS representation of  $\mathbf{z}_{-n}$  be  $(v_{-n+1}^{(n)}, v_{-n+2}^{(n)}, v_{-n+3}^{(n)}, \dots)$ . Then by (2.2) we have

$$M_{\mathbf{r}}^n \mathbf{z}_{-n} = \mathbf{x} - \sum_{j=1}^n M_{\mathbf{r}}^{n-j}(0, \dots, 0, v_{-n+j}^{(n)})^t,$$

thus  $\|M_{\mathbf{r}}^n \mathbf{z}_{-n} - \mathbf{x}\| < R$  and, hence,  $\|\mathbf{t} - \mathbf{x}\| \leq R$ . The uniform boundedness of the cardinality of  $\{\mathbf{x} \in \mathbb{Z}^d \mid \mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})\}$  and the local finiteness follow immediately.  $\square$

**Lemma 3.4.** *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced and denote by  $\delta(\cdot, \cdot)$  the Hausdorff metric induced by a norm satisfying (3.1). Then  $(M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x}))_{n \geq 0}$  is a Cauchy sequence w.r.t.  $\delta$ , in particular,*

$$\delta(M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x}), M_{\mathbf{r}}^{n+1} \tau_{\mathbf{r}}^{-n-1}(\mathbf{x})) \leq \tilde{\rho}^n \|(0, \dots, 0, 1)^t\|.$$

*Proof.* Let  $\mathbf{t} \in M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x})$ . Then

$$(3.2) \quad M_{\mathbf{r}}^{n+1} \tau_{\mathbf{r}}^{-1}(M_{\mathbf{r}}^{-n} \mathbf{t}) \subset M_{\mathbf{r}}^{n+1} \tau_{\mathbf{r}}^{-n-1}(\mathbf{x}).$$

Note that  $\tau_{\mathbf{r}}$  is surjective since  $0 < |r_0| = |\det M_{\mathbf{r}}| < 1$ . Thus there exists some  $\mathbf{t}' \in \tau_{\mathbf{r}}^{-1}(M_{\mathbf{r}}^{-n} \mathbf{t})$ . By the definition of  $\tau_{\mathbf{r}}$ , there is a  $\mathbf{v} = (0, \dots, 0, v)$  with  $v \in [0, 1)$  such that  $\tau_{\mathbf{r}}(\mathbf{t}') = M_{\mathbf{r}} \mathbf{t}' + \mathbf{v}$ . Now we have  $M_{\mathbf{r}} \mathbf{t}' + \mathbf{v} = M_{\mathbf{r}}^{-n} \mathbf{t}$ . Using (3.2), this gives  $\mathbf{t} - M_{\mathbf{r}}^n \mathbf{v} = M_{\mathbf{r}}^{n+1} \mathbf{t}' \in M_{\mathbf{r}}^{n+1} \tau_{\mathbf{r}}^{-n-1}(\mathbf{x})$ .

On the other hand, let  $\mathbf{t} \in M_{\mathbf{r}}^{n+1} \tau_{\mathbf{r}}^{-n-1}(\mathbf{x})$ . Then  $M_{\mathbf{r}}^n \tau_{\mathbf{r}}(M_{\mathbf{r}}^{-n-1} \mathbf{t}) \in M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x})$ . As there exists a  $\mathbf{v} = (0, \dots, 0, v)$  with  $v \in [0, 1)$  such that  $\tau_{\mathbf{r}}(M_{\mathbf{r}}^{-n-1} \mathbf{t}) = M_{\mathbf{r}}^{-n} \mathbf{t} + \mathbf{v}$ , we conclude that  $\mathbf{t} + M_{\mathbf{r}}^n \mathbf{v} \in M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x})$ .

Since  $\|M_{\mathbf{r}}^n \mathbf{v}\| \leq \tilde{\rho}^n \|(0, \dots, 0, 1)^t\|$  we are done.  $\square$

**Theorem 3.5** (Basic properties of SRS tiles). *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced and  $\mathbf{x} \in \mathbb{Z}^d$ . The SRS tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  can be written as*

$$\mathcal{T}_{\mathbf{r}}(\mathbf{x}) = \text{Lim}_{n \rightarrow \infty} M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x})$$

where  $\text{Lim}$  denotes the limit w.r.t. the Hausdorff metric  $\delta$ . It is a non-empty compact set that satisfies the set equation

$$(3.3) \quad \mathcal{T}_{\mathbf{r}}(\mathbf{x}) = \bigcup_{\mathbf{y} \in \tau_{\mathbf{r}}^{-1}(\mathbf{x})} M_{\mathbf{r}} \mathcal{T}_{\mathbf{r}}(\mathbf{y}).$$

*Proof.* The fact that the Hausdorff limit  $\text{Lim}_{n \rightarrow \infty} M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x})$  exists and equals  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  follows from Lemma 3.4. Moreover,  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is closed since Hausdorff limits are closed by definition. As it is also bounded by Lemma 3.3 the compactness of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is shown. The fact that  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is non-empty follows from the surjectivity of  $\tau_{\mathbf{r}}$ . It remains to prove the set equation. This follows from

$$\begin{aligned} \mathcal{T}_{\mathbf{r}}(\mathbf{x}) &= \text{Lim}_{n \rightarrow \infty} M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{x}) = M_{\mathbf{r}} \text{Lim}_{n \rightarrow \infty} \bigcup_{\mathbf{y} \in \tau_{\mathbf{r}}^{-1}(\mathbf{x})} M_{\mathbf{r}}^{n-1} \tau_{\mathbf{r}}^{-n+1}(\mathbf{y}) \\ &= M_{\mathbf{r}} \bigcup_{\mathbf{y} \in \tau_{\mathbf{r}}^{-1}(\mathbf{x})} \text{Lim}_{n \rightarrow \infty} M_{\mathbf{r}}^{n-1} \tau_{\mathbf{r}}^{-n+1}(\mathbf{y}) = M_{\mathbf{r}} \bigcup_{\mathbf{y} \in \tau_{\mathbf{r}}^{-1}(\mathbf{x})} \mathcal{T}_{\mathbf{r}}(\mathbf{y}). \end{aligned} \quad \square$$

The points in an SRS tile are characterized by the following proposition.

**Proposition 3.6.** *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced and  $\mathbf{x} \in \mathbb{Z}^d$ . Then  $\mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$  if and only if there exist some numbers  $v_{-j} \in [0, 1)$ ,  $j \in \mathbb{N}$ , such that*

$$(3.4) \quad \mathbf{t} = \mathbf{x} - \sum_{j=0}^{\infty} M_{\mathbf{r}}^j (0, \dots, 0, v_{-j})^t$$

and

$$(3.5) \quad M_{\mathbf{r}}^{-n} \left( \mathbf{x} - \sum_{j=0}^{n-1} M_{\mathbf{r}}^j (0, \dots, 0, v_{-j})^t \right) \in \mathbb{Z}^d \quad \text{for all } n \in \mathbb{N}.$$

Set  $\mathbf{z}_{-n} := M_{\mathbf{r}}^{-n} \left( \mathbf{x} - \sum_{j=0}^{n-1} M_{\mathbf{r}}^j (0, \dots, 0, v_{-j})^t \right)$ . Then condition (3.5) is equivalent to  $\mathbf{z}_0 = \mathbf{x}$ ,

$$(3.6) \quad \tau_{\mathbf{r}}(\mathbf{z}_{-n}) = \mathbf{z}_{-n+1} \quad \text{and} \quad v_{-n+1} = \{\mathbf{r} \mathbf{z}_{-n}\} \quad \text{for all } n \geq 1.$$

*Proof.* The equivalence of (3.5) and (3.6) follows from Proposition 2.6. If these conditions hold and  $\mathbf{t}$  is defined by (3.4), then it is clear from Remark 3.2 that  $\mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$ .

Now let  $\mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$ . We show that we can choose the  $\mathbf{z}_{-n}$  given in Remark 3.2 such that  $\tau_{\mathbf{r}}(\mathbf{z}_{-n}) = \mathbf{z}_{-n+1}$  for all  $n \geq 1$ . Let  $\mathbf{z}_0 = \mathbf{x}$ . By (3.3), there is some  $\mathbf{z}_{-1} \in \tau_{\mathbf{r}}^{-1}(\mathbf{z}_0)$  with  $M_{\mathbf{r}}^{-1} \mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{z}_{-1})$ , and inductively  $\mathbf{z}_{-n} \in \tau_{\mathbf{r}}^{-1}(\mathbf{z}_{-n+1})$  with  $M_{\mathbf{r}}^{-n} \mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{z}_{-n})$  for all  $n \geq 1$ . By Lemma 3.3, we have  $\|M_{\mathbf{r}}^{-n} \mathbf{t} - \mathbf{z}_{-n}\| \leq R$ , thus  $\lim_{n \rightarrow \infty} M_{\mathbf{r}}^n \mathbf{z}_{-n} = \mathbf{t}$ , and  $\mathbf{t}$  is of the form (3.4) with  $v_{-n+1} = \{\mathbf{r} \mathbf{z}_{-n}\}$ .  $\square$

It remains to show the covering property.

**Proposition 3.7.** *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced. The family of SRS tiles  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is a covering of  $\mathbb{R}^d$ , i.e.,*

$$\bigcup_{\mathbf{x} \in \mathbb{Z}^d} \mathcal{T}_{\mathbf{r}}(\mathbf{x}) = \mathbb{R}^d.$$

*Proof.* Set  $\mathcal{C} = \bigcup_{\mathbf{x} \in \mathbb{Z}^d} \mathcal{T}_{\mathbf{r}}(\mathbf{x})$ . By Lemma 3.3 and the non-emptiness of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$ , the set  $\mathcal{C}$  is relatively dense in  $\mathbb{R}^d$ . By (3.3), we have  $M_{\mathbf{r}}\mathcal{C} \subseteq \mathcal{C}$ . As  $M_{\mathbf{r}}$  is contractive, this implies that  $\mathcal{C}$  is dense in  $\mathbb{R}^d$ . We conclude by noticing that the SRS tiles are compact by Theorem 3.5 and that the family of SRS tiles  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is locally finite, according to Lemma 3.3.  $\square$

**3.3. Around the origin.** The tile associated with  $\mathbf{0}$  plays a specific role.

**Definition 3.8** (Central SRS tile). Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced. The tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{0})$  is called *central SRS tile* associated with  $\mathbf{r}$ .

Since  $\tau_{\mathbf{r}}(\mathbf{0}) = \mathbf{0}$  for every  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$ , the origin is an element of the central tile. However, in general it can be contained in finitely many other tiles of the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ . Whether or not  $\mathbf{0}$  is contained exclusively in the central tile plays an important role in numeration. Indeed, for beta-numeration,  $\mathbf{0}$  is contained exclusively in the central beta-tile (see Definition 6.5 below) if and only if the so-called finiteness property (F) is satisfied (see [3, 19]). An analogous criterion holds for CNS (cf. [11]). Now, we show that this characterizes also the SRS with finiteness property.

**Definition 3.9** (Purely periodic point). Let  $\mathbf{r} \in \mathcal{D}_d$ . A point  $\mathbf{z} \in \mathbb{Z}^d$  is called *purely periodic* if  $\tau_{\mathbf{r}}^p(\mathbf{z}) = \mathbf{z}$  for some  $p \geq 1$ .

**Theorem 3.10.** *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced and  $\mathbf{x} \in \mathbb{Z}^d$ . Then  $\mathbf{0} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$  if and only if  $\mathbf{x}$  is purely periodic. There are only finitely many purely periodic points.*

*Proof.* We first show that, if  $\mathbf{x}$  is purely periodic with period  $p$ , then  $\mathbf{0} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$ . We have  $\tau_{\mathbf{r}}^p(\mathbf{x}) = \mathbf{x}$ . Therefore  $\mathbf{x} \in \tau_{\mathbf{r}}^{-kp}(\mathbf{x})$  for all  $k \in \mathbb{N}$ , and since  $M_{\mathbf{r}}$  is contractive we gain

$$\mathbf{0} = \lim_{k \rightarrow \infty} M_{\mathbf{r}}^{kp} \mathbf{x} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x}).$$

To prove the other direction, let  $\mathbf{0} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$ ,  $\mathbf{x} \in \mathbb{Z}^d$ . Let  $\mathbf{z}_{-n} \in \tau_{\mathbf{r}}^{-n}(\mathbf{x})$  be defined as in Proposition 3.6, with  $\mathbf{t} = \mathbf{0}$ . Multiplying (3.4) by  $M_{\mathbf{r}}^{-n}$ , we gain  $\mathbf{0} \in \mathcal{T}_{\mathbf{r}}(\mathbf{z}_{-n})$  for all  $n \in \mathbb{N}$  by Proposition 3.6 because the expression in (3.5) is zero for each  $n \in \mathbb{N}$  in this case. By Lemma 3.3, the set  $\{\mathbf{z}_{-n} \mid n \in \mathbb{N}\}$  is finite, therefore we have  $\mathbf{z}_{-n} = \mathbf{z}_{-k}$  for some  $n > k \geq 0$ . Since  $\tau_{\mathbf{r}}^{n-k}(\mathbf{z}_{-n}) = \mathbf{z}_{-k}$ , we gain that  $\mathbf{z}_{-n}$  is purely periodic, and thus the same is true for  $\mathbf{x} = \tau_{\mathbf{r}}^n(\mathbf{z}_{-n})$ .

Again, by Lemma 3.3, it follows that only points  $\mathbf{x} \in \mathbb{Z}^d$  with  $\|\mathbf{x}\| \leq R$  can be purely periodic. Note that this was already proved in [5].  $\square$

A point  $\mathbf{t} \in \mathbb{R}^d$  that satisfies  $\mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x}) \setminus \bigcup_{\mathbf{y} \neq \mathbf{x}} \mathcal{T}_{\mathbf{r}}(\mathbf{y})$  for some  $\mathbf{x} \in \mathbb{Z}^d$  is called an *exclusive point* of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  according to the terminology introduced in [3] in the case of beta-tiles. Note that every exclusive point of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is an inner point of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  because SRS tiles are compact (Theorem 3.5) and because the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is locally finite (Lemma 3.3) and covers  $\mathbb{R}^d$  (Proposition 3.7). We will come back to the notion of exclusive points in Section 4. The following corollary is a consequence of Theorem 3.10.

**Corollary 3.11.** *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced. Then  $\mathbf{r} \in \mathcal{D}_d^{(0)}$  if and only if  $\mathbf{0} \in \mathcal{T}_{\mathbf{r}}(\mathbf{0}) \setminus \bigcup_{\mathbf{y} \neq \mathbf{0}} \mathcal{T}_{\mathbf{r}}(\mathbf{y})$ .*

*Proof.* Note that  $\mathbf{r} \in \mathcal{D}_d^{(0)}$  if and only if each orbit ends up in  $\mathbf{0}$ , implying that  $\mathbf{0}$  is the only purely periodic point.  $\square$

It immediately follows that for  $\mathbf{r} \in \mathcal{D}_d^{(0)}$  the central tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{0})$  has non-empty interior. One may ask if the interior of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is non-empty for each choice  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$ ,  $\mathbf{x} \in \mathbb{Z}^d$ . The answer is no, as the following example shows.

*Example 3.12.* Set  $\mathbf{r} = (\frac{9}{10}, -\frac{11}{20})$ . Consider the points

$$\mathbf{z}_1 = (-1, -1)^t, \mathbf{z}_2 = (-1, 1)^t, \mathbf{z}_3 = (1, 2)^t, \mathbf{z}_4 = (2, 1)^t, \mathbf{z}_5 = (1, -1)^t.$$

It can easily be verified that

$$\tau_{\mathbf{r}} : \mathbf{z}_1 \mapsto \mathbf{z}_2 \mapsto \mathbf{z}_3 \mapsto \mathbf{z}_4 \mapsto \mathbf{z}_5 \mapsto \mathbf{z}_1.$$

Thus, each of these points is purely periodic. Now calculate  $\tau_{\mathbf{r}}^{-1}(\mathbf{z}_1)$ :

$$\tau_{\mathbf{r}}^{-1}(\mathbf{z}_1) = \left\{ (x, -1)^t \mid x \in \mathbb{Z}, 0 \leq \frac{9}{10}x + \frac{11}{20} - 1 < 1 \right\} = \{(1, -1)^t\} = \{\mathbf{z}_5\}.$$

Similarly it can be shown that  $\tau_{\mathbf{r}}^{-1}(\mathbf{z}_i) = \{\mathbf{z}_{i-1}\}$  for  $i \in \{2, 3, 4, 5\}$ . Hence every tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{z}_i)$ ,  $i \in \{1, 2, 3, 4, 5\}$ , consists of a single point (the point  $\mathbf{0}$ ). The central tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{0})$  for this parameter is the one shown in Figure 1 on the lower right hand side.

*Example 3.13.* For  $\mathbf{r} = (\frac{3}{4}, 1)$ , the tiles  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  with  $\|\mathbf{x}\|_{\infty} \leq 2$  are depicted in Figure 3. As  $\mathbf{r} \in \mathcal{D}_d^{(0)}$ , we will see in Corollary 4.7 that the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is a weak tiling of  $\mathbb{R}^d$ .




FIGURE 3. The SRS tiles  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  with  $\|\mathbf{x}\|_{\infty} \leq 2$  corresponding to  $\mathbf{r} = (\frac{3}{4}, 1)$ .

#### 4. MULTIPLE TILINGS AND TILING CONDITIONS

According to Proposition 3.7, the family  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  of SRS tiles forms a covering of  $\mathbb{R}^d$ . Various tiling conditions concerning CNS and beta-tiles are spread in the literature (see *e.g.* the references in [13, 16]). They are of a combinatorial or dynamical nature, or they are expressed in terms of number systems. Among these conditions, the fact that  $\mathbf{0}$  is an inner point of the central tile plays an important role, which is related to the finiteness property (F) introduced and discussed in Sections 6 (see *e.g.* [2] for the case of beta-tiles).

In this section, we study tiling properties of SRS tiles. The notions of covering and tiling we will use here are discussed in Section 4.1. In Section 4.2, some facts on  $m$ -exclusive points are shown. A sufficient condition for coverings to be in fact tilings is given in Section 4.3.

**4.1. Coverings and tilings.** According to Lemma 3.3 and Proposition 3.7, the family of SRS tiles  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is a *covering with bounded degree*, that is, every point  $\mathbf{t} \in \mathbb{R}^d$  is contained in a finite and uniformly bounded number of tiles. Thus there exists a unique positive integer  $m$  such that every point  $\mathbf{t} \in \mathbb{R}^d$  is contained in at least  $m$  SRS tiles and there exists a point that is contained in exactly  $m$  tiles. Let us introduce several definitions concerning the notions of covering and tiling.

**Definition 4.1** (Covering and tiling; exclusive and inner point). Let  $\mathcal{K}$  be a locally finite collection of compact subsets covering  $\mathbb{R}^d$ .

- The *covering degree* with respect to  $\mathcal{K}$  of a point  $\mathbf{t} \in \mathbb{R}^d$  is given by  $\deg(\mathcal{K}, \mathbf{t}) := \#\{K \in \mathcal{K} \mid \mathbf{t} \in K\}$ .
- The *covering degree* of  $\mathcal{K}$  is given by  $\deg(\mathcal{K}) := \min\{\deg(\mathcal{K}, \mathbf{t}) \mid \mathbf{t} \in \mathbb{R}^d\}$ .
- The collection  $\mathcal{K}$  is a *weak  $m$ -tiling* if  $\deg(\mathcal{K}) = m$  and  $\bigcap_{i=1}^{m+1} \text{int}(K_i) = \emptyset$  for every choice of  $m+1$  pairwise different  $K_1, \dots, K_{m+1} \in \mathcal{K}$ . A weak 1-tiling is also called *weak tiling*.
- A point  $\mathbf{t} \in \mathbb{R}^d$  is  *$m$ -exclusive* with respect to  $\mathcal{K}$  if  $\deg(\mathcal{K}, \mathbf{t}) = \deg(\mathcal{K}) = m$ .
- A point  $\mathbf{t} \in \mathbb{R}^d$  is an *inner point* of the collection  $\mathcal{K}$  if  $\mathbf{t} \notin \bigcup_{K \in \mathcal{K}} \partial K$ .

In particular, the collection  $\mathcal{K}$  is a weak tiling if each inner point of  $\mathcal{K}$  belongs to exactly one element of  $\mathcal{K}$ . Moreover, the definition of 1-exclusive points with respect to  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  recovers the notion of exclusive points introduced in Section 3.3.

Let us recall that a tiling by translation is often defined as a collection of tiles having finitely many tiles up to translation, with a tile being assumed to be the closure of its interior. We study weak tilings in the sense of Definition 4.1 because of the following reasons:

- There exist choices of  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$ ,  $\mathbf{x} \in \mathbb{Z}^d$ , such that the tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is not the closure of its interior, see Example 3.12.
- There exist parameters  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  such that the family  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is not a collection of finitely many tiles up to translation, *e.g.*  $\mathbf{r} = (-2/3)$ , see Corollary 5.20. We conjecture that this holds for every  $\mathbf{r}$  related to a non-monic CNS (see Section 5) or a non-unit Pisot number (see Section 6).
- We are not able to show that the boundaries of the tiles  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  have zero  $d$ -dimensional Lebesgue measure. If  $\mathbf{r}$  is related to a monic CNS or a unit Pisot number, the fact that the boundary of each tile has zero measure is a direct consequence of the self-affine structure of the boundary of the tiles (*cf.* [3, 24, 29]). For other parameters  $\mathbf{r}$ , we have no appropriate description of the boundary, and we cannot evaluate its measure.

We are now going to prove that, for a large class of parameters  $\mathbf{r}$ , the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is a weak  $m$ -tiling, by showing that the set of  $m$ -exclusive points is dense in  $\mathbb{R}^d$ .

**4.2.  $m$ -exclusive points.** Let us first prove that  $m$ -exclusive points are always inner points of  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ .

**Lemma 4.2.** *Let  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  be reduced and let  $m$  be the covering degree of  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ . Then there exists an  $m$ -exclusive point. Every  $m$ -exclusive point  $\mathbf{t} \in \bigcap_{i=1}^m \mathcal{T}_{\mathbf{r}}(\mathbf{x}_i)$  satisfies  $\mathbf{t} \in \bigcap_{i=1}^m \text{int}(\mathcal{T}_{\mathbf{r}}(\mathbf{x}_i))$ . In particular,  $m$ -exclusive points are inner points of  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ .*

*Proof.* The existence of an  $m$ -exclusive point is a direct consequence of the definition of the covering degree.

Assume that  $\mathbf{t}$  is an  $m$ -exclusive point. Let  $\mathcal{T}_{\mathbf{r}}(\mathbf{x}_1), \dots, \mathcal{T}_{\mathbf{r}}(\mathbf{x}_m)$  be the  $m$  tiles it belongs to. Assume that there exists a sequence of points  $(\mathbf{t}_n)_{n \in \mathbb{N}}$  with values in  $\mathbb{R}^d$  with  $\lim_{n \rightarrow \infty} \mathbf{t}_n = \mathbf{t}$ , and a sequence of points  $(\mathbf{z}_n)_{n \in \mathbb{N}}$  with values in  $\mathbb{Z}^d$  such that  $\mathbf{t}_n \in \mathcal{T}_{\mathbf{r}}(\mathbf{z}_n)$  for all  $n \in \mathbb{N}$ , with  $\mathbf{z}_n$  distinct from all  $\mathbf{x}_i$ 's. Since the collection is locally finite, an infinite subsequence of  $\mathbf{z}_n$ 's is constant, say equal to  $\mathbf{z}$ . Since the tiles are compact, this implies that  $\mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{z})$ , which contradicts the  $m$ -exclusivity. Thus there exists a neighbourhood  $U$  of  $\mathbf{t}$  with  $U \cap \mathcal{T}_{\mathbf{r}}(\mathbf{z}) = \emptyset$  for each  $\mathbf{z} \in \mathbb{Z}^d \setminus \{\mathbf{x}_1, \dots, \mathbf{x}_m\}$ . By the  $m$ -covering property, we know that each point belongs to at least  $m$  tiles, which implies that  $U \subseteq \bigcap_{i=1}^m \mathcal{T}_{\mathbf{r}}(\mathbf{x}_i)$ .  $\square$

Let us now prove that an  $m$ -exclusive point  $\mathbf{t} \in \bigcap_{i=1}^m \mathcal{T}_r(\mathbf{x}_i)$  is somehow characterized by any sequence of approximations  $M_r^n \mathbf{z}_{-n}$  defined in Proposition 3.6. Note that  $\mathbf{t} = \text{Lim}_{n \rightarrow \infty} M_r^n \mathcal{T}_r(\mathbf{z}_{-n})$ , and that  $M_r^n \mathcal{T}_r(\mathbf{z}_{-n})$  is a tile in the  $n$ -fold subdivision  $\mathcal{T}_r(\mathbf{x}_i) = \bigcup_{\mathbf{z} \in \tau_r^{-n}(\mathbf{x}_i)} M_r^n \mathcal{T}_r(\mathbf{z})$  for some  $\mathbf{x}_i$ , which is given by applying (3.3)  $n$  times. The following proposition states that a point  $\mathbf{t}$  is  $m$ -exclusive provided that, for some  $n \in \mathbb{N}$ , each  $M_r^n \mathcal{T}_r(\mathbf{z}_{-n} + \mathbf{y})$  with  $\|\mathbf{y}\| \leq 2R$  occurs in the subdivision of some tile  $\mathcal{T}_r(\mathbf{x}_i)$ ,  $1 \leq i \leq m$ .

**Proposition 4.3.** *Let  $m$  be the covering degree of  $\{\mathcal{T}_r(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ ,  $\mathbf{t} \in \mathbb{R}^d$ ,  $(\mathbf{z}_{-n})_{n \in \mathbb{N}}$  as in Proposition 3.6, and  $R$  defined by (3.1). Then  $\mathbf{t}$  is  $m$ -exclusive and contained in the intersection  $\bigcap_{i=1}^m \mathcal{T}_r(\mathbf{x}_i)$  if and only if there exists some  $n \in \mathbb{N}$  such that*

$$(4.1) \quad \tau_r^n(\mathbf{z}_{-n} + \mathbf{y}) \in \{\mathbf{x}_1, \dots, \mathbf{x}_m\} \quad \text{for all } \mathbf{y} \in \mathbb{Z}^d \text{ with } \|\mathbf{y}\| \leq 2R.$$

*If, for some  $\mathbf{z} \in \mathbb{Z}^d$ ,  $n \in \mathbb{N}$ ,*

$$(4.2) \quad \#\{\tau_r^n(\mathbf{z} + \mathbf{y}) \mid \mathbf{y} \in \mathbb{Z}^d, \|\mathbf{y}\| \leq R\} = m,$$

*then  $M_r^n \mathbf{z}$  is an  $m$ -exclusive point.*

*Proof.* Assume that  $\mathbf{t}$  is an  $m$ -exclusive point. By Lemma 4.2, there exists some  $\varepsilon > 0$  such that every point  $\mathbf{t}' \in \mathbb{R}^d$  satisfying  $\|\mathbf{t}' - \mathbf{t}\| < \varepsilon$  lies only in the tiles  $\mathcal{T}_r(\mathbf{x}_1), \dots, \mathcal{T}_r(\mathbf{x}_m)$ . Let  $n \in \mathbb{N}$  satisfy  $4\tilde{\rho}^n R < \varepsilon$ . Since  $M_r^{-n} \mathbf{t} \in \mathcal{T}_r(\mathbf{z}_{-n})$ , we have  $\|M_r^{-n} \mathbf{t} - \mathbf{z}_{-n}\| \leq R$  by Lemma 3.3, thus  $\|\mathbf{t} - M_r^n(\mathbf{z}_{-n} + \mathbf{y})\| \leq 3\tilde{\rho}^n R$  if  $\|\mathbf{y}\| \leq 2R$ . By Theorem 3.5, there exists a point  $\mathbf{t}' \in \mathcal{T}_r(\tau_r^n(\mathbf{z}_{-n} + \mathbf{y}))$  with  $\|\mathbf{t}' - M_r^n(\mathbf{z}_{-n} + \mathbf{y})\| \leq \tilde{\rho}^n R$ . Since  $\|\mathbf{t}' - \mathbf{t}\| \leq 4\tilde{\rho}^n R < \varepsilon$ , we obtain  $\tau_r^n(\mathbf{z}_{-n} + \mathbf{y}) \in \{\mathbf{x}_1, \dots, \mathbf{x}_m\}$ .

For the other direction, assume that there exists some  $n \in \mathbb{N}$  such that (4.1) holds. We have to show that  $\mathbf{t} \in \mathcal{T}_r(\mathbf{z}'_0)$  implies  $\mathbf{z}'_0 \in \{\mathbf{x}_1, \dots, \mathbf{x}_m\}$ . Let  $(\mathbf{z}'_{-n})_{n \in \mathbb{N}}$  be as in Proposition 3.6. Since  $\|M_r^{-n} \mathbf{t} - \mathbf{z}'_{-n}\| \leq R$  and  $\|M_r^{-n} \mathbf{t} - \mathbf{z}_{-n}\| \leq R$ , (4.1) implies  $\tau_r(\mathbf{z}'_{-n}) \in \{\mathbf{x}_1, \dots, \mathbf{x}_m\}$ . Since  $\mathbf{z}'_0 = \tau_r(\mathbf{z}'_{-n})$ , the point  $\mathbf{t}$  is  $m$ -exclusive.

For the second statement, let  $\{\tau_r^n(\mathbf{z} + \mathbf{y}) \mid \mathbf{y} \in \mathbb{Z}^d, \|\mathbf{y}\| \leq R\} = \{\mathbf{x}_1, \dots, \mathbf{x}_m\}$ ,  $M_r^n \mathbf{z} \in \mathcal{T}_r(\mathbf{z}'_0)$ , and  $(\mathbf{z}'_{-n})_{n \in \mathbb{N}}$  be as in Proposition 3.6, with  $\mathbf{t} = M_r^n \mathbf{z}$ . Then we have  $\|\mathbf{z}'_{-n} - \mathbf{z}\| \leq R$ , thus  $\mathbf{z}'_0 \in \{\mathbf{x}_1, \dots, \mathbf{x}_m\}$  and  $M_r^n \mathbf{z}$  is  $m$ -exclusive.  $\square$

Note that Proposition 4.3 provides an easy way to show that a point is  $m$ -exclusive. However, it does not provide a finite method to prove that a point is not  $m$ -exclusive.

The following corollary of Proposition 4.3 permits us to obtain an  $m$ -exclusive point from another one, by finding a translation preserving the local configuration of tiles up to the  $n$ -th level, with  $n$  such that (4.2) holds.

**Corollary 4.4.** *Let  $m$  be the covering degree of  $\{\mathcal{T}_r(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ , and assume that  $\mathbf{z} \in \mathbb{Z}^d$ ,  $n \in \mathbb{N}$  satisfy (4.2). Let  $\mathbf{a} \in \mathbb{Z}^d$ . If there exists some  $\mathbf{b} \in \mathbb{Z}^d$  such that*

$$\tau_r^n(\mathbf{z} + \mathbf{a} + \mathbf{y}) = \tau_r^n(\mathbf{z} + \mathbf{y}) + \mathbf{b} \quad \text{for all } \mathbf{y} \in \mathbb{Z}^d \text{ with } \|\mathbf{y}\| \leq R,$$

*then  $M_r^n(\mathbf{z} + \mathbf{a})$  is an  $m$ -exclusive point.*

A simple way to obtain an infinite number of  $m$ -exclusive points from one  $m$ -exclusive point  $\mathbf{t} \neq \mathbf{0}$  is provided by the following lemma.

**Lemma 4.5.** *If  $\mathbf{t}$  is an  $m$ -exclusive point, then  $M_r \mathbf{t}$  is an  $m$ -exclusive point.*

*Proof.* If  $M_r \mathbf{t} \in \mathcal{T}_r(\mathbf{x}_i)$ , then there exists some  $\mathbf{y}_i \in \tau_r^{-1}(\mathbf{x}_i)$  such that  $\mathbf{t} \in \mathcal{T}_r(\mathbf{y}_i)$ , and all  $\mathbf{y}_i$  are mutually different if the  $\mathbf{x}_i$  are. Therefore the number of tiles to which a point  $M_r \mathbf{t}$  belongs cannot be larger than that for  $\mathbf{t}$ .  $\square$

**4.3. Weak  $m$ -tilings.** In what follows we will establish our tiling results. In order to prove these results we first show that if  $\mathbf{r}$  has certain algebraic properties then there exists a set that is relatively dense in  $\mathbb{R}^d$  containing only vectors that stabilize the configuration of tiles. Together with Corollary 4.4 and Lemma 4.5, this will prove that  $\{\mathcal{T}_r(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  forms a weak  $m$ -tiling. If  $\mathbf{r}$  also satisfies the finiteness property, it even forms a weak tiling.

**Theorem 4.6.** *Let  $\mathbf{r} = (r_0, \dots, r_{d-1}) \in \text{int}(\mathcal{D}_d)$  with  $r_0 \neq 0$ , let  $m$  be the covering degree of  $\{\mathcal{T}_r(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ , and assume that  $\mathbf{r}$  satisfies one of the following conditions:*

- $\mathbf{r} \in \mathbb{Q}^d$ ,
- $(x - \beta)(x^d + r_{d-1}x^{d-1} + \cdots + r_1x + r_0) \in \mathbb{Z}[x]$  for some  $\beta > 1$ ,
- $r_0, \dots, r_{d-1}$  are algebraically independent over  $\mathbb{Q}$ .

Then the set of  $m$ -exclusive points is dense in  $\mathbb{R}^d$ , and  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is a weak  $m$ -tiling.

*Proof.* By the definition of the covering degree (Definition 4.1), there exists an  $m$ -exclusive point  $\mathbf{t}$  with respect to  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$ . Thus the first part of Proposition 4.3 implies that there exist  $\mathbf{z} \in \mathbb{Z}^d$  and  $n \in \mathbb{N}$  satisfying (4.2). If we find a relatively dense set  $\Lambda$  of vectors  $\mathbf{a} \in \mathbb{Z}^d$  satisfying the conditions of Corollary 4.4, then the set  $\{M_{\mathbf{r}}^n(\mathbf{z} + \mathbf{a}) \mid \mathbf{a} \in \Lambda\}$  forms a set of  $m$ -exclusive inner points which is relatively dense in  $\mathbb{R}^d$ . Applying Lemma 4.5 to the elements of this set yields that the set of  $m$ -exclusive points is dense in  $\mathbb{R}^d$ . In view of Definition 4.1, this already proves that  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  forms a weak  $m$ -tiling.

It remains to find a relatively dense set  $\Lambda$  of vectors  $\mathbf{a} \in \mathbb{Z}^d$  satisfying the conditions of Corollary 4.4. This is done separately for each of the three classes of parameters given in the statement of the theorem.

**Case 1:**  $\mathbf{r} \in \mathbb{Q}^d$ . Let  $\mathbf{z} \in \mathbb{Z}^d$  and  $n \in \mathbb{N}$  satisfying (4.2) be given as above. Choose  $q \in \mathbb{N}$  in a way that  $\mathbf{r} \in q^{-1}\mathbb{Z}^d$ . Then we have for every  $\mathbf{x}, \mathbf{a} \in \mathbb{Z}^d$ ,  $k \geq 1$ ,

$$\tau_{\mathbf{r}}(\mathbf{x} + q^k \mathbf{a}) = M_{\mathbf{r}}(\mathbf{x} + q^k \mathbf{a}) + (0, \dots, 0, \{\mathbf{r}(\mathbf{x} + q^k \mathbf{a})\}) = \tau_{\mathbf{r}}(\mathbf{x}) + q^{k-1} \mathbf{a}'$$

for some  $\mathbf{a}' \in \mathbb{Z}^d$  which does not depend on  $\mathbf{x}$ . Iterating this, we get that  $\tau_{\mathbf{r}}^n(\mathbf{x} + q^n \mathbf{a}) = \tau_{\mathbf{r}}^n(\mathbf{x}) + \mathbf{b}$  for some  $\mathbf{b} \in \mathbb{Z}^d$  which does not depend on  $\mathbf{x}$ . In particular, this implies that

$$\tau_{\mathbf{r}}^n(\mathbf{x} + q^n \mathbf{a} + \mathbf{y}) = \tau_{\mathbf{r}}^n(\mathbf{x} + \mathbf{y}) + \mathbf{b} \quad \text{for all } \mathbf{y} \in \mathbb{Z}^d \text{ with } \|\mathbf{y}\| \leq R.$$

Thus each element of the set  $\Lambda := \{q^n \mathbf{a} \mid \mathbf{a} \in \mathbb{Z}^d\}$  satisfies the conditions of Corollary 4.4. As  $\Lambda$  is relatively dense in  $\mathbb{R}^d$ , we are done in this case.

**Case 2:**  $A(x) = (x - \beta)(x^d + r_{d-1}x^{d-1} + \cdots + r_1x + r_0) \in \mathbb{Z}[x]$  for some  $\beta > 1$ . Let  $\mathbf{z} \in \mathbb{Z}^d$  and  $n \in \mathbb{N}$  satisfying (4.2) be given as above. Since  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$ , all roots of  $x^d + r_{d-1}x^{d-1} + \cdots + r_1x + r_0$  have modulus less than 1. Therefore,  $A(x)$  is irreducible. (Indeed,  $\beta$  is a Pisot number.) Let

$$\varepsilon := \min_{\|\mathbf{y}\| \leq R, 0 \leq k < n} \beta^{-k} (1 - \{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{z} + \mathbf{y})\}) > 0.$$

From the definition of  $\tau_{\mathbf{r}}$  we know that for every  $\mathbf{x}, \mathbf{a} \in \mathbb{Z}^d$ , one has  $\tau_{\mathbf{r}}(\mathbf{x} + \mathbf{a}) = \tau_{\mathbf{r}}(\mathbf{x}) + \tau_{\mathbf{r}}(\mathbf{a})$  if and only if  $\{\mathbf{r}\mathbf{x} + \mathbf{r}\mathbf{a}\} = \{\mathbf{r}\mathbf{x}\} + \{\mathbf{r}\mathbf{a}\}$ . In Proposition 6.1, we will see that  $\{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{a})\} = T_{\beta}^k(\{\mathbf{r}\mathbf{a}\})$ , where  $T_{\beta}$  is the  $\beta$ -transformation defined in Section 6.1. If we choose  $\mathbf{a} \in \mathbb{Z}^d$  such that  $\{\mathbf{r}\mathbf{a}\} < \varepsilon$ , then we get  $\{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{a})\} = \beta^k \{\mathbf{r}\mathbf{a}\}$  for  $0 \leq k < n$ , and

$$(4.3) \quad \tau_{\mathbf{r}}^k(\mathbf{z} + \mathbf{a} + \mathbf{y}) = \tau_{\mathbf{r}}^k(\mathbf{z} + \mathbf{y}) + \tau_{\mathbf{r}}^k(\mathbf{a}) \quad \text{for all } \|\mathbf{y}\| \leq R, 0 \leq k \leq n.$$

Thus each element of the set  $\Lambda := \{\mathbf{a} \in \mathbb{Z}^d \mid \{\mathbf{r}\mathbf{a}\} < \varepsilon\}$  satisfies the conditions of Corollary 4.4. Since  $A(x)$  is irreducible, the coordinates of  $\mathbf{r}$  are linearly independent over  $\mathbb{Q}$ . Thus Kronecker's theorem yields that  $\Lambda$  is relatively dense in  $\mathbb{R}^d$  and we are done also in this case.

**Case 3:**  $r_0, \dots, r_{d-1}$  are algebraically independent. Let  $\mathbf{z} \in \mathbb{Z}^d$  and  $n \in \mathbb{N}$  satisfying (4.2) be given as above. Set

$$\varepsilon_k := \min_{\|\mathbf{y}\| \leq R} (1 - \{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{z} + \mathbf{y})\}) \quad \text{for } 0 \leq k < n.$$

We have  $\varepsilon_k > 0$ . Similarly to Case 2, each element of the set

$$\Lambda := \{\mathbf{a} \in \mathbb{Z}^d \mid \{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{a})\} < \varepsilon_k \text{ for } 0 \leq k < n\}$$

satisfies (4.3). Thus each  $\mathbf{a} \in \Lambda$  satisfies the conditions of Corollary 4.4. It remains to prove that  $\Lambda$  is relatively dense in  $\mathbb{R}^d$ . To this matter we need the following notations. Let  $[x]$  denote the nearest integer to  $x \in \mathbb{R}$  (with  $[z + 1/2] = z$  for  $z \in \mathbb{Z}$ ), let  $\{x\}_c = x - [x]$  be the ‘‘centralized fractional part’’ and set

$$\tilde{\tau}_{\mathbf{r}}(\mathbf{x}) = M_{\mathbf{r}}\mathbf{x} + (0, \dots, 0, \{\mathbf{r}\mathbf{x}\}_c).$$

Consider the partition  $\Lambda = \bigcup_{\eta \in \{0,1\}^n} \Lambda_{\eta}$  with

$$\Lambda_{\eta} := \{\mathbf{a} \in \mathbb{Z}^d \mid \{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{a})\} \in \eta_k \varepsilon_k / 2 + [0, \varepsilon_k / 2] \text{ for } 0 \leq k < n\} \quad (\eta \in \{0,1\}^n)$$

(here  $\eta = (\eta_0, \dots, \eta_{n-1})$ ). We will prove the following claim.

*Claim.* Let  $\mathbf{e}_j$ ,  $1 \leq j \leq d$ , be the canonical unit vectors. For each  $\eta \in \{0, 1\}^n$  and  $j \in \{1, \dots, d\}$ , there exists  $z_{\eta,j} \in \mathbb{Z}$  with

$$(4.4) \quad \{\mathbf{r}\tilde{\tau}_{\mathbf{r}}^k(z_{\eta,j}\mathbf{e}_j)\}_c \in (-1)^{\eta_k}[0, \varepsilon_k/2) \quad \text{for all } 0 \leq k < n.$$

Before we prove this claim we show that it implies the relative denseness of  $\Lambda$  in  $\mathbb{R}^d$ . Indeed, let  $\eta \in \{0, 1\}^n$  and  $j \in \{1, \dots, d\}$  be arbitrary. Then the claim yields that for each  $\mathbf{a} \in \Lambda_\eta$  we have

$$\{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{a} + z_{\eta,j}\mathbf{e}_j)\} = \{\mathbf{r}(\tau_{\mathbf{r}}^k(\mathbf{a}) + \tilde{\tau}_{\mathbf{r}}^k(z_{\eta,j}\mathbf{e}_j))\} = \{\mathbf{r}\tau_{\mathbf{r}}^k(\mathbf{a})\} + \{\mathbf{r}\tilde{\tau}_{\mathbf{r}}^k(z_{\eta,j}\mathbf{e}_j)\}_c \in [0, \varepsilon_k)$$

and thus  $\mathbf{a} + z_{\eta,j}\mathbf{e}_j \in \Lambda$ . Moreover, by analogous reasoning we see that the claim implies the following ‘‘dual’’ result: for each  $\eta \in \{0, 1\}^n$  set  $\eta' := (1, \dots, 1) - \eta \in \{0, 1\}^n$ . Then for each  $j \in \{1, \dots, d\}$  and each  $\mathbf{a} \in \Lambda_\eta$  we have  $\mathbf{a} - z_{\eta',j}\mathbf{e}_j \in \Lambda$ .

Summing up, the claim implies that from each  $\mathbf{a} \in \Lambda$  there exist other elements of  $\Lambda$  in uniformly bounded distance in all positive and negative coordinate directions. This proves that  $\Lambda$  is relatively dense in  $\mathbb{R}^d$ . Thus it remains to prove the above claim.

To prove this claim let  $\eta = (\eta_0, \dots, \eta_{n-1}) \in \{0, 1\}^n$  and  $j \in \{1, \dots, d\}$  be arbitrary but fixed. We need to find an integer  $z_{\eta,j}$  satisfying (4.4). First observe that for  $\mathbf{b} \in \mathbb{Z}^d$

$$(4.5) \quad \tilde{\tau}_{\mathbf{r}}^k(\mathbf{b}) = M_{\mathbf{r}}^k \mathbf{b} + \sum_{j=0}^{k-1} M_{\mathbf{r}}^{k-j-1} (0, \dots, 0, \{\mathbf{r}\tilde{\tau}_{\mathbf{r}}^j(\mathbf{b})\}_c)^t.$$

Let  $\gamma_k = \{\mathbf{r}M_{\mathbf{r}}^k \mathbf{e}_d\}_c$ ,  $0 \leq k < n$ . If the arguments of all centralized fractional parts are small, then multiplying (4.5) by  $\mathbf{r}$  and applying  $\{\cdot\}_c$  yields

$$(4.6) \quad \{\mathbf{r}\tilde{\tau}_{\mathbf{r}}^k(\mathbf{b})\}_c = \{\mathbf{r}M_{\mathbf{r}}^k \mathbf{b}\}_c + \sum_{j=0}^{k-1} \{\mathbf{r}\tilde{\tau}_{\mathbf{r}}^j(\mathbf{b})\}_c \gamma_{k-j-1}.$$

Inserting (4.6) iteratively in itself we gain

$$\begin{aligned} \{\mathbf{r}\tilde{\tau}_{\mathbf{r}}^k(\mathbf{b})\}_c &= \{\mathbf{r}M_{\mathbf{r}}^k \mathbf{b}\}_c + \sum_{k=j_0 > j_1 > \dots > j_\ell \geq 0, \ell \geq 1} \{\mathbf{r}M_{\mathbf{r}}^{j_\ell} \mathbf{b}\}_c \prod_{i=0}^{\ell-1} \gamma_{j_i - j_{i+1} - 1} \\ &= \sum_{h=0}^k \{\mathbf{r}M_{\mathbf{r}}^h \mathbf{b}\}_c \sum_{k=j_0 > \dots > j_\ell = h, \ell \geq 0} \prod_{i=0}^{\ell-1} \gamma_{j_i - j_{i+1} - 1}. \end{aligned}$$

Setting

$$c_k := \sum_{k=j_0 > \dots > j_\ell = 0, \ell \geq 0} \prod_{i=0}^{\ell-1} \gamma_{j_i - j_{i+1} - 1}$$

we get

$$\{\mathbf{r}\tilde{\tau}_{\mathbf{r}}^k(\mathbf{b})\}_c = \sum_{h=0}^k c_{k-h} \{\mathbf{r}M_{\mathbf{r}}^h \mathbf{b}\}_c.$$

Now we inductively choose intervals  $I_k$ ,  $0 \leq k < n$ , satisfying

$$I_k + \sum_{h=0}^{k-1} c_{k-h} I_h \subseteq (-1)^{\eta_k} [0, \varepsilon_k/2).$$

W.l.o.g., we can choose the intervals  $I_k$  sufficiently small such that (4.6) holds provided that  $\{\mathbf{r}M_{\mathbf{r}}^k \mathbf{b}\}_c \in I_k$  for  $0 \leq k < n$ .

Since  $r_0, \dots, r_{d-1}$  are algebraically independent, the numbers  $\{\mathbf{r}M_{\mathbf{r}}^k \mathbf{e}_j\}_c$ , for  $0 \leq k < n$ , are linearly independent over  $\mathbb{Q}$ . Thus Kronecker’s theorem yields the existence of an integer  $z_{\eta,j}$  satisfying  $\{z_{\eta,j} \mathbf{r}M_{\mathbf{r}}^k \mathbf{e}_j\}_c \in I_k$  for  $0 \leq k < n$ , hence,  $z_{\eta,j}$  satisfies (4.4) and we are done.  $\square$

Since Corollary 3.11 implies that for each SRS with finiteness property the origin is an exclusive point of  $\mathcal{T}_{\mathbf{r}}(\mathbf{0})$  we gain the following tiling property.

**Corollary 4.7.** *If  $\mathbf{r} \in \mathcal{D}_d^{(0)}$  is reduced and satisfies one of the conditions of Theorem 4.6, then  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is a weak tiling.*

Let us stress the fact that we have no general algorithmic criterion to check Proposition 4.3. This is mainly due to the fact that we have no IFS describing the boundary of an SRS tile. Nonetheless, Theorem 4.6 implies a tiling criterion.

**Corollary 4.8.** *If  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  is reduced and satisfies one of the conditions of Theorem 4.6, then the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \mid \mathbf{x} \in \mathbb{Z}^d\}$  is a weak tiling if and only if it has at least one exclusive point.*

We conclude this section with a result that treats the case  $d = 1$  in a very complete way. (We identify one dimensional vectors with scalars here.)

**Theorem 4.9.** *Let  $r \in \text{int}(\mathcal{D}_1)$  be reduced, i.e.,  $0 < |r| < 1$ . Then  $\{\mathcal{T}_r(x) \mid x \in \mathbb{Z}\}$  is a tiling of  $\mathbb{R}$  by (possibly degenerate) intervals. Here, tiling has to be understood in the usual sense, i.e.,*

$$\bigcup_{x \in \mathbb{Z}} \mathcal{T}_r(x) = \mathbb{R} \quad \text{with} \quad \#(\mathcal{T}_r(x) \cap \mathcal{T}_r(x')) \leq 1 \quad \text{for } x, x' \in \mathbb{Z}, x \neq x'.$$

*Proof.* Let first  $r > 0$  and  $x_0, y_0 \in \mathbb{Z}$ . Then by the definition of  $\tau_r$  we easily see that  $x_0 > y_0$  implies that  $-x_1 > -y_1$  for each  $x_1 \in \tau_r^{-1}(x_0)$ ,  $y_1 \in \tau_r^{-1}(y_0)$ . Thus, by induction on  $n$  we have

$$x_0 > y_0 \quad \text{implies that} \quad (-1)^n x_n > (-1)^n y_n \quad \text{for each } x_n \in \tau_r^{-n}(x_0), y_n \in \tau_r^{-n}(y_0).$$

From this implication the assertions follow for the case  $r > 0$ . The case  $r < 0$  can be treated similarly.  $\square$

## 5. SRS AND CANONICAL NUMBER SYSTEMS

The aim of this section is to relate SRS tiles to tiles associated with expanding polynomials.

**5.1. Expanding polynomials over  $\mathbb{Z}$ , SRS representations and canonical number systems.** Let  $A = a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0 \in \mathbb{Z}[x]$ ,  $a_0 \geq 2$ ,  $a_d \neq 0$ , and  $\mathcal{Q} := \mathbb{Z}[x]/A\mathbb{Z}[x]$  the factor ring, with  $X \in \mathcal{Q}$  being the image of  $x$  under the canonical epimorphism. Furthermore, set  $\mathcal{N} = \{0, \dots, a_0 - 1\}$ . We want to represent each element  $P \in \mathcal{Q}$  formally as

$$(5.1) \quad P = \sum_{n=0}^{\infty} b_n X^n \quad (b_n \in \mathcal{N}).$$

More precisely, we want to find a sequence  $(b_n)_{n \in \mathbb{N}}$  as in the following definition.

**Definition 5.1** ( $(A, \mathcal{N})$ -representation). Let  $P \in \mathcal{Q} = \mathbb{Z}[x]/A\mathbb{Z}[x]$ . A sequence  $(b_n)_{n \in \mathbb{N}}$  with  $b_n \in \mathcal{N}$  is called an  $(A, \mathcal{N})$ -representation of  $P$  if  $P - \sum_{i=0}^{m-1} b_i X^i \in X^m \mathcal{Q}$  for all  $m \in \mathbb{N}$ .

In order to show that such an representation exists and is unique, we introduce a *backward division mapping*  $D_A : \mathcal{Q} \rightarrow \mathcal{Q}$ .

**Lemma 5.2.** *The backward division mapping  $D_A : \mathcal{Q} \rightarrow \mathcal{Q}$  given for  $P = \sum_{i=0}^{\ell} p_i X^i$ ,  $p_i \in \mathbb{Z}$ , by*

$$D_A(P) = \sum_{i=0}^{\ell-1} p_{i+1} X^i - \sum_{i=0}^{d-1} q a_{i+1} X^i, \quad q = \left\lfloor \frac{p_0}{a_0} \right\rfloor,$$

*is well defined. Every  $P \in \mathcal{Q}$  has one and only one  $(A, \mathcal{N})$ -representation.*

*Proof.* It is easy to see that  $D_A(P)$  does not depend on the choice of the representation of  $P$ , that  $P = (p_0 - q a_0) + X D_A(P)$ , and that  $b_0 = p_0 - q a_0$  is the unique element in  $\mathcal{N}$  with  $P - b_0 \in X \mathcal{Q}$  (for the case of monic polynomials  $A$ , this is detailed in [5]; for the non-monic case, see [36]).

The  $(A, \mathcal{N})$ -representation can be obtained by successively applying  $D_A$ , which yields

$$P = \sum_{n=0}^{m-1} b_n X^n + X^m D_A^m(P)$$

with  $b_n = D_A^n(P) - X D_A^{n+1}(P) \in \mathcal{N}$ . Therefore, the  $(A, \mathcal{N})$ -representation of  $P$  is unique.  $\square$

In order to relate  $D_A$  to an SRS, we use an appropriate  $\mathbb{Z}$ -submodule of  $\mathcal{Q}$ , following [36].

**Definition 5.3** (Brunotte basis and Brunotte module). The *Brunotte basis modulo  $A$*  is defined by  $\{W_0, \dots, W_{d-1}\}$  with

$$(5.2) \quad W_0 = a_d \quad \text{and} \quad W_k = XW_{k-1} + a_{d-k} \quad \text{for } 1 \leq k \leq d-1.$$

The *Brunotte module*  $\Lambda_A$  is the  $\mathbb{Z}$ -submodule of  $\mathcal{Q}$  generated by the Brunotte basis. The representation mapping with respect to the Brunotte basis is denoted by

$$\Psi_A : \Lambda_A \rightarrow \mathbb{Z}^d, \quad P = \sum_{k=0}^{d-1} z_k W_k \mapsto (z_0, \dots, z_{d-1})^t.$$

Note that  $\Lambda_A$  is isomorphic to  $\mathcal{Q}$  if  $A$  is monic. (Here, monic means that  $a_d \in \{-1, 1\}$ .) This is easily seen by checking that the coordinate matrix of  $\{W_0, \dots, W_{d-1}\}$  w.r.t.  $\{1, X, \dots, X^{d-1}\}$  is given by

$$(5.3) \quad V := \begin{pmatrix} a_d & a_{d-1} & \cdots & \cdots & a_1 \\ 0 & \ddots & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & a_{d-1} \\ 0 & \cdots & \cdots & 0 & a_d \end{pmatrix}.$$

The restriction to Brunotte module  $\Lambda_A$  allows us to relate the SRS transformation  $\tau_{\mathbf{r}}$  to the backward division mapping  $D_A$  in the following way.

**Proposition 5.4.** *Let  $A = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0 \in \mathbb{Z}[x]$ ,  $a_0 \geq 2$ ,  $a_d \neq 0$ ,  $\mathbf{r} = (\frac{a_d}{a_0}, \dots, \frac{a_1}{a_0})$ . Then we have*

$$(5.4) \quad D_A^n \Psi_A^{-1}(\mathbf{z}) = \Psi_A^{-1} \tau_{\mathbf{r}}^n(\mathbf{z}) \quad \text{for all } \mathbf{z} \in \mathbb{Z}^d, n \in \mathbb{N}.$$

*In particular, the restriction of  $D_A$  to the Brunotte module  $\Lambda_A$  is conjugate to  $\tau_{\mathbf{r}}$ .*

*Proof.* On  $\Lambda_A$ , the mapping  $D_A$  can be written as

$$(5.5) \quad D_A \left( \sum_{k=0}^{d-1} z_k W_k \right) = \sum_{k=0}^{d-2} z_{k+1} W_k - \left\lfloor \frac{z_0 a_d + \cdots + z_{d-1} a_1}{a_0} \right\rfloor W_{d-1}$$

(see e.g. [5, Section 3]). Therefore, we have  $D_A \Psi_A^{-1}(\mathbf{z}) = \Psi_A^{-1} \tau_{\mathbf{r}}(\mathbf{z})$ , which implies (5.4). Since  $\Psi_A$  is bijective, the restriction of  $D_A$  to  $\Lambda_A$  is conjugate to  $\tau_{\mathbf{r}}$ .  $\square$

Note that in the monic case this gives a conjugacy between the backward division mapping on the full set  $\mathcal{Q}$  and the SRS transformation.

With help of the conjugacy proved in Proposition 5.4, we get a simple formula to gain the  $(A, \mathcal{N})$ -representation (5.1) for each  $P \in \Lambda_A$  using the associated transformation  $\tau_{\mathbf{r}}$ .

**Lemma 5.5.** *Let  $A = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0 \in \mathbb{Z}[x]$ ,  $a_0 \geq 2$ ,  $a_d \neq 0$ ,  $\mathbf{r} = (\frac{a_d}{a_0}, \dots, \frac{a_1}{a_0})$ . The  $(A, \mathcal{N})$ -representation of  $P \in \Lambda_A$  is given by*

$$b_n = \{\mathbf{r} \tau_{\mathbf{r}}^n \Psi_A(P)\} a_0 \quad \text{for all } n \in \mathbb{N}.$$

*Proof.* For fixed  $n$ , let  $D_A^n(P) = \sum_{k=0}^{d-1} z_k W_k$  and recall that  $\Psi_A D_A^n(P) = (z_0, \dots, z_{d-1})$ . By (5.2), (5.5) and the fact that  $XW_{d-1} + a_0 = 0$ , we obtain that

$$XD_A \left( \sum_{k=0}^{d-1} z_k W_k \right) = \sum_{k=0}^{d-2} z_{k+1} (W_{k+1} - a_{d-k-1}) + \left\lfloor \frac{z_0 a_d + \cdots + z_{d-1} a_1}{a_0} \right\rfloor a_0,$$

therefore

$$\begin{aligned}
 b_n &= D_A^n(P) - X D_A^{n+1}(P) \\
 &= z_0 W_0 + \sum_{k=0}^{d-2} z_{k+1} a_{d-k-1} - \left\lfloor \frac{z_0 a_d + \cdots + z_{d-1} a_1}{a_0} \right\rfloor a_0 = \left\{ \frac{z_0 a_d + \cdots + z_{d-1} a_1}{a_0} \right\} a_0 \\
 &= \{\mathbf{r} \Psi_A D_A^n(P)\} a_0 = \{\mathbf{r} \tau_{\mathbf{r}}^n \Psi_A(P)\} a_0. \quad \square
 \end{aligned}$$

If the  $(A, \mathcal{N})$ -representation  $(b_n)_{n \in \mathbb{N}}$  has only finitely many non-zero elements, then  $P$  can be written as a finite sum of the shape

$$P = \sum_{n=0}^{m-1} b_n X^n \quad (b_n \in \mathcal{N}).$$

We recover the following well-known notion of canonical number systems.

**Definition 5.6** (Canonical number system). Let  $A = a_d x^d + \cdots + a_1 x + a_0 \in \mathbb{Z}[x]$ ,  $a_0 \geq 2$ ,  $a_d \neq 0$ ,  $\mathcal{Q} = \mathbb{Z}[x]/AZ[x]$ , and  $\mathcal{N} = \{0, \dots, a_0 - 1\}$ . If for each  $P \in \mathcal{Q}$ , the  $(A, \mathcal{N})$ -representation  $(b_n)_{n \in \mathbb{N}}$ ,  $b_n \in \mathcal{N}$ , has only finitely many non-zero elements, then we call  $(A, \mathcal{N})$  a *canonical number system* (CNS, for short).

It is shown in [36] that it is sufficient to check the finiteness of the  $(A, \mathcal{N})$ -representations for all  $P \in \Lambda_A$  in order to check whether  $(A, \mathcal{N})$  is a CNS. Thus the conjugacy between  $D_A$  and  $\tau_{\mathbf{r}}$  is sufficient to reformulate the CNS property in terms of SRS with finiteness property.

**Proposition 5.7.** *Let  $A = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0 \in \mathbb{Z}[x]$ ,  $a_0 \geq 2$ ,  $a_d \neq 0$ , and  $\mathcal{N} = \{0, \dots, a_0 - 1\}$ . Then the following assertions hold.*

- *The polynomial  $A$  is expanding if and only if  $\mathbf{r} = \left(\frac{a_d}{a_0}, \frac{a_{d-1}}{a_0}, \dots, \frac{a_1}{a_0}\right)$  is contained in  $\text{int}(\mathcal{D}_d)$ .*
- *The pair  $(A, \mathcal{N})$  is a CNS if and only if  $\mathbf{r} = \left(\frac{a_d}{a_0}, \frac{a_{d-1}}{a_0}, \dots, \frac{a_1}{a_0}\right) \in \mathcal{D}_d^{(0)}$ .*

*Proof.* The first assertion follows from Lemma 2.3. The second assertion follows from the conjugacy given in Proposition 5.4 together with the fact that it is sufficient to check the finiteness of the  $(A, \mathcal{N})$ -representations for all  $P \in \Lambda_A$  (see also [36]).  $\square$

**5.2. Tiles associated with an expanding polynomial.** We define two classes of tiles for expanding polynomials. The first definition goes back to Kátai and Kórnyci [25] (see also [37]) for monic polynomials which give rise to a CNS. We first extend this definition to arbitrary expanding polynomials.

**Definition 5.8** (Self-affine tile associated with an expanding polynomial). Let  $A = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_1 x + a_0 \in \mathbb{Z}[x]$  be an expanding polynomial with  $a_0 \geq 2$ ,  $a_d \neq 0$ , and  $\mathcal{N} = \{0, \dots, a_0 - 1\}$ . The *self-affine tile associated with  $A$*  is defined as

$$(5.6) \quad \mathcal{F}_A := \left\{ \mathbf{t} \in \mathbb{R}^d \mid \mathbf{t} = \sum_{i=1}^{\infty} B^{-i} (c_i, 0, \dots, 0)^t, c_i \in \mathcal{N} \right\}$$

where  $B := V M_{\mathbf{r}}^{-1} V^{-1}$  with  $\mathbf{r} = \left(\frac{a_d}{a_0}, \frac{a_{d-1}}{a_0}, \dots, \frac{a_1}{a_0}\right)$ , and  $V$  is given by (5.3).

*Remark 5.9.* The matrix  $B$  is a companion matrix associated with the polynomial  $a_d^{-1} A$ . We use it instead of  $M_{\mathbf{r}}^{-1}$  in order to be consistent with the existing literature on CNS tiles.

Since  $A$  is expanding, it is easy to see that the series in (5.6) always converges. The tile  $\mathcal{F}_A$  has the following properties.

- The tile  $\mathcal{F}_A$  is compact.
- The tile  $\mathcal{F}_A$  is a self-affine set (hence the terminology) as it obeys the set equation  $B\mathcal{F}_A = \bigcup_{c \in \mathcal{N}} (\mathcal{F}_A + (c, 0, \dots, 0)^t)$ . Indeed, it is the unique non-empty compact set satisfying this equation (cf. e.g. [14, 18, 22]). Self-affine tiles have been studied extensively in a very general context in the literature. We refer the reader to the surveys by Vince [41] and Wang [42].

- If  $A$  is monic, then  $\{\mathbf{z} + \mathcal{F}_A \mid \mathbf{z} \in \mathbb{Z}^d\}$  forms a tiling of  $\mathbb{R}^d$  for many choices of  $A$  (e.g. for irreducible  $A$ , this is shown in Lagarias and Wang [30]).

For non-monic polynomials  $A$ , the above definition turns out to be not well-suited. The tiles have strong overlaps. The submodule  $\Lambda_A$  is a good tool to define a new class of tiles which forms a (multiple) tiling also for non-monic polynomials.

**Definition 5.10** (Brunotte tile associated with an expanding polynomial). Let  $A = a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0 \in \mathbb{Z}[x]$  be an expanding polynomial with  $a_0 \geq 2$ ,  $a_d \neq 0$ , and  $\mathcal{N} = \{0, \dots, a_0 - 1\}$ . For each  $P \in \Lambda_A$ , the *Brunotte tile* associated with  $A$  is defined as

$$\mathcal{G}_A(P) := \operatorname{Li}_{n \rightarrow \infty} B^{-n} V \Psi_A(D_A^{-n}(P) \cap \Lambda_A),$$

where  $\operatorname{Li}$  denotes the lower Hausdorff limit. The set  $\mathcal{G}_A(0)$  is called the *central Brunotte tile* associated with the expanding polynomial  $A$ .

**Lemma 5.11.** *We have*

$$\mathcal{F}_A = \operatorname{Lim}_{n \rightarrow \infty} B^{-n} V \Psi_A(D_A^{-n}(0))$$

and thus  $\mathcal{G}_A(P) \subseteq V \Psi_A(P) + \mathcal{F}_A$ .

*Proof.* This follows by direct calculation.  $\square$

If  $A$  is monic, then we will show that the inclusion in Lemma 5.11 becomes the equality  $\mathcal{G}_A(P) = V \Psi_A(P) + \mathcal{F}_A$  for every  $P \in \mathcal{Q}$ . Note that  $V \Psi_A(P) = (p_0, \dots, p_{d-1})$  for  $P = \sum_{j=0}^{d-1} p_j X^j$ .

**5.3. From tiles associated with expanding polynomials to SRS tiles.** Next we show that the Brunotte tiles are obtained from the SRS tiles by a linear transformation.

**Theorem 5.12.** *Let  $A = a_d x^d + a_{d-1} x^{d-1} + \dots + a_1 x + a_0 \in \mathbb{Z}[x]$ ,  $a_0 \geq 2$ ,  $a_d \neq 0$ , be an expanding polynomial, and  $\mathbf{r} = (\frac{a_d}{a_0}, \frac{a_{d-1}}{a_0}, \dots, \frac{a_1}{a_0})$ . Then*

$$\mathcal{G}_A(\Psi_A^{-1}(\mathbf{z})) = V \mathcal{T}_{\mathbf{r}}(\mathbf{z}) \quad \text{for all } \mathbf{z} \in \mathbb{Z}^d.$$

*Proof.* Set  $P = \Psi_A^{-1}(\mathbf{z})$ . We have

$$(5.7) \quad B^{-n} V \Psi_A(D_A^{-n}(P) \cap \Lambda_A) = B^{-n} V \tau_{\mathbf{r}}^{-n}(\mathbf{z}) = V M_{\mathbf{r}}^n \tau_{\mathbf{r}}^{-n}(\mathbf{z}).$$

Taking  $\operatorname{Li}$  for  $n \rightarrow \infty$  proves the result.  $\square$

The following corollary shows that the lower Hausdorff limit in the definition of the Brunotte tiles is indeed a Hausdorff limit, as for the SRS tiles. Furthermore, we show how (5.6) has to be adapted to give the Brunotte tiles.

**Corollary 5.13.** *The Brunotte tiles can be written as*

$$\mathcal{G}_A(P) = \operatorname{Lim}_{n \rightarrow \infty} B^{-n} V \Psi_A(D_A^{-n}(P) \cap \Lambda_A),$$

where  $\operatorname{Lim}$  denotes the Hausdorff limit. Moreover, if  $P = \sum_{k=0}^{d-1} p_k X^k$ , then

$$\mathcal{G}_A(P) = (p_0, \dots, p_{d-1})^t + \left\{ \mathbf{t} \in \mathbb{R}^d \mid \mathbf{t} = \sum_{i=1}^{\infty} B^{-i} (c_i, 0, \dots, 0)^t, c_i \in \mathcal{N}, \sum_{i=1}^n c_i X^{n-i} + X^n P \in \Lambda_A \text{ for all } n \in \mathbb{N} \right\}.$$

*Proof.* The first assertion follows from (5.7) and Theorem 3.5. The second assertion follows from Proposition 3.6 and the fact that

$$D_A^{-n}(P) \cap \Lambda_A = \left\{ \sum_{i=1}^n c_i X^{n-i} + X^n P \in \Lambda_A \mid c_i \in \mathcal{N} \right\}. \quad \square$$

For the monic case, we derive the following corollary.

**Corollary 5.14.** *Suppose that the polynomial  $A$  in Theorem 5.12 is monic. Then*

$$\mathcal{G}_A(P) = V\Psi_A(P) + \mathcal{F}_A \quad \text{for all } P \in \mathcal{Q}.$$

Now we can state the main result of the present section. It establishes tiling properties for Brunotte tiles which are even valid in the non-monic case.

**Theorem 5.15.** *Let  $A = a_dx^d + a_{d-1}x^{d-1} + \dots + a_1x + a_0 \in \mathbb{Z}[x]$ ,  $a_0 \geq 2$ ,  $a_d \neq 0$ , be an expanding polynomial, and  $\mathcal{N} = \{0, \dots, a_0 - 1\}$ . Then the following assertions hold.*

- *The collection  $\{\mathcal{G}_A(P) \mid P \in \Lambda_A\}$  forms a weak  $m$ -tiling of  $\mathbb{R}^d$  for some  $m \geq 1$ .*
- *If  $(A, \mathcal{N})$  is a CNS, then  $\{\mathcal{G}_A(P) \mid P \in \Lambda_A\}$  forms a weak tiling of  $\mathbb{R}^d$ .*

*Proof.* By Theorem 5.12, it is equivalent to consider the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{z}) \mid \mathbf{z} \in \mathbb{Z}^d\}$  for  $\mathbf{r} = (\frac{a_d}{a_0}, \frac{a_{d-1}}{a_0}, \dots, \frac{a_1}{a_0})$ . In view of Proposition 5.7, the first assertion follows from Theorem 4.6, while the second assertion is a consequence of Corollary 4.7.  $\square$

*Example 5.16.* Consider the monic CNS polynomial  $A = x^2 - x + 2$ . The associated SRS parameter is  $\mathbf{r} = (\frac{1}{2}, -\frac{1}{2})$ . The central SRS tile  $\mathcal{T}_{\mathbf{r}}(\mathbf{0})$  as well as its neighbors are shown in Figure 4 on the left hand side. To obtain  $\mathcal{G}_A(\Psi^{-1}(\mathbf{z})) = V\mathbf{z} + \mathcal{F}_A$ , we have to multiply the SRS tiles by the matrix  $V = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$ . The Brunotte tiles are shown on the right hand side of Figure 4.




FIGURE 4. The SRS tiles  $\mathcal{T}_{\mathbf{r}}(\mathbf{z})$  for  $\mathbf{r} = (\frac{1}{2}, -\frac{1}{2})$ ,  $\|\mathbf{z}\|_{\infty} \leq 1$ , (left) and the Brunotte tiles  $\mathcal{G}_A(\Psi^{-1}(\mathbf{z})) = V\mathbf{z} + \mathcal{F}_A$  associated with  $A = x^2 - x + 2$  (right).

An example of SRS tiles associated with the parameter  $\mathbf{r} = (3/4, 1)$  is depicted in Figure 3. This parameter is related to a non-monic CNS.

**5.4. Rational base number systems.** Akiyama *et al.* [9] considered expansions of integers in rational bases  $p/q$ , with coprime integers  $p > q \geq 1$ , of the form

$$N = \frac{1}{q} \sum_{n=0}^{\infty} b_n \left(\frac{p}{q}\right)^n \quad (b_n \in \mathcal{N} = \{0, \dots, p-1\}).$$

In our setting, the sequence  $(b_n)_{n \in \mathbb{N}}$  is the  $(A, \mathcal{N})$ -representation of  $qN$ , where  $A = -qx + p$ . The Brunotte basis modulo  $A$  is given by  $\{-q\}$ . By Lemma 5.5, the corresponding SRS  $\tau_{-q/p}(N) = -\lfloor -N \frac{q}{p} \rfloor$  yields  $b_n = \{-\frac{q}{p} \tau_{-q/p}^n(-N)\}p$ . (Here we write one-dimensional vectors as scalars.) In view of Theorem 4.9, we obtain that the collection of tiles associated with these number systems forms a tiling for each choice  $p/q$  with  $p > q \geq 1$ .

For the case  $p/q = 3/2$ , we show that the collection  $\{\mathcal{T}_{-2/3}(N) \mid N \in \mathbb{Z}\}$  consists of (possibly degenerate) intervals with infinitely many different lengths.

**Lemma 5.17.** *Let  $I$  be a finite set of consecutive integers. Then  $\tau_{-2/3}^{-n}(I)$  is a finite set of consecutive integers for all  $n \in \mathbb{N}$ , and we have*

$$(\#I - 1) \left(\frac{3}{2}\right)^n + 1 \leq \#\tau_{-2/3}^{-n}(I) \leq (\#I + 1) \left(\frac{3}{2}\right)^n - 1.$$

*Proof.* By the definition of  $\tau_{-2/3}$ , the preimage of every finite set of consecutive integers is a finite set of consecutive integers (see also the proof of Theorem 4.9). We have  $\#\tau_{-2/3}^{-1}(N) = 2$  if and only if  $N$  is even and  $\#\tau_{-2/3}^{-1}(N) = 1$  if  $N$  is odd. Therefore the inequalities hold for  $n = 1$ , and by induction for all  $n \in \mathbb{N}$ .  $\square$

**Lemma 5.18.** *For every  $k \geq 1$ , there exists some  $N_k \in \mathbb{Z}$  such that  $\#\tau_{-2/3}^{-k}(N_k) = 2$ .*

*Proof.* It follows from [9, Proposition 10] that there exists some  $L_k \in \mathbb{Z}$  such that the  $(-2x + 3, \{0, 1, 2\})$ -representation of  $2L_k$  satisfies  $b_0 = 0, b_n = 1$  for  $1 \leq n < k$ . The SRS representation of  $-L_k$  with respect to  $-2/3$  is thus given by  $(b_0/3, b_1/3, \dots) = (0, 1/3, \dots, 1/3, b_k/3, b_{k+1}/3, \dots)$ . Let  $N_k = \tau_{-2/3}^k(-L_k)$ . For  $1 \leq n < k$ , the set  $\tau_{-2/3}^{-n}(N_k)$  consists only of the number with SRS representation  $(1/3, \dots, 1/3, b_k/3, b_{k+1}/3, \dots)$  because  $(1/3 + (-2/3)\mathbb{Z}) \cap [0, 1) = \{1/3\}$ . Thus  $\tau_{-2/3}^{-k}(N_k)$  consists exactly of the two numbers  $-L_k$  and  $-L_k - 1$ , which have the SRS representations  $(0, 1/3, \dots, 1/3, b_k/3, b_{k+1}/3, \dots)$  and  $(2/3, 1/3, \dots, 1/3, b_k/3, b_{k+1}/3, \dots)$ , respectively.  $\square$

**Proposition 5.19.** *For every  $k \geq 1$ , there exists some  $N_k \in \mathbb{Z}$  such that*

$$\left(\frac{3}{2}\right)^{-k} \leq \lambda_1(\mathcal{T}_{-2/3}(N_k)) \leq 3 \left(\frac{3}{2}\right)^{-k},$$

where  $\lambda_1$  denotes the one-dimensional Lebesgue measure.

*Proof.* We have

$$\lambda_1(\mathcal{T}_{-2/3}(N_k)) = \lim_{n \rightarrow \infty} \left(\frac{2}{3}\right)^n \#\tau_{-2/3}^{-n}(N_k) = \lim_{n \rightarrow \infty} \left(\frac{2}{3}\right)^n \#\tau_{-2/3}^{k-n}(\tau_{-2/3}^{-k}(N_k))$$

for all  $N_k \in \mathbb{Z}$ . For the  $N_k$  given by Lemma 5.18, we have  $\#\tau_{-2/3}^{-k}(N_k) = 2$ . Using Lemma 5.17, the inequalities are proved.  $\square$

Summing up we get the following result.

**Corollary 5.20.** *The tiling  $\{\mathcal{T}_{-2/3}(N) \mid N \in \mathbb{Z}\}$  consists of (possibly degenerate) intervals with infinitely many different lengths.*

## 6. SRS AND BETA-EXPANSION

We prove now that there is a tight relation with beta-tiles when  $\mathbf{r}$  is related to a unit Pisot number, and that SRS tiles are new objects in the non-unit case (similarly as shown in Section 5 in the non-monic CNS case).

**6.1. Beta-expansions and SRS representations.** We start with beta-expansions, which were first studied by Rényi [35] and Parry [32]. For a real number  $\beta > 1$ , the  $\beta$ -transformation  $T_\beta : [0, 1) \rightarrow [0, 1)$  is defined by  $T_\beta(x) = \{\beta x\} = \beta x - \lfloor \beta x \rfloor$ . The  $\beta$ -expansion of  $x \in [0, 1)$  is

$$x = \sum_{n=1}^{\infty} b_n \beta^{-n} \quad \text{with} \quad b_n = \lfloor \beta T_\beta^{n-1}(x) \rfloor \quad \text{for all } n \geq 1.$$

The following relation between  $T_\beta$  and  $\tau_{\mathbf{r}}$  was shown in [5], see also [20].

**Proposition 6.1.** *Let  $\beta > 1$  be an algebraic integer with minimal polynomial*

$$(6.1) \quad x^{d+1} + a_d x^d + \dots + a_1 x + a_0 = (x - \beta)(x^d + r_{d-1} x^{d-1} + \dots + r_1 x + r_0)$$

and  $\mathbf{r} = (r_0, \dots, r_{d-1})$ . Then we have

$$(6.2) \quad \{\mathbf{r} \tau_{\mathbf{r}}^n(\mathbf{z})\} = T_\beta^n(\{\mathbf{r} \mathbf{z}\}) \quad \text{for all } \mathbf{z} \in \mathbb{Z}^d, n \in \mathbb{N}.$$

In particular, the restriction of  $T_\beta$  to  $\mathbb{Z}[\beta] \cap [0, 1)$  is conjugate to  $\tau_{\mathbf{r}}$ .

*Proof.* Let  $\mathbf{z} = (z_0, \dots, z_{d-1})^t \in \mathbb{Z}^d$ . If we set  $z_d = -\lfloor \mathbf{r}\mathbf{z} \rfloor$ , then we have

$$(6.3) \quad \{\mathbf{r}\mathbf{z}\} = (r_0, \dots, r_{d-1}, 1)(z_0, \dots, z_{d-1}, z_d)^t.$$

Furthermore,  $(r_0, \dots, r_{d-1}, 1)$  is a left eigenvector of the companion matrix  $M_{(a_0, \dots, a_d)}$ , in particular

$$(6.4) \quad (r_0, \dots, r_{d-1}, 1)M_{(a_0, \dots, a_d)} = \beta(r_0, \dots, r_{d-1}, 1).$$

Using (6.3), (6.4) and the fact that  $M_{(a_0, \dots, a_d)}(z_0, \dots, z_d)^t = (z_1, \dots, z_d, m)^t$  with  $m \in \mathbb{Z}$ , we gain

$$\{\mathbf{r}\tau_{\mathbf{r}}(\mathbf{z})\} = \{\mathbf{r}(z_1, \dots, z_d)^t\} = \{(r_0, \dots, r_{d-1}, 1)M_{(a_0, \dots, a_d)}(z_0, \dots, z_d)^t\} = \{\beta\{\mathbf{r}\mathbf{z}\}\} = T_{\beta}(\{\mathbf{r}\mathbf{z}\}),$$

Inductively, we obtain (6.2). Since the polynomial in (6.1) is irreducible,  $\{r_0, \dots, r_{d-1}, 1\}$  is a basis of  $\mathbb{Z}[\beta]$ . Therefore the map  $f : \mathbb{Z}^d \rightarrow \mathbb{Z}[\beta] \cap [0, 1)$ ,  $\mathbf{z} \mapsto \{\mathbf{r}\mathbf{z}\}$  is bijective, and we have  $f \circ \tau_{\mathbf{r}} = T_{\beta} \circ f$ .  $\square$

**Corollary 6.2.** *Let  $\beta$  and  $\mathbf{r}$  be defined as in Proposition 6.1,  $(v_1, v_2, v_3, \dots)$  be the SRS representation of  $\mathbf{z} \in \mathbb{Z}^d$  and  $\{\mathbf{r}\mathbf{z}\} = \sum_{n=1}^{\infty} b_n \beta^{-n}$  be the  $\beta$ -expansion of  $v_1 = \{\mathbf{r}\mathbf{z}\}$ . Then we have*

$$v_n = T_{\beta}^{n-1}(\{\mathbf{r}\mathbf{z}\}) \quad \text{and} \quad b_n = \beta v_n - v_{n+1} \quad \text{for all } n \geq 1.$$

*Proof.* By Definition 2.4 and (6.2), we have  $v_n = \{\mathbf{r}\tau_{\mathbf{r}}^{n-1}(\mathbf{z})\} = T_{\beta}^{n-1}(\{\mathbf{r}\mathbf{z}\})$ , which yields the first assertion. Using this equation and the definition of the  $\beta$ -expansion, we obtain

$$b_n = \lfloor \beta T_{\beta}^{n-1}(\{\mathbf{r}\mathbf{z}\}) \rfloor = \beta T_{\beta}^{n-1}(\{\mathbf{r}\mathbf{z}\}) - \{\beta T_{\beta}^{n-1}(\{\mathbf{r}\mathbf{z}\})\} = \beta v_n - T_{\beta}^n(\{\mathbf{r}\mathbf{z}\}) = \beta v_n - v_{n+1}. \quad \square$$

The eigenvalues of  $M_{\mathbf{r}}$  are exactly the Galois conjugates of  $\beta$ . It follows by Lemma 2.3 that  $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  if  $\beta$  is a Pisot number.

**Definition 6.3** (Finiteness property (F)). A number  $\beta > 1$  is said to have the *finiteness property (F)* if the  $\beta$ -expansion of every  $x \in \mathbb{Z}[\beta^{-1}] \cap [0, 1)$  is finite, i.e.,  $T_{\beta}^n(x) = 0$  for some  $n \in \mathbb{N}$ .

Frougny and Solomyak [19] proved that (F) implies that  $\beta$  is a Pisot number. By [3, Lemma 3], it is sufficient to consider  $x \in \mathbb{Z}[\beta] \cap [0, 1)$  in Definition 6.3 (note that in the present section  $\mathbb{Z}[\beta] \cap [0, 1)$  plays the same role as  $\Lambda_A$  plays in Section 5). Therefore Proposition 6.1 implies the following result (see also [5, Theorem 2.1]).

**Proposition 6.4.** *Let  $\beta > 1$  be an algebraic integer and let  $\mathbf{r}$  be as in Proposition 6.1. Then the following assertions hold.*

- $\mathbf{r} \in \text{int}(\mathcal{D}_d)$  if and only if  $\beta$  is a Pisot number.
- $\mathbf{r} \in \mathcal{D}_d^{(0)}$  if and only if  $\beta$  satisfies (F).

**6.2. Beta-tiles.** For any Pisot number  $\beta$  of degree  $d + 1$ , Akiyama [3] defined a family of tiles covering  $\mathbb{R}^d$  which is conjectured to be always a tiling if  $\beta$  is a unit Pisot number, i.e., if  $|a_0| = 1$  in (6.1). If  $\beta$  is not a unit, then this family cannot be a tiling. In analogy with the previous section, we modify the definition of the tiles so that we obtain tilings also for non-unit Pisot numbers.

Let  $\beta_1, \dots, \beta_d$  be the  $d = r + 2s$  Galois conjugates of  $\beta$ , such that  $\beta_1, \dots, \beta_r \in \mathbb{R}$  and  $\beta_{r+1}, \dots, \beta_{r+2s} \in \mathbb{C}$  with  $\beta_{r+1} = \overline{\beta_{r+s+1}}, \dots, \beta_{r+s} = \overline{\beta_{r+2s}}$ . For  $x \in \mathbb{Q}(\beta)$ ,  $1 \leq j \leq d$ , denote by  $x^{(j)} \in \mathbb{Q}(\beta_j)$  the corresponding conjugate of  $x$ , and let

$$\Phi_{\beta} : \mathbb{Q}(\beta) \rightarrow \mathbb{R}^d, \quad x \mapsto (x^{(1)}, \dots, x^{(r)}, \Re(x^{(r+1)}), \Im(x^{(r+1)}), \dots, \Re(x^{(r+s)}), \Im(x^{(r+s)}))^t.$$

**Definition 6.5** (Beta-tile; see [3]). Let  $\beta$  be a Pisot number. For  $x \in \mathbb{Z}[\beta^{-1}] \cap [0, 1)$ , the set

$$\mathcal{R}_{\beta}(x) := \text{Lim}_{n \rightarrow \infty} \Phi_{\beta}(\beta^n T_{\beta}^{-n}(x)),$$

where the limit is taken with respect to the Hausdorff metric, is called a  $\beta$ -tile. The tile  $\mathcal{R}_{\beta}(0)$  is called *central  $\beta$ -tile*.

Note that the limit in Definition 6.5 exists since  $\Phi_{\beta}(\beta^n T_{\beta}^{-n}(x)) \subseteq \Phi_{\beta}(\beta^{n+1} T_{\beta}^{-n-1}(x))$ . Indeed, if  $y \in [0, 1)$ , then  $T_{\beta}(y\beta^{-1}) = y$ , which implies  $y \in \beta T_{\beta}^{-1}(y)$ .

For unit Pisot numbers, beta-tiles have been studied extensively (mostly under the name ‘‘central tile’’) (see e.g. [2, 3, 40]) and are strongly related to Rauzy fractals associated with unimodular

Pisot substitutions. For a recent survey on these relations and on properties of Rauzy fractals and beta-tiles we refer to [16].

Beta-tiles are known to satisfy a graph-directed IFS equation to be compared with the set equation of Theorem 3.5 (see *e.g.* the survey [16]). Nonetheless the pieces obtained in the decomposition of the central tile might not be measurably disjoint. When  $\beta$  is a unit Pisot number, it is proved that these pieces are disjoint, but when  $\beta$  is not a unit, the pieces do overlap (*cf.* [38]). To make the pieces disjoint in the non-unit case, one can use two different strategies: enlarging the representation space or making the tiles smaller. The first strategy can be found in the literature (see [4, 38]). It consists in adding  $p$ -adic fields for the prime divisors  $p$  of the norm of  $\beta$  to the representation space. In the present work, we want to carry out the second strategy, leading to tiles that form a tiling of  $\mathbb{R}^d$ . To this matter, in the  $p$ -adic extension obtained with the first strategy, we have to choose among all points with the same Euclidean part, a single specific point having  $p$ -adic components with certain properties. Using Proposition 6.1, we will see that SRS tiles actually perform this choice by arithmetical means, that is, by picking points in beta-tiles that come from  $\mathbb{Z}[\beta]$ . (We mention here also Barnsley's study on *fractal tops*, which provide a method to get rid of overlaps in IFS attractors, see [15, Chapter 4].)

Note that beta-tiles can be described as

$$\mathcal{R}_\beta(x) = \lim_{n \rightarrow \infty} \{ \Phi_\beta(\beta^n y) \mid y \in \mathbb{Z}[\beta^{-1}] \cap [0, 1), T_\beta^n(y) = x \}.$$

Proposition 6.1 shows that we have to consider  $\mathbb{Z}[\beta]$  instead of  $\mathbb{Z}[\beta^{-1}]$  in this formula to get a correspondence with SRS tiles (observe again that in the present section  $\mathbb{Z}[\beta] \cap [0, 1)$  plays the same role as  $\Lambda_A$  plays in Section 5).

**Definition 6.6** (Integral beta-tile). Let  $\beta$  be a Pisot number. For  $x \in \mathbb{Z}[\beta] \cap [0, 1)$ , the set

$$\mathcal{S}_\beta(x) := \text{Li}_{n \rightarrow \infty} \Phi_\beta(\beta^n (T_\beta^{-n}(x) \cap \mathbb{Z}[\beta])),$$

where Li is the lower Hausdorff limit defined in Section 3.1, is called *integral  $\beta$ -tile*. The tile  $\mathcal{S}_\beta(0)$  is called *central integral  $\beta$ -tile*.

The difference between this definition and Definition 6.5 is that any approximation of a tile is given just by points  $y \in \mathbb{Z}[\beta] \cap [0, 1)$  with  $T_\beta^n(y) = x$ , instead of considering all points in  $T_\beta^{-n}(x)$ . The limitation to  $\mathbb{Z}[\beta]$  is the core of the selection process. However this implies that in general  $\mathcal{S}_\beta(x)$  is not a graph directed self-affine set. It is obvious that

$$\mathcal{R}_\beta(x) \supseteq \mathcal{S}_\beta(x),$$

where equality holds if and only if  $\beta$  is a unit Pisot number.

**6.3. From integral beta-tiles to SRS tiles.** In the sequel, we will see how SRS-tiles are related to integral beta-tiles by a linear transformation. We will show that SRS tiles provide a decomposition of beta-tiles into disjoint pieces: the process can be seen as selecting an integral representation in each  $p$ -adic leaf of the central tile, for each prime divisor of the norm of  $\beta$ . The main feature here is that this selection of an integral representant can be performed in a dynamical way.

**Theorem 6.7.** *Let  $\beta$  be a Pisot number with minimal polynomial  $(x - \beta)(x^d + r_{d-1}x^{d-1} + \dots + r_0)$  and  $d = r + 2s$  Galois conjugates  $\beta_1, \dots, \beta_r \in \mathbb{R}$ ,  $\beta_{r+1}, \dots, \beta_{r+2s} \in \mathbb{C}$ , ordered such that  $\beta_{r+1} = \overline{\beta_{r+s+1}}, \dots, \beta_{r+s} = \overline{\beta_{r+2s}}$ . Let*

$$x^d + r_{d-1}x^{d-1} + \dots + r_0 = (x - \beta_j)(x^{d-1} + q_{d-2}^{(j)}x^{d-2} + \dots + q_0^{(j)})$$

for  $1 \leq j \leq d$  and

$$U = \begin{pmatrix} q_0^{(1)} & q_1^{(j)} & \cdots & q_{d-2}^{(1)} & 1 \\ \vdots & \vdots & & \vdots & \vdots \\ q_0^{(r)} & q_1^{(j)} & \cdots & q_{d-2}^{(r)} & 1 \\ \Re(q_0^{(r+1)}) & \Re(q_1^{(r+1)}) & \cdots & \Re(q_{d-2}^{(r+1)}) & 1 \\ \Im(q_0^{(r+1)}) & \Im(q_1^{(r+1)}) & \cdots & \Im(q_{d-2}^{(r+1)}) & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ \Re(q_0^{(r+s)}) & \Re(q_1^{(r+s)}) & \cdots & \Re(q_{d-2}^{(r+s)}) & 1 \\ \Im(q_0^{(r+s)}) & \Im(q_1^{(r+s)}) & \cdots & \Im(q_{d-2}^{(r+s)}) & 0 \end{pmatrix} \in \mathbb{R}^{d \times d}.$$

Then we have

$$\mathcal{S}_\beta(\{\mathbf{rx}\}) = U(M_{\mathbf{r}} - \beta I_d)\mathcal{T}_{\mathbf{r}}(\mathbf{x})$$

for every  $\mathbf{x} \in \mathbb{Z}^d$ , where  $\mathbf{r} = (r_0, \dots, r_{d-1})$  and  $I_d$  is the  $d$ -dimensional identity matrix.

*Proof.* Let  $\mathbf{t} \in \mathcal{T}_{\mathbf{r}}(\mathbf{x})$  and  $\mathbf{z}_{-n}, v_{-n}, n \in \mathbb{N}$ , as in Proposition 3.6, i.e.,

$$\mathbf{t} = \mathbf{x} - \sum_{n=0}^{\infty} M_{\mathbf{r}}^n(0, \dots, 0, v_{-n})^t \quad \text{and} \quad v_{-n+1} = \{\mathbf{rz}_{-n}\} \text{ for all } n \in \mathbb{N}.$$

Set  $b_{-n} = \beta v_{-n} - v_{-n+1}$  for  $n \in \mathbb{N}$ . Then we obtain, by using  $\tau_{\mathbf{r}}(\mathbf{x}) = M_{\mathbf{r}}\mathbf{x} + \{\mathbf{rx}\}$  and  $v_1 = \{\mathbf{rx}\}$ ,

$$(6.5) \quad (M_{\mathbf{r}} - \beta I_d)\mathbf{t} = \tau_{\mathbf{r}}(\mathbf{x}) - \beta\mathbf{x} + \sum_{n=0}^{\infty} M_{\mathbf{r}}^n(0, \dots, 0, b_{-n})^t.$$

Similarly to (6.4), we see that  $(q_0^{(j)}, \dots, q_{d-2}^{(j)}, 1)$  is a left eigenvector of  $M_{\mathbf{r}}$ ,

$$(q_0^{(j)}, \dots, q_{d-2}^{(j)}, 1)M_{\mathbf{r}} = \beta_j(q_0^{(j)}, \dots, q_{d-2}^{(j)}, 1) \quad \text{for } 1 \leq j \leq d.$$

By using (6.5), we obtain

$$(6.6) \quad (q_0^{(j)}, \dots, q_{d-2}^{(j)}, 1)(M_{\mathbf{r}} - \beta I_d)\mathbf{t} = (q_0^{(j)}, \dots, q_{d-2}^{(j)}, 1)\left(\tau_{\mathbf{r}}(\mathbf{x}) - \beta\mathbf{x} + \sum_{n=0}^{\infty} \beta_j^n(0, \dots, 0, b_{-n})^t\right).$$

Since the minimal polynomial of  $\beta$  can be decomposed as

$$(x - \beta_j)(x^d + r_{d-1}^{(j)}x^{d-1} + \cdots + r_0^{(j)}) = (x - \beta)(x - \beta_j)(x^{d-1} + q_{d-2}^{(j)}x^{d-2} + \cdots + q_0^{(j)}),$$

we have

$$r_0^{(j)} = -\beta q_0^{(j)}, \quad r_k^{(j)} = q_{k-1}^{(j)} - \beta q_k^{(j)} \text{ for } 1 \leq k \leq d-2, \quad r_{d-1}^{(j)} = q_{d-2}^{(j)} - \beta,$$

and obtain

$$(6.7) \quad (q_0^{(j)}, \dots, q_{d-2}^{(j)}, 1)(\tau_{\mathbf{r}}(\mathbf{x}) - \beta\mathbf{x}) = (r_0^{(j)}, \dots, r_{d-1}^{(j)})\mathbf{x} - [\mathbf{rx}] = \{\mathbf{rx}\}^{(j)}.$$

Inserting (6.7) in (6.6) yields

$$(6.8) \quad \begin{aligned} (q_0^{(j)}, \dots, q_{d-2}^{(j)}, 1)(M_{\mathbf{r}} - \beta I_d)\mathbf{t} &= \{\mathbf{rx}\}^{(j)} + \sum_{n=0}^{\infty} b_{-n}\beta_j^n \\ &= \lim_{n \rightarrow \infty} \left( \{\mathbf{rx}\} + \sum_{k=0}^{n-1} b_{-k}\beta^k \right)^{(j)} = \lim_{n \rightarrow \infty} (\beta^n \{\mathbf{rz}_{-n}\})^{(j)}; \end{aligned}$$

here we used that  $b_{-k} = \beta v_{-k} - v_{-k+1}$ ,  $v_{-n+1} = \{\mathbf{rz}_{-n}\}$  and  $v_1 = \{\mathbf{rx}\}$ . By Proposition 6.1, we have  $\{\mathbf{rz}_{-n}\} \in T_\beta^{-n}(\{\mathbf{rx}\})$  and  $\{\mathbf{rz}_{-n}\} \in \mathbb{Z}[\beta]$ , thus

$$U(M_{\mathbf{r}} - \beta I_d)\mathbf{t} = \lim_{n \rightarrow \infty} \Phi_\beta(\beta^n \{\mathbf{rz}_{-n}\}) \in \mathcal{S}_\beta(\{\mathbf{rx}\}).$$

Now, let  $\mathbf{u} \in \mathcal{S}_\beta(\{\mathbf{rx}\})$ . Then there exists a sequence  $(\mathbf{z}_{-n})_{n \in \mathbb{N}}$  such that  $\{\mathbf{rz}_{-n}\} \in T_\beta^{-n}(\{\mathbf{rx}\}) \cap \mathbb{Z}[\beta]$  and  $\lim_{n \rightarrow \infty} \Phi_\beta(\beta^n \{\mathbf{rz}_{-n}\}) = \mathbf{u}$ . Similarly to Proposition 3.6, we can choose the sequence

$(\mathbf{z}_{-n})_{n \in \mathbb{N}}$  such that  $T_\beta(\{\mathbf{r}\mathbf{z}_{-n}\}) = \{\mathbf{r}\mathbf{z}_{-n+1}\}$  for all  $n \geq 1$ . Set  $b_{-n} = \beta\{\mathbf{r}\mathbf{z}_{-n-1}\} - \{\mathbf{r}\mathbf{z}_{-n}\}$ . Then (6.8) implies that  $\mathbf{u} \in U(M_{\mathbf{r}} - \beta I_d)\mathcal{T}_{\mathbf{r}}(\mathbf{x})$ .  $\square$

*Remark 6.8.* We would like to emphasize that  $U(M_{\mathbf{r}} - \beta I_d)\mathcal{T}_{\mathbf{r}}(\mathbf{x}) = \mathcal{S}_\beta(\{\mathbf{r}\mathbf{x}\})$  does not imply that the “center”  $\mathbf{x}$  of  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  is mapped to the “center”  $\Phi_\beta(\{\mathbf{r}\mathbf{x}\})$  of  $\mathcal{S}_\beta(\{\mathbf{r}\mathbf{x}\})$ . Indeed, by (6.7), we have

$$\Phi_\beta(\{\mathbf{r}\mathbf{x}\}) = U(\tau_{\mathbf{r}}(\mathbf{x}) - \beta\mathbf{x}) = U(M_{\mathbf{r}} - \beta I_d)\mathbf{x} + U(0, \dots, 0, \{\mathbf{r}\mathbf{x}\})^t.$$

In particular, this means that, even though there is only a finite number of shapes in the unit Pisot case, no SRS tile is obtained by a  $\mathbb{Z}^d$ -translation of another SRS tile.

**Corollary 6.9.** *Integral beta-tiles can be written as*

$$\mathcal{S}_\beta(x) = \text{Lim}_{n \rightarrow \infty} \Phi_\beta(\beta^n(T_\beta^{-n}(x) \cap \mathbb{Z}[\beta])),$$

where  $\text{Lim}$  denotes the Hausdorff limit.

We deduce the following reformulation of the set equation in Theorem 3.5.

**Corollary 6.10.** *For a Pisot number  $\beta$  and  $x \in \mathbb{Z}[\beta] \cap [0, 1)$ , we have*

$$\mathcal{S}_\beta(x) = \bigcup_{y \in T_\beta^{-1}(x) \cap \mathbb{Z}[\beta]} \Lambda_\beta \mathcal{S}_\beta(y),$$

where  $\Lambda_\beta = \text{diag}(\beta_1, \dots, \beta_r, \begin{pmatrix} \Re(\beta_{r+1}) & \Im(\beta_{r+1}) \\ -\Im(\beta_{r+1}) & \Re(\beta_{r+1}) \end{pmatrix}, \dots, \begin{pmatrix} \Re(\beta_{r+s}) & \Im(\beta_{r+s}) \\ -\Im(\beta_{r+s}) & \Re(\beta_{r+s}) \end{pmatrix})$ .

The main result of this section extends the results in [3, 23, 24] on tiling properties for unit Pisot numbers to arbitrary Pisot numbers.

**Theorem 6.11.** *Let  $\beta$  be a Pisot number. Then the following assertions hold.*

- *The collection  $\{\mathcal{S}_\beta(x) \mid x \in \mathbb{Z}[\beta]\}$  forms a weak  $m$ -tiling of  $\mathbb{R}^d$  for some  $m \geq 1$ .*
- *If  $\beta$  satisfies the finiteness property (F), then  $\{\mathcal{S}_\beta(x) \mid x \in \mathbb{Z}[\beta]\}$  forms a weak tiling of  $\mathbb{R}^d$ .*

*Proof.* By Theorem 6.7, it is equivalent to consider the collection  $\{\mathcal{T}_{\mathbf{r}}(\mathbf{z}) \mid \mathbf{z} \in \mathbb{Z}^d\}$  for  $\mathbf{r} = (r_0, \dots, r_{d-1})$  such that  $(x - \beta)(x^d + r_{d-1}x^{d-1} + \dots + r_0)$  is the minimal polynomial of  $\beta$ . In view of Proposition 6.4, the first assertion follows from Theorem 4.6, while the second assertion is a consequence of Corollary 4.7.  $\square$

*Example 6.12.* Consider the polynomial  $x^3 - 3x^2 + 1$ . Its largest root in modulus is  $\beta \approx 2.879$ , a unit Pisot number. Set  $\mathbf{r} = (r_0, r_1)$  with  $r_0 = -1/\beta$ ,  $r_1 = -1/\beta^2$ . The 25 SRS-tiles  $\mathcal{T}_{\mathbf{r}}(\mathbf{x}) \subset \mathbb{R}^2$  with  $\|\mathbf{x}\|_\infty \leq 2$  are shown in Figure 5 on the left.

Let  $\beta_1, \beta_2$  be the Galois conjugates of  $\beta$ . The numbers  $\beta_1$  and  $\beta_2$  are real numbers and thus the (integral)  $\beta$ -tiles  $\mathcal{R}_\beta(\{\mathbf{r}\mathbf{x}\}) = \mathcal{S}_\beta(\{\mathbf{r}\mathbf{x}\})$  are obtained by multiplying  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  by the matrix

$$U(M_{\mathbf{r}} - \beta I_2) = \begin{pmatrix} -\beta_2 & 1 \\ -\beta_1 & 1 \end{pmatrix} \left( \begin{pmatrix} 0 & 1 \\ -r_0 & -r_1 \end{pmatrix} - \begin{pmatrix} \beta & 0 \\ 0 & \beta \end{pmatrix} \right).$$

They are shown on the right hand side of Figure 5.

## REFERENCES

- [1] B. ADAMCZEWSKI, C. FROUGNY, A. SIEGEL, AND W. STEINER, *Rational numbers with purely periodic beta-expansion*, preprint, (2009).
- [2] S. AKIYAMA, *Self affine tilings and Pisot numeration systems*, in Number Theory and its Applications, K. Györy and S. Kanemitsu, eds., Kluwer, 1999, pp. 1–17.
- [3] S. AKIYAMA, *On the boundary of self affine tilings generated by Pisot numbers*, J. Math. Soc. Japan, 54 (2002), pp. 283–308.
- [4] S. AKIYAMA, G. BARAT, V. BERTHÉ, AND A. SIEGEL, *Boundary of central tiles associated with Pisot beta-numeration and purely periodic expansions*, Monatsh. Math., 155 (2008), pp. 377–419.
- [5] S. AKIYAMA, T. BORBÉLY, H. BRUNOTTE, A. PETHŐ, AND J. M. THUSWALDNER, *Generalized radix representations and dynamical systems. I*, Acta Math. Hungar., 108 (2005), pp. 207–238.
- [6] S. AKIYAMA, H. BRUNOTTE, A. PETHŐ, AND J. M. THUSWALDNER, *Generalized radix representations and dynamical systems. II*, Acta Arith., 121 (2006), pp. 21–61.




FIGURE 5. SRS tiles  $\mathcal{T}_{\mathbf{r}}(\mathbf{x})$  associated with  $\mathbf{r} = (-1/\beta, -1/\beta^2)$ ,  $\beta^3 = 3\beta^2 - 1$ , (left) and the corresponding  $\beta$ -tiles  $\mathcal{R}_{\beta}(\{\mathbf{r}\mathbf{x}\}) = \mathcal{S}_{\beta}(\{\mathbf{r}\mathbf{x}\})$  (right).

- [7] ———, *Generalized radix representations and dynamical systems. III*, Osaka J. Math., 45 (2008), pp. 347–374.
- [8] ———, *Generalized radix representations and dynamical systems. IV*, Indag. Math. (N.S.), to appear (2009).
- [9] S. AKIYAMA, C. FROUGNY, AND J. SAKAROVITCH, *Powers of rationals modulo 1 and rational base number systems*, Israel J. Math., 168 (2008), pp. 53–91.
- [10] S. AKIYAMA AND K. SCHEICHER, *Intersecting two dimensional fractals and lines*, Acta Sci. Math. (Szeged), 71 (2005), pp. 555–580.
- [11] S. AKIYAMA AND J. M. THUSWALDNER, *Topological properties of two-dimensional number systems*, J. Théor. Nombres Bordeaux, 12 (2000), pp. 69–79.
- [12] ———, *A survey on topological properties of tiles related to number systems*, Geom. Dedicata, 109 (2004), pp. 89–105.
- [13] G. BARAT, V. BERTHÉ, P. LIARDET, AND J. THUSWALDNER, *Dynamical directions in numeration*, Ann. Inst. Fourier (Grenoble), 56 (2006), pp. 1987–2092.
- [14] M. BARNSLEY, *Fractals Everywhere*, Academic Press INC., Orlando, 1988.
- [15] ———, *Superfractals*, Cambridge University Press, 2006.
- [16] V. BERTHÉ AND A. SIEGEL, *Tilings associated with beta-numeration and substitutions*, Integers, 5 (2005), pp. A2, 46 pp. (electronic).
- [17] N. CHEKHOVA, P. HUBERT, AND A. MESSAOUDI, *Propriétés combinatoires, ergodiques et arithmétiques de la substitution de Tribonacci*, J. Théor. Nombres Bordeaux, 13 (2001), pp. 371–394.
- [18] K. J. FALCONER, *Fractal Geometry*, John Wiley and Sons, Chichester, 1990.
- [19] C. FROUGNY AND B. SOLOMYAK, *Finite beta-expansions*, Ergodic Theory Dynam. Systems, 12 (1992), pp. 713–723.
- [20] M. HOLLANDER, *Linear Numeration Systems, Finite Beta Expansions, and Discrete Spectrum of Substitution Dynamical Systems*, PhD. thesis, Washington University, Seattle, 1996.
- [21] P. HUBERT AND A. MESSAOUDI, *Best simultaneous diophantine approximations of Pisot numbers and Rauzy fractals*, Acta Arith., 124 (2006), pp. 1–15.
- [22] J. E. HUTCHINSON, *Fractals and self-similarity*, Indiana Univ. Math. J., 30 (1981), pp. 713–747.

- [23] S. ITO AND H. RAO, *Atomic surfaces, tilings and coincidence. I. Irreducible case*, Israel J. Math., 153 (2006), pp. 129–155.
- [24] C. KALLE AND W. STEINER, *Beta-expansions, natural extensions and multiple tilings*, preprint.
- [25] I. KÁTAI AND I. KÖRNYEI, *On number systems in algebraic number fields*, Publ. Math. Debrecen, 41 (1992), pp. 289–294.
- [26] D. E. KNUTH, *The Art of Computer Programming, Vol 2: Seminumerical Algorithms*, Addison Wesley, London, 3rd ed., 1998.
- [27] B. KOVÁCS AND A. PETHŐ, *Number systems in integral domains, especially in orders of algebraic number fields*, Acta Sci. Math. (Szeged), 55 (1991), pp. 286–299.
- [28] K. KURATOWSKI, *Topology*, vol. I, Academic Press, Polish Scientific Publishers, New York, London, Warszawa, 1966.
- [29] J. LAGARIAS AND Y. WANG, *Self-affine tiles in  $\mathbb{R}^n$* , Adv. Math., 121 (1996), pp. 21–49.
- [30] ———, *Integral self-affine tiles in  $\mathbb{R}^n$  II. lattice tilings*, J. Fourier Anal. Appl., 3 (1997), pp. 83–102.
- [31] R. D. MAULDIN AND S. C. WILLIAMS, *Hausdorff dimension in graph directed constructions*, Trans. Amer. Math. Soc., 309 (1988), pp. 811–829.
- [32] W. PARRY, *On the  $\beta$ -expansions of real numbers*, Acta Math. Acad. Sci. Hungar., 11 (1960), pp. 401–416.
- [33] A. PETHŐ, *On a polynomial transformation and its application to the construction of a public key cryptosystem*, in Computational number theory (Debrecen, 1989), de Gruyter, Berlin, 1991, pp. 31–43.
- [34] G. RAUZY, *Nombres algébriques et substitutions*, Bull. Soc. Math. France, 110 (1982), pp. 147–178.
- [35] A. RÉNYI, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hungar., 8 (1957), pp. 477–493.
- [36] K. SCHEICHER, P. SURER, J. M. THUSWALDNER, AND C. VAN DE WOESTIJNE, *Digit systems over commutative rings*, preprint.
- [37] K. SCHEICHER AND J. M. THUSWALDNER, *Canonical number systems, counting automata and fractals*, Math. Proc. Cambridge Philos. Soc., 133 (2002), pp. 163–182.
- [38] A. SIEGEL, *Représentation des systèmes dynamiques substitutifs non unimodulaires*, Ergodic Theory Dynam. Systems, 23 (2003), pp. 1247–1273.
- [39] P. SURER, *Characterisation results for shift radix systems*, Math. Pannon., 18 (2007), pp. 265–297.
- [40] W. P. THURSTON, *Groups, tilings and finite state automata*. AMS Colloquium lectures, 1989.
- [41] A. VINCE, *Digit tiling of euclidean space*, in Directions in Mathematical Quasicrystals, Providence, RI, 2000, Amer. Math. Soc., pp. 329–370.
- [42] Y. WANG, *Self-affine tiles*, in Advances in Wavelet, K. S. Lau, ed., Springer, 1998, pp. 261–285.

LIRMM, CNRS UMR 5506, UNIVERSITÉ MONTPELLIER II, 161 RUE ADA, 34392 MONTPELLIER CEDEX 5, FRANCE

*E-mail address:* `berthe@lirmm.fr`

IRISA, CAMPUS DE BEAULIEU, 35042 RENNES CEDEX, FRANCE

*E-mail address:* `Anne.Siegel@irisa.fr`

LIAFA, CNRS UMR 7089, UNIVERSITÉ PARIS DIDEROT – PARIS 7, CASE 7014, 75205 PARIS CEDEX 13, FRANCE

*E-mail address:* `steiner@liafa.jussieu.fr`

CHAIR OF MATHEMATICS AND STATISTICS, UNIVERSITY OF LEOBEN, A-8700 LEOBEN, AUSTRIA

*E-mail address:* `me@palovsky.com`

CHAIR OF MATHEMATICS AND STATISTICS, UNIVERSITY OF LEOBEN, A-8700 LEOBEN, AUSTRIA

*E-mail address:* `joerg.thuswaldner@mu-leoben.at`