HAL open science

Combinatorial map as multiplication of combinatorial knots

Dainis Zeps

▶ To cite this version:

Dainis Zeps. Combinatorial map as multiplication of combinatorial knots. 2007. hal-00408153

HAL Id: hal-00408153
https://hal.science/hal-00408153v1

Preprint submitted on 29 Jul 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Combinatorial map as multiplication of combinatorial knots

Dainis ZEPS *

Abstract

We show that geometrical map can be expressed as multiplication of combinatorial maps, i.e. map P is equal to multiplication of its knot, inner knot's square and trivial knot ($= \mu \cdot \nu^2 \cdot \pi_1$).

1 Introduction

We proceed with building combinatorial map theory that from different points of view and formulations is considered in from [1] to [35].

We multiply permutations from left to right. Geometrical *combinatorial map* is pair of permutations, *vertex and face rotations*, (P, Q) acting on set of elements C if $P \cdot Q^{-1} = \rho$ *edge rotation* or *inner edge rotation* $\pi = Q^{-1}P$ is involution without fixed elements. We consider set of maps with fixed π calling them *normalized maps*. Mostly we use one particular choice of π equal to (12) ... $(2k-1\ 2k)$, $k > 1$. If so, map may be characterized with one permutation, say vertex rotation P.

In [30] we saw that particular choice of ρ by fixed π induces partitioning of the set C into to subsets C_1 and C_2 [in general in several ways] so that the *knot* $\mu = \left\{ \begin{array}{ll} C_1: & \pi \\ C_2: & \rho \end{array} \right.$ is defined. Here, knot μ as permutation has 2^k choices if k is number of cycles in it. Rightly, changing direction of some cycle of μ we get another possible value for knot μ. Moreover, ρ with choice of particular μ partitions π into $\pi_1 \cdot \pi_2$, where we call π_1 *cut edges* and π_2 *cycle edges*, so that $P \cdot \pi_1 : C_1 \mapsto C_2$ and $P \cdot \pi_2 : C_1 \mapsto C_1$. In [31] was shown that by fixing μ map P may be expressed as multiplication $\gamma_1 \cdot \gamma_2 \cdot \pi_2$, where γ_1 acts within C_1 and γ_2 acts within C_2.

In [30] was shown that normalized map always may be expressed as $P = \mu \cdot \alpha$, where α is called *knotting* and it is selfconjugate map in sense that $\alpha^\pi = \alpha$. In [32] we got formulas for μ and α, i.e.,

$$\mu = \gamma_2 \pi \gamma_1^{-1}$$

and

$$\alpha = \gamma_1 \gamma_1^\pi.$$

From [30] we know that α's form a group K_π with respect to multiplication of maps. Moreover, classes of maps with fixed ρ's, denoted as K_ρ, are cosets (left and right) of K_π.

2 Main part

We are going to regain main formulae from introduction.

Let us prove some theorems that leads us to the main result.

Theorem 1. *$\rho \cdot \pi$ [or $\pi \cdot \rho$] is equal to some combinatorial knot μ squared and one or other color cycles induced from this knot reversed.*

Proof. Let us write knot μ in the form $\left\{ \begin{array}{ll} C_1: & \pi \\ C_2: & \rho \end{array} \right.$. Then square of μ we would get applying $\pi \cdot \rho$ for one color corners and $\rho \cdot \pi$ for other color corners. □

Theorem 2. *By fixing the square of the knot it has 2^k knots in correspondence [in general for different maps] where k is the number of cycles in the knot.*

Proof. Two joined cycles of square of knot may be combined in the cycle of knot in two ways, and thus, k independent operations give 2^k results. □

*Author's address: Institute of Mathematics and Computer Science, University of Latvia, 29 Rainis blvd., Riga, Latvia. dainize@mii.lu.lv

Theorem 3. *[$\mu \cdot \pi$ is knot's half-square]*
1) $\mu \cdot \pi$ contains squared knot's cycles of only one color.
2) For vertex rotation $\mu \cdot \pi$ corresponding face rotation and knot are equal to μ, and knotting equal to π. $\gamma_1 = id$, $\gamma_2 = \mu \cdot \pi$, and $\pi_1 = \pi$, because all edges of this map are cut edges.

Proof. 1) μ expressing as $\left\{ \begin{array}{ll} C_1: & \pi \\ C_2: & \rho \end{array} \right.$, and multiplying by π, we get $\left\{ \begin{array}{ll} C_1: & \pi \cdot \pi \\ C_2: & \rho \cdot \pi \end{array} \right.$ and using theorem 1 what was to be proved.
2) Corresponding graph to this map is set of star graphs as many as cycles in μ. Direct calculation gives what is stated by theorem. □

Theorem 4. *Map P can be expressed as $P\pi_1 = \gamma_1\gamma_2\pi = \gamma_2\gamma_1\pi$ with $\mu(P) = \gamma_2\pi\gamma_1^{-1}[= \gamma_1\rho\gamma_2^{-1}]$ and π_1 as inner cut edge rotation and π_2 inner cycle edge rotation.*

Proof. Let knot $\mu = \mu(P)$ be fixed. Then set of corners is partitioned into two sets C_1 and C_2. From form of $\mu(= \gamma_2 \cdot \pi \cdot \gamma_1^{-1})$ we directly judge that γ_1 belongs to, say, C_1 and γ_2 to C_2. Thus, γ_1 and γ_2 commute by multiplying. Let us choose vertex rotation with this fixed knot and $\pi_1 = id$, i.e., with all edges being cycle edges. Then vertex rotation is alternation of corners from C_1 and C_2 respectively, and face rotation's cycles are correspondingly of one color. Then form of $\mu = \gamma_2 \cdot \pi \cdot \gamma_1^{-1}$ shows directly that $P \cdot \pi_1$ must be equal to $\gamma_1 \cdot \gamma_2(= \gamma_2 \cdot \gamma_1)$. Finally, in general we get

$$\mu = \left\{ \begin{array}{ll} C_1: & \pi \\ C_2: & \rho \end{array} \right. = \left\{ \begin{array}{l} C_2 : \gamma_2\pi\gamma_1^{-1} \\ C_1 : \gamma_1\rho\gamma_2^{-1} \end{array} \right. .$$

□

Theorem 5. *Map $P \cdot \pi_1$ can be expressed as $\left\{ \begin{array}{ll} C_1: & \beta_1 \\ C_2: & \beta_2 \end{array} \right.$, where involutions β_1 and β_2 are equal to $\beta_1 = \pi^{-\gamma_1}$ and $\beta_2 = \pi^{-\gamma_2}$. Moreover, $\beta_1 = \gamma_1\gamma_2^{-1}\mu$ and $\beta_2 = \gamma_2\gamma_1^{-1}\mu$. Moreover, $\beta_1\beta_2$ is squared knot $\mu(..., \beta_1)$ with one color cycles reversed. See theorem 1. $\delta = \pi^{\gamma_1}$. $P = \gamma_1\gamma_2\pi_2$.*

Proof.

$$P\pi_1 = \left\{ \begin{array}{ll} C_1: & \beta_1 \\ C_2: & \beta_2 \end{array} \right. = \left\{ \begin{array}{ll} C_1: & \gamma_1\pi\gamma_1^{-1} \\ C_2: & \gamma_2\pi\gamma_2^{-1} \end{array} \right. = (\left\{ \begin{array}{ll} C_1: & \gamma_1 \\ C_2: & \gamma_2 \end{array} \right.) \cdot \pi = \gamma_1\gamma_2\pi.$$

□

Corollary 6. *Map $P\pi_1$ is a knot for inner edge rotation β_1 and edge rotation β_2.*

Theorem 7. *Let for some fixed knot the map P be equal to $\mu\alpha$. Then α is equal to $\gamma_1\pi\gamma_1\pi_2$ or $\gamma_1\gamma_1^{\pi}\pi_1$.*

Proof. $\alpha = \mu^{-1}P\pi_1 = \gamma_1\pi\gamma_2^{-1}\gamma_2\gamma_1\pi = \gamma_1\pi\gamma_1\pi = \gamma_1\gamma_1^{\pi}$. This α is knotting for $P\pi_1$. For map P knotting is $\gamma_1\pi\gamma_1\pi_2$ or $\gamma_1\gamma_1^{\pi}\pi_1$. □

Corollary 8. *$\alpha^{\pi} = \alpha$. Selfconjugate α's comprise group.*

Proof. $\alpha^{\pi} = (\gamma_1\gamma_1^{\pi}\pi_1)^{\pi} = \gamma_1^{\pi}\gamma^{\pi\pi}\pi_1^{\pi} = \gamma_2\gamma_1\pi_1 = \alpha$. □

Theorem 9. *$\gamma_1\gamma_1^{\pi}$ is some knot's square.*

Proof. Let us denote this knot by ν. Direct observation shows that theorem is correct. Then fixed knot μ induces α and it determines fixed ν such that $\nu^2 = \gamma_1\gamma_1^{\pi}$. □

Theorem 10. *Every combinatorial P can be expressed as multiplication of knots in the form*

$$P = \mu \cdot \nu^2 \cdot \pi_1.$$

Proof. It directly follows from previous theorems. Really, $P = \mu\alpha = \gamma_2\pi\gamma_1^{-1}\gamma_1\gamma_1^{\pi}\pi_1 = \gamma_2\gamma_1\pi_2 = \mu(\gamma_1\pi)^2\pi_1 = \mu\nu^2\pi_1$. □

It must be noted that π_1 is some knot too. We call this knot trivial knot. Let us call knot ν map's inner knot.

Corollary 11. *Map is multiplication of its knot with its inner knot's square and with its trivial knot.*

Theorem 12. *For $P\pi_1$ μ commutes with α, i.e.,*

$$P\pi_1 = \mu \cdot \alpha = \alpha \cdot \mu.$$

In general,

$$P = \mu\alpha = \alpha\mu^{\pi_1}.$$

Proof. For $P\pi_1$, $\mu^\alpha = (\gamma_2\pi\gamma_1^{-1})^{\gamma_1\pi\gamma_1\pi}$. Further, $\gamma_2^{\gamma_1\pi\gamma_1\pi} = \gamma_2^{\pi\cdot\pi} = \gamma_2$, because corners of γ_1 and γ_2 do not intersect. The same is true for the member γ_1^{-1}. Further, $\pi^{\gamma_1\pi\gamma_1\pi} = \pi^\alpha = \pi$. Thus, we get $\mu^\alpha = \mu$. □

Theorem 13. *For partial map $[P, \mu]$ its inner edge rotation is α.*

Proof. Direct observation. □

3 Conclusions

There are four types of permutations that are used to build "all" in combinatorial map theory, i.e., knot-type, knot-square-type, knot-square-with-reversed-cycles-type, two-color-involutions. Comprehensive algebra of all these types should be ground for combinatorial map theory.

References

[1] P. Bonnington, C.H.C. Little, *Fundamentals of topological graph theory*, Springer-Verlag, N.Y.,1995.

[2] G. Burde, H. Zieschang. *Knots*, Walter de Gruiter, Berlin N.Y., 1985.

[3] R. Cori. *Un Code pour les Graphes Planaires et ses Applications.* Astérisque, 1975, vol. 27.

[4] R. Cori, A. Machi. *Maps, Hypermaps and their Automorphisms: A Survey, I, II, III.* Expositiones Mathematicae, 1992, vol. 10, 403-427, 429-447, 449-467.

[5] J. K. Edmonds. *A combinatorial representation for polyhedral surfaces*, Notices Amer. Math. Soc. (1960), 646.

[6] M. Ferri, C. Gagliardi. *Cristallisation Moves*, Pacific Journ. Math., Vol.100. No 1, 1982.

[7] P. J. Giblin. *Graphs, Surfaces, and Homology.* John Willey & Sons, 1977.

[8] L. Heffter. Über das Problem der Nachbargebiete. Math. Ann., 1891, vol. 38, 477-508.

[9] L. Heffter. *Uber metacyklische Gruppen und Nachbarconfigurationen*, Math. Ann. 50, 261- 268, 1898.

[10] A. Jacques. *Sur le genre dune paire de substitutions*, C.R.Acad. Sci.Paris ser:I Math. **367**, 625-627,1968.

[11] P. Kikusts, D. Zeps. *Graphs on surfaces*, Conf. LMS, Riga, 1994.

[12] Sergei K. Lando , Alexander K. Zvonkin. *Graphs on Surphaces and Their Applications.* Springer, 2003.

[13] S. Lins. *Graph Encoded Maps*, Journ.Comb.Theory, Series B 32, 171-181, 1982.

[14] C.H.C. Little. *Cubic combinatorial maps*, J.Combin.Theory Ser. B 44 (1988), 44-63.

[15] Yanpei A. Liu *Polyhedral Theory on Graphs*, Acta Mathematica Sinica, New Series, 1994, Vol.10,No.2, pp.136-142.

[16] G.Ringel. *Map Color Theorem*, Springer Verlag, 1974.

[17] P. Rosenstiel, R.C. Read. *On the principal edge tripartition of a graph*, Discrete Math. 3,195-226, 1978.

[18] S. Stahl. *The Embedding of a Graph - A Survey.* J.Graph Th., Vol 2 (1978), 275-298.

[19] S. Stahl. *Permutation-partition pairs: A combinatorial generalisation of graph embedding*, Trans Amer. Math. Soc. 1 (259) (1980), 129-145.

[20] S. Stahl. *A combinatorial analog of the Jordan curve theorem*, J.Combin.Theory Ser.B 35 (1983), 28-38.

[21] S. Stahl. *A duality for permutations*, Discrete Math. 71 (1988), 257-271.

[22] S. Stahl. *The Embedding of a Graph — A Survey*, Journ.Graph Theory, Vol.2, 275-298, 1978.

[23] W.T. Tutte. *What is a map? New directions in the theory of graphs.* Academic Press, NY, 1973, pp. 309-325.

[24] W.T.Tutte. *Combinatorial maps*, in *Graph theory*, chapter X, 1984.

[25] A. Vince. *Combinatorial maps*, J.Combin.Theory Ser.B 34 (1983), 1-21.

[26] T. R. S. Walsh, *Hypermaps Versus Bipartite Maps*, Journ. Comb. Math., Ser B 18, 155-163, 1975.

[27] A. T. White. *Graphs, Groups and Surfaces*, North-Holland, P.C., 1973.

[28] H. Wielandt. *Finite Permutations Groups*, Academic Press, New York, 1964.

[29] D. Zeps. *Graphs with rotation in permutation technique*, KAM Series, N 94-274, 1994, Prague, Charles University, 8pp.

[30] D. Zeps. *Graphs as rotations*, KAM Series, 96-327, Prague,1996, 9pp.

[31] D. Zeps. *Graphs with rotations:partial maps*, KAM Series, 97-365, 1996, 12pp.

[32] D. Zeps. *The use of the combinatorial map theory in graph-topological computations*, KAM Series, 97-364, Prague, 1997, 8pp.

[33] D.Zeps. *The theory of Combinatorial Maps and its Use in the Graph-topological computations*, PhD thesis, 1998.

[34] D. Zeps. *Using combinatorial maps in graph-topological computations* , KAM Series, 99-438, Prague, 1997, 11pp.

[35] D. Zeps. *Combinatorial Maps. Tutorial*, Online book, http://www.ltn.lv/ dainize/tutorial/Combinatorial Mps.Tutorial.htm, 2004.