



HAL
open science

A probabilistic study of neural complexity

Jerome Buzzi, Lorenzo Zambotti

► **To cite this version:**

| Jerome Buzzi, Lorenzo Zambotti. A probabilistic study of neural complexity. 2009. hal-00409143v2

HAL Id: hal-00409143

<https://hal.science/hal-00409143v2>

Preprint submitted on 13 Sep 2009 (v2), last revised 18 Dec 2009 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A PROBABILISTIC STUDY OF NEURAL COMPLEXITY

J. BUZZI AND L. ZAMBOTTI

ABSTRACT. G. Edelman, O. Sporns, and G. Tononi have introduced in theoretical biology the *neural complexity* of a family of random variables. They have defined it as a specific average of mutual information over subsystems. We show that their choice of weights satisfies two natural properties, namely exchangeability and additivity. This paper classifies all functionals satisfying these two properties (which we call *intricacies*) in terms of probability laws on the unit interval and studies the growth rate of maximal intricacies when the size of the system goes to infinity. For systems of a fixed size, we show that the maximizers are non-unique and that the maximal value is not approached by exchangeable laws.

1. INTRODUCTION

1.1. A measure of complexity from theoretical biology. Biology has to deal with "complex systems" in some obvious and not so obvious senses. To quantify this complexity and more specifically measure the balance between integration and differentiation which manifests itself in neural networks of the brain, G. Edelman, O. Sporns and G. Tononi [18] have introduced a quantity which they have called *neural complexity*. It is defined in terms of the mutual information between any subsystem and its complement (see below and the Appendix for the needed facts of information theory).

A series of works [1, 5, 7, 9, 10, 12, 13, 14, 15, 16, 19, 20] have used numerical experiments based on Gaussian approximations and simple examples to suggest that high values of this neural complexity are associated to non-trivial organization of the network, away from "chaos" (maximal entropy and independence of the neurons) and "order" (minimal entropy and complete determinacy).

The aim of this paper is to provide a mathematical foundation for this complexity for discrete systems. Starting with the example of the functional defined by Edelman-Sporns-Tononi we give a natural class of functionals satisfying two main properties: *exchangeability* and *weak-additivity*. The former property means that the functional is invariant under permutations of the system. The latter that it is additive over independent systems. We call these functional *intricacies* and give a unified probabilistic representation of these functionals.

They are *differences* of entropies, and, except in trivial cases, are neither convex nor concave. Hence their maximization is a non-trivial problem. We were not able

2000 *Mathematics Subject Classification.* 94A17, 92B30, 60C05.

Key words and phrases. Entropy, Mutual information, Complexity, Discrete Probability, Exchangeable vectors, Stationary processes.

to use the techniques of statistical mechanics and the methods which have been successful for frustrated systems (see [17]).

Nevertheless we are able to estimate the growth rate of the maximal intricacy of finite systems with size going to infinity, possibly subject to a further constraint on their entropy. We also show that laws maximizing intricacy are not exchangeable.

Finally we extend these considerations to stationary processes. In particular, in opposition to exchangeable systems (which have small intricacy), the mean intricacy of a stationary process can approximate the maximal value found for finite size.

The computation of the exact growth rate of the intricacy as a function of the size builds on the techniques of this paper, especially the probabilistic representation, but requires additional ideas, so is deferred to another paper [3].

1.2. Intricacy. We recall that the *entropy* of a random variable X taking values in a finite space E is defined by

$$H(X) := - \sum_{x \in E} P_X(x) \log(P_X(x)), \quad P_X(x) := \mathbb{P}(X = x).$$

Given two random variables defined over the same probability space, the *mutual information* between X and Y is

$$\text{MI}(X, Y) := H(X) + H(Y) - H(X, Y).$$

We refer to the appendix for a review of the main properties of the entropy and the mutual information and to [4] and [8] for introductions to information theory and the various roles of entropy in mathematical physics, respectively. For now, it suffices to recall that $\text{MI}(X, Y) \geq 0$ is equal to zero if and only if X and Y are independent, and therefore $\text{MI}(X, Y)$ is a measure of the dependence between X and Y .

Edelman, Sporns and Tononi consider systems formed by a finite family $X = (X_i)_{i \in I}$ and define the following concept of complexity. For any $S \subset I$, they divide the system in two families

$$X_S := (X_i, i \in S), \quad X_{S^c} := (X_i, i \in S^c),$$

where $S^c := I \setminus S$. Then they compute the mutual informations $\text{MI}(X_S, X_{S^c})$ and consider an average of these:

$$\mathcal{I}(X) := \frac{1}{|I| + 1} \sum_{S \subset I} \frac{1}{\binom{|I|}{|S|}} \text{MI}(X_S, X_{S^c}),$$

where $|I|$ denotes the cardinality of I . Note that $\mathcal{I}(X)$ is really a function of the *law* of X and not of its random values. We remark that the factor $(|I| + 1)$ in the denominator is not present in the original definition but is necessary for the additivity property to hold.

In this paper we prove that \mathcal{I} fits into a natural class of functionals, which we call **intricacies**. We shall see that these functionals have very similar, though not identical properties and admit a natural and technically very useful probabilistic representation by means of a probability measure on $[0, 1]$.

1.3. Main results. We state our main results in terms of an intricacy $\mathcal{I}^c(X)$, but, until the corresponding definition of Section 2, the reader is invited to assume that this is just the neural complexity $\mathcal{I}(X)$.

We need some notations. In this section, the system will be $X = (X_i)_{i=1,\dots,N}$ where each X_i takes value in a set with $d \geq 2$ elements (without loss of generality, $\{0, \dots, d-1\}$). $\mathcal{X}(d, N)$ is the set of such random variables and $\mathcal{M}(d, N)$ the set of their laws. If X is such a system with law μ , we denote its entropy, resp. intricacy, by $H(X) = H(\mu)$, resp. $\mathcal{I}^c(X) = \mathcal{I}^c(\mu)$. Of course, entropy and intricacy are in fact (deterministic) functions of the law μ of X and not of the (random) values of X .

Minimization of intricacies is a trivial problem. Indeed, both complete "order" (i.e., a deterministic family X) or total "chaos" (e.g., a uniform family) imply that every mutual information vanishes and therefore $\mathcal{I}^c(X) = 0$.

On the other hand, maximization of intricacies is a subtler issue, since large intricacy values require a compromise between randomness and mutual dependence. One wants X_S to be "as random as possible" and at the same time X_S and X_{S^c} to be as correlated as possible, for many subsets S . Moreover, intricacies can be expressed as a difference between entropies, see (2.5). Therefore intricacies, as differences of concave functionals, are not concave, a fact which explains why maximizing intricacies is difficult.

We shall obtain intricacies growing linearly in the size of the system by joining independent copies of a system with some positive mutual information. We shall see that this is in fact the maximal possible growth rate and that there is a well-defined maximal speed.

Theorem 1.1. *Let \mathcal{I}^c be some intricacy and let $d \geq 2$ be some integer.*

(1) *Denote by $\mathcal{I}^c(d, N)$ the supremum of $\mathcal{I}^c(X)$ over all $X \in \mathcal{X}(d, N)$. Then*

$$\mathcal{I}^c(d) := \lim_{n \rightarrow \infty} \frac{\mathcal{I}^c(d, N)}{N} \text{ exists in }]0, (\log d)/2]. \quad (1.1)$$

(2) *Let $(\delta_N)_{N \geq 1}$ be a sequence of non-negative numbers converging to zero and $x \in [0, 1]$. Then the following limit*

$$\mathcal{I}^c(d, x) := \lim_{N \rightarrow \infty} \frac{1}{N} \sup \left\{ \mathcal{I}^c(X) : X \in \mathcal{X}(d, N), \left| \frac{H(X)}{N \log d} - x \right| \leq \delta_N \right\}$$

exists and is independent of $(\delta_N)_{N \geq 1}$.

Remark 1.2. By considering a set of independent, identically distributed (i.i.d. for short) Bernoulli variables with the appropriate parameter, it is easy to see that for any $0 \leq h \leq N \log d$, there is $X \in \mathcal{X}(d, N)$ such that $H(X) = h$ and $\mathcal{I}^c(X) = 0$. Hence minimization of intricacies is a trivial problem also under fixed entropy.

Thus maximal intricacy grows linearly in the size of the system. What happens if we restrict to smaller classes of systems, enjoying particular symmetries? We consider two interesting classes: stationary sequences and (finite) exchangeable families.

We denote by $\text{EX}(d, N)$ the set of random variables $X \in \mathcal{X}(d, N)$ which are exchangeable, i.e., for all permutations σ of $\{1, \dots, N\}$, $X := (X_1, \dots, X_N)$ and

$X_\sigma := (X_{\sigma(1)}, \dots, X_{\sigma(N)})$ have the same law. We also consider stationary sequences $S \in \mathcal{S}(d, \mathbb{N}^*)$, namely sequences $(S_i)_{i \geq 1}$ such that $S_i \in \{0, \dots, d-1\}$ for all i and, for any $N \geq 1$, $(S_i)_{i \geq 1}$ and $(S_{i+N})_{i \geq 1}$ have the same law.

Theorem 1.3. *Let \mathcal{I}^c be an intricacy.*

(1) *Exchangeable systems have small intricacies. More precisely*

$$\sup_{X \in \text{EX}(d, N)} \mathcal{I}^c(X) = o(N^{2/3+\epsilon}), \quad N \rightarrow +\infty,$$

for any $\epsilon > 0$. In particular

$$\lim_{N \rightarrow \infty} \frac{1}{N} \max_{X \in \text{EX}(d, N)} \mathcal{I}^c(X) = 0.$$

(2) *For all $\varepsilon > 0$ there exists a stationary sequence $S \in \mathcal{S}(d, \mathbb{N}^*)$ such that*

$$\limsup_{N \rightarrow +\infty} \frac{\mathcal{I}^c(S_1, \dots, S_N)}{N} \geq \mathcal{I}^c(d) - \varepsilon,$$

where $\mathcal{I}^c(d)$ is defined in (1.1).

A consequence is that exact maximizers at fixed system size are not unique: for N sufficiently large, the maximal intricacy is not attained at an exchangeable law; therefore, by permuting a system with maximal intricacy we obtain different laws, all with the same maximal intricacy. However we do not know the number of exact maximizers nor if there are always finitely many of them.

1.4. Further questions. The properties of the maximizers remain mostly unknown at this point. The techniques involved in the computation of the maximal values $\mathcal{I}^c(d)$, $\mathcal{I}^c(d, x)$, in [3], yield information about their entropy and the size of their support. However it would also be interesting to have an explicit construction of maximizers and to estimate their number.

Applications in biology and/or works of previous authors suggest other directions of research. One would like to generalize this analysis to *continuous random variables* or systems with geometrical or combinatorial constraints (e.g., living on some *graphs* or invariant with respect to some groups). This would allow a comparison with the estimation of the neural complexity of some Gaussian laws satisfying such constraints as performed by several authors [1, 5, 14]. It would also be interesting to consider *infinite systems* since biological systems can be very large.

Numerical evaluation of the neural complexity is difficult. Indeed, entropies are notoriously hard to compute and neural complexity involves an exponential number of those. It is therefore important to estimate the *computational complexity* of this problem and design efficient, possibly probabilistic algorithms with good error bounds. We note that purely heuristic estimates have been used by previous authors [10, 18].

1.5. Organization of the paper. In Sec. 2, we formally define intricacies, giving some basic properties and examples. Sec. 3 relates the weak additivity of an intricacy with a property of its coefficients. As a by-product, we obtain a probabilistic representation of all intricacies. We check that neural complexity corresponds to the uniform law on $[0, 1]$. In Sec. 4 we prove Theorem 1.1 by showing the existence of the limits $\mathcal{I}^c(d)$, $\mathcal{I}^c(d, x)$. Finally in Sec. 5 we prove Theorem 1.3. An Appendix recalls some basic facts from information theory for the convenience of the reader and to fix notations.

2. INTRICACY

2.1. Definition. In this paper, a **system** is a finite collection $(X_i)_{i \in I}$ of random variables, each X_i , $i \in I$, taking value in the same finite set, say $\{0, \dots, d-1\}$ with $d \geq 2$ given. The set I is called the *support* of X . Most of the time I is a subset of the positive integers or simply $\{1, \dots, N\}$. In this case it is convenient to write N for I .

We let $\mathcal{M}(d, I)$ be the set of the corresponding laws, that is, the probability measures on $\{0, \dots, d-1\}^I$ for any finite subset. We often identify it with $\mathcal{M}(d, N) := \mathcal{M}(d, \{1, \dots, N\})$ for $N = |I|$.

For $S \subset I$, we use the abbreviation:

$$X_S := (X_i, i \in S).$$

Intricacies will be functionals over such systems (more precisely: over their laws) formalizing (and generalizing) the neural complexity of Edelman-Sporns-Tononi [18] as follows.

Definition 2.1. *A system of coefficients over some set of indices \mathbb{I} (typically $\mathbb{I} = \mathbb{N}^*$) is a collection of numbers*

$$c := (c_S^I : I \subset \subset \mathbb{I}, S \subset I)$$

(i.e., I ranges over the finite subsets of \mathbb{I}) satisfying, for all I and all $S \subset I$:

$$c_S^I \geq 0, \quad \sum_{S \subset I} c_S^I = 1, \quad \text{and } c_{S^c}^I = c_S^I \quad (2.1)$$

where $S^c := I \setminus S$. We denote the set of such systems by $\mathcal{C}(\mathbb{I})$.

The corresponding **mutual information functional** is $\mathcal{I}^c : \mathcal{M} \rightarrow \mathbb{R}$ defined by:

$$\mathcal{I}^c(X) := \sum_{S \subset I} c_S^I \text{MI}(X_S, X_{S^c}).$$

By convention, $\text{MI}(X_\emptyset, X_I) = \text{MI}(X_I, X_\emptyset) = 0$. An **intricacy**, is a mutual information function satisfying:

- (1) **exchangeability** (invariance by permutations): if $I, J \subset \subset \mathbb{I}$ and $\phi : I \rightarrow J$ is a bijection, then $\mathcal{I}^c(X) = \mathcal{I}^c(Y)$ for any $X := (X_i)_{i \in I}$, $Y := (X_{\phi^{-1}(j)})_{j \in J}$;
- (2) **weak additivity**: for any two independent sub-systems $(X_i)_{i \in I}$, $(Y_j)_{j \in J}$ (defined on the same probability space): $\mathcal{I}^c(X, Y) = \mathcal{I}^c(X) + \mathcal{I}^c(Y)$.

Let $\mathcal{I}(\mathbb{I}) \subset \mathcal{C}(\mathbb{I})$ denote the systems of coefficients defining intricacies.

\mathcal{I}^c is **non-null** if some coefficient c_S^I with $S \notin \{\emptyset, I\}$ is not zero.

Remarks 2.2. For all intricacies, and more generally exchangeable mutual information functionals, $c \in \mathcal{C}(\mathbb{I})$ readily extends to any set with cardinality at most that of \mathbb{I} .

As $\text{MI}(X_S, X_{S^c}) = \text{MI}(X_{S^c}, X_S)$, the above symmetric condition $c_{S^c}^I = c_S^I$ can always be satisfied by replacing c_S^I with $\frac{1}{2}(c_S^I + c_{S^c}^I)$ without changing the functional. Also $\sum_{S \subset I} c_S^I = 1$ is simply an irrelevant normalization when studying systems with a given index set I .

It turns out that a non-null intricacy automatically satisfies an *a priori* stronger property: it has no zero coefficient, see Lemma 3.9.

Lemma 2.3. *A system of coefficients $c \in \mathcal{C}(\mathbb{I})$ is uniquely determined by the corresponding mutual information functional $\mathcal{I}^c(X)$, $X \in \mathcal{X}(d, I)$ for $I \subset \subset \mathbb{I}$ for any given $d \geq 2$. Moreover, the following equivalences hold:*

- $\mathcal{I}^c(X)$ is exchangeable if and only if c_S^I depends only on $|I|$ and $|S|$;
- $\mathcal{I}^c(X)$ is weakly additive if and only if:

$$\forall I \subset \subset \mathbb{I}, \forall J \subset \subset \mathbb{I} \setminus I, \forall S \subset I, \quad c_S^I = \sum_{T \subset J} c_{S \sqcup T}^{I \sqcup J}. \quad (2.2)$$

We say that a system c of coefficients satisfying (2.2) above is **projective**. We defer the proof of Lemma 2.3 to the end of Sec. 3.1. Proposition 3.6 below describes all intricacies.

The following mutual information functionals will be proved to be intricacies in section 3.

Definition 2.4. *The intricacy \mathcal{I} of Edelman-Sporns-Tononi is defined by its coefficients:*

$$c_S^I = \frac{1}{|I| + 1} \frac{1}{\binom{|I|}{|S|}}. \quad (2.3)$$

For $0 < p < 1$, the p -symmetric intricacy $\mathcal{I}^p(X)$ is:

$$c_S^I = \frac{1}{2} (p^{|S|} (1-p)^{|I \setminus S|} + (1-p)^{|S|} p^{|I \setminus S|}).$$

For $p = 1/2$, this is the **uniform intricacy** $\mathcal{I}^U(X)$ with:

$$c_S^I = 2^{-|I|}.$$

It is not obvious that the three above mutual information functionals are intricacies, but this will follow easily from Lemma 3.8.

Remark 2.5. The coefficients of the Edelman-Sporns-Tononi intricacy \mathcal{I} ensure that subsystems of all sizes contribute significantly to the intricacy. This is in sharp contrast to the p -symmetric coefficients for which subsystems of size far from pN or $(1-p)N$ give a vanishing contribution when N gets large.

Remark 2.6. The global $1/(|I| + 1)$ factor in \mathcal{I} is not present in [18], which did not compare systems of different sizes. However it is required for weak additivity.

2.2. Basic Properties. We prove some general and easy properties of intricacies. Recall that $\mathcal{X}(d, N)$ is the set $\Lambda_{d,N}$ -valued random variables, where $\Lambda_{d,N} = \{0, \dots, d-1\}^N$.

Lemma 2.7. *Let \mathcal{I}^c be a mutual information functional. For each $d \geq 2$ and $N \geq 1$, $\mathcal{I}^c : \mathcal{M}(d, N) \rightarrow \mathbf{R}$ is continuous. In particular, the following supremum, introduced in Theorem 1.1, is achieved:*

$$\mathcal{I}^c(d, N) := \sup_{X \in \mathcal{X}(d, N)} \mathcal{I}^c(X) = \sup_{\mu \in \mathcal{M}(d, N)} \mathcal{I}^c(\mu), \quad (2.4)$$

If \mathcal{I}^c is non-null, then \mathcal{I}^c is neither convex nor concave.

Proof. Continuity is obvious and existence of the maximum follows from the compactness of the finite-dimensional simplex $\mathcal{M}(d, N)$.

To disprove convexity and concavity we use the following examples. Pick I with at least two elements, say 1 and 2. Observe that $K := c_{\{1\}}^I + c_{\{2\}}^I$ is positive by the non-degeneracy of \mathcal{I}^c . Fix $d \geq 2$.

First, for $i = 0, 1$, let μ_i over $\{0, \dots, d-1\}^I$ be defined by $\mu_i(i, i, 0, \dots, 0) = 1$. We have:

$$\mathcal{I}^c\left(\frac{\mu_0 + \mu_1}{2}\right) \geq K \cdot \log d > \frac{\mathcal{I}^c(\mu_0) + \mathcal{I}^c(\mu_1)}{2} = 0.$$

Second, let ν_0 be defined by $\nu_0(0, 0, 0, \dots, 0) = \nu_0(1, 1, 0, \dots, 0) = 1/2$ and ν_1 by $\nu_1(0, 1, 0, \dots, 0) = \nu_1(1, 0, 0, \dots, 0) = 1/2$. We have:

$$\mathcal{I}^c\left(\frac{\nu_0 + \nu_1}{2}\right) = 0 < K \cdot \log d \leq \frac{\mathcal{I}^c(\nu_0) + \mathcal{I}^c(\nu_1)}{2}.$$

□

The following expression of an intricacy as a non-convex combination of the entropy of subsystems is crucial to its understanding.

Lemma 2.8. *For any intricacy \mathcal{I}^c and $X \in \mathcal{X}(d, N)$*

$$\mathcal{I}^c(X) = \sum_S c_S^I \text{MI}(X_S, X_{S^c}) = 2 \left(\sum_S c_S^I \text{H}(X_S) \right) - \text{H}(X). \quad (2.5)$$

Proof. The result readily follows from: $\text{MI}(X, Y) = \text{H}(X) + \text{H}(Y) - \text{H}(X, Y)$, $c_S^I = c_{S^c}^I$, and $\sum_S c_S^I = 1$. □

We introduce the notation

$$\text{MI}(S) := \text{MI}(X_S, X_{I \setminus S})$$

which will be used only when the understood dependence on X and I is clear.

Lemma 2.9. *For any intricacy \mathcal{I}^c and $X \in \mathcal{X}(d, N)$*

$$0 \leq \mathcal{I}^c(X) \leq \frac{N}{2} \log d.$$

Finally, if \mathcal{I}^c is non-null, then $\mathcal{I}^c(X) = 0$ if and only if $X = (X_1, \dots, X_N)$ is an independent family.

Proof. The inequalities follow from basic properties of the mutual information (see the Appendix):

$$0 \leq \text{MI}(S) \leq \min\{\text{H}(X_S), \text{H}(X_{S^c})\} \leq \min\{|S|, N - |S|\} \log d \leq \frac{N}{2} \log d.$$

To prove the final remark, it is enough to show that: $\mathcal{I}^c(X) = 0 \iff \text{H}(X) = \sum_{i \in I} \text{H}(X_i)$. But observe that $\text{MI}(S) = 0$ iff $\text{H}(X) = \text{H}(X_S) + \text{H}(X_{S^c})$. An easy induction yields the claim. \square

2.3. Simple examples. We give some examples of finite systems and their intricacies both for illustrative purposes and use in some proofs below.

Let X_i take values in $\{0, \dots, d-1\}$ for all $i \in I$, a finite subset of \mathbb{N}^* . The first two examples show that total order and total disorder make the intricacy vanish.

Example 2.10. If the variables X_i are independent then each mutual information is zero and therefore: $\mathcal{I}^c(X) = 0$. \square

Example 2.11. If each X_i is a.s. equal to a constant c_i in $\{0, \dots, d-1\}$, then, for any $S \neq \emptyset$, $\text{H}(X_S) = 0$. Hence, $\mathcal{I}^c(X) = 0$. \square

A global constraint between random variables gives a non zero intricacy:

Example 2.12. If X_1 is uniform on $\{0, \dots, d-1\}$ and $X_i = X_1$ for all $i \in I$, then, for any $S \neq \emptyset$, $\text{H}(X_S) = \log d$ and, if additionally $S^c \neq \emptyset$, $\text{H}(X_S | X_{S^c}) = 0$ so that each mutual information $\text{MI}(X_S, X_{S^c})$ is $\log d$. Hence,

$$\mathcal{I}^c(X) = \sum_{S \subset I \setminus \{\emptyset, I\}} c_S^I \cdot \log d = (1 - c_\emptyset^I - c_I^I) \log d \leq \log d. \quad \square$$

The intricacy over $\mathcal{X}(d, N)$ can indeed reach the order N of Lemma 2.9:

Example 2.13. Let $d \geq 2$ and $N \geq 1$ be two integers. We are going to define a system $(X_i)_{i \in I}$, $I = \{1, \dots, N\}$, over alphabet $\{0, \dots, d^2-1\}$ for which $\mathcal{I}^U(X)/N \rightarrow \log d^2/4$; see the computation in Example 3.10 for an arbitrary intricacy: it yields the same result, except for the factor $1/4$.

Introduce Y_1, \dots, Y_N , i.i.d. random variables, each uniform over $\{0, \dots, d-1\}$ and define $X_i = Y_i + d(Y_{i+1} - 1)$ for $i = 1, \dots, N$. For $S \subset I$, set $\Delta := \{k = 1, \dots, N-1 : 1_S(k) \neq 1_S(k+1)\}$. Observe that $\text{MI}(S) = |\Delta| \log d$ and that we have a bijection:

$$S \in \{0, 1\}^{\{1, \dots, N\}} \mapsto (\mathbb{1}_S(1), \Delta) \in \{0, 1\} \times \{0, 1\}^{\{1, \dots, N-1\}}.$$

Hence:

$$I^U(X) = 2^{-N} \sum_{S \subset I} |\Delta| \log d = 2^{-N} \times 2 \sum_{\Delta \subset \{1, \dots, N-1\}} |\Delta| \log d$$

so:

$$I^U(X) = 2^{-N+1} \sum_{k=0}^{N-1} \binom{N-1}{k} k \log d = 2^{-N+1} \times (N-1) 2^{N-2} \log d = \frac{N-1}{4} \log d^2.$$

The following example will be useful to show that an intricacy \mathcal{I}^c determines its coefficients $c \in \mathcal{C}(\mathbb{N}^*)$ (Lemma 3.3).

Example 2.14. We consider a system of uniform variables, with a subset of equal ones and the remainder independent. More precisely, let $I \subset \mathbb{N}^*$, $K \subset I$ and fix $i_0 \in K$. $(X_i)_{i \in I} \in \mathcal{X}(d, I)$ is the system satisfying:

- (i) the family $X_{K^c \cup \{i_0\}}$ is uniform on $\{0, \dots, d-1\}^{K^c \cup \{i_0\}}$;
- (ii) $X_i = X_{i_0}$ for all $i \in K$.

It follows that $\text{MI}(S) = 0$ unless S and S^c both intersect K and then $\text{MI}(S) = \log d$. Thus

$$\mathcal{I}^c(X) = \log d \sum_{\substack{S \subset I \\ \emptyset \neq S \cap K \subsetneq K}} c_S^I \leq \log d, \quad \text{H}(X) = (|K^c| + 1) \log d.$$

3. WEAK ADDITIVITY, PROJECTIVITY AND REPRESENTATION

In this section we study the additivity of mutual information functionals, characterize it in terms of the coefficients. We establish a probabilistic representation of all intricacies and check that the neural complexity is indeed an intricacy, that defined by the uniform measure on the interval. We conclude this section by some useful consequences of this representation.

Throughout this section, $X = (X_i)_{i \in I}$ and $Y = (Y_i)_{i \in J}$, will be two systems defined on the same probability space and we shall consider the joint family $(X, Y) = \{X_i, Y_j : i \in I, j \in J\}$. (X, Y) is again a system and its index set is the disjoint union $I \sqcup J$ of I and J .

3.1. Projectivity and Additivity. We show that weak additivity can be read off the coefficients and that non-null intricacies are neither sub-additive nor super-additive.

Proposition 3.1. *Let \mathcal{I}^c be a mutual information functional. Then*

- (1) \mathcal{I}^c is weakly additive if and only if the coefficients are **projective**, i.e., satisfy (2.2).
- (2) Let \mathcal{I}^c be an intricacy. Then, for non-necessarily independent systems X, Y , we have: $\mathcal{I}^c(X, Y) \geq \max\{\mathcal{I}^c(X), \mathcal{I}^c(Y)\}$ and the approximate additivity:

$$|\mathcal{I}^c(X) + \mathcal{I}^c(Y) - \mathcal{I}^c(X, Y)| \leq \text{MI}(X, Y);$$

- (3) \mathcal{I}^c can fail to be super-additive or sub-additive.

Remark 3.2. *A direct, if slightly tedious computation, can show that $\mathcal{I}(X)$ and all $\mathcal{I}^p(X)$ satisfy weak-additivity. However, it will more easily follow from the representation theorem in the next section.*

To prove the equivalence in (1) we shall need the following fact:

Lemma 3.3. *Let $d \geq 2$ and I be an arbitrary support. The data $\mathcal{I}^c(X)$ for $X \in \mathcal{X}(d, J)$ for all $J \subset \subset I$ determines $c \in \mathcal{C}(I)$.*

Proof. The map is onto by definition. Let $c \in \mathcal{C}(I)$ so that \mathcal{I}^c is some mutual information functional. Using $c_{S^c}^I = c_S^I$, we restrict ourselves to coefficients with $|S| \leq |S^c|$, i.e., $|S| \leq |I|/2$.

Let us first consider a system $(X_i)_{i \in I} \in \mathcal{X}(d, I)$ where all variables are equal: $X_i = X_j$ for all $i, j \in I$ and X_i is uniform on $\{0, \dots, d-1\}$. Then $\text{MI}(S) := \text{MI}(X_S, X_{S^c}) = 0$ for $S = \emptyset$ or $S = I$, otherwise $\text{MI}(S) = \log d$. Hence, using the normalization $1 = \sum_S c_S^I$:

$$1 - \frac{\mathcal{I}^c(X)}{\log d} = \sum_S c_S^I - \sum_{\emptyset \subsetneq S \subsetneq I} c_S^I = c_\emptyset^I + c_I^I.$$

In particular, $c_\emptyset^I = c_I^I = (1 - \mathcal{I}^c(X)/\log d)/2$.

For each $K \subset I$, let X^K be the system as in Example 2.14. Fix $i_0 \in K$. Recall that $\text{MI}(S) := \text{MI}(X_S, X_{S^c})$ is 0 if $S \supset K$ or $S^c \supset K$, and is $\log d$ otherwise. Assume by induction that, for $1 \leq s \leq |I|/2$, c_S^I is determined for $|S| < s$. Picking $K \subset I$ with $|K| = |I| - s \geq |I|/2 \geq |K^c| = s$, we get:

- if $|S| < s$, we say nothing of $\text{MI}(S)$ but will use the inductive assumption;
- if $S = K$ or $S = K^c$, then $\text{MI}(S) = 0$;
- if $s \leq |S| \leq |K|$, $S \supset K$ implies $S = K$, $S \subset K^c$ implies $S = K^c$ since $s = |K^c|$. In all other cases: $\text{MI}(S) = \log d$.

Therefore,

$$\begin{aligned} \frac{\mathcal{I}^c(X^K)}{\log d} &= 2 \sum_{S \subset I} c_S^I \frac{\text{MI}(S)}{\log d} - \frac{\text{H}(X)}{\log d} \\ &= 4 \sum_{|S| < |I|/2} c_S^I \frac{\text{MI}(S)}{\log d} + 2 \sum_{|S| = |I|/2} c_S^I \frac{\text{MI}(S)}{\log d} - \frac{\text{H}(X)}{\log d} \\ &= 4 \sum_{|S| < s} c_S^I \frac{\text{MI}(S)}{\log d} + 4 \sum_{s \leq |S| < |I|/2} c_S^I + 2 \sum_{|S| = |I|/2} c_S^I - 2(c_K^I + c_{K^c}^I) - \frac{\text{H}(X)}{\log d} \end{aligned}$$

(the sum over $|S| = |I|/2$ is non-zero only if $|I|$ is even). Using $\sum_S c_S^I = 1$ and $c_S^I = c_{S^c}^I$, we get:

$$\frac{\mathcal{I}^c(X^K)}{\log d} - 2 \frac{\text{H}(X)}{\log d} = 2 \sum_{S \subset I} c_S^I \left(\frac{\text{MI}(S)}{\log d} - 1 \right) = 4 \sum_{|S| < s} c_S^I \left(\frac{\text{MI}(S)}{\log d} - 1 \right) - 4c_{K^c}^I.$$

It follows that $c_K^I = c_{K^c}^I$ is determined for any K with $|K| = s$. This completes the induction step and the proof of the lemma. \square

Proof of Proposition 3.1. We first check that weak additivity implies projectivity. For any $X \in \mathcal{X}(d, I)$ with $I \subset \subset \mathbb{N}^*$ and $J \subset \subset \mathbb{N}^* \setminus I$, we have:

$$\mathcal{I}^c(X) = \mathcal{I}^c(X, Z) = \sum_{S \subset I} \sum_{T \subset J} c_{S \cup T}^{I \cup J} \text{MI}(X_S, X_{S^c})$$

for $Z = (Z_j)_{j \in J}$ with each Z_j a.s. constant and therefore independent of X . Lemma 3.3 then implies that (2.2) holds. Moreover, (A.6) yields the monotonicity claim of point (2).

For the approximate additivity, we consider (A.7) for any $S \subset I$, $T \subset J$:

$$\text{MI}(X_S, Y_T, X_{S^c}, Y_{T^c}) = \text{MI}(X_S, X_{S^c}) + \text{MI}(Y_T, Y_{T^c}) \pm \text{MI}(X, Y)$$

where $\pm \text{MI}(X, Y)$ denotes a term belonging to $[-\text{MI}(X, Y), \text{MI}(X, Y)]$. The projectivity now gives:

$$\begin{aligned} \mathcal{I}^c(X, Y) &= \sum_{S \subseteq I, T \subseteq J} c_{S \sqcup T}^{I \sqcup J} \text{MI}(S \sqcup T) \\ &= \sum_{S \subseteq I, T \subseteq J} c_{S \sqcup T}^{I \sqcup J} (\text{MI}(X_S, X_{S^c}) + \text{MI}(Y_T, Y_{T^c}) \pm \text{MI}(X, Y)) \\ &= \mathcal{I}^c(X) + \mathcal{I}^c(Y) \pm \text{MI}(X, Y). \end{aligned}$$

This is the approximate additivity of point (2). If X and Y are independent, then $\text{MI}(X, Y) = 0$, proving the weak additivity.

We finally give the counter-examples. For sub-additivity, it is enough to assume the intricacy to be non-null and to consider $X = Y$ a single random variable uniform on $\{1, 2\}$ and compute:

$$\mathcal{I}^c(X) = \mathcal{I}^c(Y) = 0 \text{ whereas } \mathcal{I}^c(X, Y) = 2c_1^2 \log 2 > 0.$$

For super-additivity, we assume $c_\emptyset^I + c_I^I < \frac{1}{2} + \frac{c_\emptyset^{I \sqcup I} + c_I^{I \sqcup I}}{2}$ and take $X = Y$ a collection of $N = |I|$ copies of the same variable taking two values with probability $(\frac{1}{2}, \frac{1}{2})$. Then $\text{MI}(S) = \log 2$ except if $S \in \{\emptyset, I\}$, in which case $\text{MI}(S) = 0$. By example 2.12

$$\frac{\mathcal{I}^c(X, Y)}{\log 2} = 1 - c_\emptyset^{I \sqcup I} - c_I^{I \sqcup I} < 2(1 - c_\emptyset^I - c_I^I) = \frac{\mathcal{I}^c(X) + \mathcal{I}^c(Y)}{\log 2}.$$

□

Proof of Lemma 2.3. The Lemma follows easily: we have just seen the equivalence of projectivity and weak additivity whereas the characterization of exchangeability is a direct consequence of Lemma 3.3. □

3.2. Representation of intricacies. In this section, we give a probabilistic representation for all intricacies. This will provide us with a way to estimate the maximal value of intricacy for large systems in [3]. For notational convenience, we consider intricacies over the positive integers \mathbb{N}^* .

We say that a random variable W over $[0, 1]$ is **symmetric** if W and $1 - W$ have the same law. A measure on $[0, 1]$ is symmetric if it is the law of a symmetric random variable.

Proposition 3.4. *Let \mathcal{I}^c be a mutual information functional defined by some system of coefficients $c \in \mathcal{C}(\mathbb{N}^*)$ over some infinite index set, which we assume to be \mathbb{N}^* for notational convenience.*

- (1) \mathcal{I}^c is an intricacy, i.e., it is exchangeable and weakly additive, if and only if there exists a symmetric random variable W_c over $[0, 1]$ such that:

$$\forall I \subset \subset \mathbb{N}^*, \forall S \subset I, \quad c_S^I = \mathbb{P}(\mathcal{Z} \cap I = S) \tag{3.1}$$

where \mathcal{Z} is the random subset of \mathbb{N}^*

$$\mathcal{Z} := \{i \in \mathbb{N}^* : Y_i \geq W_c\}. \tag{3.2}$$

with $(Y_i)_{i \geq 1}$ an i.i.d. sequence of uniform random variables on $[0, 1]$, independent of W_c .

(2) If \mathcal{I}^c is an intricacy, then the law λ_c of W_c is uniquely determined by \mathcal{I}^c and for all $X \in \mathcal{X}(d, I)$ independent of \mathcal{Z}

$$\mathcal{I}^c(X) = \mathbb{E}(\text{MI}(\mathcal{Z} \cap I)), \quad \text{MI}(S) := \text{MI}(X_S, X_{I \setminus S}).$$

Remark 3.5. The definition (3.2) of the random set \mathcal{Z} is a particular case of the so-called *Kingman paintbox* construction, see [2]. In this setting, it yields a random exchangeable partition of \mathbb{N}^* into a subset \mathcal{Z} and its complement, each with asymptotic density a.s. equal to W_c , respectively $1 - W_c$.

After the proof of the proposition we give the measures μ, μ^U, μ^p representing respectively $\mathcal{I}, \mathcal{I}^U$ and \mathcal{I}^p .

We start with the following Lemma

Lemma 3.6. *Let $\mathcal{C}(\mathbb{N}^*)$ be the set of systems of coefficients of general (i.e. exchangeable and weakly additive) intricacies. Let $\mathcal{PS}([0, 1])$ be the set of symmetric probability measures λ on $[0, 1]$. Then, the map $\lambda \mapsto c$ defined from $\mathcal{PS}([0, 1])$ to $\mathcal{C}(\mathbb{N}^*)$ according to:*

$$c_k^n = \int_{[0,1]} x^k (1-x)^{n-k} \lambda(dx), \quad \forall 0 \leq k \leq n, \quad (3.3)$$

is a bijection.

Proof of Lemma 3.6. We first show that for an exchangeable projective \mathcal{I}^c , there exists a probability measure λ on $[0, 1]$ such that

$$c_n^{n+k} = \int_{[0,1]} x^n (1-x)^k \lambda(dx), \quad n \geq 1, k \geq 0 \quad (3.4)$$

i.e., the main claim of the Lemma, up to a convenient renumbering. We need the following classical moment result (see [6, VII.3]).

Lemma 3.7. *Let $(a_n)_{n \geq 1}$ be a sequence of numbers in $[0, 1]$. We define $(Da)_n := a_n - a_{n+1}$, $n \geq 1$. There exists a probability measure λ on $[0, 1]$ such that $a_n = \int x^n \lambda(dx)$ if and only if*

$$(D^k a)_n \geq 0, \quad \forall n \geq 1, \forall k \geq 1.$$

Remark that, setting $N = |I|$ and $M = |J|$, projectivity is equivalent to

$$c_k^N = \sum_{\ell=0}^M c_{k+\ell}^{M+N} \binom{M}{\ell}, \quad \forall 0 \leq k \leq N. \quad (3.5)$$

For $M = 1$ we obtain

$$c_k^{N+1} + c_{k+1}^{N+1} = c_k^N, \quad \forall 1 \leq k \leq N. \quad (3.6)$$

Let us set $m_n := c_n^n$. One proves easily by (3.6) and recurrence on k that

$$(D^k m)_n = c_n^{n+k} \in [0, 1], \quad \forall k, n \geq 1.$$

Therefore $(m_n)_{n \geq 1}$ defines a unique measure λ satisfying (3.4) for $k = 0$. The general case follows by induction from:

$$c_n^{n+k+1} = c_n^{n+k} - c_{n+1}^{n+k+1} = \int_0^1 x^n(1-x)^k - x^{n+1}(1-x)^k d\lambda = \int_0^1 x^n(1-x)^{k+1} d\lambda.$$

Thus λ is the unique solution to the claim of the Lemma. This uniqueness together with $c_k^n = c_{n-k}^n$, implies that λ is symmetric. Thus any intricacy defines a measure as claimed.

We turn to the converse, considering a symmetric measure λ on $[0, 1]$ and defining c by means of (3.3). The coefficients depending only on the cardinalities, \mathcal{I}^c is trivially exchangeable. The symmetry of λ yields immediately $c_k^n = c_{n-k}^n$, and the normalization condition is given by

$$\sum_{k=0}^N \binom{N}{k} c_k^N = \int_{[0,1]} \sum_{k=0}^N \binom{N}{k} x^k(1-x)^{n-k} \lambda(dx) = 1,$$

i.e. $c \in \mathcal{C}(\mathbb{N}^*)$. To prove the projectivity of c , namely (3.5), we compute:

$$\begin{aligned} \sum_{\ell=0}^M c_{k+\ell}^{M+N} \binom{M}{\ell} &= \int_0^1 \left[\sum_{\ell=0}^M \binom{M}{\ell} x^\ell(1-x)^{M-\ell} \right] x^k(1-x)^{N-k} \lambda(dx) \\ &= \int_0^1 x^k(1-x)^{N-k} \lambda(dx) = c_k^N. \end{aligned}$$

Thus (3.5) and projectivity follow. The Lemma is proved. \square

Proof of Proposition 3.4. First, let \mathcal{I}^c be an intricacy. Lemma 3.4 yields a symmetric probability measure λ_c on $[0, 1]$. Let W_c be a random variable with law λ_c and let Y_1, Y_2, \dots be defined as in the statement.

Each $i \in \mathbb{N}^*$ belongs to the random set \mathcal{Z} if and only if $Y_i \geq W_c$. The probability of this event is therefore $1 - W_c$ conditionally to W_c . As the variables Y_1, Y_2, \dots are independent:

$$\mathbb{P}(\mathcal{Z} \cap I = S \mid W_c) = (1 - W_c)^{|I \setminus S|} W_c^{|S|}.$$

Averaging over the values of W_c we obtain

$$\mathbb{P}(\mathcal{Z} \cap I = S) = \mathbb{E}((1 - W_c)^{|I \setminus S|} W_c^{|S|}) = \int_{[0,1]} x^{|S|}(1-x)^{|I \setminus S|} \lambda_c(dx) = c_S^I.$$

by (3.3) and this yields (3.1).

Conversely, suppose that $c = (c_S^I)_{S \subset I}$ has the form (3.1) for some probability \mathbb{P} defined by W_c, Y_1, Y_2, \dots as in the statement. Obviously $c_S^I \geq 0$ and $\sum_{S \subset I} c_S^I = 1$. $c_S^I = c_{S^c}^I$ follows from the symmetry of W_c . Thus c is a system of coefficients. Exchangeability of c follows from exchangeability of the random variables $\mathbb{1}_{(Y_i \geq W_c)}$, $i \in I$. By (3.1) we know that

$$c_S^I = c_{|S|}^{|I|} = \mathbb{E}((1 - W_c)^{|I \setminus S|} W_c^{|S|}) = \int_{[0,1]} x^{|S|}(1-x)^{|I \setminus S|} \lambda_c(dx).$$

Therefore, by Lemma 3.6 the functional \mathcal{I}^c is an intricacy. The Proposition is proved. \square

3.3. Examples of intricacies. We show that the Edelman-Sporns-Tononi intricacy and the uniform and p -symmetric intricacies correspond to natural probability laws on $[0, 1]$. In particular, they are weakly additive and really intricacies:

Lemma 3.8. *In the setting of Lemma 3.6*

- (1) *If W_c is uniform on $[0, 1]$ then \mathcal{I}^c is the Edelman-Sporns-Tononi coefficients.*
- (2) *If W_c is uniform on $\{p, 1-p\}$ then \mathcal{I}^c is the p -symmetric intricacy \mathcal{I}^p ; in the case $p = 1/2$, $W_c = \frac{1}{2}$ a.s. yields the uniform intricacy \mathcal{I}^U .*

Proof. Let W_c be uniform on $[0, 1]$. Then

$$\begin{aligned} \mathbb{P}(\mathcal{Z} \cap I = \{1, \dots, k\}) &= \mathbb{P}(Z_1 = \dots = Z_k = 1, Z_{k+1} = \dots = Z_N = 0) \\ &= \int_0^1 x^k (1-x)^{N-k} dx =: a(k, N-k). \end{aligned}$$

We claim now that for all $k \geq 1$ and $j \geq 0$

$$a(k, j) = \frac{j!}{(k+1) \cdots (k+j+1)} = \frac{1}{(k+j+1) \binom{k+j}{k}},$$

i.e., the Edelman-Sporns-Tononi coefficient c_j^{k+j} .

Indeed, for $j = 0$ this reduces to $\int_0^1 x^k dx = 1/(k+1)$. To prove the general case, one fixes k and uses recurrence on j . Indeed, suppose we have the result for $j \geq 0$. Then

$$\begin{aligned} \int_0^1 x^k (1-x)^{j+1} dx &= \int_0^1 x^k (1-x)^j dx - \int_0^1 x^{k+1} (1-x)^j dx \\ &= \frac{1}{(k+j+1) \binom{k+j}{k}} - \frac{1}{(k+j+2) \binom{k+j+1}{k+1}} = \frac{1}{(k+j+2) \binom{k+j+1}{k}}. \end{aligned}$$

If W_c is uniform over $\{p, 1-p\}$ then

$$\int_0^1 x^k (1-x)^{N-k} \frac{1}{2} (\delta_p + \delta_{1-p})(dx) = \frac{1}{2} (p^k (1-p)^{N-k} + (1-p)^k p^{N-k}),$$

which is the coefficient c_k^N of \mathcal{I}^p . \square

3.4. Further properties. We deduce two useful facts from the above representation.

Lemma 3.9. *The following are equivalent for an intricacy \mathcal{I}^c with associated measure λ_c as in Lemma 3.6.*

- (1) \mathcal{I}^c is non-null, i.e. $c_k^N > 0$ for at least one choice of $N \geq 2$ and $1 \leq k < N$;
- (2) $c_k^N > 0$ for all $N \geq 2$ and $1 \leq k < N$;
- (3) $\lambda_c([0, 1]) > 0$.

Proof. We have:

$$c_j^n = \int_{[0,1]} x^j (1-x)^{n-j} \lambda_c(dx)$$

with $x^j(1-x)^{n-j}$ zero exactly at $x \in \{0, 1\}$ whenever $0 < j < n$ and strictly positive on $]0, 1[$. Thus (1) \implies (3) \implies (2) \implies (1). \square

Example 3.10. We generalize the computation in Example 2.13 from \mathcal{I}^U to a non-null intricacy \mathcal{I}^c . Following the analysis done there, we get by Proposition 3.4

$$\frac{\mathcal{I}^c(X)}{\log d} = \sum_{S \subset I} c_S^I |\Delta(S)| = \sum_{k=1}^{N-1} \mathbb{E}(\mathbb{1}_{\mathcal{Z}}(k) \neq \mathbb{1}_{\mathcal{Z}}(k+1)) = (N-1) \mathbb{E}(\mathbb{1}_{\mathcal{Z}}(1) \neq \mathbb{1}_{\mathcal{Z}}(2))$$

Recall the probabilistic representation by a random variable W_c with law λ_c on $[0, 1]$. Compute:

$$c_0 := \mathbb{E}(\mathbb{1}_{\mathcal{Z}}(1) \neq \mathbb{1}_{\mathcal{Z}}(2)) = \int_0^1 2x(1-x) \lambda_c(dx) \in]0, 1/2].$$

Then:

$$\begin{aligned} \mathcal{I}^c(X) &= \mathbb{E}(\text{MI}(\mathbb{Z} \cap I)) = \sum_{i=1}^{N-1} \mathbb{E}(\mathbb{1}_{\mathcal{Z}}(i) \neq \mathbb{1}_{\mathcal{Z}}(i+1)) \\ &= (N-1) \mathbb{E}(\mathbb{1}_{\mathcal{Z}}(1) \neq \mathbb{1}_{\mathcal{Z}}(2)) \log d = c_0(N-1) \log d. \end{aligned}$$

4. BOUNDS FOR MAXIMAL INTRICACIES

In this section we prove Theorem 1.1. \mathcal{I}^c is some non-null intricacy and we define using (3.3)

$$\kappa_c := 1 - (c_0^2 + c_2^2) = 2c_1^2 = 2 \int_{[0,1]} x(1-x) \lambda_c(dx) > 0. \quad (4.1)$$

We recall that $\mathcal{M}(d, N)$ denotes the set of probability measures over the finite set $\Lambda_{d,N} := \{0, \dots, d-1\}^N$. Then $\mathcal{M}(d, N)$ is a compact subset of an affine space of dimension d^N . Any $\mu \in \mathcal{M}(d, N)$ can be interpreted as the law of a random vector $X \in \mathcal{X}(d, N)$. An intricacy \mathcal{I}^c is in fact a function of the law μ of X , i.e., a functional $\mathcal{I}^c : \mathcal{M}(d, N) \rightarrow \mathbb{R}_+$. Recall that $\mathcal{I}^c(d, N)$ (2.4) denotes its maximum over $\mathcal{M}(d, N)$.

We are going to show the following

Proposition 4.1. *Consider some intricacy general \mathcal{I}^c . Let $d \geq 2$. For all $N \geq 2$,*

$$\frac{\kappa_c \cdot \log d}{2} \cdot \left(1 - \frac{1}{N}\right) \leq \frac{\mathcal{I}^c(d, N)}{N} \leq \frac{1}{2} \log d \quad (4.2)$$

where $\kappa_c > 0$ is defined in (4.1).

Proof. The upper bound for $\mathcal{I}^c(d, N)/N$ follows from Lemma 2.9. We show now that by the weak additivity property one can construct vectors $X \in \mathcal{X}(d, N)$ such that $\mathcal{I}^c(X)$ grows linearly in N . More precisely, we shall obtain:

$$\mathcal{I}^c(d, N) \geq \frac{\kappa_c \cdot \log d}{2} \cdot (N-1). \quad (4.3)$$

Let X_1 be a uniform variable on $\{0, \dots, d-1\}$ and $X_2 = X_1$. Then by the second example in section 2.3, we know that for $I = \{1, 2\}$:

$$\mathcal{I}^c(X) = (c_{\{1\}}^I - c_{\{2\}}^I) \log d = \kappa_c \cdot \log d,$$

which is positive by (4.1). In particular, (4.3) holds for $N = 2$.

Let now $(X_{2i+1})_{i \geq 0}$ an i.i.d. family of uniform variables on $\{0, \dots, d-1\}$, and set $X_{2(i+1)} := X_{2i+1}$ for all $i \geq 0$. Then, for $M \geq 1$, $(X_i)_{i=1, \dots, 2M}$ is the product of M independent copies of (X_1, X_2) and by weak additivity

$$\mathcal{I}^c(X_1, \dots, X_{2M}) = M \cdot \mathcal{I}^c(X_1, X_2) = \frac{\kappa_c \cdot \log d}{2} \cdot 2M.$$

Analogously, by Proposition 3.1

$$\mathcal{I}^c(X_1, \dots, X_{2M+1}) \geq \mathcal{I}^c(X_1, \dots, X_{2M}) = \frac{\kappa_c \cdot \log d}{2} \cdot 2M,$$

and setting $N = 2M$ or $N = 2M + 1$ we obtain (4.3). \square

4.1. Super-additivity of $N \mapsto \mathcal{I}^c(d, N)$. Let us first remark that the following suprema are achieved by continuity and compactness:

$$\mathcal{I}^c(d, N) = \sup_{\mu \in \mathcal{M}(d, N)} \mathcal{I}^c(\mu), \quad \mathcal{I}^c(d, N, x) = \sup_{\substack{\mu \in \mathcal{M}(d, N) \\ \mathbf{H}(\mu) = xN \log d}} \mathcal{I}^c(\mu) \quad (4.4)$$

For the second supremum, observe that $\mathbf{H}(\mathcal{M}(d, N))$ is an interval containing 0 and $N \log d$. Hence the second supremum is also taken over a non-empty compact subset.

The measures achieving the first supremum are called *maximal intricacy measures*. Since the functional \mathcal{I}^c is not concave, we do not expect uniqueness of maximizers.

Lemma 4.2. *For any intricacy \mathcal{I}^c and $d \geq 2$, the following limits exist. First,*

$$\mathcal{I}^c(d) := \lim_{N \rightarrow \infty} \frac{\mathcal{I}^c(d, N)}{N} = \sup_{N \geq 1} \frac{\mathcal{I}^c(d, N)}{N} \in]0, +\infty[\quad (4.5)$$

and, for each $x \in [0, 1]$,

$$\mathcal{I}^c(d, x) := \lim_{N \rightarrow \infty} \frac{\mathcal{I}^c(d, N, x)}{N} = \sup_{N \geq 1} \frac{\mathcal{I}^c(d, N, x)}{N} \in]0, +\infty[. \quad (4.6)$$

Proof. We prove (4.6), (4.5) being similar and simpler. Fix $x \in [0, 1]$.

For each $N \geq 1$, let $a_N := \mathcal{I}^c(d, N, x)$. We claim that this sequence is *super-additive*, i.e.,

$$a_{N+M} \geq a_N + a_M, \quad \forall N, M \geq 1.$$

Indeed, let X^N and X^M such that $\mathcal{I}^c(X^N) = \mathcal{I}^c(d, N, x)$, $\mathcal{I}^c(X^M) = \mathcal{I}^c(d, M, x)$ and $\mathbf{H}(X^N) = xN \log d$, $\mathbf{H}(X^M) = xM \log d$. Assume that X^N and X^M are independent (if necessary consider two independent copies). Now, $\mathcal{I}^c(X^N, X^M) = \mathcal{I}^c(X^N) + \mathcal{I}^c(X^M)$ and $\mathbf{H}(X^N, X^M) = \mathbf{H}(X^N) + \mathbf{H}(X^M) = x(N+M) \log d$. Thus,

$$\begin{aligned} a_N + a_M &= \mathcal{I}^c(d, N, x) + \mathcal{I}^c(d, M, x) = \mathcal{I}^c(X^N) + \mathcal{I}^c(X^M) \\ &= \mathcal{I}^c(X^N, X^M) \leq \mathcal{I}^c(d, N+M, x) = a_{N+M}. \end{aligned}$$

Moreover, by Proposition 4.1, we have $\sup_{N \geq 1} a_N/N \in [\kappa_c \log d/2, \log d/2]$. Therefore, the existence of the limit and its coincidence with the supremum follows by a standard argument. \square

4.2. Adjusting Entropy. To strengthen the previous result to obtain the second assertion of Theorem 1.1, we must adjust the entropy without significantly changing the intricacy.

Lemma 4.3. *Let $X^{(1)}, \dots, X^{(r)} \in \mathcal{X}(d, N)$. Let U be a random variable over $\{1, \dots, r\}$, independent of $\{X^{(1)}, \dots, X^{(r)}\}$. Let $Y := X^{(U)} \in \mathcal{X}(d, N)$, i.e., $Y = X^{(u)}$ whenever $U = u$. Then:*

$$0 \leq H(Y_S) - \sum_{u=1}^r \mathbb{P}(U = u) H(X_S^{(u)}) \leq \log r, \quad \forall S \subset \{1, \dots, N\}, \quad (4.7)$$

$$-\log r \leq \mathcal{I}^c(Y) - \sum_{u=1}^r \mathbb{P}(U = u) \mathcal{I}^c(X^{(u)}) \leq 2 \log r. \quad (4.8)$$

Proof. We first prove (4.7). By (A.2),

$$H(Y_S|U) \leq H(Y_S) \leq H(Y_S, U) = H(Y_S|U) + H(U).$$

Now $H(U) \leq \log r$. Also (4.7) now follows from:

$$H(Y_S|U) = \sum_{u=1}^r \mathbb{P}(U = u) H(Y_S|U = u) = \sum_{u=1}^r \mathbb{P}(U = u) H(X_S^{(u)}).$$

Then (4.8) follows immediately, using (2.5) and (4.7). \square

Lemma 4.4. *Let $0 < x < 1$ and $\epsilon > 0$ and \mathcal{I}^c be some non-null intricacy. Then there exists $\delta_0 > 0$ and $N_0 < \infty$ with the following property for all $0 < \delta < \delta_0$ and $N \geq N_0$. For any $X \in \mathcal{X}(d, N)$ such that $\left| \frac{H(X)}{N \log d} - x \right| \leq \delta$, there exists $Y \in \mathcal{X}(d, N)$ satisfying:*

$$H(Y) = xN \log d, \quad |\mathcal{I}^c(Y) - \mathcal{I}^c(X)| \leq \epsilon N \log d.$$

Proof. We fix $\delta_0 = \delta_0(\epsilon, x) > 0$ so small that:

$$\frac{\delta_0}{\min\{1 - x - \delta_0, x - \delta_0\}} < \epsilon/4$$

and $N_0 = N_0(\epsilon, x, \delta_0)$ so large that:

$$\frac{\log 2}{N_0 \min\{1 - x - \delta_0, x - \delta_0\} \log d} < \epsilon/4.$$

Let $N \geq N_0$ and $X \in \mathcal{X}(d, N)$ be such that $\left| \frac{H(X)}{N \log d} - x \right| \leq \delta \leq \delta_0$. There are two similar cases, depending on whether $H(X)$ is greater or less than $xN \log d$. We assume $h := H(X)/N \log d < x$ and shall explain at the end the necessary modifications for the other case.

Let $Z = (Z_i, i = 1, \dots, N)$ be i.i.d. random variables, uniform over $\{0, \dots, d-1\}$. We consider $Y^t \in \mathcal{X}(d, N)$ defined by

$$Y^t := X \mathbb{1}_{(U \leq 1-t)} + Z \mathbb{1}_{(U > 1-t)},$$

where U is a uniform random variable over $[0, 1]$ independent of X and Z . $\mathcal{I}^c(Y^0) = \mathcal{I}^c(X)$ and $\mathcal{I}^c(Y^1) = \mathcal{I}^c(Z) = 0$. Hence, by the continuity of the intricacy, we get that there is some $0 < t_0 < 1$ such that $\mathbb{H}(Y^{t_0}) = xN \log d$. Let us check that t_0 is small.

By (4.7)

$$0 \leq \mathbb{H}(Y^t) - (1-t)\mathbb{H}(X) - t\mathbb{H}(Z) = \mathbb{H}(Y^t) - (1-t)hN \log d - tN \log d \leq \log 2.$$

so that, for some $\alpha \in [0, 1]$,

$$0 < t_0 = \frac{x-h}{1-h} + \frac{\alpha \log 2}{N(1-h) \log d} \leq \frac{\delta}{1-x-\delta} + \frac{\log 2}{N(1-x-\delta) \log d} < \frac{\epsilon}{2},$$

since $N \geq N_0$ and $\delta \leq \delta_0$. Thus, by (4.8), setting $Y := Y^{t_0}$,

$$|\mathcal{I}^c(Y) - (1-t_0)\mathcal{I}^c(X) - t_0\mathcal{I}^c(Z)| = |\mathcal{I}^c(Y) - (1-t_0)\mathcal{I}^c(X)| \leq 2 \log 2,$$

and therefore by (4.2)

$$|\mathcal{I}^c(Y) - \mathcal{I}^c(X)| \leq t_0\mathcal{I}^c(X) + 2 \log 2 \leq \frac{\epsilon}{2} N \log d + 2 \log 2.$$

Dividing by $N \log d$ we obtain the desired estimate.

For the case $h > x$, we use instead a system Z with constant variables, so that $\mathbb{H}(Z) = 0 = \mathcal{I}^c(Z)$ and a similar argument gives the result. \square

4.3. Proof of Theorem 1.1. Assertion (1) is already established: see (4.5) and Proposition 5.2. It remains to complete the proof of the second assertion.

We first observe that (4.6) gives the existence of the limit $\mathcal{I}^c(d, x)$ if $\delta_n = 0$, for all $n \geq 1$. Consider now a general sequence of non-negative numbers δ_n converging to zero. Obviously, $\mathcal{I}^c(d, N, x, \delta_N) \geq \mathcal{I}^c(d, N, x, 0)$, so that

$$\liminf_{N \rightarrow \infty} \frac{1}{N} (\mathcal{I}^c(d, N, x, \delta_N) - \mathcal{I}^c(d, N, x, 0)) \geq 0.$$

Let us prove the reverse inequality for the lim sup. Let $\epsilon > 0$. Let $X^N \in \mathcal{X}(d, N)$ realize $\mathcal{I}^c(d, N, x, \delta_N)$. Let δ_0 and N_0 be as in Lemma 4.4. We may assume that $N \geq N_0$ and $\delta_N < \delta_0$. It follows that there is some $Y^N \in \mathcal{X}(d, N)$ with the entropy $Nx \log d$ such that $\mathcal{I}^c(Y^N) \geq \mathcal{I}^c(X^N) - \epsilon N$. Hence, $\mathcal{I}^c(d, N, x, 0) \geq \mathcal{I}^c(d, N, x, \delta_N) - \epsilon N$. We obtain

$$\limsup_{N \rightarrow \infty} \frac{1}{N} (\mathcal{I}^c(d, N, x, \delta_n) - \mathcal{I}^c(d, N, x, 0)) \leq \epsilon,$$

Assertion (2) follows by letting $\epsilon \rightarrow 0$.

5. STATIONARY AND EXCHANGEABLE SYSTEMS

In this section we prove Theorem 1.3, i.e., that stationarity does not constrain intricacy but exchangeability does.

5.1. Stationary sequences. A sequence $(S_n)_{n \geq 1}$ of random variables is stationary if it has the same law as $(S_{n+1})_{n \geq 1}$. We prove the following result

Proposition 5.1. *Let \mathcal{I}^c be an intricacy. For all $\varepsilon > 0$ there exists a stationary sequence $S \in \mathcal{S}(d, \mathbb{N}^*)$ such that the following limit exists and satisfies:*

$$\lim_{N \rightarrow +\infty} \frac{\mathcal{I}^c(S_1, \dots, S_N)}{N} \geq \mathcal{I}^c(d) - \varepsilon,$$

where $\mathcal{I}^c(d)$ is defined in (4.5).

We do not know whether there exists a stationary sequence such that the above limit is exactly $\mathcal{I}^c(d)$.

Proof. Let $\varepsilon > 0$ and $N_\varepsilon \in \mathbb{N}$ be such that

$$\frac{\mathcal{I}^c(d, N_\varepsilon)}{N_\varepsilon} > \mathcal{I}^c(d) - \varepsilon.$$

Then there exists $X \in \mathcal{X}(d, N_\varepsilon)$ such that $\frac{\mathcal{I}^c(X)}{N_\varepsilon} > \mathcal{I}^c(d) - \varepsilon$. Let us consider an i.i.d. sequence $(X^k)_{k \geq 0}$ such that each $X^k \in \mathcal{X}(d, N_\varepsilon)$ has the same distribution as X . Let M be a uniform variable on $\{0, \dots, N_\varepsilon - 1\}$, independent of $(X^k)_{k \geq 0}$, and let us define

$$Y_{kN_\varepsilon+i} := X_i^k, \quad i = 1, \dots, N_\varepsilon, \quad k \geq 0,$$

and finally

$$S_i := Y_{i+M}, \quad i \geq 1.$$

Let us check that the sequence $(S_i)_{i \geq 1}$ is stationary. Let $N \geq 1$. Let f be an arbitrary measurable and bounded function over $\{0, \dots, d-1\}^p$, for some $p \geq 1$.

Observe that the laws of the random variables $f(Y_{N+1+m}, \dots, Y_{N+p+m})$, $m = 0, \dots, N_\varepsilon$, are the same, up to a permutation, that those of $f(Y_{1+m}, \dots, Y_{p+m})$. Hence,

$$\begin{aligned} \mathbb{E}(f(S_{N+1}, \dots, S_{N+p})) &= \frac{1}{N_\varepsilon} \sum_{m=0}^{N_\varepsilon-1} \mathbb{E}(f(Y_{N+1+m}, \dots, Y_{N+p+m})) \\ &= \frac{1}{N_\varepsilon} \sum_{m=0}^{N_\varepsilon-1} \mathbb{E}(f(Y_m, \dots, Y_{p+m})) = \mathbb{E}(f(S_1, \dots, S_p)). \end{aligned}$$

Obviously, $\mathcal{I}^c(S_1, \dots, S_N) \leq \mathcal{I}^c(d, N)$ so that $\limsup_{N \rightarrow \infty} \mathcal{I}^c(S_1, \dots, S_N)/N \leq \mathcal{I}^c(d)$. We turn to the non-trivial inequality, the lower bound on the lim inf.

Let $n \geq 1$ be an integer and set $n = kN_\varepsilon + r$ with $0 \leq r < N_\varepsilon$. Using (4.8) and weak additivity, we compute:

$$\begin{aligned} \mathcal{I}^c(S_1, \dots, S_n) &\geq \mathcal{I}^c(S_{N_\varepsilon}, \dots, S_{kN_\varepsilon}) \geq \frac{1}{N_\varepsilon} \sum_{m=1}^{N_\varepsilon} \mathcal{I}^c(Y_{N_\varepsilon+m}, \dots, Y_{kN_\varepsilon+m}) - \log N_\varepsilon \\ &= (k-1)\mathcal{I}^c(X) - \log N_\varepsilon > (k-1)N_\varepsilon(\mathcal{I}^c(d) - \varepsilon) - \log N_\varepsilon. \end{aligned}$$

Dividing by $n \rightarrow \infty$, we obtain:

$$\liminf_{n \rightarrow +\infty} \frac{\mathcal{I}^c(S_1, \dots, S_n)}{n} \geq \mathcal{I}^c(d) - 2\varepsilon.$$

□

5.2. Finite exchangeable families. We prove that exchangeable systems have small intricacy. In particular, one cannot approach the maximal intricacy with such systems.

Proposition 5.2. *Let \mathcal{I}^c be any mutual information functional and $d \geq 2$. Then for all $\varepsilon > 0$ there exists a constant $C = C(\varepsilon, d)$ such that for all exchangeable $X \in \mathcal{X}(d, N)$*

$$\mathcal{I}^c(X) \leq CN^{\frac{2}{3}+\varepsilon}, \quad N \geq 2. \quad (5.1)$$

In particular

$$\lim_{N \rightarrow \infty} \frac{1}{N} \max_{X \in \text{EX}(d, N)} \mathcal{I}^c(X) = 0.$$

Proof. Fix $\varepsilon > 0$. Throughout the proof, we denote by C constants which only depend on d and ε and which may change value from line to line. Also $\mathbf{k} = (k_1, \dots, k_d) \in \mathbb{N}^d$, $\mathbf{x} := \frac{1}{n}\mathbf{k}$ and $|\mathbf{k}| := k_1 + \dots + k_d = n$ and the multinomial coefficients and the entropy function are denoted by:

$$\binom{n}{\mathbf{k}} = \frac{n!}{k_1!k_2!\dots k_d!}, \quad h(\mathbf{x}) = -\sum_{i=1}^d x_i \log x_i.$$

We are going to use the following version of Stirling's formula

$$n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\zeta_n}, \quad \frac{1}{12n+1} < \zeta_n < \frac{1}{12n}, \quad n \geq 1.$$

Therefore, for all $\mathbf{k} \in \mathbb{N}^d$ such that $|\mathbf{k}| = n$

$$\binom{n}{\mathbf{k}} = \left[e^{nh(\mathbf{x})} (2\pi n)^{1/2} \prod_{x_i \neq 0} (2\pi n x_i)^{-1/2} \right] g(\mathbf{k}, n),$$

where $g(\mathbf{k}, n) := \exp(\zeta_n - \zeta_{k_1} - \dots - \zeta_{k_d})$ and therefore

$$\exp(-d) \leq g(\mathbf{k}, n) \leq \exp(1).$$

In particular, as all non-zero x_i satisfy $x_i \geq 1/n$,

$$\left| \frac{1}{n} \log \binom{n}{\mathbf{k}} - h(\mathbf{x}) \right| \leq C \frac{\log n}{n}. \quad (5.2)$$

Let $X \in \text{EX}(d, N)$. We set for $0 \leq n \leq N$ and $|\mathbf{k}| = n$

$$p_{n, \mathbf{k}} = \mathbb{P}(X_1 = \dots = X_{k_1} = 1, \dots, X_{k_1+\dots+k_{d-1}+1} = \dots = X_n = d).$$

These $\binom{n+d-1}{d-1}$ numbers determine the law of any subsystem X_S of size $|S| = n$. It is convenient to define also $Y_i := \#\{1 \leq j \leq n : X_j = i\}$ for $i = 0, \dots, d-1$ and

$$q_{n, \mathbf{k}} := \mathbb{P}(Y_i = k_i, i = 0, \dots, d-1) = \binom{n}{\mathbf{k}} p_{n, \mathbf{k}}.$$

Since the vector $(q_{n,\mathbf{k}})_{|\mathbf{k}|=n}$ gives the law of the vector (Y_1, \dots, Y_d) we have in particular

$$\sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} = 1.$$

Second, we observe that for $|S| = n$

$$\left| \frac{\mathbb{H}(X_S)}{n} - \frac{1}{n} \sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} h(\mathbf{x}) \right| \leq C \frac{\log n}{n}. \quad (5.3)$$

Indeed

$$\begin{aligned} \frac{\mathbb{H}(X_S)}{n} &= -\frac{1}{n} \sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} \log \frac{q_{n,\mathbf{k}}}{\binom{n}{\mathbf{k}}} = \sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} \frac{1}{n} \log \binom{n}{\mathbf{k}} - \frac{1}{n} \sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} \log q_{n,\mathbf{k}} \\ &= \frac{1}{n} \sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} h(\mathbf{x}) + G(n), \quad |G(n)| \leq C \frac{\log n}{n}, \end{aligned}$$

where we use (5.2) and the fact that

$$-\sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} \log q_{n,\mathbf{k}} = \mathbb{H}(Y_1, \dots, Y_d) \leq d \log n,$$

since the support of the random vector (Y_1, \dots, Y_d) has cardinality at most n^d .

Third, we claim that, for $\varepsilon > 0$, there exists a constant C such that for all N and all $X \in \text{EX}(d, N)$, for all $n \in [\tilde{N}, N]$ with $\tilde{N} := \lfloor N^{\frac{2}{3}+\varepsilon} + 1 \rfloor$,

$$\left| \sum_{|\mathbf{k}|=n} q_{n,\mathbf{k}} h(\mathbf{x}) - \sum_{|\mathbf{K}|=N} q_{N,\mathbf{K}} h(\mathbf{X}) \right| \leq C N^{-\frac{1}{3}+\varepsilon}, \quad (5.4)$$

where $\mathbf{X} := \frac{1}{N} \mathbf{K}$ (no relation with the random variable X). By (5.3) and (5.4) we obtain for all $n \in [\tilde{N}, N]$ and $|S| = n$

$$\left| \frac{\mathbb{H}(X_S)}{n} - \frac{\mathbb{H}(X)}{N} \right| \leq C N^{-\frac{1}{3}+\varepsilon}. \quad (5.5)$$

Let us show how (5.5) implies (5.1). Using $\mathbb{H}(X_S) \leq \log d \cdot |S|$, $\sum_{S \subset I} c_S^I = 1$, we get

$$\sum_{|S| < \tilde{N}} c_S^I \text{MI}(S) \leq \sum_{S \subset I} c_S^I \times \log d \cdot \tilde{N} = \log d \cdot \tilde{N}.$$

Using (2.5), the exchangeability of X , $\sum_{n=0}^N c_n^N \binom{N}{n} = 1$ and (5.5), we estimate

$$\begin{aligned} \mathcal{I}^c(X) &\leq 2 \cdot \log d \cdot \tilde{N} + 2 \sum_{n=\tilde{N}}^N \binom{N}{n} c_n^N \mathbb{H}(X_{\{1,\dots,n\}}) - \mathbb{H}(X) \\ &\leq 2 \sum_{n=0}^N \binom{N}{n} c_n^N n \left(\frac{\mathbb{H}(X)}{N} + C N^{-\frac{1}{3}+\varepsilon} \right) - \mathbb{H}(X) + C \tilde{N}. \end{aligned}$$

Finally, using $c_n^N \binom{N}{n} = c_{N-n}^N \binom{N}{N-n}$ and $\sum_{n=0}^N c_n^N \binom{N}{n} = 1$

$$\begin{aligned} \mathcal{I}^c(X) &\leq \left(2 \sum_{n=0}^N c_n^N \binom{N}{n} \frac{n}{N} - 1 \right) \mathbb{H}(X) + CN \times N^{-\frac{1}{3}+\varepsilon} + C\tilde{N} \\ &\leq \left(\sum_{n=0}^N c_n^N \binom{N}{n} \left(\frac{n}{N} + \frac{N-n}{N} \right) - 1 \right) \mathbb{H}(X) + CN^{\frac{2}{3}+\varepsilon} = CN^{\frac{2}{3}+\varepsilon} \end{aligned}$$

and (5.1) is proved.

We turn now to the proof of (5.4). We claim first that

$$p_{n,\mathbf{k}} = \sum_{|\mathbf{K}|=N, \mathbf{K} \geq \mathbf{k}} p_{N,\mathbf{K}} \binom{N-n}{\mathbf{K}-\mathbf{k}}. \quad (5.6)$$

Indeed, notice that

$$p_{n,\mathbf{k}} = \sum_{j=1}^d p_{n+1,\mathbf{k}+\delta^j}, \quad \forall 0 \leq n < N, \quad \forall |\mathbf{k}| = n,$$

where $\delta^j := (\delta_1^j, \dots, \delta_d^j)$ with $\delta_i^j = 1$ if $i = j$, 0 otherwise. This in particular yields (5.6) for $N = n + 1$. Moreover if $|\mathbf{K}| = n + 1$ then

$$\binom{n+1}{\mathbf{K}} = \sum_{j=1}^d \binom{n}{\mathbf{K}-\delta^j} \mathbb{1}_{(\mathbf{K} \geq \delta^j)}.$$

Then, arguing by induction on $N \geq n$

$$\begin{aligned} p_{n,\mathbf{k}} &= \sum_{|\mathbf{K}|=N, \mathbf{K} \geq \mathbf{k}} p_{N,\mathbf{K}} \binom{N-n}{\mathbf{K}-\mathbf{k}} = \sum_{|\mathbf{K}|=N, \mathbf{K} \geq \mathbf{k}} \sum_{j=1}^d p_{N+1,\mathbf{K}+\delta^j} \binom{N-n}{\mathbf{K}-\mathbf{k}} \\ &= \sum_{|\mathbf{K}'|=N+1} p_{N+1,\mathbf{K}'} \sum_{j=1}^d \binom{N-n}{\mathbf{K}'-\mathbf{k}-\delta^j} \mathbb{1}_{(\mathbf{K}-\mathbf{k} \geq \delta^j)} \\ &= \sum_{|\mathbf{K}'|=N+1, \mathbf{K}' \geq \mathbf{k}} p_{N+1,\mathbf{K}'} \binom{N+1-n}{\mathbf{K}'-\mathbf{k}}. \end{aligned}$$

We recall that $q_{n,\mathbf{k}} = \binom{n}{\mathbf{k}} p_{n,\mathbf{k}}$. Notice that it is enough to prove Claim (5.4) in the case $q_{N,\mathbf{k}'} = \delta_{\mathbf{k}',\mathbf{K}}$, i.e., $p_{N,\mathbf{k}'} = \binom{N}{\mathbf{k}'}^{-1}$ for $\mathbf{k}' = \mathbf{K}$ and zero otherwise, if we find a constant C which does not depend on (N, n, \mathbf{K}) . Indeed, the two expressions are linear and the average of $CN^{-1/3+\varepsilon}$ will remain of the same order. Thus, we need to estimate:

$$a(N, \mathbf{K}, n, \mathbf{k}) := q_{n,\mathbf{k}} = \binom{n}{\mathbf{k}} \times \binom{N}{\mathbf{K}}^{-1} \binom{N-n}{\mathbf{K}-\mathbf{k}}.$$

Let $\mathbf{x} := \mathbf{k}/n \in [0, 1]^d$, $\mathbf{X} := \mathbf{K}/N \in [0, 1]^d$ and $\nu := n/(N - n)$. Formula (5.2) implies that $\frac{1}{n} \log a(N, \mathbf{K}, n, \mathbf{k})$ is equal to:

$$\underbrace{h(\mathbf{x}) - (1 + \nu^{-1})h(\mathbf{X}) + \nu^{-1}h(\mathbf{X} + \nu(\mathbf{X} - \mathbf{x}))}_{=: \phi_{\nu, \mathbf{x}}(\mathbf{x})} + G(N, n),$$

where $|G(N, n)| \leq \kappa(\log N)/n$, for some $\kappa = \kappa(d)$.

Let us now write for all $(x_1, \dots, x_{d-1}) \in [0, 1]^{d-1}$ such that $\sum_i x_i \leq 1$

$$H(x_1, \dots, x_{d-1}) := h(x_1, \dots, x_d), \quad x_d := 1 - x_1 - \dots - x_{d-1}.$$

Observe that for $i, j \leq d - 1$

$$\frac{\partial H}{\partial x_i} = \log\left(\frac{x_d}{x_i}\right), \quad \frac{\partial^2 H}{\partial x_i \partial x_j} = -\frac{1}{x_d} - \frac{1}{x_i} \mathbb{1}_{(i=j)}.$$

In particular the Hessian of H is negative-definite, since for all $a \in \mathbb{R}^{d-1} \setminus \{0\}$

$$\sum_{i,j=1}^{d-1} a_i a_j \frac{\partial^2 H}{\partial x_i \partial x_j} = -\frac{1}{x_d} \left(\sum_{i=1}^{d-1} a_i \right)^2 - \sum_{i=1}^{d-1} \frac{1}{x_i} a_i^2 \leq -\sum_{i=1}^{d-1} a_i^2$$

where we use the fact that $x_i \leq 1$. Hence, h is concave and we obtain

$$\phi_{\nu, \mathbf{x}}(\mathbf{x}) = \frac{\nu + 1}{\nu} \left[\frac{\nu}{\nu + 1} h(\mathbf{x}) + \frac{1}{\nu + 1} h((1 + \nu)\mathbf{X} - \nu\mathbf{x}) - h(\mathbf{X}) \right] \leq 0,$$

so that the maximum of $\phi_{\nu, \mathbf{x}}$ is $0 = \phi_{\nu, \mathbf{x}}(\mathbf{X})$. The second order derivative estimate gives:

$$\phi_{\nu, \mathbf{x}}(\mathbf{x}) \leq -2\|\mathbf{x} - \mathbf{X}\|^2 \quad \text{where } \|\mathbf{x}\| := \sqrt{x_1^2 + \dots + x_d^2}.$$

Combining with the bound $|G(N, n)| \leq \kappa(\log N)/n$ above, we get, for all $n < N$:

$$a(N, \mathbf{K}, n, \mathbf{k}) \leq N^\kappa \times e^{-2n\|\mathbf{x} - \mathbf{X}\|^2}.$$

Recall $n \geq \tilde{N} = N^{\frac{2}{3} + \varepsilon}$ and set $\delta := N^{-\frac{1}{3}}$ and

$$\omega := \sup_{\|\mathbf{x} - \mathbf{X}\| < \delta} \|h(\mathbf{X}) - h(\mathbf{x})\| \leq C \delta \log \frac{1}{\delta}.$$

Finally, using $h(\mathbf{x}) \leq \log d$,

$$\begin{aligned} \left| \sum_{|\mathbf{k}|=n} q_{n, \mathbf{k}} h(\mathbf{x}) - h(\mathbf{X}) \right| &\leq \omega \sum_{\|\mathbf{x} - \mathbf{X}\| < \delta} q_{n, \mathbf{k}} + 2 \log d \sum_{\|\mathbf{x} - \mathbf{X}\| \geq \delta} q_{n, \mathbf{k}} \\ &\leq C \delta \log \frac{1}{\delta} + C n^d N^\kappa e^{-2\tilde{N}\delta^2} \leq C(\log N) N^{-\frac{1}{3}} + C N^{\kappa+d} e^{-2N^\varepsilon} \leq C N^{-\frac{1}{3} + \varepsilon}. \end{aligned}$$

Then (5.4) and the proposition are proved. \square

APPENDIX A. ENTROPY

In this Appendix, we recall needed facts from basic information theory. The main object is the entropy functional which may be said to quantify the randomness of a random variable.

Let X be a random variable taking values in a finite space E . We define the *entropy* of X

$$H(X) := - \sum_{x \in E} P_X(x) \log(P_X(x)), \quad P_X(x) := \mathbb{P}(X = x),$$

where we adopt the convention

$$0 \cdot \log(0) = 0 \cdot \log(+\infty) = 0.$$

We recall that

$$0 \leq H(X) \leq \log |E|, \tag{A.1}$$

More precisely, $H(X)$ is minimal iff X is a constant, it is maximal iff X is uniform over E . To prove (A.1), just notice that since $\varphi \geq 0$ and $\varphi(x) = 0$ if and only if $x \in \{0, 1\}$, and by strict convexity of $x \mapsto \varphi(x) = x \log x$ and Jensen's inequality

$$\begin{aligned} \log |E| - H(X) &= \frac{1}{|E|} \sum_{x \in E} P_X(x) |E| (\log(P_X(x)) + \log |E|) \\ &= \frac{1}{|E|} \sum_{x \in E} \varphi(P_X(x) |E|) \geq \varphi \left(\frac{1}{|E|} \sum_{x \in E} P_X(x) |E| \right) = \varphi(1) = 0, \end{aligned}$$

with $\log |E| - H(X) = 0$ if and only if $P_X(x) |E|$ is constant in $x \in E$.

If we have a E -valued random variable X and a F -valued random variable Y defined on the same probability space, with E and F finite, we can consider the vector (X, Y) as a $E \times F$ -valued random variable. The entropy of (X, Y) is then

$$H(X, Y) := - \sum_{x, y} P_{(X, Y)}(x, y) \log(P_{(X, Y)}(x, y)), \quad P_{(X, Y)}(x, y) := \mathbb{P}(X = x, Y = y).$$

This entropy $H(X, Y)$ does not only depends on the (separate) laws of X and Y but on the extent to which the "randomness of the two variables is shared". The following notions formalize this idea.

A.1. Conditional Entropy. The *conditional entropy* of X given Y is:

$$H(X | Y) := H(X, Y) - H(Y).$$

We claim that

$$0 \leq H(X | Y) \leq H(X) \leq H(X, Y). \tag{A.2}$$

Remark that $P_X(x)$ and $P_Y(y)$, defined in the obvious way, are the marginals of $P_{(X, Y)}(x, y)$, i.e.

$$P_X(x) = \sum_y P_{(X, Y)}(x, y), \quad P_Y(y) = \sum_x P_{(X, Y)}(x, y).$$

In particular, $P_X(x) \geq P_{(X,Y)}(x, y)$ for all x, y . Therefore

$$\sum_{x,y} P_{(X,Y)}(x, y) \log \left(\frac{P_{(X,Y)}(x, y)}{P_X(x)} \right) \leq 0$$

which yields

$$H(X, Y) = - \sum_{x,y} P_{(X,Y)}(x, y) \log P_{(X,Y)}(x, y) \geq - \sum_x P_X(x) \log P_X(x) = H(X),$$

i.e. $H(X, Y) \geq H(X)$ and $H(X|Y) \geq 0$. Therefore

$$H(X, Y) \geq \max\{H(X), H(Y)\}. \quad (\text{A.3})$$

Moreover $H(X, Y) = H(X)$, i.e. $H(Y|X) = 0$, if and only if $P_{(X,Y)}(x, y) = P_X(x)$ whenever $P_{(X,Y)}(x, y) \neq 0$, which means that Y is a function of X .

On the other hand,

$$H(X, Y) \leq H(X) + H(Y) \quad (\text{A.4})$$

with equality, i.e., $H(Y|X) = H(Y)$, if and only if X and Y are independent. This shows that $H(X|Y) \leq H(X)$ and completes the proof of (A.2).

Formula (A.4) can be shown by considering the Kullback-Leibler divergence or relative entropy:

$$I := \sum_{x,y} P_{(X,Y)}(x, y) \log \left(\frac{P_{(X,Y)}(x, y)}{P_X(x) P_Y(y)} \right).$$

Since $\log(\cdot)$ is concave, by Jensen's inequality

$$-I \leq \log \left(\sum_{x,y} P_{(X,Y)}(x, y) \frac{P_X(x) P_Y(y)}{P_{(X,Y)}(x, y)} \right) = \log \left(\sum_{x,y} P_X(x) P_Y(y) \right) = 0.$$

By strict concavity, $I = 0$ if and only if $P_{(X,Y)}(x, y) = P_X(x) P_Y(y)$ for all x, y , i.e., whenever X and Y are independent.

By the above considerations, $H(X|Y) \in [0, H(X)]$ is a measure of the uncertainty associated with X if Y is known. It is minimal iff X is a function of Y and it maximal iff X and Y are independent.

A.2. Adding information decreases uncertainty. Let us consider three random variables $(X, Y, Z) \mapsto E \times F \times G$ with E, F, G finite. Then we have that

$$H(X | (Y, Z)) \leq H(X | Y). \quad (\text{A.5})$$

Indeed, this is equivalent to

$$H(X, Y, Z) + H(Y) \leq H(X, Y) + H(Y, Z).$$

Consider the quantity

$$J := \sum_{x,y,z} P_{(X,Y,Z)}(x, y, z) \log \left(\frac{P_{(X,Y,Z)}(x, y, z) P_Y(y)}{P_{(X,Y)}(x, y) P_{(Y,Z)}(y, z)} \right).$$

Since $-\log(\cdot)$ is convex, by Jensen's inequality

$$J \geq -\log \left(\sum_{x,y} \frac{P_{(X,Y)}(x,y) \sum_z P_{(Y,Z)}(y,z)}{P_Y(y)} \right) = -\log \left(\sum_{x,y} P_{(X,Y)}(x,y) \right) = 0,$$

and the inequality follows.

A.3. Mutual Information. Finally, we recall the notion of *mutual information* between two random variables X and Y defined on the same probability space:

$$\begin{aligned} \text{MI}(X, Y) &:= H(X) + H(Y) - H(X, Y) \\ &= H(X) - H(X | Y) = H(Y) - H(Y | X) \\ &= \sum_{x,y} P_{(X,Y)}(x,y) \log \left(\frac{P_{(X,Y)}(x,y)}{P_X(x) P_Y(y)} \right). \end{aligned}$$

This quantity is a measure of the common randomness of X and Y . By (A.3) and (A.4) we have $\text{MI}(X, Y) \in [0, \min\{H(X), H(Y)\}]$. $\text{MI}(X, Y)$ is minimal (zero) iff X, Y are independent and maximal, i.e. equal to $\min\{H(X), H(Y)\}$, iff one variable is a function of the other.

Mutual information is non-decreasing. Let $X, X', Y, Y', \hat{X}, \hat{Y}$ be random variables such that X, X' , resp. Y, Y' , are (deterministic) functions of \hat{X} , resp. \hat{Y} . Then:

$$\text{MI}(X, Y) \leq \text{MI}(\hat{X}, \hat{Y}) \quad (\text{A.6})$$

Mutual information is almost additive:

$$|\text{MI}((X, Y), (X', Y')) - (\text{MI}(X, X') + \text{MI}(Y, Y'))| \leq \text{MI}(\hat{X}, \hat{Y}). \quad (\text{A.7})$$

These properties follow from the properties of conditional entropy. First,

$$\begin{aligned} \text{MI}(\hat{X}, \hat{Y}) &= H(\hat{X}) + H(\hat{Y}) - H(\hat{X}, \hat{Y}) \\ &= H(X) + H(\hat{X}|X) + H(Y) + H(\hat{Y}|Y) - H(X, Y) - H(\hat{X}|X, Y) - H(\hat{Y}|\hat{X}, Y) \\ &= \text{MI}(X, Y) + (H(\hat{X}|X) - H(\hat{X}|X, Y)) + (H(\hat{Y}|Y) - H(\hat{Y}|\hat{X}, Y)). \end{aligned}$$

(A.6) now follows from (A.5). Second,

$$\begin{aligned} \text{MI}((X, Y), (X', Y')) &= H(X, Y) + H(X', Y') - H(X, X', Y, Y') \\ &= H(X) + H(Y) - \text{MI}(X, Y) + H(X') + H(Y') - \text{MI}(X', Y') \\ &\quad - H(X, X') - H(Y, Y') + \text{MI}((X, X'), (Y, Y')) \\ &= H(X) + H(X') - H(X, X') + H(Y) + H(Y') - H(Y, Y') \\ &\quad + (\text{MI}((X, X'), (Y, Y')) - \text{MI}(X, Y) - \text{MI}(X', Y')) \\ &= \text{MI}(X, X') + \text{MI}(Y, Y') + (\text{MI}((X, X'), (Y, Y')) - \text{MI}(X, Y) - \text{MI}(X', Y')). \end{aligned}$$

The nonnegativity of mutual information and (A.6) yields

$$\begin{aligned} -\min(\text{MI}(X, Y), \text{MI}(X', Y')) &\leq \text{MI}((X, Y), (X', Y')) - (\text{MI}(X, X') + \text{MI}(Y, Y')) \\ &\leq \text{MI}((X, X'), (Y, Y')). \end{aligned}$$

(A.7) follows.

REFERENCES

- [1] L. Barnett, C. L. Buckley, S. Bullock, Neural complexity and structural connectivity, *Phys. Rev. E* **79** (2009), 051914.
- [2] J. Bertoin, *Random Fragmentation And Coagulation Processes*, Cambridge University Press (2006).
- [3] J. Buzzi, L. Zambotti, *Approximate Maximizers of Intricacy Functionals*, preprint (2009), <http://arxiv.org/abs/0909.2120>.
- [4] T. Cover, J. Thomas, *Elements of Information Theory*, John Wiley & Sons, 2006.
- [5] M. De Lucia, M. Bottaccio, M. Montuori, L. Pietronero, A topological approach to neural complexity, *Phys. Rev. E* **71**, 016114 (2005) — arXiv:nlin/0411011v1.
- [6] W. Feller, *An introduction to probability theory and its applications. Vol. 2*, John Wiley & Sons, 1971.
- [7] G. Edelman, J. Gally, Degeneracy and complexity in biological systems, *Proc. Natl. Acad. Sci. USA*, **98** (2001), 13763–13768.
- [8] A. Greven, G. Keller, G. Warnecke, *Entropy*, Princeton University Press, 2003.
- [9] K. Holthausen, O. Breidbach, Analytical description of the evolution of neural networks: learning rules and complexity, *Biol. Cybern.* **81** (1999), 169–176.
- [10] J. Krichmar, D. Nitz, J. Gally, G. Edelman, Characterizing functional hippocampal pathways in a brain-based device as it solves a spatial memory task, *Proc. Natl. Acad. Sci. USA*, **102** (2005), 2111–2116.
- [11] A. Seth, E. Izhikevich, G. Reeke, G. Edelman, Theories and measures of consciousness: an extended framework, *Proc. Natl. Acad. Sci. USA*, **103** (2006), 10799–10804.
- [12] Anil K Seth (2007), Models of consciousness, *Scholarpedia*, 2(1):1328.
- [13] M. P. Shanahan, Dynamical complexity in small-world networks of spiking neurons, *Phys. Rev. E* **78**, 041924 (2008).
- [14] O. Sporns, G. Tononi, G. Edelman, Connectivity and complexity: the relationship between neuroanatomy and brain dynamics, *Neural Netw.* 2000 Oct-Nov;13(8-9):909-22.
- [15] O. Sporns, Networks analysis, complexity, and brain function, *Complexity* **8** (2002), 56 - 60.
- [16] O. Sporns (2007), Complexity, *Scholarpedia*, 2(10):1623.
- [17] M. Talagrand, *Spin Glasses: A Challenge for Mathematicians*, Springer, 2003.
- [18] G. Tononi, O. Sporns, G. Edelman, A measure for brain complexity: relating functional segregation and integration in the nervous system, *Proc. Natl. Acad. Sci. USA*, **91** (1994), 5033–5037.
- [19] G. Tononi, O. Sporns, G. Edelman, A complexity measure for selective matching of signals by the brain, *Proc. Natl. Acad. Sci. USA*, **93** (1996), 3422–3427.
- [20] G. Tononi, O. Sporns, G. Edelman, Measures of degeneracy and redundancy in biological networks, *Proc. Natl. Acad. Sci. USA*, **96** (1999), 3257–3262.

LABORATOIRE DE MATHÉMATIQUE D'ORSAY - C.N.R.S. (U.M.R. 8628) & UNIVERSITÉ PARIS-SUD, UNIVERSITÉ PARIS-SUD, F-91405 ORSAY CEDEX, FRANCE

E-mail address: jerome.buzzi@math.u-psud.fr

LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES (CNRS U.M.R. 7599) AND UNIVERSITÉ PARIS 6 – PIERRE ET MARIE CURIE, U.F.R. MATHÉMATIQUES, CASE 188, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address: lorenzo.zambotti@upmc.fr