



HAL
open science

Stochastic Cahn-Hilliard equation with double singular nonlinearities and two reflections

Arnaud Debussche, Ludovic Goudenège

► **To cite this version:**

Arnaud Debussche, Ludovic Goudenège. Stochastic Cahn-Hilliard equation with double singular nonlinearities and two reflections. *SIAM Journal on Mathematical Analysis*, 2011, 43 (3), pp.1473-1494. 10.1137/090769636 . hal-00411599

HAL Id: hal-00411599

<https://hal.science/hal-00411599v1>

Submitted on 28 Aug 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stochastic Cahn-Hilliard equation with double singular nonlinearities and two reflections

Arnaud Debussche, Ludovic Goudenège

ENS Cachan - Antenne de Bretagne, Campus de Ker-Lann, 35170 Bruz, France
arnaud.debussche@bretagne.ens-cachan.fr, ludovic.goudeneg@bretagne.ens-cachan.fr

Résumé

On considère une équation aux dérivées partielles stochastique possédant deux non-linéarités de type logarithmique, avec deux réflexions en 1 et -1 sous la contrainte de conservation de masse. L'équation, dirigée par un bruit blanc en espace et en temps, contient un double Laplacien. L'absence de principe de maximum pour le double Laplacien pose des difficultés pour l'utilisation d'une méthode classique de pénalisation, pour laquelle une importante propriété de monotonie est utilisée. Etant inspiré par les travaux de Debussche, Goudenège et Zambotti, on démontre l'existence et l'unicité de solutions pour des données initiales entre -1 et 1 . Enfin, on démontre que l'unique mesure invariante est ergodique, et on énonce un résultat de mélange exponentiel.

Abstract

We consider a stochastic partial differential equation with two logarithmic nonlinearities, with two reflections at 1 and -1 and with a constraint of conservation of the space average. The equation, driven by the derivative in space of a space-time white noise, contains a bi-Laplacian in the drift. The lack of the maximum principle for the bi-Laplacian generates difficulties for the classical penalization method, which uses a crucial monotonicity property. Being inspired by the works of Debussche, Goudenège and Zambotti, we obtain existence and uniqueness of solution for initial conditions in the interval $(-1, 1)$. Finally, we prove that the unique invariant measure is ergodic, and we give a result of exponential mixing.

Introduction and main results

The Cahn-Hilliard-Cook equation is a model to describe phase separation in a binary alloy (see [6], [7] and [8]) in the presence of thermal fluctuations (see [11] and [27]). It takes the form:

$$\begin{cases} \partial_t u = -\frac{1}{2} \Delta (\Delta u - \psi(u)) + \dot{\xi}, & \text{on } \Omega \subset \mathbb{R}^n, \\ \nabla u \cdot \nu = 0 = \nabla (\Delta u - \psi(u)) \cdot \nu, & \text{on } \partial\Omega, \end{cases} \quad (0.1)$$

where t denotes the time variable and Δ is the Laplace operator. Also $u \in [-1, 1]$ represents the ratio between the two species and the noise term $\dot{\xi}$ accounts for the thermal fluctuations. The nonlinear term ψ has the double-logarithmic form:

$$\psi : u \mapsto \frac{\theta}{2} \ln \left(\frac{1+u}{1-u} \right) - \theta_c u, \quad (0.2)$$

AMS 2000 subject classifications. 60H15, 60H07, 37L40.

Key words and phrases : Cahn-Hilliard, stochastic partial differential equations, integration by parts formulae, reflection measures, invariant measures, singular nonlinearity, double, two.

where θ and θ_c are temperatures with $\theta < \theta_c$.

The study of this equation presents several difficulties. First, the singularities at ± 1 have to be treated carefully. Also, since it is a fourth order equation, no comparison principle holds.

The deterministic equation where ψ is replaced by a polynomial function have first been studied (see [7], [27] and [32]). Then non smooth ψ have been considered (see [5] and [17]).

Phase separation have been analysed thanks to this model: see for example the survey [31], and the references therein, or others recent results on spinodal decomposition and nucleation in [1, 4, 24, 28, 29, 35, 36, 37].

In the case of a polynomial nonlinearity, some results have been obtained in the stochastic case (see [2, 3, 9, 10, 12, 19]).

Note that the solutions of the equation with polynomial nonlinearity do not remain in $[-1, 1]$ in general, and their physical interpretation is not clear.

To our knowledge, the case of the logarithmic nonlinearity in the presence of noise have never been studied. The presence of noise has a strong effect and equation (0.1) cannot have a solution. Indeed, a solution should remain in $[-1, 1]$ which is impossible with an additive noise. Two reflection measures have to be added to the model to remedy this problem. In this article, we propose to study:

$$\begin{cases} \partial_t u = -\frac{1}{2}\Delta(\Delta u - \psi(u) + \eta_- - \eta_+) + \dot{\xi}, & \text{with } \theta \in [0, 1] = \Omega, \\ \nabla u \cdot \nu = 0 = \nabla(\Delta u) \cdot \nu, & \text{on } \partial\Omega, \end{cases} \quad (0.3)$$

where the measures are subject to the contact conditions almost surely:

$$\int (1 + u)d\eta_- = \int (1 - u)d\eta_+ = 0. \quad (0.4)$$

The stochastic heat equation with reflection, *i.e.* when the fourth order operator is replaced by the Laplace operator, is a model for the evolution of random interfaces near a hard wall. It has been extensively studied in the literature (see [16], [21], [22], [33] [38], [39] and [40]). Essential tools in these articles are the comparison principle and the fact that the underlying Dirichlet form is symmetric so that the invariant measure is known explicitly.

In our case, we consider a noise which is obtained as the space derivative of the space-time white noise. In other words, the noise is the time derivative of a cylindrical Wiener process in $H^{-1}(0, 1)$. This is physically reasonable since the Cahn-Hilliard equation can be interpreted as a gradient system in this space. With such noise, the system is still symmetric and the invariant measure is known explicitly. As in the second order case, we use this fact in an essential way.

However, as already mentioned, no comparison principle holds and new techniques have to be developed. The equation (0.3) has been studied with a single reflection and when no nonlinear term is taken into account in [18]. The reflection is introduced to enforce positivity of the solution. Various techniques have been introduced to overcome this lack of comparison principle. Moreover, as in the second order case, an integration by part formula for the invariant measure has been derived. Then, in [23], a singular nonlinearity of the form $u^{-\alpha}$ or $\ln u$ have been considered. Existence and uniqueness of solutions have been obtained and using the integration by parts formula as in [39], it has been proved that the reflection measure vanishes if and only if $\alpha \geq 3$. In particular, for a logarithmic nonlinearity, the reflection is active.

Here, we consider the original Cahn-Hilliard-Cook model (0.1) with the double-logarithmic nonlinear term (0.2). The noise is as in the above mentioned articles and we still have an explicit invariant measure. Our method mixes ideas from [18], [23] and [39]. Additional difficulties are overcome, the main one being to understand how to deal with the nonlinear term. Indeed, in [23], the positivity of the nonlinear term was essential. We overcome this difficulty thanks to a delicate a priori estimate. Our main results state that equations (0.3), (0.4) together with an initial condition have a unique solution (see Proposition 2.3 and Theorem 1.1). As in [18], it is constructed thanks to the gradient structure of (0.3) and strong Feller property. Moreover, we prove that this solution is the limit of the solution of the Cahn-Hilliard-Cook equation with polynomial nonlinearity without reflections. This justifies the use of the polynomial models. We also prove that the invariant measure is unique and ergodic. Such property is very easy to obtain if θ_c is small (see [18]) or in

the polynomial case (see [12]). Finally, a stronger result of exponential mixing is given in the last Theorem 3.1. It is based on coupling and arguments developed by Odasso in [34].

In future studies, we shall generalize the integration by part formula obtain in [18] to prove that the reflection measure does not vanish. The presence of two reflection measures introduces additional difficulties. In the second order case, this has been studied in [20].

1 Preliminaries

We denote by $\langle \cdot, \cdot \rangle$ the scalar product in $L^2(0, 1)$; A is the realization in $L^2(0, 1)$ of the Laplace operator with Neumann boundary condition, i.e.:

$$D(A) = \text{Domain of } A = \{h \in W^{2,2}(0, 1) : h'(0) = h'(1) = 0\}$$

where we use $W^{n,p}$ and $\|\cdot\|_{W^{n,p}}$ to denote the Sobolev space $W^{n,p}(0, 1)$ and its associated norm. Remark that A is self-adjoint on $L^2(0, 1)$ and we have a complete orthonormal system of eigenvectors $(e_i)_{i \in \mathbb{N}}$ in $L^2(0, 1)$ for the eigenvalues $(\lambda_i)_{i \in \mathbb{N}}$. We denote by \bar{h} the mean of $h \in L^2(0, 1)$:

$$\bar{h} = \int_0^1 h(\theta) d\theta.$$

We remark that A is invertible on the space of functions with 0 average. In general, we define $(-A)^{-1}h = (-A)^{-1}(h - \bar{h}) + \bar{h}$.

For $\gamma \in \mathbb{R}$, we define $(-A)^\gamma$ by classical interpolation. We set $V_\gamma := D((-A)^{\gamma/2})$. It is endowed with the classical seminorm and norm :

$$|h|_\gamma = \left(\sum_{i=1}^{+\infty} (-\lambda_i)^\gamma h_i^2 \right)^{1/2}, \quad \|h\|_\gamma = (|h|_\gamma^2 + \bar{h}^2)^{1/2}, \quad \text{for } h = \sum_{i \in \mathbb{N}} h_i e_i.$$

$|\cdot|_\gamma$ is associated to the scalar product $(\cdot, \cdot)_\gamma$. To lighten notations, we set $(\cdot, \cdot) := (\cdot, \cdot)_{-1}$ and $H := V_{-1}$. The average can be defined in any V_γ by $\bar{h} = (h, e_0)$. It plays an important role and we often work with functions with a fixed average $c \in \mathbb{R}$. We define $H_c = \{h \in H : \bar{h} = c\}$ for all $c \in \mathbb{R}$.

We use the following regularization operators:

$$Q_N x = \frac{1}{N} \sum_{n=0}^N \sum_{i=0}^n (x, e_i) e_i.$$

It is defined on $L^2(0, 1)$ and can be extended to any V_γ . Clearly $Q_N x$ converges to x in V_γ if $x \in V_\gamma$. Moreover, it is well known that if $x \in C([0, 1]; \mathbb{R})$, then the convergence holds in $C([0, 1]; \mathbb{R})$. Note also that Q_N is self-adjoint in V_γ and commutes with A .

The covariance operator of the noise is the operator B defined by

$$B = \frac{\partial}{\partial \theta}, \quad D(B) = W_0^{1,2}(0, 1).$$

Note that

$$B^* = -\frac{\partial}{\partial \theta}, \quad D(B^*) = W^{1,2}(0, 1), \quad BB^* = -A.$$

We denote by $\mathcal{B}_b(H_c)$ the space of all Borel bounded functions on H_c . We set $O_{s,t} := [s, t] \times [0, 1]$ for $s, t \in [0, T]$ with $s < t$ and $T > 0$, and $O_t = O_{0,t}$ for $0 \leq t \leq T$. Given a measure ζ on $O_{s,t}$ and a continuous function v on $O_{s,t}$, we write

$$\langle v, \zeta \rangle_{O_{s,t}} := \int_{O_{s,t}} v \, d\zeta.$$

For $\lambda \in \mathbb{R}$, we define:

$$f(x) := \begin{cases} +\infty, & \text{for all } x \leq -1, \\ \ln\left(\frac{1-x}{1+x}\right) + \lambda x, & \text{for all } x \in (-1, 1), \\ -\infty, & \text{for all } x \geq 1, \end{cases} \quad (1.1)$$

and the following antiderivative F of $-f$:

$$F(x) = (1+x)\ln(1+x) + (1-x)\ln(1-x) - \frac{\lambda}{2}x^2, \text{ for all } x \in (-1, 1).$$

With these notations, we rewrite (0.3) in the abstract form:

$$\begin{cases} dX = -\frac{1}{2}A(AX + f(X) + \eta_- - \eta_+) dt + BdW, \\ \langle (1+X), \eta_- \rangle_{O_T} = \langle (1-X), \eta_+ \rangle_{O_T} = 0, \\ X(0, x) = x \text{ for } x \in V_{-1}, \end{cases} \quad (1.2)$$

where W is a cylindrical Wiener process on $L^2(0, 1)$.

Definition 1.1 *Let $x \in \mathcal{C}([0, 1]; [-1, 1])$. We say that $((X(t, x))_{t \in [0, T]}, \eta_+, \eta_-, W)$, defined on a filtered complete probability space $(\Omega, \mathbb{P}, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]})$, is a weak solution to (0.3) on $[0, T]$ for the initial condition x if:*

- (a) a.s. $X \in \mathcal{C}((0, T] \times [0, 1]; [-1, 1]) \cap \mathcal{C}([0, T]; H)$ and $X(0, x) = x$,
- (b) a.s. η_{\pm} are two positive measures on $(0, T] \times [0, 1]$, such that $\eta_{\pm}(O_{\delta, T}) < +\infty$ for all $\delta \in (0, T]$,
- (c) W is a cylindrical Wiener process on $L^2(0, 1)$,
- (d) the process $(X(\cdot, x), W)$ is (\mathcal{F}_t) -adapted,
- (e) a.s. $f(X(\cdot, x)) \in L^1(O_T)$,
- (f) for all $h \in D(A^2)$ and for all $0 < \delta \leq t \leq T$:

$$\begin{aligned} \langle X(t, x), h \rangle &= \langle X(\delta, x), h \rangle - \frac{1}{2} \int_{\delta}^t \langle X(s, x), A^2 h \rangle ds - \frac{1}{2} \int_{\delta}^t \langle Ah, f(X(s, x)) \rangle ds \\ &\quad - \frac{1}{2} \langle Ah, \eta_+ \rangle_{O_{\delta, t}} + \frac{1}{2} \langle Ah, \eta_- \rangle_{O_{\delta, t}} - \int_{\delta}^t \langle Bh, dW \rangle, \quad \text{a.s.}, \end{aligned}$$

- (g) a.s. the contact properties hold :
 $\text{supp}(\eta_-) \subset \{(t, \theta) \in O_T / X(t, x)(\theta) = -1\}$ and $\text{supp}(\eta_+) \subset \{(t, \theta) \in O_T / X(t, x)(\theta) = 1\}$,
that is,

$$\langle (1+X), \eta_- \rangle_{O_T} = \langle (1-X), \eta_+ \rangle_{O_T} = 0.$$

Finally, a weak solution (X, η_+, η_-, W) is a strong solution if the process $t \mapsto X(t, x)$ is adapted to the filtration $t \mapsto \sigma(W(s, \cdot), s \in [0, t])$

Remark 1.1 *In (f), the only term where we use the function f is well defined. Indeed, by (e) we have $f(X(\cdot, x)) \in L^1(O_T)$ and by Sobolev embedding $Ah \in D(A) \subset L^\infty(O_T)$. Hence the notation $\langle \cdot, \cdot \rangle$ should be interpreted as a duality between L^∞ and L^1 .*

The solution of the linear equation with initial data $x \in H$ is given by

$$Z(t, x) = e^{-tA^2/2}x + \int_0^t e^{-(t-s)A^2/2}BdW_s.$$

As easily seen this process is in $\mathcal{C}([0, +\infty[; H)$ (see [14]). In particular, the mean of Z is constant and the law of the process $Z(t, x)$ is the Gaussian measure:

$$Z(t, x) \sim \mathcal{N}(e^{-tA^2/2}x, Q_t), \quad Q_t = \int_0^t e^{-sA^2/2}BB^*e^{-sA^2/2}ds = (-A)^{-1}(I - e^{-tA^2}).$$

If we let $t \rightarrow +\infty$, the law of $Z(t, x)$ converges to the Gaussian measure on L^2 :

$$\mu_c := \mathcal{N}(ce_0, (-A)^{-1}), \quad \text{where } c = \bar{x}.$$

Notice that μ_c is concentrated on $H_c \cap C([0, T])$.

In order to solve equation (1.2), we use polynomial approximations of this equation. We denote by $\{f^n\}_{n \in \mathbb{N}}$ the sequence of polynomial functions which converges to the function f on $(-1, 1)$, defined for $n \in \mathbb{N}$ by:

$$f^n(x) = -2 \sum_{k=0}^n \frac{x^{2k+1}}{(2k+1)} + \lambda x, \quad \text{for all } x \in \mathbb{R}.$$

We use the following antiderivative F^n of $-f^n$ defined by:

$$F^n(x) = 2 \sum_{k=0}^n \frac{x^{2k+2}}{(2k+2)(2k+1)} - \frac{\lambda}{2}x^2, \quad \text{for all } x \in \mathbb{R}.$$

Then for $n \in \mathbb{N}$, we study for the following polynomial approximation of (1.2) with an initial condition $x \in H$:

$$\begin{cases} dX^n + \frac{1}{2}(A^2X^n + Af^n(X^n))dt = BdW, \\ X^n(0, x) = x. \end{cases} \quad (1.3)$$

This equation has been studied in [12] in the case $B = I$. The results generalize immediately and it can be proved that for any $x \in H$, there exists a unique solution $X^n(\cdot, x)$ a.s. in $\mathcal{C}([0, T]; H) \cap L^{2n+2}((0, T) \times (0, 1))$. It is a solution in the mild or weak sense. Moreover the average of $X^n(t, x)$ does not depend on t .

For each $c \in \mathbb{R}$, (1.2) defines a transition semigroup $(P_t^{n,c})_{t \geq 0}$:

$$P_t^{n,c}\phi(x) = \mathbb{E}[\phi(X^n(t, x))], \quad t \geq 0, x \in H_c, \phi \in \mathcal{B}_b(H_c), n \in \mathbb{N}^*.$$

Existence of an invariant measure can be proved as in [12].

Using Galerkin approximation and Bismut-Elworthy-Li formula, it can be seen that $(P_t^{n,c})_{t \geq 0}$ is Strong Feller. More precisely, for all $\phi \in \mathcal{B}_b(H_c)$, $n \in \mathbb{N}$ and $t > 0$:

$$|P_t^{n,c}\phi(x) - P_t^{n,c}\phi(y)| \leq \frac{2e^{\lambda^2 t/4}}{\lambda\sqrt{t}} \|\phi\|_\infty |x - y|_{-1}, \quad \text{for all } x, y \in H_c. \quad (1.4)$$

Irreducibility follows from a control argument. By Doob Theorem we deduce that there exists an unique and ergodic invariant measure ν_c^n .

It is classical that equation (1.3) is a gradient system in H_c and can be rewritten as:

$$\begin{cases} dX^n + \frac{1}{2}A(AX^n - \nabla U^n(X^n))dt = BdW, \\ X^n(0, x) = x \in L^2(0, 1), \end{cases} \quad (1.5)$$

where ∇ denotes the gradient in the Hilbert space $L^2(0, 1)$, and:

$$U^n(x) := \int_0^1 F^n(x(\theta))d\theta, \quad x \in L^2(0, 1).$$

The measure ν_c^n is therefore given by:

$$\nu_c^n(\mathrm{d}x) = \frac{1}{Z_c^n} \exp(-U^n(x)) \mu_c(\mathrm{d}x),$$

where Z_c^n is a normalization constant.

We prove in section 2 that, for $c \in (-1, 1)$, the sequence $(\nu_c^n)_{n \in \mathbb{N}}$ converges to the measure

$$\nu_c(\mathrm{d}x) = \frac{1}{Z_c} \exp(-U(x)) \mathbb{1}_{x \in K} \mu_c(\mathrm{d}x),$$

where

$$U(x) := \int_0^1 F(x(\theta)) \mathrm{d}\theta, \quad x \in L^2(0, 1).$$

and

$$K = \{x \in L^2 : 1 \geq x \geq -1\}.$$

In section 2, we prove the following result.

Theorem 1.1 *Let $c \in (-1, 1)$. Let $x \in K$ such that $\bar{x} = c$, then there exists a continuous process denoted $(X(t, x))_{t \geq 0}$ and two nonnegative measures η_+^x and η_-^x such that $((X(t, x))_{t \geq 0}, \eta_+^x, \eta_-^x, W)$ is the unique strong solution of (0.3) with $X(0, x) = x$ a.s.*

The Markov process $(X(t, x), t \geq 0, x \in K \cap H_c)$ is continuous and has P^c for transition semigroup which is strong Feller on H_c .

For all $x \in K \cap H_c$ and $0 = t_0 < t_1 < \dots < t_m$, $(X(t_i, x), i = 1, \dots, m)$ is the limit in distribution of $(X^n(t_i, x))_{i=1, \dots, m}$.

Finally ν_c is an invariant measure for P^c .

In all the article, C denotes a constant which may depend on T and its value may change from one line to another.

2 Proof of Theorem 1.1

2.1 Pathwise uniqueness

We first prove that for any pair $(X^i, \eta_+^i, \eta_-^i, W), i = 1, 2$, of weak solutions of (0.3) defined on the same probability space with the same driving noise W and with $X^1(0) = X^2(0)$, we have $(X^1, \eta_+^1, \eta_-^1) = (X^2, \eta_+^2, \eta_-^2)$. This pathwise uniqueness will be used in the next subsection to construct stationary strong solutions of (0.3).

Proposition 2.1 *Let $x \in \mathcal{C}([0, 1]; [-1, 1])$. Let $(X^i, \eta_+^i, \eta_-^i, W), i = 1, 2$ be two weak solutions of (0.3) with $X^1(0) = X^2(0) = x$. Then $(X^1, \eta_+^1, \eta_-^1) = (X^2, \eta_+^2, \eta_-^2)$.*

Proof : We use the following Lemma from [23].

Lemma 2.1 *Let ζ be a finite measure on $O_{\delta, T}$ and $V \in \mathcal{C}(O_{\delta, T})$. Suppose that there exists a positive continuous function $c_T : [0, T] \rightarrow \mathbb{R}^+$ such that :*

$$i) \text{ for all } r \in [\delta, T], \text{ for all } h \in \mathcal{C}([0, 1]), \text{ such that } \bar{h} = 0, \langle h, \zeta \rangle_{O_{r, T}} = 0,$$

$$ii) \text{ for all } r \in [\delta, T], \overline{V(r, \cdot)} = c_T(r) \text{ with } \langle V, \zeta \rangle_{O_{r, T}} = 0,$$

then ζ is the null measure.

Let $Y(t) = X^1(t, x) - X^2(t, x)$, $\zeta_+ = \eta_+^1 - \eta_+^2$ and $\zeta_- = \eta_-^1 - \eta_-^2$, Y is the solution of the following equation:

$$\begin{cases} \mathrm{d}Y = -\frac{1}{2}A(AY + (f(X^1) - f(X^2))) + \zeta_- - \zeta_+ \mathrm{d}t, \\ Y(0) = 0. \end{cases} \quad (2.1)$$

Taking the scalar product in H with $Y^N = Q_N Y$ and integrating in time, we obtain since Y has zero average:

$$|Y^N(t)|_{-1}^2 - |Y^N(\delta)|_{-1}^2 = - \int_{\delta}^t (|Y^N(s)|_1^2 - \langle f(X^1) - f(X^2), Y^N \rangle) ds + \langle \zeta_- - \zeta_+, Y^N \rangle_{O_{\delta,t}}. \quad (2.2)$$

For all $s \in [\delta, t]$,

$$\begin{aligned} & \langle Y^N(s) - Y(s), f(X^1(s, x)) - f(X^2(s, x)) \rangle \\ & \leq \|Y^N(s) - Y(s)\|_{L^\infty([0,1])} \|f(X^1(s, x)) - f(X^2(s, x))\|_{L^1([0,1])}, \end{aligned}$$

where $\|\cdot\|_{L^\infty([0,1])}$ and $\|\cdot\|_{L^1([0,1])}$ are the classical norms on the space $[0, 1]$. The latter term converges to zero since $Y^N(s)$ converges uniformly to $Y(s)$ on $[0, 1]$. Since $f(x) - \lambda x$ is nonincreasing,

$$\begin{aligned} (\langle Y(s), f(X^1(s, x)) - f(X^2(s, x)) \rangle) &= (\langle Y(s), f(X^1(s, x)) - f(X^2(s, x)) - \lambda Y(s) \rangle) \\ &\quad + (\langle Y(s), \lambda Y(s) \rangle) \\ &\leq \lambda |Y(s)|_0^2. \end{aligned}$$

Taking the limit in (2.2) as N grows to infinity, we obtain:

$$|Y(t)|_{-1}^2 - |Y(\delta)|_{-1}^2 \leq \langle Y, \zeta_- - \zeta_+ \rangle_{O_{\delta,t}} + \lambda \int_{\delta}^t |Y(s)|_0^2 ds.$$

We now write

$$\begin{aligned} & \langle Y, \zeta_- - \zeta_+ \rangle_{O_{\delta,t}} \\ &= \langle 1 + X^1, \eta_-^1 \rangle_{O_{\delta,t}} - \langle 1 + X^2, \eta_-^1 \rangle_{O_{\delta,t}} - \langle 1 + X^1, \eta_-^2 \rangle_{O_{\delta,t}} + \langle 1 + X^2, \eta_-^2 \rangle_{O_{\delta,t}} \\ &+ \langle 1 - X^1, \eta_+^1 \rangle_{O_{\delta,t}} - \langle 1 - X^2, \eta_+^1 \rangle_{O_{\delta,t}} - \langle 1 - X^1, \eta_+^2 \rangle_{O_{\delta,t}} + \langle 1 - X^2, \eta_+^2 \rangle_{O_{\delta,t}} \\ &\leq 0 \end{aligned}$$

by the contact condition and the positivity of the measures. It follows:

$$|Y(t)|_{-1}^2 - |Y(\delta)|_{-1}^2 \leq \lambda \int_{\delta}^t |Y(s)|_0^2 ds.$$

By Gronwall Lemma, and letting $\delta \rightarrow 0$, we have $|Y(t)|_{-1} = 0$ for all $t \geq 0$. Since $\bar{Y}(t) = 0$, we deduce $X^1(t, x) = X^2(t, x)$ for all $t \geq 0$. Moreover, with the definition of a weak solution, we see that :

$$\text{for all } h \in D(A^2), \quad \langle Ah, \zeta_+ - \zeta_- \rangle_{O_{\delta,t}} = 0.$$

By density, we obtain that $\zeta := \zeta_- - \zeta_+$ and $V := (1 - X^1)(1 + X^1) = (1 - X^2)(1 + X^2)$ satisfy the hypothesis of Lemma 2.1, and therefore $\zeta = \zeta_- - \zeta_+$ is the null measure. And since ζ_- and ζ_+ have disjoint supports, then ζ_- and ζ_+ are the null measure, i.e. $\eta_-^1 = \eta_-^2$ and $\eta_+^1 = \eta_+^2$. \square

2.2 Convergence of invariants measures

We know (see [18]) that μ_c is the law of $Y^c = \mathbf{B} - \bar{\mathbf{B}} + c$, where B is brownian motion. Then for $0 \leq c < 1$, we remark the following inclusion :

$$\{\mathbf{B}_\theta \in \left[\frac{c-1}{2}, \frac{1-c}{2} \right], \text{ for all } \theta \in [0, 1]\} \subset \{Y^c \in K\},$$

and we have a similar result for $-1 < c \leq 0$. Therefore $\mu_c(K) > 0$ with $-1 < c < 1$. Let us define U the potential associated to the function f :

$$U(x) = \begin{cases} \int_0^1 F(x(\theta)) d\theta & \text{if } x \in K, \\ +\infty & \text{else.} \end{cases}$$

We have the following result :

Proposition 2.2 For $-1 < c < 1$,

$$\nu_c^n \rightarrow \nu_c := \frac{1}{Z_c} \exp^{-U(x)} \mathbb{1}_{x \in K} \mu_c(dx), \text{ when } n \rightarrow +\infty,$$

where Z_c is a normalization constant.

Proof : Let $\psi \in \mathcal{C}_b(L^2, \mathbb{R})$. We want to prove that

$$\int_H \psi(x) \exp(-U^n(x)) \mu_c(dx) \xrightarrow{n \rightarrow +\infty} \int_H \psi(x) \exp(-U(x)) \mathbb{1}_{x \in K} \mu_c(dx). \quad (2.3)$$

We first prove,

$$\exp(-U^n(x)) \xrightarrow{n \rightarrow +\infty} \exp(-U(x)) \mathbb{1}_{x \in K}, \mu_c \text{ a.s.} \quad (2.4)$$

Since $\mu_c(C([0, 1])) = 1$, we can restrict our attention to $x \in C([0, 1])$. Then if $x \notin K$ there exists $\delta_x > 0$ such that $m(\{\theta \in [0, 1] : x(\theta) \leq -1 - \delta_x\}) > 0$ or $m(\{\theta \in [0, 1] : x(\theta) \geq 1 + \delta_x\}) > 0$, m being the Lebesgue measure. Suppose $m(\{\theta \in [0, 1] / x(\theta) \leq -1 - \delta_x\}) > 0$, then we have since $\tilde{F}^n(x) = F^n(x) + \frac{\lambda}{2}x^2$ is positive and non increasing on $(-\infty, -1)$

$$\begin{aligned} 0 \leq \exp(-U^n(x)) &\leq \exp\left(-\int_0^1 \tilde{F}^n(x(\theta)) \mathbb{1}_{\{x \leq -1 - \delta_x\}} - \frac{\lambda}{2}x(\theta)^2 d\theta\right) \\ &\leq \exp\left(-\int_0^1 \tilde{F}^n(-1 - \delta_x) \mathbb{1}_{\{x \leq -1 - \delta_x\}} - \frac{\lambda}{2}x(\theta)^2 d\theta\right) \\ &\leq \exp\left(-\tilde{F}^n(-1 - \delta_x)m(\{x \leq -1 - \delta_x\}) + \int_0^1 \frac{\lambda}{2}x(\theta)^2 d\theta\right). \end{aligned}$$

And this latter term converges to zero as n grows to infinity.

Now for $x \in K$, $F^n(x(\theta))$ converges to $F(x(\theta))$ almost everywhere as n grows to infinity. Moreover $-\frac{\lambda}{2}x(\theta)^2 \leq F^n(x(\theta)) \leq \ln 2$, and by the dominated convergence Theorem, we deduce (2.4). Finally, (2.3) follows again by dominated convergence Theorem.

2.3 Existence of stationary solutions

In this section, we prove the existence of stationary solutions of equation (1.2) and that they are limits of stationary solutions of (1.3), in some suitable sense. Fix $-1 < c < 1$ and consider the unique (in law) stationary solution of (1.3) denote \hat{X}_c^n in H_c . We are going to prove that the laws of \hat{X}_c^n weakly converge as n grows to infinity to a stationary strong solution of (0.3).

Proposition 2.3 Let $-1 < c < 1$ and $T > 0$, \hat{X}_c^n converges in probability as n grows to infinity to a process \hat{X}_c in $\mathcal{C}(O_T)$. Moreover $f(\hat{X}_c) \in L^1(O_T)$ almost surely, and setting

$$d\eta_+^n = -f^n(\hat{X}_c^n(t, \theta)) \mathbb{1}_{\hat{X}_c^n(t, \theta) > 0} dt d\theta + f(\hat{X}_c(t, \theta)) \mathbb{1}_{0 < \hat{X}_c(t, \theta) \leq 1} dt d\theta,$$

and

$$d\eta_-^n = f^n(\hat{X}_c^n(t, \theta)) \mathbb{1}_{\hat{X}_c^n(t, \theta) \leq 0} dt d\theta - f(\hat{X}_c(t, \theta)) \mathbb{1}_{-1 \leq \hat{X}_c(t, \theta) \leq 0} dt d\theta,$$

then (η_+^n, η_-^n) converges in probability to (η_+, η_-) such that $(\hat{X}_c, \eta_+, \eta_-, W)$ is a stationary strong solution of (0.3).

Proof : Proceeding exactly as in [18] (see Lemma 5.2), we prove that the laws of $(\hat{X}_c^n, W^n)_{n \in \mathbb{N}}$ are tight in $\mathcal{C}(O_T) \times \mathcal{C}([0, T]; V_\gamma)$, $\gamma < -1/2$. We have set $W^n = W$, $n \in \mathbb{N}$. We therefore can extract convergent subsequences. Let $(\hat{X}_c^{n_k}, W^{n_k})_{k \in \mathbb{N}}$ be such a subsequence. Using Skohorod theorem, one may find a probability space and a sequence of random variables $(\tilde{X}_c^k, \mathcal{W}^k)_{k \in \mathbb{N}}$ on this probability space with the same laws as $(\hat{X}_c^{n_k}, W^{n_k})_{k \in \mathbb{N}}$ which converge almost surely.

Below, we show in Step 1 that its limit \tilde{X}_c satisfies $f(\tilde{X}_c) \in L^1(O_T)$ almost surely. Then in Step 2, we prove that the measures $\tilde{\eta}_\pm^k$, defined as above with $\hat{X}_c^{n_k}$ replaced by \tilde{X}_c^k , converges to two positive measures $\tilde{\eta}_\pm$ and that $(\tilde{X}_c, \tilde{\eta}_+, \tilde{\eta}_-)$ is a weak solution in the probabilistic sense. It then remains to use pathwise uniqueness to conclude in Step 3. In this proof, we only treat the case $\lambda = 0$. This assumption is not essential at all but lightens the computations. For $\lambda \neq 0$, an extra term has to be taken into account. It is very easy to deal with.

Step 1.

Applying Ito formula to $|Q_N \hat{X}_c^n(t)|_{-1}^2$, we obtain

$$\begin{aligned} & |Q_N \hat{X}_c^n(T)|_{-1}^2 - |Q_N \hat{X}_c^n(0)|_{-1}^2 + \int_0^T |Q_N \hat{X}_c^n(t)|_1^2 dt - 2 \int_{O_T} f_n(\hat{X}_c^n) (Q_N \hat{X}_c^n - c) \, dsd\theta \\ &= 2 \int_0^T (Q_N \hat{X}_c^n, BdW(s)) + T \operatorname{Tr}(Q_N B) \end{aligned}$$

Note that

$$\mathbb{E} \left(\left(\int_0^T (Q_N \hat{X}_c^n, BdW(s)) \right)^2 \right) = \mathbb{E} \int_0^T |Q_N \hat{X}_c^n|_{-1}^2 ds = T \int_H |Q_N x|_{-1}^2 \nu_c^n(dx) \leq CT$$

We set

$$\begin{aligned} \varphi_n^N &= |Q_N \hat{X}_c^n(T)|_{-1}^2 - |Q_N \hat{X}_c^n(0)|_{-1}^2 + \int_0^T |Q_N \hat{X}_c^n(t)|_1^2 dt \\ &\quad - 2 \int_{O_T} f_n(\hat{X}_c^n) (Q_N \hat{X}_c^n - c) \, dsd\theta - T \operatorname{Tr}(Q_N B) \end{aligned}$$

and deduce

$$\mathbb{P}(|\varphi_n^N| \geq M) \leq \frac{CT}{M^2}.$$

Thus, for all $N \in \mathbb{N}$, the laws of $(\varphi_n^N)_{n \in \mathbb{N}}$ are tight. Therefore the laws of $(\hat{X}_c^n, W^n, (\varphi_n^N)_{n \in \mathbb{N}})_{n \in \mathbb{N}}$ are tight and using Skohorod theorem on this sequence, we can assume that $\tilde{X}_c^k, \mathcal{W}^k$ and, for $N \in \mathbb{N}$, $\tilde{\varphi}_k^N$ converge almost surely. We have defined $\tilde{\varphi}_k^N$ as above with \tilde{X}_c^k instead of \hat{X}_c^n . In particular, $\tilde{\varphi}_k^N$ is bounded almost surely:

$$\begin{aligned} & |Q_N \tilde{X}_c^k(T)|_{-1}^2 - |Q_N \tilde{X}_c^k(0)|_{-1}^2 + 2 \int_0^T |Q_N \tilde{X}_c^k(t)|_1^2 dt \\ &\quad - 2 \int_{O_T} f_k(\tilde{X}_c^k) (Q_N \tilde{X}_c^k - c) \, dsd\theta - T \operatorname{Tr}(Q_N B) \\ &\leq C(N, T, c) \end{aligned}$$

where $C(N, T, c)$ is random. The first three terms are clearly also bounded almost surely. This uses the fact that Q_N is a bounded operator from H to V_1 . Since Q_N has finite dimensional range, we obtain

$$- \int_{O_T} f_{nk}(\tilde{X}_c^k) (Q_N \tilde{X}_c^k - c) \, dsd\theta \leq C(N, T, c) \tag{2.5}$$

for a different random constant $C(N, T, c)$.

Let us choose $\epsilon_0 = \min \left\{ \frac{1-c}{4}, \frac{1+c}{4} \right\}$ and take $N \in \mathbb{N}$ such that

$$|Q_N \tilde{X}_c - \tilde{X}_c|_{C(O_T)} \leq \frac{1}{2} \epsilon_0$$

and K_0 such that for $k \geq K_0$

$$|\tilde{X}_c^k - \tilde{X}_c|_{C(O_T)} \leq \frac{1}{4} \epsilon_0.$$

Then, for all $k \geq K_0$,

$$|Q_N \tilde{X}_c^k - \tilde{X}_c^k|_{C(O_T)} \leq \epsilon_0.$$

Moreover, if $\tilde{X}_c^k \geq \frac{1+c}{2}$ then $f_{n_k}(\tilde{X}_c^k) \leq 0$ and

$$Q_N \tilde{X}_c^k - c \geq -\epsilon_0 + \frac{1+c}{2} - c \geq \frac{1-c}{4} \geq \epsilon_0.$$

Similarly, if $\tilde{X}_c^k \leq \frac{-1+c}{2}$ then $f_{n_k}(\tilde{X}_c^k) \geq 0$ and

$$Q_N \tilde{X}_c^k - c \leq -\epsilon_0.$$

Finally, noticing that f_n is uniformly bounded by a constant $K(c)$ on $[\frac{-1+c}{2}, \frac{1+c}{2}]$, we deduce

$$\begin{aligned} \int_{O_T} |f_{n_k}(\tilde{X}_c^k)| \, dsd\theta &\leq -\frac{1}{\epsilon_0} \int_{\tilde{X}_c^k \geq \frac{1+c}{2}} f_{n_k}(\tilde{X}_c^k) (Q_N \tilde{X}_c^k - c) \, dsd\theta \\ &\quad -\frac{1}{\epsilon_0} \int_{\tilde{X}_c^k \leq \frac{-1+c}{2}} f_{n_k}(\tilde{X}_c^k) (Q_N \tilde{X}_c^k - c) \, dsd\theta + K(c) \\ &\leq -\frac{1}{\epsilon_0} \int_{O_T} f_{n_k}(\tilde{X}_c^k) (Q_N \tilde{X}_c^k - c) \, dsd\theta \\ &\quad + \frac{1}{8} (\max\{1-c, 1+c\})^2 K(c) + K(c). \end{aligned}$$

Thanks to (2.5), we obtain

$$\int_{O_T} |f_{n_k}(\tilde{X}_c^k)| \, dsd\theta \leq C(N, T, c), \quad (2.6)$$

where the value of the random constant $C(N, T, c)$ has again changed. It easily deduced from this uniform bound that $|\tilde{X}_c| \leq 1$ almost everywhere with respect to t and ω and by Fatou Lemma that $f(\tilde{X}_c) \in L^1(O_T)$ almost surely. □

Step 2.

Let now ξ^k be the following measure on O_T :

$$d\xi^k := -f^{n_k}(\tilde{X}_c^k(t, \theta)) \, dt d\theta.$$

and ξ_+^k and ξ_-^k the positive and negative parts:

$$d\xi_+^k := -f^{n_k}(\tilde{X}_c^k(t, \theta)) \mathbb{1}_{\tilde{X}_c^k > 0} \, dt d\theta, \quad d\xi_-^k := f^{n_k}(\tilde{X}_c^k(t, \theta)) \mathbb{1}_{\tilde{X}_c^k \leq 0} \, dt d\theta.$$

By step 1, $f(\tilde{X}_c) \in L^1(O_T)$ and we can define the following measure:

$$d\lambda := -f((\tilde{X}_c(t, \theta)) \mathbb{1}_{-1 \leq \tilde{X}_c \leq 1} \, dt d\theta,$$

and the positive and negative parts:

$$d\lambda_+ := -f((\tilde{X}_c(t, \theta)) \mathbb{1}_{0 < \tilde{X}_c \leq 1} \, dt d\theta, \quad d\lambda_- := f((\tilde{X}_c(t, \theta)) \mathbb{1}_{-1 \leq \tilde{X}_c < 0} \, dt d\theta.$$

By (2.6), $f^{n_k}(\tilde{X}_c^k) - f(\tilde{X}_c)$ is bounded in $L^1(O_T)$. We deduce that ξ^k has a subsequence ξ^{k_ℓ} which converges to a measure ζ . Note that this subsequence may depend on the random parameter ω . We set $\tilde{\eta} = \zeta - \lambda$.

Thanks to Fatou Lemma we have the following inequality for all $h \in \mathcal{C}(O_T)$ nonnegative:

$$\begin{aligned} \int_{O_T} h(s, \theta) [-f(\tilde{X}_c(s, \theta)) \mathbb{1}_{0 < \tilde{X}_c \leq 1}] ds d\theta &= \int_{O_T} \liminf_{\ell \rightarrow +\infty} [-h(s, \theta) f^{n_{k_\ell}}(\tilde{X}_c^{k_\ell}(s, \theta)) \mathbb{1}_{0 < \tilde{X}_c^{k_\ell} \leq 1}] ds d\theta \\ &\leq \liminf_{\ell \rightarrow +\infty} \int_{O_T} [-h(s, \theta) f^{n_{k_\ell}}(\tilde{X}_c^{k_\ell}(s, \theta)) \mathbb{1}_{0 < \tilde{X}_c^{k_\ell} \leq 1}] ds d\theta. \end{aligned}$$

Therefore $\tilde{\eta}_+^{k_\ell} = \xi_+^{k_\ell} - \lambda_+$ converges to a positive measure. Similarly, $\tilde{\eta}_-^{k_\ell} = \xi_-^{k_\ell} - \lambda_-$ converges to a positive measure. It follows:

$$\xi_+^{k_\ell} - \lambda_+ \rightharpoonup \tilde{\eta}_+ \quad \text{and} \quad \xi_-^{k_\ell} - \lambda_- \rightharpoonup \tilde{\eta}_-,$$

where $\tilde{\eta}_+$ and $\tilde{\eta}_-$ are the positive and negative parts of $\tilde{\eta}$.

Let us now show that the contact conditions holds for $(1 - \tilde{X}_c, \tilde{\eta}_+)$ and $(1 + \tilde{X}_c, \tilde{\eta}_-)$. Let us define the following measures for $\varepsilon > 0$ and $k \in \mathbb{N}$.

$$\begin{aligned} d\xi_{+, \varepsilon}^k &:= -f^{n_k}(\tilde{X}_c^k(t, \theta)) \mathbb{1}_{1-\varepsilon \leq \tilde{X}_c^k} dt d\theta, & d\tau_{+, \varepsilon}^k &:= -f^{n_k}(\tilde{X}_c^k(t, \theta)) \mathbb{1}_{0 < \tilde{X}_c^k < 1-\varepsilon} dt d\theta, \\ d\lambda_{+, \varepsilon} &:= -f(\tilde{X}_c(t, \theta)) \mathbb{1}_{1-\varepsilon \leq \tilde{X}_c} dt d\theta, & d\tau_{+, \varepsilon} &:= -f(\tilde{X}_c(t, \theta)) \mathbb{1}_{0 < \tilde{X}_c < 1-\varepsilon} dt d\theta. \end{aligned}$$

Clearly $\tau_{+, \varepsilon}^k$ converges to $\tau_{+, \varepsilon}$, it follows

$$\begin{aligned} \limsup_{\ell \rightarrow +\infty} \langle 1 - \tilde{X}_c^{k_\ell}, \xi_+^{k_\ell} - \lambda_+ \rangle_{O_T} &= \limsup_{\ell \rightarrow +\infty} \left(\langle 1 - \tilde{X}_c^{k_\ell}, \xi_{+, \varepsilon}^{k_\ell} \rangle_{O_T} - \langle 1 - \tilde{X}_c^{k_\ell}, \lambda_{+, \varepsilon} \rangle_{O_T} \right. \\ &\quad \left. + \langle 1 - \tilde{X}_c^{k_\ell}, \tau_{+, \varepsilon}^{k_\ell} \rangle_{O_T} - \langle 1 - \tilde{X}_c^{k_\ell}, \tau_{+, \varepsilon} \rangle_{O_T} \right) \\ &= \limsup_{\ell \rightarrow +\infty} \left(\int_{O_T} (\tilde{X}_c^{k_\ell} - 1) f^{n_{k_\ell}}(\tilde{X}_c^{k_\ell}) \mathbb{1}_{1-\varepsilon \leq \tilde{X}_c^{k_\ell}} dt d\theta \right. \\ &\quad \left. + \int_{O_T} (1 - \tilde{X}_c^{k_\ell}) f(\tilde{X}_c) \mathbb{1}_{1-\varepsilon \leq \tilde{X}_c} dt d\theta \right) \\ &\leq \limsup_{\ell \rightarrow +\infty} \left(\int_{O_T} (\tilde{X}_c^{k_\ell} - 1) f^{n_{k_\ell}}(\tilde{X}_c^{k_\ell}) \mathbb{1}_{1-\varepsilon \leq \tilde{X}_c^{k_\ell} \leq 1} dt d\theta \right) \\ &\quad + \limsup_{\ell \rightarrow +\infty} \left(\int_{O_T} (1 - \tilde{X}_c^{k_\ell})^- f(\tilde{X}_c) \mathbb{1}_{1-\varepsilon \leq \tilde{X}_c} dt d\theta \right) \end{aligned}$$

Since $(1 - \tilde{X}_c^{k_\ell})^-$ converges uniformly to zero, we deduce:

$$\begin{aligned} \limsup_{\ell \rightarrow +\infty} \langle 1 - \tilde{X}_c^{k_\ell}, \xi_+^{k_\ell} - \lambda_+ \rangle_{O_T} &\leq T \sup_{x \in [1-\varepsilon, 1]} |(x-1)f(x)| \\ &\leq -T\varepsilon \ln \left(\frac{\varepsilon}{2-\varepsilon} \right). \end{aligned}$$

Letting $\varepsilon \rightarrow 0$, we obtain the first contact condition since the left hand side clearly converges to $\langle 1 - \tilde{X}_c, \tilde{\eta}_+ \rangle$. The second is obtained similarly.

We now prove that $\xi^k - \lambda$ does not have more than one limit point so that in fact the whole sequence converge to $\tilde{\eta}$. Let $\tilde{\eta}_i$, $i = 1, 2$ be two limit points.

For all $h \in D(A^2)$ and for all $0 \leq t \leq T$:

$$\begin{aligned} \langle Ah, \xi^k - \lambda \rangle_{O_t} &= \langle \tilde{X}_c^k(t, \cdot), h \rangle - \langle x, h \rangle + \int_{O_t} \tilde{X}_c^k(s, \theta) A^2 h(\theta) ds d\theta \\ &\quad + \int_0^t \langle Bh, d\mathcal{W}^k \rangle + \int_{O_t} f(\tilde{X}_c(s, \theta)) Ah(\theta) ds d\theta. \end{aligned}$$

We deduce

$$\begin{aligned}\langle Ah, \tilde{\eta}_i \rangle_{O_t} &= -\langle \tilde{X}_c(t, \cdot), h \rangle + \langle x, h \rangle - \int_{O_t} \tilde{X}_c(s, \theta) A^2 h(\theta) ds d\theta \\ &\quad - \int_0^t \langle Bh, d\mathcal{W} \rangle - \int_{O_t} f(\tilde{X}_c(s, \theta)) Ah(\theta) ds d\theta.\end{aligned}$$

And by density

$$\langle h, \tilde{\eta}_1 \rangle_{O_t} = \langle h, \tilde{\eta}_2 \rangle_{O_t}$$

for any $h \in C([0, 1])$ such that $\bar{h} = 0$. Since by the contact condition

$$\langle (1 - \tilde{X}_c)(1 + \tilde{X}_c), \tilde{\eta}_1 \rangle_{O_t} = \langle (1 - \tilde{X}_c)(1 + \tilde{X}_c), \tilde{\eta}_2 \rangle_{O_t}.$$

We deduce from Lemma 2.1 that $\tilde{\eta}_1 = \tilde{\eta}_2$. □

Step 3.

We use a result from [25] that allows to get the convergence of the approximated solutions in probability in any space in which these approximated solutions are tight.

Lemma 2.2 *Let $\{Z_n\}_{n \geq 1}$ be a sequence of random elements on a Polish space E endowed by its borel σ -algebra. Then $\{Z_n\}_{n \geq 1}$ converges in probability to an E -valued random element if and any if from every pair of subsequences $\{(Z_{n_k^1}, Z_{n_k^2})_{k \geq 1}$, one can extract a subsequence which converges weakly to a random element supported on the diagonal $\{(x, y) \in E \times E, x = y\}$.*

Assume $(n_k^1)_{k \in \mathbb{N}}$ and $(n_k^2)_{k \in \mathbb{N}}$ are two arbitrary subsequences. Clearly, the process $(\hat{X}_c^{n_k^1}, \hat{X}_c^{n_k^2}, W^k)$ is tight in a suitable space. By Skorohod's theorem, we can find a probability space and a sequence of processes $(\tilde{X}_c^{1,k}, \tilde{X}_c^{2,k}, \mathcal{W}^k)$ such that $(\tilde{X}_c^{1,k}, \tilde{X}_c^{2,k}, \mathcal{W}^k) \rightarrow (\tilde{X}_c^1, \tilde{X}_c^2, \mathcal{W})$ almost surely and $(\tilde{X}_c^{1,k}, \tilde{X}_c^{2,k}, \mathcal{W}^k)$ has the same distribution as $(\hat{X}_c^{n_k^1}, \hat{X}_c^{n_k^2}, W^k)$ for all $k \in \mathbb{N}$. In the Skorohod's space, the approximated measures respectively converge to two contact measures $\tilde{\eta}_1$ and $\tilde{\eta}_2$. By the second step, $(\tilde{X}_c^1, \tilde{\eta}_1, \mathcal{W})$ and $(\tilde{X}_c^2, \tilde{\eta}_2, \mathcal{W})$ are both weak solutions of (0.3). By uniqueness, necessarily $\tilde{X}_c^1 = \tilde{X}_c^2$ and $\tilde{\eta}_1 = \tilde{\eta}_2$. Therefore the subsequence $(\hat{X}_c^{n_k^1}, \hat{X}_c^{n_k^2})_{k \in \mathbb{N}}$ converges in distribution to a process supported on the diagonal. We use Lemma 2.2 to prove that the sequence (\hat{X}_c^n) converges in probability to a process \hat{X}_c . Clearly \hat{X}_c is stationary. Reproducing the argument of Step 1 and Step 2, we prove that it is a strong solution of (0.3) and the convergence of the contact measures. □

2.4 Convergence of the semigroup

First we state the following result which is a corollary of Proposition 2.3.

Corollary 2.1 *Let $c > 0$.*

- i) *There exists a continuous process $(X(t, x), t \geq 0, x \in K \cap H_c)$ with $X(0, x) = x$ and a set K_0 dense in $K \cap H_c$, such that for all $x \in K_0$ there exists a unique strong solution of equation (0.3) given by $((X(t, x))_{t \geq 0}, \eta_+^x, \eta_-^x, W)$.*
- ii) *The law of $(X(t, x)_{t \geq 0}, \eta_+^x, \eta_-^x)$ is a regular conditional distribution of the law of $(\hat{X}_c, \eta_+, \eta_-)$ given $\hat{X}_c(0) = x \in K \cap H_c$.*

Proof : By Proposition 2.3, we have a stationary strong solution \hat{X}_c in H_c , such that W and $\hat{X}_c(0)$ are independent. Conditioning $(\hat{X}_c, \eta_+, \eta_-)$ on the value of $\hat{X}_c(0) = x$, with $c = \bar{x}$, we obtain for ν_c -almost every x a strong solution that we denote $(X(t, x), \eta_+^x, \eta_-^x)$ for all $t \geq 0$ and for all $x \in K \cap H_c$. This process is the desired process. Indeed, since the support of ν_c is $K \cap H_c$, we have a strong solution for a dense set K_0 in $K \cap H_c$.

Notice that all processes $(X(t, x))_{t \geq 0}$ with $x \in K_0$ are driven by the same noise W and are continuous with values in H . Moreover, we have the following obvious identity:

$$|X^n(t, x) - X^n(t, y)|_{-1} \leq e^{\lambda t} |x - y|_{-1}, \quad x, y \in L_c^2, \quad t \geq 0,$$

and by density we obtain a continuous process $(X(t, x))_{t \geq 0}$ in H_c for all $x \in K \cap H_c$. □

Proposition 2.4 *Let $c > 0$, for all $\phi \in \mathcal{C}_b(H)$ and $x \in K \cap H_c$:*

$$\lim_{n \rightarrow +\infty} P_t^{n,c} \phi(x) = \mathbb{E}[\phi(X(t, x))] =: P_t^c \phi(x). \quad (2.7)$$

Moreover the Markov process $(X(t, x), t \geq 0, x \in K \cap H_c)$ is strong Feller and its transition semigroup P^c is such that:

$$|P_t^c \phi(x) - P_t^c \phi(y)| \leq \frac{2e^{\lambda^2 t/4}}{\lambda \sqrt{t}} |x - y|_{-1}, \quad \text{for all } x, y \in K \cap H_c, \text{ for all } t > 0. \quad (2.8)$$

Proof : Since $(\nu_c^n)_{n \geq 1}$ is tight in H_c , then there exists an increasing sequence of compact sets $(J^p)_{p \in \mathbb{N}}$ in H such that:

$$\lim_{p \rightarrow +\infty} \sup_{n \geq 1} \nu_c^n(H \setminus J^p) = 0.$$

Set $J := \bigcup_{p \in \mathbb{N}} J^p \cap K$. Since the support of ν_c is in $K \cap H_c$ and $\nu_c(J) = 1$, then J is dense in $K \cap H_c$.

Fix $t > 0$, by (1.4), for any $\phi \in \mathcal{C}_b(H)$:

$$\sup_{n \in \mathbb{N}} (\|P_t^{n,c} \phi\|_\infty + [P_t^{n,c} \phi]_{Lip(H_c)}) < +\infty.$$

Let $(n_j)_{j \in \mathbb{N}}$ be any sequence in \mathbb{N} . With a diagonal procedure, by Arzelà-Ascoli Theorem, there exists $(n_{j_l})_{l \in \mathbb{N}}$ a subsequence and a function $\Theta_t : J \rightarrow \mathbb{R}$ such that:

$$\lim_{l \rightarrow +\infty} \sup_{x \in J^p} |P_t^{n_{j_l}, c} \phi(x) - \Theta_t(x)| = 0, \quad \text{for all } p \in \mathbb{N}.$$

By density, Θ_t can be extended uniquely to a bounded Lipschitz function $\tilde{\Theta}_t$ on $K \cap H_c$ such that

$$\tilde{\Theta}_t(x) = \lim_{l \rightarrow +\infty} P_t^{n_{j_l}, c} \phi(x), \quad \text{for all } x \in K \cap H_c.$$

Note that the subsequence depends on t . Therefore, we have to prove that the limit defines a semigroup and does not depend on the chosen subsequence.

By Proposition 2.3, we have for all $\phi, \psi \in \mathcal{C}_b(H)$:

$$\begin{aligned} \mathbb{E} \left[\psi \left(\hat{X}_c(0) \right) \phi \left(\hat{X}_c(t) \right) \right] &= \lim_{l \rightarrow +\infty} \mathbb{E} \left[\psi \left(\hat{X}_c^{n_{j_l}}(0) \right) \phi \left(\hat{X}_c^{n_{j_l}}(t) \right) \right] \\ &= \lim_{l \rightarrow +\infty} \int_H \psi(y) \mathbb{E} \left[\phi \left(\hat{X}_c^{n_{j_l}}(t) \right) \middle| \hat{X}_c^{n_{j_l}}(0) = y \right] \nu_c^{n_{j_l}}(dy) \\ &= \lim_{l \rightarrow +\infty} \int_H \psi(y) P_t^{n_{j_l}, c} \phi(y) \nu_c^{n_{j_l}}(dy) \\ &= \int_H \psi(y) \tilde{\Theta}_t(y) \nu_c(dy). \end{aligned}$$

Thus, by Corollary 2.1, we have the following equality:

$$\mathbb{E}[\phi(X(t, x))] = \tilde{\Theta}_t(x), \quad \text{for } \nu_c\text{-almost every } x. \quad (2.9)$$

Since $\mathbb{E}[\phi(X(t, \cdot))]$ and $\tilde{\Theta}_t$ are continuous on $K \cap H_c$, and $\nu_c(K \cap H_c) = 1$, the equality (2.9) is true for all $x \in K \cap H_c$. Moreover the limit does not depend on the chosen subsequence, and we obtain (2.7). Letting $n \rightarrow \infty$ in (1.4), we deduce (2.8). □

2.5 End of the proof of Theorem 1.1

We have proved that there exists a continuous process X which is a strong solution of equation (0.3) for an x in a dense space. In this section, we prove existence for an initial condition in $K \cap H_c$ with $c > 0$.

By Corollary 2.1 we have a process $(X(t, x), t \geq 0, x \in K \cap H_c)$, such that for all x in a set K_0 dense in $K \cap H_c$ we have a strong solution $\left((X(t, x))_{t \geq 0}, \tilde{\eta}_+^x, \tilde{\eta}_-^x, W \right)$ of (0.3) with initial condition x . By Proposition 2.3, the Markov process X has transition semigroup P^c on H_c .

The strong Feller property of P^c implies that for all $x \in K \cap H_c$ and $s > 0$ the law of $X(s, x)$ is absolutely continuous with respect to the invariant measure ν_c . Indeed, if $\nu_c(\Gamma) = 0$, then $\nu_c(P_s^c(\mathbb{1}_\Gamma)) = \nu_c(\Gamma) = 0$. So $P_s^c(\mathbb{1}_\Gamma)(x) = 0$ for ν_c -almost every x and by continuity for all $x \in K \cap H_c$.

Therefore almost surely $X(s, x) \in K_0$ for all $s > 0$ and $x \in K \cap H_c$. Fix $s > 0$, denote for all $\theta \in [0, 1]$:

$$\tilde{X} := t \mapsto X(t + s, x), \tilde{W}(\cdot, \theta) := t \mapsto W(t + s, \theta) - W(s, \theta),$$

and the measures $\tilde{\eta}_\pm^x$ such that for all $T > 0$, and for all $h \in \mathcal{C}(O_T)$:

$$\langle h, \tilde{\eta}_\pm^x \rangle_{O_T} := \int_{O_s^{T+s}} h(t - s, \theta) \eta_\pm^x(dt, d\theta).$$

So we have a process $\tilde{X} \in \mathcal{C}([0, T]; H) \cap \mathcal{C}(O_T)$ and two measures $\tilde{\eta}_+^x$ and $\tilde{\eta}_-^x$ on O_T which is finite on $[\delta, T] \times [0, 1]$ for all $\delta \geq 0$, such that $\left((\tilde{X}(t, x))_{t \geq 0}, \tilde{\eta}_+^x, \tilde{\eta}_-^x, \tilde{W} \right)$ is a strong solution of (0.3) with initial condition $X(s, x)$. By continuity $X(s, x) \rightarrow x$ in H as $s \rightarrow 0$, so $\left((X(t, x))_{t \geq 0}, \eta_+^x, \eta_-^x, W \right)$ is a strong solution of (0.3) with initial condition x in the sense of the definition 1.1.

3 Ergodicity and mixing

When λ is small, it can be easily shown that ν_c is the unique invariant measure and is ergodic. We now prove that this is in fact true for any λ . Note that since $(P_t^c)_{t \geq 0}$ is Strong Feller, the results follows from Doob theorem if we prove that $(P_t^c)_{t \geq 0}$ is irreducible (see for instance [15]). For additive noise driven SPDEs, this is often proved by a control argument and continuity with respect to the noise. This latter property is not completely trivial in our situation but we are able to adapt the argument.

Proposition 3.1 *For any $c \in (-1, 1)$, the semigroup $(P_t^c)_{t \geq 0}$ is irreducible.*

Proof :

Let $x, y \in C^\infty([0, 1])$ be such that $|x|_{L^\infty(0,1)} \leq 1 - \delta$ and $|y|_{L^\infty(0,1)} \leq 1 - \delta$ for some $\delta > 0$ and $\bar{x} = \bar{y} = c$. We set

$$u(t) = \frac{t}{T}y + \left(1 - \frac{t}{T}\right)x$$

and define g_0 by

$$g_0(\theta, t) = \int_0^\theta \left(\frac{1}{T}(y - x) + \frac{1}{2}A(Au + f(u)) \right) (\vartheta, t) d\vartheta$$

Then g_0 is in $C^\infty([0, T] \times [0, 1])$, $g_0(t) \in D(B)$, $t \in [0, T]$, and:

$$\frac{d}{dt}u = -\frac{1}{2}A(Au + f(u)) + Bg_0.$$

Moreover

$$\frac{d}{dt}u = -\frac{1}{2}A(Au + f_\delta(u)) + Bg_0 \tag{3.1}$$

where f_δ is any Lipschitz function equal to f on $[-1 + \delta/2, 1 - \delta/2]$.

Let $X^\delta(\cdot, x)$ be the solution of (0.3) with f replaced by f_δ and set $Y^\delta(\cdot, x) = X^\delta(\cdot, x) - Z$, where $Z = Z(\cdot, 0)$ is the solution of the linear equation with 0 as initial data. Then

$$\frac{d}{dt}Y^\delta = -\frac{1}{2}A(AY^\delta + f_\delta(Y^\delta + Z)), \quad Y^\delta(0, x) = x.$$

Let also

$$z_0(t) = \int_0^t e^{-A^2(t-s)/2} B g_0(s) ds.$$

Since the gaussian process Z is almost surely continuous and has a non degenerate covariance, we clearly have

$$\mathbb{P}(|Z - z_0|_{C(O_T)} \leq \varepsilon) > 0$$

for any $\varepsilon > 0$. Let us denote by Y^z the solution of

$$\frac{d}{dt}Y^z = -\frac{1}{2}A(AY^z + f_\delta(Y^z + z)), \quad Y^z(0, x) = x. \quad (3.2)$$

We prove below that the mapping

$$\Phi_\delta : z \mapsto Y^z$$

is continuous from $C(O_T)$ into $C(O_T)$. Since $u = \Phi_\delta(z_0) + z_0$ and $X^\delta = \Phi_\delta(Z) + Z$, we deduce that there exists ε such that

$$\mathbb{P}(|X^\delta - u|_{C(O_T)} \leq \delta/2) \geq \mathbb{P}(|Z - z_0|_{C(O_T)} \leq \varepsilon) > 0$$

Let us now observe that $|X^\delta - u|_{C(O_T)} \leq \delta/2$ implies $|X^\delta|_{C(O_T)} \leq 1 - \delta/2$ so that $f_\delta(X^\delta) = f(X^\delta)$ and X^δ is fact solution of (0.3). By pathwise uniqueness, we deduce that $|X^\delta - u|_{C(O_T)} \leq \delta/2$ implies $X^\delta = X$. It follows

$$\mathbb{P}(|X - u|_{C(O_T)} \leq \delta/2) \geq \mathbb{P}(|X^\delta - u|_{C(O_T)} \leq \delta/2) > 0$$

In particular

$$\mathbb{P}(|X(T, x) - y| \leq \delta/2) > 0.$$

If we assume now that $x, y \in H_c$, we choose $\tilde{x}, \tilde{y} \in C^\infty([0, 1])$ such that

$$|x - \tilde{x}| \leq \delta, \quad |y - \tilde{y}| \leq \delta, \quad |\tilde{x}|_{L^\infty(0,1)} \leq 1 - \delta \quad \text{and} \quad |\tilde{y}|_{L^\infty(0,1)} \leq 1 - \delta,$$

and $\tilde{\tilde{x}} = \tilde{\tilde{y}} = c$. We have

$$|X(T, x) - X(T, \tilde{x})| \leq e^{\lambda T} |x - \tilde{x}|.$$

Therefore

$$\mathbb{P}(|X(T, x) - y| \leq \delta/2 + (1 + e^{\lambda T})\delta) \geq \mathbb{P}(|X(T, \tilde{x}) - \tilde{y}| \leq \delta/2) > 0.$$

This proves the results.

It remains to prove that Φ_δ is continuous. This follows from the mild form of equation (3.2):

$$Y^z(t) = e^{-tA^2/2}x + \int_0^t e^{-(t-s)A^2/2} A f_\delta(Y^z(s) + z(s)) ds.$$

It is classical that, for $t > 0$, $Ae^{-tA^2/2}$ maps $C([0, 1])$ into itself and

$$\left| Ae^{-tA^2/2} \right|_{\mathcal{L}(C([0,1]))} \leq Ct^{-1/2}.$$

This can be seen from the formula

$$Ae^{-tA^2/2}u = - \sum_{i \in \mathbb{N}} \lambda_i e^{-\lambda_i^2 t/2} \langle u, e_i \rangle e_i,$$

where $(e_i)_{i \in \mathbb{N}}$ and $(\lambda_i)_{i \in \mathbb{N}}$ are the eigenvectors and eigenvalues of $-A$. Since $|e_i|_{C([0,1])}$ are equi-bounded, we deduce

$$\begin{aligned} \left| Ae^{-tA^2/2}u \right|_{C([0,1])} &\leq C \left(\sum_{i \in \mathbb{N}} \lambda_i e^{-\lambda_i^2 t/2} \right) |u|_{L^1(0,1)} \\ &\leq C t^{-1/2} |u|_{C([0,1])} \end{aligned}$$

We deduce

$$\begin{aligned} &|Y^{z_1}(t) - Y^{z_2}(t)|_{C([0,1])} \\ &\leq C L_\delta \int_0^t (t-s)^{-1/2} \left(|Y^{z_1}(s) - Y^{z_2}(s)|_{C([0,1])} + |z_1(s) - z_2(s)|_{C([0,1])} \right) ds \end{aligned}$$

where L_δ is the Lipschitz constant of f_δ . Gronwall Lemma implies the result for T sufficiently small. Iterating the argument we obtain the continuity of Φ_δ . \square

Corollary 3.1 *For every $c \in (-1, 1)$, ν_c is the unique invariant measure of the transition semi-group $(P_t^c)_{t \geq 0}$. Moreover it is ergodic.*

Using classical arguments, it is easily seen that, for $\lambda = 0$, ν_c^n satisfies a log-Sobolev inequality and therefore a Poincaré inequality. The constant in these inequality do not depend on n so that we have the same result for ν_c . For $\lambda \neq 0$, we can argue as in [13] and prove that this is still true.

We now want to prove a stronger result : exponential mixing. We use coupling arguments developed by Odasso in [34].

Theorem 3.1 *For every $c \in (-1, 1)$, there exist a small $\beta > 0$ and a constant $C > 0$ such that for all $\varphi \in \mathcal{B}_b(K \cap H_c)$, $t > 0$ and $x \in H_c$*

$$|\mathbb{E}[\varphi(X(t, x))] - \nu_c(\varphi)| \leq C \|\varphi\|_\infty e^{-\beta t}. \quad (3.3)$$

Proof : By (2.8), we know that for any $\varphi \in \mathcal{B}_b(K \cap H_c)$, $T > 0$, $\varepsilon > 0$,

$$|P_T^c \varphi(x) - P_T^c \varphi(y)| \leq \frac{4e^{\lambda^2 T/4}}{\lambda \sqrt{T}} \varepsilon \|\varphi\|_\infty$$

if $x, y \in H_c$, $|x|_{-1} \leq \varepsilon$ and $|y|_{-1} \leq \varepsilon$. By definition of the total variation norm, we deduce

$$\| (P_T^c)^* \delta_x - (P_T^c)^* \delta_y \|_{var} = \sup_{\|\varphi\|_\infty \leq 1} |P_T^c \varphi(x) - P_T^c \varphi(y)| \leq \frac{4e^{\lambda^2 T/4}}{\lambda \sqrt{T}} \varepsilon \quad (3.4)$$

for $T > 0$, $x, y \in H_c$, $|x|_{-1} \leq \varepsilon$ and $|y|_{-1} \leq \varepsilon$. We have denoted by δ_x the Dirac mass at $x \in H_c$ so that $(P_T^c)^* \delta_x$ is the law of $X(T, x)$.

Recall that a coupling of $((P_T^c)^* \delta_x, (P_T^c)^* \delta_y)$ is a couple of random variable (X_1, X_2) such that the law of X_1 is $(P_T^c)^* \delta_x$ and the law of X_2 is $(P_T^c)^* \delta_y$. By standard results on couplings (see for instance [26] section 4, or [30]), we know there exists a maximal coupling of $((P_T^c)^* \delta_x, (P_T^c)^* \delta_y)$. Let us denote by $(Y_1(x, y), Y_2(x, y))$ this maximal coupling, it satisfies

$$\mathbb{P}(Y_1(x, y) \neq Y_2(x, y)) = \| (P_T^c)^* \delta_x - (P_T^c)^* \delta_y \|_{var}. \quad (3.5)$$

Moreover $(Y_1(x, y), Y_2(x, y))$ depends measurably on (x, y) .

By the Strong Feller property, we know that $x \mapsto \mathbb{P}(|X(T, x)|_{-1} \leq \varepsilon)$ is continuous on H_c . Therefore, thanks to Proposition 3.1, for any $x \in K \cap H_c$, there exists a $\eta_x > 0$ and a $\kappa_x > 0$ such that

$$\mathbb{P}(|X(T, y)|_{-1} \leq \varepsilon) > \kappa_x$$

for all $y \in K \cap H_c$ such that $|x - y|_{-1} \leq \eta_x$. By compactness of $K \cap H_c$ in H_c , we deduce that that there exists $\kappa_0 > 0$ such that

$$\mathbb{P}(|X(T, y)|_{-1} \leq \varepsilon) > \kappa_0 \quad (3.6)$$

for all $y \in K \cap H_c$.

Let \tilde{W} a cylindrical Wiener process independent on W and denote by \tilde{X} the associated solution of the stochastic Cahn-Hilliard equation which has the same law as X . For arbitrary $x, y \in K \cap H_c$, we define the coupling $(Z_1(x, y), Z_2(x, y))$ of $((P_T^c)^* \delta_x, (P_T^c)^* \delta_y)$ as follows

$$(Z_1(x, y), Z_2(x, y)) = \begin{cases} (X(T, x), X(T, y)) & \text{if } x = y, \\ (Y_1(x, y), Y_2(x, y)) & \text{if } |x|_{-1} \leq \varepsilon, |y|_{-1} \leq \varepsilon \text{ and } x \neq y, \\ (X(T, x), \tilde{X}(T, y)) & \text{otherwise.} \end{cases}$$

We now construct recursively $(X_1(kT, x, y), X_2(kT, x, y))$ a coupling of $((P_{kT}^c)^* \delta_x, (P_{kT}^c)^* \delta_y)$, the laws of $X(kT, x)$ and $X(kT, y)$. For $k = 0$, we set $(X_1(kT, x, y), X_2(kT, x, y)) = (x, y)$. For $k \geq 0$, we define $(X_1((k+1)T, x, y), X_2((k+1)T, x, y))$ by

$$\begin{aligned} X_1((k+1)T, x, y) &= Z_1(X_1(kT, x, y), X_2(kT, x, y)), \\ X_2((k+1)T, x, y) &= Z_2(X_1(kT, x, y), X_2(kT, x, y)). \end{aligned}$$

Let us define

$$\tau = \inf\{kT : |X_1(kT, x, y)|_{-1} \leq \varepsilon, |X_2(kT, x, y)|_{-1} \leq \varepsilon\}$$

If $|x|_{-1} \leq \varepsilon$ and $|y|_{-1} \leq \varepsilon$, then $\tau = 0$ and $\mathbb{E}(e^{\alpha\tau}) = 1$.

If $\tau \neq 0$ i.e. if $|x|_{-1} \geq \varepsilon$ or $|y|_{-1} \geq \varepsilon$, then by construction of the coupling and (3.6)

$$\mathbb{P}(\tau > T) < 1 - \kappa_0^2.$$

More generally

$$\mathbb{P}(\tau > kT | \tau \geq kT) < 1 - \kappa_0^2.$$

We deduce

$$\mathbb{P}(\tau > kT) < (1 - \kappa_0^2)^k$$

and

$$\mathbb{E}(e^{\alpha\tau}) = \sum_{k \in \mathbb{N}} e^{\alpha kT} \mathbb{P}(\tau = kT) \leq \sum_{k \in \mathbb{N}} e^{\alpha kT} (1 - \kappa_0^2)^{k-1} = M < \infty$$

for α small enough. Similarly, if we define

$$\tau_n = \inf\{kT > \tau_{n-1} : |X_1(kT, x, y)|_{-1} \leq \varepsilon, |X_2(kT, x, y)|_{-1} \leq \varepsilon\},$$

for all $n \geq 2$ and with $\tau_1 := \tau$. We have

$$\mathbb{E}(e^{\alpha(\tau_n - \tau_{n-1})}) \leq M$$

so that

$$\mathbb{E}(e^{\alpha\tau_n}) \leq M^n.$$

Define

$$k_0 = \inf\{n \geq 1 : X_1(\tau_n + T, x, y) = X_2(\tau_n + T, x, y)\}.$$

By (3.4), (3.5), for all $n \geq 1$

$$\mathbb{P}(k_0 = n) \leq \mathbb{P}(k_0 > n - 1) \leq \left(\frac{4e^{\lambda^2 T/4}}{\lambda\sqrt{T}} \varepsilon \right)^{n-1}.$$

We choose ε small enough such that

$$\left(\frac{4e^{\lambda^2 T/4}}{\lambda\sqrt{T}} \varepsilon \right) < 1.$$

Then we write

$$\begin{aligned} \mathbb{E}(e^{\beta\tau_{k_0}}) &= \sum_{n \geq 1} \mathbb{E}(e^{\beta\tau_n} \mathbb{1}_{k_0=n}) \leq \sum_{n \geq 1} (\mathbb{E}(e^{2\beta\tau_n}))^{1/2} (\mathbb{P}(k_0 = n))^{1/2} \\ &\leq \sum_{n \geq 1} M^{n\beta/\alpha} \left(\frac{4e^{\lambda^2 T/4}}{\lambda\sqrt{T}} \varepsilon \right)^{(n-1)/2} = \mathcal{M} < \infty \end{aligned}$$

for β small enough.

By Markov's inequality, we conclude that for all $k \geq 1$

$$\begin{aligned} &|\mathbb{E}(\varphi(X(kT, x))) - \mathbb{E}(\varphi(X(kT, y)))| \\ &= |\mathbb{E}(\varphi(X_1(kT, x, y))) - \mathbb{E}(\varphi(X_2(kT, x, y)))| \\ &\leq 2\|\varphi\|_\infty \mathbb{P}(X_1(kT, x, y) \neq X_2(kT, x, y)) \\ &\leq 2\|\varphi\|_\infty \mathbb{P}(kT > \tau_{k_0} + T) \\ &\leq 2\|\varphi\|_\infty \mathcal{M} e^{-\beta(k-1)T}. \end{aligned}$$

We define $k := \left\lfloor \frac{t}{T} \right\rfloor$ such that we have $P_t^c = P_{kT}^c P_{t-kT}^c$. Thus we can write

$$\begin{aligned} |\mathbb{E}[\varphi(X(t, x))] - \nu_c(\varphi)| &= \left| P_t^c \varphi(x) - \int_{H_c} \varphi(y) \nu_c(dy) \right| \\ &= \left| \int_{H_c} P_t^c \varphi(x) \nu_c(dy) - \int_{H_c} P_t^c \varphi(y) \nu_c(dy) \right| \\ &\leq \left| \int_{H_c} (P_{kT}^c P_{t-kT}^c \varphi(x) - P_{kT}^c P_{t-kT}^c \varphi(y)) \nu_c(dy) \right| \\ &\leq \int_{H_c} 2\|P_{t-kT}^c \varphi\|_\infty \mathcal{M} e^{-\beta(k-1)T} \nu_c(dy) \\ &\leq 2\|\varphi\|_\infty \mathcal{M} e^{-\beta(k-1)T} \\ &\leq C\|\varphi\|_\infty e^{-\beta t}. \end{aligned}$$

□

References

- [1] P. W. Bates and P. C. Fife. The dynamics of nucleation for the Cahn-Hilliard equation. *SIAM J. Appl. Math.*, 53(4):990–1008, 1993.
- [2] D. Blömker, S. Maier-Paape, and T. Wanner. Spinodal decomposition for the Cahn-Hilliard-Cook equation. *Communications in Mathematical Physics*, 223(3):553–582, 2001.
- [3] D. Blömker, S. Maier-Paape, and T. Wanner. Phase separation in stochastic Cahn-Hilliard models. *Mathematical Methods and Models in Phase Transitions*, pages 1–41, 2005.
- [4] D. Blömker, S. Maier-Paape, and T. Wanner. Second phase spinodal decomposition for the Cahn-Hilliard-Cook equation. *Trans. Amer. Math. Soc.*, 360(1):449–489 (electronic), 2008.

- [5] J. F. Blowey and C. M. Elliott. The Cahn-Hilliard gradient theory for phase separation with nonsmooth free energy. I. Mathematical analysis. *European J. Appl. Math.*, 2(3):233–280, 1991.
- [6] J. W. Cahn. On spinodal decomposition. *Acta Metallurgica*, 9(9):795–801, September 1961.
- [7] J. W. Cahn and J. E. Hilliard. Free energy of a nonuniform system. i. interfacial free energy. *Journal of Chemical Physics*, 28(258), February 1958.
- [8] J. W. Cahn and J. E. Hilliard. Spinodal decomposition: a reprise. *Acta Metallurgica*, 19(2):151–161, February 1971.
- [9] C. Cardon-Weber. Cahn-Hilliard stochastic equation: existence of the solution and of its density. *Bernoulli*, 7(5):777–816, 2001.
- [10] C. Cardon-Weber. Cahn-Hilliard stochastic equation: strict positivity of the density. *Stoch. Stoch. Rep.*, 72(3-4):191–227, 2002.
- [11] H. Cook. Brownian motion in spinodal decomposition. *Acta Metallurgica*, 18:297–306, 1970.
- [12] G. Da Prato and A. Debussche. Stochastic Cahn-Hilliard equation. *Nonlinear Anal.*, 26(2):241–263, 1996.
- [13] G. Da Prato, A. Debussche, and B. Goldys. Some properties of invariant measures of non symmetric dissipative stochastic systems. *Probab. Theory Related Fields*, 123(3):355–380, 2002.
- [14] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1992.
- [15] G. Da Prato and J. Zabczyk. *Ergodicity for infinite-dimensional systems*, volume 229 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1996.
- [16] R. C. Dalang, C. Mueller, and L. Zambotti. Hitting properties of parabolic s.p.d.e.’s with reflection. *Ann. Probab.*, 34(4):1423–1450, 2006.
- [17] A. Debussche and L. Dettori. On the Cahn-Hilliard equation with a logarithmic free energy. *Nonlinear Anal.*, 24(10):1491–1514, 1995.
- [18] A. Debussche and L. Zambotti. Conservative stochastic Cahn-Hilliard equation with reflection. *Ann. Probab.*, 35(5):1706–1739, 2007.
- [19] N. Elezović and A. Mikelić. On the stochastic Cahn-Hilliard equation. *Nonlinear Anal.*, 16(12):1169–1200, 1991.
- [20] T. Funaki and K. Ishitani. Integration by parts formulae for Wiener measures on a path space between two curves. *Probab. Theory Related Fields*, 137(3-4):289–321, 2007.
- [21] T. Funaki and S. Olla. Fluctuations for $\nabla\phi$ interface model on a wall. *Stochastic Process. Appl.*, 94(1):1–27, 2001.
- [22] G. Giacomin, S. Olla, and H. Spohn. Equilibrium fluctuations for $\nabla\phi$ interface model. *Ann. Probab.*, 29(3):1138–1172, 2001.
- [23] L. Goudenège. Stochastic Cahn-Hilliard equation with singular nonlinearity and reflection. *Stochastic Processes and their Applications*, 2009.
- [24] C. P. Grant. Spinodal decomposition for the Cahn-Hilliard equation. *Comm. Partial Differential Equations*, 18(3-4):453–490, 1993.
- [25] I. Gyöngy and N. Krylov. Existence of strong solutions for Itô’s stochastic equations via approximations. *Probab. Theory Related Fields*, 105(2):143–158, 1996.

- [26] S. Kuksin and A. Shirikyan. A coupling approach to randomly forced nonlinear PDE's. I. *Comm. Math. Phys.*, 221(2):351–366, 2001.
- [27] J. S. Langer. Theory of spinodal decomposition in alloys. *Annals of Physics*, 65:53–86, 1971.
- [28] S. Maier-Paape and T. Wanner. Spinodal decomposition for the Cahn-Hilliard equation in higher dimensions. I. Probability and wavelength estimate. *Comm. Math. Phys.*, 195(2):435–464, 1998.
- [29] S. Maier-Paape and T. Wanner. Spinodal decomposition for the Cahn-Hilliard equation in higher dimensions: nonlinear dynamics. *Arch. Ration. Mech. Anal.*, 151(3):187–219, 2000.
- [30] J. C. Mattingly. Exponential convergence for the stochastically forced Navier-Stokes equations and other partially dissipative dynamics. *Comm. Math. Phys.*, 230(3):421–462, 2002.
- [31] A. Novick-Cohen. The Cahn-Hilliard equation: mathematical and modeling perspectives. *Adv. Math. Sci. Appl.*, 8(2):965–985, 1998.
- [32] A. Novick-Cohen and L. A. Segel. Nonlinear aspects of the Cahn-Hilliard equation. *Phys. D*, 10(3):277–298, 1984.
- [33] D. Nualart and É. Pardoux. White noise driven quasilinear SPDEs with reflection. *Probab. Theory Related Fields*, 93(1):77–89, 1992.
- [34] C. Odasso. Exponential mixing for the 3D stochastic Navier-Stokes equations. *Comm. Math. Phys.*, 270(1):109–139, 2007.
- [35] E. Sander and T. Wanner. Monte Carlo simulations for spinodal decomposition. *J. Statist. Phys.*, 95(5-6):925–948, 1999.
- [36] E. Sander and T. Wanner. Unexpectedly linear behavior for the Cahn-Hilliard equation. *SIAM J. Appl. Math.*, 60(6):2182–2202 (electronic), 2000.
- [37] T. Wanner. Maximum norms of random sums and transient pattern formation. *Trans. Amer. Math. Soc.*, 356(6):2251–2279 (electronic), 2004.
- [38] L. Zambotti. Integration by parts formulae on convex sets of paths and applications to SPDEs with reflection. *Probab. Theory Related Fields*, 123(4):579–600, 2002.
- [39] L. Zambotti. Integration by parts on δ -Bessel bridges, $\delta > 3$ and related SPDEs. *Ann. Probab.*, 31(1):323–348, 2003.
- [40] L. Zambotti. Fluctuations for a $\nabla\phi$ interface model with repulsion from a wall. *Probab. Theory Related Fields*, 129(3):315–339, 2004.