



Bioaccumulation of inorganic Hg by the juvenile cuttlefish *Sepia officinalis* exposed to ^{203}Hg radiolabelled seawater and food

Thomas Lacoue-Labarthe, Michel Warnau, François Oberhänsli, Jean-Louis Teyssié, Paco Bustamante

► To cite this version:

Thomas Lacoue-Labarthe, Michel Warnau, François Oberhänsli, Jean-Louis Teyssié, Paco Bustamante. Bioaccumulation of inorganic Hg by the juvenile cuttlefish *Sepia officinalis* exposed to ^{203}Hg radiolabelled seawater and food. *Aquatic Biology*, 2009, 6, pp.91-98. 10.3354/ab00172 . hal-00415807

HAL Id: hal-00415807

<https://hal.science/hal-00415807>

Submitted on 11 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Bioaccumulation of inorganic Hg by the juvenile cuttlefish *Sepia officinalis*
exposed to ²⁰³Hg radiolabelled seawater and food**

Lacoue-Labarthe T.¹⁻², Warnau M.¹⁻², Oberhänsli F.², Teyssié J.-L.², Bustamante P.^{1*}

1- Littoral Environnement et Sociétés (LIENSs), UMR 6250, CNRS-Université de La
Rochelle, 2 rue Olympe de Gouges, F-17042 La Rochelle Cedex 01, France

2- International Atomic Energy Agency – Marine Environment Laboratories, 4 Quai Antoine
I^{er}, MC-98000 Principality of Monaco

* Corresponding author: Prof. Paco Bustamante
Littoral Environnement et Sociétés (LIENSs)
UMR 6250, CNRS-Université de La Rochelle
2 rue Olympe de Gouges
F-17042 La Rochelle Cedex 01
France

Phone: +33-546-507-625

Fax: +33-546-458-264

E-mail: pbustama@univ-lr.fr

ABSTRACT: Uptake and depuration kinetics of inorganic mercury (Hg) was investigated in the juvenile common cuttlefish *Sepia officinalis* following exposures via seawater and food using sensitive radiotracer technique (^{203}Hg). The cuttlefish readily concentrated ^{203}Hg when exposed via seawater, with whole body CF > 260 after only 10d of exposure. The total Hg accumulated from seawater was depurated relatively fast with a $\text{Tb}_{1/2}$ of 17d. During both exposure and depuration periods, accumulated Hg was mainly (> 70%) associated with the muscular parts of the cuttlefish. However, the proportion of the whole body Hg content associated with the digestive gland increased during exposure and depuration phases, suggesting that the metal was transferred from the muscles towards this organ for detoxification. When fed with radiolabelled food, cuttlefish displayed high assimilation efficiency (AE > 90%) and the metal was found to be mainly located in the digestive gland (60% of the whole Hg content). Nevertheless, high depuration rates resulted in short $\text{Tb}_{1/2}$ (i.e., 4 d), suggesting that this organ has a major role in Hg detoxification and depuration. Whatever the exposure pathway was, low proportion of Hg (< 2%) was found in the cuttlebone. Assessment of the relative contribution of the dietary and dissolved exposure pathways to inorganic Hg bioaccumulation in juvenile cuttlefish revealed that Hg was mainly accumulated from food that contributed for $77 \pm 16\%$ of the global metal bioaccumulation.

Keywords: mercury; bioaccumulation; kinetics; body distribution; cephalopod; relative contribution.

INTRODUCTION

Mercury (Hg) is one of the metals of highest concern in the marine environment as it is readily methylated by micro-organisms, bioaccumulates in marine biota and consistently biomagnifies along the food chain (Cossa 1990). Among marine organisms, most of the available information on Hg is related to fish, mainly because of their importance as a food source for human. In fish, most of the Hg (i.e. > 95%) is methylated and is therefore bioavailable for upper trophic levels (Bloom 1992). Hence, fish consumption is an important source of Hg for human (Svensson et al. 1992) and is of particular health concern (Clarkson 1990).

In contrast to fish, information on Hg in cephalopods tissues is scarce despite the fact that these molluscs represent an increasing component of the world fisheries (Boyle & Rodhouse, 2006, FAO 2007). In addition, information on Hg in cephalopod is essentially limited to the main species targeted by fisheries and mainly reports metals levels in edible tissues, i.e. mantle muscle, arms and fins (e.g. Buzina et al. 1989, Sapunar et al. 1989, Plessi et al. 2001). Recently, a study on a large range of cephalopod species from the North Eastern Atlantic waters reported that Hg concentrations varied on two orders of magnitude among cephalopod species and that the metal was mainly stored under organic form in the muscular tissues (Bustamante et al. 2006). Several other studies on cephalopods from the Mediterranean suggest that these molluscs are able to accumulate high Hg concentrations in their tissues (Renzoni et al. 1973, Rossi et al. 1993, Storelli & Marcotrigiano 1999). Various factors are likely to influence Hg concentrations in cephalopods among which the size seems to be of primary importance (Monteiro et al. 1992, Rossi et al. 1993, Pierce et al. 2008).

Although it was suggested that food would be the main source for Hg accumulation in cephalopod tissues (Bustamante et al., 2006), the relative contribution of dietary and waterborne pathways has not been assessed in cephalopods. Moreover, as cephalopods are

short-lived species, they might be interesting as short-term indicator species of the variation of Hg concentrations in the environment (Seixas et al. 2005, Pierce et al. 2008).

For these reasons, the aim of this work was to investigate the biokinetics of Hg uptake and depuration in cephalopods in order to better characterize its bioaccumulation, tissue distribution and retention capacity. The common cuttlefish *Sepia officinalis* was selected as a model to study Hg transfer in cephalopods from seawater and food.

MATERIALS AND METHODS

Organisms and radiotracer

Adult cuttlefish were collected by net fishing off Monaco in March and April 2006. They were acclimated and maintained in open-circuit tanks in the IAEA-MEL premises. After mating, the eggs laid by a single female were separated to optimise their oxygenation and kept in a separate aquarium during the whole embryonic development (constantly aerated open circuit; flux: 50 l h⁻¹; salinity: 37 p.s.u; temperature: 17 ± 0.5°C; pH: 8.0 ± 0.1; light/dark cycle: 12 h/12 h). Hatching occurred approx. 50 d after the spawning. Young cuttlefish were then maintained in the same aquarium and fed with brine shrimp (*Artemia* sp.) for 5 days before the experiments.

The radiotracer, ²⁰³Hg [as ²⁰³HgNO₃; t_{1/2} = 46.59 d], was purchased from Isotope Product Laboratory, USA. Stock solutions were prepared in 1 N nitric acid to obtain final radioactivity allowing the use of spikes of only a few microliters (typically 5 µl).

Experimental procedure

Contamination from seawater: juveniles (n = 23; mean weight ± SD, 0.258 ± 0.009 g) were placed for 10 d in a 20-l glass aquarium containing 0.45 µm filtered natural seawater (constantly aerated closed circuit; temperature 17°C; 37 p.s.u.; light/dark cycle 12h/12h)

99 spiked with ^{203}Hg (0.6 kBq l^{-1}). In terms of stable metal, this concentration corresponded to 20
100 ng l^{-1} . To facilitate the recurrent counting of each individual during the experiment, the
101 juveniles were held individually in separate circular plastics boxes (10 cm diameter, 5 cm
102 height) covered up and down with a meshed plastic net to allow for free water circulation.
103 Radiotracer and seawater were renewed every second day to maintain water quality and
104 radiotracer activity constant. Radiotracer activities in seawater were checked before and after
105 each water renewal in order to determine the time-integrated radiotracer activities (Rodriguez
106 y Baena et al. 2006a, Warnau et al. 1996). Juveniles were separated in two groups: the first
107 group contained 16 tag-identified individuals and the second was composed of unidentified
108 animals for the body distribution analyses. At different time intervals, radiotracer activities
109 were counted in the same tag-identified juveniles ($n = 16$) all along the experiment.
110 According to their physiological states, the individuals should be removed from the
111 experiment leading the following sampling plan: $n=16$ from days 0 to 3 and $n = 14, 7, 6$ at
112 days 6, 9 10, respectively. In addition, after 3 and 9 d of exposure, 3 juveniles of the second
113 group were counted and dissected to determine the radiotracer distribution among the
114 digestive gland, the cuttlebone and the remaining tissues.
115 After this exposure period, the 6 remaining identified and radiolabelled juveniles ($n= 6$ from
116 days 0 to 4, 4 at day 6 and 1 at day 11) were held for 11 d in clean flowing water (open circuit
117 with constant aeration; seawater flux 50 l h^{-1} ; temperature 17°C ; 37 p.s.u.; light/dark cycle
118 12h/12h). At different time intervals during the depuration period, the same identified
119 juveniles ($n = 6$) were counted to establish the depuration kinetics of the radiotracer.

120
121 ***Contamination through the food:*** brine shrimp (*Artemia* sp.) were exposed for 5 d in a
122 plastic aquarium containing 4 l of natural seawater spiked with ^{203}Hg until pools of 25 brine

shrimp reached 3.1 kBq g^{-1} wet weight. The organisms were subsequently used as food for the juvenile cuttlefish.

As detailed in the seawater experimental procedure, juveniles were separated in two groups of identified and non-identified animals, which were devoted for the depuration kinetics and the body distribution studies, respectively. Hence, sixteen newly hatched cuttlefish (mean weight \pm SD, $0.297 \pm 0.011 \text{ g}$) were placed in individual plastic containers (10 cm diameter, 5 cm height), and held in a 20-l aquarium under the same conditions as in the previous experiment. Each juvenile was fed for 1 h with 25 of the previously radiolabelled *Artemia* sp. At the end of the feeding period, the cuttlefish were immediately counted. From that time on, the cuttlefish were fed twice a day with uncontaminated *Artemia* sp. for one month and regularly counted to determine radiotracer depuration kinetics and assimilation efficiency. As mentioned above, juveniles showing poor health condition were removed leading to a sample number decrease along the experiment: $n = 16$ from days 0 to 2, 14 from days 3 to 14, 4 from days 17 to 22 and 1 from days 24. Throughout the depuration period, faeces were removed twice a day to reduce possible radiotracer recycling through leaching from the faeces. In addition, after 3 h, 9 d, and 22 d of exposure, 3 others juveniles were counted and dissected to determine the radiotracer distribution among the digestive gland, the cuttlebone and the remaining tissues.

Radioanalyses and data treatment

Radioactivities were measured using a high-resolution γ -spectrometry system consisting of four coaxial Germanium (N- or P-type) detectors (EGNC 33-195-R, Canberra[®] and Eurysis[®]) connected to a multi-channel analyzer and a computer equipped with a spectra analysis software (Interwinner[®] 6). The detectors were calibrated with an appropriate standard for each counting geometry used and measurements were corrected for background and physical decay

of the radiotracer. Counting times were adapted to obtain relative propagated errors less than 5% (Rodriguez y Baena, 2006b). They ranged from 10 min to 30 min for whole juveniles and from 10 min to 24h for the dissected tissues.

Uptake of the ^{203}Hg from seawater was expressed as change in concentration factors (CF; ratio between radiotracer content in the juvenile – Bq g^{-1} – and time-integrated activity in seawater – Bq g^{-1}) along time (Warnau et al. 1996). Uptake kinetic was best described by a saturation equation (Eq.1):

$$\text{CF}_t = \text{CF}_{ss} (1 - e^{-k_e t}) \quad (\text{Eq. 1})$$

where CF_t and CF_{ss} are the concentration factors at time t (d) and at steady state, respectively, k_e is the biological depuration rate constants (d^{-1}) (Whicker & Schultz 1982).

Radiotracer depuration kinetics were expressed in terms of change of percentage of remaining activity (i.e., radioactivity at time t divided by initial radioactivity measured in the organisms or in the tissue at the beginning of the depuration period * 100) along with time.

The depuration kinetic was best fitted by a mono-exponential equation (Eq. 2):

$$A_t = A_0 e^{-k_e t} \quad (\text{Eq. 2})$$

where A_t and A_0 are the remaining activities (%) at time t (d) and 0, respectively, and k_e is the biological depuration rate constant (d^{-1}). The determination of k_e allows the calculation of the radiotracer biological half-life ($T_{b/2} = \ln 2 / k_e$). In the context of seawater and feeding experiment, A_0 represents the absorption ($A_{0,w}$) and the assimilation (AE) efficiencies, respectively.

Bioaccumulation model

The relative contribution of each uptake pathway was determined using the bioaccumulation model originally proposed by Thomann (1981) and revised by Thomann et al. (1995) and Metian et al. (2008). In this model, the total concentration of radiotracers in the juveniles, C_t

(Bq g⁻¹) is equal to the sum of each concentration resulting from the uptake by the different pathways (Eq. 3):

$$C_t = C_{f,ss} + C_{w,ss} \quad (\text{Eq. 3})$$

where $C_{f,ss}$ is the food-derived radiotracer concentration (Bq g⁻¹) in juveniles at steady state (Eq. 4) and C_w is the water-derived radiotracer concentration (Bq g⁻¹) in juveniles at steady state (Eq. 5).

$$C_{f,ss} = (AE \times IR \times C_f) / k_{e,f} \quad (\text{Eq. 4})$$

$$C_{w,ss} = (A_{0,w} \times k_{u,w} \times C_w) / k_{e,w} \quad (\text{Eq. 5})$$

where $A_{0,w}$ is the absorption efficiencies (%) of the radiotracer from seawater, AE is the assimilation efficiency (%) of the radiotracer from food, C_f and C_w are the radiotracer activities in food and seawater (Bq g⁻¹ and Bq ml⁻¹, respectively), respectively, IR is the ingestion rate (g g⁻¹ d⁻¹), $k_{u,w}$ is the uptake rate constants (d⁻¹) from seawater, $k_{e,f}$ and $k_{e,w}$ are the biological depuration rate constants (d⁻¹), for food and water pathways, respectively.

The relative contribution (%) of each uptake pathway is then assessed from the relation:

$$\% \text{ food} = C_{f,ss} / (C_{f,ss} + C_{w,ss})$$

$$\% \text{ seawater} = C_{w,ss} / (C_{f,ss} + C_{w,ss})$$

Constants (and their statistics) of the best fitting equations (decision based on ANOVA tables for two fitted model objects) were estimated by iterative adjustment of the models using the *nls* curve-fitting routine in R freeware. The level of significance for statistical analysis was always set at $\alpha = 0.05$.

Results

Contamination through seawater

Uptake activity of ^{203}Hg in whole-body *S. officinalis* was best fitted by a saturation exponential equation with a calculated CF_{ss} of 480 (Fig. 1; Table 1). The CF actually measured at the end of the uptake period ($\text{CF}_{10\text{d}}$) of ^{203}Hg was 260 ± 70 (Table 2). Calculated $\text{CF}_{10\text{d}}$ for the different organs indicated that ^{203}Hg was concentrated according to the following decreasing order: digestive gland (1460 ± 480) > remaining tissues (290 ± 80) > cuttlebone (47 ± 9). In terms of body distribution, ^{203}Hg was mainly found in the remaining tissues all along the exposure period; this compartment accounted for 89 ± 3 and 80 ± 4 % of the whole body load of ^{203}Hg after 3 d and 10 d of exposure, respectively (Table 2). The cuttlebone presented very low Hg activity ($< 15 \text{ Bq g}^{-1}$) and loads ($< 3\%$) whereas the Hg activity in the digestive gland was increasing with time, varying from $24 \pm 4 \text{ Bq g}^{-1}$ at day 3 to $520 \pm 170 \text{ Bq g}^{-1}$ at day 9 corresponding to 8% and 20% of the total radioactivity loads, respectively.

After the 10 d exposure period, non-contaminating conditions were restored and depuration kinetic of ^{203}Hg was followed for 11 d. The whole-body depuration kinetic of ^{203}Hg in *S. officinalis* was best described by a mono-exponential model (Fig. 2; Table 1). This result indicated that 95% of the ^{203}Hg previously accumulated were depurated with a relatively biological half-life relatively short (i.e. 17 d, Table 1). After 11 d of depuration conditions, most of the ^{203}Hg body load was associated with the remaining tissues (72%) while the digestive gland proportion increase up to 28% (Table 2).

Contamination through food

The depuration kinetic of ^{203}Hg ingested with food in *S. officinalis* was best fitted by a mono-exponential model, characterized by a biological half-life of 4 d (Fig.2; Table 1) and allowed an estimated assimilation efficiency of 91%.

During the depuration period, the digestive gland displayed highest proportion of the total body burden of ^{203}Hg , i.e. 68, 60, 64 % after 3 h, 9 d and 22 d of depuration, respectively (Table 2). The distribution of ^{203}Hg remained unchanged all along the loss experiment, with the cuttlebone always showing the lowest proportion of the radiotracer (Table 2).

Bioaccumulation model

In order to assess the relative contribution of each uptake pathway to the global Hg accumulation in *S. officinalis*, the different kinetic parameters obtained for seawater and food experiments were used to feed the bioaccumulation model, along with other parameters such as the ^{203}Hg concentration in seawater and food (C_w : 0.364 Bq ml^{-1} and C_f : 3100 Bq g^{-1} , respectively, present study) and the ingestion rate (IR: $0.07 \text{ g g}^{-1} \text{ d}^{-1}$, present study value being congruent with IR value determined by Koueta et al., 1999). Modelling showed that food represented the main pathway for ^{203}Hg bioaccumulation in the juvenile of cuttlefish, contributing to $77 \pm 16\%$ of the global metal bioaccumulation vs. $23 \pm 14\%$ for the seawater pathway.

Discussion

Cephalopods are an increasing marine resource for world fisheries (Boyle & Rodhouse 2006). In the 1990s alone there was a 40% increase in squid catches worldwide (FAO 2007). Cephalopods are well-known for their capacity to accumulate high levels of non-essential metals, especially Ag and Cd, in their tissues (e.g. Martin & Flegal 1975, Bustamante et al. 1998, 2008). Hence, the intake of contaminants such as metals by human through cephalopod consumption is a matter of concern (e.g. Pierce et al. 2008). Some studies also reported high concentrations of Hg in cephalopods from areas naturally contaminated by cinnabar such the Tyrrhenian and Adriatic Seas (e.g. Renzoni et al. 1973, Rossi et al. 1993, Storelli &

Marcotrigiano 1999). However, the dynamic of Hg incorporation and its metabolism in cephalopods remain poorly understood. In the field, Hg could be methylated that generally increases its toxicity as a result of its enhanced capacity of penetration across cell membranes (Boudou et al. 1983). The known methylHg uptake pathways in the biota are the transfer from the sediment where sediment-associated bacterial floras able to methylate part of the inorganic Hg (Compeau & Bartha 1985), and the consumption of the prey such as juveniles of shrimp, crab or fish where Hg is stored under the methylated form in variable proportions. Considering that sand burying is a transient cryptic behaviour of cuttlefish (Poirier et al. 2004) and the grain size of the sediment selected by these species did not favour the methylation of Hg, the inorganic form of the metal could also be a significant source of accumulation for this nectobenthic cephalopod. In this study, uptake and depuration biokinetics of Hg were determined using inorganic carrier-free ^{203}Hg in order to measure metal fluxes in real time at environmentally realistic contaminant concentrations (Warnau et al. 1996).

As a typical cephalopod, the common cuttlefish *Sepia officinalis* has high food intake requirements to sustain its elevated growth rate. Being active predators, cephalopods have a high digestion efficiency (Boucher-Rodoni et al. 1987) and food has been shown to constitute an important source of uptake for various trace elements such as Am, Cd, Co and Zn (e.g., Guary & Fowler 1982, Koyama et al. 2000, Bustamante et al. 2002ab, 2004). However, seawater could be an important bioaccumulation pathway as well, as elements can be taken up efficiently through the skin and the gills. For instance, it has been shown that seawater is the main intake pathway for Ag in adult *S. officinalis* (Bustamante et al. 2004). Therefore, there was a need to provide insights on the bioaccumulation of Hg under controlled experimental conditions in order to delineate the contribution of its uptake via the dissolved and dietary pathways.

After 10 d of exposure to dissolved ^{203}Hg , cuttlefish displayed quite elevated whole-body activities ($\text{CF} = 260 \pm 70$), indicating that Hg was efficiently bioconcentrated from seawater (Table 2). The estimated steady-state equilibrium ($\text{CF}_{\text{ss}} = 480 \pm 150$) would be reached after 2 weeks of exposure. Among the three considered compartments, the digestive gland displayed the highest concentration capacity ^{203}Hg ($\text{CF} = 1460 \pm 480$; Table 2) whereas the cuttlebone showed the lowest CF (i.e. less than 50; Table 2). This pattern is very similar to that for experimental data reported for other metals such as Ag, Cd, Co and Zn (Bustamante et al. 2002a, 2004). However, in terms of body burden distribution, ^{203}Hg was mainly stored in the remaining tissues (up to 80 % of the total Hg load; Table 2), which are mainly composed of muscles although they include the skin and the respiratory organs (i.e. gills) corresponding to the tissues directly exposed to the contaminated seawater. Consistently, Bustamante et al. (2006) reported that 70-90% of the total Hg body burden was stored in the muscular tissues in different cephalopod species collected from the North East Atlantic. This could be explained by the fact that 1) that these tissues represents more than 70% of the total body weight and 2) that Hg has a stronger affinity for the sulphydryl groups of muscular proteins rather than of fat tissue found in fish and/or cephalopod (Bloom 1992, Bustamante et al. 2006).

When *Sepia officinalis* was exposed to dissolved Hg, the digestive gland contained 8% and 20% of the total Hg burden after 3 d and 10 d exposure, respectively (Table 2). Therefore, this increasing Hg proportion found in this organ implied that the digestive gland accumulated more efficiently than the others. In a second time, in depuration conditions, ^{203}Hg was release of following a mono-exponential model and whole-body depuration was relatively rapid with a $T_{b/2}$ of approx. 17 d. After 11 d of depuration under running seawater, the tissues in direct contact with seawater contained significantly less ^{203}Hg than at the beginning of the depuration period varying from $135 \pm 15 \text{ Bq g}^{-1}$ and to $95 \pm 17 \text{ Bq g}^{-1}$ at d 0 and d 11 of the depuration period, respectively, whereas the digestive gland activity remained unchanged

(i.e., from $520 \pm 170 \text{ Bq g}^{-1}$ at day 0 to $560 \pm 30 \text{ Bq g}^{-1}$ at day 11; Table 2). These results suggest that 1) the digestive gland shows a stronger retention capacity for Hg than the other compartments, and/or that 2) a metal translocation occurred from the remaining tissues towards the digestive gland. In cephalopods, the digestive gland plays obviously a major role in the digestive processes but also in the detoxification of xenobiotics. Indeed, this organ has already been shown to retain translocated metals from the other tissues (Bustamante et al. 2002a, 2004). The digestive gland is involved in the storage and detoxification of several metals such as Ag, Cd, Cu, or Zn (e.g. Miramand & Bentley 1992, Bustamante et al. 2002b, Dorneles et al. 2007) and persistent organic pollutants such as PCBs (Ueno et al. 2003, Danis et al. 2005, Storelli et al., 2006). However, due to the fact this organ does not store Hg in large amounts as shown in the field study (Bustamante et al. 2006), the digestive gland might be also involved in the depuration of Hg when cuttlefish is exposed to the dissolved metal. In the case of dietary exposure, the AE of Hg ingested with food was found to be nearly 100% (Table 1). This high degree of Hg assimilation might be due to the very efficient digestive metabolism that characterizes the juvenile cuttlefish (Mangold 1989). Indeed, this early life stage is characterized by a predominant intracellular digestion process (compared to the extracellular digestion which is dominant in adults) (Boucaud-Camou & Roper 1995), which could favour the metal assimilation. Such extreme assimilation efficiency has already been documented in cuttlefish: for instance, 90% and almost 100% AE for Cd and Co, respectively, were reported in juvenile fed brine shrimp (Bustamante et al. 2002a, 2004). Nevertheless, the assimilated Hg was rapidly depurated with a $T_{b1/2}$ of 4 d, which suggests that the processes governing Hg elimination are particularly efficient as well. Indeed, the ^{203}Hg activities in the digestive gland dropped from $310 \pm 80 \text{ Bq g}^{-1}$ to $36 \pm 6 \text{ Bq g}^{-1}$ during the 22 d depuration period after the feeding (Table 2). These results highlight the efficient excretion capacity of the digestive gland for Hg. Because following dietary exposure the main fraction of the whole

Hg body burden was associated with this organ, it is not surprising that the retention time of Hg following dietary exposure was 4 times lower than following seawater exposure, i.e. 4 d vs. 17 d (Table 1).

In our experimental conditions, it appeared that the exposure of cuttlefish to contaminated seawater lead to an accumulation of the Hg in the juvenile remaining tissues (> 80%) mainly composed of muscular tissues. In the same time, Hg was translocated to the digestive gland and subsequently eliminated from the organism. Following dietary exposure, inorganic Hg was assimilated via the digestive gland and then rapidly eliminated. Consequently, the storage and/or redistribution of the bioaccumulated metal towards the remaining tissues is limited in cuttlefish. Nonetheless, considering both seawater and dietary exposure, food appears as the predominant pathway for inorganic Hg bioaccumulation in juvenile cuttlefish. This result is not surprising considering the relative high ingestion rate and the efficient digestive metabolism of cephalopods (Lee 1994).

Acknowledgements

Authors are grateful to M. Metian (IAEA-MEL) for skilful assistance during the experiments. MW is an Honorary Senior Research Associate of the National Fund for Scientific Research (NFSR, Belgium). The IAEA is grateful for the support provided to its Marine Environment Laboratories by the Government of the Principality of Monaco. This work was financially supported by the IAEA and by LIENSs (University of La Rochelle).

References

346 Bloom NS (1992) On the chemical form of mercury in edible fish and marine invertebrate
 347 tissue. Can J Fish Aquat Sci 49: 1010-1017

348 Boucaud-Camou E, Roper CFE (1995) Digestive enzymes in paralarval cephalopods. Bull
 349 Mar Sci 57: 313-327

350 Boucher-Rodoni R, Boucaud-Camou E, Mangold K (1987) Feeding and digestion. In: Boyle
 351 PR (ed) Cephalopod Life Cycle, Vol 2. Academic Press, London, p 85-108

352 Boudou A, Georgescauld D, Desmazes JP (1983) Ecotoxicological role of the membrane
 353 barriers in transport and bioaccumulation of mercury compounds. In: Nriagu JO (ed)
 354 Aquatic Toxicology, J. Wiley & Sons, New York, p 118-136

355 Boyle PR, Rodhouse PG (2006) Cephalopods: Ecology and Fisheries. Blackwell Science,
 356 Oxford, United Kingdom

357 Bustamante P, Caurant F, Fowler SW, Miramand P (1998) Cephalopods as a vector for the
 358 transfer of cadmium to top marine predators in the north-east Atlantic Ocean. Sci Total
 359 Environ 220: 71-80

360 Bustamante P, Teyssié J-L, Fowler SW, Cotret O, Danis B, Miramand P, Warnau M (2002a)
 361 Biokinetics of zinc and cadmium accumulation and depuration at different stages in the
 362 life cycle of the cuttlefish *Sepia officinalis*. Mar Ecol Prog Ser 231: 167-177

363 Bustamante P, Cosson RP, Gallien I, Caurant F, Miramand P (2002b) Cadmium
 364 detoxification processes in the digestive gland of cephalopods in relation to
 365 accumulated cadmium concentrations. Mar Environ Res 53: 227-241

366 Bustamante P, Teyssié J-L, Danis B, Fowler SW, Miramand P, Cotret O, Warnau M (2004)
 367 Uptake, transfer and distribution of silver and cobalt in tissues of the common cuttlefish
 368 *Sepia officinalis* at different stages of its life cycle. Mar Ecol Prog Ser 269: 185-195

369 Bustamante P, Lahaye V, Durnez C, Churlaud C, Caurant F (2006) Total and organic Hg
 370 concentrations in cephalopods from the North Eastern Atlantic waters: Influence of

geographical origin and feeding ecology. *Sci Total Environ* 368: 585-596

Bustamante P, Gonzalez AF, Rocha F, Miramand P, Guerra A (2008) Metal and metalloid concentrations in the giant squid *Architeuthis dux* from Iberian waters. *Mar Environ Res* 66: 278-287

Buzina R, Suboticaneć K, Vukusić J, Sapunar J, Antonić K, Zorica M (1989) Effect of industrial pollution on seafood content and dietary intake of total and methylmercury. *Sci Total Environ* 78: 45-57

Clarkson TW (1990) Human health risks from methylmercury in fish. *Environ Toxicol Chem* 9:957-961

Compeau GC, Bartha R (1985) Sulphate-reducing bacteria: principal methylators of mercury in anoxic estuarine sediment. *Appl Environ Microbiol* 50: 498-502

Cossa D, Thibaud Y, Roméo M, Gnassia-Barelli M (1990) Le mercure en milieu marin. Biogéochimie et écotoxicologie. Rapport scientifiques et techniques de l'IFREMER, N°19, Brest

Danis B, Bustamante P, Cotret O, Teyssié JL, Fowler SW, Warnau M (2005) Bioaccumulation of PCBs in the cuttlefish *Sepia officinalis* from seawater, sediment and food pathways. *Environ Pollut* 134: 113-122

Dorneles PR, Lailson-Brito J, dos Santos RA, Silva da Costa PA, Malm O, Azevedo AF, Machado Torres JP (2007) Cephalopods and cetaceans as indicators of offshore bioavailability of cadmium off Central South Brazil Bight. *Environ Pollut* 148: 352-359

FAO Fisheries and Aquaculture Information and Statistics Service (2007) Fisheries commodities production and trade 1976-2005. FISHSTAT Plus - Universal software for fishery statistical time series [online or CD-ROM]. Food and Agriculture Organization of the United Nations. Available at:

<http://www.fao.org/fi/statist/FISOFT/FISHPLUS.asp>

396 Guary J-C, Fowler SW (1982) Experimental studies on the biokinetics of plutonium and
 397 americium in the cephalopod *Octopus vulgaris*. Mar Ecol Prog Ser 7: 37-43
 398 Koueta N, Boucaud-Camou E (1999) Food intake and growth in reared early juvenile
 399 cuttlefish *Sepia officinalis* L. (Mollusca Cephalopoda). J Exp Mar Biol Ecol 240: 93-109
 400 Koyama J, Nanamori N, Segawa S (2000) Bioaccumulation of waterborne and dietary
 401 cadmium by oval squid, *Sepioteuthis lessoniana*, and its distribution among organs. Mar
 402 Pollut Bull 40: 961-967
 403 Lee PG (1994) Nutrition of cephalopods: fueling the system. In: Pörtner HO, O'Dor RK,
 404 Macmillan DL (eds) Physiology of cephalopod molluscs: lifestyle and performance
 405 adaptations. Gordon and Breach Science Publishers, Basel, p 35-51
 406 Mangold K (1989) Reproduction, croissance et durée de vie. In: Grassé P-P (ed) Traité de
 407 Zoologie : Céphalopodes, Vol V. Masson, Paris, p 493-552
 408 Martin JH, Flegal AR (1975) High copper concentrations in squid livers in association with
 409 elevated levels of silver, cadmium, and zinc. Mar Biol 30: 51-55
 410 Metian M, Bustamante P, Cosson RP, Hédouin L, Warnau M (2008) Investigation of Ag in
 411 the king scallop *Pecten maximus* using field and laboratory approaches. J Exp Mar Biol
 412 Ecol 367: 53-60
 413 Miramand P, Bentley D (1992) Concentration and distribution of heavy metals in tissues of
 414 two cephalopods, *Eledone cirrhosa* and *Sepia officinalis* from the French coast of the
 415 English Channel. Mar Biol 114: 407-414
 416 Monteiro LR, Porteiro, FM, Gonçalves JM (1992) Inter- and intra-specific variation of
 417 mercury levels in muscle of cephalopods from the Azores Archipelago. Life Earth Sci
 418 10: 13-22
 419 Plessi M, Bertelli D, Monzani A (2001) Mercury and selenium content in selected seafood. J
 420 Food Compos Anal 14: 461-467

421 Pierce GJ, Stowasser G, Hastie LC, Bustamante P (2008) Geographic, seasonal and
 422 ontogenetic variation in cadmium and mercury concentrations in squid (Cephalopoda:
 423 Teuthoidea) from UK waters. *Ecotoxicol Environ Saf* 70: 422-432
 424 Poirier R, Chichery R, Dickel L (2004) Effects of rearing conditions on sand digging
 425 efficiency in juvenile cuttlefish. *Behav Process* 67: 273-279
 426 Renzoni A, Bacci E, Falciai L (1973) Mercury concentration in the water, sediments and
 427 fauna of an area of the Tyrrhenian coast. *Revue Int Océan Méd* 31-32: 17-45
 428 Rodriguez y Baena AM, Metian M, Teyssié JL, De Broyer C, Warnau M (2006a)
 429 Experimental evidence for ^{234}Th bioaccumulation in three Antarctic crustaceans:
 430 Potential implications for particle flux studies. *Mar Chem* 100: 354-365
 431 Rodriguez y Baena AM, Miquel JC, Masqué P, Povinec PP, La Rosa J (2006b) A single vs.
 432 double spike approach to improve the accuracy of ^{234}Th measurements in small-volume
 433 seawater samples. *Mar Chem* 100: 269-281
 434 Rossi A, Pellegrini D, Belcari P, Barghigiani C (1993) Mercury in *Eledone cirrhosa* from the
 435 northern Tyrrhenian sea: Contents and relations with life cycle. *Mar Pollut Bull* 26:
 436 683-686
 437 Sapunar J, Jusic M, Bazulic D (1989) Total mercury and cadmium in some cephalopods and
 438 fish from the Adriatic sea. *Mar Pollut Bull* 20: 576-577
 439 Seixas S, Bustamante P, Pierce G (2005) Accumulation of mercury in the tissues of the
 440 common octopus *Octopus vulgaris* (L.) in two localities on the Portuguese coast. *Sci*
 441 *Total Environ* 340: 113-122
 442 Storelli MM, Marcotrigiano GO (1999) Cadmium and total mercury in some cephalopods
 443 from the South Adriatic Sea (Italy). *Food Addit Contam* 16: 261-265
 444 Storelli MM, Barone G, D'Addabbo R, Marcotrigiano GO (2006) Concentrations and
 445 composition of organochlorine contaminants in different species of cephalopod

446 molluscs from the Italian waters (Adriatic Sea). *Chemosphere* 64: 129-134

447 Svensson BG, Schütz A, Nilsson A, Åkesson I, Åkesson B, Skerfving S (1992) Fish as a
 448 source of exposure to mercury and selenium. *Sci Total Environ* 126: 61-74.

449 Thomann RV (1981) Equilibrium model of fate of microcontaminants in diverse aquatic food
 450 chains. *Can J Fish Aquat Sci* 38: 280–296.

451 Thomann RV, Mahony JD, Mueller R (1995) Steady-state model of biota sediment
 452 accumulation factor for metals in two marine bivalves. *Environ Toxicol Chem* 14:
 453 1989-1998

454 Ueno D, Inoue S, Ikeda K, Tanaka H, Yamada H, Tanabe S (2003) Specific accumulation of
 455 polychlorinated biphenyls and organochlorine pesticides in Japanese common squid as
 456 a bioindicator. *Environ Pollut* 125: 227-235

457 Warnau M, Teyssié J-L, Fowler SW (1996) Biokinetics of selected heavy metals and
 458 radionuclides in the common Mediterranean echinoid *Paracentrotus lividus*: sea water
 459 and food exposures. *Mar Ecol Prog Ser* 141: 83-94

460 Whicker FW, Schultz V (1982) *Radioecology: nuclear energy and the environment*, Vol 2.
 461 CRC Press, Boca Raton, FL




Fig. 1: *Sepia officinalis*. Whole-body uptake kinetics of ^{203}Hg in juvenile cuttlefish exposed for 10 days to the radiotracer dissolved in seawater (concentration factors, CF; mean \pm SD, n=16 from days 0 to 3 and n = 14, 7, 6 at days 6, 9 10, respectively).





Fig. 2: *Sepia officinalis*. Whole-body loss kinetics of ^{203}Hg (% of remaining activity; mean \pm SD) in juvenile cuttlefish A) previously exposed to radiolabelled seawater for 10 days (n= 6 from days 0 to 4, 4 at day 6 and 1 at day 11); B) previously fed with radiolabelled brine shrimp (n = 16 from days 0 to 2, 14 from days 3 to 14, 4 from days 17 to 22 and 1 from days 24). Parameters for the best fitting equations are given in Table 1.

Table 1. *Sepia officinalis*. Whole-body uptake and loss kinetic parameters of ^{203}Hg in whole cuttlefish following different exposure experiments:

1) individuals (n = 16) were exposed 10-d to the radiotracer in seawater then placed 11-d in depuration conditions (n = 6);

2) individuals fed on radiolabelled brine shrimp *Artemia* sp. were placed in depuration conditions for 30 d (n = 16);

Uptake parameters: CF_{ss} concentration and transfer factors at steady state; k_u : uptake rate constant (d^{-1})

All loss kinetics followed a mono-exponential depuration fit.

Depuration parameters: A_0 : activity (%) lost according to the exponential component; ASE: asymptotic standard error; k_e : depuration rate constant (d^{-1}), $T_{b/2}$: biological half-life (d); R^2 : determination coefficient of the uptake or loss kinetics; p: probability of the model adjustment:

*** and **: p-values < 0.001 and < 0.01, respectively.

Condition	Uptake			Loss			
	CF_{ss}	k_e	R^2	A_0 (SE)	k_e	$T_{b/2} \pm \text{SE}$ (d)	R^2
1) uptake seawater	480 ± 150	0.083 ± 0.036	0.703	-	-	-	-
2) loss seawater	-	-	-	95.3 (2.8) ***	0.041 ***	16.9 ± 3.9	0.472
3) loss food	-	-	-	90.7 (2.6) ***	0.180 ***	3.9 ± 0.3	0.804

1 Table 2. *Sepia officinalis*. Concentration Factors (mean CF \pm SD) and tissue distribution (mean % \pm SD) of ^{203}Hg during seawater and feeding
2 experiments.

3

Compartments	Seawater contamination					Food contamination		
	Uptake (3d, n = 3)		Uptake (10d, n=3)		Loss (11d, n=1)	Loss (3h, n=3)	Loss (9d, n=3)	Loss (22d, n=3)
	Concentration Factor	Distribution (%)	Concentration Factor	Distribution (%)	Distribution (%)	Distribution (%)	Distribution (%)	Distribution (%)
Digestive gland	110 \pm 20	8.3 \pm 2.6	1460 \pm 480	19.7 \pm 3.8	28.2	67.9 \pm 10.6	59.9 \pm 3.4	63.6 \pm 5.8
Cuttlebone	29.7 \pm 3.7	2.8 \pm 1.0	47.0 \pm 8.8	<1	<1	1.4 \pm 0.9	2.4 \pm 1.9	9.9 \pm 3.2
Remaining tissues	110 \pm 10	88.9 \pm 3.4	290 \pm 80	79.5 \pm 3.7	71.8	30.7 \pm 9.9	37.7 \pm 4.5	26.5 \pm 3.3
Whole body	96 \pm 15	100	260 \pm 70	100	100	100	100	100