



HAL
open science

Bioaccumulation of inorganic Hg by the juvenile cuttlefish *Sepia officinalis* exposed to ²⁰³Hg radiolabelled seawater and food

Thomas Lacoue-Labarthe, Michel Warnau, François Oberhänsli, Jean-Louis Teyssié, Paco Bustamante

► **To cite this version:**

Thomas Lacoue-Labarthe, Michel Warnau, François Oberhänsli, Jean-Louis Teyssié, Paco Bustamante. Bioaccumulation of inorganic Hg by the juvenile cuttlefish *Sepia officinalis* exposed to ²⁰³Hg radiolabelled seawater and food. *Aquatic Biology*, 2009, 6, pp.91-98. 10.3354/ab00172 . hal-00415807

HAL Id: hal-00415807

<https://hal.science/hal-00415807>

Submitted on 11 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Bioaccumulation of inorganic Hg by the juvenile cuttlefish *Sepia officinalis***
2 **exposed to ²⁰³Hg radiolabelled seawater and food**

3
4
5 Lacoue-Labarthe T.¹⁻², Warnau M.¹⁻², Oberhänsli F.², Teyssié J.-L.², Bustamante P.^{1*}

6
7
8 1- Littoral Environnement et Sociétés (LIENSs), UMR 6250, CNRS-Université de La
9 Rochelle, 2 rue Olympe de Gouges, F-17042 La Rochelle Cedex 01, France

10
11 2- International Atomic Energy Agency – Marine Environment Laboratories, 4 Quai Antoine
12 I^{er}, MC-98000 Principality of Monaco

13
14
15
16 * Corresponding author: Prof. Paco Bustamante

17 Littoral Environnement et Sociétés (LIENSs)

18 UMR 6250, CNRS-Université de La Rochelle

19 2 rue Olympe de Gouges

20 F-17042 La Rochelle Cedex 01

21 France

22
23 Phone: +33-546-507-625

24 Fax: +33-546-458-264

25 E-mail: pbustama@univ-lr.fr

26

27 **ABSTRACT:** Uptake and depuration kinetics of inorganic mercury (Hg) was investigated in
28 the juvenile common cuttlefish *Sepia officinalis* following exposures via seawater and food
29 using sensitive radiotracer technique (^{203}Hg). The cuttlefish readily concentrated ^{203}Hg when
30 exposed via seawater, with whole body CF > 260 after only 10d of exposure. The total Hg
31 accumulated from seawater was depurated relatively fast with a $T_{b_{1/2}}$ of 17d. During both
32 exposure and depuration periods, accumulated Hg was mainly (> 70%) associated with the
33 muscular parts of the cuttlefish. However, the proportion of the whole body Hg content
34 associated with the digestive gland increased during exposure and depuration phases,
35 suggesting that the metal was transferred from the muscles towards this organ for
36 detoxification. When fed with radiolabelled food, cuttlefish displayed high assimilation
37 efficiency (AE > 90%) and the metal was found to be mainly located in the digestive gland
38 (60% of the whole Hg content). Nevertheless, high depuration rates resulted in short $T_{b_{1/2}}$ (i.e.,
39 4 d), suggesting that this organ has a major role in Hg detoxification and depuration.
40 Whatever the exposure pathway was, low proportion of Hg (< 2%) was found in the
41 cuttlebone. Assessment of the relative contribution of the dietary and dissolved exposure
42 pathways to inorganic Hg bioaccumulation in juvenile cuttlefish revealed that Hg was mainly
43 accumulated from food that contributed for $77 \pm 16\%$ of the global metal bioaccumulation.

44

45

46 **Keywords:** mercury; bioaccumulation; kinetics; body distribution; cephalopod; relative
47 contribution.

48

49 INTRODUCTION

50 Mercury (Hg) is one of the metals of highest concern in the marine environment as it is
51 readily methylated by micro-organisms, bioaccumulates in marine biota and consistently
52 biomagnifies along the food chain (Cossa 1990). Among marine organisms, most of the
53 available information on Hg is related to fish, mainly because of their importance as a food
54 source for human. In fish, most of the Hg (i.e. > 95%) is methylated and is therefore
55 bioavailable for upper trophic levels (Bloom 1992). Hence, fish consumption is an important
56 source of Hg for human (Svensson et al. 1992) and is of particular health concern (Clarkson
57 1990).

58 In contrast to fish, information on Hg in cephalopods tissues is scarce despite the fact that
59 these molluscs represent an increasing component of the world fisheries (Boyle & Rodhouse,
60 2006, FAO 2007). In addition, information on Hg in cephalopod is essentially limited to the
61 main species targeted by fisheries and mainly reports metals levels in edible tissues, i.e.
62 mantle muscle, arms and fins (e.g. Buzina et al. 1989, Sapunar et al. 1989, Plessi et al. 2001).
63 Recently, a study on a large range of cephalopod species from the North Eastern Atlantic
64 waters reported that Hg concentrations varied on two orders of magnitude among cephalopod
65 species and that the metal was mainly stored under organic form in the muscular tissues
66 (Bustamante et al. 2006). Several other studies on cephalopods from the Mediterranean
67 suggest that these molluscs are able to accumulate high Hg concentrations in their tissues
68 (Renzoni et al. 1973, Rossi et al. 1993, Storelli & Marcotrigiano 1999). Various factors are
69 likely to influence Hg concentrations in cephalopods among which the size seems to be of
70 primary importance (Monteiro et al. 1992, Rossi et al. 1993, Pierce et al. 2008).

71 Although it was suggested that food would be the main source for Hg accumulation in
72 cephalopod tissues (Bustamante et al., 2006), the relative contribution of dietary and
73 waterborne pathways has not been assessed in cephalopods. Moreover, as cephalopods are

74 short-lived species, they might be interesting as short-term indicator species of the variation
75 of Hg concentrations in the environment (Seixas et al. 2005, Pierce et al. 2008).

76 For these reasons, the aim of this work was to investigate the biokinetics of Hg uptake and
77 depuration in cephalopods in order to better characterize its bioaccumulation, tissue
78 distribution and retention capacity. The common cuttlefish *Sepia officinalis* was selected as a
79 model to study Hg transfer in cephalopods from seawater and food.

80

81 **MATERIALS AND METHODS**

82 **Organisms and radiotracer**

83 Adult cuttlefish were collected by net fishing off Monaco in March and April 2006. They
84 were acclimated and maintained in open-circuit tanks in the IAEA-MEL premises. After
85 mating, the eggs laid by a single female were separated to optimise their oxygenation and kept
86 in a separate aquarium during the whole embryonic development (constantly aerated open
87 circuit; flux: 50 l h⁻¹; salinity: 37 p.s.u; temperature: 17 ± 0.5°C; pH: 8.0 ± 0.1; light/dark
88 cycle: 12 h/12 h). Hatching occurred approx. 50 d after the spawning. Young cuttlefish were
89 then maintained in the same aquarium and fed with brine shrimp (*Artemia* sp.) for 5 days
90 before the experiments.

91 The radiotracer, ²⁰³Hg [as ²⁰³HgNO₃; t_{1/2} = 46.59 d], was purchased from Isotope Product
92 Laboratory, USA. Stock solutions were prepared in 1 N nitric acid to obtain final radioactivity
93 allowing the use of spikes of only a few microliters (typically 5 µl).

94

95 **Experimental procedure**

96 **Contamination from seawater:** juveniles (n = 23; mean weight ± SD, 0.258 ± 0.009 g) were
97 placed for 10 d in a 20-l glass aquarium containing 0.45 µm filtered natural seawater
98 (constantly aerated closed circuit; temperature 17°C; 37 p.s.u.; light/dark cycle 12h/12h)

99 spiked with ^{203}Hg (0.6 kBq l^{-1}). In terms of stable metal, this concentration corresponded to 20
100 ng l^{-1} . To facilitate the recurrent counting of each individual during the experiment, the
101 juveniles were held individually in separate circular plastics boxes (10 cm diameter, 5 cm
102 height) covered up and down with a meshed plastic net to allow for free water circulation.
103 Radiotracer and seawater were renewed every second day to maintain water quality and
104 radiotracer activity constant. Radiotracer activities in seawater were checked before and after
105 each water renewal in order to determine the time-integrated radiotracer activities (Rodriguez
106 y Baena et al. 2006a, Warnau et al. 1996). Juveniles were separated in two groups: the first
107 group contained 16 tag-identified individuals and the second was composed of unidentified
108 animals for the body distribution analyses. At different time intervals, radiotracer activities
109 were counted in the same tag-identified juveniles ($n = 16$) all along the experiment.
110 According to their physiological states, the individuals should be removed from the
111 experiment leading the following sampling plan: $n=16$ from days 0 to 3 and $n = 14, 7, 6$ at
112 days 6, 9 10, respectively. In addition, after 3 and 9 d of exposure, 3 juveniles of the second
113 group were counted and dissected to determine the radiotracer distribution among the
114 digestive gland, the cuttlebone and the remaining tissues.
115 After this exposure period, the 6 remaining identified and radiolabelled juveniles ($n= 6$ from
116 days 0 to 4, 4 at day 6 and 1 at day 11) were held for 11 d in clean flowing water (open circuit
117 with constant aeration; seawater flux 50 l h^{-1} ; temperature 17°C ; 37 p.s.u.; light/dark cycle
118 12h/12h). At different time intervals during the depuration period, the same identified
119 juveniles ($n = 6$) were counted to establish the depuration kinetics of the radiotracer.

120

121 ***Contamination through the food:*** brine shrimp (*Artemia* sp.) were exposed for 5 d in a
122 plastic aquarium containing 4 l of natural seawater spiked with ^{203}Hg until pools of 25 brine

123 shrimp reached 3.1 kBq g^{-1} wet weight. The organisms were subsequently used as food for the
124 juvenile cuttlefish.

125 As detailed in the seawater experimental procedure, juveniles were separated in two groups of
126 identified and non-identified animals, which were devoted for the depuration kinetics and the
127 body distribution studies, respectively. Hence, sixteen newly hatched cuttlefish (mean weight
128 \pm SD, $0.297 \pm 0.011 \text{ g}$) were placed in individual plastic containers (10 cm diameter, 5 cm
129 height), and held in a 20-l aquarium under the same conditions as in the previous experiment.
130 Each juvenile was fed for 1 h with 25 of the previously radiolabelled *Artemia* sp. At the end
131 of the feeding period, the cuttlefish were immediately counted. From that time on, the
132 cuttlefish were fed twice a day with uncontaminated *Artemia* sp. for one month and regularly
133 counted to determine radiotracer depuration kinetics and assimilation efficiency. As
134 mentioned above, juveniles showing poor health condition were removed leading to a sample
135 number decrease along the experiment: $n = 16$ from days 0 to 2, 14 from days 3 to 14, 4 from
136 days 17 to 22 and 1 from days 24. Throughout the depuration period, faeces were removed
137 twice a day to reduce possible radiotracer recycling through leaching from the faeces. In
138 addition, after 3 h, 9 d, and 22 d of exposure, 3 others juveniles were counted and dissected to
139 determine the radiotracer distribution among the digestive gland, the cuttlebone and the
140 remaining tissues.

141

142 **Radioanalyses and data treatment**

143 Radioactivities were measured using a high-resolution γ -spectrometry system consisting of
144 four coaxial Germanium (N- or P-type) detectors (EGNC 33-195-R, Canberra[®] and Eurysis[®])
145 connected to a multi-channel analyzer and a computer equipped with a spectra analysis
146 software (Interwinner[®] 6). The detectors were calibrated with an appropriate standard for each
147 counting geometry used and measurements were corrected for background and physical decay

148 of the radiotracer. Counting times were adapted to obtain relative propagated errors less than
149 5% (Rodriguez y Baena, 2006b). They ranged from 10 min to 30 min for whole juveniles and
150 from 10 min to 24h for the dissected tissues.

151 Uptake of the ^{203}Hg from seawater was expressed as change in concentration factors (CF;
152 ratio between radiotracer content in the juvenile – Bq g^{-1} – and time-integrated activity in
153 seawater – Bq g^{-1}) along time (Warnau et al. 1996). Uptake kinetic was best described by a
154 saturation equation (Eq.1):

$$155 \quad \text{CF}_t = \text{CF}_{\text{ss}} (1 - e^{-k_e t}) \quad (\text{Eq. 1})$$

156 where CF_t and CF_{ss} are the concentration factors at time t (d) and at steady state, respectively,
157 k_e is the biological depuration rate constants (d^{-1}) (Whicker & Schultz 1982).

158 Radiotracer depuration kinetics were expressed in terms of change of percentage of remaining
159 activity (i.e., radioactivity at time t divided by initial radioactivity measured in the organisms
160 or in the tissue at the beginning of the depuration period * 100) along with time.

161 The depuration kinetic was best fitted by a mono-exponential equation (Eq. 2):

$$162 \quad A_t = A_0 e^{-k_e t} \quad (\text{Eq. 2})$$

163 where A_t and A_0 are the remaining activities (%) at time t (d) and 0, respectively, and k_e is the
164 biological depuration rate constant (d^{-1}). The determination of k_e allows the calculation of the
165 radiotracer biological half-life ($T_{b/2} = \ln 2 / k_e$). In the context of seawater and feeding
166 experiment, A_0 represents the absorption ($A_{0,w}$) and the assimilation (AE) efficiencies,
167 respectively.

168

169 **Bioaccumulation model**

170 The relative contribution of each uptake pathway was determined using the bioaccumulation
171 model originally proposed by Thomann (1981) and revised by Thomann et al. (1995) and
172 Metian et al. (2008). In this model, the total concentration of radiotracers in the juveniles, C_t

173 (Bq g⁻¹) is equal to the sum of each concentration resulting from the uptake by the different
174 pathways (Eq. 3):

$$175 \quad C_t = C_{f,ss} + C_{w,ss} \quad (\text{Eq. 3})$$

176 where $C_{f,ss}$ is the food-derived radiotracer concentration (Bq g⁻¹) in juveniles at steady state
177 (Eq. 4) and C_w is the water-derived radiotracer concentration (Bq g⁻¹) in juveniles at steady
178 state (Eq. 5).

$$179 \quad C_{f,ss} = (AE \times IR \times C_f) / k_{e,f} \quad (\text{Eq. 4})$$

$$180 \quad C_{w,ss} = (A_{0,w} \times k_{u,w} \times C_w) / k_{e,w} \quad (\text{Eq. 5})$$

181 where $A_{0,w}$ is the absorption efficiencies (%) of the radiotracer from seawater, AE is the
182 assimilation efficiency (%) of the radiotracer from food, C_f and C_w are the radiotracer
183 activities in food and seawater (Bq g⁻¹ and Bq ml⁻¹, respectively), respectively, IR is the
184 ingestion rate (g g⁻¹ d⁻¹), $k_{u,w}$ is the uptake rate constants (d⁻¹) from seawater, $k_{e,f}$ and $k_{e,w}$ are
185 the biological depuration rate constants (d⁻¹), for food and water pathways, respectively.

186 The relative contribution (%) of each uptake pathway is then assessed from the relation:

$$187 \quad \% \text{ food} = C_{f,ss} / (C_{f,ss} + C_{w,ss})$$

$$188 \quad \% \text{ seawater} = C_{w,ss} / (C_{f,ss} + C_{w,ss})$$

189

190 Constants (and their statistics) of the best fitting equations (decision based on ANOVA tables
191 for two fitted model objects) were estimated by iterative adjustment of the models using the
192 *nls* curve-fitting routine in R freeware. The level of significance for statistical analysis was
193 always set at $\alpha = 0.05$.

194

195 **Results**

196 **Contamination through seawater**

197 Uptake activity of ^{203}Hg in whole-body *S. officinalis* was best fitted by a saturation
198 exponential equation with a calculated CF_{ss} of 480 (Fig. 1; Table 1). The CF actually
199 measured at the end of the uptake period ($\text{CF}_{10\text{d}}$) of ^{203}Hg was 260 ± 70 (Table 2). Calculated
200 $\text{CF}_{10\text{d}}$ for the different organs indicated that ^{203}Hg was concentrated according to the following
201 decreasing order: digestive gland (1460 ± 480) > remaining tissues (290 ± 80) > cuttlebone
202 (47 ± 9). In terms of body distribution, ^{203}Hg was mainly found in the remaining tissues all
203 along the exposure period; this compartment accounted for 89 ± 3 and 80 ± 4 % of the whole
204 body load of ^{203}Hg after 3 d and 10 d of exposure, respectively (Table 2). The cuttlebone
205 presented very low Hg activity ($< 15 \text{ Bq g}^{-1}$) and loads ($< 3\%$) whereas the Hg activity in the
206 digestive gland was increasing with time, varying from $24 \pm 4 \text{ Bq g}^{-1}$ at day 3 to $520 \pm 170 \text{ Bq}$
207 g^{-1} at day 9 corresponding to 8% and 20% of the total radioactivity loads, respectively.

208

209 After the 10 d exposure period, non-contaminating conditions were restored and depuration
210 kinetic of ^{203}Hg was followed for 11 d. The whole-body depuration kinetic of ^{203}Hg in *S.*
211 *officinalis* was best described by a mono-exponential model (Fig. 2; Table 1). This result
212 indicated that 95% of the ^{203}Hg previously accumulated were depurated with a relatively
213 biological half-life relatively short (i.e. 17 d, Table 1). After 11 d of depuration conditions,
214 most of the ^{203}Hg body load was associated with the remaining tissues (72%) while the
215 digestive gland proportion increase up to 28% (Table 2).

216

217 **Contamination through food**

218 The depuration kinetic of ^{203}Hg ingested with food in *S. officinalis* was best fitted by a mono-
219 exponential model, characterized by a biological half-life of 4 d (Fig.2; Table 1) and allowed
220 an estimated assimilation efficiency of 91%.

221 During the depuration period, the digestive gland displayed highest proportion of the total
222 body burden of ^{203}Hg , i.e. 68, 60, 64 % after 3 h, 9 d and 22 d of depuration, respectively
223 (Table 2). The distribution of ^{203}Hg remained unchanged all along the loss experiment, with
224 the cuttlebone always showing the lowest proportion of the radiotracer (Table 2).

225

226 **Bioaccumulation model**

227 In order to assess the relative contribution of each uptake pathway to the global Hg
228 accumulation in *S. officinalis*, the different kinetic parameters obtained for seawater and food
229 experiments were used to feed the bioaccumulation model, along with other parameters such
230 as the ^{203}Hg concentration in seawater and food (C_w : 0.364 Bq ml^{-1} and C_f : 3100 Bq g^{-1} ,
231 respectively, present study) and the ingestion rate (IR: $0.07 \text{ g g}^{-1} \text{ d}^{-1}$, present study value being
232 congruent with IR value determined by Koueta et al., 1999). Modelling showed that food
233 represented the main pathway for ^{203}Hg bioaccumulation in the juvenile of cuttlefish,
234 contributing to $77 \pm 16\%$ of the global metal bioaccumulation vs. $23 \pm 14\%$ for the seawater
235 pathway.

236

237 **Discussion**

238 Cephalopods are an increasing marine resource for world fisheries (Boyle & Rodhouse 2006).
239 In the 1990s alone there was a 40% increase in squid catches worldwide (FAO 2007).
240 Cephalopods are well-known for their capacity to accumulate high levels of non-essential
241 metals, especially Ag and Cd, in their tissues (e.g. Martin & Flegal 1975, Bustamante et al.
242 1998, 2008). Hence, the intake of contaminants such as metals by human through cephalopod
243 consumption is a matter of concern (e.g. Pierce et al. 2008). Some studies also reported high
244 concentrations of Hg in cephalopods from areas naturally contaminated by cinnabar such the
245 Tyrrhenian and Adriatic Seas (e.g. Renzoni et al. 1973, Rossi et al. 1993, Storelli &

246 Marcotrigiano 1999). However, the dynamic of Hg incorporation and its metabolism in
247 cephalopods remain poorly understood. In the field, Hg could be methylated that generally
248 increases its toxicity as a result of its enhanced capacity of penetration across cell membranes
249 (Boudou et al. 1983). The known methylHg uptake pathways in the biota are the transfer from
250 the sediment where sediment-associated bacterial floras able to methylate part of the
251 inorganic Hg (Compeau & Bartha 1985), and the consumption of the prey such as juveniles of
252 shrimp, crab or fish where Hg is stored under the methylated form in variable proportions.
253 Considering that sand burying is a transient cryptic behaviour of cuttlefish (Poirier et al.
254 2004) and the grain size of the sediment selected by these species did not favour the
255 methylation of Hg, the inorganic form of the metal could also be a significant source of
256 accumulation for this nectobenthic cephalopod. In this study, uptake and depuration
257 biokinetics of Hg were determined using inorganic carrier-free ^{203}Hg in order to measure
258 metal fluxes in real time at environmentally realistic contaminant concentrations (Warnau et
259 al. 1996).

260 As a typical cephalopod, the common cuttlefish *Sepia officinalis* has high food intake
261 requirements to sustain its elevated growth rate. Being active predators, cephalopods have a
262 high digestion efficiency (Boucher-Rodoni et al. 1987) and food has been shown to constitute
263 an important source of uptake for various trace elements such as Am, Cd, Co and Zn (e.g.,
264 Guary & Fowler 1982, Koyama et al. 2000, Bustamante et al. 2002ab, 2004). However,
265 seawater could be an important bioaccumulation pathway as well, as elements can be taken up
266 efficiently through the skin and the gills. For instance, it has been shown that seawater is the
267 main intake pathway for Ag in adult *S. officinalis* (Bustamante et al. 2004). Therefore, there
268 was a need to provide insights on the bioaccumulation of Hg under controlled experimental
269 conditions in order to delineate the contribution of its uptake via the dissolved and dietary
270 pathways.

271 After 10 d of exposure to dissolved ^{203}Hg , cuttlefish displayed quite elevated whole-body
272 activities ($\text{CF} = 260 \pm 70$), indicating that Hg was efficiently bioconcentrated from seawater
273 (Table 2). The estimated steady-state equilibrium ($\text{CF}_{\text{ss}} = 480 \pm 150$) would be reached after 2
274 weeks of exposure. Among the three considered compartments, the digestive gland displayed
275 the highest concentration capacity ^{203}Hg ($\text{CF} = 1460 \pm 480$; Table 2) whereas the cuttlebone
276 showed the lowest CF (i.e. less than 50; Table 2). This pattern is very similar to that for
277 experimental data reported for other metals such as Ag, Cd, Co and Zn (Bustamante et al.
278 2002a, 2004). However, in terms of body burden distribution, ^{203}Hg was mainly stored in the
279 remaining tissues (up to 80 % of the total Hg load; Table 2), which are mainly composed of
280 muscles although they include the skin and the respiratory organs (i.e. gills) corresponding to
281 the tissues directly exposed to the contaminated seawater. Consistently, Bustamante et al.
282 (2006) reported that 70-90% of the total Hg body burden was stored in the muscular tissues in
283 different cephalopod species collected from the North East Atlantic. This could be explained
284 by the fact that 1) that these tissues represents more than 70% of the total body weight and 2)
285 that Hg has a stronger affinity for the sulphhydryl groups of muscular proteins rather than of fat
286 tissue found in fish and/or cephalopod (Bloom 1992, Bustamante et al. 2006).

287 When *Sepia officinalis* was exposed to dissolved Hg, the digestive gland contained 8% and
288 20% of the total Hg burden after 3 d and 10 d exposure, respectively (Table 2). Therefore, this
289 increasing Hg proportion found in this organ implied that the digestive gland accumulated
290 more efficiently than the others. In a second time, in depuration conditions, ^{203}Hg was release
291 of following a mono-exponential model and whole-body depuration was relatively rapid with
292 a $T_{b/2}$ of approx. 17 d. After 11 d of depuration under running seawater, the tissues in direct
293 contact with seawater contained significantly less ^{203}Hg than at the beginning of the
294 depuration period varying from $135 \pm 15 \text{ Bq g}^{-1}$ and to $95 \pm 17 \text{ Bq g}^{-1}$ at d 0 and d 11 of the
295 depuration period, respectively, whereas the digestive gland activity remained unchanged

296 (i.e., from $520 \pm 170 \text{ Bq g}^{-1}$ at day 0 to $560 \pm 30 \text{ Bq g}^{-1}$ at day 11; Table 2). These results
297 suggest that 1) the digestive gland shows a stronger retention capacity for Hg than the other
298 compartments, and/or that 2) a metal translocation occurred from the remaining tissues
299 towards the digestive gland. In cephalopods, the digestive gland plays obviously a major role
300 in the digestive processes but also in the detoxification of xenobiotics. Indeed, this organ has
301 already been shown to retain translocated metals from the other tissues (Bustamante et al.
302 2002a, 2004). The digestive gland is involved in the storage and detoxification of several
303 metals such as Ag, Cd, Cu, or Zn (e.g. Miramand & Bentley 1992, Bustamante et al. 2002b,
304 Dorneles et al. 2007) and persistent organic pollutants such as PCBs (Ueno et al. 2003, Danis
305 et al. 2005, Storelli et al., 2006). However, due to the fact this organ does not store Hg in
306 large amounts as shown in the field study (Bustamante et al. 2006), the digestive gland might
307 be also involved in the depuration of Hg when cuttlefish is exposed to the dissolved metal.
308 In the case of dietary exposure, the AE of Hg ingested with food was found to be nearly 100%
309 (Table 1). This high degree of Hg assimilation might be due to the very efficient digestive
310 metabolism that characterizes the juvenile cuttlefish (Mangold 1989). Indeed, this early life
311 stage is characterized by a predominant intracellular digestion process (compared to the
312 extracellular digestion which is dominant in adults) (Boucaud-Camou & Roper 1995), which
313 could favour the metal assimilation. Such extreme assimilation efficiency has already been
314 documented in cuttlefish: for instance, 90% and almost 100% AE for Cd and Co, respectively,
315 were reported in juvenile fed brine shrimp (Bustamante et al. 2002a, 2004). Nevertheless, the
316 assimilated Hg was rapidly depurated with a $T_{b/2}$ of 4 d, which suggests that the processes
317 governing Hg elimination are particularly efficient as well. Indeed, the ^{203}Hg activities in the
318 digestive gland dropped from $310 \pm 80 \text{ Bq g}^{-1}$ to $36 \pm 6 \text{ Bq g}^{-1}$ during the 22 d depuration
319 period after the feeding (Table 2). These results highlight the efficient excretion capacity of
320 the digestive gland for Hg. Because following dietary exposure the main fraction of the whole

321 Hg body burden was associated with this organ, it is not surprising that the retention time of
322 Hg following dietary exposure was 4 times lower than following seawater exposure, i.e. 4 d
323 vs. 17 d (Table 1).

324 In our experimental conditions, it appeared that the exposure of cuttlefish to contaminated
325 seawater lead to an accumulation of the Hg in the juvenile remaining tissues (> 80%) mainly
326 composed of muscular tissues. In the same time, Hg was translocated to the digestive gland
327 and subsequently eliminated from the organism. Following dietary exposure, inorganic Hg
328 was assimilated via the digestive gland and then rapidly eliminated. Consequently, the storage
329 and/or redistribution of the bioaccumulated metal towards the remaining tissues is limited in
330 cuttlefish. Nonetheless, considering both seawater and dietary exposure, food appears as the
331 predominant pathway for inorganic Hg bioaccumulation in juvenile cuttlefish. This result is
332 not surprising considering the relative high ingestion rate and the efficient digestive
333 metabolism of cephalopods (Lee 1994).

334

335

336 **Acknowledgements**

337

338 Authors are grateful to M. Metian (IAEA-MEL) for skilful assistance during the experiments.

339 MW is an Honorary Senior Research Associate of the National Fund for Scientific Research

340 (NFSR, Belgium). The IAEA is grateful for the support provided to its Marine Environment

341 Laboratories by the Government of the Principality of Monaco. This work was financially

342 supported by the IAEA and by LIENSs (University of La Rochelle).

343

344 **References**

345

346 Bloom NS (1992) On the chemical form of mercury in edible fish and marine invertebrate
347 tissue. *Can J Fish Aquat Sci* 49: 1010-1017

348 Boucaud-Camou E, Roper CFE (1995) Digestives enzymes in paralarval cephalopods. *Bull*
349 *Mar Sci* 57: 313-327

350 Boucher-Rodoni R, Boucaud-Camou E, Mangold K (1987) Feeding and digestion. In: Boyle
351 PR (ed) *Cephalopod Life Cycle*, Vol 2. Academic Press, London, p 85-108

352 Boudou A, Georgescauld D, Desmazes JP (1983) Ecotoxicological role of the membrane
353 barriers in transport and bioaccumulation of mercury compounds. In: Nriagu JO (ed)
354 *Aquatic Toxicology*, J. Wiley & Sons, New York, p 118-136

355 Boyle PR, Rodhouse PG (2006) *Cephalopods: Ecology and Fisheries*. Blackwell Science,
356 Oxford, United Kingdom

357 Bustamante P, Caurant F, Fowler SW, Miramand P (1998) Cephalopods as a vector for the
358 transfer of cadmium to top marine predators in the north-east Atlantic Ocean. *Sci Total*
359 *Environ* 220: 71-80

360 Bustamante P, Teyssié J-L, Fowler SW, Cotret O, Danis B, Miramand P, Warnau M (2002a)
361 Biokinetics of zinc and cadmium accumulation and depuration at different stages in the
362 life cycle of the cuttlefish *Sepia officinalis*. *Mar Ecol Prog Ser* 231: 167-177

363 Bustamante P, Cosson RP, Gallien I, Caurant F, Miramand P (2002b) Cadmium
364 detoxification processes in the digestive gland of cephalopods in relation to
365 accumulated cadmium concentrations. *Mar Environ Res* 53: 227-241

366 Bustamante P, Teyssié J-L, Danis B, Fowler SW, Miramand P, Cotret O, Warnau M (2004)
367 Uptake, transfer and distribution of silver and cobalt in tissues of the common cuttlefish
368 *Sepia officinalis* at different stages of its life cycle. *Mar Ecol Prog Ser* 269: 185-195

369 Bustamante P, Lahaye V, Durnez C, Churlaud C, Caurant F (2006) Total and organic Hg
370 concentrations in cephalopods from the North Eastern Atlantic waters: Influence of

371 geographical origin and feeding ecology. *Sci Total Environ* 368: 585-596

372 Bustamante P, Gonzalez AF, Rocha F, Miramand P, Guerra A (2008) Metal and metalloid
373 concentrations in the giant squid *Architeuthis dux* from Iberian waters. *Mar Environ Res*
374 66: 278-287

375 Buzina R, Suboticane K, Vukusic J, Sapunar J, Antonic K, Zorica M (1989) Effect of
376 industrial pollution on seafood content and dietary intake of total and methylmercury.
377 *Sci Total Environ* 78: 45-57

378 Clarkson TW (1990) Human health risks from methylmercury in fish. *Environ Toxicol Chem*
379 9:957-961

380 Compeau GC, Bartha R (1985) Sulphate-reducing bacteria: principal methylators of mercury
381 in anoxic estuarine sediment. *Appl Environ Microbiol* 50: 498-502

382 Cossa D, Thibaud Y, Roméo M, Gnassia-Barelli M (1990) Le mercure en milieu marin.
383 Biogéochimie et écotoxicologie. Rapport scientifiques et techniques de l'IFREMER,
384 N°19, Brest

385 Danis B, Bustamante P, Cotret O, Teyssié JL, Fowler SW, Warnau M (2005)
386 Bioaccumulation of PCBs in the cuttlefish *Sepia officinalis* from seawater, sediment and
387 food pathways. *Environ Pollut* 134: 113-122

388 Dorneles PR, Lailson-Brito J, dos Santos RA, Silva da Costa PA, Malm O, Azevedo AF,
389 Machado Torres JP (2007) Cephalopods and cetaceans as indicators of offshore
390 bioavailability of cadmium off Central South Brazil Bight. *Environ Pollut* 148: 352-359

391 FAO Fisheries and Aquaculture Information and Statistics Service (2007) Fisheries
392 commodities production and trade 1976-2005. FISHSTAT Plus - Universal software for
393 fishery statistical time series [online or CD-ROM]. Food and Agriculture Organization
394 of the United Nations. Available at:
395 <http://www.fao.org/fi/statist/FISOFT/FISHPLUS.asp>

396 Guary J-C, Fowler SW (1982) Experimental studies on the biokinetics of plutonium and
397 americium in the cephalopod *Octopus vulgaris*. Mar Ecol Prog Ser 7: 37-43

398 Koueta N, Boucaud-Camou E (1999) Food intake and growth in reared early juvenile
399 cuttlefish *Sepia officinalis* L. (Mollusca Cephalopoda). J Exp Mar Biol Ecol 240: 93-109

400 Koyama J, Nanamori N, Segawa S (2000) Bioaccumulation of waterborne and dietary
401 cadmium by oval squid, *Sepioteuthis lessoniana*, and its distribution among organs. Mar
402 Pollut Bull 40: 961-967

403 Lee PG (1994) Nutrition of cephalopods: fueling the system. In: Pörtner HO, O'Dor RK,
404 Macmillan DL (eds) Physiology of cephalopod molluscs: lifestyle and performance
405 adaptations. Gordon and Breach Science Publishers, Basel, p 35-51

406 Mangold K (1989) Reproduction, croissance et durée de vie. In: Grassé P-P (ed) Traité de
407 Zoologie : Céphalopodes, Vol V. Masson, Paris, p 493-552

408 Martin JH, Flegal AR (1975) High copper concentrations in squid livers in association with
409 elevated levels of silver, cadmium, and zinc. Mar Biol 30: 51-55

410 Metian M, Bustamante P, Cosson RP, Hédouin L, Warnau M (2008) Investigation of Ag in
411 the king scallop *Pecten maximus* using field and laboratory approaches. J Exp Mar Biol
412 Ecol 367: 53-60

413 Miramand P, Bentley D (1992) Concentration and distribution of heavy metals in tissues of
414 two cephalopods, *Eledone cirrhosa* and *Sepia officinalis* from the French coast of the
415 English Channel. Mar Biol 114: 407-414

416 Monteiro LR, Porteiro, FM, Gonçalves JM (1992) Inter- and intra-specific variation of
417 mercury levels in muscle of cephalopods from the Azores Arquipelago. Life Earth Sci
418 10: 13-22

419 Plessi M, Bertelli D, Monzani A (2001) Mercury and selenium content in selected seafood. J
420 Food Compos Anal 14: 461-467

421 Pierce GJ, Stowasser G, Hastie LC, Bustamante P (2008) Geographic, seasonal and
422 ontogenetic variation in cadmium and mercury concentrations in squid (Cephalopoda:
423 Teuthoidea) from UK waters. *Ecotoxicol Environ Saf* 70: 422-432

424 Poirier R, Chichery R, Dickel L (2004) Effects of rearing conditions on sand digging
425 efficiency in juvenile cuttlefish. *Behav Process* 67: 273-279

426 Renzoni A, Bacci E, Falciai L (1973) Mercury concentration in the water, sediments and
427 fauna of an area of the Tyrrhenian coast. *Revue Int Océan Méd* 31-32: 17-45

428 Rodriguez y Baena AM, Metian M, Teyssié JL, De Broyer C, Warnau M (2006a)
429 Experimental evidence for ^{234}Th bioaccumulation in three Antarctic crustaceans:
430 Potential implications for particle flux studies. *Mar Chem* 100: 354-365

431 Rodriguez y Baena AM, Miquel JC, Masqué P, Povinec PP, La Rosa J (2006b) A single vs.
432 double spike approach to improve the accuracy of ^{234}Th measurements in small-volume
433 seawater samples. *Mar Chem* 100: 269-281

434 Rossi A, Pellegrini D, Belcari P, Barghigiani C (1993) Mercury in *Eledone cirrhosa* from the
435 northern Tyrrhenian sea: Contents and relations with life cycle. *Mar Pollut Bull* 26:
436 683-686

437 Sapunar J, Jusic M, Bazulic D (1989) Total mercury and cadmium in some cephalopods and
438 fish from the Adriatic sea. *Mar Pollut Bull* 20: 576-577

439 Seixas S, Bustamante P, Pierce G (2005) Accumulation of mercury in the tissues of the
440 common octopus *Octopus vulgaris* (L.) in two localities on the Portuguese coast. *Sci*
441 *Total Environ* 340: 113-122

442 Storelli MM, Marcotrigiano GO (1999) Cadmium and total mercury in some cephalopods
443 from the South Adriatic Sea (Italy). *Food Addit Contam* 16: 261-265

444 Storelli MM, Barone G, D'Addabbo R, Marcotrigiano GO (2006) Concentrations and
445 composition of organochlorine contaminants in different species of cephalopod

446 molluscs from the Italian waters (Adriatic Sea). *Chemosphere* 64: 129-134

447 Svensson BG, Schütz A, Nilsson A, Åkesson I, Åkesson B, Skerfving S (1992) Fish as a
448 source of exposure to mercury and selenium. *Sci Total Environ* 126: 61-74.


449 Thomann RV (1981) Equilibrium model of fate of microcontaminants in diverse aquatic food
450 chains. *Can J Fish Aquat Sci* 38: 280–296.

451 Thomann RV, Mahony JD, Mueller R (1995) Steady-state model of biota sediment
452 accumulation factor for metals in two marine bivalves. *Environ Toxicol Chem* 14:
453 1989-1998

454 Ueno D, Inoue S, Ikeda K, Tanaka H, Yamada H, Tanabe S (2003) Specific accumulation of
455 polychlorinated biphenyls and organochlorine pesticides in Japanese common squid as
456 a bioindicator. *Environ Pollut* 125: 227-235


457 Warnau M, Teyssié J-L, Fowler SW (1996) Biokinetics of selected heavy metals and
458 radionuclides in the common Mediterranean echinoid *Paracentrotus lividus*: sea water
459 and food exposures. *Mar Ecol Prog Ser* 141: 83-94

460 Whicker FW, Schultz V (1982) *Radioecology: nuclear energy and the environment*, Vol 2.
461 CRC Press, Boca Raton, FL




462
 463
 464
 465
 466
 467
 468

Fig. 1: *Sepia officinalis*. Whole-body uptake kinetics of ^{203}Hg in juvenile cuttlefish exposed for 10 days to the radiotracer dissolved in seawater (concentration factors, CF; mean \pm SD, n=16 from days 0 to 3 and n = 14, 7, 6 at days 6, 9 10, respectively).



469



470
471

472 Fig. 2: *Sepia officinalis*. Whole-body loss kinetics of ^{203}Hg (% of remaining activity; mean \pm SD) in juvenile
 473 cuttlefish A) previously exposed to radiolabelled seawater for 10 days (n= 6 from days 0 to 4, 4 at day 6 and 1 at
 474 day 11); B) previously fed with radiolabelled brine shrimp (n = 16 from days 0 to 2, 14 from days 3 to 14, 4 from
 475 days 17 to 22 and 1 from days 24).

476 Parameters for the best fitting equations are given in Table 1.

Table 1. *Sepia officinalis*. Whole-body uptake and loss kinetic parameters of ^{203}Hg in whole cuttlefish following different exposure experiments:

1) individuals (n = 16) were exposed 10-d to the radiotracer in seawater then placed 11-d in depuration conditions (n = 6);

2) individuals fed on radiolabelled brine shrimp *Artemia* sp. were placed in depuration conditions for 30 d (n = 16);

Uptake parameters: CF_{ss} concentration and transfer factors at steady state; k_u : uptake rate constant (d^{-1})

All loss kinetics followed a mono-exponential depuration fit.

Depuration parameters: A_0 : activity (%) lost according to the exponential component; ASE: asymptotic standard error; k_e : depuration rate constant (d^{-1}), $T_{b/2}$: biological half-life (d); R^2 : determination coefficient of the uptake or loss kinetics; p: probability of the model adjustment:

*** and **: p-values < 0.001 and < 0.01, respectively.

Condition	Uptake			Loss			
	CF_{ss}	k_e	R^2	A_0 (SE)	k_e	$T_{b/2} \pm \text{SE}$ (d)	R^2
1) uptake seawater	480 ± 150	0.083 ± 0.036	0.703	-	-	-	-
2) loss seawater	-	-	-	95.3 (2.8) ***	0.041 ***	16.9 ± 3.9	0.472
3) loss food	-	-	-	90.7 (2.6) ***	0.180 ***	3.9 ± 0.3	0.804

1 Table 2. *Sepia officinalis*. Concentration Factors (mean CF \pm SD) and tissue distribution (mean % \pm SD) of ^{203}Hg during seawater and feeding
 2 experiments.

3

Compartments	Seawater contamination				Food contamination			
	Uptake (3d, n = 3)		Uptake (10d, n=3)		Loss (11d, n=1)	Loss (3h, n=3)	Loss (9d, n=3)	Loss (22d, n=3)
	Concentration Factor	Distribution (%)	Concentration Factor	Distribution (%)	Distribution (%)	Distribution (%)	Distribution (%)	Distribution (%)
Digestive gland	110 \pm 20	8.3 \pm 2.6	1460 \pm 480	19.7 \pm 3.8	28.2	67.9 \pm 10.6	59.9 \pm 3.4	63.6 \pm 5.8
Cuttlebone	29.7 \pm 3.7	2.8 \pm 1.0	47.0 \pm 8.8	<1	<1	1.4 \pm 0.9	2.4 \pm 1.9	9.9 \pm 3.2
Remaining tissues	110 \pm 10	88.9 \pm 3.4	290 \pm 80	79.5 \pm 3.7	71.8	30.7 \pm 9.9	37.7 \pm 4.5	26.5 \pm 3.3
Whole body	96 \pm 15	100	260 \pm 70	100	100	100	100	100