



HAL
open science

Phenoloxidase activation in the embryo of the common cuttlefish *Sepia officinalis* and responses to the Ag and Cu exposure

Thomas Lacoue-Labarthe, Paco Bustamante, Edouard Hörlin, Andrea Luna Acosta, Anne Bado-Nilles, Hélène Thomas-Guyon

► **To cite this version:**

Thomas Lacoue-Labarthe, Paco Bustamante, Edouard Hörlin, Andrea Luna Acosta, Anne Bado-Nilles, et al.. Phenoloxidase activation in the embryo of the common cuttlefish *Sepia officinalis* and responses to the Ag and Cu exposure. *Fish and Shellfish Immunology*, 2009, 27 (3), pp.516-521. 10.1016/j.fsi.2009.07.002 . hal-00415866

HAL Id: hal-00415866

<https://hal.science/hal-00415866>

Submitted on 11 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Phenoloxidase activation in the embryo of the common cuttlefish *Sepia***
2 ***officinalis* and responses to the Ag and Cu exposure**

3
4
5 T. Lacoue-Labarthe, P. Bustamante*, E. Hörlin, A. Luna-Acosta, A. Bado-Nilles[†], H.
6 Thomas-Guyon

7
8 Littoral, Environnement et Sociétés (LIENSs), UMR 6250, CNRS-Université de La
9 Rochelle, 2 rue Olympe de Gouges, F-17042 La Rochelle Cedex 01, France

10
11
12
13 (*) Corresponding author. Littoral, Environnement et Sociétés (LIENSs), UMR 6250,
14 CNRS-Université de La Rochelle, 2 rue Olympe de Gouges, F-17042 La Rochelle Cedex
15 01, France.

16 Tel: +33-546507625. Fax: +33-546458264. E-mail address: pbustama@univ-lr.fr (P.
17 Bustamante).

18
19 ^(†) Present address: AFSSA site de Ploufragan-Plouzané. Laboratoire d'études et de
20 recherches avicoles, porcines et piscicoles - Technopôle Brest-Iroise F-29 280 Plouzané,
21 France

22
23

24 **Abstract:** The prophenoloxidase (proPO) system catalysing the melanin production is
25 considered as implicated in the innate immune system in invertebrates. The phenoloxidase
26 (PO)-like activity was detected in the cuttlefish embryo sampled at the end of the
27 organogenesis and few hours before hatching. Various modulators of the PO activity were
28 used to assess the triggering of the proPO activating system. The results demonstrated the
29 evidence of a true phenoloxidase activity in the cuttlefish embryo. However, SDS and LPS
30 granted contrasting effects on the PO-like activity between the developmental stages
31 suggesting a progressive maturation of the proPO system from the embryonic to the
32 juvenile stages. In eggs exposed to dissolved trace metals all along the embryonic
33 development, Ag ($1.2 \mu\text{g L}^{-1}$) inhibited the PO-like activity in the cuttlefish embryo except
34 at hatching time, suggesting the synthesis of a new "juvenile" form of the phenoloxidase
35 enzyme. In similar conditions as for Ag, Cu ($230 \mu\text{g L}^{-1}$) stimulated and then inhibited the
36 PO-like activity according to a progressive metal accumulation within the egg and
37 suggesting the occurrence of a threshold, above which the toxicity of the essential metal
38 reduced the PO activity.

39

40

41 **Key words:** Cephalopod; Egg; Developmental Stage; Metal; Phenoloxidase; Modulation

42

43 1. Introduction

44

45 In invertebrates, the phenoloxidase (PO) is a Cu-dependent enzyme involved in the innate
46 defence mechanism, capable of catalyzing the oxidation of phenols such as of L-3, 4-
47 dihydroxyphenylalanine (L-Dopa) to quinones, which subsequently are transformed
48 nonenzymatically to a bactericide pigment, *i.e.* the melanin [1,2]. The prophenoloxidase
49 (proPO) system consists to the activation of the inactive proenzyme proPO to PO by
50 proteolytic cleavage [3] following the induction of a complex cascade by endogenous
51 activators or exogenous agents such as lipopolysaccharides (LPS), bacterial peptidoglycans
52 and β -1,3 glycans from fungi [2, 4]. Others compounds have been defined as elicitors, e.g.
53 trypsin, laminarin or SDS [4, 5, 6], or inhibitors, e.g. β -2-mercaptoethanol, sodium
54 diethylthiocarbonate (DETC) and tropolone [7, 8] of the proPO activating system.
55 Nevertheless, some modulators of the proPO activating system have been demonstrated as
56 specific to the species or the phylum probably linked with the various molecular
57 characteristics (*i.e.* size, amino-acids sequences) of the PO enzymes found among the
58 living being [9]. For instance, laminarin leads to the stimulation of the PO activity in the
59 bivalve *Crassostrea gigas* [6,7] whereas it has no effect in the ascidian *Styela plicata* [8].
60 In arthropods and molluscs, PO activity has been detected in the haemolymph [e.g. 10-13]
61 and has been described to be implicated in intra-cellular defence mechanisms [14]. This
62 PO activity has been attributed in some cases to another important family of type-3 copper
63 proteins, the haemocyanins (Hcs), Indeed, Hcs are structurally and phylogenetically related
64 to POs and in many organisms an intrinsic PO activity has been detected [for review, see
65 15], e.g. in cephalopods, the haemocyanin of the common octopus *Octopus vulgaris* [16-
66 17] and a sub-unit of the haemocyanin of the common cuttlefish *Sepia officinalis* [18]
67 exhibit intrinsic PO activity. While POs use one molecule of O₂ for chemical

68 transformation of phenolic compounds into quinones, Hcs are oxygen carriers that
69 transport O₂ in the body fluid [19]. However, because haemocyanins from several species
70 of arthropods and molluscs 1) exhibit phenoloxidase activity, 2) their basal PO activity can
71 be increased under certain artificial conditions such as limited proteolysis [20, 21],
72 interaction with lipids or antimicrobial peptides [22, 23] and denaturing agents like SDS
73 [9, 19] and because, in some cases, 3) endogenous components of the haemolymph can
74 lead to their activation [22, 24, 25], it has been suggested that the haemocyanin-derived
75 phenoloxidase activity participates in the immune response of arthropods and molluscs
76 [26]. Moreover, very few studies assessed the PO activity in early life stage of marine
77 organisms [e.g. 27, 28] but, to the best of our knowledge, none in the cephalopod embryo
78 and/or paralarvae. Nevertheless, in bivalves, no PO activity was detected in the larvae
79 homogenates of the Pacific oyster *Crassostrea gigas*, of the scallops *Argopecten*
80 *ventricosus* and *Nodipecten subnodosus* and of the Chinese pen-shell *Atrina maura* [6]
81 whereas PO activity was reported in the common mussel *Mytilus edulis* larvae, particularly
82 at the veliger stage [29]. More precisely, Thomas-Guyon et al. [30] recently observed that
83 the PO activity decreased from the “embryo 6h” to low value at the “larvae 11 day old”
84 stages in the Pacific oyster, and then increased to reach maximal value at the juvenile
85 stage. These results suggest that the proPO system could be progressively set off along the
86 early ontogenetic stages in molluscs.

87 When common cuttlefish mate in spring, females laid their eggs in the shallow waters
88 along the coast. These latter could be therefore subjected to various contaminants, which
89 are released from human activities in the coastal marine environment. Many pollutants
90 have been shown to modulate the PO activity in marine organisms. For example, PCBs
91 inhibit PO activity in the shrimp *Crangon crangon* [31], whereas fluoranthrene stimulates
92 the PO expression in *Mytilus edulis* [32]. Recently, Bado-Nilles et al. showed the

93 polycyclic aromatic hydrocarbons could induce a reduction of the PO activity in
94 *Crassostrea gigas* *in vitro* and *in vivo* [33]. Concerning trace metals, the proPO system
95 activation in the Norway lobster *Nephrops norvegicus* was blocked by manganese [34] and
96 *in vitro* experiments showed that mercury suppressed the L-Dopa transformation in the
97 haemolymph of *Crassostrea gigas* [13].
98 In the common cuttlefish *Sepia officinalis*, Ag, Co, Hg, Mn and Zn trace metals have
99 shown to penetrate through the eggshell during the developmental time [35, 36, 37].
100 Moreover, since haemocyanin/PO catalytic domains are copper-dependent [19, 26], *i.e.*
101 two copper atoms are essential to the oxygen binding in the catalytic active site, copper
102 and other metals may act as inhibitory surrogates by blocking oxygen and/or substrate
103 binding to the active site [38, 39]. Therefore, the PO-like activity of cuttlefish embryo
104 could be modulated in natural condition by these trace metals.
105 In this study, *in vivo* experiments on the detection and the modulation of the PO-like
106 activity have been conducted in the cuttlefish embryo at two developmental stages: *i.e.* the
107 end of the organogenesis occurring 32 days after the egg laying (d32) and a few hours
108 before hatching, *i.e.* after 50 days of development (d50). Several specific inhibitors and
109 activators of the PO activity were used to modulate the activation of the proPO system in
110 order to characterise the processes of PO activation and the potential evolution of the
111 proPO system during the embryonic development. Finally, PO-like activity was measured
112 from d32 to d50 in cuttlefish embryos from eggs exposed since the spawning time to one
113 toxic metal (Ag), which highly accumulate in embryonic tissues [40] and to one essential
114 (Cu) which is a co-factor of the PO enzyme.

115

116 **2. Materials and methods**

117 **2.1. Animals and experimental procedure**

118 The cuttlefish eggs were collected on pots from the west coast of Cotentin, France, by local
119 fishermen. Because pots were picked up every day, sampled eggs were laid in the previous
120 24-48 h. In the laboratory, eggs were separated for optimal oxygenation and placed into
121 floating sieves in a rearing structure as described by Koueta and Boucaud-Camou [41].
122 Few days after the collection, 700 eggs were randomly placed in tanks containing 11 L
123 seawater (constantly aerated closed circuit; temperature 17°C; 34 p.s.u.; light/dark cycle
124 12h/12h).

125 Cuttlefish eggs were then exposed to Ag ($2 \mu\text{g AgCl}_2 \text{ l}^{-1}$, *i.e.* $1.2 \mu\text{g Ag l}^{-1}$), Cu ($500 \mu\text{g}$
126 $\text{CuCl}_2 \text{ l}^{-1}$, *i.e.* $230 \mu\text{g Cu l}^{-1}$) and placed in control conditions all along their development
127 (50 d). All these conditions were made in duplicates. The control condition was repeated
128 for the Ag and Cu experiments. The Ag concentration was selected according to the
129 thresholds concentration defined by US-EPA [42] as the criterion for the protection of the
130 marine life (*i.e.*, $2.3 \mu\text{g l}^{-1}$). Regarding Cu, the tested concentration was chosen according
131 to the higher Cu concentrations recorded in the UK estuaries ($176 \mu\text{g.l}^{-1}$) [43], known as a
132 spawning grounds and nursery area for the cuttlefish from the English Channel [44].

133 Metals and seawater were renewed daily to maintain water quality and metal
134 concentrations constant. After one month, *i.e.* when development allowed distinguishing
135 and separating the egg compartments, eggs were sampled from control conditions for the
136 detection of the PO-like activity in the embryo ($n = 4$) and to test the effects of several
137 modulators on it ($n = 3$ for each modulator test). Finally, 2 and 3 eggs were collected at
138 different times from each duplicated tank (4 and 6 eggs per condition, respectively) to
139 determine the effect of metal on PO-like activity and the metal concentration in the
140 embryonic tissues, respectively. Eggs were weighed (wet wt), frozen in liquid nitrogen,
141 and stored at -80°C before further analyses. Non-contaminated eggs were also sampled for
142 the characterisation of PO-like activity assays.

143

144 **2.2. Phenoloxidase-like (PO-like) activity detection**

145 Eggs were dissected to remove the embryos from the others compartments, *i.e.* the
146 eggshell, the vitellus and the peri-vitelline fluid [45]. The embryos were weighted and
147 homogenized with mortar and pestle in cacodylate buffer (10 mM sodium cacodylate, 100
148 mM trisodium citrate, 0.45 M NaCl, 10 mM CaCl₂, 26 mM MgCl₂, pH 7.0) corresponding
149 to 8/1 v:v. This homogenate was then centrifuged (1500 g, 20 min, 4°C) and the resulting
150 supernatant was collected for PO-like activity measurements.

151 The detection of the PO-like activity was carried out by measurement of L-3,4-
152 dihydroxyphenylalanine (L-Dopa, C₉H₁₁NO₄, Sigma) transformation to dopachromes as
153 described by Thomas-Guyon et al. [30]. Transformation to dopachromes was monitored by
154 spectrophotometry at 490 nm. Samples were distributed in 96-well microplates. The assay
155 was run at 20°C. Control wells and negative control contained only 120 µl of CAC buffer
156 and 100 µl of CAC buffer plus 20 µl of L-Dopa, respectively.

157 The activation kinetic of the proPO system was determined by following the absorbance at
158 490 nm (Abs₄₉₀) at regular times during an incubation period of 125 h. Several well-known
159 modulators of the PO-like activity were then tested on embryos separated from one-month
160 old eggs and on embryos sampled a few hours before hatching: Lipopolysacharides
161 (*Escherichia coli* 0111: B4, 1 g L⁻¹), SDS (Sodium Dodecyl Sulfate, 6 mg mL⁻¹), purified
162 trypsin TPCK (N-Tosyl-L-phenylalanine chloromethyl ketone, 1 g. L⁻¹) as elicitors [4, 7].
163 Compounds such as β-2-mercaptoethanol (3,5 mM), Benzamidine (2,2 mM), EDTA
164 (Ethylenediamine Tetra-acetic acid, 5 mM), 100 % ethanol, PTU (N-Phenylthiourea, 20
165 mM), Tropolone (10 mM), and Zymosan A (1 g. L⁻¹) were also tested for their inhibitory
166 effects on the PO-like activity [7]. All chemicals were purchased from Sigma, France.

167 Sixty microliters of cacodylate buffer (CAC), 20 μL of a PO activity modulator, 20 μL of
168 L-Dopa (L-3,4-dihydroxyphenylalanine, 3 mg mL^{-1} , Sigma) and 20 μL of sample were
169 added in each well. Each sample was tested in triplicate wells and (Abs_{490}) was measured
170 after 96 h incubation period at room temperature.

171

172 **2.3. Metal concentrations in embryo tissue**

173 Embryo was dried at 70°C for two days and weighed (dry wt). Dried samples were
174 digested with 4 ml 65% ultrapur HNO_3 for several days at 100°C . After evaporation, the
175 residues were dissolved in 2 ml of 0.3 N ultrapur HNO_3 . Ag and Cu were determined by
176 flame and furnace atomic absorption spectrophotometry using a Z-5000 Hitachi
177 spectrophotometer with Zeeman background correction. Before use, all plastic- and
178 glassware were cleaned overnight with a mixture of 0.45 N HNO_3 and 0.9 N HCl in milli-
179 Ro quality water and rinsed 3 times with Milli-Q water. Blanks and certified reference
180 material (DOLT-3 CRM dogfish liver, NRCC) were treated and analysed in the same way
181 as the samples. Results for the CRM were in good agreement with certified values ($1.27 \pm$
182 0.01 vs. 1.20 ± 0.07 and 32.3 ± 1.4 vs. 31.2 ± 1.0 for Ag and Cu, respectively; Table 1).
183 Metal concentrations were expressed as $\mu\text{g g}^{-1}$ dry wt.

184

185 **2.4. Statistical analysis**

186 PO-like activity detection in all the samples was performed in triplicates. A one-way
187 analysis of variance (ANOVA) was applied to determine the differences among the
188 different developmental stages. A Kruskal-Wallis test was applied to determine the
189 significant differences among treatments and controls at different development time. P
190 values lower than 0.05 were used to identify significant differences.

191

192 **3. Results**

193 **3.1. Detection of PO-like activity in the cuttlefish embryo**

194 The transformation of the L-Dopa to dopachromes was followed during 120 h incubation
195 period (Fig. 1). Measures of Abs₄₉₀ increased progressively with incubation time and then
196 reached steady state equilibrium after 96 h suggesting that the proPO system was totally
197 activated after this incubation period. Therefore, PO-like activity values were considered in
198 this study after 96h of incubation with L-Dopa.

199 Table 2 shows the effects of different well-known inhibitors of the PO activity, which were
200 tested on embryo at the end of the embryogenesis (d32) and at the end of the embryonic
201 development (d50). PO-like activity was not modulated by benzamidine but totally
202 inhibited by EDTA, β-2-mercaptoethanol, PTU and Tropolone in all samples. Surprisingly,
203 SDS showed a contrasted effect between the two studied developmental stages. Indeed,
204 SDS totally suppressed Abs₄₉₀ at d32, whereas the PO-like activity few hours before
205 hatching were not significantly different from the control (69 ± 19 vs. 100 ± 13 %; $P >$
206 0.05).

207 A significant stimulation of PO-like activity was induced with the TPCK and Zymosan
208 (Table 3; $P < 0.05$) in the one-month embryo old (207 ± 53 and 154 ± 21 % of the control
209 activity, respectively). In embryos reaching the last developmental stage, the TPCK,
210 Zymosan and LPS enhanced significantly the PO-like activity (316 ± 190 , 167 ± 84 and
211 173 ± 51 %, respectively).

212

213 **3.2. Metal effects on PO-like activity**

214 The effect of metal was also tested on PO-like activity, which was measured in the control
215 embryos and in embryos exposed to $1.2 \mu\text{g Ag L}^{-1}$, from the end of the organogenesis
216 (d32) to the hatching time (d50) (Fig. 2). At 32, 36, 40 and 44 days of embryonic

217 development, the Abs₄₉₀ measured in the exposed embryos was 2-, 2-, 3-, and 4-fold
218 lower, respectively, than in the control. However, a few hours before hatching, PO-like
219 activity in controls decreased whereas it increased significantly in the exposed embryo
220 (0.64 ± 0.03 vs. 0.44 ± 0.04 in the Ag-exposed and control embryos, respectively; $P <$
221 0.05).

222 PO-like activity in the embryo exposed to Cu showed a two-phase kinetic compared to the
223 control group. Indeed, between d32 and d40, Abs₄₉₀ in the exposed-group were
224 significantly higher (d32 and d40; $P < 0.05$) or similar (d36; $P > 0.05$) than these measured
225 in the control embryos. Then, the PO-like activity in the Cu exposed embryos decreased
226 and remained lower compared to control values ($P < 0.05$) until hatching time. Statistical
227 analysis confirmed the effect of metal on PO-like activity detection.

228

229 **3.3. Metal concentration in embryos**

230 For both metals, exposed embryos showed higher concentrations compared to the control
231 (Table 1), 1) proving that the experimental procedure of contamination succeed and 2)
232 indicating that dissolved trace metals in seawater accumulated in the embryonic tissues
233 during the egg development. As a result of the dilution effect due to the increasing egg
234 weight of the embryo, metal concentrations decreased or remained stable between the end
235 of the embryogenesis (d32) and the end of the embryonic development (d50). However,
236 increasing Ag and Cu contents in the exposed embryo suggested a progressive metal
237 accumulation all along this development period.

238

239 **4. Discussion**

240

241 The present study revealed the detection of a PO-like activity during the post-
242 organogenesis period of the common cuttlefish embryo. Enzymes such as peroxidases may
243 act on the same substrates than POs. Therefore, the potential substrate of peroxidases and
244 specific inhibitor of PO activity, tropolone, was used in this study [7, 30]. The suppression
245 of Abs₄₉₀ in the wells by tropolone, confirmed that only PO-like activity was detected.
246 Moreover, additions of PTU, EDTA, or β-2-mercaptoethanol, leading to copper chelation,
247 induced a suppression of Abs₄₉₀. These results clearly show that the L-Dopa oxidation
248 was catalyzed by a Cu-dependant enzyme, *i.e.* the PO, and not by a Fe-dependant enzyme
249 such as a peroxidase [8].

250 The detection of the PO-like activity in the embryos of the common cuttlefish was
251 consistent with the results reported by Declair et al. [46] who demonstrated the appearance
252 of a brown colour in embryo haemolymph as a proof for a PO activity. More recently,
253 Siddiqui et al. [18] determined and localised a sub-unit of the haemocyanin responsible of
254 the PO activity in adult cuttlefish. In consistence with the present study, this sub-unit was
255 sensitive to *in vitro* inhibition by PTU. Thus, our results indicate that the cuttlefish embryo
256 is provided with enzymatic functions implicated in the innate defence system, at least from
257 the end of organogenesis.

258 The maximal absorbance observed after 96 h incubation period in embryo samples was
259 congruent with the pro-PO system activation after 96 h determined in the mussel *Mytilus*
260 *edulis* [47]. In the Pacific oyster *Crassostrea gigas*, the maximal PO activity in the
261 acellular fraction of the haemolymph was detected after a 21 h incubation time [7, 13, 30].
262 Thus, the delay observed in the cuttlefish embryo suggests the presence of an inactive form
263 of PO-like enzyme which is progressively activated [13]. Moreover, stimulation of the PO-
264 like activity by trypsin (TPCK) suggests that PO requires a proteolytic cleavage of the pro-
265 enzyme. Among other elicitors, *i.e.* zymosan, ethanol and LPS, described as specific of the

266 PO [4, 8, 48], only zymosan enhanced the PO-like activity in the embryos of both
267 developmental stages (*i.e.* d32 and d50). This result is in agreement with the proPO
268 activating system property in non-self recognition [15]. In haemocyanins, SDS induces a
269 minor conformational change leading to exposure of a phenoloxidase active site. In the
270 pro-PO system, activation is produced by proteases [9] *in vivo* and may be produced by
271 SDS *in vitro*, *e.g.* in the green mussel *Perna viridis*, in the crayfish *Panulirus interruptus*,
272 in the Pacific oyster *Crassostrea gigas* and in the shrimp *Penaeus californiensis* [4, 6, 12].
273 In our experimental conditions, SDS showed a contrasting effect, suppressing the PO
274 activity in the one-month old embryo. Nevertheless, this inhibition significantly decreased
275 with the embryonic development until hatching time. This result suggests that 1) SDS
276 blocks the active site of the PO in the one-month old embryos and 2) that the interaction
277 between SDS and PO enzyme decreases with the embryonic development, potentially
278 linked to a change of the PO protein form. Moreover, among the elicitors tested in the
279 present study, only LPS enhanced Abs₄₉₀ and this only in the embryo sampled a few hours
280 before hatching. Thus, the contrasting effect of SDS and LPS on PO-like activity between
281 both developmental stages (*i.e.* d32 and d50), suggests that the PO protein form
282 progressively changes with development. Indeed, although the cuttlefish development is
283 considered as direct [49], organs and tissues undergo a strong physiological maturation
284 from the last embryonic stages and the first month of the juvenile life [50]. Similar
285 observation was reported in the Pacific oyster *C. gigas* larvae, in which PO activity
286 increased at the end of the development probably due to a structural change of the PO
287 enzyme or of the active site [30]. Considering that a haemocyanin sub-unit was responsible
288 of the PO activity [18], Declair et al. [51] identified pre-haemocyanin at the embryonic
289 stage with higher oxygen-affinity than the juvenile one [52]. Thus, different haemocyanin
290 forms followed one another (*i.e.* eleven different protein fractions) progressively all along

291 the ontogenesis until the adult stage [53]. In this context, changes of the haemocyanin
292 forms, responsible of the PO-like activity during the embryonic development could be one
293 potential explanation for 1) the contrasting response of the PO to modulators between the
294 end of the organogenesis (embryonic Hc form) and the last stage until hatching (juvenile
295 Hc form), and 2) the inhibitor effect of EDTA in embryos whereas no effect was
296 highlighted in adult cuttlefish [18].

297 Few works studied the effect of trace metals on the PO-like activity in molluscs, as well *in*
298 *vivo* as *in vitro* conditions [e.g. 13, 54]. In this study, Ag repressed the spontaneous activity
299 in the embryo exposed to the metal. Indeed, Lacoue-Labarthe et al. [40] demonstrated that
300 Ag highly accumulated in the embryonic tissues from this stage and could lead to a toxic
301 effect on enzymatic functions. To the best of our knowledge, this is the first time that Ag
302 effect on the PO activity was assessed. Nevertheless, it is known that Ag could disturb, *in*
303 *vitro*, the activity of another Cu-dependant enzyme, *i.e.* the cathepsin from the cuttlefish
304 digestive gland [55]. Moreover, since haemocyanin/PO catalytic domains are copper-
305 dependent [19, 26], a biochemical interaction of Ag with molecular oxygen, the catalytic
306 active site and/or with the substrate may impede active site accessibility to oxygen and/or
307 the substrate [38, 39]. However, it is surprising that, at the last developmental stage, the
308 PO-like activity detected in exposed embryo was higher than the one found in the control
309 samples. This may arise for two reasons: 1) the new “juvenile” form of the haemocyanin
310 was predominant on the old “embryonic” one few hours before hatching [51, 53] and 2) its
311 activity was not altered by Ag exposition. Further studies should be carried out to
312 determine the effect of Ag exposure on the PO-like activity in juvenile cuttlefish and verify
313 the sensitivity of PO juvenile form to the metal. Following Cu-exposure, no effect or a
314 slight stimulation of the PO-like activity was observed between d32 and d40. In
315 cephalopod, Cu concentration in the egg varied slightly all along the embryonic

316 development [46]. Indeed, it is known that essential elements could be transferred from the
317 mother to the egg by metal incorporation into the vitellus [56, 57] and thus supplied the
318 needs of the embryo. Nevertheless, the higher Cu tissue burdens in the exposed embryo
319 suggest that Cu could progressively penetrate through the eggshell during the development
320 as shown for others essential metals such as Zn [58]. The behaviour of Cu could explain
321 the slight PO stimulation observed between d32 and d40, followed by an inhibition of the
322 enzyme activity when the metal reached a toxicity threshold in the embryonic tissues [54].

323

324 **Acknowledgments**

325

326 We are very grateful to Dr. N. Koueta from the LBBM of the University of Caen who
327 allowed the egg rearing experiment to be conducted in CREC at Luc sur Mer (France).

328 Authors thank MM. Lapie & Lapie, M. Guenon, M. Le Monnier and M. Longuet,
329 fisherman of the West-Cotentin coast, Normandy, France, who providing us the cuttlefish
330 eggs needed for this study. Authors are also grateful to M. Morinière and M. Maggiorani
331 of the Aquarium of La Rochelle for the technical assistance during the experiments.

332

333 **References**

334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381

- [1] Carballal MJ, Lopez C, Azevedo C, Villalba A. Enzymes involved in defense functions of hemocytes of mussel *Mytilus galloprovincialis*. *J Invertebr Pathol* 1997;70:96-105.
- [2] Söderhäll K, Cerenius L. Role of the prophenoloxidase-activating system in invertebrate immunity. *Curr Opin Immunol* 1998;10:23-28.
- [3] Smith VJ, Söderhäll K. A Comparison of phenoloxidase activity in the blood of marine invertebrates. *Dev Comp Immunol* 1991;15:251-261.
- [4] Asokan R, Arumugam M, Mullainadhan P. Activation of prophenoloxidase in the plasma and haemocytes of the marine mussel *Perna viridis* Linnaeus. *Dev Comp Immunol* 1997;21:1-12.
- [5] Bado-Nilles A, Thomas-Guyon H. Phenoloxidase activity in the haemolymph and haemocytes of the pacific oyster *Crassostrea gigas*. *Eur J Biochem* 2003;3:3-26.
- [6] Luna-Gonzalez A, Maeda-Martinez AN, Vargas-Albores F, Ascencio-Valle F, Robles-Mungaray M. Phenoloxidase activity in larval and juvenile homogenates and adult plasma and haemocytes of bivalve molluscs. *Fish Shellfish Immunol* 2003;15:275-282.
- [7] Hellio C, Bado-Nilles A, Gagnaire B, Renault T, Thomas-Guyon H. Demonstration of a true phenoloxidase activity and activation of a ProPO cascade in Pacific oyster, *Crassostrea gigas* (Thunberg) in vitro. *Fish Shellfish Immunol* 2007;22:433-440.
- [8] Arizza V, Cammarata M, Tomasino MC, Parrinello N. Phenoloxidase Characterization in Vacuolar Hemocytes from the Solitary Ascidian *Styela plicata*. *J Invertebr Pathol* 1995;66:297-302.
- [9] Decker H, Jaenicke E. Recent findings on phenoloxidase activity and antimicrobial activity of hemocyanins. *Dev Comp Immunol* 2004;28:673-687.
- [10] Söderhäll K, Smith VJ. Separation of the haemocyte populations of *Carcinus maenas* and other marine decapods, and prophenoloxidase distribution. *Dev Comp Immunol* 1983;7:229-239.
- [11] Johansson MW, Söderhäll K. Cellular immunity in Crustaceans and the proPO system. *Parasitol Today* 1989;5:171-176.
- [12] Hernandez-Lopez J, Gollas-Galvan T, Vargas-Albores F. Activation of the prophenoloxidase system of the brown shrimp (*Penaeus californiensis* Holmes). *Comp Biochem Physiol* 1996;113C:61-66.
- [13] Gagnaire B, Thomas-Guyon H, Renault T. In vitro effects of cadmium and mercury on Pacific oyster, *Crassostrea gigas* (Thunberg), haemocytes. *Fish Shellfish Immunol* 2004;16:501-512.

- 382 [14] Butt, D., Raftos, D. Phenoloxidase-associated cellular defence in the Sydney rock
383 oyster, *Saccostrea glomerata*, provides resistance against QX disease infections.
384 Dev Comp Immunol 2008; 32:299-306
385
- 386 [15] Cerenius L, Lee BL, Söderhäll K. The proPO-system: pros and cons for its role in
387 invertebrate immunity. Trends Immunol 2008;29:263-271.
388
- 389 [16] Salvato B, Santamaria M, Beltramini M, Alzuet G, Casella L. The enzymatic properties of
390 *Octopus vulgaris* hemocyanin: *o*-diphenol oxidase activity. Biochemistry
391 1008;37:14065-14077.
392
- 393 [17] Campello S, Beltramini M, Giordano G, Di Muro P, Marino SM, Bubacco L. Role of the
394 tertiary structure in the diphenol oxidase activity of *Octopus vulgaris* hemocyanin.
395 Arch Biochem Biophys 2008;471:159-167.
396
- 397 [18] Siddiqui NL, Akosung RF, Gielens C. Location of intrinsic and inducible phenoloxidase
398 activity in molluscan hemocyanin. Biochem Biophys Res Commun 2006;348:1138-
399 1144.
400
- 401 [19] Decker H, Tuzcek F. Tyrosinase/catecholase activity of hemocyanins: structural basis
402 and molecular mechanism, Trends Biochem Sci 2000;25:392–397.
403
- 404 [20] Decker H, Rimke T. Tarantula hemocyanin shows phenoloxidase activity, J Biol
405 Chem 1998;273:25889–25892.
406
- 407 [21] Siddiqui NI, Préaux G, Gielens C. Intrinsic and induced *o*-diphenoloxidase activity of
408 *b*-hemocyanin of *Helix pomatia*, Micron 2004;35:91–92.
409
- 410 [22] Nagai T, Osaki T, Shun-Ichiro K, Functional conversion of hemocyanin to
411 phenoloxidase by horseshoe crab antimicrobial peptides, J Biol Chem
412 2001;276:27166–27170.
413
- 414 [23] Jaenicke E, Decker H, Tyrosinases from crustaceans form hexamers, Biochem J
415 2003;371:515–523.
416
- 417 [24] Nagai T, Kawabata S. A link between blood coagulation and prophenoloxidase
418 activation in arthropod host defense, J Biol Chem 2000;275:29264–29267.
419
- 420 [25] Adachi K., Hirata T, Nishioka T, Sakaguchi M, Hemocyte components in crustacean
421 convert hemocyanin into a phenoloxidase-like enzyme, Comp Biochem Physiol
422 2003;134B: 135–141.
423
- 424 [26] Perdomo-Morales R, Montero-Alejo V, Perera E, Pardo-Ruiz Z, Alonso-Jimenez E.
425 Hemocyanin-derived phenoloxidase activity in the spiny lobster *Panulirus argus*
426 (Latreille, 1804), Biochim Biophys Acta 2008;1780:652
427
- 428 [27] Bai G, Brown JF, Watson C, Yoshino TP. Isolation and characterization of phenoloxidase
429 from egg masses of the Gastropod Mollusc, *Biomphalaria glabatra*. Comp Biochem
430 Physiol 1997;118B:463-469.
431

- 432 [28] Locci MT, Masetti M, Cecchettini A, Giorgi F. Cells released *in vitro* from the embryonic
433 yolk sac of the stick insect *Carausius morosus* (Br.) (Phasmatodea: Heteronemiidae)
434 may include embryonic hemocytes. *Int J Insect Morphol Embryol* 1998;27:325-331.
435
- 436 [29] Dyrzynda EA, Pipe RK, Ratcliffe NA. Host defence mechanisms in marine invertebrate
437 larvae. *Fish Shellfish Immunol* 1995;5:569-580.
438
- 439 [30] Thomas-Guyon H, Gagnaire B, Bado-Nilles A, Bouilly K, Lapègue S, Renault T.
440 Detection of phenoloxidase activity in early stages of the Pacific oyster *Crassostrea*
441 *gigas* (Thunberg). *Dev Comp Immunol* 2009, doi:10.1016/j.dci.2008.11.011
442
- 443 [31] Smith VJ, Johnston PA. Differential haemotoxic effect of PCB congeners in the common
444 shrimp, *Crangon crangon*. *Comp Biochem Physiol* 1992;101C:641-649.
445
- 446 [32] Coles JA, Farley SR, Pipe RK. Effects of fluoranthene on the immunocompetence of the
447 common marine mussel, *Mytilus edulis*. *Aquat Toxicol* 1994;30:367-379.
448
- 449 [33] Bado-Nilles A., Renault T., Gagnaire B., Auffret M., Le Floch S., Thomas-Guyon H. *In*
450 *vitro* and *in vivo* effects of polycyclic aromatic hydrocarbons on Pacific Oyster,
451 *Crassostrea gigas* (Thunberg), hemocyte parameters. *Comp Biochem Physiol* 2007;
452 146A, S71.
453
- 454 [34] Hernroth B, Baden SP, Holm K, André T, Söderhäll I. Manganese induced immune
455 suppression of the lobster *Nephrops norvegicus*. *Aquat Toxicol* 2004;70:223-231.
456
- 457 [35] Bustamante P, Teyssié J-L, Fowler SW, Cotret O, Danis B, Miramand P, Warnau M.
458 Biokinetics of zinc and cadmium accumulation and depuration at different stages in
459 the life cycle of the cuttlefish *Sepia officinalis*. *Mar Ecol Prog Ser* 2002;231:167-177.
460
- 461 [36] Bustamante P, Teyssié J-L, Danis B, Fowler SW, Miramand P, Cotret O, Warnau M.
462 Uptake, transfer and distribution of silver and cobalt in tissues of the common
463 cuttlefish *Sepia officinalis* at different stages of its life cycle. *Mar Ecol Prog Ser*
464 2004;269:185-195.
465
- 466 [37] Villanueva R, Bustamante P (2006) Composition in essential and non-essential elements
467 of early stages of cephalopods and dietary effects on the elemental profiles of *Octopus*
468 *vulgaris* paralarvae. *Aquaculture* 261:225-240.
469
- 470 [38] Tujula N, Radford J, Nair S V, Raftos D A. Effects of tributyltin and other metals on
471 the phenoloxidase activating system of the tunicate, *Styela plicata*, *Aquat Toxicol*
472 2001;55:191-201
473
- 474 [39] Floch C., Alarcon-Gutierrez E, Stéven C. Metal effects on phenoloxidase activities of
475 soils, *Ecotoxicol Environ Saf* 2009;72:108-114
476
477
- 478 [40] Lacoue-Labarthe T, Oberhänsli FR, Teyssié J-L, Warnau M, Koueta N, Bustamante P.
479 Differential bioaccumulation behaviour of Ag and Cd during the early development of
480 the cuttlefish *Sepia officinalis*. *Aquat Toxicol* 2008;86:437-446.
481

- 482 [41] Koueta N, Boucaud-Camou E. Food intake and growth in reared early juvenile cuttlefish
483 *Sepia officinalis* L. (Mollusca Cephalopoda). J Exp Mar Biol Ecol 1999;240:93-109.
484
- 485 [42] US-EPA. Ambient water quality criteria for silver. EPA-440/5-80-071. Office of water
486 regulations and standards. Criteria and Standards Division, US EPA, Washington, DC,
487 USA, 1980.
488
- 489 [43] Bryan GW, Langston WJ. Bioavailability, accumulation and effects of heavy metals
490 in sediments with special reference to United Kingdom estuaries. Environ Pollut
491 1992;76:89-131
492
- 493 [44] Boucaud-Camou E, Boismery J. The migrations of the cuttlefish (*Sepia officinalis* L)
494 in the English Channel. In: Boucaud-Camou E (ed) The Cuttlefish. Centre de
495 publication de l'Université de Caen, Caen 1991 ; pp 179-189
496
- 497 [45] Boletzky SV. Cephalopod eggs and egg masses. Oceanography and Marine Biology:
498 Annual Review 1998 ;36:341-371
499
- 500 [46] Declair W, Richard A. A study of the blood proteins in *Sepia officinalis* L. with special
501 reference to embryonic hemocyanin. Comp Biochem Physiol 1970;34:203-211.
502
- 503 [47] Coles JA, Pipe RK. Phenoloxidase activity in the haemolymph and haemocytes of the
504 marine mussel *Mytilus edulis*. Fish Shellfish Immunol 1994;4:337-352.
505
- 506 [48] Hernandez-Lopez J, Gollas-Galvan T, Gomez-Jimenez S, Portillo-Clark G, Vargas-
507 Albores F. In the spiny lobster (*Panulirus interruptus*) the prophenoloxidase is located
508 in plasma not in hemocytes. Fish Shellfish Immunol 2003;14:105-114.
509
- 510 [49] Lemaire J. Table de développement embryonnaire de *Sepia officinalis* L. (Mollusque
511 Céphalopode). Bull Soc Zool Fr 1970;95:773-782.
512
- 513 [50] Boucaud-Camou E, Yim M, Tresgot A. Feeding and digestion of young *Sepia officinalis*
514 L. (Mollusca: Cephalopoda) during post-hatching development. Vie Milieu
515 1985;35:263-266.
516
- 517 [51] Declair W, Lemaire J, Richard A. The differentiation of blood proteins during ontogeny in
518 *Sepia officinalis*. Comp Biochem Physiol 1971;40B:923-928.
519
- 520 [52] Wolf G, Verheyen E, Vlaeminck A, Lemaire J, Declair W. Respiration of *Sepia officinalis*
521 during the embryonic and early juvenile life. Mar Biol 1985 ;90:35-39
522
- 523 [53] De Wachter B, Wolf G, Richard A, Declair W. Regulation of respiration during the
524 juvenile development of *Sepia officinalis* (Mollusca: Cephalopoda). Mar Biol
525 1988;97:365-371.
526
- 527 [54] Pipe RK, Coles JA, Carissan FMM, Ramanathan K. Copper induced immunomodulation
528 in the marine mussel, *Mytilus edulis*. Aquat Toxicol 1999;46:43-54.
529
- 530 [55] Le Bihan E, Perrin A, Koueta N. Development of a bioassay from isolated digestive gland
531 cells of the cuttlefish *Sepia officinalis* L. (Mollusca Cephalopoda): effect of Cu, Zn,

- 532 and Ag on enzymes activities and cell viability. J Exp Mar Biol Ecol 2004;309:47-66.
533
- 534 [56] Lacoue-Labarthe T, Warnau M, Oberhänsli F, Teyssié J-L, Jeffree RA, Bustamante P.
535 First experiments on the maternal transfer of metals in the cuttlefish *Sepia officinalis*.
536 Mar Pollut Bull 2008;57:826-831.
537
- 538 [57] Craig S, Overnell J. Metals in squid, *Loligo forbesi*, adults, eggs, and hatchlings. No
539 evidence for a role for Cu- or Zn-metallothionein. Comp Biochem Physiol
540 2003;34C:311-317.
541
- 542 [58] Lacoue-Labarthe T, Martin S, Oberhänsli F, Teyssié J-L, Markich SJ, Jeffree R,
543 Bustamante P. Effects of increased pCO₂ and temperature on trace element (Ag, Cd
544 and Zn) bioaccumulation in the eggs of the common cuttlefish, *Sepia officinalis*,
545 Biogeosciences Discuss 2009 ; 6 :4865-4894.
546

547 Caption to figures


548

549 Fig 1. PO-like activity assessed by spectrophotometry in the embryo of *Sepia officinalis*
550 after 120 h of incubation. Values represent the average of four measures per egg (n= 4). a ≠
551 b ≠ c ≠ d; statistical differences for P < 0.05.

552

553 Fig 2. PO-like activity assessed by spectrophotometry in the embryo of *Sepia officinalis*
554 sampled in eggs exposed to (A) Ag and (B) Cu from the end of the organogenesis stage (d
555 32) to the hatching time (d 50). Values represent the average of six measures per egg
556 (n=4). * = statistical difference among treatments and controls for P < 0.05.

557




558


559

560

Figure 1



561



562
563
564

Figure 2

Table 1. Metal concentrations (mean \pm SD, $\mu\text{g g}^{-1}$ dwt; n=6) and metal content (mean \pm SD, ng for Ag and Cd, μg for Cu; n=6) in embryos from eggs sampled in non-contaminated condition (control) and exposed to $1.2 \mu\text{g l}^{-1}$ and $230 \mu\text{g l}^{-1}$ of dissolved Ag and Cu, respectively, at the end of the organogenesis (d32) and at the end of the development (d50). DOLT-3: Comparison of Ag and Cu certified concentrations (mean \pm SD, $\mu\text{g g}^{-1}$ dwt; n=5) in the certified reference material (dogfish liver DOLT-3, NRCC) with those obtained in the present study.

Embryo	Ag				Cu			
	Control		Exposed		Control		Exposed	
	Concentration	Content	Concentration	Content	Concentration	Content	Concentration	Content
End of the organogenesis (d32)	0.1 ± 0.03	0.8 ± 0.3	7.9 ± 5.1	21.5 ± 5.8	83 ± 1	1.0 ± 0.1	138 ± 20	1.9 ± 0.1
End of the development (d50)	0.1 ± 0.02	4.2 ± 1.6	2.0 ± 0.5	95.1 ± 40.0	38 ± 11	4.3 ± 0.9	145 ± 21	14.5 ± 1.0
DOLT-3								
Certified values			1.20 ± 0.07				31.2 ± 1.0	
Measured values			1.27 ± 0.01				32.3 ± 1.4	
References	Lacoue-Labarthe et al. (2008a)				Present study			

Table 2. PO-like activity (expressed as a percentage of the control activity; mean \pm SD; %) with different inhibitors in the embryo of the common cuttlefish *Sepia officinalis* at the end of the organogenesis (d32) and at the end of the embryonic development (d50).

Treatments	d 32		d 50	
	PO-like activity	Significance	PO-like activity	Significance
Control	100.0 \pm 6.2		100.0 \pm 13.3	
EDTA	0 \pm 0	***	0 \pm 0	***
β -mercaptoethanol	0 \pm 0	***	0 \pm 0	***
PTU	0 \pm 0	***	2.8 \pm 8.3	***
Tropolone	0 \pm 0	***	8.7 \pm 15.5	***
SDS	0 \pm 0	***	69.1 \pm 18.9	
Benzamidine	136.0 \pm 67.9		130.1 \pm 38.8	

Values represent the average of three measures per egg (n=3). *** Significance difference from control, P < 0.001.

Table 3. PO-like activity (expressed as a percentage of the control activity; mean \pm SD; %) with different elicitors in the embryo of *Sepia officinalis* at the end of the organogenesis (d32) and at the end of the embryonic development (d50).

Treatments	d 32		d 50	
	PO-like activity	Significance	PO-like activity	Significance
Control	100.0 \pm 6.2		100.0 \pm 13.4	
Zymosan	154.4 \pm 20.5	**	166.8 \pm 83.9	*
Ethanol	176.4 \pm 72.4		96.5 \pm 31.5	
TPCK	207.5 \pm 52.7	***	315.9 \pm 190.1	***
LPS	128.2 \pm 16.6		172.5 \pm 50.9	***

Values represent the average of three measures per egg (n=3). *** Significance difference from the control, P < 0.001.