



HAL
open science

Bursting bubbles

Henri Lhuissier, Emmanuel Villermaux

► **To cite this version:**

Henri Lhuissier, Emmanuel Villermaux. Bursting bubbles. *Physics of Fluids*, 2009, 21 (9), pp.091111. 10.1063/1.3200933 . hal-00426717

HAL Id: hal-00426717

<https://hal.science/hal-00426717v1>

Submitted on 31 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




FIG. 1.

Bursting bubbles

Henri Lhuissier¹ and Emmanuel Villermaux^{1,2}

¹IRPHE, Aix-Marseille Université, 13384 Marseille Cedex 13, France

²Institut Universitaire de France, France

(Received 28 May 2009; published online 11 September 2009)

[doi:10.1063/1.3200933]

A young bubble is a piece of bare spherical liquid shell of radius R , with uniform thickness h , formed from an air volume rising underneath a water pool. It bursts by nucleating a hole, opening at the constant tangential velocity $V = \sqrt{2\sigma/\rho h}$ balancing inertia with surface tension forces, and collecting liquid in its rim. The centripetal acceleration $\gamma = V^2/R$ exerted on the rim perpendicular to the shell surface induces its Rayleigh–Taylor destabilization whose wavelength $\lambda_{\perp} \sim \sqrt{\sigma/\rho\gamma}$ scales as the geometrical mean of the only two length scales characterizing the bubble,

$$\lambda_{\perp} \sim \sqrt{Rh}. \quad (1)$$

Examples shown here include: [Fig. 1(a)] a snapshot of the rim instability and ligaments expulsion from the full

bursting sequence [Fig. 1(b)] of a $R=10$ mm radius bubble. $\Delta t=2$ ms between frames. Two water bubbles with distinct thicknesses h lead to different wavelengths λ_{\perp} : [Fig. 1(c)] $R=14$ mm, $\Delta t=5$ ms between frames and $\lambda_{\perp}=2$ mm; and [Fig. 1(d)] $R=11.5$ mm, $\Delta t=1.33$ ms between frames and $\lambda_{\perp}=0.8$ mm.

Ligaments emerge from the rim unstable crests and are stretched by centrifugation. They ultimately break by a capillary instability setting the resulting drop size distribution in the spray, as seen in Fig. 1(e). The distribution width is fixed by the relative ligament radius corrugations.¹ The drops thus formed are called “film drops.”² They originate in nature from bubbles entrained below breaking waves,³ and are believed to contribute significantly to the net water evaporation from the sea.⁴

¹E. Villermaux, “Fragmentation,” *Annu. Rev. Fluid Mech.* **39**, 419 (2007).

²D. C. Blanchard and L. D. Sysdek, “Film drop production as a function of bubble size,” *J. Geophys. Res.* **93**, 3649, DOI:10.1029/JC093iC04p03649 (1988).

³G. B. Deane and D. Stokes, “Scale dependence of bubble creation mechanisms in breaking waves,” *Nature (London)* **418**, 839 (2002).

⁴E. L. Andreas, J. B. Edson, E. C. Monahan, M. P. Rouault, and S. D. Smith, “The spray contribution to net evaporation from the sea: A review of recent progress,” *Boundary-Layer Meteorol.* **72**, 3 (1995).