



HAL
open science

The new molecular biology of granulosa cell tumors of the ovary.

N. Kalfa, Ra Veitia, Ba Benayoun, B. Boizet-Bonhoure, C. Sultan

► **To cite this version:**

N. Kalfa, Ra Veitia, Ba Benayoun, B. Boizet-Bonhoure, C. Sultan. The new molecular biology of granulosa cell tumors of the ovary.. *Genome Medicine*, 2009, 1 (8), pp.81. 10.1186/gm81 . hal-00429361

HAL Id: hal-00429361

<https://hal.science/hal-00429361>

Submitted on 3 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Minireview

The new molecular biology of granulosa cell tumors of the ovary

Nicolas Kalfa*†, Reiner A Veitia‡§, Bérénice A Benayoun‡§, Brigitte Boizet-Bonhoure¶
and Charles Sultan**

Addresses: *Service d'Hormonologie, Hôpital Lapeyronie, CHU Montpellier et UM1, Montpellier, France. †Service de Chirurgie Pédiatrique, Hôpital Lapeyronie, CHU Montpellier, Montpellier, France. ‡Institut Jacques Monod, CNRS-UMR 7592, Bâtiment Buffon, 15 Rue Hélène Brion, 75013 Paris, France. §Université Paris Diderot, Paris 7, 75013 Paris, France. ¶Institut de Génétique Humaine, CNRS UPR1142, Montpellier, France. **Unité d'Endocrinologie-Gynécologie Pédiatriques, CHU Montpellier et UM1, Montpellier, France

Correspondence: Charles Sultan. Email: c-sultan@chu-montpellier.fr

Abstract

Granulosa cell tumors (GCTs) of the ovary belong to the group of ovarian sex-cord stromal tumors and represent 5 to 10% of ovarian malignancies. GCTs exhibit several morphological, biochemical and hormonal features of normal proliferating pre-ovulatory granulosa cells, such as estrogen biosynthesis. Prognostic factors of this condition are lacking, and alternative treatment options to preserve future fertility are needed. Several groups have shown that two genetic factors implicated in GCTs are of particular interest. The *gsp* oncogene is a constitutive activating mutation of the *Gsα* subunit and is correlated with the prognosis of the tumor. *FOXL2* is a transcription factor gene involved in ovarian development and function, whose expression is reduced and which is mutated in the majority of GCTs. *FOXL2* appears to play a major role in cell cycle regulation. These recent findings open new pathophysiological insights into GCT development as well as revisitation of granulosa cell and ovarian function.

Natural history of granulosa cell tumors of the ovary

Granulosa cell tumors (GCTs) of the ovary are relatively uncommon neoplasms, representing approximately 5 to 10% of all ovarian malignancies [1,2]; the others are germ cell tumors (teratomas and yolk sac tumors, 60 to 70%) and epithelial adenomas (10 to 20%) [3]. GCTs of the ovary belong to the group of ovarian sex-cord stromal tumors. Other tumors in this class are thecoma-fibromas, Sertoli cell tumors, sex cord tumors with annular tubules, and gynandroblastomas. The majority of patients with GCTs are adults, but 5% are pre- or peri-pubertal [4]. Consequently, two histological types of GCTs can be distinguished on a histopathological basis: juvenile and adult. GCTs exhibit several morphological, biochemical and hormonal features of normal proliferating pre-ovulatory granulosa cells, including both estrogen and inhibin biosynthesis [5]. This tumoral hyperestrogenism induces precocious puberty in children (premature breast development, vaginal bleeding)

and advanced growth and bone maturation. The prognosis of juvenile GCTs is excellent overall: tumor recurrence and metastasis are rare and usually occur early [6]. On the other hand, adult GCTs are low-grade indolent malignant neoplasms that display a significant propensity for recurrence and metastasis, and they may cause menorrhagia or intermenstrual bleeding. The cornerstone of treatment remains surgery. The disease in young patients is often confined to one ovary and thus, in order to preserve fertility, a unilateral salpingo-oophorectomy is preferable. In the case of postmenopausal women, a total abdominal hysterectomy and bilateral salpingo-oophorectomy should be performed [7].

Challenges in GCT management

Nevertheless, management of patients with ovarian GCTs needs to be improved. First, the recurrent or metastatic tumor may manifest many years after removal of the primary neoplasm, with intervals of 10 or even 20 years being not uncommon, especially in the adult disease [8,9]. Clinical follow-up of these patients is thus critical, but prognostic factors are lacking and assessment of the recurrence risk remains imprecise. For instance, in International Federation of Obstetricians and Gynecologists (FIGO) stage I disease (that is, intra-ovarian tumor with no extension, the most frequent stage at diagnosis according to FIGO), no association between outcome and the clinical signs or tumor histology (mitotic, nuclear atypia, ploidy, Ki67 expression) has been undisputedly demonstrated [10,11]. Furthermore, in more advanced disease, aggressive debulking surgery and postoperative chemotherapy may be required [8]. Alternative treatment options and the use of molecular markers are thus necessary, especially for young women whose fertility should be preserved.

FOXL2 and GCTs of the ovary

FOXL2 is a recent candidate gene in the pathophysiology of GCTs and could prove useful to evaluate their prognosis.

FIGO, International Federation of Obstetricians and Gynaecologists; FOXL2, forkhead transcription factor FOXL2; FSH, follicle-stimulating hormone; GCT, granulosa cell tumor; PKA, cAMP-dependent protein kinase; RFLP, restriction fragment length polymorphism; WT1, Wilms' tumor suppressor gene.

Indeed, *FOXL2* is a winged helix/forkhead transcription factor gene involved in ovarian development and function [12,13]. *FOXL2* ovarian expression in mammals starts before the morphological differentiation of the gonad is recognizable, and persists until adulthood, mainly in granulosa cells [14,15]. *FOXL2* is thought to be a key factor in the early development and maintenance of the vertebrate ovary [12,15]. Its expression in granulosa cells is suppressed, or at least lowered, in the majority of juvenile GCTs, particularly in those with the most aggressive pattern of progression [16]. Similar expression studies have yet to be performed in adult GCTs.

Interestingly, Shah *et al.* [17] recently studied four adult GCTs using whole-transcriptome paired-end RNA sequencing. After removing the previously described germline genetic variants and non-specific mutations present in non-GCTs, the authors identified the somatic missense point mutation c.402C→G (p.C134W) in all four specimens. Subsequently, they used direct sequencing, restriction fragment length polymorphism analyses (RFLPs), and a real-time PCR-based allelic discrimination assay to genotype *FOXL2* in a large collection of tumors. The p.C134W mutation was found in 97% of 89 additional adult GCTs studied, and was detected in only one case out of eight juvenile GCTs. The mutation was also found in 21% of thecomas, that is, 3 out of 14, in the study of Shah *et al.* Further studies are needed to determine the link between this mutation and GCTs, and whether this mutation alone is sufficient to induce tumor progression.

Other genetic factors implicated in ovarian GCTs

Genetic defects in other signaling pathways have been investigated in GCTs. Although Peutz-Jeghers syndrome is associated with an increased risk for GCTs, neither allele loss at the disease locus (19p13.3) nor mutations in the *LKB1* gene are associated with sporadic GCTs [18,19]. Similarly, despite the association of epithelial ovarian cancer with the familial breast cancer genes *BRCA1* and *BRCA2*, sporadic mutations have not been reported for GCTs.

Other interesting candidate pathways are those involving the mitogen-activated protein kinases [20] and growth factors, because they are known to induce increased oncogene functioning or a loss-of-anti-oncogene function in several solid tumors. However, King *et al.* [21] failed to identify any prognostic value for the oncogenes *c-myc*, *p21-ras*, *c-erb B2*, and *p53* in a group of 40 GCTs. Similarly, the Wilms' tumor suppressor gene (*WT1*) and *TP53* genes, which play a role in follicular development, have been tested [22]. *WT1* is expressed in GCT, but neither mutations nor loss of heterozygosity have been identified [23]. *TP53* mutations are frequent in numerous cancers, especially in epithelial tumors of the ovary, but there is neither hyperexpression [24] nor mutation [25] of this gene in GCT.

The role of the follicle-stimulating hormone (FSH) signaling pathway is also strongly suspected in GCTs for two reasons: firstly, normal granulosa cells are under the control of FSH and, secondly, the gene expression profile of GCTs is consistent with a constitutive activation of FSH signaling [26]. However, there is no evidence of activating mutations in the gene encoding the FSH receptor [27-29]. FSH signaling involves the coupling of heterotrimeric G proteins to activate intracellular second messenger systems, mainly the cAMP-dependent protein kinase (PKA).

Conflicting data have been reported regarding the presence of a constitutive activating mutation of the G α subunit in GCTs, the so-called *gsp* oncogene. It has rarely been described in adult GCTs [30,31], but our group has identified it frequently in the juvenile type, where mutations in hot spot position 201 (p.R201C or p.R201H) were found in 30% of patients [32]. The precise role of this mutation in the transformation of ovarian cells into malignant cells remains unclear, but it has been demonstrated in other tissues that the rate of cell proliferation and invasiveness can be influenced by the constitutive activation of G α s [33]. The oncological stages are indeed significantly different according to the *gsp* oncogene status. Patients with a hyperactivated G α exhibit significantly more advanced tumors, and almost 80% of *gsp*-positive patients present with an extra-ovarian extension (stage Ic) or have a recurrence [32].

Potential diagnostic and therapeutic applications

The clinical applications of these findings may be promising in the near future, for several reasons. Firstly, identification of a recurrent mutation in *FOXL2* may be used as another diagnostic tool for adult GCTs, in addition to the classical pathological and immunohistochemical features. It may be particularly useful to explore granulosa tumor participation in heterogeneous ovarian non-germinal tumors with components of different origins. Also, the low frequency of the *FOXL2* mutation in juvenile GCTs compared with adult cases, as reported by Shah and collaborators [17], supports the distinction made between the juvenile type and the adult type based on clinical presentation, pathological features and natural progression [1]. Clearly, differentiating the two types of tumor is clinically relevant, since relapses of adult GCTs are more frequent and may occur later than juvenile GCTs. The mutational or expressional status of *FOXL2* could thus be of importance in adapting the length of postoperative follow-up. Another genetic abnormality that could be critical in the follow-up of these patients is the *gsp* oncogene status, especially for juvenile GCTs. A constitutively activated G α s is indeed involved in the natural history of the tumor and is a prognostic factor of its extension. Finally, the finding of both extinction [16] and mutation [17] of *FOXL2* in GCTs highlights the role of this

gene in the regulation of cell proliferation. These findings increase the body of evidence implicating the *FOX* family genes, whether as oncogenes or anti-oncogenes, in malignant processes such as rhabdomyosarcomas [34], secondary acute myeloblastic leukemia [35], and laryngeal, lung, breast [36-38] and pancreatic cancers [39]. *FOXL2* may act as a transcriptional regulator and a coordinator of SMAD3 downstream targets [40,41] that, with SMAD2 and the TGF- β superfamily ligands, regulate granulosa cell proliferation [42]. These recently identified steps could be the targets of the next generation of therapies.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We would like to acknowledge Dr and Pr F Paris, Unité d'endocrinologie pédiatrique, Hôpital Aranud-de-Villeneuve, CHU Montpellier, France; P Philibert and A Ecochard, Service d'Hormonologie, Hôpital Lapeyronie, CHU Montpellier, France; C Patte, Département d'Oncologie, Institut Gustave Roussy, Villejuif, France; E Thibaud, Service d'Endocrinologie Pédiatrique, Hôpital Necker, Paris, France; C Pienkowski, Service d'Endocrinologie Pédiatrique, Hôpital des Enfants, CHU Toulouse, France; M Fellous, Génétique Humaine, Inserm 709, Université Paris 7, Hôpital Cochin, Paris, France; and F Jaubert, Service d'anatopathologie, Hôpital Necker, Paris, France, for their contribution in this work. This work was supported by a national grant from the Association de Recherche contre le Cancer (ARC: JR/MLD/MDV-P05/5).

References

- Schumer ST, Cannistra SA: **Granulosa cell tumor of the ovary.** *J Clin Oncol* 2003, **21**:1180-1189.
- Colombo N, Parma G, Zanagnolo V, Insinga A: **Management of ovarian stromal cell tumors.** *J Clin Oncol* 2007, **25**:2944-2951.
- Merras-Salmio L, Vettenranta K, Mottonen M, Heikinheimo M: **Ovarian granulosa cell tumors in childhood.** *Pediatr Hematol Oncol* 2002, **19**:145-156.
- Young RH, Dickersin GR, Scully RE: **Juvenile granulosa cell tumor of the ovary. A clinicopathological analysis of 125 cases.** *Am J Surg Pathol* 1984, **8**:575-596.
- Fuller PJ, Chu S, Fikret S, Burger HG: **Molecular pathogenesis of granulosa cell tumours.** *Mol Cell Endocrinol* 2002, **191**:89-96.
- Kalfa N, Patte C, Orbach D, Lecointre C, Pienkowski C, Philippe F, Thibault E, Plantaz D, Brauner R, Rubie H, Guedj AM, Ecochard A, Paris F, Jeandel C, Baldet P, Sultan C: **A nationwide study of granulosa cell tumors in pre- and postpubertal girls: missed diagnosis of endocrine manifestations worsens prognosis.** *J Pediatr Endocrinol Metab* 2005, **18**:25-31.
- Evans AT, 3rd, Gaffey TA, Malkasian GD, Jr., Annegers JF: **Clinicopathologic review of 118 granulosa and 82 theca cell tumors.** *Obstet Gynecol* 1980, **55**:231-238.
- Pectasides D, Papaxoinis G, Fountzilias G, Aravantinos G, Pectasides E, Mouratidou D, Economopoulos T, Andreadis C: **Adult granulosa cell tumors of the ovary: a clinicopathological study of 34 patients by the Hellenic Cooperative Oncology Group (HeCOG).** *Anticancer Res* 2008, **28(2B)**:1421-1427.
- Hasiakos D, Papakonstantinou K, Karvouni E, Fotiou S: **Recurrence of granulosa cell tumor 25 years after initial diagnosis. Report of a case and review of the literature.** *Eur J Gynaecol Oncol* 2008, **29**:86-88.
- Leuvenink EM, Brennan BA, Crook ML, Doherty DA, Hammond IG, Ruba S, Stewart CJ: **Prognostic value of mitotic counts and Ki-67 immunoreactivity in adult-type granulosa cell tumour of the ovary.** *J Clin Pathol* 2008, **61**:914-919.
- Kalfa N, Sultan C: **Juvenile ovarian granulosa cell tumor: a benign or malignant condition?** *Gynecol Endoc* 2009, **25**:299.
- Schmidt D, Ovitt CE, Anlag K, Fehsenfeld S, Gredsted L, Treier AC, Treier M: **The murine winged-helix transcription factor Foxl2 is required for granulosa cell differentiation and ovary maintenance.** *Development* 2004, **131**:933-942.
- Uda M, Ottolenghi C, Crisponi L, Garcia JE, Deiana M, Kimber W, Forabosco A, Cao A, Schlessinger D, Pilia G: **Foxl2 disruption causes mouse ovarian failure by pervasive blockage of follicle development.** *Hum Mol Genet* 2004, **13**:1171-1181.
- Cocquet J, Pailhoux E, Jaubert F, Servel N, Xia X, Pannetier M, De Baere E, Messiaen L, Cotinot C, Fellous M, Veitia RA: **Evolution and expression of FOXL2.** *J Med Genet* 2002, **39**:916-921.
- Cocquet J, De Baere E, Gareil M, Pannetier M, Xia X, Fellous M, Veitia RA: **Structure, evolution and expression of the FOXL2 transcription unit.** *Cytogenet Genome Res* 2003, **101**:206-211.
- Kalfa N, Philibert P, Patte C, Ecochard A, Duviard P, Baldet P, Jaubert F, Fellous M, Sultan C: **Extinction of FOXL2 expression in aggressive ovarian granulosa cell tumors in children.** *Fertil Steril* 2007, **87**:896-901.
- Shah SP, Köbel M, Senz J, Morin RD, Clarke BA, Wiegand KC, Leung G, Zayed A, Mehl E, Kalloger SE, Sun M, Giuliany R, Yorlida E, Jones S, Varhol R, Swenerton KD, Miller D, Clement PB, Crane C, Madore J, Provencher D, Leung P, DeFazio A, Khattra J, Turashvili G, Zhao Y, Zeng T, Glover JN, Vanderhyden B, Zhao C, et al.: **Mutation of FOXL2 in granulosa-cell tumors of the ovary.** *N Engl J Med* 2009, **360**:2719-2729.
- Wang ZJ, Churchman M, Campbell IG, Xu WH, Yan ZY, McCluggage WG, Foulkes WD, Tomlinson IP: **Allele loss and mutation screen at the Peutz-Jeghers (LKB1) locus (19p13.3) in sporadic ovarian tumours.** *Br J Cancer* 1999, **80**:70-72.
- Kato N, Romero M, Catusus L, Prat J: **The STK11/LKB1 Peutz-Jegher gene is not involved in the pathogenesis of sporadic sex cord-stromal tumors, although loss of heterozygosity at 19p13.3 indicates other gene alteration in these tumors.** *Hum Pathol* 2004, **35**:1101-1104.
- Kim K, Lindstrom MJ, Gould MN: **Regions of H- and K-ras that provide organ specificity/potency in mammary cancer induction.** *Cancer Res* 2002, **62**:1241-1245.
- King LA, Okagaki T, Gallup DG, Twiggs LB, Messing MJ, Carson LF: **Mitotic count, nuclear atypia, and immunohistochemical determination of Ki-67, c-myc, p21-ras, c-erbB2, and p53 expression in granulosa cell tumors of the ovary: mitotic count and Ki-67 are indicators of poor prognosis.** *Gynecol Oncol* 1996, **61**:227-232.
- Makrigiannakis A, Amin K, Coukos G, Tilly JL, Coutifaris C: **Regulated expression and potential roles of p53 and Wilms' tumor suppressor gene (WT1) during follicular development in the human ovary.** *J Clin Endocrinol Metab* 2000, **85**:449-459.
- Coppes MJ, Ye Y, Rackley R, Zhao XL, Liefers GJ, Casey G, Williams BR: **Analysis of WT1 in granulosa cell and other sex cord-stromal tumors.** *Cancer Res* 1993, **53**:2712-2714.
- Liu FS, Ho ES, Lai CR, Chen JT, Shih RT, Yang CH, Tsao CM: **Overexpression of p53 is not a feature of ovarian granulosa cell tumors.** *Gynecol Oncol* 1996, **61**:50-53.
- Kappes S, Milde-Langosch K, Kressin P, Passlack B, Dockhorn-Dworniczak B, Rohlke P, Loning T: **p53 mutations in ovarian tumors, detected by temperature-gradient gel electrophoresis, direct sequencing and immunohistochemistry.** *Int J Cancer* 1995, **64**:52-59.
- Chu S, Rushdi S, Zumpe ET, Mamers P, Healy DL, Jobling T, Burger HG, Fuller PJ: **FSH-regulated gene expression profiles in ovarian tumours and normal ovaries.** *Mol Hum Reprod* 2002, **8**:426-433.

27. Fuller PJ, Verity K, Shen Y, Mamers P, Jobling T, Burger HG: **No evidence of a role for mutations or polymorphisms of the follicle-stimulating hormone receptor in ovarian granulosa cell tumors.** *J Clin Endocrinol Metab* 1998, **83**:274-279.
28. Kotlar T, Young RH, Albanese C, Crowley WF Jr, Scully RE, Jameson JL: **Absence of mutations in the FSH receptor in ovarian granulosa cell tumors.** *J Clin Endocrinol Metab* 1998, **83**:3001.
29. Bas F, Pescovitz OH, Steinmetz R: **No activating mutations of FSH receptor in four children with ovarian juvenile granulosa cell tumors and the association of these tumors with central precocious puberty.** *J Pediatr Adolesc Gynecol* 2009, **22**:173-179.
30. Fragoso MC, Latronico AC, Carvalho FM, Zerbini MC, Marcondes JA, Araujo LM, Lando VS, Frazzatto ET, Mendonca BB, Villares SM: **Activating mutation of the stimulatory G protein (gsp) as a putative cause of ovarian and testicular human stromal Leydig cell tumors.** *J Clin Endocrinol Metab* 1998, **83**:2074-2078.
31. Ligtenberg MJ, Siers M, Themmen AP, Hanselaar TG, Willemsen W, Brunner HG: **Analysis of mutations in genes of the follicle-stimulating hormone receptor signaling pathway in ovarian granulosa cell tumors.** *J Clin Endocrinol Metab* 1999, **84**:2233-2234.
32. Kalfa N, Ecochard A, Patte C, Duvillard P, Audran F, Pienkowski C, Thibaud E, Brauner R, Lecointre C, Plantaz D, Guedj AM, Paris F, Baldet P, Lumbroso S, Sultan C: **Activating mutations of the stimulatory g protein in juvenile ovarian granulosa cell tumors: a new prognostic factor?** *J Clin Endocrinol Metab* 2006, **91**:1842-1847.
33. Chien J, Wong E, Nikes E, Noble MJ, Pantazis CG, Shah GV: **Constitutive activation of stimulatory guanine nucleotide binding protein (G(S)alphaQL)-mediated signaling increases invasiveness and tumorigenicity of PC-3M prostate cancer cells.** *Oncogene* 1999, **18**:3376-3382.
34. Barr FG: **Gene fusions involving PAX and FOX family members in alveolar rhabdomyosarcoma.** *Oncogene* 2001, **20**:5736-5746.
35. Hillion J, Le Coniat M, Jonveaux P, Berger R, Bernard OA: **AF6q21, a novel partner of the MLL gene in t(6;11)(q21;q23), defines a forkhead transcriptional factor subfamily.** *Blood* 1997, **90**:3714-3719.
36. Mani H, Shilo K, Galvin JR, Stocker JT, Franks TJ: **Spectrum of precursor and invasive neoplastic lesions in type 1 congenital pulmonary airway malformation: case report and review of the literature.** *Histopathology* 2007, **51**:561-565.
37. Mani N, Lowe D, Pope L, El-Daly H, Pfeleiderer A: **An unusual case of laryngeal spindle cell carcinoma metastasising to the orbit and heart.** *J Laryngol Otol* 2007, **121**:e19.
38. Mani SA, Yang J, Brooks M, Schwaninger G, Zhou A, Miura N, Kutok JL, Hartwell K, Richardson AL, Weinberg RA: **Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and is associated with aggressive basal-like breast cancers.** *Proc Natl Acad Sci USA* 2007, **104**:10069-10074.
39. Nakamura T, Furukawa Y, Nakagawa H, Tsunoda T, Ohgashi H, Murata K, Ishikawa O, Ohgaki K, Kashimura N, Miyamoto M, Hirano S, Kondo S, Katoh H, Nakamura Y, Katagiri T: **Genome-wide cDNA microarray analysis of gene expression profiles in pancreatic cancers using populations of tumor cells and normal ductal epithelial cells selected for purity by laser microdissection.** *Oncogene* 2004, **23**:2385-2400.
40. Blount AL, Schmidt K, Justice NJ, Vale WW, Fischer WH, Bilezikjian LM: **FoxL2 and Smad3 coordinately regulate follistatin gene transcription.** *J Biol Chem* 2009, **284**:7631-7645.
41. Ellsworth BS, Burns AT, Escudero KW, Duval DL, Nelson SE, Clay CM: **The gonadotropin releasing hormone (GnRH) receptor activating sequence (GRAS) is a composite regulatory element that interacts with multiple classes of transcription factors including Smads, AP-1 and a forkhead DNA binding protein.** *Mol Cell Endocrinol* 2003, **206**:93-111.
42. Dragovic RA, Ritter LJ, Schulz SJ, Amato F, Thompson JG, Armstrong DT, Gilchrist RB: **Oocyte-secreted factor activation of SMAD 2/3 signaling enables initiation of mouse cumulus cell expansion.** *Biol Reprod* 2007, **76**:848-857.

Published: 25 August 2009
doi:10.1186/gm81
© 2009 BioMed Central Ltd