



HAL
open science

Simultaneous Identification of the Diffusion Coefficient and the Potential for the Schrödinger Operator with only one Observation

Laure Cardoulis, Patricia Gaitan

► **To cite this version:**

Laure Cardoulis, Patricia Gaitan. Simultaneous Identification of the Diffusion Coefficient and the Potential for the Schrödinger Operator with only one Observation. 2009. hal-00432821v1

HAL Id: hal-00432821

<https://hal.science/hal-00432821v1>

Preprint submitted on 17 Nov 2009 (v1), last revised 1 Feb 2010 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Simultaneous Identification of the Diffusion Coefficient and the Potential for the Schrödinger Operator with only one Observation

Laure Cardoulis¹ and Patricia Gaitan²

¹ Université de Toulouse, UT1 CEREMATH, CNRS, Institut de Mathématiques de Toulouse, UMR 5219, 21 Allées de Brienne 31042 Toulouse, France

² Universités d'Aix-Marseille, IUT Aix-en-Provence Avenue Gaston Berger 413 av Gaston Berger, 13625 Aix-en-Provence et LATP, UMR CNRS 6632, 39, rue Joliot Curie, 13453 Marseille Cedex 13, France

E-mail: laure.cardoulis@univ-tlse1.fr and patricia.gaitan@univmed.fr

Abstract. This article is devoted to prove a stability result for two independent coefficients for a Schrödinger operator in an unbounded strip. The result is obtained with only one observation on an unbounded subset of the boundary and the data of the solution at a fixed time on the whole domain.

1. Introduction

Let $\Omega = \mathbb{R} \times (d, 2d)$ be an unbounded strip of \mathbb{R}^2 with a fixed width $d > 0$. Let ν be the outward unit normal to Ω on $\Gamma = \partial\Omega$. We denote $x = (x_1, x_2)$ and $\Gamma = \Gamma^+ \cup \Gamma^-$, where $\Gamma^+ = \{x \in \Gamma; x_2 = 2d\}$ and $\Gamma^- = \{x \in \Gamma; x_2 = d\}$. We consider the following Schrödinger equation

$$\begin{cases} Hq := i\partial_t q + a\Delta q + bq = 0 & \text{in } \Omega \times (0, T), \\ q(x, t) = F(x, t) & \text{on } \partial\Omega \times (0, T), \\ q(x, 0) = q_0(x) & \text{in } \Omega, \end{cases} \quad (1.1)$$

where a and b are real-valued functions such that $a \in \mathcal{C}^3(\overline{\Omega})$, $b \in \mathcal{C}^2(\overline{\Omega})$ and $a(x) \geq a_{min} > 0$. Moreover, we assume that a is bounded and b and all its derivatives up to order two are bounded. If we assume that q_0 belongs to $H^4(\Omega)$ and $F \in H^2(0, T, H^2(\partial\Omega)) \cap H^1(0, T, H^4(\partial\Omega)) \cap H^3(0, T, L^2(\partial\Omega))$, then (1.1) admits a solution in $H^1(0, T, H^2(\Omega)) \cap H^2(0, T, L^2(\Omega))$.

Our problem can be stated as follows:

Is it possible to determine the coefficients a and b from the measurement of $\partial_\nu(\partial_t^2 q)$ on Γ^+ ?

Let q (resp. \tilde{q}) be a solution of (1.1) associated with (a, b, F, q_0) (resp. $(\tilde{a}, \tilde{b}, F, q_0)$). We assume that q_0 is a real valued function.

Our main result is

$$\begin{aligned} \|a - \tilde{a}\|_{L^2(\Omega)}^2 + \|b - \tilde{b}\|_{L^2(\Omega)}^2 &\leq C \|\partial_\nu(\partial_t^2 q) - \partial_\nu(\partial_t^2 \tilde{q})\|_{L^2((-T, T) \times \Gamma^+)}^2 \\ &\quad + C \sum_{i=0}^2 \|\partial_t^i (q - \tilde{q})(\cdot, 0)\|_{H^2(\Omega)}^2, \end{aligned}$$

where C is a positive constant which depends on (Ω, Γ, T) and where the above norms are weighted Sobolev norms.

This paper is an improvement of the work [7] in the sense that we simultaneously determine with only one observation, two independent coefficients, the diffusion coefficient and the potential. We use for that two important tools: Carleman estimate (2.5) and Lemma 2.4.

Carleman inequalities constitute a very efficient tool to derive observability estimates. The method of Carleman estimates has been introduced in the field of inverse problems by Bukhgeim and Klivanov (see [5], [6], [10], [11]). Carleman estimates techniques are presented in [12] for standard coefficients inverse problems for both linear and non-linear partial differential equations. These methods give a local Lipschitz stability around a single known solution.

A lot of works using the same strategy concern the wave equation (see [13], [3], [2]) and the heat equation (see [15], [9], [4]). For the determination of a time-independent potential in Schrödinger evolution equation, we can refer to [1] for bounded domains and [7] for unbounded domains. We can also cite [14] where the authors use

weight functions satisfying a relaxed pseudo-convexity condition which allows to prove Carleman inequalities with less restrictive boundary observations.

Up to our knowledge, there are few results concerning the simultaneous identification of two coefficients with only one observation. In [8] a stability result is given for the particular case where each coefficient only depends on one variable ($a = a(x_2)$ and $b = b(x_1)$) for the operator $i\partial_t q + \nabla \cdot (a\nabla q) + bq$ in an unbounded strip of \mathbb{R}^2 . The authors give a stability result for the diffusion coefficient a and the potential b with only one observation in an unbounded part of the boundary.

This paper is organized as follows. Section 2 is devoted to some useful estimates. We first give an adapted global Carleman estimate for the operator H . We then recall the crucial Lemma given in [12]. In Section 3 we state and prove our main result.

2. Some Useful Estimates

2.1. Global Carleman Inequality

Let a be a real-valued function in $\mathcal{C}^3(\overline{\Omega})$ and b be a real-valued function in $\mathcal{C}^2(\overline{\Omega})$ such that

- Assumption 2.1**
- $a \geq a_{\min} > 0$, a and all its derivatives up to order three are bounded,
 - b and its derivatives up to order two are bounded.

Let $q(x, t)$ be a function equals to zero on $\partial\Omega \times (-T, T)$ and solution of the Schrödinger equation

$$i\partial_t q + a\Delta q + bq = f.$$

We prove here a global Carleman-type estimate for q with a single observation acting on a part Γ^+ of the boundary Γ in the right-hand side of the estimate.

Note that this estimate is quite similar to the one obtained in [7], but the computations are different. Indeed, the weight function β does not satisfy the same pseudo-convexity assumptions (see Assumption 2.2) and the decomposition of the operator H is different (see (2.3)).

Let $\tilde{\beta}$ be a $\mathcal{C}^4(\overline{\Omega})$ positive function such that there exists positive constants C_0, C_{pc} which satisfy

- Assumption 2.2**
- $|\nabla\tilde{\beta}| \geq C_0 > 0$ in Ω , $\partial_\nu\tilde{\beta} \leq 0$ on Γ^- ,
 - $\tilde{\beta}$ and all its derivatives up to order four are bounded in $\overline{\Omega}$,
 - $2\Re(D^2\tilde{\beta}(\zeta, \bar{\zeta})) - \nabla a \cdot \nabla\tilde{\beta}|\zeta|^2 + 2a^2|\nabla\tilde{\beta} \cdot \zeta|^2 \geq C_{pc}|\zeta|^2$, for all $\zeta \in \mathbb{C}$

where

$$D^2\tilde{\beta} = \begin{pmatrix} \partial_{x_1}(a^2\partial_{x_1}\tilde{\beta}) & \partial_{x_1}(a^2\partial_{x_2}\tilde{\beta}) \\ \partial_{x_2}(a^2\partial_{x_1}\tilde{\beta}) & \partial_{x_2}(a^2\partial_{x_2}\tilde{\beta}) \end{pmatrix}.$$

Note that the last assertion of Assumption 2.2 expresses the pseudo-convexity condition for the function β . This Assumption imposes restrictive conditions for the choice of the

diffusion coefficient a in connection with the function $\tilde{\beta}$ as in [7].

Note that there exist functions satisfying such assumptions. Indeed if we assume that $\tilde{\beta}(x) := \tilde{\beta}(x_2)$, these conditions can be written in the following form:

$$A = 2\partial_{x_2}(a^2\partial_{x_2}\tilde{\beta}) - \partial_{x_2}a \partial_{x_2}\tilde{\beta} + 2a^2(\partial_{x_2}\tilde{\beta})^2 \geq cst > 0$$

and

$$-\frac{(\partial_{x_1}(a^2\partial_{x_2}\tilde{\beta}))^2}{A} - \partial_{x_2}a \partial_{x_2}\tilde{\beta} \geq cst > 0.$$

For example $\tilde{\beta}(x) = e^{-x_2}$ with $a(x) = \frac{1}{2}(x_2^2 + 5)$ satisfy the previous conditions (with $x_2 \in (d, 2d)$).

Then, we define $\beta = \tilde{\beta} + K$ with $K = m\|\tilde{\beta}\|_\infty$ and $m > 1$. For $\lambda > 0$ and $t \in (-T, T)$, we define the following weight functions

$$\varphi(x, t) = \frac{e^{\lambda\beta(x)}}{(T+t)(T-t)}, \quad \eta(x, t) = \frac{e^{2\lambda K} - e^{\lambda\beta(x)}}{(T+t)(T-t)}.$$

We set $\psi = e^{-s\eta}q$, $M\psi = e^{-s\eta}H(e^{s\eta}\psi)$ for $s > 0$. Let H be the operator defined by

$$Hq := i\partial_t q + a\Delta q + bq \text{ in } \Omega \times (-T, T). \quad (2.2)$$

Following [1], we introduce the operators :

$$\begin{aligned} M_1\psi &:= i\partial_t\psi + a\Delta\psi + s^2a|\nabla\eta|^2\psi + (b - s\nabla\eta \cdot \nabla a)\psi, \\ M_2\psi &:= is\partial_t\eta\psi + 2as\nabla\eta \cdot \nabla\psi + s\nabla \cdot (a\nabla\eta)\psi. \end{aligned} \quad (2.3)$$

Then

$$\begin{aligned} \int_{-T}^T \int_{\Omega} |M\psi|^2 dx dt &= \int_{-T}^T \int_{\Omega} |M_1\psi|^2 dx dt + \int_{-T}^T \int_{\Omega} |M_2\psi|^2 dx dt \\ &\quad + 2\Re\left(\int_{-T}^T \int_{\Omega} M_1\psi \overline{M_2\psi} dx dt\right), \end{aligned}$$

where \bar{z} is the conjugate of z , $\Re(z)$ its real part and $\Im(z)$ its imaginary part. Then the following result holds.

Theorem 2.3 *Let H , M_1 , M_2 be the operators defined respectively by (2.2), (2.3). We assume that Assumptions 2.1 and 2.2 are satisfied. Then there exist $\lambda_0 > 0$, $s_0 > 0$ and a positive constant $C = C(\Omega, \Gamma, T)$ such that, for any $\lambda \geq \lambda_0$ and any $s \geq s_0$, the next inequality holds:*

$$\begin{aligned} s^3\lambda^4 \int_{-T}^T \int_{\Omega} e^{-2s\eta}|q|^2 dx dt + s\lambda \int_{-T}^T \int_{\Omega} e^{-2s\eta}|\nabla q|^2 dx dt + \|M_1(e^{-s\eta}q)\|_{L^2(\Omega \times (-T, T))}^2 \\ + \|M_2(e^{-s\eta}q)\|_{L^2(\Omega \times (-T, T))}^2 \leq Cs\lambda \int_{-T}^T \int_{\Gamma^+} e^{-2s\eta}|\partial_\nu q|^2 \partial_\nu \beta d\sigma dt \quad (2.4) \\ + \int_{-T}^T \int_{\Omega} e^{-2s\eta} |Hq|^2 dx dt, \end{aligned}$$

for all q satisfying $q \in L^2(-T, T; H_0^1(\Omega) \cap H^2(\Omega)) \cap H^1(-T, T; L^2(\Omega))$, $\partial_\nu q \in L^2(-T, T; L^2(\Gamma))$. Moreover we have

$$\begin{aligned} & s^3 \lambda^4 \int_{-T}^T \int_{\Omega} e^{-2s\eta} |q|^2 dx dt + s\lambda \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla q|^2 dx dt + \|M_1(e^{-s\eta} q)\|_{L^2(\Omega \times (-T, T))}^2 \\ & + \|M_2(e^{-s\eta} q)\|_{L^2(\Omega \times (-T, T))}^2 + s^{-1} \lambda^{-1} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |i\partial_t q + a\Delta q|^2 dx dt \quad (2.5) \\ & \leq C \left[s\lambda \int_{-T}^T \int_{\Gamma^+} e^{-2s\eta} |\partial_\nu q|^2 \partial_{\nu, \beta} d\sigma dt + \int_{-T}^T \int_{\Omega} e^{-2s\eta} |Hq|^2 dx dt \right]. \end{aligned}$$

Proof:

We have to estimate the scalar product

$$\Re \left(\int_{-T}^T \int_{\Omega} M_1 \psi \overline{M_2 \psi} dx dt \right) = \sum_{i=1}^4 \sum_{j=1}^3 I_{ij}$$

with

$$\begin{aligned} I_{11} &= \Re \left(\int_{-T}^T \int_{\Omega} (i\partial_t \psi)(-is\partial_t \eta \overline{\psi}) dx dy dt \right), \quad I_{12} = \Re \left(\int_{-T}^T \int_{\Omega} (i\partial_t \psi)(2as\nabla \eta \cdot \nabla \overline{\psi}) dx dy dt \right), \\ I_{13} &= \Re \left(\int_{-T}^T \int_{\Omega} (i\partial_t \psi)(s\nabla \cdot (a\nabla \eta) \overline{\psi}) dx dy dt \right), \quad I_{21} = \Re \left(\int_{-T}^T \int_{\Omega} (a\Delta \psi)(-is\partial_t \eta \overline{\psi}) dx dy dt \right), \\ I_{22} &= \Re \left(\int_{-T}^T \int_{\Omega} (a\Delta \psi)(2as\nabla \eta \cdot \nabla \overline{\psi}) dx dy dt \right), \quad I_{23} = \Re \left(\int_{-T}^T \int_{\Omega} (a\Delta \psi)(s\nabla \cdot (a\nabla \eta) \overline{\psi}) dx dy dt \right), \\ I_{31} &= \Re \left(\int_{-T}^T \int_{\Omega} (s^2 a |\nabla \eta|^2 \psi)(-is\partial_t \eta \overline{\psi}) dx dy dt \right), \quad I_{32} = \Re \left(\int_{-T}^T \int_{\Omega} (s^2 a |\nabla \eta|^2 \psi)(2as\nabla \eta \cdot \nabla \overline{\psi}) dx dy dt \right), \\ I_{33} &= \Re \left(\int_{-T}^T \int_{\Omega} (s^2 a |\nabla \eta|^2 \psi)(s\nabla \cdot (a\nabla \eta) \overline{\psi}) dx dy dt \right), \quad I_{41} = \Re \left(\int_{-T}^T \int_{\Omega} ((b - s\nabla \eta \cdot \nabla a) \psi)(-is\partial_t \eta \overline{\psi}) dx dy dt \right), \\ I_{42} &= \Re \left(\int_{-T}^T \int_{\Omega} ((b - s\nabla \eta \cdot \nabla a) \psi)(2as\nabla \eta \cdot \nabla \overline{\psi}) dx dy dt \right), \quad I_{43} = \Re \left(\int_{-T}^T \int_{\Omega} ((b - s\nabla \eta \cdot \nabla a) \psi)(s\nabla \cdot (a\nabla \eta) \overline{\psi}) dx dy dt \right). \end{aligned}$$

Following [1], using integrations by part and Young estimates, we get (2.4). Moreover from (2.3) we have:

$$i\partial_t q + a\Delta q = M_1 q - s^2 a |\nabla \eta|^2 q + (b - s\nabla \eta \cdot \nabla a) q.$$

So

$$\begin{aligned} i\partial_t q + a\Delta q &= e^{s\eta} M_1(e^{-s\eta} q) + is\partial_t \eta q - ae^{s\eta} \Delta(e^{-s\eta} q) - 2ae^{s\eta} \nabla(e^{-s\eta}) \cdot \nabla q \\ &\quad - s^2 a |\nabla \eta|^2 q + (b - s\nabla \eta \cdot \nabla a) q. \end{aligned}$$

And we deduce (2.5) from (2.4). ■

2.2. The Crucial Lemma

We recall in this section the proof of a very important lemma proved by Klivanov and Timonov (see for example [11], [12]).

Lemma 2.4 *There exists a positive constant κ such that*

$$\int_{-T}^T \int_{\Omega} \left| \int_0^t q(x, \xi) d\xi \right|^2 e^{-2s\eta} dx dt \leq \frac{\kappa}{s} \int_{-T}^T \int_{\Omega} |q(x, t)|^2 e^{-2s\eta} dx dt,$$

for all $s > 0$.

Proof :

By the Cauchy-Schwartz inequality, we have

$$\begin{aligned} & \int_{-T}^T \int_{\Omega} \left| \int_0^t q(x, \xi) d\xi \right|^2 e^{-2s\eta} dx dt \leq \int_{\Omega} \int_{-T}^T |t| \left| \int_0^t |q(x, \xi)|^2 d\xi \right| e^{-2s\eta} dx dt \quad (2.6) \\ & \leq \int_{\Omega} \int_0^T t \left(\int_0^t |q(x, \xi)|^2 d\xi \right) e^{-2s\eta} dx dt + \int_{\Omega} \int_{-T}^0 (-t) \left(\int_t^0 |q(x, \xi)|^2 d\xi \right) e^{-2s\eta} dx dt. \end{aligned}$$

Note that

$$\partial_t (e^{-2s\eta(x,t)}) = -2s(e^{2\lambda K} - e^{\lambda\beta(x)}) \frac{2t}{(T^2 - t^2)^2} e^{-2s\eta(x,t)}.$$

So, if we denote by $\alpha(x) = e^{2\lambda K} - e^{\lambda\beta(x)}$, we have

$$te^{-2s\eta(x,t)} = -\frac{(T^2 - t^2)^2}{4s\alpha(x)} \partial_t (e^{-2s\eta(x,t)}).$$

For the first integral of the right hand side of (2.6), by integration by parts we have

$$\begin{aligned} & \int_{\Omega} \int_0^T t \left(\int_0^t |q(x, \xi)|^2 d\xi \right) e^{-2s\eta} dx dt = \int_{\Omega} \int_0^T \left(\int_0^t |q(x, \xi)|^2 d\xi \right) \frac{(T^2 - t^2)^2}{-4s\alpha(x)} \partial_t (e^{-2s\eta}) dt dx \\ & = \int_{\Omega} \left[\left(\int_0^t |q(x, \xi)|^2 d\xi \right) \frac{(T^2 - t^2)^2}{-4s\alpha(x)} e^{-2s\eta} \right]_{t=0}^{t=T} dx + \int_{\Omega} \int_0^T |q(x, t)|^2 \frac{(T^2 - t^2)^2}{4s\alpha(x)} e^{-2s\eta} dt dx \\ & \quad + \int_{\Omega} \int_0^T \left(\int_0^t |q(x, \xi)|^2 d\xi \right) \frac{t(t^2 - T^2)}{s\alpha(x)} e^{-2s\eta} dt dx. \end{aligned}$$

Here we used $\alpha(x) > 0$ for all $x \in \overline{\Omega}$ and we obtain

$$\int_{\Omega} \int_0^T t \left(\int_0^t |q(x, \xi)|^2 d\xi \right) e^{-2s\eta} dx dt \leq \frac{1}{4s} \sup_{x \in \overline{\Omega}} \left(\frac{1}{\alpha(x)} \right) \int_{\Omega} \int_0^T |q(x, t)|^2 e^{-2s\eta} (T^2 - t^2)^2 dx dt.$$

Similarly for the second integral of the right hand side of (2.6)

$$\int_{\Omega} \int_{-T}^0 (-t) \left(\int_t^0 |q(x, \xi)|^2 d\xi \right) e^{-2s\eta} dx dt \leq \frac{1}{4s} \sup_{x \in \overline{\Omega}} \left(\frac{1}{\alpha(x)} \right) \int_{\Omega} \int_{-T}^0 |q(x, t)|^2 e^{-2s\eta} (T^2 - t^2)^2 dx dt.$$

Thus the proof of Lemma 2.4 is completed. ■

3. Stability result

In this section, we establish a stability inequality for the diffusion coefficient a and the potential b .

Let q be solution of

$$\begin{cases} i\partial_t q + a\Delta q + bq = 0 & \text{in } \Omega \times (0, T), \\ q(x, t) = F(x, t) & \text{on } \partial\Omega \times (0, T), \\ q(x, 0) = q_0(x) & \text{in } \Omega, \end{cases}$$

and \tilde{q} be solution of

$$\begin{cases} i\partial_t \tilde{q} + \tilde{a}\Delta \tilde{q} + \tilde{b}\tilde{q} = 0 & \text{in } \Omega \times (0, T), \\ \tilde{q}(x, t) = F(x, t) & \text{on } \partial\Omega \times (0, T), \\ \tilde{q}(x, 0) = q_0(x) & \text{in } \Omega, \end{cases}$$

where (a, b) and (\tilde{a}, \tilde{b}) both satisfy Assumption 2.1.

Assumption 3.1 • *All the time-derivatives up to order three and the space-derivatives up to order four for \tilde{q} exist and are bounded.*

- *There exists a positive constant $C > 0$ such that $|\tilde{q}| \geq C$, $|\partial_t(\frac{\Delta \tilde{q}}{\tilde{q}})| \geq C$, $|\Delta \tilde{q}| \geq C$, $|\partial_t(\frac{\tilde{q}}{\Delta \tilde{q}})| \geq C$.*
- *q_0 is a real-valued function.*

Since q_0 is a real-valued function, we can extend the function q (resp. \tilde{q}) on $\Omega \times (-T, T)$ by the formula $q(x, t) = \bar{q}(x, -t)$ for every $(x, t) \in \Omega \times (-T, 0)$. Note that this extension satisfies the previous Carleman estimate. Our main stability result is

Theorem 3.2 *Let q and \tilde{q} be solutions of (1.1). We assume that Assumptions 2.1, 2.2, 3.1 are satisfied. Then there exists a positive constant $C = C(\Omega, \Gamma, T)$ such that for s and λ large enough,*

$$\begin{aligned} \int_{-T}^T \int_{\Omega} e^{-2s\eta} (|\tilde{a} - a|^2 + |\tilde{b} - b|^2) dx dt &\leq Cs\lambda^2 \int_{-T}^T \int_{\Gamma^+} \varphi e^{-2s\eta} \partial_\nu \beta |\partial_\nu (\partial_t^2 q - \partial_t^2 \tilde{q})|^2 d\sigma dt \\ &+ C\lambda \int_{-T}^T \int_{\Omega} e^{-2s\eta} \left(\sum_{i=0}^2 |\partial_t^i (q - \tilde{q})(\cdot, 0)|^2 + |\nabla(q - \tilde{q})(\cdot, 0)|^2 \right. \\ &\quad \left. + |\partial_t \nabla(q - \tilde{q})(\cdot, 0)|^2 + |\partial_t \Delta(q - \tilde{q})(\cdot, 0)|^2 \right) dx dt. \end{aligned}$$

Therefore

$$\begin{aligned} \|a - \tilde{a}\|_{L^2(\Omega)}^2 + \|b - \tilde{b}\|_{L^2(\Omega)}^2 &\leq C \|\partial_\nu (\partial_t^2 q) - \partial_\nu (\partial_t^2 \tilde{q})\|_{L^2((-T, T) \times \Gamma^+)}^2 \\ &+ C \sum_{i=0}^2 \|\partial_t^i (q - \tilde{q})(\cdot, 0)\|_{H^2(\Omega)}^2, \end{aligned}$$

Proof:

We denote by $u = q - \tilde{q}$, $\alpha = \tilde{a} - a$ and $\gamma = \tilde{b} - b$, so we get:

$$\begin{cases} i\partial_t u + a\Delta u + bu = \alpha\Delta\tilde{q} + \gamma\tilde{q} & \text{in } \Omega \times (-T, T), \\ u(x, t) = 0 & \text{on } \partial\Omega \times (-T, T), \\ u(x, 0) = 0 & \text{in } \Omega. \end{cases} \quad (3.7)$$

The proof will be done in two steps: in a first step we prove an estimation for α and in a second step for γ .

First step: We set $u_1 = \frac{u}{\tilde{q}}$. Then from (3.7)

$$\begin{cases} i\partial_t u_1 + a\Delta u_1 + bu_1 + A_{11}u_1 + B_{11} \cdot \nabla u_1 = \alpha \frac{\Delta\tilde{q}}{\tilde{q}} + \gamma & \text{in } \Omega \times (-T, T), \\ u_1(x, t) = 0 & \text{on } \partial\Omega \times (-T, T) \end{cases}$$

where $A_{11} = i\frac{\partial_t\tilde{q}}{\tilde{q}} + a\frac{\Delta\tilde{q}}{\tilde{q}}$ and $B_{11} = \frac{2a}{\tilde{q}}\nabla\tilde{q}$.

Then defining $u_2 = \partial_t u_1$ we get

$$\begin{cases} i\partial_t u_2 + a\Delta u_2 + bu_2 + \sum_{i=1}^2 A_{i2}u_i + \sum_{i=1}^2 B_{i2} \cdot \nabla u_i = \alpha\partial_t\left(\frac{\Delta\tilde{q}}{\tilde{q}}\right) & \text{in } \Omega \times (-T, T), \\ u_2(x, t) = 0 & \text{on } \partial\Omega \times (-T, T) \end{cases}$$

where $A_{12} = \partial_t A_{11}$, $A_{22} = A_{11}$, $B_{12} = \partial_t B_{11}$, $B_{22} = B_{11}$.

Now let $u_3 = \frac{u_2}{\partial_t\left(\frac{\Delta\tilde{q}}{\tilde{q}}\right)}$, we have

$$\begin{cases} i\partial_t u_3 + a\Delta u_3 + bu_3 + \sum_{i=1}^3 A_{i3}u_i + \sum_{i=1}^3 B_{i3} \cdot \nabla u_i = \alpha & \text{in } \Omega \times (-T, T), \\ u_3(x, t) = 0 & \text{on } \partial\Omega \times (-T, T) \end{cases} \quad (3.8)$$

where A_{i3} and B_{i3} are bounded functions. If we denote by $g = \partial_t\left(\frac{\Delta\tilde{q}}{\tilde{q}}\right)$, then $A_{13} = \frac{1}{g}A_{12}$, $A_{23} = \frac{1}{g}A_{22}$, $A_{33} = \frac{1}{g}(i\partial_t g + \Delta g)$, $B_{13} = \frac{1}{g}B_{12}$, $B_{23} = \frac{1}{g}B_{22}$, $B_{33} = \frac{2a}{g}\nabla g$.

At last we define $u_4 = \partial_t u_3$ and we get

$$\begin{cases} i\partial_t u_4 + a\Delta u_4 + bu_4 + \sum_{i=1}^4 A_{i4}u_i + \sum_{i=1}^4 B_{i4} \cdot \nabla u_i = 0 & \text{in } \Omega \times (-T, T), \\ u_4(x, t) = 0 & \text{on } \partial\Omega \times (-T, T) \end{cases}$$

where A_{i4} and B_{i4} are still bounded functions. Note that $A_{14} = \partial_t A_{13}$, $A_{24} = \partial_t A_{23} + A_{13}$, $A_{34} = \partial_t A_{33} + A_{23}\partial_t g + B_{23} \cdot \nabla(\partial_t g)$, $A_{44} = A_{23}g + A_{33} + B_{23} \cdot \nabla g$, $B_{14} = \partial_t B_{13}$, $B_{24} = \partial_t B_{23} + B_{13}$, $B_{34} = \partial_t B_{33} + \partial_t g B_{23}$, $B_{44} = B_{33} + gB_{23}$.

Applying the Carleman inequality (2.5) for u_4 we obtain (for s and λ sufficiently large):

$$\begin{aligned} & s^3\lambda^4 \int_{-T}^T \int_{\Omega} e^{-2s\eta}|u_4|^2 dx dt + s\lambda \int_{-T}^T \int_{\Omega} e^{-2s\eta}|\nabla u_4|^2 dx dt \\ & \quad + s^{-1}\lambda^{-1} \int_{-T}^T \int_{\Omega} e^{-2s\eta}|i\partial_t u_4 + a\Delta u_4|^2 dx dt \\ & \leq C \left[s\lambda \int_{-T}^T \int_{\Gamma^+} e^{-2s\eta}|\partial_\nu u_4|^2 \partial_\nu \beta d\sigma dt + \sum_{i=1}^3 \int_{-T}^T \int_{\Omega} e^{-2s\eta} (|u_i|^2 + |\nabla u_i|^2) dx dt \right]. \end{aligned} \quad (3.9)$$

Note that $\int_{-T}^T \int_{\Omega} e^{-2s\eta} |u_1|^2 dx dt = \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\int_0^t \partial_t u_1|^2 dx dt$, so from Lemma 2.4 we get

$$\begin{aligned} & \int_{-T}^T \int_{\Omega} e^{-2s\eta} |u_1|^2 dx dt \leq \frac{C}{s} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |u_3|^2 dx dt \\ & \leq \frac{C}{s^2} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |u_4|^2 dx dt + \frac{C}{s} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |u_3(\cdot, 0)|^2 dx dt. \end{aligned}$$

By the same way, we have

$$\begin{aligned} & \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla u_1|^2 dx dt \leq \frac{C}{s^2} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla u_4|^2 dx dt \\ & + \frac{C}{s} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla u_3(\cdot, 0)|^2 dx dt + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla u_1(\cdot, 0)|^2 dx dt. \end{aligned}$$

So (3.9) becomes

$$\begin{aligned} & s^3 \lambda^4 \int_{-T}^T \int_{\Omega} e^{-2s\eta} |u_4|^2 dx dt + s\lambda \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla u_4|^2 dx dt \quad (3.10) \\ & + s^{-1} \lambda^{-1} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |i\partial_t u_4 + a\Delta u_4|^2 dx dt \leq Cs\lambda \int_{-T}^T \int_{\Gamma^+} e^{-2s\eta} |\partial_\nu u_4|^2 \partial_\nu \beta d\sigma dt \\ & + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} (|u_3(\cdot, 0)|^2 + |\nabla u_3(\cdot, 0)|^2 + |\nabla u_1(\cdot, 0)|^2) dx dt. \end{aligned}$$

Furthermore from (3.8) we have (with C a positive constant)

$$|\alpha|^2 \leq C \left(|i\partial_t u_3 + a\Delta u_3|^2 + \sum_{i=1}^3 (|u_i|^2 + |\nabla u_i|^2) \right).$$

Therefore for s sufficiently large, from Lemma 2.4

$$\begin{aligned} & \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\alpha|^2 dx dt \leq \frac{C}{s} \int_{-T}^T \int_{\Omega} e^{-2s\eta} (|i\partial_t u_4 + a\Delta u_4|^2 + |u_4|^2 + |\nabla u_4|^2) dx dt \\ & + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} |(i\partial_t u_3 + a\Delta u_3)(\cdot, 0)|^2 dx dt + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla u_1(\cdot, 0)|^2 dx dt \\ & + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} (|u_3(\cdot, 0)|^2 + |\nabla u_3(\cdot, 0)|^2) dx dt. \end{aligned}$$

Using (3.10) we get

$$\begin{aligned} & \frac{1}{\lambda} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\alpha|^2 dx dt \leq Cs\lambda \int_{-T}^T \int_{\Gamma^+} e^{-2s\eta} |\partial_\nu u_4|^2 \partial_\nu \beta d\sigma dt \\ & + \frac{C}{\lambda} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |(i\partial_t u_3 + a\Delta u_3)(\cdot, 0)|^2 dx dt \\ & + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\nabla u_1(\cdot, 0)|^2 dx dt \\ & + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} (|u_3(\cdot, 0)|^2 + |\nabla u_3(\cdot, 0)|^2) dx dt \end{aligned}$$

and then

$$\begin{aligned} & \frac{1}{\lambda} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\alpha|^2 dx dt \leq Cs\lambda \int_{-T}^T \int_{\Gamma^+} e^{-2s\eta} |\partial_{\nu} u_4|^2 \partial_{\nu} \beta d\sigma dt \quad (3.11) \\ & + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} \left(\sum_{i=0}^2 |\partial_t^i u(\cdot, 0)|^2 + |\nabla u(\cdot, 0)|^2 + |\partial_t \nabla u(\cdot, 0)|^2 + |\partial_t \Delta u(\cdot, 0)|^2 \right) dx dt. \end{aligned}$$

Second step: By the same way we obtain an estimation of γ . We set

$$v_1 = \frac{u}{\Delta \tilde{q}}, \quad v_2 = \partial_t v_1, \quad v_3 = \frac{v_2}{\partial_t \left(\frac{\tilde{q}}{\Delta \tilde{q}} \right)}.$$

Following the same methodology as in the first step, we obtain:

$$\begin{aligned} & \frac{1}{\lambda} \int_{-T}^T \int_{\Omega} e^{-2s\eta} |\alpha|^2 dx dt \leq Cs\lambda \int_{-T}^T \int_{\Gamma^+} e^{-2s\eta} |\partial_{\nu} u_4|^2 \partial_{\nu} \beta d\sigma dt \quad (3.12) \\ & + C \int_{-T}^T \int_{\Omega} e^{-2s\eta} \left(\sum_{i=0}^2 |\partial_t^i u(\cdot, 0)|^2 + |\nabla u(\cdot, 0)|^2 + |\partial_t \nabla u(\cdot, 0)|^2 + |\partial_t \Delta u(\cdot, 0)|^2 \right) dx dt. \end{aligned}$$

From (3.11) and (3.12) we can conclude. ■

Remark 3.3 (i) Note that the following function $\tilde{q}(x, t) = e^{-it} + x_2^2 + 5$ with $\tilde{a}(x) = \frac{x_2^2 + 5}{2}$, $\tilde{b}(x) = -1$ satisfies Assumption 3.1.

(ii) This method works for the Schrödinger operator in the divergential form:

$$i\partial_t q + \nabla \cdot (a\nabla q) + bq.$$

We still obtain the same stability result but with more restrictive hypotheses on the regularity of the function \tilde{q} .

References

- [1] L. BAUDOIN AND J.P. PUEL, 2002, Uniqueness and stability in an inverse problem for the Schrödinger equation, *Inverse Problems*, 18, 1537–1554.
- [2] L. BAUDOIN, A. MERCADO AND A. OSSES, 2007, A global Carleman estimate in a transmission wave equation and application to a one-measurement inverse problem, *Inverse Problems*, 23, 257–278.
- [3] M. BELLASSOUED, 2004, Uniqueness and stability in determining the speed of propagation of second order hyperbolic equation with variable coefficient, *Appl. An.*, 83, 983–1014.
- [4] A. BENABDALLAH, P. GAITAN AND J. LE ROUSSEAU, 2007, Stability of discontinuous diffusion coefficients and initial conditions in an inverse problem for the heat equation, *SIAM J. Control Optim.*, 46, 1849–1881.
- [5] A.L. BUKHGEIM, 1999, Volterra Equations and Inverse Problems, *Inverse and Ill-Posed Problems Series, VSP, Utrecht*.
- [6] A.L. BUKHGEIM AND M.V. KLIBANOV, 1981, Uniqueness in the large of a class of multidimensional inverse problems, *Soviet Math. Dokl.*, 17, 244–247.

- [7] L. CARDOULIS, M. CRISTOFOL AND P. GAITAN, 2008, Inverse problem for the Schrödinger operator in an unbounded strip, *J. Inverse and Ill-posed Problems*, 16, no 2, 127–146.
- [8] L. CARDOULIS AND P. GAITAN, 2009, Identification of two independent coefficients with one observation for the Schrödinger operator in an unbounded strip, to appear in *Comptes Rendus Académie des Sciences*.
- [9] O. YU. IMMANUVILOV AND M. YAMAMOTO, 2001, Global uniqueness and stability in determining coefficients of wave equations, *Comm. Partial Diff. Equat.*, 26, 1409–1425.
- [10] M.V. KLIBANOV, 1984, Inverse problems in the large and Carleman bounds, *Differential Equations*, 20, 755–760.
- [11] M.V. KLIBANOV, 1992, Inverse problems and Carleman estimates, *Inverse Problems*, 8, 575–596.
- [12] M.V. KLIBANOV AND A. TIMONOV, 2004, Carleman estimates for coefficient inverse problems and numerical applications, *Inverse and Ill-posed series, VSP, Utrecht*.
- [13] I. LASIECKA, R. TRIGGIANI AND P.F. YAO, 1999, Inverse/observability estimates for second order hyperbolic equations with variable coefficients, *J. Math. Anal. Appl.*, 235, 13–57.
- [14] A. MERCADO, A. OSSES AND L. ROSIER, 2008, Inverse problems for the Schrödinger equations via Carleman inequalities with degenerate weights, *Inverse Problems*, 24, 015017.
- [15] J.P. PUEL AND M. YAMAMOTO, 1997, Generic well-posedness in a multidimensional hyperbolic inverse problem, *J. Inverse Ill-Posed Probl.*, 1, 53–83.