



HAL
open science

Stationary states and fractional dynamics in systems with long range interactions

Tineke L. van den Berg, Duccio Fanelli, Xavier Leoncini

► **To cite this version:**

Tineke L. van den Berg, Duccio Fanelli, Xavier Leoncini. Stationary states and fractional dynamics in systems with long range interactions. 2009. hal-00441021v1

HAL Id: hal-00441021

<https://hal.science/hal-00441021v1>

Preprint submitted on 14 Dec 2009 (v1), last revised 29 Mar 2010 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stationary states and fractional dynamics in systems with long range interactions

Tineke Van den Berg¹, Duccio Fanelli², Xavier Leoncini³

1. *IM2NP, Aix-Marseille Université, Centre de Saint Jérôme, Case 142, 13397 Marseille Cedex 20, France*

2. *Dipartimento di Energetica "Sergio Stecco", Università di Firenze, via s. Marta 3, 50139 Firenze, Italia and Centro interdipartimentale per lo Studio delle Dinamiche Complesse (CSDC) and INFN*

3. *Centre de Physique Théorique, Aix-Marseille Université, CNRS, Luminy, Case 907, F-13288 Marseille cedex 9, France*

Dynamics of many-body Hamiltonian systems with long range interactions is studied, in the context of the so called α -HMF model. Building on the analogy with the related mean field model, we construct stationary states of the α -HMF model for which the spatial organization satisfies a fractional equation. At variance, the microscopic dynamics turns out to be regular and explicitly known. As a consequence, dynamical regularity is achieved at the price of strong spatial complexity, namely a microscopic inhomogeneity which locally displays scale invariance.

The out-of-equilibrium dynamics of many body systems subject to long range couplings defines a fascinating field of investigations, which can potentially impact different domains of applications. In a long range system every constituent is simultaneously solicited by the ensemble of microscopic actors, resulting in a complex dynamical picture. This latter scenario applies to a vast realm of fundamental problems, including gravity and plasma physics, and also extends to a rich variety of cross disciplinary studies, classically falling within the boundaries of biological and social sciences, where an intricate network of mutual interactions between agents has to be accommodated for [1]. In particular, and with specific emphasis on the peculiar non equilibrium features, long range systems often display a slow relaxation to equilibrium. They are in fact trapped in long-lasting out of equilibrium regimes, termed in the literature Quasi Stationary States (QSS) which bear distinct characteristics, as compared to the corresponding deputed equilibrium configuration. A paradigmatic representative of long range interactions, sharing the mean field viewpoint, is the so called Hamiltonian Mean Field model, which describes the coupled evolution of N rotators, populating the unitary circle and interacting via a cosines like potential. In the limit of infinite system size the discrete HMF model becomes a Vlasov equation for the evolution of the single particle distribution function[2]. This leads to a statistical based treatment, inspired to the seminal work of Lynden-Bell, which revealed the existence of two different classes of QSS, spatially homogeneous or magnetized[3]. More recently, and still with reference to the HMF case study, stationary states have been constructed using a dynamical scheme which exploited the formal analogy with a set of uncoupled pendula [4]. This represented a substantial leap forward in understanding of the dynamical properties of the QSS in the HMF model, beyond the aforementioned statistical approach. Indeed, it was understood that the microscopic dynamics in the magnetized stationary state is regular and explicitly known, an observation which contributed to shed light onto the puzzling abundance of emerging regular orbits as revealed in [5].

These last results have been though obtained in the

framework of mean field systems: The actual distance between particles does not explicitly appear in the HMF potential. In this letter we aim at bridging this gap, by focusing on the so called α -HMF model [6]. In its long range version ($\alpha < 1$ as discussed below), this model behaves at equilibrium as the HMF, see for instance [3, 7, 8]. We may then ask if the same correspondence applies to the out of equilibrium dynamics. QSS exists as depicted in[9], but, can one still appreciate the asymptotic trend towards regularity? How does the spatial organization impact on the aforementioned features? It was also shown recently that fractional calculus may be a crucial ingredient when dealing with long range systems [10], and we shall see how this point enters the picture in the considered case. To anticipate our findings, we shall here show that all stationary states of the HMF model are shared by the α -HMF model: Particles still exhibit regular orbits, at the price of a degree of enhanced microscopic spatial complexity, which materializes in a small scale inhomogeneity being locally scale invariant. Let us start by introducing the governing Hamiltonian which can be cast in the form:

$$H = \sum_{i=1}^N \left[\frac{p_i^2}{2} + \frac{1}{2\tilde{N}} \sum_{j \neq i}^N \frac{1 - \cos(q_i - q_j)}{\|i - j\|^\alpha} \right], \quad (1)$$

where q_i stands for the orientation of the rotor occupying the lattice position i , while p_i labels the conjugate momentum. The quantity $\|i - j\|$ denotes the shortest distance on the circle of perimeter $N - 1$. The coupling constant between classical rotators decays as a power law of the sites distance. The HMF limit is recovered for $\alpha = 0$. For N even, we have

$$\tilde{N} = \left(\frac{2}{N} \right)^\alpha + 2 \sum_{i=1}^{N/2-1} \frac{1}{i^\alpha}, \quad (2)$$

which guarantees extensivity of the system. The equation of motions of element i are derived from the above Hamiltonian and can be written as follows

$$\dot{p}_i = -\sin(q_i)C_i + \cos(q_i)S_i = M_i \sin(q_i - \varphi_i). \quad (3)$$

where use has been made of the following global quantities:

$$C_i = \frac{1}{\tilde{N}} \sum_{j \neq i} \frac{\cos q_j}{\|i - j\|^\alpha} \quad (4)$$

$$S_i = \frac{1}{\tilde{N}} \sum_{j \neq i} \frac{\sin q_j}{\|i - j\|^\alpha}, \quad (5)$$

These identify the two components of a non-local magnetization per site, with modulus $M_i = \sqrt{C_i^2 + S_i^2}$, and phase $\varphi_i = \arctan(S_i/C_i)$. In doing so, one brings into evidence the formal analogy with the HMF setting. Indeed we notice that each individual α -HMF particle obeys a dynamical equation which closely resembles that of a pendulum. This observation represented the starting point of the analysis carried out in [4], where stationary states were constructed from first principles. More specifically, the authors of [4] imagined that the system of coupled rotators reached a given equilibrium state, characterized by a constant magnetization in the limit for $N \rightarrow \infty$. Then, the HMF model is mapped into a set of N uncoupled pendula, constrained to collectively return a global magnetization identical to that driving their individual dynamics. Self-consistency is hence a crucial ingredient explicitly accommodated for the formulation proposed in [4]. Technically, the stationary state is built by exploiting the ergodic measure on the torus originating from the pendulum motion, a working ansatz that we cannot invoke in the context of the α -HMF, due to site localization. Eventually we will overcome this difficulty by considering the continuous limit. We notice that, for large N , and assuming $0 < \alpha < 1$, the following estimate applies

$$\tilde{N} \approx \frac{2}{1-\alpha} (N/2)^{1-\alpha}. \quad (6)$$

We can then use expression (6) in Eq.(4) and, as $N \rightarrow \infty$, introduce the continuous variables $x = i/N$ and $y = j/N$ and arrive to the following Riemann integral

$$C(x) = \frac{1-\alpha}{2^\alpha} \int_{-1/2}^{1/2} \frac{\cos(q(y))}{\|x-y\|^\alpha} dy, \quad (7)$$

where $\|x-y\|$ represents the minimal distance on a circle of perimeter one. By invoking the Riemann-Liouville formalism on the circle, we recognize the fractional integral $I^{1-\alpha}$ and consequently write

$$C(x) = \frac{1-\alpha}{2^\alpha} \Gamma(1-\alpha) I^{1-\alpha} (\cos q(x)). \quad (8)$$

A similar relation holds for the $S(x)$ component. Studying the α -HMF dynamics implies characterizing the evolution of the scalar fields $q(x,t)$ and $p(x,t)$ which are ruled by the fractional (non-local) partial differential

equations

$$\begin{aligned} \frac{\partial q}{\partial t} &= p(x,t) \\ \frac{\partial p}{\partial t} &= \frac{\mu}{2^\alpha} \Gamma(\mu) (-\sin(q) I^\mu (\cos q) + \cos(q) I^\mu (\sin q)). \end{aligned}$$

where $\mu = 1 - \alpha$. At variance with the simpler HMF ($\alpha = 0$) model, the spatial organization $q(x)$ matters within the general setting $\alpha > 0$, an observation which, as anticipated above, poses technical problems to a straightforward extension of the analysis in [4]. At first sight, it would be tempting to require $C(x) = \langle C \rangle = M$, if conditioning, at the same time, the system to $I^\mu (\sin q) = 0$. The translation invariance along the lattice is also likely to statistically lead to such a state[8]. For finite size systems, the latter identity can be solely matched by the trivial state where all q are equal. Conversely, in the infinite N limit, and recalling expression (8), we require that:

$$D^\alpha \cos q = \frac{d^\alpha \cos q}{dx^\alpha} = 0. \quad (9)$$

where the operator D^α stands for the fractional derivative. Trivial states as evidenced in the finite size approximation are also solutions of this equation. However, as we will argue in the following, the continuum limit enables us to compute an independent set of admissible solutions. This task is accomplished by exploiting the fact that $\alpha < 1$, which makes the integral $\int 1/\|x\|^\alpha dx$ convergent near 0 and that the function $1/\|x-y\|^\alpha$ is smooth. Noticing that we rewrite Eq. (7) as:

$$C(x) = \frac{1-\alpha}{2^\alpha} \sum_{k=0}^{L-1} \int_{k/L}^{(k+1)/L} \frac{\cos q(y)}{\|x-y\|^\alpha} dy,$$

using the regularity of $1/\|x-y\|^\alpha$, we can extract it from the integral:

$$C(x) \approx \frac{1-\alpha}{2^\alpha} \sum_{k=0}^{L-1} \frac{1}{\|x-y_k\|^\alpha} \int_{k/L}^{(k+1)/L} \cos q(y) dy,$$

with $y_k \in [k/L, (k+1)/L]$ [14]. We now assign the q values on a circle so that in any small interval the average of $\cos q(x)$ is constant and equal to the magnetization of the system. This procedure implies a peculiar spatial organization which returns a constant coarse grained image of the function $\cos q$, equal in turn to M . We have now a configuration which gives $C(x) = M$, a constant function on the circle. However we still need stationarity. This can be achieved by referring to the available stationary distribution for the HMF. The recipe goes as follows: we pick up q and p values as originating from this distribution, so as to ensure that the time evolution will only consist of a local reshuffling of the actual phase space coordinate, without affecting their associated coarse grained value. In doing so, we hence obtain a family of stationary solutions of the α -HMF model. [15]




Figure 1: QSS for the α -HMF model built using a stationary solution of the HMF model corresponding to a one particle PDF of the type $f(I, \theta) = \delta(I - I_0)/2\pi$ (one torus $M = 0.5$). The system is evolved via the α -HMF dynamics with $\alpha = 0.25$. The figure displays the resulting distribution of actions I at $t = 200$, for different sizes. The integration uses the fifth order optimal symplectic integrator [11] and a time step $\delta t = 0.05$.

The microscopic dynamics of the particles in such states can be mapped into a pendulum motion and is hence integrable. Interestingly, the spatial organization is locally scale free. The functions $q(x, t)$ and $p(x, t)$ are thus “very complicated” along the spatial direction, while displaying a regular time evolution and no chaos. In order to validate this result for a finite size sample on the original lattice we proceed as follows. We consider a stationary state of the HMF consisting of just one torus with associated magnetization $M = 0.5$. In pendulum action-angle variable we refer hence to a one particle distribution of the type $f(I, \theta) = \delta(I - I_0)/2\pi$. We are focusing our attention on an “individual component” belonging to the extended set of a linearly independent elements, which define the QSS basis. In order to be as close as possible to the stationary state of the α -HMF model, we simply distribute randomly on the lattice the values picked from such, analytically accessible, distribution. The analysis for different values of the number of sites and $\alpha = 0.25$, is then performed by monitoring the values of the action and shows as expected a trend towards the stabilization of such a state as N increases (see Fig. 1). We also checked that as expected increasing the value of α , which weakens the coupling strength, implies inducing a more pronounced destabilization of the state, which can be effectively opposed by increasing the number of simulated rotors. The solutions here constructed are hence stable versus the α -HMF dynamics, provided the continuum limit is being performed and so represent a consistent analytical prediction for the existence of quasi-stationary states, beyond the original HMF setting.

To further scrutinize the dynamics of the α -HMF model we turn to direct simulations. We initialize the system in $q = 0$ and a Gaussian distribution for the conjugate momenta p [7]. The system state is monitored by estimating the average magnetization amount as a




Figure 2: Magnetization vs. energy per particle E/N for different α , for a system made of $N = 4096$ particles. The magnetization values here reported follow from a time average over the window $400 < T < 800$.




Figure 3: Left: “Poincaré section” of a QSS for $E/N = 1.2$, $N = 65536$. Initial conditions are Gaussian in p and $q = 0$. The section is computed for $150 < t < 1000$ and $\alpha = 0.25$. Right: Magnetization versus time.

function of the energy, see Fig. 2. For energies larger than 0.75 one would expect the homogeneous solution to prevail, as dictated by the statistical mechanics calculation. However, the system gets confined into an inhomogeneous state, the residual time averaged magnetization being large and persistent in time. It is therefore tempting to interpret those states as QSS, and so analyze their associated dynamical features in light of the above conclusions. In particular, we expect the microscopic dynamics to resemble that of a pendulum, bearing some degree of intrinsic regularity. To unravel the phase space characteristics we compute the Poincaré sections, following the recipe in [5, 12] and so visualizing the single particle stroboscopic dynamics, with a rate of acquisition imposed by the self-consistent mean field evolution. The averages of C_i and S_i refer to the two components of the magnetization per site. The Poincaré sections are drawn by recording the positions p_i and $q_i - \varphi_i$ in phase space each time the equality $M_i = M$ is verified. Results for a specific initial conditions are depicted in Fig. 3 where one hundred trajectories are retained. The phase portrait shares many similarities with that obtained for a simple one and a half degree of freedom Hamiltonian (see for instance [13]), with many resonances and invariant tori. Clearly, and in agreement with the above scenario, a large number of particles exhibit regular dynamics. However as the nature of phase space reveals, these QSSs are




Figure 4: Top: $M(x) = \sqrt{C(x)^2 + S(x)^2}$ as a function of time. Bottom: individual particle's action as a function of time. The system is initialized to get a QSS with $E/N = 1.2$ and $\alpha = 0.25$ (see Fig. 3). Total number of particle is $N = 8192$, $x = i/N$, with $1 \leq i \leq N$. The actions are those of a pendulum corresponding to Eq.(3). $M(x)$ is almost uniform in space, while the individual spatial organization is complex. The time evolution of the actions appears quite regular.

steady state of the discrete dynamics, not stationary solutions. Nevertheless, we set to analyze the spatial organization of the identified steady state to test whether stationary state features are present in this configuration. To this end, we computed the values of the local magnetization $M(x, t) = \sqrt{C(x, t)^2 + S(x, t)^2}$ versus time and estimated an individual action, stemming from a Hamiltonian pendulum, which would give rise to an equation of motion formally identical to the Eq.(3). Results of the analysis are depicted in Fig. 4. One clearly sees that the function $M(x, t)$ is homogeneous and presents a modest

dependence on t , thus suggesting that the distribution of $q(x, t)$ is a solution of Eq.(9). The plot of action versus time as depicted in Fig. 4, clearly indicates a degree of enhanced spatial complexity, nearby particles not belonging to the same tori. We find in these simulations and in this (N - finite) steady state the same distinctive features of the stationary solutions as depicted earlier.

To conclude, in this Letter we have shown that stationary solutions for the mean field HMF model are as well stationary ones of the α -HMF model. Microscopic dynamics in these stationary states is regular, and explicitly known. The price to pay for this microscopic regularity in time is a complex, self similar, spatial organization corresponding to the solution of a fractional equation. When turning to direct numerical investigations we have identified a series of quasi-stationary states, which corresponds to steady states. Still, such states share many of the features of their stationary counterparts. The importance of these conclusions are manifold: On the one side, we confirm that QSS do exist in a generalized non mean field setting [9]. Also, we validate a theoretical method to construct, from first principle, (out of) equilibrium stationary solutions. Finally, the fact that in long range systems stationary states (among which we may of course count the equilibrium) which display regular microscopic dynamics do exist, allows us to dispose of an enormous amount of information regarding the intimate dynamics of a system frozen in such state. Looking forward, we may speculate that the knowledge of the microscopic dynamics can somehow enables one to challenge thermodynamics by setting up appropriate Maxwell daemons. Note also, that the regularity of the dynamics in the large N limit, induces from a statistical stand point a freezing of the system in a zero entropy state.

Acknowledgments

We are very grateful to S. Ruffo for pointing us to this problem and providing useful comments and remarks. X. L. would also like to thank F. Bouchet for fruitful and inspiring discussions.

-
- [1] *Dynamics and Thermodynamics of Systems with Long Range Interactions*, Vol. 602 of *Lect. Not. Phys.*, edited by T. Dauxois, S. Ruffo, E. Arimondo, and M. Wilkens (Springer-Verlag, Berlin, 2002).
 - [2] P. H. Chavanis, G. D. Ninno, D. Fanelli, and S. Ruffo, in *Chaos, Complexity and Transport*, edited by C. Chandre, X. Leoncini, and G. Zaslavsky (World Scientific, Singapore, 2008), pp. 3-26.
 - [3] A. Campa, T. Dauxois, and S. Ruffo, *Phys. Rep.* **480**, 57 (2009).
 - [4] X. Leoncini, T. L. Van den Berg, and D. Fanelli, *EPL* **86**, (2009).
 - [5] R. Bachelard *et al.*, *Phys. Rev. Lett.* **101**, (2008).
 - [6] C. Anteneodo and C. Tsallis, *Phys. Rev. Lett.* **80**, 5313 (1998).
 - [7] F. Tamarit and C. Anteneodo, *Phys. Rev. Lett.* **84**, 208 (2000).
 - [8] A. Campa, A. Giansanti, and D. Moroni, *J. Phys. A: Math. Gen.* **36**, 6897 (2003).
 - [9] A. Campa, A. Giansanti, and D. Moroni, *Physica A* **305**, 137 (2002).
 - [10] G. M. Zaslavsky, M. Edelman, and V. E. Tarasov, *Chaos* **17**, 043124 (2007).
 - [11] R. I. McLachlan and P. Atela, *Nonlinearity* **5**, 541 (1992).
 - [12] R. Bachelard *et al.*, *Eur. Phys. J. D* **42**, 125 (2007).
 - [13] G. M. Zaslavsky, *The Physics of Chaos in Hamiltonian*

Systems (Imperial College Press, London, 2007), second Edition.

- [14] The above approximation can be rigorously justified via a detailed expansion not reported here.
- [15] We note that many distinct stationary distributions of the original HMF model are in principle allowed, yielding to identical values of M . It should then be possible to place different distributions in distinct regions of space.

Thus stationary distributions of the α -HMF model can correspond to either a stationary distribution of the HMF model put scale free on the circle, or to decomposition of it in a sum of different stationary distributions, which all have the same magnetization and are put in a scale free form in different regions of the circle.