



Canonical bases and affine Hecke algebras of type D

Peng Shan, Michela Varagnolo, Eric Vasserot

► To cite this version:

Peng Shan, Michela Varagnolo, Eric Vasserot. Canonical bases and affine Hecke algebras of type D. 2009. hal-00442405v1

HAL Id: hal-00442405

<https://hal.science/hal-00442405v1>

Preprint submitted on 21 Dec 2009 (v1), last revised 29 Mar 2010 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CANONICAL BASES AND AFFINE HECKE ALGEBRAS OF TYPE D

P. SHAN, M. VARAGNOLO, E. VASSEROT

ABSTRACT. We prove a conjecture of Miemietz and Kashiwara on canonical bases and branching rules of affine Hecke algebras of type D. The proof is similar to the proof of the type B case in [VV].

INTRODUCTION

Let \mathbf{f} be the negative part of the quantized enveloping algebra of type $A^{(1)}$. Lusztig's description of the canonical basis of \mathbf{f} implies that this basis can be naturally identified with the set of isomorphism classes of simple objects of a category of modules of the affine Hecke algebras of type A. This identification was mentioned in [G], and was used in [A]. More precisely, there is a linear isomorphism between \mathbf{f} and the Grothendieck group of finite dimensional modules of the affine Hecke algebras of type A, and it is proved in [A] that the induction/restriction functors for affine Hecke algebras are given by the action of the Chevalley generators and their transposed operators with respect to some symmetric bilinear form on \mathbf{f} .

The branching rules for affine Hecke algebras of type B have been investigated quite recently, see [E], [EK1,2,3], [M] and [VV]. In [E], [EK1,2,3] a similar behavior is conjectured and studied for affine Hecke algebras of type B. Here \mathbf{f} is replaced by an explicit module ${}^\theta\mathbf{V}(\lambda)$ over an explicit algebra ${}^\theta\mathcal{B}$. More precisely it was conjectured that ${}^\theta\mathbf{V}(\lambda)$ admits a canonical basis which is naturally identified with the set of isomorphism classes of simple objects of a category of modules of the affine Hecke algebras of type B. Further, in this identification the branching rules of the affine Hecke algebras of type B should be given by the ${}^\theta\mathcal{B}$ -action on ${}^\theta\mathbf{V}(\lambda)$. This conjecture has been proved [VV]. It uses both the geometric picture introduced in [E] (to prove a part of the conjecture) and a new kind of graded algebras similar to the KLR algebras from [KL], [R].

A similar description of the branching rules for affine Hecke algebras of type D has also been conjectured in [KM]. In this case \mathbf{f} is replaced by another module ${}^\circ\mathbf{V}$ over the algebra ${}^\theta\mathcal{B}$ (the same algebra as in the type B case). The purpose of this paper is to prove this conjecture. The method of the proof is the same as in [VV]. First we introduce a family of graded algebras ${}^\circ\mathbf{R}_m$, $m \in \mathbb{Z}_{>0}$. They can be viewed as the Ext-algebras of some complex of constructible sheaves naturally attached to the Lie algebra of the group $SO(2m)$, see Remark 2.8. This complex enters in the Kazhdan-Lusztig classification of the simple modules of the affine Hecke algebra of the group $Spin(2m)$. Then we identify ${}^\circ\mathbf{V}$ with the sum of the Grothendieck groups of the graded algebras ${}^\circ\mathbf{R}_m$.

2000 *Mathematics Subject Classification*. Primary ??; Secondary ??.

The plan of the paper is the following. In Section 1 we introduce the graded algebra ${}^\circ\mathbf{R}(\Gamma)_\nu$. It is associated with a quiver Γ with an involution θ and with a dimension vector ν . In Section 2 we consider a particular choice of pair (Γ, θ) . Then the graded algebras ${}^\circ\mathbf{R}(\Gamma)_\nu$ specialize to the family of graded algebras ${}^\circ\mathbf{R}_m$. Next we introduce the affine Hecke algebra of type D and we prove that it is Morita equivalent to ${}^\circ\mathbf{R}_m$. In Section 3 we categorify the module ${}^\circ\mathbf{V}$ from [KM] using the graded algebras ${}^\circ\mathbf{R}_m$, see Theorem 3.28. The main result of the paper is Theorem 3.33.

0. NOTATION

0.1. Graded modules over graded algebras. Let \mathbf{k} be an algebraically closed field of characteristic 0. By a graded \mathbf{k} -algebra $\mathbf{R} = \bigoplus_i \mathbf{R}_i$ we'll always mean a \mathbb{Z} -graded associative \mathbf{k} -algebra. Let $\mathbf{R}\text{-mod}$ be the category of finitely generated graded \mathbf{R} -modules, $\mathbf{R}\text{-mod}^f$ be the full subcategory of finite-dimensional graded modules and $\mathbf{R}\text{-proj}$ be the full subcategory of projective objects. Unless specified otherwise all modules are left modules. We'll abbreviate

$$K(\mathbf{R}) = [\mathbf{R}\text{-proj}], \quad G(\mathbf{R}) = [\mathbf{R}\text{-mod}^f].$$

Here $[\mathcal{C}]$ denotes the Grothendieck group of an exact category \mathcal{C} . Assume that the \mathbf{k} -vector spaces \mathbf{R}_i are finite dimensional for each i . Then $K(\mathbf{R})$ is a free Abelian group with a basis formed by the isomorphism classes of the indecomposable objects in $\mathbf{R}\text{-proj}$, and $G(\mathbf{R})$ is a free Abelian group with a basis formed by the isomorphism classes of the simple objects in $\mathbf{R}\text{-mod}^f$. Given an object M of $\mathbf{R}\text{-proj}$ or $\mathbf{R}\text{-mod}^f$ let $[M]$ denote its class in $K(\mathbf{R})$, $G(\mathbf{R})$ respectively. When there is no risk of confusion we abbreviate $M = [M]$. We'll write $[M : N]$ for the composition multiplicity of the \mathbf{R} -module N in the \mathbf{R} -module M . Consider the ring $\mathcal{A} = \mathbb{Z}[v, v^{-1}]$. If the grading of \mathbf{R} is bounded below then the \mathcal{A} -modules $K(\mathbf{R})$, $G(\mathbf{R})$ are free. Here \mathcal{A} acts on $G(\mathbf{R})$, $K(\mathbf{R})$ as follows

$$vM = M[1], \quad v^{-1}M = M[-1].$$

For any M, N in $\mathbf{R}\text{-mod}$ let

$$\text{hom}_{\mathbf{R}}(M, N) = \bigoplus_d \text{Hom}_{\mathbf{R}}(M, N[d])$$

be the \mathbb{Z} -graded \mathbf{k} -vector space of all \mathbf{R} -module homomorphisms. If $\mathbf{R} = \mathbf{k}$ we'll omit the subscript \mathbf{R} in hom 's and in tensor products.

0.2. Quivers with involutions. Recall that a quiver Γ is a tuple $(I, H, h \mapsto h', h \mapsto h'')$ where I is the set of vertices, H is the set of arrows and for each $h \in H$ the vertices $h', h'' \in I$ are the origin and the goal of h respectively. Note that the set I may be infinite. We'll assume that no arrow may join a vertex to itself. For each $i, j \in I$ we write

$$H_{i,j} = \{h \in H; h' = i, h'' = j\}.$$

We'll abbreviate $i \rightarrow j$ if $H_{i,j} \neq \emptyset$. Let $h_{i,j}$ be the number of elements in $H_{i,j}$ and set

$$i \cdot j = -h_{i,j} - h_{j,i}, \quad i \cdot i = 2, \quad i \neq j.$$

An involution θ on Γ is a pair of involutions on I and H , both denoted by θ , such that the following properties hold for each h in H

- (a) $\theta(h)' = \theta(h'')$ and $\theta(h)'' = \theta(h')$,
- (b) $\theta(h') = h''$ iff $\theta(h) = h$.

We'll always assume that θ has no fixed points in I , i.e., there is no $i \in I$ such that $\theta(i) = i$. To simplify we'll say that θ has no fixed point. Let

$${}^\theta \mathbb{N}I = \{\nu = \sum_i \nu_i i \in \mathbb{N}I : \nu_{\theta(i)} = \nu_i, \forall i\}.$$

For any $\nu \in {}^\theta \mathbb{N}I$ set $|\nu| = \sum_i \nu_i$. It is an even integer. Write $|\nu| = 2m$ with $m \in \mathbb{N}$. We'll denote by ${}^\theta I^\nu$ the set of sequences

$$\mathbf{i} = (i_{1-m}, \dots, i_{m-1}, i_m)$$

of elements in I such that $\theta(i_l) = i_{1-l}$ and $\sum_k i_k = \nu$. For any such sequence \mathbf{i} we'll abbreviate $\theta(\mathbf{i}) = (\theta(i_{1-m}), \dots, \theta(i_{m-1}), \theta(i_m))$. Finally, we set

$${}^\theta I^m = \bigcup_{\nu} {}^\theta I^\nu, \quad \nu \in {}^\theta \mathbb{N}I, \quad |\nu| = 2m.$$

0.3. The wreath product. Given a positive integer m , let \mathfrak{S}_m be the symmetric group, and $\mathbb{Z}_2 = \{-1, 1\}$. Consider the wreath product $W_m = \mathfrak{S}_m \wr \mathbb{Z}_2$. Write s_1, \dots, s_{m-1} for the simple reflections in \mathfrak{S}_m . For each $l = 1, 2, \dots, m$ let $\varepsilon_l \in (\mathbb{Z}_2)^m$ be -1 placed at the l -th position. There is a unique action of W_m on the set $\{1-m, \dots, m-1, m\}$ such that \mathfrak{S}_m permutes $1, 2, \dots, m$ and such that ε_l fixes k if $k \neq l, 1-l$ and switches l and $1-l$. The group W_m acts also on ${}^\theta I^\nu$. Indeed, view a sequence \mathbf{i} as the map

$$\{1-m, \dots, m-1, m\} \rightarrow I, \quad l \mapsto i_l.$$

Then we set $w(\mathbf{i}) = \mathbf{i} \circ w^{-1}$ for $w \in W_m$. For each ν we fix once for all a sequence

$$\mathbf{i}_e = (i_{1-m}, \dots, i_m) \in {}^\theta I^\nu.$$

Let W_e be the centralizer of \mathbf{i}_e in W_m . Then there is a bijection

$$W_e \backslash W_m \rightarrow {}^\theta I^\nu, \quad W_e w \mapsto w^{-1}(\mathbf{i}_e).$$

Now, assume that $m > 1$. We set $s_0 = \varepsilon_1 s_1 \varepsilon_1$. Let ${}^\circ W_m$ be the subgroup of W_m generated by s_0, \dots, s_{m-1} . We'll regard it as a Weyl group of type D_m such that s_0, \dots, s_{m-1} are the simple reflections. Note that W_e is a subgroup of ${}^\circ W_m$. Indeed, if $W_e \not\subset {}^\circ W_m$ there should exist l such that ε_l belongs to W_e . This would imply that $i_l = \theta(i_l)$, contradicting the fact that θ has no fixed point. Therefore ${}^\theta I^\nu$ decomposes into two ${}^\circ W_m$ -orbits. We'll denote them by ${}^\theta I_+^\nu$ and ${}^\theta I_-^\nu$. For $m = 1$ we set ${}^\circ W_1 = \{e\}$.

1. THE GRADED \mathbf{k} -ALGEBRA ${}^\circ\mathbf{R}(\Gamma)_\nu$

Fix a quiver Γ with set of vertices I and set of arrows H . Fix an involution θ on Γ . Assume that Γ has no 1-loops and that θ has no fixed points. Fix a dimension vector $\nu \neq 0$ in ${}^\theta\mathbb{N}I$. Set $|\nu| = 2m$.

1.1. Definition of the graded \mathbf{k} -algebra ${}^\circ\mathbf{R}(\Gamma)_\nu$. Assume that $m > 1$. We define a graded \mathbf{k} -algebra ${}^\circ\mathbf{R}(\Gamma)_\nu$ with 1 generated by $1_{\mathbf{i}}$, \varkappa_l , σ_k , with $\mathbf{i} \in {}^\theta T^\nu$, $l = 1, 2, \dots, m$, $k = 0, 1, \dots, m-1$ modulo the following defining relations

$$\begin{aligned}
(a) \quad & 1_{\mathbf{i}} 1_{\mathbf{i}'} = \delta_{\mathbf{i}, \mathbf{i}'} 1_{\mathbf{i}}, \quad \sigma_k 1_{\mathbf{i}} = 1_{s_k(\mathbf{i})} \sigma_k, \quad \varkappa_l 1_{\mathbf{i}} = 1_{\mathbf{i}} \varkappa_l, \\
(b) \quad & \varkappa_l \varkappa_{l'} = \varkappa_{l'} \varkappa_l, \\
(c) \quad & \sigma_k^2 1_{\mathbf{i}} = Q_{i_k, i_{s_k(k)}}(\varkappa_{s_k(k)}, \varkappa_k) 1_{\mathbf{i}}, \\
(d) \quad & \sigma_k \sigma_{k'} = \sigma_{k'} \sigma_k \text{ if } 1 \leq k < k' - 1 < m - 1 \text{ or } 0 = k < k' \neq 2, \\
(e) \quad & (\sigma_{s_k(k)} \sigma_k \sigma_{s_k(k)} - \sigma_k \sigma_{s_k(k)} \sigma_k) 1_{\mathbf{i}} = \\
& = \begin{cases} \frac{Q_{i_k, i_{s_k(k)}}(\varkappa_{s_k(k)}, \varkappa_k) - Q_{i_k, i_{s_k(k)}}(\varkappa_{s_k(k)}, \varkappa_{s_k(k)+1})}{\varkappa_k - \varkappa_{s_k(k)+1}} 1_{\mathbf{i}} & \text{if } i_k = i_{s_k(k)+1}, \\ 0 & \text{else,} \end{cases} \\
(f) \quad & (\sigma_k \varkappa_l - \varkappa_{s_k(l)} \sigma_k) 1_{\mathbf{i}} = \begin{cases} -1_{\mathbf{i}} & \text{if } l = k, i_k = i_{s_k(k)}, \\ 1_{\mathbf{i}} & \text{if } l = s_k(k), i_k = i_{s_k(k)}, \\ 0 & \text{else.} \end{cases}
\end{aligned}$$

Here we have set $\varkappa_{1-l} = -\varkappa_l$ and

$$(1.1) \quad Q_{i,j}(u, v) = \begin{cases} (-1)^{h_{i,j}} (u - v)^{-i \cdot j} & \text{if } i \neq j, \\ 0 & \text{else.} \end{cases}$$

If $m = 0$ we set ${}^\circ\mathbf{R}(\Gamma)_0 = \mathbf{k} \oplus \mathbf{k}$. If $m = 1$ then we have $\nu = i + \theta(i)$ with $i \in I$. Write $\mathbf{i} = i\theta(i)$. We set

$${}^\circ\mathbf{R}(\Gamma)_\nu = \mathbf{k}[\varkappa_1] 1_{\mathbf{i}} \oplus \mathbf{k}[\varkappa_1] 1_{\theta(\mathbf{i})}.$$

We'll abbreviate $\sigma_{\mathbf{i},k} = \sigma_k 1_{\mathbf{i}}$ and $\varkappa_{\mathbf{i},l} = \varkappa_l 1_{\mathbf{i}}$. The grading on ${}^\circ\mathbf{R}(\Gamma)_0$ is the trivial one. For $m \geq 1$ the grading on ${}^\circ\mathbf{R}(\Gamma)_\nu$ is given by the following rules :

$$\begin{aligned}
\deg(1_{\mathbf{i}}) &= 0, \\
\deg(\varkappa_{\mathbf{i},l}) &= 2, \\
\deg(\sigma_{\mathbf{i},k}) &= -i_k \cdot i_{s_k(k)}.
\end{aligned}$$

We define ω to be the unique involution of the graded \mathbf{k} -algebra ${}^\circ\mathbf{R}(\Gamma)_\nu$ which fixes $1_{\mathbf{i}}$, \varkappa_l , σ_k . We set ω to be identity on ${}^\circ\mathbf{R}(\Gamma)_0$.

1.2. Relation with the graded \mathbf{k} -algebra ${}^\theta\mathbf{R}(\Gamma)_\nu$. A family of graded \mathbf{k} -algebra ${}^\theta\mathbf{R}(\Gamma)_{\lambda,\nu}$ was introduced in [VV, sec. 5.1], where λ is an arbitrary dimension vector. Here we'll only consider the special case $\lambda = 0$ and abbreviate ${}^\theta\mathbf{R}(\Gamma)_\nu = {}^\theta\mathbf{R}(\Gamma)_{0,\nu}$. Recall that ${}^\theta\mathbf{R}(\Gamma)_\nu$ is the graded \mathbf{k} -algebra with 1 generated by $1_{\mathbf{i}}, \varkappa_l, \sigma_k, \pi_1$, with $\mathbf{i} \in {}^\theta I^\nu$, $l = 1, 2, \dots, m$, $k = 1, \dots, m-1$ such that $1_{\mathbf{i}}, \varkappa_l$ and σ_k satisfy the same relations as before and

$$\pi_1^2 = 1, \quad \pi_1 1_{\mathbf{i}} \pi_1 = 1_{\varepsilon_1 \mathbf{i}}, \quad \pi_1 \varkappa_l \pi_1 = \varkappa_{\varepsilon_1(l)},$$

$$(\pi_1 \sigma_1)^2 = (\sigma_1 \pi_1)^2, \quad \pi_1 \sigma_k \pi_1 = \sigma_k \text{ if } k \neq 1.$$

If $\nu = 0$ then ${}^\theta\mathbf{R}(\Gamma)_0 = \mathbf{k}$. The grading is given by setting $\deg(1_{\mathbf{i}})$, $\deg(\varkappa_{l,l})$, $\deg(\sigma_{\mathbf{i},k})$ to be as before and $\deg(\pi_1 1_{\mathbf{i}}) = 0$. In the rest of Section 1 we'll assume $m > 0$. There is a canonical inclusion of graded \mathbf{k} -algebras

$$(1.2) \quad {}^\circ\mathbf{R}(\Gamma)_\nu \subset {}^\theta\mathbf{R}(\Gamma)_\nu$$

such that $1_{\mathbf{i}}, \varkappa_l, \sigma_k \mapsto 1_{\mathbf{i}}, \varkappa_l, \sigma_k$ for $\mathbf{i} \in {}^\theta I^\nu$, $l = 1, \dots, m$, $k = 1, \dots, m-1$ and such that $\sigma_0 \mapsto \pi_1 \sigma_1 \pi_1$. From now on we'll write $\sigma_0 = \pi_1 \sigma_1 \pi_1$ for all $m > 1$. The assignment $x \mapsto \pi_1 x \pi_1$ defines an involution of the graded \mathbf{k} -algebra ${}^\theta\mathbf{R}(\Gamma)_\nu$ which normalizes ${}^\circ\mathbf{R}(\Gamma)_\nu$. Thus it yields an involution

$$\gamma : {}^\circ\mathbf{R}(\Gamma)_\nu \rightarrow {}^\circ\mathbf{R}(\Gamma)_\nu.$$

Let $\langle \gamma \rangle$ be the group of two elements generated by γ . The smash product ${}^\circ\mathbf{R}(\Gamma)_\nu \rtimes \langle \gamma \rangle$ is a graded \mathbf{k} -algebra such that $\deg(\gamma) = 0$. There is an unique isomorphism of graded \mathbf{k} -algebras

$$(1.3) \quad {}^\circ\mathbf{R}(\Gamma)_\nu \rtimes \langle \gamma \rangle \rightarrow {}^\theta\mathbf{R}(\Gamma)_\nu$$

which is identity on ${}^\circ\mathbf{R}(\Gamma)_\nu$ and which takes γ to π_1 .

1.3. The polynomial representation and the PBW theorem. For any \mathbf{i} in ${}^\theta I^\nu$ let ${}^\theta\mathbf{F}_{\mathbf{i}}$ be the subalgebra of ${}^\circ\mathbf{R}(\Gamma)_\nu$ generated by $1_{\mathbf{i}}$ and $\varkappa_{l,l}$ with $l = 1, 2, \dots, m$. It is a polynomial algebra. Let

$${}^\theta\mathbf{F}_\nu = \bigoplus_{\mathbf{i} \in {}^\theta I^\nu} {}^\theta\mathbf{F}_{\mathbf{i}}.$$

The group W_m acts on ${}^\theta\mathbf{F}_\nu$ via $w(\varkappa_{l,l}) = \varkappa_{w(\mathbf{i}),w(l)}$ for any $w \in W_m$. Write

$${}^\circ\mathbf{S}_\nu = ({}^\theta\mathbf{F}_\nu)^{{}^\circ W_m}.$$

Regard ${}^\theta\mathbf{R}(\Gamma)_\nu$ and $\text{End}({}^\theta\mathbf{F}_\nu)$ as ${}^\theta\mathbf{F}_\nu$ -algebras via the left multiplication. In [VV, prop. 5.4] is given an injective graded ${}^\theta\mathbf{F}_\nu$ -algebra morphism ${}^\theta\mathbf{R}(\Gamma)_\nu \rightarrow \text{End}({}^\theta\mathbf{F}_\nu)$. It restricts via (1.2) to an injective graded ${}^\theta\mathbf{F}_\nu$ -algebra morphism

$${}^\circ\mathbf{R}(\Gamma)_\nu \rightarrow \text{End}({}^\theta\mathbf{F}_\nu).$$

Recall that ${}^\circ W_m$ is the Weyl group of type D_m with simple reflections s_0, \dots, s_{m-1} . For each w in ${}^\circ W_m$ we choose a reduced decomposition \dot{w} of w . It has the following form

$$w = s_{k_1} s_{k_2} \cdots s_{k_r}, \quad 0 \leq k_1, k_2, \dots, k_r \leq m-1.$$

We define an element $\sigma_{\dot{w}}$ in ${}^\circ\mathbf{R}(\Gamma)_\nu$ by

$$(1.4) \quad \sigma_{\dot{w}} = \sum_{\mathbf{i}} 1_{\mathbf{i}} \sigma_{\dot{w}}, \quad 1_{\mathbf{i}} \sigma_{\dot{w}} = \begin{cases} 1_{\mathbf{i}} & \text{if } r = 0 \\ 1_{\mathbf{i}} \sigma_{k_1} \sigma_{k_2} \cdots \sigma_{k_r} & \text{else,} \end{cases}$$

Observe that $\sigma_{\dot{w}}$ may depend on the choice of the reduced decomposition \dot{w} .

1.4. Proposition. *The \mathbf{k} -algebra ${}^\circ\mathbf{R}(\Gamma)_\nu$ is a free (left or right) ${}^\theta\mathbf{F}_\nu$ -module with basis $\{\sigma_{\dot{w}}; w \in {}^\circ W_m\}$. Its rank is $2^{m-1}m!$. The operator $1_{\mathbf{i}}\sigma_{\dot{w}}$ is homogeneous and its degree is independent of the choice of the reduced decomposition \dot{w} .*

Proof: This is the same proof as in [VV, prop. 5.5]. The algebra ${}^\circ\mathbf{R}(\Gamma)_\nu$ is filtered with $1_{\mathbf{i}}, \varkappa_{\mathbf{i},l}$ in degree 0 and $\sigma_{\mathbf{i},k}$ in degree 1. The Nil Hecke algebra of type D_m is the \mathbf{k} -algebra ${}^\circ\mathbf{NH}_m$ generated by $\bar{\sigma}_0, \bar{\sigma}_1, \dots, \bar{\sigma}_{m-1}$ with relations

$$\bar{\sigma}_k \bar{\sigma}_{k'} = \bar{\sigma}_{k'} \bar{\sigma}_k \text{ if } 1 \leq k < k' - 1 < m - 1 \text{ or } 0 = k < k' \neq 2,$$

$$\bar{\sigma}_{s_k(k)} \bar{\sigma}_k \bar{\sigma}_{s_k(k)} = \bar{\sigma}_k \bar{\sigma}_{s_k(k)} \bar{\sigma}_k, \quad \bar{\sigma}_k^2 = 0.$$

We can form the semidirect product ${}^\theta\mathbf{F}_\nu \rtimes {}^\circ\mathbf{NH}_m$, which is generated by $1_{\mathbf{i}}, \bar{\varkappa}_l, \bar{\sigma}_k$ with the relations above and

$$\bar{\sigma}_k \bar{\varkappa}_l = \bar{\varkappa}_{s_k(l)} \bar{\sigma}_k, \quad \bar{\varkappa}_l \bar{\varkappa}_{l'} = \bar{\varkappa}_{l'} \bar{\varkappa}_l.$$

One proves as in [VV, prop. 5.5] that the map

$${}^\theta\mathbf{F}_\nu \rtimes {}^\circ\mathbf{NH}_m \rightarrow \text{gr}({}^\circ\mathbf{R}(\Gamma)_\nu), \quad 1_{\mathbf{i}} \mapsto 1_{\mathbf{i}}, \quad \bar{\varkappa}_l \mapsto \varkappa_l, \quad \bar{\sigma}_k \mapsto \sigma_k.$$

is an isomorphism of \mathbf{k} -algebras. \square

Let ${}^\theta\mathbf{F}'_\nu = \bigoplus_{\mathbf{i}} {}^\theta\mathbf{F}'_{\mathbf{i}}$, where ${}^\theta\mathbf{F}'_{\mathbf{i}}$ is the localization of the ring ${}^\theta\mathbf{F}_{\mathbf{i}}$ with respect to the multiplicative system generated by

$$\{\varkappa_{\mathbf{i},l} \pm \varkappa_{\mathbf{i},l'}; 1 \leq l \neq l' \leq m\} \cup \{\varkappa_{\mathbf{i},l}; l = 1, 2, \dots, m\}.$$

1.5. Corollary. *There is an inclusion ${}^\circ\mathbf{R}(\Gamma)_\nu \subset \text{End}({}^\theta\mathbf{F}_\nu)$ which yields an isomorphism of ${}^\theta\mathbf{F}'_\nu$ -algebras ${}^\theta\mathbf{F}'_\nu \otimes_{{}^\theta\mathbf{F}_\nu} {}^\circ\mathbf{R}(\Gamma)_\nu \rightarrow {}^\theta\mathbf{F}'_\nu \rtimes {}^\circ W_m$, such that for each \mathbf{i} and each $l = 1, 2, \dots, m, k = 0, 1, 2, \dots, m-1$ we have*

$$(1.5) \quad \begin{aligned} 1_{\mathbf{i}} &\mapsto 1_{\mathbf{i}}, \\ \varkappa_{\mathbf{i},l} &\mapsto \varkappa_l 1_{\mathbf{i}}, \\ \sigma_{\mathbf{i},k} &\mapsto \begin{cases} (\varkappa_k - \varkappa_{s_k(k)})^{-1} (s_k - 1) 1_{\mathbf{i}} & \text{if } i_k = i_{s_k(k)}, \\ (\varkappa_k - \varkappa_{s_k(k)})^{h_{i_{s_k(k)}, i_k}} s_k 1_{\mathbf{i}} & \text{if } i_k \neq i_{s_k(k)}. \end{cases} \end{aligned}$$

Proof: This follows from [VV, cor. 5.6] and Proposition 1.4. \square

Restricting the ${}^\theta\mathbf{F}_\nu$ -action on ${}^\circ\mathbf{R}(\Gamma)_\nu$ to the \mathbf{k} -subalgebra ${}^\circ\mathbf{S}_\nu$ we get a structure of graded ${}^\circ\mathbf{S}_\nu$ -algebra on ${}^\circ\mathbf{R}(\Gamma)_\nu$.

1.6. Proposition. (a) ${}^\circ\mathbf{S}_\nu$ is isomorphic to the center of ${}^\circ\mathbf{R}(\Gamma)_\nu$.

(b) ${}^\circ\mathbf{R}(\Gamma)_\nu$ is a free graded module over ${}^\circ\mathbf{S}_\nu$ of rank $(2^{m-1}m!)^2$.

Proof: Part (a) follows from Corollary 1.5. Part (b) follows from (a) and Proposition 1.4. \square

2. AFFINE HECKE ALGEBRAS OF TYPE D

2.1. Affine Hecke algebras of type D. Fix p in \mathbf{k}^\times . For any integer $m \geq 0$ we define the extended affine Hecke algebra \mathbf{H}_m of type D_m as follows. If $m > 1$ then \mathbf{H}_m is the \mathbf{k} -algebra with 1 generated by

$$T_k, \quad X_l^{\pm 1}, \quad k = 0, 1, \dots, m-1, \quad l = 1, 2, \dots, m$$

satisfying the following defining relations :

- (a) $X_l X_{l'} = X_{l'} X_l$,
- (b) $T_k T_{s_k(k)} T_k = T_{s_k(k)} T_k T_{s_k(k)}$, $T_k T_{k'} = T_{k'} T_k$ ($1 \leq k < k' - 1$ or $k = 0, k' \neq 2$),
- (c) $(T_k - p)(T_k + p^{-1}) = 0$,
- (d) $T_0 X_1^{-1} T_0 = X_2$, $T_k X_k T_k = X_{s_k(k)}$ ($k \neq 0$), $T_k X_l = X_l T_k$ ($k \neq 0, l, l-1$ or $k = 0, l \neq 1, 2$).

We set $\mathbf{H}_0 = \mathbf{k} \oplus \mathbf{k}$ and $\mathbf{H}_1 = \mathbf{k}[X_1^{\pm 1}]$.

2.2. Remarks. (a) The extended affine Hecke algebra \mathbf{H}_m^B of type B_m with parameters $p \in \mathbf{k}^\times$ and $q = 1$ is generated by $P, T_k, X_l^{\pm 1}$, $k = 1, \dots, m-1, l = 1, \dots, m$ such that $T_k, X_l^{\pm 1}$ satisfy the relations as above and

$$P^2 = 1, \quad (PT_1)^2 = (T_1 P)^2, \quad PT_k = T_k P \text{ if } k \neq 1, \\ PX_1^{-1} P = X_1, \quad PX_l = X_l P \text{ if } l \neq 1.$$

There is an obvious \mathbf{k} -algebra embedding $\mathbf{H}_m \subset \mathbf{H}_m^B$. Let γ denote also the involution $\mathbf{H}_m \rightarrow \mathbf{H}_m$, $a \mapsto PaP$. We have a canonical isomorphism of \mathbf{k} -algebras

$$\mathbf{H}_m \rtimes \langle \gamma \rangle \simeq \mathbf{H}_m^B.$$

(b) Note that \mathbf{H}_m is the extended affine Hecke algebra associated with the group $SO(2m)$. Let \mathbf{H}_m^e be the extended affine Hecke algebra associated with $Spin(2m)$. It is generated by \mathbf{H}_m and an element X_0 such that

$$X_0^2 = X_1 X_2 \dots X_m, \quad T_k X_0 = X_0 T_k \text{ if } k \neq 0, \quad T_0 X_0 X_1^{-1} X_2^{-1} T_0 = X_0.$$

Thus \mathbf{H}_m is the fixed point subset of the \mathbf{k} -algebra automorphism of \mathbf{H}_m^e taking T_k, X_l to $T_k, (-1)^{\delta_{l,0}} X_l$ for all k, l . If p is not a root of 1 the simple \mathbf{H}_m -modules can be recovered from the Kazhdan-Lusztig classification of the simple \mathbf{H}_m^e -modules via Clifford theory, see e.g., [Re].

2.3. Intertwiners and blocks of \mathbf{H}_m . We define

$$\mathbf{A} = \mathbf{k}[X_1^{\pm 1}, X_2^{\pm 1}, \dots, X_m^{\pm 1}], \quad \mathbf{A}' = \mathbf{A}[\Sigma^{-1}], \quad \mathbf{H}'_m = \mathbf{A}' \otimes_{\mathbf{A}} \mathbf{H}_m,$$

where Σ is the multiplicative set generated by

$$1 - X_l X_{l'}^{\pm 1}, \quad 1 - p^2 X_l^{\pm 1} X_{l'}^{\pm 1}, \quad l \neq l'.$$

For $k = 0, \dots, m-1$ the intertwiner φ_k in \mathbf{H}'_m is given by the following formulas

$$(2.1) \quad \varphi_k - 1 = \frac{X_k - X_{s_k(k)}}{pX_k - p^{-1}X_{s_k(k)}} (T_k - p)$$

The group ${}^\circ W_m$ acts on \mathbf{A}' as follows

$$(s_k a)(X_1, \dots, X_m) = a(X_1, \dots, X_{k+1}, X_k, \dots, X_m) \text{ if } k \neq 0$$

$$(s_0 a)(X_1, \dots, X_m) = a(X_2^{-1}, X_1^{-1}, \dots, X_m).$$

There is an isomorphism of \mathbf{A}' -algebras

$$\mathbf{A}' \rtimes {}^\circ W_m \rightarrow \mathbf{H}'_m, \quad s_k \mapsto \varphi_k.$$

The semi-direct product group $\mathbb{Z} \rtimes \mathbb{Z}_2 = \mathbb{Z} \rtimes \{-1, 1\}$ acts on \mathbf{k}^\times by $(n, \varepsilon) : z \mapsto z^\varepsilon p^{2n}$. Given a $\mathbb{Z} \rtimes \mathbb{Z}_2$ -invariant subset I of \mathbf{k}^\times we denote by $\mathbf{H}_m\text{-Mod}_I$ the category of all \mathbf{H}_m -modules such that the action of X_1, X_2, \dots, X_m is locally finite with eigenvalues in I . We associate to the set I and to the element $p \in \mathbf{k}^\times$ a quiver Γ as follows. The set of vertices is I and there is one arrow $i \rightarrow p^2 i$ whenever $i, p^2 i$ lie in I . We equip Γ with an involution θ such that $\theta(i) = i^{-1}$ for each vertex i and such that θ takes the arrow $i \rightarrow p^2 i$ to the arrow $p^{-2} i^{-1} \rightarrow i^{-1}$. *We'll assume that the set I does not contain 1 nor -1 and that $p \neq 1, -1$.* Thus the involution θ has no fixed points and no arrow may join a vertex of Γ to itself.

2.4. Remark. We may assume that $I = \pm\{p^n; n \in \mathbb{Z}_{\text{odd}}\}$. See the discussion in [KM]. Then Γ is of type A_∞ if p has infinite order and Γ is of type $A_r^{(1)}$ if p^2 is a primitive r -th root of unity.

2.5. \mathbf{H}_m -modules versus ${}^\circ \mathbf{R}_m$ -modules. Assume that $m \geq 1$. We define the graded \mathbf{k} -algebra

$${}^\circ \mathbf{R}_{I,m} = \bigoplus_{\nu} {}^\circ \mathbf{R}_{I,\nu}, \quad {}^\circ \mathbf{R}_{I,\nu} = {}^\circ \mathbf{R}(\Gamma)_\nu, \quad \theta_I^m = \coprod_{\nu} \theta_{I^\nu},$$

where ν runs over the set of all dimension vectors in ${}^\theta \mathbb{N}I$ such that $|\nu| = 2m$. When there is no risk of confusion we abbreviate ${}^\circ \mathbf{R}_\nu = {}^\circ \mathbf{R}_{I,\nu}$ and ${}^\circ \mathbf{R}_m = {}^\circ \mathbf{R}_{I,m}$. Note that the \mathbf{k} -algebra ${}^\circ \mathbf{R}_m$ may not have 1, because the set I may be infinite. We define ${}^\circ \mathbf{R}_m\text{-Mod}_0$ as the category of all (non-graded) ${}^\circ \mathbf{R}_m$ -modules such that the elements $\varkappa_1, \varkappa_2, \dots, \varkappa_m$ act locally nilpotently. Let ${}^\circ \mathbf{R}_m\text{-Mod}_0^f$ and $\mathbf{H}_m\text{-Mod}_I^f$ be the full subcategories of finite dimensional modules in ${}^\circ \mathbf{R}_m\text{-Mod}_0$ and $\mathbf{H}_m\text{-Mod}_I$ respectively.

2.6. Theorem. *We have an equivalence of categories*

$${}^\circ \mathbf{R}_m\text{-Mod}_0 \rightarrow \mathbf{H}_m\text{-Mod}_I, \quad M \mapsto M$$

which is given by

- (a) X_l acts on $1_i M$ by $i_l^{-1} e^{\varkappa_l}$ for each $l = 1, 2, \dots, m$,

(b) if $m > 1$ then T_k acts on $1_1 M$ as follows for each $k = 0, 1, \dots, m-1$,

$$\begin{aligned} & \frac{(pe^{\mathfrak{x}_k} - p^{-1}e^{\mathfrak{x}_{s_k(k)}})(\mathfrak{x}_k - \mathfrak{x}_{s_k(k)})}{e^{\mathfrak{x}_k} - e^{\mathfrak{x}_{s_k(k)}}} \sigma_k + p & \text{if } i_k = i_{s_k(k)}, \\ & \frac{e^{\mathfrak{x}_k} - e^{\mathfrak{x}_{s_k(k)}}}{(p^{-1}e^{\mathfrak{x}_k} - pe^{\mathfrak{x}_{s_k(k)}})(\mathfrak{x}_k - \mathfrak{x}_{s_k(k)})} \sigma_k - \frac{(p^2 - 1)e^{\mathfrak{x}_{s_k(k)}}}{p^{-1}e^{\mathfrak{x}_k} - pe^{\mathfrak{x}_{s_k(k)}}} & \text{if } i_k = p^2 i_{s_k(k)}, \\ & \frac{pi_{s_k(k)}e^{\mathfrak{x}_k} - p^{-1}i_k e^{\mathfrak{x}_{s_k(k)}}}{i_{s_k(k)}e^{\mathfrak{x}_k} - i_k e^{\mathfrak{x}_{s_k(k)}}} (\sigma_k - 1) + p & \text{if } i_k \neq i_{s_k(k)}, p^2 i_{s_k(k)}. \end{aligned}$$

Proof : This follows from [VV, thm. 6.5] by Section 1.2 and Remark 2.2(a). One can also prove it by reproducing the arguments in loc. cit. by using (1.5) and (2.1). \square

2.7. Corollary. *There is an equivalence of categories*

$$\Psi : {}^\circ \mathbf{R}_m\text{-Mod}_0^f \rightarrow \mathbf{H}_m\text{-Mod}_I^f, \quad M \mapsto M.$$

2.8. Remark. Let \mathfrak{g} be the Lie algebra of $G = SO(2m)$. Fix a maximal torus $T \subset G$. The group of characters of T is the lattice $\bigoplus_{l=1}^m \mathbb{Z} \varepsilon_l$, with Bourbaki's notation. Fix a dimension vector $\nu \in {}^\theta \mathbb{N}I$. Recall the sequence $\mathbf{i}_e = (i_{1-m}, \dots, i_{m-1}, i_m)$ from Section 0.3. Let $g \in T$ be the element such that $\varepsilon_l(g) = i_l^{-1}$ for each $l = 1, 2, \dots, m$. Recall the notation ${}^\theta \mathcal{V}_\nu$, \mathbf{V} , ${}^\theta E_\mathbf{V}$, ${}^\theta G_\mathbf{V}$ from [VV]. Here \mathbf{V} is an object of ${}^\theta \mathcal{V}_\nu$. Then ${}^\theta G_\mathbf{V} = G_g$, the centralizer of g in G , and

$${}^\theta E_\mathbf{V} = \mathfrak{g}_{g,p}, \quad \mathfrak{g}_{g,p} = \{x \in \mathfrak{g}; \text{ad}_g(x) = p^2 x\}.$$

Let F_g be the set of all Borel Lie subalgebras of \mathfrak{g} fixed by the adjoint action of g . It is non connected manifold whose connected components are labelled by ${}^\theta I_+^\nu$. In Section 3.14 we introduce two central idempotents $1_{\nu,+}$, $1_{\nu,-}$ of ${}^\circ \mathbf{R}_\nu$. This yields a graded \mathbf{k} -algebra decomposition ${}^\circ \mathbf{R}_\nu = {}^\circ \mathbf{R}_\nu 1_{\nu,+} \oplus {}^\circ \mathbf{R}_\nu 1_{\nu,-}$. By [VV, thm. 5.8] the graded \mathbf{k} -algebra ${}^\circ \mathbf{R}_\nu 1_{\nu,+}$ is isomorphic to

$$\text{Ext}_{G_g}^*(\mathcal{L}_{g,p}, \mathcal{L}_{g,p}),$$

where $\mathcal{L}_{g,p}$ is the direct image of the constant perverse sheaf by the projection

$$\{(\mathfrak{b}, x) \in F_g \times \mathfrak{g}_{g,p}; x \in \mathfrak{b}\} \rightarrow \mathfrak{g}_{g,p}, \quad (\mathfrak{b}, x) \mapsto x.$$

The perverse sheaves on $\mathfrak{g}_{g,p}$ have been extensively studied by Lusztig, see e.g., [L1], [L2]. We hope to come back to this elsewhere.

2.9. Induction and restriction of \mathbf{H}_m -modules. For $i \in I$ we define functors

$$(2.2) \quad \begin{aligned} E_i &: \mathbf{H}_{m+1}\text{-Mod}_I^f \rightarrow \mathbf{H}_m\text{-Mod}_I^f, \\ F_i &: \mathbf{H}_m\text{-Mod}_I^f \rightarrow \mathbf{H}_{m+1}\text{-Mod}_I^f, \end{aligned}$$

where $E_i M \subset M$ is the generalized $\theta(i)$ -eigenspace of the X_{m+1} -action, and where

$$F_i M = \text{Ind}_{\mathbf{H}_m \otimes \mathbf{k}[X_{m+1}^{\pm 1}]}^{\mathbf{H}_{m+1}} (M \otimes \mathbf{k}_i).$$

Here \mathbf{k}_i is the 1-dimensional representation of $\mathbf{k}[X_{m+1}^{\pm 1}]$ defined by $X_{m+1} \mapsto \theta(i)$.

3. GLOBAL BASES OF ${}^\circ\mathbf{V}$ AND PROJECTIVE GRADED ${}^\circ\mathbf{R}$ -MODULES

3.1. The Grothendieck groups of ${}^\circ\mathbf{R}_m$. The graded \mathbf{k} -algebra ${}^\circ\mathbf{R}_m$ is free of finite rank over its center, a commutative graded \mathbf{k} -subalgebra. Therefore any simple object of ${}^\circ\mathbf{R}_m\text{-mod}$ is finite-dimensional and there is a finite number of isomorphism classes of simple modules in ${}^\circ\mathbf{R}_m\text{-mod}$. The Abelian group $G({}^\circ\mathbf{R}_m)$ is free with a basis formed by the classes of the simple objects of ${}^\circ\mathbf{R}_m\text{-mod}$. The Abelian group $K({}^\circ\mathbf{R}_m)$ is free with a basis formed by the classes of the indecomposable projective objects. Both are free \mathcal{A} -modules where v shifts the grading by 1. We consider the following \mathcal{A} -modules

$$\begin{aligned} {}^\circ\mathbf{K}_I &= \bigoplus_{m \geq 0} {}^\circ\mathbf{K}_{I,m}, & {}^\circ\mathbf{K}_{I,m} &= K({}^\circ\mathbf{R}_m), \\ {}^\circ\mathbf{G}_I &= \bigoplus_{m \geq 0} {}^\circ\mathbf{G}_{I,m}, & {}^\circ\mathbf{G}_{I,m} &= G({}^\circ\mathbf{R}_m). \end{aligned}$$

We'll also abbreviate

$${}^\circ\mathbf{K}_{I,*} = \bigoplus_{m > 0} {}^\circ\mathbf{K}_{I,m}, \quad {}^\circ\mathbf{G}_{I,*} = \bigoplus_{m > 0} {}^\circ\mathbf{G}_{I,m}.$$

From now on, to unburden the notation we may abbreviate ${}^\circ\mathbf{R} = {}^\circ\mathbf{R}_m$, hoping it will not create any confusion. For any M, N in ${}^\circ\mathbf{R}\text{-mod}$ we set

$$(M : N) = \text{gdim}(M^\omega \otimes_{{}^\circ\mathbf{R}} N), \quad \langle M : N \rangle = \text{gdim} \text{hom}_{{}^\circ\mathbf{R}}(M, N).$$

The Cartan pairing is the perfect \mathcal{A} -bilinear form

$${}^\circ\mathbf{K}_I \times {}^\circ\mathbf{G}_I \rightarrow \mathcal{A}, \quad (P, M) \mapsto \langle P : M \rangle.$$

First, we concentrate on the \mathcal{A} -module ${}^\circ\mathbf{G}_I$. Consider the duality

$${}^\circ\mathbf{R}\text{-mod}^f \rightarrow {}^\circ\mathbf{R}\text{-mod}^f, \quad M \mapsto M^\flat = \text{hom}(M, \mathbf{k}),$$

with the action and the grading given by

$$(xf)(m) = f(\omega(x)m), \quad (M^\flat)_i = \text{hom}(M_{-i}, \mathbf{k}).$$

This duality functor yields an \mathcal{A} -antilinear map

$${}^\circ\mathbf{G}_I \rightarrow {}^\circ\mathbf{G}_I, \quad M \mapsto M^\flat.$$

Let ${}^\circ B$ denote the set of isomorphism classes of simple objects of ${}^\circ\mathbf{R}\text{-Mod}_0^f$. We can now define the upper global basis of ${}^\circ\mathbf{G}_I$ as follows. The proof is given in Section 3.21.

3.2. Proposition/Definition. *For each b in ${}^\circ B$ there is a unique selfdual irreducible graded ${}^\circ\mathbf{R}$ -module ${}^\circ G^{\text{up}}(b)$ which is isomorphic to b as a (non graded) ${}^\circ\mathbf{R}$ -module. We set ${}^\circ G^{\text{up}}(0) = 0$ and ${}^\circ\mathbf{G}^{\text{up}} = \{{}^\circ G^{\text{up}}(b); b \in {}^\circ B\}$. Hence ${}^\circ\mathbf{G}^{\text{up}}$ is a \mathcal{A} -basis of ${}^\circ\mathbf{G}_I$.*

Now, we concentrate on the \mathcal{A} -module ${}^\circ\mathbf{K}_I$. We equip ${}^\circ\mathbf{K}_I$ with the symmetric \mathcal{A} -bilinear form

$$(3.1) \quad {}^\circ\mathbf{K}_I \times {}^\circ\mathbf{K}_I \rightarrow \mathcal{A}, \quad (M, N) \mapsto (M : N).$$

Consider the duality

$${}^\circ\mathbf{R}\text{-}\mathbf{proj} \rightarrow {}^\circ\mathbf{R}\text{-}\mathbf{proj}, \quad P \mapsto P^\sharp = \text{hom}_{{}^\circ\mathbf{R}}(P, {}^\circ\mathbf{R}),$$

with the action and the grading given by

$$(xf)(p) = f(p)\omega(x), \quad (P^\sharp)_i = \text{hom}(P_i, \mathbf{k}).$$

This duality functor yields an \mathcal{A} -antilinear map

$${}^\circ\mathbf{K}_I \rightarrow {}^\circ\mathbf{K}_I, \quad P \mapsto P^\sharp.$$

Set $\mathcal{K} = \mathbb{Q}(v)$. Let $\mathcal{K} \rightarrow \mathcal{K}$, $f \mapsto \bar{f}$ be the unique involution such that $\bar{v} = v^{-1}$.

3.3. Definition. For each b in ${}^\circ B$ let ${}^\circ G^{\text{low}}(b)$ be the unique indecomposable graded module in ${}^\circ\mathbf{R}\text{-}\mathbf{proj}$ whose top is isomorphic to ${}^\circ G^{\text{up}}(b)$. We set ${}^\circ G^{\text{low}}(0) = 0$ and ${}^\circ \mathbf{G}^{\text{low}} = \{{}^\circ G^{\text{low}}(b); b \in {}^\circ B\}$, a \mathcal{A} -basis of ${}^\circ\mathbf{K}_I$.

3.4. Proposition. (a) We have $\langle {}^\circ G^{\text{low}}(b) : {}^\circ G^{\text{up}}(b') \rangle = \delta_{b,b'}$ for each b, b' in ${}^\circ B$.

(b) We have $\langle P^\sharp : M \rangle = \overline{\langle P : M^\flat \rangle}$ for each P, M .

(c) We have ${}^\circ G^{\text{low}}(b)^\sharp = {}^\circ G^{\text{low}}(b)$ for each b in ${}^\circ B$.

The proof is the same as in [VV, prop. 8.4].

3.5. Example. Set $\nu = i + \theta(i)$ and $\mathbf{i} = i\theta(i)$. Consider the graded ${}^\circ\mathbf{R}_\nu$ -modules

$${}^\circ\mathbf{R}_\mathbf{i} = {}^\circ\mathbf{R}1_\mathbf{i} = 1_\mathbf{i}{}^\circ\mathbf{R}, \quad {}^\circ\mathbf{L}_\mathbf{i} = \text{top}({}^\circ\mathbf{R}_\mathbf{i}).$$

The global bases are given by

$${}^\circ\mathbf{G}_\nu^{\text{low}} = \{{}^\circ\mathbf{R}_\mathbf{i}, {}^\circ\mathbf{R}_{\theta(\mathbf{i})}\}, \quad {}^\circ\mathbf{G}_\nu^{\text{up}} = \{{}^\circ\mathbf{L}_\mathbf{i}, {}^\circ\mathbf{L}_{\theta(\mathbf{i})}\}.$$

For $m = 0$ we have ${}^\circ\mathbf{R}_0 = \mathbf{k} \oplus \mathbf{k}$. Set $\phi_+ = \mathbf{k} \oplus 0$ and $\phi_- = 0 \oplus \mathbf{k}$. We have

$${}^\circ\mathbf{G}_0^{\text{low}} = {}^\circ\mathbf{G}_0^{\text{up}} = \{\phi_+, \phi_-\}.$$

3.6. Definition of the operators e_i, f_i, e'_i, f'_i . In this section we'll always assume $m > 0$ unless specified otherwise. First, let us introduce the following notation. Let $D_{m,1}$ be the set of minimal representative in ${}^\circ W_{m+1}$ of the cosets in ${}^\circ W_m \backslash {}^\circ W_{m+1}$. Write

$$D_{m,1;m,1} = D_{m,1} \cap (D_{m,1})^{-1}.$$

For each element w of $D_{m,1;m,1}$ we set

$$W(w) = {}^\circ W_m \cap w({}^\circ W_m)w^{-1}.$$

Let \mathbf{R}_1 be the \mathbf{k} -algebra generated by elements $1_i, \varkappa_i, i \in I$, satisfying the defining relations $1_i 1_{i'} = \delta_{i,i'} 1_i$ and $\varkappa_i = 1_i \varkappa_i 1_i$. We equip \mathbf{R}_1 with the grading such that $\deg(1_i) = 0$ and $\deg(\varkappa_i) = 2$. Let

$$\mathbf{R}_i = 1_i \mathbf{R}_1 = \mathbf{R}_1 1_i, \quad \mathbf{L}_i = \text{top}(\mathbf{R}_i) = \mathbf{R}_i / (\varkappa_i).$$

Then \mathbf{R}_i is a graded projective \mathbf{R}_1 -module and \mathbf{L}_i is simple. We abbreviate

$${}^\theta \mathbf{R}_{m,1} = {}^\theta \mathbf{R}_m \otimes \mathbf{R}_1, \quad {}^\circ \mathbf{R}_{m,1} = {}^\circ \mathbf{R}_m \otimes \mathbf{R}_1.$$

There is an unique inclusion of graded \mathbf{k} -algebras

$$(3.2) \quad \begin{aligned} {}^\theta \mathbf{R}_{m,1} &\rightarrow {}^\theta \mathbf{R}_{m+1}, \\ 1_{\mathbf{i}} \otimes 1_i &\mapsto 1_{\mathbf{i}'}, \\ 1_{\mathbf{i}} \otimes \varkappa_{i,l} &\mapsto \varkappa_{\mathbf{i}',m+l}, \\ \varkappa_{\mathbf{i},l} \otimes 1_i &\mapsto \varkappa_{\mathbf{i}',l}, \\ \pi_{\mathbf{i},1} \otimes 1_i &\mapsto \pi_{\mathbf{i}',1}, \\ \sigma_{\mathbf{i},k} \otimes 1_i &\mapsto \sigma_{\mathbf{i}',k}, \end{aligned}$$

where, given $\mathbf{i} \in {}^\theta I^m$ and $i \in I$, we have set $\mathbf{i}' = \theta(i)\mathbf{i}i$, a sequence in ${}^\theta I^{m+1}$. This inclusion restricts to an inclusion ${}^\circ \mathbf{R}_{m,1} \subset {}^\circ \mathbf{R}_{m+1}$.

3.7. Lemma. *The graded ${}^\circ \mathbf{R}_{m,1}$ -module ${}^\circ \mathbf{R}_{m+1}$ is free of rank $2(m+1)$.*

Proof : For each w in $D_{m,1}$ we have the element $\sigma_{\dot{w}}$ in ${}^\circ \mathbf{R}_{m+1}$ defined in (1.5). Using filtered/graded arguments it is easy to see that

$${}^\circ \mathbf{R}_{m+1} = \bigoplus_{w \in D_{m,1}} {}^\circ \mathbf{R}_{m,1} \sigma_{\dot{w}}.$$

□

We define a triple of adjoint functors $(\psi_!, \psi^*, \psi_*)$ where

$$\psi^* : {}^\circ \mathbf{R}_{m+1}\text{-mod} \rightarrow {}^\circ \mathbf{R}_m\text{-mod} \times \mathbf{R}_1\text{-mod}$$

is the restriction and $\psi_!, \psi_*$ are given by

$$\begin{aligned} \psi_! : & \begin{cases} {}^\circ \mathbf{R}_m\text{-mod} \times \mathbf{R}_1\text{-mod} \rightarrow {}^\circ \mathbf{R}_{m+1}\text{-mod}, \\ (M, M') \mapsto {}^\circ \mathbf{R}_{m+1} \otimes_{{}^\circ \mathbf{R}_{m,1}} (M \otimes M'), \end{cases} \\ \psi_* : & \begin{cases} {}^\circ \mathbf{R}_m\text{-mod} \times \mathbf{R}_1\text{-mod} \rightarrow {}^\circ \mathbf{R}_{m+1}\text{-mod}, \\ (M, M') \mapsto \text{hom}_{{}^\circ \mathbf{R}_{m,1}}({}^\circ \mathbf{R}_{m+1}, M \otimes M'). \end{cases} \end{aligned}$$

The functor ψ^* is exact, it commutes with the shift of the grading, and it takes finite dimensional modules to finite dimensional modules. Thus $\psi_!$ takes projectives to

projectives and it commutes with the shift of the grading. Since the graded ${}^\circ\mathbf{R}_{m,1}$ -module ${}^\circ\mathbf{R}_{m+1}$ is free of finite rank, the functor $\psi_!$ is exact and it takes finite dimensional modules to finite dimensional ones. For the same reason ψ_* is exact and ψ^* takes projective modules to projective ones. To summarize, the functors $\psi_!$, ψ^* , ψ_* are exact, they commute with the shift of the grading, they take projective modules to projective ones and finite dimensional modules to finite dimensional ones. For any graded ${}^\circ\mathbf{R}_m$ -module M and any graded ${}^\circ\mathbf{R}_{m+1}$ -module N we write

$$(3.3) \quad f_i(M) = {}^\circ\mathbf{R}_{m+1}1_{m,i} \otimes_{{}^\circ\mathbf{R}_m} M, \quad e'_i(N) = 1_{m,i}N.$$

Here and below, for any graded ${}^\circ\mathbf{R}_{m+1}$ -module N we set

$$1_{m,i}N = \bigoplus_{\mathbf{i}} 1_{\theta(\mathbf{i})\mathbf{i}i}N, \quad \mathbf{i} \in {}^\theta I^m.$$

Note that $f_i(M)$ is a graded ${}^\circ\mathbf{R}_{m+1}$ -module, while $e'_i(N)$ is a graded ${}^\circ\mathbf{R}_m$ -module.

3.8. Definition. Let e'_i, f_i be the \mathcal{A} -linear operators on ${}^\circ\mathbf{K}_I$ such that e'_i, f_i on ${}^\circ\mathbf{K}_{I,*}$ by the formula (3.3) and

$$f_i(\phi_+) = {}^\circ\mathbf{R}_{\theta(\mathbf{i})i}, \quad f_i(\phi_-) = {}^\circ\mathbf{R}_{i\theta(\mathbf{i})}, \quad e'_i(\mathbf{R}_{\theta(\mathbf{j})j}) = \delta_{i,j}\phi_+ + \delta_{i,\theta(\mathbf{j})}\phi_-.$$

Let e_i, f'_i be the \mathcal{A} -linear operators on ${}^\circ\mathbf{G}_I$ which are the transpose of f_i, e'_i with respect to the Cartan pairing. We will also write $e_i = (1-v^2)e'_i$ and $f_i = (1-v^2)f'_i$.

The proof of the following lemma is the same as in [VV, lem. 8.9].

3.9. Lemma. (a) For each $M \in {}^\circ\mathbf{R}_m\text{-mod}^f$, $N \in {}^\circ\mathbf{R}_{m+1}\text{-mod}^f$ we have

$$e_i(N) = 1_{m,i}N, \quad f'_i(M) = \text{hom}_{{}^\circ\mathbf{R}_{m,1}}({}^\circ\mathbf{R}_{m+1}, M \otimes \mathbf{R}_i).$$

(b) For each $M \in {}^\circ\mathbf{R}_m\text{-mod}$, $N \in {}^\circ\mathbf{R}_{m+1}\text{-mod}$ we have

$$(e'_i(N) : M) = (N : f_i(M)).$$

(c) We have $f_i(P)^\sharp = f_i(P^\sharp)$ for each $P \in {}^\circ\mathbf{R}\text{-proj}$.

(d) We have $e_i(M)^\flat = e_i(M^\flat)$ for each $M \in {}^\circ\mathbf{R}\text{-mod}^f$.

3.10. Induction of \mathbf{H}_m -modules versus induction of ${}^\circ\mathbf{R}_m$ -modules. Recall the functors E_i, F_i on $\mathbf{H}\text{-Mod}_I^f$ defined in (2.2). We have also the functors

$$\mathbf{for} : {}^\circ\mathbf{R}_m\text{-mod}^f \rightarrow {}^\circ\mathbf{R}_m\text{-Mod}_0^f, \quad \Psi : {}^\circ\mathbf{R}_m\text{-Mod}_0^f \rightarrow \mathbf{H}_m\text{-Mod}_I^f,$$

where \mathbf{for} is the forgetting of the grading. Finally we define functors

$$(3.4) \quad \begin{aligned} E_i : {}^\circ\mathbf{R}_m\text{-Mod}_0^f &\rightarrow {}^\circ\mathbf{R}_{m-1}\text{-Mod}_0^f, & E_i M &= 1_{m-1,i}M, \\ F_i : {}^\circ\mathbf{R}_m\text{-Mod}_0^f &\rightarrow {}^\circ\mathbf{R}_{m+1}\text{-Mod}_0^f, & F_i M &= \psi_!(M, \mathbf{L}_i). \end{aligned}$$

3.11. Proposition. *There are canonical isomorphisms of functors*

$$E_i \circ \Psi = \Psi \circ E_i, \quad F_i \circ \Psi = \Psi \circ F_i, \quad E_i \circ \mathbf{for} = \mathbf{for} \circ e_i, \quad F_i \circ \mathbf{for} = \mathbf{for} \circ f_{\theta(i)}.$$

Proof: The proof is the same as in [VV, prop. 8.17]. \square

3.12. Proposition. (a) *The functor Ψ yields an isomorphism of Abelian groups*

$$\bigoplus_{m \geq 0} [{}^\circ \mathbf{R}_m\text{-}\mathbf{Mod}_0^f] = \bigoplus_{m \geq 0} [\mathbf{H}_m\text{-}\mathbf{Mod}_I^f].$$

The functors E_i, F_i yield endomorphisms of both sides which are intertwined by Ψ .

(b) *The functor \mathbf{for} factors to a group isomorphism*

$${}^\circ \mathbf{G}_I / (v - 1) = \bigoplus_{m \geq 0} [{}^\circ \mathbf{R}_m\text{-}\mathbf{Mod}_0^f].$$

Proof: Claim (a) follows from Corollary 2.7 and Proposition 3.11. Claim (b) follows from Proposition 3.2. \square

3.13. Type D versus type B. We can compare the previous constructions with their analogues in type B. Let ${}^\theta \mathbf{K}, {}^\theta B, {}^\theta G^{\text{low}}$, etc, denote the type B analogues of ${}^\circ \mathbf{K}, {}^\circ B, {}^\circ G^{\text{low}}$, etc. See [VV] for details. We'll use the same notation for the functors $\psi^*, \psi!, \psi_*, e'_i, f_i$, etc, on the type B side and on the type D side. Fix $m > 0$ and $\nu \in {}^\theta \mathbf{NI}$ such that $|\nu| = 2m$. The inclusion of graded \mathbf{k} -algebras ${}^\circ \mathbf{R}_\nu \subset {}^\theta \mathbf{R}_\nu$ in (1.2) yields a restriction functor

$$\text{res} : {}^\theta \mathbf{R}_\nu\text{-mod} \rightarrow {}^\circ \mathbf{R}_\nu\text{-mod}$$

and an induction functor

$$\text{ind} : {}^\circ \mathbf{R}_\nu\text{-mod} \rightarrow {}^\theta \mathbf{R}_\nu\text{-mod}, \quad M \mapsto {}^\theta \mathbf{R}_\nu \otimes_{{}^\circ \mathbf{R}_\nu} M.$$

Both functors are exact, they map finite dimensional modules to finite dimensional ones, and map projectives to projectives. Thus, they yield morphisms of \mathcal{A} -modules

$$\text{res} : {}^\theta \mathbf{K}_{I,m} \rightarrow {}^\circ \mathbf{K}_{I,m}, \quad \text{res} : {}^\theta \mathbf{G}_{I,m} \rightarrow {}^\circ \mathbf{G}_{I,m},$$

$$\text{ind} : {}^\circ \mathbf{K}_{I,m} \rightarrow {}^\theta \mathbf{K}_{I,m}, \quad \text{ind} : {}^\circ \mathbf{G}_{I,m} \rightarrow {}^\theta \mathbf{G}_{I,m}.$$

Moreover, for any $P \in {}^\theta \mathbf{K}_{I,m}$ and any $L \in {}^\theta \mathbf{G}_{I,m}$ we have

$$(3.5) \quad \begin{aligned} \text{res}(P^\sharp) &= (\text{res} P)^\sharp, & \text{ind}(P^\sharp) &= (\text{ind} P)^\sharp \\ \text{res}(L^\flat) &= (\text{res} L)^\flat, & \text{ind}(L^\flat) &= (\text{ind} L)^\flat. \end{aligned}$$

Note also that ind and res are left and right adjoint functors, because

$${}^\theta \mathbf{R}_\nu \otimes_{{}^\circ \mathbf{R}_\nu} M = \text{hom}_{{}^\circ \mathbf{R}_\nu}({}^\theta \mathbf{R}_\nu, M)$$

as graded ${}^\theta \mathbf{R}_\nu$ -modules.

3.14. Definition. For any graded ${}^\circ\mathbf{R}_\nu$ -module M we define the graded ${}^\circ\mathbf{R}_\nu$ -module M^γ with the same underlying graded \mathbf{k} -vector space as M such that the action of ${}^\circ\mathbf{R}_\nu$ is twisted by γ , i.e., the graded \mathbf{k} -algebra ${}^\circ\mathbf{R}_\nu$ acts on M^γ by $a \cdot m = \gamma(a)m$ for each $a \in {}^\circ\mathbf{R}_\nu$, $m \in M$. Note that $(M^\gamma)^\gamma = M$, and that M^γ is simple (resp. projective, indecomposable) if M has the same property.

For any graded ${}^\circ\mathbf{R}_m$ -module M and any graded ${}^\circ\mathbf{R}_{m+1}$ -module N we have canonical isomorphisms of ${}^\circ\mathbf{R}$ -modules

$$(3.6) \quad (f_i(M))^\gamma = f_i(M^\gamma), \quad (e'_i(N))^\gamma = e'_i(N^\gamma).$$

The first isomorphism is given by

$${}^\circ\mathbf{R}_{m+1}1_{m,i} \otimes_{{}^\circ\mathbf{R}_m} M \rightarrow {}^\circ\mathbf{R}_{m+1}1_{m,i} \otimes_{{}^\circ\mathbf{R}_m} M, \quad a \otimes m \mapsto \gamma(a) \otimes m.$$

The second one is the identity map on the vector space $1_{m,i}N$. Recall that ${}^\theta I^\nu$ is the disjoint union of ${}^\theta I_+^\nu$ and ${}^\theta I_-^\nu$. Set

$$1_{\nu,+} = \sum_{\mathbf{i} \in {}^\theta I_+^\nu} 1_{\mathbf{i}}, \quad 1_{\nu,-} = \sum_{\mathbf{i} \in {}^\theta I_-^\nu} 1_{\mathbf{i}}.$$

The following holds.

3.15. Lemma. Let M be a graded ${}^\circ\mathbf{R}_\nu$ -module.

- (a) Both $1_{\nu,+}$ and $1_{\nu,-}$ are central idempotents in ${}^\circ\mathbf{R}_\nu$. We have $1_{\nu,+} = \gamma(1_{\nu,-})$.
- (b) There is a decomposition of graded ${}^\circ\mathbf{R}_\nu$ -modules $M = 1_{\nu,+}M \oplus 1_{\nu,-}M$.
- (c) We have a canonical isomorphism of ${}^\circ\mathbf{R}_\nu$ -modules $\text{res} \circ \text{ind}(M) = M \oplus M^\gamma$.
- (d) Suppose that there exists $a \in \{+, -\}$ such that $0 = 1_{\nu,-a}M$, then there are canonical isomorphisms of graded ${}^\circ\mathbf{R}$ -modules

$$M = 1_{\nu,a}M, \quad 0 = 1_{\nu,a}M^\gamma, \quad M^\gamma = 1_{\nu,-a}M^\gamma.$$

Proof: Part (a) is given by Proposition 1.6 and the fact that ε_1 maps ${}^\theta I_+^\nu$ to ${}^\theta I_-^\nu$. Part (b) follows from (a), (c) is given by definition, and (d) follows from (a), (b). \square

Now, let m and ν be as before. Given $i \in I$, we set $\nu' = \nu + i + \theta(i)$. There is an obvious inclusion $W_m \subset W_{m+1}$. Thus the group W_m acts on ${}^\theta I^{\nu'}$, and the map

$${}^\theta I^\nu \rightarrow {}^\theta I^{\nu'}, \quad \mathbf{i} \mapsto \theta(i)\mathbf{i}$$

is W_m -equivariant. This implies that there exists $a_i \in \{+, -\}$ such that the image of ${}^\theta I_+^\nu$ is contained in ${}^\theta I_{a_i}^{\nu'}$, and the image of ${}^\theta I_-^\nu$ is contained in ${}^\theta I_{-a_i}^{\nu'}$.

3.16. Lemma. Let M be a graded ${}^\circ\mathbf{R}_\nu$ -module such that $1_{\nu,-a}M = 0$ with $a \in \{+, -\}$. Then we have

$$1_{\nu',-a_i a} f_i(M) = 0, \quad 1_{\nu',a_i a} f_{\theta(i)}(M) = 0.$$

Proof: We have

$$\begin{aligned} 1_{\nu', -a_i a} f_i(M) &= 1_{\nu', -a_i a} \circ \mathbf{R}_{\nu'} 1_{\nu, i} \otimes_{\circ \mathbf{R}_{\nu}} M \\ &= \circ \mathbf{R}_{\nu'} 1_{\nu', -a_i a} 1_{\nu, i} 1_{\nu, a} \otimes_{\circ \mathbf{R}_{\nu}} M. \end{aligned}$$

Here we have identified $1_{\nu, a}$ with the image of $(1_{\nu, a}, 1_i)$ via the inclusion (3.2). The definition of this inclusion and that of a_i yield that

$$1_{\nu', a_i a} 1_{\nu, i} 1_{\nu, a} = 1_{\nu, a}, \quad 1_{\nu', -a_i a} 1_{\nu, i} 1_{\nu, a} = 0.$$

The first equality follows. Next, note that for any $\mathbf{i} \in {}^\theta I^\nu$, the sequences $\theta(i)\mathbf{i}\mathbf{i}$ and $\mathbf{i}\mathbf{i}\theta(i) = \varepsilon_{m+1}(\theta(i)\mathbf{i}\mathbf{i})$ always belong to different ${}^\circ W_{m+1}$ -orbits. Thus, we have $a_{\theta(i)} = -a_i$. So the second equality follows from the first. \square

Consider the following diagram

$$\begin{array}{ccc} {}^\circ \mathbf{R}_{\nu'}\text{-mod} \times \mathbf{R}_i\text{-mod} & \begin{array}{c} \xrightarrow{\psi_!} \\ \xleftarrow{\psi^*} \end{array} & {}^\circ \mathbf{R}_{\nu'}\text{-mod} \\ \begin{array}{c} \text{res} \times \text{id} \updownarrow \text{ind} \times \text{id} \end{array} & & \begin{array}{c} \text{res} \updownarrow \text{ind} \end{array} \\ {}^\theta \mathbf{R}_{\nu'}\text{-mod} \times \mathbf{R}_i\text{-mod} & \begin{array}{c} \xrightarrow{\psi_!} \\ \xleftarrow{\psi^*} \end{array} & {}^\theta \mathbf{R}_{\nu'}\text{-mod}. \end{array}$$

3.17. Lemma. *There are canonical isomorphisms of functors*

$$\text{ind} \circ \psi_! = \psi_! \circ (\text{ind} \times \text{id}), \quad \psi^* \circ \text{ind} = (\text{ind} \times \text{id}) \circ \psi^*, \quad \text{ind} \circ \psi_* = \psi_* \circ (\text{ind} \times \text{id}),$$

$$\text{res} \circ \psi_! = \psi_! \circ (\text{res} \times \text{id}), \quad \psi^* \circ \text{res} = (\text{res} \times \text{id}) \circ \psi^*, \quad \text{res} \circ \psi_* = \psi_* \circ (\text{res} \times \text{id}).$$

Proof: The functor ind is left and right adjoint to res . Therefore it is enough to prove the first two isomorphisms in the first line. The isomorphism

$$\text{ind} \circ \psi_! = \psi_! \circ (\text{ind} \times \text{id})$$

comes from the associativity of the induction. Let us prove that

$$\psi^* \circ \text{ind} = (\text{ind} \times \text{id}) \circ \psi^*.$$

For any M in ${}^\circ \mathbf{R}_{\nu'}\text{-mod}$, the obvious inclusion ${}^\theta \mathbf{R}_{\nu} \otimes \mathbf{R}_i \subset {}^\theta \mathbf{R}_{\nu'}$ yields a map

$$(\text{ind} \times \text{id}) \psi^*(M) = ({}^\theta \mathbf{R}_{\nu} \otimes \mathbf{R}_i) \otimes_{\circ \mathbf{R}_{\nu} \otimes \mathbf{R}_i} \psi^*(M) \rightarrow \psi^*({}^\theta \mathbf{R}_{\nu'} \otimes_{\circ \mathbf{R}_{\nu} \otimes \mathbf{R}_i} M).$$

Combining it with the obvious map

$${}^\theta \mathbf{R}_{\nu'} \otimes_{\circ \mathbf{R}_{\nu} \otimes \mathbf{R}_i} M \rightarrow {}^\theta \mathbf{R}_{\nu'} \otimes_{\circ \mathbf{R}_{\nu'}} M$$

we get a morphism of ${}^\theta \mathbf{R}_{\nu} \otimes \mathbf{R}_i$ -modules

$$(\text{ind} \times \text{id}) \psi^*(M) \rightarrow \psi^* \text{ind}(M).$$

We need to show that it is bijective. This is obvious because at the level of vector spaces, the map above is given by

$$M \oplus (\pi_{1, \nu} \otimes M) \rightarrow M \oplus (\pi_{1, \nu'} \otimes M), \quad m + \pi_{1, \nu} \otimes n \mapsto m + \pi_{1, \nu'} \otimes n.$$

Here $\pi_{1, \nu}$ and $\pi_{1, \nu'}$ denote the element π_1 in ${}^\theta \mathbf{R}_{\nu}$ and ${}^\theta \mathbf{R}_{\nu'}$ respectively. \square

3.18. Corollary. (a) The operators e'_i, f_i on ${}^\circ\mathbf{K}_{I,*}$ and on ${}^\theta\mathbf{K}_{I,*}$ are intertwined by the maps ind, res , i.e., we have

$$e'_i \circ \text{ind} = \text{ind} \circ e'_i, \quad f_i \circ \text{ind} = \text{ind} \circ f_i, \quad e'_i \circ \text{res} = \text{res} \circ e'_i, \quad f_i \circ \text{res} = \text{res} \circ f_i.$$

(b) The same result holds for the operators e_i, f'_i on ${}^\circ\mathbf{G}_{I,*}$ and on ${}^\theta\mathbf{G}_{I,*}$.

3.19. Now, we concentrate on non graded irreducible modules. First, let

$$\text{Res} : {}^\theta\mathbf{R}_\nu\text{-Mod} \rightarrow {}^\circ\mathbf{R}_\nu\text{-Mod}, \quad \text{Ind} : {}^\circ\mathbf{R}_\nu\text{-Mod} \rightarrow {}^\theta\mathbf{R}_\nu\text{-Mod}$$

be the (non graded) restriction and induction functors. We have

$$\text{for} \circ \text{res} = \text{Res} \circ \text{for}, \quad \text{for} \circ \text{ind} = \text{Ind} \circ \text{for}.$$

3.20. Lemma. Let L, L' be irreducible ${}^\circ\mathbf{R}_\nu$ -modules.

(a) The ${}^\circ\mathbf{R}_\nu$ -modules L and L^γ are not isomorphic.

(b) $\text{Ind}(L)$ is an irreducible ${}^\theta\mathbf{R}_\nu$ -module.

(c) Every irreducible ${}^\theta\mathbf{R}_\nu$ -module is obtained in this way.

(d) $\text{Ind}(L) \simeq \text{Ind}(L')$ iff $L' \simeq L$ or L^γ .

Proof: For any ${}^\theta\mathbf{R}_\nu$ -module $M \neq 0$, both $1_{\nu,+}M$ and $1_{\nu,-}M$ are nonzero. Indeed, we have $M = 1_{\nu,+}M \oplus 1_{\nu,-}M$, and we may suppose that $1_{\nu,+}M \neq 0$. The automorphism $M \rightarrow M, m \mapsto \pi_1 m$ takes $1_{\nu,+}M$ to $1_{\nu,-}M$ by Lemma 3.15(a). Hence $1_{\nu,-}M \neq 0$.

Now, to prove part (a), suppose that $\phi : L \rightarrow L^\gamma$ is an isomorphism of ${}^\circ\mathbf{R}_\nu$ -modules. We can regard ϕ as a γ -antilinear map $L \rightarrow L$. Since L is irreducible, by Schur's lemma we may assume that $\phi^2 = \text{id}_L$. Then L admits a ${}^\theta\mathbf{R}_\nu$ -module structure such that the ${}^\circ\mathbf{R}_\nu$ -action is as before and π_1 acts as ϕ . Thus, by the discussion above, neither $1_{\nu,+}L$ nor $1_{\nu,-}L$ is zero. This contradicts the fact that L is an irreducible ${}^\circ\mathbf{R}_\nu$ -module.

Parts (b), (c), (d) follows from (a) by Clifford theory, see e.g., [RR, appendix]. \square

We can now prove Proposition 3.2.

3.21. Proof of Proposition 3.2. Let $b \in {}^\circ B$. We may suppose that $b = 1_{\nu,+}b$. By Lemma 3.20(b) the module ${}^\theta b = \text{Ind}(b)$ lies in ${}^\theta B$. By [VV, prop. 8.2] there exists a unique selfdual irreducible graded ${}^\theta\mathbf{R}$ -module ${}^\theta G^{\text{up}}({}^\theta b)$ which is isomorphic to ${}^\theta b$ as a non graded module. Set

$${}^\circ G^{\text{up}}(b) = 1_{\nu,+}\text{res}({}^\theta G^{\text{up}}({}^\theta b)).$$

By Lemma 3.15(d) we have ${}^\circ G^{\text{up}}(b) = b$ as a non graded ${}^\circ\mathbf{R}$ -module, and by (3.5) it is selfdual. This proves existence part of the proposition. To prove the uniqueness, suppose that M is another module with the same properties. Then $\text{ind}(M)$ is a selfdual graded ${}^\theta\mathbf{R}$ -modules which is isomorphic to ${}^\theta b$ as a non graded ${}^\theta\mathbf{R}$ -module. Thus we have $\text{ind}(M) = {}^\theta G^{\text{up}}({}^\theta b)$ by loc. cit. By Lemma 3.15(d) we have also

$$M = 1_{\nu,+}\text{res}({}^\theta G^{\text{up}}({}^\theta b)).$$

So M is isomorphic to ${}^\circ G^{\text{up}}(b)$. \square

3.22. The crystal operators on ${}^\circ\mathbf{G}_I$ and ${}^\circ B$. Fix a vertex i in I . For each irreducible graded ${}^\circ\mathbf{R}_m$ -module M we define

$$\tilde{e}_i(M) = \text{soc}(e_i(M)), \quad \tilde{f}_i(M) = \text{top } \psi_1(M, \mathbf{L}_i), \quad \varepsilon_i(M) = \max\{n \geq 0; e_i^n(M) \neq 0\}.$$

3.23. Lemma. *Let M be an irreducible graded ${}^\circ\mathbf{R}$ -module such that $e_i(M)$, $f_i(M)$ belong to ${}^\circ\mathbf{G}_{I,*}$. We have*

$$\text{ind}(\tilde{e}_i(M)) = \tilde{e}_i(\text{ind}(M)), \quad \text{ind}(\tilde{f}_i(M)) = \tilde{f}_i(\text{ind}(M)), \quad \varepsilon_i(M) = \varepsilon_i(\text{ind}(M)).$$

In particular, both $\tilde{e}_i(M)$ and $\tilde{f}_i(M)$ are irreducible.

Proof: By Corollary 3.18 we have $\text{ind}(e_i(M)) = e_i(\text{ind}(M))$. Thus, since ind is an exact functor we have $\text{ind}(\tilde{e}_i(M)) \subset e_i(\text{ind}(M))$. Since ind is an additive functor, by Lemma 3.20(b) we have indeed

$$\text{ind}(\tilde{e}_i(M)) \subset \tilde{e}_i(\text{ind}(M)).$$

Note that $\text{ind}(M)$ is irreducible by Lemma 3.20(b), thus $\tilde{e}_i(\text{ind}(M))$ is irreducible by [VV, prop. 8.21]. Since $\text{ind}(\tilde{e}_i(M))$ is nonzero, the inclusion is an isomorphism. The fact that $\text{ind}(\tilde{e}_i(M))$ is irreducible implies in particular that $\tilde{e}_i(M)$ is simple. The proof of the second isomorphism is similar. The third equality is obvious. \square

For each irreducible module b in ${}^\circ B$ we define

$$(3.6) \quad \tilde{E}_i(b) = \text{soc}(E_i(b)), \quad \tilde{F}_i(b) = \text{top}(F_i(b)), \quad \varepsilon_i(b) = \max\{n \geq 0; E_i^n(b) \neq 0\}.$$

Hence we have **for** $\circ \tilde{e}_i = \tilde{E}_i \circ \mathbf{for}$, **for** $\circ \tilde{f}_i = \tilde{F}_i \circ \mathbf{for}$, and $\varepsilon_i = \varepsilon_i \circ \mathbf{for}$.

3.24. Proposition. *For each b, b' in ${}^\circ B$ we have*

- (a) $\tilde{F}_i(b) \in {}^\circ B$,
- (b) $\tilde{E}_i(b) \in {}^\circ B \cup \{0\}$,
- (c) $\tilde{F}_i(b) = b' \iff \tilde{E}_i(b') = b$,
- (d) $\varepsilon_i(b) = \max\{n \geq 0; \tilde{E}_i^n(b) \neq 0\}$,
- (e) $\varepsilon_i(\tilde{F}_i(b)) = \varepsilon_i(b) + 1$,
- (f) if $\tilde{E}_i(b) = 0$ for all i then $b = \phi_\pm$.

Proof: Part (c) follows from adjunction. The other parts follow from [VV, prop. 3.14] and Lemma 3.23. \square

3.25. Remark. For any $b \in {}^\circ B$ and any $i \neq j$ we have $\tilde{F}_i(b) \neq \tilde{F}_j(b)$. This is obvious if $j \neq \theta(i)$. Because in this case $\tilde{F}_i(b)$ and $\tilde{F}_j(b)$ are ${}^\circ\mathbf{R}_\nu$ -modules for different ν . Now, consider the case $j = \theta(i)$. We may suppose that $\tilde{F}_i(b) = 1_{\nu,+} \tilde{F}_i(b)$ for certain ν . Then by Lemma 3.16 we have $1_{\nu,+} \tilde{F}_{\theta(i)}(b) = 0$. In particular $\tilde{F}_i(b)$ is not isomorphic to $\tilde{F}_{\theta(i)}(b)$.

3.26. The algebra ${}^\theta\mathcal{B}$ and the ${}^\theta\mathcal{B}$ -module ${}^\circ\mathbf{V}$. Following [EK1,2,3] we define a \mathcal{K} -algebra ${}^\theta\mathcal{B}$ as follows.

3.27. Definition. Let ${}^\theta\mathcal{B}$ be the \mathcal{K} -algebra generated by e_i, f_i and invertible elements $t_i, i \in I$, satisfying the following defining relations

- (a) $t_i t_j = t_j t_i$ and $t_{\theta(i)} = t_i$ for all i, j ,
- (b) $t_i e_j t_i^{-1} = v^{i \cdot j + \theta(i) \cdot j} e_j$ and $t_i f_j t_i^{-1} = v^{-i \cdot j - \theta(i) \cdot j} f_j$ for all i, j ,
- (c) $e_i f_j = v^{-i \cdot j} f_j e_i + \delta_{i,j} + \delta_{\theta(i),j} t_i$ for all i, j ,
- (d) $\sum_{a+b=1-i \cdot j} (-1)^a e_i^{(a)} e_j e_i^{(b)} = \sum_{a+b=1-i \cdot j} (-1)^a f_i^{(a)} f_j f_i^{(b)} = 0$ if $i \neq j$.

Here and below we use the following notation

$$\theta^{(a)} = \theta^a / \langle a \rangle!, \quad \langle a \rangle = \sum_{l=1}^a v^{a+1-2l}, \quad \langle a \rangle! = \prod_{l=1}^m \langle l \rangle.$$

We can now construct a representation of ${}^\theta\mathcal{B}$ as follows. The \mathcal{K} -vector space

$${}^\circ\mathbf{V} = \mathcal{K} \otimes_{\mathcal{A}} {}^\circ\mathbf{K}_I$$

is equipped with \mathcal{K} -linear operators e_i, e'_i, f_i and with a \mathcal{K} -bilinear form in the obvious way.

3.28. Theorem. (a) The operators e_i, f_i define a representation of ${}^\theta\mathcal{B}$ on ${}^\circ\mathbf{V}$. The ${}^\theta\mathcal{B}$ -module ${}^\circ\mathbf{V}$ is generated by linearly independent vectors ϕ_+ and ϕ_- such that for each $i \in I$ we have

$$e_i \phi_{\pm} = 0, \quad t_i \phi_{\pm} = \phi_{\mp}, \quad \{x \in {}^\circ\mathbf{V}; e_j x = 0, \forall j\} = \mathbf{k} \phi_+ \oplus \mathbf{k} \phi_-.$$

(b) The symmetric bilinear form on ${}^\circ\mathbf{V}$ is non-degenerate. We have $(\phi_a : \phi_{a'}) = \delta_{a,a'}$ for $a, a' \in \{+, -\}$ and $(e'_i x : y) = (x : f_i y)$ for all i and all $x, y \in {}^\circ\mathbf{V}$.

Proof: For each i in I we define the \mathcal{A} -linear operator t_i on ${}^\circ\mathbf{K}_I$ by setting

$$t_i \phi_{\pm} = \phi_{\mp} \quad \text{and} \quad t_i P = v^{-\nu \cdot (i + \theta(i))} P^\gamma, \quad \forall P \in {}^\circ\mathbf{R}_{\nu}\text{-proj}.$$

We must prove that the operators e_i, f_i , and t_i satisfy the relations of ${}^\theta\mathcal{B}$. The relations (a), (b) are obvious. The relation (d) is standard. It remains to check (c). For this we need a version of the Mackey's induction-restriction theorem. Note that for $m > 1$ we have

$$D_{m,1;m,1} = \{e, s_m, \varepsilon_{m+1} \varepsilon_1\},$$

$$W(e) = {}^\circ W_m, \quad W(s_m) = {}^\circ W_{m-1}, \quad W(\varepsilon_{m+1} \varepsilon_1) = {}^\circ W_m.$$

Recall also that for $m = 1$ we have set ${}^\circ W_1 = \{e\}$.

3.29. Lemma. Fix i, j in I . Let μ, ν in ${}^\theta\mathbb{N}I$ be such that $\nu + i + \theta(i) = \mu + j + \theta(j)$. Put $|\nu| = |\mu| = 2m$. The graded $({}^\circ\mathbf{R}_{m,1}, {}^\circ\mathbf{R}_{m,1})$ -bimodule $1_{\nu,i} {}^\circ\mathbf{R}_{m+1} 1_{\mu,j}$ has a filtration by graded bimodules whose associated graded is isomorphic to

$$\delta_{i,j}({}^\circ\mathbf{R}_\nu \otimes \mathbf{R}_i) \oplus \delta_{\theta(i),j}(({}^\circ\mathbf{R}_\nu)^\gamma \otimes \mathbf{R}_{\theta(i)})[d'] \oplus A[d],$$

where A is equal to

$$\begin{aligned} & ({}^\circ\mathbf{R}_m 1_{\nu',i} \otimes \mathbf{R}_i) \otimes_{\mathbf{R}'} (1_{\nu',i} {}^\circ\mathbf{R}_m \otimes \mathbf{R}_i) && \text{if } m > 1, \\ & ({}^\circ\mathbf{R}_{\theta(j)} \otimes \mathbf{R}_i \otimes {}^\circ\mathbf{R}_1 \otimes \mathbf{R}_1 {}^\circ\mathbf{R}_{\theta(i)} \otimes \mathbf{R}_j) \oplus ({}^\circ\mathbf{R}_j \otimes \mathbf{R}_i \otimes {}^\circ\mathbf{R}_1 \otimes \mathbf{R}_1 {}^\circ\mathbf{R}_i \otimes \mathbf{R}_j) && \text{if } m = 1. \end{aligned}$$

Here we have set $\nu' = \nu - j - \theta(j)$, $\mathbf{R}' = {}^\circ\mathbf{R}_{m-1,1} \otimes \mathbf{R}_1$, $\mathbf{i} = i\theta(i)$, $\mathbf{j} = j\theta(j)$, $d = -i \cdot j$, and $d' = -\nu \cdot (i + \theta(i))/2$.

The proof is standard and is left to the reader. Now, recall that for $m > 1$ we have

$$f_j(P) = {}^\circ\mathbf{R}_{m+1} 1_{m,j} \otimes {}^\circ\mathbf{R}_{m,1} (P \otimes \mathbf{R}_1), \quad e'_i(P) = 1_{m-1,i} P,$$

where $1_{m-1,i} P$ is regarded as a ${}^\circ\mathbf{R}_{m-1}$ -module. Therefore we have

$$\begin{aligned} e'_i f_j(P) &= 1_{m,i} {}^\circ\mathbf{R}_{m+1} 1_{m,j} \otimes {}^\circ\mathbf{R}_{m,1} (P \otimes \mathbf{R}_1), \\ f_j e'_i(P) &= {}^\circ\mathbf{R}_m 1_{m-1,j} \otimes {}^\circ\mathbf{R}_{m-1,1} (1_{m-1,i} P \otimes \mathbf{R}_1). \end{aligned}$$

Therefore, up to some filtration we have the following identities

- (a) $e'_i f_i(P) = P \otimes \mathbf{R}_i + f_i e'_i(P)[-2]$,
- (b) $e'_i f_{\theta(i)}(P) = P^\gamma \otimes \mathbf{R}_{\theta(i)}[-\nu \cdot (i + \theta(i))/2] + f_{\theta(i)} e'_i(P)[-i \cdot \theta(i)]$,
- (c) $e'_i f_j(P) = f_j e'_i(P)[-i \cdot j]$ if $i \neq j, \theta(j)$.

These identities also hold for $m = 1$ and $P = {}^\circ\mathbf{R}_{\theta(i)i}$ for any $i \in I$. The first claim of part (a) follows because $\mathbf{R}_i = \mathbf{k} \oplus \mathbf{R}_i[2]$. The fact that ${}^\circ\mathbf{V}$ is generated by ϕ_\pm is a corollary of Proposition 3.31 below. Part (b) of the theorem follows from [KM, prop. 2.2(ii)] and Lemma 3.9(b). \square

3.30. Remarks. (a) The ${}^\theta\mathcal{B}$ -module ${}^\circ\mathbf{V}$ is the same as the ${}^\theta\mathcal{B}$ -module V_θ from [KM, prop. 2.2]. The involution $\sigma : {}^\circ\mathbf{V} \rightarrow {}^\circ\mathbf{V}$ in [KM, rem. 2.5(ii)] is given by $\sigma(P) = P^\gamma$. It yields an involution of ${}^\circ B$ in the obvious way. Note that Lemma 3.20(a) yields $\sigma(b) \neq b$ for any $b \in {}^\circ B$.

(b) Let ${}^\theta\mathbf{V}$ be the ${}^\theta\mathcal{B}$ -module $\mathcal{K} \otimes_{\mathcal{A}} {}^\theta\mathbf{K}_I$ and let ϕ be the class of the trivial ${}^\theta\mathbf{R}_0$ -module \mathbf{k} , see [VV, thm. 8.30]. We have an inclusion of ${}^\theta\mathcal{B}$ -modules

$${}^\theta\mathbf{V} \rightarrow {}^\circ\mathbf{V}, \quad \phi \mapsto \phi_+ \oplus \phi_-, \quad P \mapsto \text{res}(P).$$

3.31. Proposition. For any $b \in {}^\circ B$ the following holds

$$(a) \quad \begin{cases} f_i({}^\circ G^{\text{low}}(b)) = \langle \varepsilon_i(b) + 1 \rangle {}^\circ G^{\text{low}}(\tilde{F}_i b) + \sum_{b'} f_{b,b'} {}^\circ G^{\text{low}}(b'), \\ b' \in {}^\circ B, \quad \varepsilon_i(b') > \varepsilon_i(b) + 1, \quad f_{b,b'} \in v^{2-\varepsilon_i(b')} \mathbb{Z}[v], \end{cases}$$

$$(b) \quad \begin{cases} e_i({}^\circ G^{\text{low}}(b)) = v^{1-\varepsilon_i(b)} {}^\circ G^{\text{low}}(\tilde{E}_i b) + \sum_{b'} e_{b,b'} {}^\circ G^{\text{low}}(b'), \\ b' \in {}^\circ B, \quad \varepsilon_i(b') \geq \varepsilon_i(b), \quad e_{b,b'} \in v^{1-\varepsilon_i(b')} \mathbb{Z}[v]. \end{cases}$$

Proof: We prove part (a), the proof for (b) is similar. If ${}^\circ G^{\text{low}}(b) = \phi_\pm$ this is obvious. So we assume that ${}^\circ G^{\text{low}}(b)$ is a ${}^\circ \mathbf{R}_m$ -module for $m \geq 1$. Fix $\nu \in {}^\theta \mathbb{N}I$ such that $f_i({}^\circ G^{\text{low}}(b))$ is a ${}^\circ \mathbf{R}_\nu$ -module. We'll abbreviate $1_{\nu,a} = 1_a$ for $a \in \{+, -\}$. Since ${}^\circ G^{\text{low}}(b)$ is indecomposable, it fulfills the condition of Lemma 3.16. So there exists $a \in \{+, -\}$ such that $1_{-a} f_i({}^\circ G^{\text{low}}(b)) = 0$. Thus, by Lemma 3.15(c), (d) and Corollary 3.18 we have

$$f_i({}^\circ G^{\text{low}}(b)) = 1_a \text{res ind } f_i({}^\circ G^{\text{low}}(b)) = 1_a \text{res } f_i \text{ ind } ({}^\circ G^{\text{low}}(b)).$$

Note that ${}^\theta b = \text{Ind}(b)$ belongs to ${}^\theta B$ by Lemma 3.20(b). Hence (3.5) yields

$$\text{ind}({}^\circ G^{\text{low}}(b)) = {}^\theta G^{\text{low}}({}^\theta b).$$

We deduce that

$$f_i({}^\circ G^{\text{low}}(b)) = 1_a \text{res } f_i({}^\theta G^{\text{low}}({}^\theta b)).$$

Now, write

$$f_i({}^\theta G^{\text{low}}({}^\theta b)) = \sum f_{\theta b, \theta b'} {}^\theta G^{\text{low}}({}^\theta b'), \quad \theta b' \in {}^\theta B.$$

Then we have

$$f_i({}^\circ G^{\text{low}}(b)) = \sum f_{\theta b, \theta b'} 1_a \text{res}({}^\theta G^{\text{low}}({}^\theta b')).$$

For any $\theta b' \in {}^\theta B$ the ${}^\circ \mathbf{R}$ -module $1_a \text{Res}({}^\theta b')$ belongs to ${}^\circ B$. Thus we have

$$1_a \text{res}({}^\theta G^{\text{low}}({}^\theta b')) = {}^\circ G^{\text{low}}(1_a \text{Res}({}^\theta b')).$$

If $\theta b' \neq \theta b''$ then $1_a \text{Res}({}^\theta b') \neq 1_a \text{Res}({}^\theta b'')$, because $\theta b' = \text{Ind}(1_a \text{Res}({}^\theta b'))$. Thus

$$f_i({}^\circ G^{\text{low}}(b)) = \sum f_{\theta b, \theta b'} {}^\circ G^{\text{low}}(1_a \text{Res}({}^\theta b')),$$

and this is the expansion of the lhs in the lower global basis. Finally, we have

$$\varepsilon_i(1_a \text{Res}({}^\theta b')) = \varepsilon_i({}^\theta b')$$

by Lemma 3.23. So part (a) follows from [VV, prop. 10.11(b), 10.16]. \square

3.32. The global bases of ${}^\circ \mathbf{V}$. Since the operators e_i, f_i on ${}^\circ \mathbf{V}$ satisfy the relations $e_i f_i = v^{-2} f_i e_i + 1$, we can define the modified root operators $\tilde{\mathbf{e}}_i, \tilde{\mathbf{f}}_i$ on the ${}^\theta \mathcal{B}$ -module ${}^\circ \mathbf{V}$ as follows. For each u in ${}^\circ \mathbf{V}$ we write

$$u = \sum_{n \geq 0} f_i^{(n)} u_n \text{ with } e_i u_n = 0,$$

$$\tilde{\mathbf{e}}_i(u) = \sum_{n \geq 1} f_i^{(n-1)} u_n, \quad \tilde{\mathbf{f}}_i(u) = \sum_{n \geq 0} f_i^{(n+1)} u_n.$$

Let $\mathcal{R} \subset \mathcal{K}$ be the set of functions which are regular at $v = 0$. Let ${}^\circ \mathbf{L}$ be the \mathcal{R} -submodule of ${}^\circ \mathbf{V}$ spanned by the elements $\tilde{\mathbf{f}}_{i_1} \dots \tilde{\mathbf{f}}_{i_l}(\phi_\pm)$ with $l \geq 0, i_1, \dots, i_l \in I$. The following is the main result of the paper.

3.33. Theorem. (a) *We have*

$${}^\circ\mathbf{L} = \bigoplus_{b \in {}^\circ B} \mathcal{R}({}^\circ G^{\text{low}}(b)), \quad \tilde{e}_i({}^\circ\mathbf{L}) \subset {}^\circ\mathbf{L}, \quad \tilde{f}_i({}^\circ\mathbf{L}) \subset {}^\circ\mathbf{L},$$

$$\tilde{e}_i({}^\circ G^{\text{low}}(b)) = {}^\circ G^{\text{low}}(\tilde{E}_i(b)) \bmod v {}^\circ\mathbf{L}, \quad \tilde{f}_i({}^\circ G^{\text{low}}(b)) = {}^\circ G^{\text{low}}(\tilde{F}_i(b)) \bmod v {}^\circ\mathbf{L}.$$

(b) *The assignment $b \mapsto {}^\circ G^{\text{low}}(b) \bmod v {}^\circ\mathbf{L}$ yields a bijection from ${}^\circ B$ to the subset of ${}^\circ\mathbf{L}/v {}^\circ\mathbf{L}$ consisting of the $\tilde{f}_{i_1} \dots \tilde{f}_{i_l}(\phi_\pm)$'s. Further ${}^\circ G^{\text{low}}(b)$ is the unique element $x \in {}^\circ\mathbf{V}$ such that $x^\sharp = x$ and $x = {}^\circ G^{\text{low}}(b) \bmod v {}^\circ\mathbf{L}$.*

(c) *For each b, b' in ${}^\circ B$ let $E_{i,b,b'}, F_{i,b,b'} \in \mathcal{A}$ be the coefficients of ${}^\circ G^{\text{low}}(b')$ in $e_{\theta(i)}({}^\circ G^{\text{low}}(b))$, $f_i({}^\circ G^{\text{low}}(b))$ respectively. Then we have*

$$E_{i,b,b'}|_{v=1} = [F_i \Psi \mathbf{for}({}^\circ G^{\text{up}}(b')) : \Psi \mathbf{for}({}^\circ G^{\text{up}}(b))],$$

$$F_{i,b,b'}|_{v=1} = [E_i \Psi \mathbf{for}({}^\circ G^{\text{up}}(b')) : \Psi \mathbf{for}({}^\circ G^{\text{up}}(b))].$$

Proof: Part (a) follows from [EK3, thm. 4.1, cor. 4.4], [E, Section 2.3], and Proposition 3.31. The first claim in (b) follows from (a). The second one is obvious. Part (c) follows from Proposition 3.11. More precisely, by duality we can regard $E_{i,b,b'}$, $F_{i,b,b'}$ as the coefficients of ${}^\circ G^{\text{up}}(b)$ in $f_{\theta(i)}({}^\circ G^{\text{up}}(b'))$ and $e_i({}^\circ G^{\text{up}}(b'))$ respectively. Therefore, by Proposition 3.11 we can regard $E_{i,b,b'}|_{v=1}$, $F_{i,b,b'}|_{v=1}$ as the coefficients of $\Psi \mathbf{for}({}^\circ G^{\text{up}}(b))$ in $F_i \Psi \mathbf{for}({}^\circ G^{\text{up}}(b'))$ and $E_i \Psi \mathbf{for}({}^\circ G^{\text{up}}(b'))$ respectively. \square

REFERENCES

- [A] Ariki, S., *On the decomposition numbers of the Hecke algebra of $G(m, 1, n)$* , J. Math. Kyoto Univ. **36** (1996), 789-808.
- [E] Enomoto, N., *A quiver construction of symmetric crystals*, Int. Math. Res. Notices **12** (2009), 2200-2247.
- [EK1] Enomoto, N., Kashiwara, M., *Symmetric Crystals and the affine Hecke algebras of type B*.
- [EK2] Enomoto, N., Kashiwara, M., *Symmetric Crystals and LLT-Ariki type conjectures for the affine Hecke algebras of type B*.
- [EK3] Enomoto, N., Kashiwara, M., *Symmetric Crystals for \mathfrak{gl}_∞* , Publications of the Research Institute for Mathematical Sciences, Kyoto University **44** (2008), 837-891.
- [G] Grojnowski, I., *Representations of affine Hecke algebras (and affine quantum GL_n) at roots of unity*, Internat. Math. Res. Notices **5** (1994), 215.
- [KL] Khovanov, M., Lauda, A. D., *A diagrammatic approach to categorification of quantum groups I*, arXiv:0803.4121.
- [KM] Kashiwara, M., Miemietz, V., *Crystals and affine Hecke algebras of type D*.
- [L1] Lusztig, G., *Study of perverse sheaves arising from graded Lie algebras*, Advances in Math **112** (1995), 147-217.
- [L2] Lusztig, G., *Graded Lie algebras and intersection cohomology*, arXiv:0604.535.
- [Re] Reeder, M., *Isogenies of Hecke algebras and a Langlands correspondence for ramified principal series representations*, Representation Theory **6** (2002), 101-126.
- [R] Rouquier, R., *2-Kac-Moody algebras*, arXiv:0812.5023.

- [RR] Ram, A., Ramagge, J., *Affine Hecke algebras, cyclotomic Hecke algebras and Clifford theory.*
- [VV] Varagnolo, M., Vasserot, E., *Canonical bases and affine Hecke algebras of type B.*

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ PARIS 7, 175 RUE DU CHEVALERET, F-75013 PARIS, FAX : 01 44 27 78 18

E-mail address: `shan@math.jussieu.fr`

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE CERGY-PONTOISE, UMR CNRS 8088, F-95000 CERGY-PONTOISE, FAX : 01 34 25 66 45

E-mail address: `michela.varagnolo@math.u-cergy.fr`

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ PARIS 7, 175 RUE DU CHEVALERET, F-75013 PARIS, FAX : 01 44 27 78 18

E-mail address: `vasserot@math.jussieu.fr`