



HAL
open science

On the classification of singular flat structures on surfaces

Ousama Malouf

► **To cite this version:**

Ousama Malouf. On the classification of singular flat structures on surfaces. *Balkan Journal of Geometry and Its Applications*, 2010, 15 (1), pp.90-103. hal-00442654

HAL Id: hal-00442654

<https://hal.science/hal-00442654>

Submitted on 22 Dec 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the classification of singular flat structures on surfaces

Ousama Malouf

Abstract

We study in this work flat surfaces with conical singularities, that is, surfaces provided with a flat structure with conical singular points. Finding good parameters for these surfaces in the general case is an open question. We give an answer to this question in the case of flat structures on pairs of pants with one singular point. The question of decomposability of an arbitrary flat surface into flat pairs of pants is discussed.

Mathematics Subject Classifications (2010): 57M50, 32G15.

Key words: Flat structure, Teichmüller space, conical singularity.

1 Introduction

In this article, we consider flat surfaces with a finite number of conical singularities, that is, surfaces provided with a flat structure with conical singular points. A conical singular point has a total angle different from 2π .

Finding good parameters for these surfaces is still an open question (one should mention here that there are well-known good parameters for the subset of flat surfaces defined by quadratic differentials on Riemann surfaces). We treat here the question of classifying flat pairs of pants with one singularity. The decomposition of surfaces into pairs of pants is a common practice to study various structures on surfaces (see for instance [6]). The idea, which is usually attributed to Grothendieck, and which is developed by Feng Luo in [6], is to provide building blocks (generators) which permit to reconstruct any surface endowed with the studied structure following some specific rules (relations). It should be noted however that there exist flat surfaces which are not decomposable into flat pairs of pants by disjoint simple closed geodesics (for an example of flat surface of genus 3 with one singularity, see Example 5.1). But, it is possible to decompose such a surface into pairs of pants if we sacrifice some rules of decomposition, such as the simplicity of geodesics in the decomposition.

In Section 2, we give a definition of flat surface, we recall the formula of Gauss-Bonnet and we give some examples. In Section 3, we classify flat pairs of pants with one singularity and we study their space of parameters. In Section 4, we present the Teichmüller space of flat pairs of pants with one singularity and we study its topology. In Section 5, we discuss the decomposability of a surface into pairs of pants.

This work uses ideas from [2, 3] in which the authors consider the similar question for hyperbolic surfaces with singular points.

2 Preliminaries

Definition 2.1. Let $M_{g,k}$ (denoted by M if there is no confusion) be a surface of genus g with k boundary components, provided with a flat metric $d(.,.)$ with finitely many conical singularities $\Sigma = \{s_1, \dots, s_n\}$. We assume all the boundary components of M are closed curves (homeomorphic to circles). We denote by ∂M the boundary of M and by $\text{Int}(M) = M \setminus \partial M$ the interior of M . More precisely:

- For each point $x \in \text{Int}(M) \setminus \Sigma$ there is a neighborhood $U(x)$ isometric to a disc in the Euclidean plane.
- For each point $s \in \Sigma \cap \text{Int}(M)$ there is a neighborhood isometric to a Euclidean cone of total angle $0 < \theta < +\infty$, with $\theta \neq 2\pi$.
- For each point $x \in \partial M \setminus \Sigma$ there is a neighborhood isometric to a Euclidean sector with angle measure π at the image of x .
- For each point $s \in \Sigma \cap \partial(M)$ there is a neighborhood isometric to a Euclidean sector of angle $0 < \theta < +\infty$, with $\theta \neq \pi$.

The metric $d(.,.)$ will be called a *flat structure* on M , and M will be called a *flat surface (with boundary)* if it exists).

Definition 2.2. The curvature κ at a conical singularity s of total angle θ is $\kappa = 2\pi - \theta$. The curvature at a singular point on the boundary of angle θ is $\kappa = \pi - \theta$.

This definition of curvature is motivated by the following formula:

Proposition 2.1 (Gauss-Bonnet formula for closed surfaces [5, p. 113], [1, p. 190] or [7, p. 85-86] for a proof). *Let M be a closed flat surface of genus g with n conical singularities. Let $\theta_i, i = 1, \dots, n$ be the total angles at the singularities. The formula of Gauss-Bonnet which connects the number of singularities with their total angles and the genus is:*

$$\sum_{i=1}^n (2\pi - \theta_i) = (4 - 4g)\pi \quad (2.1)$$

The Euler characteristic of this surface is given by $\chi(M) = 2 - 2g$, and the formula of Gauss-Bonnet in terms of the Euler characteristic is:

$$\sum_{i=1}^n (2\pi - \theta_i) = 2\pi\chi(M).$$

Corollary 1 (Gauss-Bonnet formula for a disc with b holes). *Let M be a flat surface of genus 0 with boundary, with n singularities in the interior and m singularities on the boundary, and let b be the number of boundary components. Let $\theta_i, i = 1, \dots, n$ be the total angles of conical singularities in the interior, and let $\tau_j, j = 1, \dots, m$ be the total angles of singularities on the boundary. Then,*

$$\sum_{i=1}^n (2\pi - \theta_i) + \sum_{j=1}^m (\pi - \tau_j) = (4 - 2b)\pi. \quad (2.2)$$

Proof. By taking the double of M (in this operation, each singularity on the boundary is glued to its copy) we obtain a closed flat surface of genus $g = b - 1$ with $2n + m$ singularities. By the formula of Gauss-Bonnet (2.1) we get the result. \square

More generally, the Euler characteristic of a surface M of genus g with b boundary components is given by $\chi(M) = 2 - 2g - b$, and the formula of Gauss-Bonnet in terms of the Euler characteristic is:

$$\sum_{i=1}^n (2\pi - \theta_i) + \sum_{j=1}^m (\pi - \tau_j) = 2\pi\chi(M).$$

Example 2.1. Let M be a pair of pants (a surface homeomorphic to a disc with two holes), equipped with a flat structure with one conical singularity s in its interior and such that each component of the boundary $\partial M = c_1 \cup c_2 \cup c_3$ is geodesic without singularities.

From s we take three geodesic segments $d_i, i = 1, 2, 3$ which realize the distances between s and c_i respectively, Figure 3. We denote by l_i, r_i the lengths of c_i, d_i respectively. We cut and open M along the geodesic segments d_i . Then we obtain a connected surface P which can be decomposed into three rectangles $R_i, i = 1, 2, 3$ whose lengths of sides are l_i, r_i respectively, and one triangle whose sides have lengths l_i , as shown in Figure 4.

Following the inverse procedure, we consider a triangle T of side lengths l_i , and three rectangles R_i each of which shares a side with T and the other side being of length r_i as shown in Figure 4.

Now, by identifying the sides of equal length r_i , we obtain a flat pair of pants with one conical singularity. Note that the total angle of the singularity is $\theta = 4\pi$.

This example will be studied later in Section 3.

Example 2.2. We cannot build a flat pair of pants with no singularities at all. Indeed, if we assume that such a pair of pants exists, by the formula (2.2) we find $0 = -2\pi$ which is impossible.

Finally, by gluing along their boundaries $2g - 2$ flat pair of pants, each one with a single singularity, we obtain a closed flat surface of genus g with $2g - 2$ conical singularities.

3 The geometry of a flat pair of pants with a conical singularity

In this section we assume that M is a flat (singular) surface with boundary homeomorphic to a disc with two holes. We assume further that M has only one conical singularity s . We call such a surface a *flat pair of pants with one singularity* (or just *flat pair of pants*). An example of such structure is obtained by taking the metric associated to two transverse measured foliations with one singularity on M (see Figure 1). It is clear from this picture that the parameter space for the transverse measures of these two transverse foliations has dimension four, and it produces a space of flat structures of dimension four on the pair of pants, with one singular point. It will follow from the discussion below




Figure 1: The two transverse measured foliations on the pair of pants induce a flat structure with one singular point




Figure 2: We can change the isometry type of a flat pair of pants without changing the boundary curve lengths

(see the remark after Definition 4.1) that there are examples of flat structures on M which do not arise from pairs of transverse measured foliations. The aim of this section is to find a set of real parameters which determine the geometry of M .

The first natural guess is that the lengths of boundary geodesics might be good parameters, in analogy with the case of hyperbolic pairs of pants (described for instance in [4]). This is easily seen to be false, as one can glue Euclidean cylinders to boundaries of flat pairs of pants, changing the isometry type, without changing the boundary component length (see Figure 2). Thus, the parameter space for flat structures on pair of pants is more complicated than the parameter space for hyperbolic structure on such surfaces.

Let $\partial M = c_1 \cup c_2 \cup c_3$ denote the boundary and denote by $d(\cdot, \cdot)$ the distance function on M . Let d_i be a geodesic segment which realizes the distance $d(s, c_i)$ between the singularity s and the boundary component c_i . This segment intersects the boundary component with a right angle. We denote by l_i the length of c_i and by r_i the length of d_i . Obviously, the parameters r_i , $i = 1, 2, 3$ depend on the position of s . We have the following proposition.

Proposition 3.1. *The real parameters $l_i, r_i, i = 1, 2, 3$ determine a unique flat pair of pants M with one singularity up to isometry, where the l_i are the lengths of the boundary components c_i , and the r_i are the lengths of the geodesic segments between s and the c_i .*

Proof. Every boundary component c_i , $i \in \{1, 2, 3\}$ is geodesic, even if the singu-




Figure 3: Flat pair of pants with one singularity in the interior




Figure 4: After cut. l_i, r_i are the lengths of c_i, d_i respectively

larity is on a boundary component, since by the formula of Gauss-Bonnet, the angle at s will be 3π . By our assumption on the uniqueness of the singularity, d_i does not meet another singularity on the component c_i of the boundary, and so, since c_i is geodesic, d_i is orthogonal to c_i . For the same reasons, d_i does not share any of its interior points with boundary components.

First case: We assume that $s \in \text{Int}(M)$. This implies that the lengths of d_i are different from zero, $r_i \neq 0, i = 1, 2, 3$. We cut M along $d_1 \cup d_2 \cup d_3$, see Figure 3, to obtain a polygonal flat surface P homeomorphic to a disc without singularities in the interior, see Figure 4.

None of the parameters $l_i, i = 1, 2, 3$ can be zero, because if one of them is zero, the surface M loses one of its boundary components. This fact implies that we always have three copies of s after the cut. Let us denote by $s_i, i = 1, 2, 3$ the copy which lies between $d_{(i+1) \bmod 3}$ and $d_{(i+2) \bmod 3}$, Figure 4. We can draw in P the geodesics between these copies. These geodesic segments bound a triangle $T = (s_1 s_2 s_3)$, Figure 5, which can be degenerate, with none of its angles greater than π . For instance, in Figure 6 the angle $\angle(s_2 s_1 s_3) = \pi$. This angle cannot be greater than π , because this implies that d_1 , along which the cut was made, was not a geodesic segment which represents the distance between s and c_1 . This contradicts the assumption.

We recognize easily, Figure 4, in addition to T , three rectangles $R_i, i = 1, 2, 3$. A rectangle $R_i, i \in \{1, 2, 3\}$ is bounded by a side of T , c_i and the two copies of d_i . P is composed of T and $R_i, i = 1, 2, 3$.

A rectangle is uniquely determined up to isometry by its length and height, and a triangle is uniquely determined up to isometry by its lengths of sides. Then, given the parameters $l_i, r_i, i = 1, 2, 3$, where the $l_i, i = 1, 2, 3$ satisfy the triangle inequalities, we can construct a unique pair of pants with one singularity by gluing together the three rectangles R_i and the triangle T given by the parameters $l_i, r_i, i = 1, 2, 3$.




Figure 5: Here we see the triangle on the pair of pants




Figure 6: The triangle is degenerate

The triangle could be degenerate, and this case lies on the boundary of the space of triangles. This happens only when

$$l_i = l_{(i+1) \bmod 3} + l_{(i+2) \bmod 3} \quad (3.1)$$

for some $i \in \{1, 2, 3\}$. Figure 6.

Second case: We assume that $s \in \partial M$. Then at least one of r_i , $i = 1, 2, 3$ equals zero. More precisely, s belongs to only one c_i , $i \in \{1, 2, 3\}$, and so only one r_i equals zero, because if s belongs to more than one boundary component, it does not have a neighborhood homeomorphic to a disc. In such a case we call this a *degenerate pair of pants*. We will outline these cases later.




Figure 7: Singularity on the boundary




Figure 8: The resulting heptagon after cut

Without loss of generality, assume $s \in c_1$, as in Figure 7. Cut along $d_2 \cup d_3$ (d_1 being reduced to s) to obtain a polygonal flat surface P homeomorphic to a disc without singularities in the interior. For the same reason as before, we have three copies of s . The geodesic segment between s_2, s_3 is c_1 itself since c_1 is geodesic.

The geodesic segments between the copies of s bound a triangle $T = (s_1s_2s_3)$ which can be degenerate, with none of its angles greater than π (for the same reason as in the first case). It is easy to recognize, in addition to T , two rectangles $R_i, i = 2, 3$. Figure 8.

Knowing the parameters $l_i, r_i, i = 1, 2, 3$ we can determine if the singularity is on the boundary and on which component. The rectangles R_i and the triangle T are uniquely determined by the parameters and so is the flat structure.

If Relation (3.1) is satisfied, the triangle is degenerate. In this case we notice that only one $r_i, i \in \{1, 2, 3\}$ that correspond to the boundaries on the right hand side of this identity can take the value zero. The other cases are degenerate.

We proved that to each flat structure on M with one singular point we may associate unique set of values $l_i, r_i, i = 1, 2, 3$. Conversely, to each choice of values, we may associate a unique flat structure on M .


We deduce that each flat pair of pants M is uniquely determined, up to isometry, by the parameters l_i and r_i . And vice versa.

□


Let us give the limits of parameters in view of the last proof. The length parameters can take any values satisfying these conditions:

- $0 < l_i < \infty, i = 1, 2, 3,$
- $l_i \leq l_{(i+1) \bmod 3} + l_{(i+2) \bmod 3}, i = 1, 2, 3,$
- $0 \leq r_i < \infty, i = 1, 2, 3,$
- $0 < r_i + r_{(i+1) \bmod 3}, i = 1, 2, 3$
- If $l_i = l_{(i+1) \bmod 3} + l_{(i+2) \bmod 3}$, for some $i \in \{1, 2, 3\}$ then $0 < r_i < \infty$

Remark. Let M be a flat surface with one singularity s , which is homeomorphic to a disc with n holes, $n \geq 3$. We denote by c_i the boundary components of M and by d_i the geodesic segments from s to c_i . If we set l_i the lengths of c_i and r_i the lengths of d_i , then the parameters l_i, r_i do not characterize M in the sense of Proposition 3.1. This follows because if we cut M along $d_1 \cup d_2 \cup \dots \cup d_n$ then instead of a triangle $(s_1s_2s_3)$ we get a $(n+1)$ -gon which, of course, is not uniquely determined by the lengths of its edges.



Two rectangles are degenerate



Three rectangles are degenerate

Figure 9: Neighborhood of singularity




Figure 10: The triangle and two rectangles are degenerate

We return to the case where M is a pair of pants and $\partial M = c_1 \cup c_2 \cup c_3$. By the previous discussion, where we saw how a flat pair of pants with one singularity can be cut into pieces, three rectangles and one triangle, we find it convenient to introduce these terms: We say that *the triangle is degenerate* if the condition:

$$l_i = l_{(i+1) \bmod 3} + l_{(i+2) \bmod 3}, \text{ for some } i \in \{1, 2, 3\}$$

is satisfied. If not, we say that *the triangle is non-degenerate*. We say that *the rectangle $R_i, i \in \{1, 2, 3\}$ is degenerate* if $r_i = 0$ for some $i \in \{1, 2, 3\}$. If not, we say that *the rectangle R_i is non-degenerate*.

Remark (Degenerate cases). A degenerate pair of pants appears when the singularity does not have a neighborhood homeomorphic to a disc in the Euclidean plane. From the previous proof, we can conclude the following degenerate cases:

- In the case when the triangle is non-degenerate, if two or three rectangles $R_i, i \in \{1, 2, 3\}$ are degenerate we have a degenerate pair of pants. A neighborhood of the singularity is similar to one of the forms in Figure 9.
- In the case when the triangle is degenerate, that is, for some $i \in \{1, 2, 3\}$ identity (3.1) is satisfied, if both rectangles $R_{(i+1) \bmod 3}, R_{(i+2) \bmod 3}$ are degenerate we have a degenerate pair of pants. A neighborhood of the singularity is similar to Figure 10.

We should finally pay attention that in this case when R_i is degenerate, the resulting surface is not a pair of pants since it has four holes. Figure 11.

After the previous discussion, we will introduce new parameters for flat pairs of pants. These parameters seem to be more convenient. For this, let now k_i be a geodesic segment which realizes the distance between $c_{(i+1) \bmod 3}$ and $c_{(i+2) \bmod 3}, i = 1, 2, 3$, see Figure 12. Denote by l_i the length of c_i and by a_i the length of k_i .

Definition 3.1. The six non-negative parameters $l_i, a_i, i = 1, 2, 3$ will be called the *distance parameters* of M .




Figure 11: Impossible case




Figure 12: The geodesic k_i represents the distance a_i between $c_{(i+1) \bmod 3}$ and $c_{(i+2) \bmod 3}$

We have the following theorem.

Proposition 3.2. *The distance parameters $l_i, a_i, i = 1, 2, 3$ determine a unique flat pair of pants M with one singularity.*

Proof. By the proof of Proposition 3.1 we see that flat pairs of pants with one singularity are uniquely determined by the parameters $l_i, r_i, i = 1, 2, 3$, and correspond to one of the following cases:

1. If the triangle is non-degenerate, we distinguish two cases.
 - (a) If all the rectangles R_i are non-degenerate, it is easy, after the cut, to see that geodesic segments between boundaries pass all by the singularity. Figure 13.
So we have:

$$r_i + r_{(i+1) \bmod 3} = a_{(i+2) \bmod 3}, i = 1, 2, 3. \quad (3.2)$$




Figure 13: After cut, l_i, a_i are the lengths of c_i, k_i respectively




Figure 14: Permitted gluing of two rectangles

These relations prove that knowing $l_i, a_i, i = 1, 2, 3$ we can determine $r_i, i = 1, 2, 3$, and so, all the parameters needed to determine the flat pair of pants in this case, by Proposition 3.1.

- (b) If only one rectangle is degenerate then the pair of pants can be cut into two rectangles and one non-degenerate triangle as in Figure 8. By the same relations (3.2) we can determine the flat pair of pants. The singularity, in this case, is on the boundary component which corresponds to the degenerate rectangle.

2. If the triangle is degenerate, we also distinguish two cases.

- (a) If all the rectangles $R_i, i = 1, 2, 3$ are non-degenerate, we see that the relations (3.2) also hold here and so the flat pair of pants is well determined, in this case, by the distance parameters. Figure 6.
- (b) If one of $R_{(i+1) \bmod 3}, R_{(i+2) \bmod 3}$ is degenerate then the pair of pants can be cut into two rectangles. Again, it is determined by relations (3.2). Figure 14.

□

Here we give the limits of the new parameters in view of the last proof. The distance parameters can take any values in the limit of these conditions:

- 1) $0 < l_i < \infty, i = 1, 2, 3,$
- 2) $l_i \leq l_{(i+1) \bmod 3} + l_{(i+2) \bmod 3}, i = 1, 2, 3,$
- 3) $0 < a_i < \infty, i = 1, 2, 3,$
- 4) $a_i \leq a_{(i+1) \bmod 3} + a_{(i+2) \bmod 3}, i = 1, 2, 3,$ (Using Relations (3.2)),
- 5) If $l_i = l_{(i+1) \bmod 3} + l_{(i+2) \bmod 3},$ for some $i \in \{1, 2, 3\},$ then $a_i < a_{(i+1) \bmod 3} + a_{(i+2) \bmod 3}.$ (Using Relations (3.2)).

It is easy to see that these conditions are equivalent to preceding ones. In addition, it is interesting to see that these conditions are equivalent to the fact that we have two Euclidean triangles (first four conditions) for which a degenerate case of one triangle prevents a degenerate case of the other (fifth condition).

We denote by \mathcal{C} the set of all flat structures defined by the parameters $l_i, a_i, i = 1, 2, 3$ under the previous conditions, and by \mathcal{B} the set of all 6-tuples

$(l_1, l_2, l_3, a_1, a_2, a_3) \in \mathbb{R}^6$ which satisfy these conditions. We can define a one-to-one mapping $\Phi : \mathcal{C} \rightarrow \mathcal{B}$ such that to a flat structure $f \in \mathcal{C}$ corresponds the unique 6-tuple $(l_1, l_2, l_3, a_1, a_2, a_3)$ of \mathcal{B} which determines f . Obviously we have $\Phi(\mathcal{C}) = \mathcal{B}$.

4 The Teichmüller space of flat pairs of pants with one singularity

Let us denote by $\mathcal{F}(M)$ the space of all flat structures with one singularity on a pair of pants M . We fix an orientation on M and let $\text{Homeo}^+(M, \partial)$ be the set of homeomorphisms of M which preserve the orientation and each boundary component of M (setwise). It is well known that each element of $\text{Homeo}^+(M, \partial)$ is isotopic to the identity (see Exposé 2 in [4]). The space $\text{Homeo}^+(M, \partial)$ acts on $\mathcal{F}(M)$ as follows: If $h \in \text{Homeo}^+(M, \partial)$ and $f \in \mathcal{F}(M)$ then $(h, f) \mapsto h * f$ where $h * f(x, y) := f(h(x), h(y))$.

Definition 4.1. We define the Teichmüller space $\mathcal{T}(M)$ of M as the quotient $\mathcal{F}(M)/\text{Homeo}^+(M, \partial)$.

Obviously, $\mathcal{T}(M)$ consists of all flat structures which belong to \mathcal{C} , the set of all flat structures defined by the parameters $l_i, a_i, i = 1, 2, 3$. To each flat structure with one singularity on M we may associate a unique configuration, which consists of a triangle T and three rectangles $R_i, i = 1, 2, 3$, glued as in Figure 4. Conversely, to each configuration we may associate a unique flat structure with one singularity on M . This defines a mapping $\Phi : \mathcal{T}(M) \rightarrow \mathcal{B}$ which is one-to-one, and $\Phi(\mathcal{T}(M)) = \mathcal{B}$.

Remark (Due to Athanase Papadopoulos). Let $\mathcal{T}_{\mathcal{M}}(M)$ be the Teichmüller space of flat structures with one singularity on a pair of pants defined by a pair of transverse measured foliations. This space is of dimension 4. Since $\mathcal{T}(M)$, the Teichmüller space of all flat structures with one singularity on a pair of pants, is of dimension 6, as we showed here, then $\mathcal{T}_{\mathcal{M}}(M) \subsetneq \mathcal{T}(M)$. Thus, the space of flat structures we counter here is larger than the space of flat structures induced by quadratic differentials with one zero.

Proposition 4.1. *The mapping $\Phi : \mathcal{T}(M) \rightarrow \mathcal{B}$ is continuous and open. Thus, Φ is homeomorphism.*

Proof. As explained in Proposition 3.2, Φ is a bijection. If we consider the Euclidean distance between tuples of parameters $(l_1, l_2, l_3, a_1, a_2, a_3)$ and the following distance between two flat structures $f_1, f_2 \in \mathcal{T}(M)$:

$$d(f_1, f_2) = \sup_{x, y \in M} |f_1(x, y) - f_2(x, y)|$$

we can see easily that both Φ and Φ^{-1} are continuous. □

Obviously, \mathcal{B} is a non bounded convex subset of \mathbb{R}^6 . In fact, the set of parameters $l_i, i = 1, 2, 3$ can be seen as the space of all triangles which could be degenerate, but without those which have a side of length zero. This is, a non bounded pyramid of three faces with its sides and apex deleted. This space is

convex in \mathbb{R}^3 . The same can be said about the set of parameters a_i , $i = 1, 2, 3$. Thus, \mathcal{B} is convex in \mathbb{R}^6 .

The boundary $\partial\mathcal{B}$ has six components which are non bounded convex subsets of \mathbb{R}^5 . They correspond to cases of equality in the triangle inequality. Let us denote by $\partial_i\mathcal{B}_l$, $i \in \{1, 2, 3\}$ the boundary component defined by

$$l_i = l_{(i+1) \bmod 3} + l_{(i+2) \bmod 3}.$$

The sets $\partial_i\mathcal{B}_l$, $i = 1, 2, 3$ are pairwise disjoint subsets of $\partial\mathcal{B}$. Similarly, we denote by $\partial_j\mathcal{B}_a$, $j \in \{1, 2, 3\}$ the boundary component defined by

$$a_j = a_{(j+1) \bmod 3} + a_{(j+2) \bmod 3}.$$

The sets $\partial_j\mathcal{B}_a$, $j = 1, 2, 3$ are also pairwise disjoint subsets of $\partial\mathcal{B}$.

The intersection $\partial_{i,j}(\partial\mathcal{B}) = \partial_i\mathcal{B}_l \cap \partial_j\mathcal{B}_a$, is homeomorphic to \mathbb{R}^4 if $i \neq j$, and empty if $i = j$. This means that each boundary component of \mathcal{B} has two convex connected boundary components homeomorphic to \mathbb{R}^4 .

Then, by the homeomorphism Φ , we have the following description of the Teichmüller space:

Corollary 2. *$\mathcal{T}(M)$ is homeomorphic to a non-compact submanifold of \mathbb{R}^6 of dimension 6, with a natural cell-structure, having six cells of codimension one on its boundary.*

We know that any convex subset of Euclidean n-space \mathbb{E}^n is contractible. This gives us the following result:

Theorem 4.2. *$\mathcal{T}(M)$ is a contractible space.*

In the next section we start a discussion of the decomposition into pairs of pants of closed surfaces with one singularity. Details will be given in subsequent work.

5 Closed flat surfaces with one singularity

In general, a flat surface with one singularity is not decomposable by disjoint simple closed geodesics into pairs of pants. For this, we have the following example:

Example 5.1. Take a closed flat surface M_3 of genus 3 with one singularity s . Suppose that we are able to decompose it by disjoint simple closed geodesics into pairs of pants. By the fact that there is no flat pair of pants without any singularity, we conclude that every pair of pants resulting from the decomposition has the singularity s on its boundary. We know that a decomposition of M_3 by disjoint simple closed geodesics gives rise to four pairs of pants. This means that four boundary components should share the singularity. This is impossible under the present rules of identification (one boundary component is identified to one boundary component). Then, the decomposition is impossible.

The decomposition becomes possible if we change the rules of composition. For example, admitting geodesics to be non-simple or non-disjoint. That is, admitting the identification of parts of boundary components rather than boundary components entirely. I will not discuss here these rules, their study can be dealt with a separate work later.

The result in Example 5.1 can be generalized as follows:

Let M be a closed flat surface of genus $g \geq 3$ with a single conical singularity s . Then M cannot be decomposed into pairs of pants by disjoint simple closed geodesics.

Acknowledgements

I would like to thank Athanase Papadopoulos for all the corrections and discussions during the preparation of this article. I also thank Daniele Alesandrini for his time.

References

- [1] A. Aleksandrov and V. Zalgaller. *Intrinsic geometry of surfaces*. Translations of Mathematical Monographs. 15. Providence, RI: American Mathematical Society (AMS). VI, 327 p. , 1967.
- [2] C. Charitos and I. Papadoperakis. Parameters for generalized Teichmüller spaces. Papadopoulos, Athanase (ed.), Handbook of Teichmüller theory. Volume I. Zürich: European Mathematical Society (EMS). IRMA Lectures in Mathematics and Theoretical Physics 11, 471-506 (2007).
- [3] C. Charitos and I. Papadoperakis. On the geometry of hyperbolic surfaces with a conical singularity. *Ann. Global Anal. Geom.*, 23(4):323–357, 2003.
- [4] A. Fathi, F. Laudenbach, and V. Poenaru. *Travaux de Thurston sur les surfaces*. *Séminaire Orsay*. 2nd ed. Centre National de la Recherche Scientifique. Astérisque, 66-67. Paris: Société Mathématique de France. 286 p., 1991.
- [5] H. Hopf. *Differential geometry in the large*. *Seminar lectures New York University 1946 and Stanford University 1956*. With a preface by S. S. Chern. 2nd ed. Lecture Notes in Mathematics, 1000. Berlin etc.: Springer-Verlag. vi, 184 p., 1989.
- [6] F. Luo. Grothendieck’s reconstruction principle and 2-dimensional topology and geometry. Papadopoulos, Athanase (ed.), Handbook of Teichmüller theory. Volume II. Zürich: European Mathematical Society (EMS). IRMA Lectures in Mathematics and Theoretical Physics 13, 733-765 (2009).
- [7] M. Troyanov. Les surfaces euclidiennes à singularités coniques. (Euclidean surfaces with cone singularities). *L’Enseignement Mathématique*, II(32):79–94, 1986.

Author’s address:

Ousama Malouf
Institut de Recherche Mathématique Avancée
7 rue René Descartes
F-67084 Strasbourg Cedex
France
malouf@math.u-strasbg.fr