



**HAL**  
open science

# Transfer of Fourier multipliers into Schur multipliers and sumsets in a discrete group

Stefan Neuwirth, Éric Ricard

► **To cite this version:**

Stefan Neuwirth, Éric Ricard. Transfer of Fourier multipliers into Schur multipliers and sumsets in a discrete group. 2010. hal-00451399v1

**HAL Id: hal-00451399**

**<https://hal.science/hal-00451399v1>**

Preprint submitted on 29 Jan 2010 (v1), last revised 22 Jun 2012 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Transfer of Fourier multipliers into Schur multipliers and sumsets in a discrete group

Stefan Neuwirth      Éric Ricard

## Abstract

We inspect the relationship between relative Fourier multipliers on noncommutative Lebesgue-Orlicz spaces of a discrete group  $\Gamma$  and relative Toeplitz-Schur multipliers on Schatten-von-Neumann-Orlicz classes. Four applications are given: lacunary sets; unconditional Schauder bases for the subspace of a Lebesgue space determined by a given spectrum  $\Lambda \subseteq \Gamma$ ; the norm of the Hilbert transform and the Riesz projection on Schatten-von-Neumann classes with exponent a power of 2; the norm of Toeplitz Schur multipliers on Schatten-von-Neumann classes with exponent less than 1.

## 1 Introduction

Let  $\Lambda$  be a subset of  $\mathbb{Z}$  and let  $x$  be a bounded measurable function on the circle  $\mathbb{T}$  with Fourier spectrum in  $\Lambda$ : we write  $x \in L^\infty_\Lambda$ ,  $x \sim \sum_{k \in \Lambda} x_k z^k$ . The matrix of the associated operator  $y \mapsto xy$  on  $L^2$  with respect to its trigonometric basis is the Toeplitz matrix

$$(x_{r-c})_{(r,c) \in \mathbb{Z} \times \mathbb{Z}} = \begin{matrix} & \dots & 1 & 0 & -1 & \dots \\ \vdots & \left( \begin{array}{ccccc} \ddots & \ddots & \ddots & \ddots & \ddots \\ \ddots & x_0 & x_1 & x_2 & \ddots \\ \ddots & \ddots & x_{-1} & x_0 & x_1 & \ddots \\ \ddots & \ddots & x_{-2} & x_{-1} & x_0 & \ddots \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \end{array} \right) \\ 1 & & & & & \\ 0 & & & & & \\ -1 & & & & & \\ \vdots & & & & & \end{matrix}$$

with support in  $\check{\Lambda} = \{(r, c) : r - c \in \Lambda\}$ .

This is an entry point to the interplay between harmonic analysis and operator theory. In the general case of a discrete group  $\Gamma$ , the counterpart to

---

This work is partially supported by ANR 06-BLAN-0015.

2000 *Mathematics subject classification*: Primary 47B49; Secondary 43A22, 43A46, 46B28.

*Key words and phrases*: Fourier multiplier, Toeplitz Schur multiplier, lacunary set, unconditional approximation property, Hilbert transform, Riesz projection.

a bounded measurable function is defined as a bounded operator on  $\ell_\Gamma^2$  whose matrix has the form  $(x_{rc^{-1}})_{(r,c) \in \Gamma \times \Gamma}$  for some sequence  $(x_\gamma)_{\gamma \in \Gamma}$ . This will be the framework of the body of this article, while the introduction sticks to the case  $\Gamma = \mathbb{Z}$ .

We are concerned with two kinds of multipliers. A sequence  $\varphi = (\varphi_k)_{k \in \Lambda}$  defines

- the relative Fourier multiplication operator on trigonometric polynomials with spectrum in  $\Lambda$  by

$$\sum_{k \in \Lambda} x_k z^k \mapsto \sum_{k \in \Lambda} \varphi_k x_k z^k; \quad (1)$$

- the relative Schur multiplication operator on finite matrices with support in  $\check{\Lambda}$  by

$$(x_{r,c})_{(r,c) \in \mathbb{Z} \times \mathbb{Z}} \mapsto (\check{\varphi}_{r,c} x_{r,c})_{(r,c) \in \mathbb{Z} \times \mathbb{Z}} \quad (2)$$

where  $\check{\varphi}_{r,c} = \varphi_{r-c}$ .

Marek Bożejko and Gero Fendler proved that these two multipliers have the same norm. The operator (1) is nothing but the restriction of (2) to Toeplitz matrices. They noted that it is automatically *completely* bounded: it has the same norm when acting on trigonometric series with operator coefficients  $x_k$ , and this permits to remove this restriction. Schur multiplication is also automatically completely bounded.

This observation has been extended by Gilles Pisier to multipliers acting on a translation invariant Lebesgue space  $L_A^p$  and on the subspace  $S_A^p$  of elements of a Schatten-von-Neumann class supported by  $\check{\Lambda}$ , respectively: it yields that the complete norm of a relative Schur multiplier (2) remains bounded by the complete norm of the relative Fourier multiplier (1).

But  $L_A^p$  is not a subspace of  $S_A^p$ , so that a relative Fourier multiplier may not be viewed anymore as the restriction of a relative Schur multiplier to Toeplitz matrices. We point out that this difficulty may be overcome by using Szegő's limit theorem: a bounded measurable real function is the weak\* limit of the normalised counting measure of eigenvalues of finite truncates of its Toeplitz matrix. Note that other types of approximation are also available, as the completely positive approximation property and Reiter sequences combined with complex interpolation: they are compared in Section 3 in terms of local embeddings of  $L^p$  into  $S^p$ . They are more canonical than Szegő's limit theorem, but give no access to general Orlicz norms.

**Theorem 1.1.** *Let  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be a continuous nondecreasing function vanishing only at 0. The norm of the following operators is equal:*

- the relative Fourier multiplication operator (1) on the Lebesgue-Orlicz space  $L_A^\psi(S^\psi)$  of  $S^\psi$ -valued trigonometric series with spectrum in  $\Lambda$ ;
- the relative Schur multiplication operator (2) on the Schatten-von-Neumann-Orlicz class  $S_A^\psi(S^\psi)$  of  $S^\psi$ -valued matrices with support in  $\check{\Lambda}$ .

Look at Theorem 2.5 for the precise statement in the general case of an amenable group  $\Gamma$ , for which a block matrix variant of Szegő's limit theorem in the style of Erik Bédos [2], Theorem 2.1, is available.

An application of this theorem to the class of all unimodular Fourier multipliers yields a transfer of lacunary subsets into lacunary matrix patterns. Call  $\Lambda$  *unconditional in  $L^p$*  if  $(z^k)_{k \in \Lambda}$  is an unconditional basis of  $L^p_\Lambda$ , and  $\check{\Lambda}$  *unconditional in  $S^p$*  if the sequence  $(e_q)_{q \in \check{\Lambda}}$  of elementary matrices is an unconditional basis of  $S^p_{\check{\Lambda}}$ . These properties are also known as  $\Lambda(\max(2, p))$  and  $\sigma(p)$ , respectively; they have natural “complete” counterparts that are also known as  $\Lambda(p)_{cb}$  ( $K(p)_{cb}$  if  $p \leq 2$ ) and  $\sigma(p)_{cb}$ , respectively: see Definitions 4.1 and 4.2.

**Corollary 1.2.** *If  $\check{\Lambda}$  is unconditional in  $S^p$ , then  $\Lambda$  is unconditional in  $L^p$ .  $\check{\Lambda}$  is completely unconditional in  $S^p$  if and only if  $\Lambda$  is completely unconditional in  $L^p$ .*

Look at Proposition 4.3 for the precise statement in the general case of a discrete group  $\Gamma$ .

The two most prominent multipliers are the Riesz projection and the Hilbert transform. The first consists in letting  $\varphi$  be the indicator function of nonnegative integers and transfers into the upper triangular truncation of matrices. The second corresponds to the sign function and transfers into the Hilbert matrix transform. We obtain the following partial results.

**Theorem 1.3.** *The norm of the matrix Riesz projection and of the matrix Hilbert transform on  $S^\psi(S^\psi)$  coincide with their norm on  $S^\psi$ .*

- If  $p$  is a power of 2, the norm of the matrix Hilbert transform on  $S^p$  is  $\cot(\pi/2p)$ .
- The norm of the matrix Riesz projection on  $S^4$  is  $\sqrt{2}$ .

The transfer technique lends itself naturally to the case where  $\Lambda$  contains a sumset  $R+C$ : if subsets  $R'$  and  $C'$  are extracted so that the  $r+c$  with  $r \in R'$  and  $c \in C'$  are pairwise distinct, they may play the rôle of rows and columns. Here are consequences of the conditionality of the sequence of elementary matrices  $e_{r,c}$  in  $S^p$  for  $p \neq 2$  and of the unboundedness of the Riesz transform on  $S^1$  and  $S^\infty$ , respectively.

**Theorem 1.4.** *If  $(z^k)_{k \in \Lambda}$  is a completely unconditional basis of  $L^p_\Lambda$  with  $p \neq 2$ , then  $\Lambda$  does not contain sumsets  $R+C$  of arbitrarily large sets.*

- If  $L^1_\Lambda$  admits some completely unconditional approximating sequence, or
- if the space  $C_\Lambda$  of continuous functions with spectrum in  $\Lambda$  admits some unconditional approximating sequence,

then  $\Lambda$  does not contain the sumset  $R+C$  of two infinite sets.

The proof of the second part of this theorem consists in constructing infinite subsets  $R'$  and  $C'$ , and skipped block sums  $\sum(T_{k_{j+1}} - T_{k_j})$  of a given approximating sequence that act like the projection on the “upper triangular” part of  $R' + C'$ . Look at Proposition 4.8 and Theorem 7.4 for the precise statement in the general case of a discrete group  $\Gamma$ .

In the case of quasi-normed Schatten-von-Neumann classes  $S^p$  with  $p < 1$ , the transfer technique yields a new proof for the following result of Alexey Alexandrov and Vladimir Peller.

**Theorem 1.5.** *Let  $0 < p < 1$ . The Fourier multiplier  $\varphi$  is contractive on  $L^p$  or on  $L^p(S^p)$  if and only if the Schur multiplier  $\check{\varphi}$  is contractive on  $S^p$  or on  $S^p(S^p)$ , if and only if the sequence  $\varphi$  is the Fourier transform of an atomic measure of the form  $\sum a_g \delta_g$  on  $\mathbb{T}$  with  $\sum |a_g|^p \leq 1$ .*

The emphasis put on *relative* Schur multipliers motivates the natural question of how the norm of an elementary Schur multiplier, that is a rank 1 matrix  $(\varrho_{r,c}) = (x_r y_c)$ , gets affected when the action of  $\varrho$  is restricted to matrices with a given support. The surprising answer is the following.

**Theorem 1.6.** *Let  $I \subseteq R \times C$  and consider  $(x_r)_{r \in R}$  and  $(y_c)_{c \in C}$ . The relative Schur multiplier on  $S_I^\infty$  given by  $(x_r y_c)_{(r,c) \in I}$  has norm  $\sup_{(r,c) \in I} |x_r y_c|$ .*

Let us finally describe the content of this article. Section 2 is devoted to transfer techniques for Fourier and Schur multipliers provided by a block matrix Szegő limit theorem. This theorem provides local embeddings of  $L^\psi$  into  $S^\psi$ ; Section 3 shows how interpolation may be used to define such embeddings for the scale of  $L^p$  spaces. Section 4 is devoted to the transfer of lacunary sets into lacunary matrix patterns; the unconditional constant of a set  $A$  is related to the size of the sumsets it contains. Section 5 deals with Toeplitz Schur multipliers for  $p < 1$  and comments on the case  $p \geq 1$ . The Riesz projection and the Hilbert transform are studied in Section 6. In Section 7, the presence of sumsets in a spectrum  $\Lambda$  is shown to be an obstruction for the existence of completely unconditional bases for  $L_A^p$ . The last section provides a norm-preserving extension for partially specified rank 1 Schur multipliers.

**Notation and terminology.** Let  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ .

Given an index set  $C$  and  $c \in C$ ,  $e_c$  is the sequence defined on  $C$  as the indicator function  $\chi_{\{c\}}$  of the singleton  $\{c\}$ , so that  $(e_c)_{c \in C}$  is the canonical Schauder basis of the Hilbert space of square summable sequences indexed by  $C$ , denoted by  $\ell_C^2$ . We will use the notation  $\ell_n^2 = \ell_{\{1,2,\dots,n\}}^2$  and  $\ell^2 = \ell_{\mathbb{N}}^2$ .

Given a product set  $I = R \times C$  and  $q = (r, c)$ , the indicator function  $e_q = e_{r,c}$  is the *elementary matrix* identified with the linear operator from  $\ell_C^2$  to  $\ell_R^2$  that maps  $e_c$  on  $e_r$  and all other basis vectors on 0. The *matrix coefficient* at coordinate  $q$  of a linear operator  $x$  from  $\ell_C^2$  to  $\ell_R^2$  is  $x_q = \text{tr } e_q^* x$  and its *matrix representation* is  $(x_q)_{q \in R \times C} = \sum_{q \in R \times C} x_q e_q$ . The *support* or *pattern* of  $x$  is  $\{q \in R \times C : x_q \neq 0\}$ .

The space of all bounded operators from  $\ell_C^2$  to  $\ell_R^2$  is denoted by  $\mathbb{B}(\ell_C^2, \ell_R^2)$ , and its subspace of compact operators is denoted by  $S^\infty$ .

Let  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be a continuous nondecreasing function vanishing only at 0. The *Schatten-von-Neumann-Orlicz class*  $S^\psi$  is the space of those compact operators  $x$  from  $\ell_C^2$  to  $\ell_R^2$  such that  $\text{tr } \psi(|x|/a) < \infty$  for some  $a > 0$ . If  $\psi$  is convex, then  $S^\psi$  is a Banach space for the norm given by  $\|x\|_{S^\psi} = \inf\{a > 0 : \text{tr } \psi(|x|/a) \leq 1\}$ . Otherwise,  $S^\psi$  is a Fréchet space for the F-norm given by  $\|x\|_{S^\psi} = \inf\{a > 0 : \text{tr } \psi(|x|/a) \leq a\}$  (see [26, Chapter 3].) This space may also be constructed as the noncommutative Lebesgue-Orlicz space  $L^\psi(\text{tr})$  associated to the von Neumann algebra  $\mathbb{B}(\ell_C^2, \ell_R^2)$  endowed with the normal faithful semi-finite trace  $\text{tr}$ . If  $\psi$  is the power function  $t \mapsto t^p$ , this space is denoted  $S^p$ : if  $p \geq 1$ , then  $\|x\|_{S^p} = (\text{tr } |x|^p)^{1/p}$ ; if  $p < 1$ , then  $\|x\|_{S^p} = (\text{tr } |x|^p)^{1/(1+p)}$ .

If  $\#C = \#R = n$ , then  $\mathbb{B}(\ell_C^2, \ell_R^2)$  identifies with the space of  $n \times n$  matrices denoted  $S_n^\infty$ , and we write  $S_n^\psi$  for  $S^\psi$ . Let  $(R_n \times C_n)$  be a sequence of finite sets such that each element of  $R \times C$  eventually is in  $R_n \times C_n$ . Then the sequence of operators  $P_n: x \mapsto \sum_{q \in R_n \times C_n} x_q e_q$  tends pointwise to the identity on  $S^\psi$ .

For  $I \subseteq R \times C$ , we define the space  $S_I^\psi$  as the closed subspace of  $S^\psi$  spanned by  $(e_q)_{q \in I}$ : this coincides with the subspace of those  $x \in S^\psi$  whose support is a subset of  $I$ .

A *relative Schur multiplier* on  $S_I^\psi$  is a sequence  $\varrho = (\varrho_q)_{q \in I} \in \mathbb{C}^I$  such that the associated *Schur multiplication operator*  $M_\varrho$  defined by  $e_q \mapsto \varrho_q e_q$  for  $q \in I$  is bounded on  $S_I^\psi$ . The norm  $\|\varrho\|_{M(S_I^\psi)}$  of  $\varrho$  is defined as the norm of  $M_\varrho$ . This norm is the supremum of the norm of its restrictions to finite rectangle sets  $R' \times C'$ . We used [30, 31] as a reference.

Let  $\Gamma$  be a discrete group with identity  $\epsilon$ . The *reduced C\*-algebra* of  $\Gamma$  is the closed subspace spanned by the left translations  $\lambda_\gamma$  (the linear operators defined on  $\ell_\Gamma^2$  by  $\lambda_\gamma e_\beta = e_{\gamma\beta}$ ) in  $\mathbb{B}(\ell_\Gamma^2)$ ; we denote it by  $C$ , set in roman type. The *von Neumann algebra* of  $\Gamma$  is its weak\* closure, endowed with the normal faithful normalised finite trace  $\tau$  defined by  $\tau(x) = x_{\epsilon, \epsilon}$ ; we denote it by  $L^\infty$ . Let  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be a continuous nondecreasing function vanishing only at 0: we define the noncommutative Lebesgue-Orlicz space  $L^\psi$  of  $\Gamma$  as the completion of  $L^\infty$  with respect to the norm given by  $\|x\|_{L^\psi} = \inf\{a > 0 : \tau(\psi(|x|/a)) \leq 1\}$  if  $\psi$  is convex and with respect to the F-norm given by  $\|x\|_{L^\psi} = \inf\{a > 0 : \tau(\psi(|x|/a)) \leq a\}$  otherwise. If  $\psi$  is the power function  $t \mapsto t^p$ , this space is denoted  $L^p$ : if  $p \geq 1$ , then  $\|x\|_{L^p} = \tau(|x|^p)^{1/p}$ ; if  $p < 1$ , then  $\|x\|_{L^p} = \tau(|x|^p)^{1/(1+p)}$ . The *Fourier coefficient* of  $x$  at  $\gamma$  is  $x_\gamma = \tau(\lambda_\gamma^* x) = x_{\gamma, \epsilon}$  and its *Fourier series* is  $\sum_{\gamma \in \Gamma} x_\gamma \lambda_\gamma$ . The *spectrum* of an element  $x$  is  $\{\gamma \in \Gamma : x_\gamma \neq 0\}$ . Let  $X$  be the C\*-algebra  $C$  or the space  $L^\psi$  and let  $A \subseteq \Gamma$ : then we define  $X_A$  as the closed subspace of  $X$  spanned by the  $\lambda_\gamma$  with  $\gamma \in A$ . We skip the general question for which spaces  $X$  this coincides with the subspace of those  $x \in X$  whose spectrum is a subset of  $A$ , but note that this is the case if  $\Gamma$  is an amenable group (or if  $L^\infty$  has the QWEP by [15, Theorem 4.4]) and  $\psi$  is the power function  $t \mapsto t^p$ ; note also that our definition of  $X_A$  makes it a subspace of the *heart* of  $X$ : if  $x \in X_A$ , then  $\tau(\psi(|x|/a))$  is finite for all  $a > 0$ .

A *relative Fourier multiplier* on  $X_A$  is a sequence  $\varphi = (\varphi_\gamma)_{\gamma \in A} \in \mathbb{C}^A$  such that the associated Fourier multiplication operator  $M_\varphi$  defined by  $\lambda_\gamma \mapsto \varphi_\gamma \lambda_\gamma$

for  $\gamma \in \Lambda$  is bounded on  $X_\Lambda$ . The norm  $\|\varphi\|_{M(X_\Lambda)}$  of  $\varphi$  is defined as the norm of  $M_\varphi$ . Fourier multipliers on the whole of the  $C^*$ -algebra  $C$  are also called *multipliers of the Fourier algebra*  $A(\Gamma)$  (which may be identified with  $L^1$ ); they form the set  $M(A(\Gamma))$ .

The space  $S^\psi(S^\psi)$  is the space of those compact operators  $x$  from  $\ell^2 \otimes \ell_C^2$  to  $\ell^2 \otimes \ell_R^2$  such that  $\|x\|_{S^\psi(S^\psi)} = \inf\{a : \text{tr} \otimes \text{tr} \psi(|x|/a) \leq 1\}$ : it is the noncommutative Lebesgue-Orlicz space  $L^\psi(\text{tr} \otimes \text{tr})$  associated to the von Neumann algebra  $\mathbb{B}(\ell^2) \otimes \mathbb{B}(\ell_C^2, \ell_R^2)$ . One may think of  $S^\psi(S^\psi)$  as the  $S^\psi$ -valued Schatten-von-Neumann class: we define the matrix coefficient of  $x$  at  $q$  by  $x_q = (\text{Id}_{S^\psi} \otimes \text{tr})((\text{Id}_{\ell^2} \otimes e_q^*)x) \in S^\psi$  and its matrix representation by  $\sum_{q \in R \times C} x_q \otimes e_q$ . The support of  $x$  and the subspace  $S_I^\psi(S^\psi)$  are defined in the same way as  $S_I^\psi$ .

Similarly, the space  $L^\psi(\text{tr} \otimes \tau)$  is the noncommutative Lebesgue-Orlicz space associated to the von Neumann algebra  $\mathbb{B}(\ell^2) \otimes L^\infty = L^\infty(\text{tr} \otimes \tau)$ . One may think of  $L^\psi(\text{tr} \otimes \tau)$  as the  $S^\psi$ -valued noncommutative Lebesgue space: we define the Fourier coefficient of  $x$  at  $\gamma$  by  $x_\gamma = (\text{Id}_{S^\psi} \otimes \tau)((\text{Id}_{\ell^2} \otimes \lambda_\gamma^*)x) \in S^\psi$  and its Fourier series by  $\sum_{\gamma \in \Gamma} x_\gamma \otimes \lambda_\gamma$ ; the spectrum of  $x$  is defined accordingly. The subspace  $L_\Lambda^\psi(\text{tr} \otimes \tau)$  is the closed subspace of  $L^\psi(\text{tr} \otimes \tau)$  spanned by the  $x \otimes \lambda_\gamma$  with  $x \in S^\psi$  and  $\gamma \in \Lambda$ .

An operator  $T$  on  $S_I^\psi$  is *bounded on*  $S_I^\psi(S^\psi)$  if the linear operator  $\text{Id}_{S^\psi} \otimes T$  defined by  $x \otimes y \mapsto x \otimes T(y)$  for  $x \in S^\psi$  and  $y$  in  $S_I^\psi$  on finite tensors extends to a bounded operator  $\text{Id}_{S^\psi} \otimes T$  on  $S_I^\psi(S^\psi)$ . The norm of a Schur multiplier  $\varrho$  on  $S_I^\psi(S^\psi)$  is defined as the norm of  $\text{Id}_{S^\psi} \otimes M_\varrho$ . Similar definitions hold for an operator  $T$  on  $L_\Lambda^\psi$ ; the norm of a Fourier multiplier  $\varphi$  on  $L_\Lambda^\psi(\text{tr} \otimes \tau)$  is the norm of  $\text{Id}_S^\psi \otimes M_\varphi$  on  $L_\Lambda^\psi(\text{tr} \otimes \tau)$ .

Let  $\psi$  be the power function  $t \mapsto t^p$  with  $p \geq 1$ : the norms on  $S^p(S^p)$  and  $L^p(\text{tr} \otimes \tau)$  describe the canonical operator space structure on  $S^p$  and  $L^p$ , respectively: see [30, Corollary 1.4]; we should rather use the notation  $S^p[S^p]$  and  $S^p[L^p]$ . This explains the following terminology. An operator  $T$  on  $S_I^p$  is *completely bounded* (c.b. for short) if  $\text{Id}_{S^p} \otimes T$  is bounded on  $S_I^p(S^p)$ ; the norm of  $\text{Id}_{S^p} \otimes T$  is the *complete norm* of  $T$  (compare [30, Lemma 1.7].) The complete norm  $\|\varrho\|_{M_{\text{cb}}(S_I^p)}$  of a Schur multiplier  $\varrho$  is defined as the complete norm of  $M_\varrho$ . Note that the complete norm of a Schur multiplier  $\varrho$  on  $S_I^\infty$  is equal to its norm [28, Theorem 3.2]:  $\|\varrho\|_{M_{\text{cb}}(S_I^\infty)} = \|\varrho\|_{M(S_I^\infty)}$ . The complete norm  $\|\varphi\|_{M_{\text{cb}}(L_\Lambda^p)}$  of a Fourier multiplier  $\varphi$  is defined as the complete norm of  $M_\varphi$ . The complete norm of an operator  $T$  on  $C_\Lambda$  is the norm of  $\text{Id}_{S^\infty} \otimes T$  on the subspace of  $S^\infty \otimes C$  spanned by the  $x \otimes \lambda_\gamma$  with  $x \in S^\infty$  and  $\gamma \in \Lambda$ ; in the case  $\Lambda = \Gamma$ ,  $\varphi$  is also called a *c.b. multiplier of the Fourier algebra*  $A(\Gamma)$  and one writes  $\varphi \in M_{\text{cb}}(A(\Gamma))$ ; if  $\Gamma$  is amenable, the complete norm of a Fourier multiplier  $\varphi$  on  $C_\Lambda$  is equal to its norm [7, Corollary 1.8]:  $\|\varphi\|_{M_{\text{cb}}(C_\Lambda)} = \|\varphi\|_{M(C_\Lambda)}$ .

An element whose norm is at most 1 is *contractive*, and if its complete norm is at most 1, it is *completely contractive*.

If  $\Gamma$  is abelian, let  $G$  be its dual group and endow it with its unique normalised Haar measure  $m$ : then the Fourier transform identifies the  $C^*$ -algebra  $C$  as the space of continuous functions on  $G$ ,  $L^\infty$  as the space of classes of bounded

measurable functions on  $(G, m)$ ,  $L^\psi$  as the Lebesgue-Orlicz space of classes of  $\psi$ -integrable functions on  $(G, m)$ ,  $\tau(x)$  as  $\int_G x(g) dm(g)$ ,  $L^\psi(\text{tr} \otimes \tau)$  as the  $S^\psi$ -valued Lebesgue-Orlicz space  $L^\psi(S^\psi)$  and  $x_\gamma$  as  $\hat{x}(\gamma)$ .

## 2 Transfer between Fourier and Schur multipliers

Let  $A$  be a subset of a discrete group  $\Gamma$  and let  $\varphi$  be a relative Fourier multiplier on  $C_A$ , the closed subspace spanned by  $(\lambda_\gamma)_{\gamma \in A}$  in the reduced  $C^*$ -algebra of  $\Gamma$ . Let  $x \in C_A$ : the matrix of  $x$  is *constant down the diagonals* in the sense that for every  $(r, c) \in \Gamma \times \Gamma$ ,  $x_{r,c} = x_{rc^{-1}, \epsilon} = x_{rc^{-1}}$ ; we say that  $x$  is a *Toeplitz operator* on  $\ell_\Gamma^2$ . Furthermore, the matrix of the Fourier product  $M_\varphi x$  of  $\varphi$  with  $x$  is given by  $(M_\varphi x)_{r,c} = \varphi_{rc^{-1}} x_{r,c}$ . This shows that if we set  $\check{A} = \{(r, c) \in \Gamma \times \Gamma : rc^{-1} \in A\}$  and  $\check{\varphi}_{r,c} = \varphi_{rc^{-1}}$ , then  $M_\varphi x$  is the Schur product  $M_{\check{\varphi}} x$  of  $\check{\varphi}$  with  $x$ . We have transferred the Fourier multiplier  $\varphi$  into the Schur multiplier  $\check{\varphi}$ : this shows at once that the norm of the Fourier multiplier  $\varphi$  on  $C_A$  is the norm of the Schur multiplier  $\check{\varphi}$  on the subspace of Toeplitz elements of  $\mathbb{B}(\ell_\Gamma^2)$  with support in  $\check{A}$ , and that the same holds for complete norms.

We shall now give us the means to generalise this identification to the setting of Lebesgue-Orlicz spaces  $L^\psi$ : we shall bypass the main obstacle, that  $L^\psi$  may not be considered as a subspace of  $S^\psi$ , by the Szegő limit theorem as stated by Erik Bédos [2, Theorem 10].

As we want to compute complete norms of multipliers, we shall generalise the Szegő limit theorem to the block matrix case, not considered in [2]. Let us first recall the scalar case. Consider a discrete amenable group  $\Gamma$ : it admits a *Følner averaging net of sets*  $(\Gamma_\iota)$ , that is,

- each  $\Gamma_\iota$  is a finite subset of  $\Gamma$ ;
- $\#(\gamma\Gamma_\iota\Delta\Gamma_\iota) = o(\#\Gamma_\iota)$  for each  $\gamma \in \Gamma$ .

Each set  $\Gamma_\iota$  corresponds to the orthogonal projection  $p_\iota$  of  $\ell_\Gamma^2$  onto its  $(\#\Gamma_\iota)$ -dimensional subspace of sequences supported by  $\Gamma_\iota$ . The *truncate* of a selfadjoint operator  $y \in \mathbb{B}(\ell_\Gamma^2)$  with respect to  $\Gamma_\iota$  is  $y_\iota = p_\iota y p_\iota$ : it has  $\#\Gamma_\iota$  eigenvalues  $\alpha_j$ , counted with multiplicities, and its *normalised counting measure of eigenvalues* is

$$\mu_\iota = \frac{1}{\#\Gamma_\iota} \sum_{j=1}^{\#\Gamma_\iota} \delta_{\alpha_j}.$$

If  $y$  is a Toeplitz operator, that is, if  $y \in L^\infty$ , Erik Bédos [2, Theorem 10] proves that  $(\mu_\iota)$  converges weak\* to the *spectral measure* of  $y$  with respect to  $\tau$ , which is the unique Borel probability measure  $\mu$  on  $\mathbb{R}$  such that

$$\tau(\psi(y)) = \int_{\mathbb{R}} \psi(\alpha) d\mu(\alpha)$$



for every continuous function  $\psi$  on  $\mathbb{R}$  that tends to zero at infinity. If  $\Gamma$  is abelian, then  $y$  may be identified as the class of a real-valued bounded measurable function on the group  $G$  dual to  $\Gamma$  and  $\mu$  is the distribution of  $y$ .

The matrix Szegő limit theorem is the analogue of this result for selfadjoint elements  $y \in S_n^\infty \otimes L^\infty$ , whose  $S_n^\infty$ -valued spectral measure  $\mu$  is defined by

$$\int_{\mathbb{R}} \psi(\alpha) d\mu(\alpha) = \text{Id}_{\ell_n^2} \otimes \tau(\psi(y)).$$

The orthogonal projection  $\tilde{p}_i = \text{Id}_{\ell_n^2} \otimes p_i$  defines the truncate  $y_i = \tilde{p}_i y \tilde{p}_i \in S_n^\infty \otimes \mathbb{B}(\ell_{\Gamma_i}^2)$ , and the  $S_n^\infty$ -valued normalised counting measure of eigenvalues  $\mu_i$  by

$$\int_{\mathbb{R}} \psi(\alpha) d\mu_i(\alpha) = \text{Id}_{\ell_n^2} \otimes \frac{\text{tr}}{\#\Gamma_i}(\psi(y_i)).$$

**Theorem 2.1** (Matrix Szegő limit theorem). *Let  $\Gamma$  be a discrete amenable group and let  $(\Gamma_i)$  be a Følner averaging net for  $\Gamma$ . Let  $y$  be a selfadjoint element of  $S_n^\infty \otimes L^\infty$ . The net  $(\mu_i)$  of  $S_n^\infty$ -valued normalised counting measures of eigenvalues of the truncates of  $y$  with respect to  $\Gamma_i$  converges in the weak\* topology to the spectral measure of  $y$ :*

$$\int_{\mathbb{R}} \psi(\alpha) d\mu_i(\alpha) \rightarrow \text{Id}_{\ell_n^2} \otimes \tau(\psi(y))$$

for every continuous function  $\psi$  on  $\mathbb{R}$  that tends to zero at infinity.

*Sketch of proof.* Let us first suppose that  $y \in S_n^\infty \otimes \mathbb{C}$ : we may suppose that  $y = \sum_{\gamma \in \Gamma} y_\gamma \otimes \lambda_\gamma$  with only a finite number of the  $y_\gamma \in S_n^\infty$  nonzero: the  $S_n^\infty$ -valued matrix of  $y_i$  for the canonical basis of  $\ell_{\Gamma_i}^2$  is  $(y_{rc^{-1}})_{(r,c) \in \Gamma_i \times \Gamma_i}$ . It suffices to prove that

$$\text{Id} \otimes \frac{\text{tr}}{\#\Gamma_i}(y_i^k) \rightarrow \text{Id} \otimes \tau(y^k) \tag{3}$$

for every  $k$ . This is trivial if  $k = 0$ . If  $k = 1$ , then

$$\text{Id} \otimes \frac{\text{tr}}{\#\Gamma_i}(y_i) = \frac{1}{\#\Gamma_i} \sum_{c \in \Gamma_i} y_{c,c} = \text{Id} \otimes \tau(y)$$

as  $y_{c,c} = y_{cc^{-1}} = y_\epsilon$ . If  $k \geq 2$ , the same formula holds with  $y^k$  instead of  $y$ :

$$\text{Id} \otimes \tau(y^k) = \text{Id} \otimes \frac{\text{tr}}{\#\Gamma_i}(\tilde{p}_i y^k \tilde{p}_i),$$

so that we wish to prove

$$\text{Id} \otimes \text{tr}(\tilde{p}_i y^k \tilde{p}_i - (\tilde{p}_i y \tilde{p}_i)^k) = o(\#\Gamma_i).$$

Note that

$$\|\text{Id} \otimes \text{tr}(\tilde{p}_i y^k \tilde{p}_i - (\tilde{p}_i y \tilde{p}_i)^k)\|_{S_n^1} \leq \|\tilde{p}_i y^k \tilde{p}_i - (\tilde{p}_i y \tilde{p}_i)^k\|_{S^1(S_n^1)}.$$

Lemma 5 in [2] provides the following estimate: as

$$\tilde{p}_i y^k \tilde{p}_i - (\tilde{p}_i y \tilde{p}_i)^k = \tilde{p}_i y^{k-1} (y \tilde{p}_i - \tilde{p}_i y \tilde{p}_i) + (\tilde{p}_i y^{k-1} \tilde{p}_i - (\tilde{p}_i y \tilde{p}_i)^{k-1}) y \tilde{p}_i,$$

an induction yields

$$\|\tilde{p}_i y^k \tilde{p}_i - (\tilde{p}_i y \tilde{p}_i)^k\|_{S^1(S_n^1)} \leq (k-1) \|y\|_{S_n^\infty \otimes L^\infty}^{k-1} \|y \tilde{p}_i - \tilde{p}_i y \tilde{p}_i\|_{S^1(S_n^1)}.$$

It suffices to consider the very last norm for each term  $y_\gamma \otimes \lambda_\gamma$  of  $y$ : let  $h \in \ell_n^2$  and  $\beta \in \Gamma$ ; as

$$((y_\gamma \otimes \lambda_\gamma) \tilde{p}_i - \tilde{p}_i (y_\gamma \otimes \lambda_\gamma) \tilde{p}_i)(h \otimes e_\beta) = \begin{cases} y_\gamma(h) e_{\gamma\beta} & \text{if } \beta \in \Gamma_i \text{ and } \gamma\beta \notin \Gamma_i \\ 0 & \text{otherwise,} \end{cases}$$

the definition of a Følner averaging net yields

$$\|(y_\gamma \otimes \lambda_\gamma) \tilde{p}_i - \tilde{p}_i (y_\gamma \otimes \lambda_\gamma) \tilde{p}_i\|_{S^1(S_n^1)} \leq \#(\Gamma_i \setminus \gamma^{-1} \Gamma_i) \|y_\gamma\|_{S_n^1} = o(\#\Gamma_i).$$

An approximation argument as in [2, proof of Proposition 4] permits to conclude for  $y \in L^\infty$ .  $\square$

Let us now describe and prove the  $L^\psi$  version of the transfer described at the beginning of this section.

**Lemma 2.2.** *Let  $\Gamma$  be a discrete amenable group and  $p > 0$ . Let  $\Lambda \subseteq \Gamma$  and  $\varphi \in \mathbb{C}^\Lambda$ . Consider the associated Toeplitz set  $\dot{\Lambda} = \{(r, c) \in \Gamma \times \Gamma : rc^{-1} \in \Lambda\}$  and the Toeplitz matrix defined by  $\check{\varphi}_{r,c} = \varphi_{rc^{-1}}$ . Let  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be a continuous nondecreasing function vanishing only at 0.*

- (a) *The norm of the relative Fourier multiplier  $\varphi$  on  $L_\Lambda^\psi$  is bounded by the norm of the relative Schur multiplier  $\check{\varphi}$  on  $S_\Lambda^\psi$ .*
- (b) *The norm of the relative Fourier multiplier  $\varphi$  on  $L_\Lambda^\psi(\text{tr} \otimes \tau)$  is bounded by the norm of the relative Schur multiplier  $\check{\varphi}$  on  $S_\Lambda^\psi(S^\psi)$ .*

Theorem 2.5 (a) below will provide the full picture of case (b): both norms are in fact equal. This is not so in case (a): see Remark 5.2.

*Proof.* A Toeplitz matrix has the form  $(x_{rc^{-1}})_{(r,c) \in \dot{\Lambda}}$ . Our definition of the space  $L_\Lambda^\psi$  (cf. section on Notation and terminology) ensures that we may suppose that only a finite number of the  $x_\gamma$  are nonzero for the computation of the norm of  $\varphi$ . Then  $(x_{rc^{-1}})_{(r,c) \in \dot{\Lambda}}$  is the matrix of the operator  $x = \sum_{\gamma \in \Lambda} x_\gamma \lambda_\gamma$  for the canonical basis of  $\ell_\Gamma^2$ .

Let  $y = x^* x$  and let us use the notation of Theorem 2.1. Let  $\tilde{\psi}$  be a continuous function with compact support such that  $\tilde{\psi}(t) = \psi(t)$  on  $[0, \|x\|^2]$ .

- (a). By Szegő's limit theorem (as given in [2, Theorem 1]),

$$\frac{1}{\#\Gamma_i} \text{tr} \psi(p_i y p_i) = \frac{1}{\#\Gamma_i} \text{tr} \tilde{\psi}(p_i y p_i) \rightarrow \tau(\tilde{\psi}(y)) = \tau(\psi(y))$$

Let us describe how  $\check{\varphi}$  acts on  $xp_\iota$ . Schur multiplication with  $\check{\varphi}$  transforms the matrix of  $xp_\iota$ , that is the truncated Toeplitz matrix  $(x_{rc^{-1}})_{(r,c) \in \check{A} \cap \Gamma \times \Gamma_\iota}$ , into the matrix  $(\check{\varphi}_{rc^{-1}} x_{rc^{-1}})_{(r,c) \in \check{A} \cap \Gamma \times \Gamma_\iota}$ , so that it transforms  $xp_\iota$  into  $(M_\varphi x)p_\iota$ .

(b). Combine the argument in (a) with the matrix Szegő limit theorem.  $\square$

In the case of a finite abelian group, no limit theorem is needed: this case has been considered in [22, Proposition 2.5 (b)].

The following well-known argument has been used (first in [5], see [6, Proposition D.6]) to show that the complete norm of the Fourier multiplier  $\varphi$  on  $L_A^\infty$  bounds the complete norm of the Schur multiplier  $\check{\varphi}$  on  $S_A^\infty$ , so that we have in full generality  $\|\varphi\|_{\text{Meb}(C_A)} = \|\check{\varphi}\|_{\text{Meb}(S_A^\infty)}$ . This argument permits to strengthen Lemma 2.2 (b).

**Lemma 2.3.** *Let  $\Gamma$  be a discrete group and let  $R$  and  $C$  be subsets of  $\Gamma$ . To  $A \subseteq \Gamma$  associate  $\check{A} = \{(r, c) \in R \times C : rc^{-1} \in A\}$ ; given  $\varphi \in \mathbb{C}^A$  define  $\check{\varphi} \in \mathbb{C}^{\check{A}}$  by  $\check{\varphi}_{r,c} = \varphi_{rc^{-1}}$ . Let  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be a continuous nondecreasing function vanishing only at 0. The norm of the relative Schur multiplier  $\check{\varphi}$  on  $S_A^\psi(S^\psi)$  is bounded by the norm of the relative Fourier multiplier  $\varphi$  on  $L_A^\psi(\text{tr} \otimes \tau)$ .*

*Proof.* We adapt the argument in [30, Lemma 8.1.4]. Let  $x_q \in S^\psi$ , of which only a finite number are nonzero. The space  $L^\psi(\text{tr} \otimes \text{tr} \otimes \tau)$  is a left and right  $L^\infty(\text{tr} \otimes \text{tr} \otimes \tau)$ -module and  $\sum_{\gamma \in \Gamma} e_{\gamma\gamma} \otimes \lambda_\gamma$  is a unitary in  $L^\infty(\text{tr} \otimes \tau)$ , so that

$$\begin{aligned} & \left\| \sum_{q \in \check{A}} x_q \otimes e_q \right\|_{S_A^\psi(S^\psi)} \\ &= \left\| \left( \text{Id} \otimes \sum_{r \in R} e_{r,r} \otimes \lambda_r \right) \left( \sum_{q \in \check{A}} x_q \otimes e_q \otimes \lambda_\epsilon \right) \left( \text{Id} \otimes \sum_{c \in C} e_{c,c} \otimes \lambda_c^* \right) \right\|_{L^\psi(\text{tr} \otimes \text{tr} \otimes \tau)} \\ &= \left\| \sum_{(r,c) \in \check{A}} x_{r,c} \otimes e_{r,c} \otimes \lambda_{rc^{-1}} \right\|_{L^\psi(\text{tr} \otimes \text{tr} \otimes \tau)} \\ &= \left\| \sum_{\gamma \in A} \left( \sum_{rc^{-1}=\gamma} x_{r,c} \otimes e_{r,c} \right) \otimes \lambda_\gamma \right\|_{L_A^\psi(\text{tr} \otimes \text{tr} \otimes \tau)}. \end{aligned}$$

This yields an isometric embedding of  $S_A^\psi(S^\psi)$  in  $L_A^\psi(\text{tr} \otimes \text{tr} \otimes \tau)$ . As  $S^\psi(S^\psi)$  is the Schatten-von-Neumann-Orlicz class for the Hilbert space  $\ell^2 \otimes \ell_F^2$  which may be identified with  $\ell^2$ ,

$$\begin{aligned} \left\| \sum_{q \in \check{A}} x_q \otimes \check{\varphi}_q e_q \right\|_{S_A^\psi(S^\psi)} &= \left\| \sum_{\gamma \in A} \left( \sum_{rc^{-1}=\gamma} x_{r,c} \otimes e_{r,c} \right) \otimes \varphi_\gamma \lambda_\gamma \right\|_{L_A^\psi(\text{tr} \otimes \text{tr} \otimes \tau)} \\ &\leq \| \text{Id}_{S^\psi} \otimes M_\varphi \| \left\| \sum_{q \in \check{A}} x_q \otimes e_q \right\|_{S_A^\psi(S^\psi)}. \quad \square \end{aligned}$$

*Remark 2.4.* The proof of Lemma 2.3 shows also the following transfer. Let  $(r_i)$  and  $(c_j)$  be sequences in  $\Gamma$ , consider  $\check{\Lambda} = \{(i, j) \in \mathbb{N} \times \mathbb{N} : r_i c_j \in \Lambda\}$  and define  $\check{\varphi} \in \mathbb{C}^{\check{\Lambda}}$  by  $\check{\varphi}(i, j) = \varphi(r_i c_j)$ . Then the norm of the relative Schur multiplier  $\check{\varphi}$  on  $S_{\check{\Lambda}}^{\psi}(S^{\psi})$  is bounded by the norm of the relative Fourier multiplier  $\text{Id}_{S^{\psi}} \otimes M_{\varphi}$  on  $L_{\check{\Lambda}}^{\psi}(\text{tr} \otimes \tau)$  (compare with [31, Theorem 6.4].) In particular, if the  $r_i c_j$  are pairwise distinct, this permits to transfer every Schur multiplier, not just the Toeplitz ones. See [22, Section 11] for applications of this transfer.

Here is the announced strengthening of Lemma 2.2.

**Theorem 2.5.** *Let  $\Gamma$  be a discrete amenable group. Let  $\Lambda \subseteq \Gamma$  and  $\varphi \in \mathbb{C}^{\Lambda}$ . Consider the associated Toeplitz set  $\check{\Lambda} = \{(r, c) \in \Gamma \times \Gamma : rc^{-1} \in \Lambda\}$  and the Toeplitz matrix defined by  $\check{\varphi}_{r,c} = \varphi_{rc^{-1}}$ .*

- (a) *Let  $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be a continuous nondecreasing function vanishing only at 0. The norm of the relative Fourier multiplier  $\varphi$  on  $L_{\Lambda}^{\psi}(\text{tr} \otimes \tau)$  and the norm of the relative Schur multiplier  $\check{\varphi}$  on  $S_{\check{\Lambda}}^{\psi}(S^{\psi})$  are equal.*
- (b) *Let  $p \geq 1$ . The complete norm of the relative Fourier multiplier  $\varphi$  on  $L_{\Lambda}^p$  and the complete norm of the relative Schur multiplier  $\check{\varphi}$  on  $S_{\check{\Lambda}}^p$  are equal:*

$$\|\varphi\|_{M_{\text{cb}}(L_{\Lambda}^p)} = \|\check{\varphi}\|_{M_{\text{cb}}(S_{\check{\Lambda}}^p)}.$$

- (c) *The norm of the relative Fourier multiplier  $\varphi$  on  $C_{\Lambda}$ , its complete norm, the norm of the relative Schur multiplier  $\check{\varphi}$  on  $S_{\check{\Lambda}}^{\infty}$  and its complete norm are equal:*

$$\|\varphi\|_{M(C_{\Lambda})} = \|\varphi\|_{M_{\text{cb}}(C_{\Lambda})} = \|\check{\varphi}\|_{M_{\text{cb}}(S_{\check{\Lambda}}^{\infty})} = \|\check{\varphi}\|_{M(S_{\check{\Lambda}}^{\infty})}.$$

- (d) *Suppose that  $\Lambda = \Gamma$ . The norm of the Fourier algebra multiplier  $\varphi$ , its complete norm, the norm of the Schur multiplier  $\check{\varphi}$  on  $S^{\infty}$  and its complete norm are equal:*

$$\|\varphi\|_{M(A(\Gamma))} = \|\varphi\|_{M_{\text{cb}}(A(\Gamma))} = \|\check{\varphi}\|_{M_{\text{cb}}(S^{\infty})} = \|\check{\varphi}\|_{M(S^{\infty})}.$$

*Proof.* Combine Proposition 2.2 (b) with Lemma 2.3. Recall that if  $\Gamma$  is amenable, the norm of a Fourier multiplier  $\varphi$  on  $C_{\Lambda}$  is equal to its complete norm [7, Corollary 1.8] and that the complete norm of a Schur multiplier  $\check{\varphi}$  on  $S_{\check{\Lambda}}^{\infty}$  is equal to its norm [28, Theorem 3.2].  $\square$

### 3 Local embeddings of $L^p$ into $S^p$

The proof of Lemma 2.2 can be interpreted as an embedding of  $L^{\psi}$  into an ultraproduct of finite-dimensional spaces  $S_n^{\psi}$  that intertwines Fourier and Toeplitz Schur multipliers. If we restrict ourselves to power functions  $\psi: t \mapsto t^p$  with  $p \geq 1$ , such embeddings are well known and the proof of Lemma 2.2 does not

need the full strength of the Matrix Szegő limit theorem but only the existence of such embeddings. In this section, we explain two ways to obtain them by interpolation.

The first way is to extend the classical result that the reduced C\*-algebra  $C$  of a discrete group  $\Gamma$  has the completely contractive approximation property if  $\Gamma$  is amenable. We follow the approach of [6, Theorem 2.6.8]. Let  $\Gamma$  be a discrete amenable group and  $\Gamma_\iota$  be a Følner averaging net of sets. As above, we denote by  $p_\iota$  the orthogonal projection from  $\ell_\Gamma^2$  to  $\ell_{\Gamma_\iota}^2$ . Define the compression  $\varphi_\iota$  and the embedding  $\psi_\iota$  by

$$\begin{aligned} \varphi_\iota: C &\rightarrow \mathbb{B}(\ell_{\Gamma_\iota}^2) & \text{and} & & \psi_\iota: \mathbb{B}(\ell_{\Gamma_\iota}^2) &\rightarrow C \\ x &\mapsto p_\iota x p_\iota & & & e_{r,c} &\mapsto (1/\#\Gamma_\iota)\lambda_r \lambda_{c^{-1}}. \end{aligned} \quad (4)$$

If we endow  $\mathbb{B}(\ell_{\Gamma_\iota}^2)$  with the normalised trace, these maps are unital completely positive, trace preserving (and normal) and the net  $(\psi_\iota \varphi_\iota)$  converges pointwise to the identity of  $C$ . One can therefore extend them by interpolation to completely positive contractions on the respective noncommutative Lebesgue spaces: recall that  $L^p(\mathbb{B}(\ell_{\Gamma_\iota}^2), (1/\#\Gamma_\iota)\text{tr})$  is  $(\#\Gamma_\iota)^{-1/p} S_{\#\Gamma_\iota}^p$ . We get a net of complete contractions

$$\tilde{\varphi}_\iota: L^p \rightarrow (\#\Gamma_\iota)^{-1/p} S_{\#\Gamma_\iota}^p \quad \text{and} \quad \tilde{\psi}_\iota: (\#\Gamma_\iota)^{-1/p} S_{\#\Gamma_\iota}^p \rightarrow L^p,$$

such that  $(\tilde{\psi}_\iota \tilde{\varphi}_\iota)$  converges pointwise to the identity of  $L^p$ . Moreover, the definitions (4) show that these maps also intertwine Fourier and Toeplitz Schur multipliers.

*Remark 3.1.* This approach is more canonical as it allows to extend the transfer to the vector-valued spaces in the sense of [30, Chapter 3]. Recall that for any hyperfinite semifinite von Neumann algebra  $M$  and any operator space  $E$ , one can define  $L^p(M, E)$ : for  $p = \infty$ , this space is defined as  $M \otimes_{\min} E$ ; for  $p = 1$ , this space is defined as  $M_*^{\text{op}} \hat{\otimes} E$ ; these spaces form an interpolation scale for the complex method when  $1 \leq p \leq \infty$ . For us,  $M$  will be  $\mathbb{B}(\ell^2)$  or the group von Neumann algebra  $L^\infty$ . As the maps  $\psi_\iota$  and  $\varphi_\iota$  are unital completely positive and trace preserving and normal, they define simultaneously complete contractions on  $M$  and  $M_*$ . By interpolation, the maps  $\psi_\iota \otimes \text{Id}_E$  and  $\varphi_\iota \otimes \text{Id}_E$  are still complete contractions on the spaces  $L_p(E)$  and  $S^p[E]$ . Let  $\varphi: \Gamma \rightarrow \mathbb{C}$ : the transfer shows that the norm of  $\text{Id}_E \otimes M_\varphi$  on  $L^p(E)$  is bounded by the norm of  $\text{Id}_E \otimes M_\varphi$  on  $S^p[E]$ , and that their complete norms coincide. In formulas,

$$\begin{aligned} \|\text{Id}_E \otimes M_\varphi\|_{\mathbb{B}(L^p(E))} &\leq \|\text{Id}_E \otimes M_\varphi\|_{\mathbb{B}(S^p[E])}, \\ \|\text{Id}_E \otimes M_\varphi\|_{\text{cb}(L^p(E))} &= \|\text{Id}_E \otimes M_\varphi\|_{\text{cb}(S^p[E])}. \end{aligned}$$

This approximation is two-sided whereas the proof of Lemma 2.2 uses only a one-sided approximation. This subtlety makes a difference if one tries to give a direct proof by complex interpolation, as we shall do now.

**Proposition 3.2.** *Let  $\Gamma$  be a discrete amenable group and let  $(\mu_\iota)$  be a Reiter net of means for  $\Gamma$ :*

- each  $\mu_\iota$  is a positive sequence summing to 1 with finite support  $\Gamma_\iota \subseteq \Gamma$  and viewed as a diagonal operator from  $\ell_{\Gamma_\iota}^2$  to  $\ell_\Gamma^2$ , so that

$$\|\mu_\iota\|_{S^1} = \sum_{\gamma \in \Gamma_\iota} (\mu_\iota)_\gamma = 1;$$

- the net  $(\mu_\iota)$  satisfies, for each  $\gamma \in \Gamma$ , Reiter's Property  $P_1$ :

$$\sum_{\beta \in \Gamma} |(\mu_\iota)_{\gamma^{-1}\beta} - (\mu_\iota)_\beta| \rightarrow 0. \quad (5)$$

Let  $x \in S_n^\infty \otimes L^\infty = L^\infty(\text{tr} \otimes \tau)$  and  $p \geq 1$ . Then

$$\limsup \|x\mu_\iota^{1/p}\|_{S^p(S_n^2)} = \|x\|_{L^p(\text{tr} \otimes \tau)}.$$

*Proof.* Consider  $x = \sum_{\gamma \in \Gamma} x_\gamma \otimes \lambda_\gamma$  with only a finite number of the  $x_\gamma \in S_n^\infty$  nonzero. As

$$\sum_{\beta \in \Gamma} |(\mu_\iota)_{\gamma^{-1}\beta}^{1/2} - (\mu_\iota)_\beta^{1/2}|^2 \leq \sum_{\beta \in \Gamma} |(\mu_\iota)_{\gamma^{-1}\beta} - (\mu_\iota)_\beta|,$$

Property  $P_1$  implies Property  $P_2$ :

$$\|\lambda_\gamma \mu_\iota^{1/2} - \mu_\iota^{1/2} \lambda_\gamma\|_{S^2} \rightarrow 0,$$

so that

$$\|x\mu_\iota^{1/2} - \mu_\iota^{1/2}x\|_{S^2(S_n^2)} \rightarrow 0.$$

As the  $S_n^\infty$ -valued matrix of  $x$  for the canonical basis of  $\ell_\Gamma^2$  is  $(x_{rc^{-1}})_{(r,c) \in \Gamma \times \Gamma}$ ,

$$\begin{aligned} \|x\mu_\iota^{1/2}\|_{S^2(S_n^2)}^2 &= \sum_{(r,c) \in \Gamma \times \Gamma} \|x_{rc^{-1}}\|_{S_2^n}^2 (\mu_\iota)_c \\ &= \sum_{c \in \Gamma} (\mu_\iota)_c \sum_{r \in \Gamma} \|x_{rc^{-1}}\|_{S_2^n}^2 \\ &= \sum_{c \in \Gamma} (\mu_\iota)_c \|x\|_{L^2(\text{tr} \otimes \tau)}^2 = \|x\|_{L^2(\text{tr} \otimes \tau)}^2. \end{aligned}$$

By density and continuity, the result extends to all  $x \in L^2(\text{tr} \otimes \tau)$ .

Let us prove now that for  $x \in L^\infty(\text{tr} \otimes \tau)$

$$\limsup \|x\mu_\iota\|_{S^1(S_n^1)} \leq \|x\|_{L^1(\text{tr} \otimes \tau)}.$$

The polar decomposition  $x = u|x|$  yields a factorisation  $x = ab$  with  $a = u|x|^{1/2}$  and  $b = |x|^{1/2}$  in  $L^\infty(\text{tr} \otimes \tau)$  such that

$$\begin{aligned} \|a\|_{L^2(\text{tr} \otimes \tau)} &= \|b\|_{L^2(\text{tr} \otimes \tau)} = \|x\|_{L^1(\text{tr} \otimes \tau)}^{1/2} \\ \|a\|_{L^\infty(\text{tr} \otimes \tau)} &= \|x\|_{L^\infty(\text{tr} \otimes \tau)}^{1/2}. \end{aligned}$$

Then  $x\mu_i = a(b\mu_i^{1/2} - \mu_i^{1/2}b)\mu_i^{1/2} + a\mu_i^{1/2}b\mu_i^{1/2}$ , so that the Cauchy-Schwarz inequality yields

$$\begin{aligned} \|x\mu_i\|_{S^1(S_n^1)} &\leq \|a\|_{L^\infty(\text{tr} \otimes \tau)} \|(b\mu_i^{1/2} - \mu_i^{1/2}b)\mu_i^{1/2}\|_{S^1(S_n^1)} + \|a\mu_i^{1/2}b\mu_i^{1/2}\|_{S^1(S_n^1)} \\ &\leq \|a\|_{L^\infty(\text{tr} \otimes \tau)} \|b\mu_i^{1/2} - \mu_i^{1/2}b\|_{S^2(S_n^2)} + \|a\|_{L^2(\text{tr} \otimes \tau)} \|b\|_{L^2(\text{tr} \otimes \tau)} \end{aligned}$$

and therefore our claim. Now complex interpolation yields

$$\limsup \|x\mu_i^{1/p}\|_{S^p(S_n^p)} \leq \|x\|_{L^p(\text{tr} \otimes \tau)} \quad (6)$$

for  $x \in L^\infty(\text{tr} \otimes \tau)$  and  $p \in [1, \infty]$ . Indeed, let  $u$  be the unitary appearing in the polar decomposition of  $x$ . Consider the function  $f(z) = u|x|^{pz}\mu_i^z$  analytic in the strip  $0 < \Im z < 1$  and continuous on its closure: then  $f(it)$  is a product of unitaries for  $t \in \mathbb{R}$ , so that

$$\|f(it)\|_{L^\infty(\text{tr} \otimes \tau)} = 1.$$

Also

$$\|f(1+it)\|_{S^1(S_n^1)} = \| |x|^p \mu_i \|_{S^1(S_n^1)}$$

As  $S^p(S_n^p)$  is the complex interpolation space  $(S^\infty(S_n^\infty), S^1(S_n^1))_{1/p}$ ,

$$\|x\mu_i^{1/p}\|_{S^p(S_n^p)} = \|f(1/p)\|_{S^p(S_n^p)} \leq \| |x|^p \mu_i \|_{S^1(S_n^1)}^{1/p}.$$

Then, taking the upper limit and using the estimate on  $S^1(S_n^1)$

$$\begin{aligned} \limsup \|x\mu_i^{1/p}\|_{S^p(S_n^p)} &\leq \limsup \| |x|^p \mu_i \|_{S^1(S_n^1)}^{1/p} \\ &\leq \| |x|^p \|_{L^1(\text{tr} \otimes \tau)}^{1/p} = \|x\|_{L^p(\text{tr} \otimes \tau)}. \end{aligned}$$

The reverse inequality is obtained by duality; first note that for  $y \in L^\infty(\text{tr} \otimes \tau)$ ,

$$\lim \text{tr } y\mu_i = \tau(y).$$

With the above notation and the inequality for  $p'$ ,

$$\begin{aligned} \|x\|_{L^p}^p &= \tau(|x|^p) = \lim \text{tr } |x|^p \mu_i = \lim \text{tr } \mu_i^{1-1/p} |x|^{1-p} u^* x \mu_i^{1/p} \\ &\leq \limsup \| \mu_i^{1-1/p} |x|^{1-p} \|_{S^{p'}(S_n^{p'})} \|x\mu_i^{1/p}\|_{S^p(S_n^p)} \\ &= \limsup \| |x|^{1-p} \mu_i^{1-1/p} \|_{S^{p'}(S_n^{p'})} \|x\mu_i^{1/p}\|_{S^p(S_n^p)} \\ &\leq \| |x|^{1-p} \|_{L^{p'}} \limsup \|x\mu_i^{1/p}\|_{S^p(S_n^p)} \end{aligned}$$

so that

$$\limsup \|x\mu_i^{1/p}\|_{S^p(S_n^p)} = \|x\|_{L^p(\text{tr} \otimes \tau)}^p. \quad \square$$

*Remark 3.3.* Let  $\mu$  be any positive diagonal operator with  $\text{tr } \mu = 1$  and  $p \geq 2$ : then  $\|x\mu^{1/p}\|_{S^p(S_n^p)} \leq \|x\|_{L^p}$  for all  $x \in L^\infty(\text{tr} \otimes \tau)$ . The Reiter condition is only necessary to go below exponent 2.

In the same way, using interpolation, we can come back to approximation on both sides using Reiter means, that is

$$\|\mu_t^{1/2p} x \mu_t^{1/2p}\|_{S^p(S_n^p)} \leq \|x\|_{L^p}$$

and we have

$$\limsup \|\mu_t^{1/2p} x \mu_t^{1/2p}\|_{S^p(S_n^p)} = \|x\|_{L^p(\text{tr} \otimes \tau)}.$$

Note that this formula is in the same spirit as the first approach of this section.

## 4 Transfer of lacunary sets into lacunary matrix patterns

As a first application of Theorem 2.5, let us mention that it provides a shortcut for some arguments in [12] as it permits to transfer lacunary subsets of discrete group  $\Gamma$  into lacunary matrix patterns in  $\Gamma \times \Gamma$ . Let us first introduce the following terminology.

**Definition 4.1.** Let  $\Gamma$  be a discrete group and  $\Lambda \subseteq \Gamma$ . Let  $X$  be the reduced  $C^*$ -algebra  $C$  of  $\Gamma$  or its noncommutative Lebesgue space  $L^p$  for  $p \in [1, \infty[$ .

- (a) The set  $\Lambda$  is *unconditional* in  $X$  if the Fourier series of every  $x \in X_\Lambda$  converges unconditionally: there is a constant  $D$  such that

$$\left\| \sum_{\gamma \in \Lambda'} x_\gamma \varepsilon_\gamma \lambda_\gamma \right\|_X \leq D \|x\|_X$$

for finite  $\Lambda' \subseteq \Lambda$  and  $\varepsilon_\gamma \in \mathbb{T}$ . The minimal constant  $D$  is the *unconditional constant of  $\Lambda$  in  $X$* .

- (b) If  $X = C$ , let  $\tilde{X} = S^\infty \otimes C$ ; if  $X = L^p$ , let  $\tilde{X} = L^p(\text{tr} \otimes \tau)$ . The set  $\Lambda$  is *completely unconditional* in  $X$  if the Fourier series of every  $x \in \tilde{X}_\Lambda$  converges unconditionally: there is a constant  $D$  such that

$$\left\| \sum_{\gamma \in \Lambda'} x_\gamma \otimes \varepsilon_\gamma \lambda_\gamma \right\|_{\tilde{X}} \leq D \|x\|_{\tilde{X}}$$

for finite  $\Lambda' \subseteq \Lambda$  and  $\varepsilon_\gamma \in \mathbb{T}$ . The minimal constant  $D$  is the *complete unconditional constant of  $\Lambda$  in  $X$* .

Unconditional sets in  $L^p$  have been introduced as “ $\Lambda(p)$  sets” in [12, Definition 1.1] for  $p > 2$ : if  $\Gamma$  is abelian, they are Walter Rudin’s  $\Lambda(p)$  sets if  $p > 2$  and his  $\Lambda(2)$  sets if  $p < 2$ : see [35, 3]. Asma Harcharras [12, Definition 1.5, Comments 1.9] termed completely unconditional sets in  $L^p$  “ $\Lambda(p)_{\text{cb}}$  sets” if  $p \in ]2, \infty[$ , and “ $\text{K}(p)_{\text{cb}}$  sets” if  $p \in ]1, 2]$ ; her definitions are equivalent to ours by the noncommutative Khinchin inequality.

Sets that are unconditional in  $C$  have been introduced as “unconditional Sidon sets” in [4]. If  $\Gamma$  is amenable, Fourier multipliers are automatically c.b. on



$C_A$ , so that such sets are automatically completely unconditional in  $C$ , and there are at least three more equivalent definitions for the counterpart of Sidon sets in an abelian group. If  $\Gamma$  is nonamenable, these definitions are not all equivalent anymore and our notion of completely unconditional sets in  $C$  corresponds to Marek Bożejko's "c.b. Sidon sets."

**Definition 4.2.** Let  $1 \leq p \leq \infty$  and  $I \subseteq R \times C$ .

- (a) The set  $I$  is *unconditional* in  $S^p$  if the matrix representation of every  $x \in S_I^p$  converges unconditionally: there is a constant  $D$  such that

$$\left\| \sum_{q \in I'} x_q \varepsilon_q e_q \right\|_p \leq D \|x\|_p$$

for finite  $I' \subseteq I$  and  $\varepsilon_q \in \mathbb{T}$ . The minimal constant  $D$  is the *unconditional constant of  $I$  in  $S^p$* .

- (b) The set  $I$  is *completely unconditional* in  $S^p$  if the matrix representation of every  $x \in S_I^p(S^p)$  converges unconditionally: there is a constant  $D$  such that

$$\left\| \sum_{q \in I'} x_q \otimes \varepsilon_q e_q \right\|_p \leq D \|x\|_p$$

for finite  $I' \subseteq I$  and  $\varepsilon_q \in \mathbb{T}$ . The minimal constant  $D$  is the *complete unconditional constant of  $I$  in  $S^p$* .

Harcharras [12, Definitions 4.1 and 4.4, Remarks 4.6 (iv)] termed unconditional and completely unconditional sets in  $S^p$  " $\sigma(p)$  sets" and " $\sigma(p)_{\text{cb}}$  sets," respectively; her definitions are equivalent by the noncommutative Khinchin inequality.

**Proposition 4.3.** Let  $\Gamma$  be a discrete group. Let  $\Lambda \subseteq \Gamma$  and consider the associated Toeplitz set  $\check{\Lambda} = \{(r, c) \in \Gamma \times \Gamma : rc^{-1} \in \Lambda\}$ . Let  $p \in [1, \infty[$ .

- (a) If  $\Gamma$  is amenable, then  $\Lambda$  is unconditional in  $L^p$  if  $\check{\Lambda}$  is unconditional in  $S^p$ .
- (b) If  $\Lambda$  is completely unconditional in  $L^p$ , then  $\check{\Lambda}$  is completely unconditional in  $S^p$ . The converse holds if  $\Gamma$  is amenable.

*Proof.* The first part of (b) follows by the argument of the proof of [12, Proposition 4.7]: let us sketch it. Consider the isometric embedding of the space  $S_{\check{\Lambda}}^p(S^p)$  in  $L_{\check{\Lambda}}^p(\text{tr} \otimes \text{tr} \otimes \tau)$  that is given in the proof of Lemma 2.3 and apply the equivalent Definition 1.5 in [12] of the complete unconditionality of  $\Lambda$ : this gives the complete unconditionality of  $\check{\Lambda}$  in the equivalent Definition 4.4 in [12].

Unconditionality in  $L^p$  expresses the uniform boundedness of relative unimodular Fourier multipliers on  $L_{\Lambda}^p$ ; complete unconditionality expresses their uniform complete boundedness. Unconditionality in  $S^p$  expresses the uniform boundedness of relative unimodular Schur multipliers on  $S_{\check{\Lambda}}^p$ ; complete unconditionality expresses their uniform complete boundedness. The second part of (b) follows therefore from Theorem 2.5 (b) and (a) follows from Lemma 2.2 (a).  $\square$

*Remark 4.4.* This transfer does not pass to the limit  $p = \infty$ : Nicholas Varopoulos ([36, Theorem 4.2], see [22, § 5] for a reader’s guide) proved that unconditional sets in  $S^\infty$  are finite unions of patterns whose rows or whose columns contain at most one element, and this excludes sets of the form  $\overset{\wedge}{A}$  for any infinite  $A$ .

*Remark 4.5.* See [22, Remark 11.3] for an illustration of Proposition 4.3 (b) in a particular context.

*Remark 4.6.* Let  $p$  be an even integer greater or equal to 4. The existence of a  $\sigma(p)_{\text{cb}}$  set that is not a  $\sigma(q)$  set for any  $q > p$  [12, Theorem 4.9] becomes a direct consequence of Walter Rudin’s construction [35, Theorem 4.8] of a  $\Lambda(p)$  set that is not a  $\Lambda(q)$  set for any  $q > p$ , because this set has property  $B(p/2)$  [12, Definition 2.4] and is therefore  $\Lambda(p)_{\text{cb}}$  by [12, Theorem 1.13] (in fact, it is even “1-unconditional” in  $L^p$  because  $B(p/2)$  is “ $p/2$ -independence” [22, § 11].)

*Remark 4.7.* In the same way, Theorem 5.2 in [12] becomes a mere reformulation of [12, Proposition 3.6] if one remembers that the Toeplitz Schur multipliers are 1-complemented in the Schur multipliers for an amenable discrete group and for all classical norms. Basically results on  $\Lambda(p)_{\text{cb}}$  sets produce results on  $\sigma(p)_{\text{cb}}$  sets.

Let us now estimate the complete unconditional constant of sumsets. In the case  $\Gamma = \mathbb{Z}$ , Harcharras [12, Prop. 2.8] proved that a completely unconditional set in  $L^p$  cannot contain the sumset of characters  $A + A$  for arbitrary large finite sets  $A$ : in particular, if  $\Lambda \supseteq A + A$  with  $A$  infinite, then  $\Lambda$  is not a completely unconditional set. Her proof provided thus examples of  $\Lambda(p)$  sets that are not  $\Lambda(p)_{\text{cb}}$  sets.

We generalise Harcharras’ result in two directions. Compare [18, § 1.4].

**Proposition 4.8.** *Let  $\Gamma$  be a discrete group and  $p \neq 2$ . A completely unconditional set in  $L^p$  cannot contain the sumset of two arbitrarily large sets. More precisely, let  $R$  and  $C$  be subsets of  $\Gamma$  with  $\#R \geq n$  and  $\#C \geq n^3$ . Then, for any  $p \geq 1$ , the complete unconditional constant of the sumset  $RC$  in  $L^p$  is at least  $n^{|1/2-1/p|}$ .*

*Proof.* Let  $r_1, \dots, r_n$  be pairwise distinct elements in  $R$ . We shall select inductively elements  $c_1, \dots, c_n$  in  $C$  such that the  $r_i c_j$  are pairwise distinct. Assume there are  $c_1, \dots, c_{m-1}$  such that the induction hypothesis

$$\forall i, k \leq n \forall j, l \leq m-1 \quad (i, j) \neq (k, l) \Rightarrow r_i c_j \neq r_k c_l.$$

holds. We are looking for an element  $c_m \in C$  such that

$$\forall i, k \leq n \forall l \leq m-1 \quad r_i c_m \neq r_k c_l.$$

Such an element exists as long as  $m \leq n$  because the set  $\{r_i^{-1} r_k c_l : i, k \leq n, l \leq m-1\}$  has at most  $(n(n-1)+1)(m-1) < n^3$  elements.

The end of the proof is the same as Harcharras’. The unconditional constant of the canonical basis of elementary matrices in  $S_n^p$  is  $n^{|1/2-1/p|}$ ; in particular,

there is an unimodular Schur multiplier  $\check{\varphi}$  on  $S_n^p$  of norm  $n^{|1/2-1/p|}$  (which is also its complete norm, by the way): see [30, Lemma 8.1.5]. Let  $\Lambda$  be the sumset  $\{r_i c_j : i, j \leq n\}$ ; as the  $r_i c_j$  are pairwise distinct, we may define a sequence  $\varphi \in \mathbb{C}^\Lambda$  by  $\varphi_{r_i c_j} = \check{\varphi}_{i,j}$ . By Remark 2.4, the complete norm of the Fourier multiplier  $\varphi$  on  $L_\Lambda^p$  is bounded below by the complete norm of the Schur multiplier  $\check{\varphi}$  on  $S_n^p$ .  $\square$

*Example 4.9.*  $\Lambda = \{2^i - 2^j : i > j\}$  is not a complete  $\Lambda(p)$  set for any  $p \neq 2$ . Indeed,  $\{2^i - 2^j\} = \Lambda \cup -\Lambda$  does not and if  $\Lambda$  did, then also  $-\Lambda$  and  $\Lambda \cup -\Lambda$ .

## 5 Toeplitz Schur multipliers on $S^p$ for $p < 1$

When  $0 < p < 1$ , a complete characterisation of bounded Schur multiplier of Toeplitz type has been obtained by Alexey Alexandrov and Vladimir Peller in [1, Theorem 5.1]. This result was an easy consequence of their deep results on Hankel Schur multipliers. The transfer approach provides a direct proof.

**Corollary 5.1.** *Let  $0 < p < 1$ . Let  $\Gamma$  be a discrete abelian group with dual group  $G$ . Let  $\varphi$  be a sequence indexed by  $\Gamma$  and define the associated Toeplitz matrix  $\check{\varphi} \in \mathbb{C}^\Lambda$  by  $\check{\varphi}(r, c) = \varphi(rc^{-1})$  for  $(r, c) \in \Gamma \times \Gamma$ . Then the following are equivalent:*

- (a) *The sequence  $\varphi$  is the Fourier transform of an atomic measure  $\mu = \sum a_g \delta_g$  on  $G$  with  $\sum |a_g|^p \leq 1$ ;*
- (b) *The Fourier multiplier  $\varphi$  is contractive on  $L^p$ ;*
- (c) *The Fourier multiplier  $\varphi$  is contractive on  $L^p(S^p)$ ;*
- (d) *The Schur multiplier  $\check{\varphi}$  is contractive on  $S^p$ ;*
- (e) *The Schur multiplier  $\check{\varphi}$  is contractive on  $S^p(S^p)$ .*

*Proof.* The implication (d)  $\Rightarrow$  (b) follows from Lemma 2.2(a). The equivalence (c)  $\Leftrightarrow$  (e) follows from Theorem 2.5(a). The characterisation (a)  $\Leftrightarrow$  (b) is an old result of Daniel Oberlin [23]. It is plain that (e)  $\Rightarrow$  (d). At last, (a)  $\Rightarrow$  (c) is obvious by the  $p$ -triangular inequality.  $\square$

*Remark 5.2.* As a consequence, we get that the norm of a Toeplitz Schur multiplier on  $S^p(S^p)$  coincides with its norm on  $S^p$  when  $p < 1$ . If  $p \in \{1, 2, \infty\}$ , this holds for every Schur multiplier. Let  $p \in ]1, 2[ \cup ]2, \infty[$ . Then we still do not know whether Schur multipliers are automatically c.b. on  $S^p$ . But from [30, Proposition 8.1.3], we know that (b) and (c) are not equivalent: if  $\Gamma$  is an infinite abelian group, there is a bounded Fourier multiplier on  $L^p$  that is not c.b. This example is easy to describe: if an infinite set  $A \subseteq \Gamma$  is lacunary enough, the sumset  $A + A$  is unconditional in  $L^p$  (see [18, Theorem 5.13]); by Proposition 4.8, it cannot be completely unconditional. In particular, this shows that in Lemma 2.2(a) we cannot remove the restriction to truncated Toeplitz matrices

in the computation of the Schur multiplier norm, that is, (b)  $\Rightarrow$  (d) does not hold.

*Remark 5.3.* Our questions may also be addressed in the case of a compact group: a measurable function  $\varphi$  on  $\mathbb{T}$  defines

- the Fourier multiplier on measurable functions on  $\mathbb{T}$  by  $x \mapsto \varphi x$ ;
- the Schur multiplier on integral operators on  $L^2(\mathbb{T})$  with kernel a measurable function  $x$  on  $\mathbb{T} \times \mathbb{T}$  by  $x \mapsto \check{\varphi}x$ , where  $\check{\varphi}(z, w) = \varphi(zw^{-1})$ .

Victor Olevskii [25] constructed a continuous function  $\varphi$  that defines a bounded Fourier multiplier on the space of functions with  $p$ -summable Fourier series endowed with the norm given by  $\|x\| = (\sum |\hat{x}(n)|^p)^{1/p}$  for every  $p \in ]1, \infty[$ , while the corresponding Schur multiplier is not bounded on the Schatten-von-Neumann class  $S^p$  of operators on  $L^2(\mathbb{T})$  for any  $p \in ]1, 2[ \cup ]2, \infty[$ .

## 6 The Riesz projection and the Hilbert transform

In this section, we concentrate on  $\Gamma = \mathbb{Z}$ , the dual group of  $\mathbb{T}$ .

**Proposition 6.1.** *Let  $\varrho$  be a linear combination of the identity and the upper triangular projection of  $\mathbb{N} \times \mathbb{N}$ : there are  $z, w \in \mathbb{C}$  so that  $\varrho_{i,j} = z$  if  $i \leq j$  and  $\varrho_{i,j} = w$  if  $i > j$ . Then the norm of the Schur multiplier  $\varrho$  on  $S^\psi$  coincides with the norm of the Schur multiplier  $\varrho$  on  $S^\psi(S^\psi)$ .*

*Proof.* Let  $a \in S_m^\psi(S_n^\psi)$ :  $a$  may be considered as an  $m \times m$  matrix  $(a_{ij})$  whose entries  $a_{ij}$  are  $n \times n$  matrices, and  $a$  may be identified with the block matrix

$$\tilde{a} = \begin{pmatrix} 0 & a_{11} & 0 & a_{12} & \cdots \\ 0 & 0 & 0 & 0 & \cdots \\ 0 & a_{21} & 0 & a_{22} & \cdots \\ 0 & 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

In this identification,  $\text{Id}_{S_n^\psi} \otimes M_\varrho(a)$  is  $M_\varrho(\tilde{a})$ . □

The Hilbert transform  $\mathcal{H}$  is the Schur multiplier obtained by choosing  $z = -1$  and  $w = 1$ . The upper triangular operators in  $S^p$  can be seen as a noncommutative  $H^p$  space, and  $\mathcal{H}$  corresponds exactly to the Hilbert transform in this setting (see [32, 19]). Using classical results on  $H^p$  spaces, all Hilbert transforms are c.b. for  $1 < p < \infty$  (see [37, 32, 19]).

On the torus  $\mathbb{T}$ , the classical Hilbert transform  $H$  corresponds to the Fourier multiplier given by the sign function (with the convention  $\text{sgn}(0) = 1$ ) and its norm on  $L^p$  is  $\cot(\pi/2 \max(p, p')) = \csc(\pi/p) + \cot(\pi/p)$  for  $1 < p < \infty$ . The story of the computation of this norm starts with a paper by Israel Gohberg

and Naum Krupnik [10] for  $p$  a power of 2. The remaining cases were handled by Stylianos Pichorides [29] and Brian Cole (see [8]) independently. The most achieved results are those of Brian Hollenbeck, Nigel Kalton and Igor Verbitsky [13], but they rely on complex variable methods that are not available in the operator-valued case. When  $p$  is a power of 2 (or its conjugate), a combination of arguments of Gohberg and Krupnik [9] with some of László Zsidó [37] yields the following result.

**Theorem 6.2.** *Let  $p \in ]1, \infty[$ . The norm and the complete norm of the Hilbert transform  $\mathcal{H}$  on  $S^p$  coincide with the complete norm of the Hilbert transform  $H$  on  $L^p$ : if  $\text{s\grave{g}n}(i, j) = \text{sgn}(i - j)$  for  $i, j \geq 1$ ,*

$$\|\text{s\grave{g}n}\|_{M(S^p)} = \|\text{s\grave{g}n}\|_{M_{\text{cb}}(S^p)} = \|\text{sgn}\|_{M_{\text{cb}}(L^p)}.$$

*If  $p$  is a power of 2, then these norms coincide with the norm of  $H$  on  $L^p$ :*

$$\|\text{s\grave{g}n}\|_{M(S^p)} = \|\text{s\grave{g}n}\|_{M_{\text{cb}}(S^p)} = \|\text{sgn}\|_{M_{\text{cb}}(L^p)} = \|\text{sgn}\|_{M(L^p)} = \cot(\pi/2p).$$

*Proof.* Let  $p \geq 2$ . The norm of  $H$  on  $L^p$  is  $\cot(\pi/2p)$  and the three other norms are equal by the transfer theorem 2.5 and the above proposition: we only need to compute the complete norm of  $H$ . Let  $\tilde{H} = \text{Id}_{S^p} \otimes H$  be the Hilbert transform on  $L^p(\text{tr} \otimes \tau)$ . We shall use Mischa Cotlar's trick to go from  $L^p$  to  $L^{2p}$ : the equality  $\text{sgn } i \text{sgn } j + 1 = \text{sgn}(i + j)(\text{sgn } i + \text{sgn } j)$  shows that

$$(\tilde{H}f)(\tilde{H}g) + fg = \tilde{H}((\tilde{H}f)g + f(\tilde{H}g)). \quad (7)$$

*First step.* The function  $\text{sgn}$  is not odd because of its value in 0: this can be fixed in the following way. Let  $\Lambda = 2\mathbb{Z} + 1$ . The norm of  $\tilde{H}$  on  $L^p(\text{tr} \otimes \tau)$  is equal to its norm on  $L^p_\Lambda(\text{tr} \otimes \tau)$ . In fact, let  $D$  be defined by  $Df(z) = zf(z^2)$ :  $D$  is a complete isometry on  $L^p$  with range  $L^p_\Lambda$  that commutes with  $H$ .

*Second step.* Let  $S$  be the real subspace of  $L^p_\Lambda(\text{tr} \otimes \tau)$  consisting of functions with values in  $S^p$  so that  $f(z)$  is selfadjoint for almost all  $z \in \mathbb{T}$ . Let us apply Vern Paulsen's off-diagonal trick [27, Lemma 8.1] to show that the norm of  $\tilde{H}$  on  $L^p$  is equal to its norm on  $S$ . Let  $f \in L^p_\Lambda(\text{tr} \otimes \tau)$ : identifying  $S^p_2(S^p)$  with  $S^p$ ,

$$g(z) = \begin{pmatrix} 0 & f(z) \\ f(z)^* & 0 \end{pmatrix}$$

defines an element of  $S$ . As adjoining is isometric on  $S^p$ ,

$$\|g\|_S = 2^{1/p} \|f\|_{L^p(\text{tr} \otimes \tau)}.$$

Let us now consider

$$\tilde{H}g = \begin{pmatrix} 0 & \tilde{H}f \\ \tilde{H}(f^*) & 0 \end{pmatrix}.$$

As  $0 \notin \Lambda$  by step 1, the equality  $\text{sgn}(-i) = -\text{sgn } i$  holds for  $i \in \Lambda$ : this yields that  $\tilde{H}(f^*) = -(\tilde{H}f)^*$ . Therefore

$$\|\tilde{H}g\|_S = 2^{1/p} \|\tilde{H}f\|_{L^p(\text{tr} \otimes \tau)}.$$

*Third step.* Let  $u_p$  be the norm of  $\tilde{H}$  on  $L^p(\text{tr} \otimes \tau)$ : then  $u_{2p} \leq u_p + \sqrt{1 + u_p}$ . It suffices to prove this estimate for  $f \in S$ , and by approximation we may suppose that  $f$  is a finite linear combination of terms  $a_i \otimes z^i + a_i^* \otimes z^{-i}$  with  $a_i$  finite matrices. Note that  $\tilde{H}f = -(\tilde{H}f)^*$ . Formula (7) with  $f = g$  yields, combined with Hölder's inequality,

$$\|(\tilde{H}f)^2\|_{L^p(\text{tr} \otimes \tau)} \leq \|f^2\|_{L^p(\text{tr} \otimes \tau)} + 2u_p \|f\|_{L^{2p}(\text{tr} \otimes \tau)} \|\tilde{H}f\|_{L^{2p}(\text{tr} \otimes \tau)}.$$

Since  $f$  and  $\tilde{H}f$  take normal values,

$$\begin{aligned} \|f^2\|_{L^p(\text{tr} \otimes \tau)} &= \|f\|_{L^{2p}(\text{tr} \otimes \tau)}^2 \\ \|(\tilde{H}f)^2\|_{L^p(\text{tr} \otimes \tau)} &= \|\tilde{H}f\|_{L^{2p}(\text{tr} \otimes \tau)}^2. \end{aligned}$$

Therefore, if  $\|f\|_{L^{2p}(\text{tr} \otimes \tau)} = 1$ ,  $\|\tilde{H}f\|_{L^{2p}(\text{tr} \otimes \tau)}$  must be smaller than the bigger root of  $t^2 - 2u_p t - 1$ , that is

$$\|\tilde{H}f\|_{L^{2p}(\text{tr} \otimes \tau)}^2 \leq u_p + \sqrt{u_p^2 + 1}$$

and  $u_{2p} \leq u_p + \sqrt{u_p^2 + 1}$ .

*Fourth step.* The multiplier  $H$  is an isometry on  $L^2(\text{tr} \otimes \tau)$ , so that  $u_2 = 1 = \cot(\pi/4)$ . As  $\cot(\vartheta/2) = \cot \vartheta + \sqrt{\cot^2 \vartheta + 1}$  for  $\vartheta \in ]0, \pi[$ , we conclude by an induction.  $\square$

Unfortunately, we cannot deal with other values of  $p > 2$  by this method.

The Riesz projection  $\mathcal{T}$  is the Schur multiplier obtained by choosing  $z = 0$  and  $w = 1$  in Proposition 6.1: it is the projection on the upper triangular part. On the torus, the classical Riesz projection  $T$ , that is the projection onto the analytic part, corresponds to the Fourier multiplier given by the indicator function  $\chi_{\mathbb{Z}^+}$  of nonnegative integers; its norm on  $L^p$  has been computed by Hollenbeck and Verbitsky [14]: it is  $\csc(\pi/p)$ . As for the Hilbert transform, we know that the norm and the complete norm of  $\mathcal{T}$  on  $S^p$  are equal and coincide with the complete norm of  $T$  on  $L^p$ ; but, to the best of our knowledge, there is no simple formula like (7) to go from exponent  $p$  to  $2p$ . We only obtained the following computation.

**Proposition 6.3.** *Let  $p \in ]1, \infty[$ . The norm and the complete norm of the Riesz projection  $\mathcal{T}$  on  $S^p$  coincide with the complete norm of the Riesz projection  $T$  on  $L^p$ : if  $\check{\chi}_{\mathbb{Z}^+}(i, j) = \chi_{\mathbb{Z}^+}(i - j)$  for  $i, j \geq 1$ ,*

$$\|\check{\chi}_{\mathbb{Z}^+}\|_{M(S^p)} = \|\check{\chi}_{\mathbb{Z}^+}\|_{M_{\text{cb}}(S^p)} = \|\chi_{\mathbb{Z}^+}\|_{M_{\text{cb}}(L^p)}.$$

*If  $p = 4$ , then these norms coincide with the norm of  $T$  on  $L^p$ :*

$$\|\check{\chi}_{\mathbb{Z}^+}\|_{M(S^4)} = \|\check{\chi}_{\mathbb{Z}^+}\|_{M_{\text{cb}}(S^4)} = \|\chi_{\mathbb{Z}^+}\|_{M_{\text{cb}}(L^4)} = \|\chi_{\mathbb{Z}^+}\|_{M(L^4)} = \sqrt{2}.$$

*Proof.* We shall compute the norm of  $\mathcal{T}$  on  $S^4$ . Let  $x$  be a finite upper triangular matrix and let  $y$  be a finite strictly lower triangular matrix. We have to prove that

$$\sqrt{2}\|x + y\|_{S^4} \geq \|x\|_{S^4}.$$

Let us make the obvious estimates on  $S^2$  and use the fact that adjoining is isometric:

$$\|\mathcal{T}(xx^*)\|_{S^2} = \|\mathcal{T}((x + y)x^*)\|_{S^2} \leq \|x + y\|_{S^4}\|x\|_{S^4}$$

and similarly,

$$\|(\text{Id} - \mathcal{T})(xx^*)\|_{S^2} = \|(\text{Id} - \mathcal{T})(x(x + y)^*)\|_{S^2} \leq \|x\|_{S^4}\|x + y\|_{S^4}.$$

As  $\mathcal{T}$  and  $\text{Id} - \mathcal{T}$  have orthogonal ranges,

$$\|x\|_{S^4}^4 = \|xx^*\|_{S^2}^2 = \|(\text{Id} - \mathcal{T})(xx^*)\|_{S^2}^2 + \|\mathcal{T}(xx^*)\|_{S^2}^2 \leq 2\|x\|_{S^4}^2\|x + y\|_{S^4}^2. \quad \square$$

## 7 Unconditional approximating sequences

The following definition makes sense for general operator spaces, but we chose to state it only in our specific context.

**Definition 7.1.** Let  $\Gamma$  be a discrete group and  $\Lambda \subseteq \Gamma$ . Let  $X$  be the reduced  $C^*$ -algebra of  $\Gamma$  or its noncommutative Lebesgue space  $L^p$  for  $p \in [1, \infty[$ .

- (a) A sequence  $(T_k)$  of operators on  $X_\Lambda$  is an *approximating sequence* if each  $T_k$  has finite rank and  $T_k x \rightarrow x$  for every  $x \in X_\Lambda$ . It is a *complete approximating sequence* if the  $T_k$  are uniformly c.b. If  $X$  admits a complete approximating sequence, then  $X_\Lambda$  enjoys the *c.b. approximation property*.
- (b) The *difference sequence*  $(\Delta T_k)$  of a sequence  $(T_k)$  is given by  $\Delta T_1 = T_1$  and  $\Delta T_k = T_k - T_{k-1}$  for  $k \geq 2$ . An approximating sequence  $(T_k)$  is *unconditional* if the operators

$$\sum_{k=1}^n \varepsilon_k \Delta T_k \quad \text{with } n \geq 1 \text{ and } \varepsilon_k \in \{-1, 1\} \quad (8)$$

are uniformly bounded on  $X_\Lambda$ : then  $X_\Lambda$  enjoys the *unconditional approximation property*.

- (c) An approximating sequence  $(T_k)$  is *completely unconditional* if the operators in (8) are uniformly c.b. on  $X_\Lambda$ : then  $X_\Lambda$  enjoys the *complete unconditional approximation property*. The minimal uniform bound of these operators is the *complete unconditional constant* of  $X_\Lambda$ .

We may always suppose that a complete approximating sequence on  $C_\Lambda$  is a Fourier multiplier sequence: see [11, Theorem 2.1]. We may also do so on  $L^p_\Lambda$  if  $L^\infty$  has the so-called QWEP: see [15, Theorem 4.4]. More precisely, the following proposition holds.

**Proposition 7.2.** *Let  $\Gamma$  be a discrete group and  $\Lambda \subseteq \Gamma$ . Let  $X$  either be its reduced  $C^*$ -algebra or its noncommutative Lebesgue space  $L^p$ , where  $p \in [1, \infty[$  and  $L^\infty$  has the QWEP. If  $X_\Lambda$  enjoys the completely unconditional approximation property with constant  $D$ , then, for every  $D' > D$ , there is a complete approximating sequence of Fourier multipliers  $(\varphi_k)$  that realises the completely unconditional approximation property with constant  $D'$ : the Fourier multipliers  $\sum_{k=1}^n \varepsilon_k \Delta \varphi_k$  are uniformly completely bounded by  $D'$  on  $X_\Lambda$ .*

Let us now describe how to skip blocks in an approximating sequence in order to construct an operator that acts like the Riesz projection on the sumset of two infinite sets. The following trick will be used in the induction below: compare [20, proof of Theorem 4.2]:

$$\left( \begin{array}{cc|c} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right) - \left( \begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right) + \left( \begin{array}{ccc} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{array} \right) = \left( \begin{array}{ccc} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{array} \right)$$

**Lemma 7.3.** *Let  $\Gamma$  be a discrete group and  $\Lambda \subseteq \Gamma$ . Suppose that  $\Lambda$  contains the sumset  $RC$  of two infinite sets  $R$  and  $C$ . Let  $(T_k)$  be either an approximating sequence on  $L^p_\Lambda$  with  $p \in [1, \infty[$ , or an approximating sequence of Fourier multipliers on  $C_\Lambda$ . Let  $\varepsilon > 0$ . There is a sequence  $(r_i)$  in  $R$ , a sequence  $(c_i)$  in  $C$  and there are indices  $l_1 < k_2 < l_2 < k_3 < \dots$  such that, for every  $n$ , the skipped block sum*

$$U_n = T_{l_1} + (T_{l_2} - T_{k_2}) + \dots + (T_{l_n} - T_{k_n}) \quad (9)$$

acts, up to  $\varepsilon$ , as the Riesz projection on the sumset  $\{r_i c_j\}_{i,j \leq n}$ :

$$\begin{cases} \|U_n(\lambda_{r_i c_j}) - \lambda_{r_i c_j}\| < \varepsilon & \text{if } i \leq j \leq n, \\ \|U_n(\lambda_{r_i c_j})\| < \varepsilon & \text{if } j < i \leq n. \end{cases} \quad (10)$$

*Proof.* Let us construct the sequences and indices by induction. If  $n = 1$ , let  $r_1$  and  $c_1$  be arbitrary; there is  $l_1$  such that  $\|T_{l_1}(\lambda_{r_1 c_1}) - \lambda_{r_1 c_1}\| < \varepsilon$ . Suppose that  $r_1, \dots, r_n, c_1, \dots, c_n, l_1, \dots, k_n, l_n$  have been constructed. Let  $\delta > 0$  to be chosen later.

- The operator  $U_n$  defined by Equation (9) has finite rank. If it is a Fourier multiplier, one can choose an element  $r_{n+1} \in R$  such that  $U_n(\lambda_{r_{n+1} c_j}) = 0$  for  $j \leq n$ . If it acts on  $L^p_\Lambda$  with  $p \in [1, \infty[$ , one can choose an element  $r_{n+1} \in R$  such that  $\|U_n(\lambda_{r_{n+1} c_j})\| < \delta$  for  $j \leq n$  because  $(\lambda_\gamma)_{\gamma \in \Gamma}$  is weakly null in  $L^p$ .
- There is  $k_{n+1} > l_n$  such that  $\|T_{k_{n+1}}(\lambda_\gamma) - \lambda_\gamma\| < \delta$  for  $\gamma \in \{r_i c_j : 1 \leq i \leq n+1, 1 \leq j \leq n\}$ .
- Again, choose  $c_{n+1} \in C$  such that  $\|(U_n - T_{k_{n+1}})(\lambda_{r_i c_{n+1}})\| < \delta$  for  $i \leq n+1$ .
- Again, choose  $l_{n+1} > k_{n+1}$  such that  $\|T_{l_{n+1}}(\lambda_\gamma) - \lambda_\gamma\| < \delta$  for  $\gamma \in \{r_i c_j : 1 \leq i, j \leq n+1\}$ .



Let  $U_{n+1} = U_n + (T_{l_{n+1}} - T_{k_{n+1}})$ . If  $i \leq n+1$  and  $j \leq n$ , then

$$\|\Delta U_{n+1}(\lambda_{r_i c_j})\| \leq \|T_{l_{n+1}}(\lambda_{r_i c_j}) - \lambda_{r_i c_j}\| + \|\lambda_{r_i c_j} - T_{k_{n+1}}(\lambda_{r_i c_j})\| < 2\delta,$$

so that

$$\begin{aligned} \|U_{n+1}(\lambda_{r_i c_j}) - \lambda_{r_i c_j}\| &< \varepsilon + 2\delta & \text{if } i \leq j \leq n \\ \|U_{n+1}(\lambda_{r_i c_j})\| &< \varepsilon + 2\delta & \text{if } j < i \leq n \\ \|U_{n+1}(\lambda_{r_{n+1} c_j})\| &< 3\delta & \text{if } j \leq n. \end{aligned}$$

If  $i \leq n+1$ , then

$$\begin{aligned} &\|U_{n+1}(\lambda_{r_i c_{n+1}}) - \lambda_{r_i c_{n+1}}\| \\ &\leq \|(U_n - T_{k_{n+1}})(\lambda_{r_i c_{n+1}})\| + \|T_{l_{n+1}}(\lambda_{r_i c_{n+1}}) - \lambda_{r_i c_{n+1}}\| < 2\delta. \end{aligned}$$

This shows that our choice of  $r_{n+1}, c_{n+1}, k_{n+1}$  and  $l_{n+1}$  is adequate if  $\delta$  is small enough.  $\square$

This construction will provide an obstacle to the unconditionality of sumsets.

**Theorem 7.4.** *Let  $\Gamma$  be a discrete group and  $\Lambda \subseteq \Gamma$ . Suppose that  $\Lambda$  contains the sumset  $RC$  of two infinite sets  $R$  and  $C$ .*

- (a) *Let  $1 < p < \infty$ . The complete unconditional constant of any approximating sequence for  $L^p$  is bounded below by the norm of the Riesz projection on  $S^p$ , and thus by  $\csc \pi/p$ .*
- (b) *The spaces  $L^1_\Lambda$  and  $C_\Lambda$  do not enjoy the complete unconditional approximation property.*
- (c) *If  $\Gamma$  is amenable, then the space  $C_\Lambda$  does not enjoy the unconditional approximation property.*

*Proof.* Let  $(T_k)$  be an approximating sequence on  $L^p_\Lambda$ . By Lemma 7.3, for every  $\varepsilon > 0$  and every  $n$ , there are elements  $r_1, \dots, r_n \in R$ ,  $c_1, \dots, c_n \in C$  such that the Fourier multiplier  $\varphi$  given by the indicator function of  $\{r_i c_j\}_{i \leq j}$  is near to a skipped block sum  $U_n$  of  $(T_k)$  in the sense that  $\|U_n(\lambda_{r_i c_j}) - \varphi_{r_i c_j} \lambda_{r_i c_j}\| < \varepsilon$ . But  $U_n$  is the mean of two operators of the form (8): its complete norm will provide a lower bound for the complete unconditional constant of  $X_\Lambda$ . Let us repeat the argument of Lemma 2.3 with  $x \in S^n_p$ : as

$$\begin{aligned} &\left\| \sum_{i,j=1}^n x_{i,j} e_{i,j} \right\|_{S^n_p} \\ &= \left\| \left( \sum_{i=1}^n e_{i,i} \otimes \lambda_{r_i} \right) \left( \sum_{i,j=1}^n x_{i,j} e_{i,j} \otimes \lambda_\varepsilon \right) \left( \sum_{j=1}^n e_{j,j} \otimes \lambda_{c_j} \right) \right\|_{L^p(\text{tr} \otimes \tau)} \\ &= \left\| \sum_{i=1}^n x_{i,j} e_{i,j} \otimes \lambda_{r_i c_j} \right\|_{L^p(\text{tr} \otimes \tau)} \end{aligned}$$

and

$$\left\| \sum_{i=1}^n x_{i,j} e_{i,j} \otimes (U_n(\lambda_{r_i c_j}) - \varphi_{r_i c_j} \lambda_{r_i c_j}) \right\|_{L^p(\text{tr} \otimes \tau)} < n^2 \varepsilon \|x\|_{S_n^p},$$

the complete norm of  $U_n$  is nearly bounded below by the norm of the Riesz projection on  $S_n^p$ :

$$\begin{aligned} \left\| \sum_{i=1}^n x_{i,j} e_{i,j} \otimes U_n(\lambda_{r_i c_j}) \right\|_{L^p(\text{tr} \otimes \tau)} &> \left\| \sum_{i \leq j} x_{i,j} e_{i,j} \otimes \lambda_{r_i c_j} \right\|_{L^p(\text{tr} \otimes \tau)} - n^2 \varepsilon \|x\|_{S_n^p} \\ &= \|\mathcal{T}(x)\|_{S_n^p} - n^2 \varepsilon \|x\|_{S_n^p}. \end{aligned}$$

This proves as well (a) as the first assertion in (b), because the Riesz projection is unbounded on  $S^1$ . Let  $(T_k)$  be an approximating sequence on  $C_A$ : by Lemma 7.2, we may suppose that  $(T_k)$  is a sequence of Fourier multipliers. Thus the second assertion in (b) follows from Lemma 7.3 combined with the preceding argument (where  $S_n^p$  is replaced by  $S_n^\infty$  and  $L^p(\text{tr} \otimes \tau)$  by  $S_n^\infty \otimes \mathbb{C}$ ) and the unboundedness of the Riesz projection on  $S^\infty$ . For (c), note that the Fourier multipliers  $T_k$  are automatically c.b. on  $C_A$  if  $\Gamma$  is amenable [7, Corollary 1.8].  $\square$

Theorem 7.4 (b) has been devised originally to prove that the Hardy space  $H^1$ , corresponding to the case  $A = \mathbb{N} \subseteq \mathbb{Z}$  and  $p = 1$ , admits no completely unconditional basis: see [33, 34]. Theorem 7.4 (c) both generalises the fact that a sumset cannot be a Sidon set (see [18, §§ 1.4, 6.6] for two proofs and historical remarks, or [17, Proposition IV.7]) and Daniel Li's result [16, Corollary 13] that the space  $C_A$  does not have the ‘‘metric’’ unconditional approximation property if  $\Gamma$  is abelian and  $A$  contains a sumset. Li [16, Theorem 10] also constructed a set  $A \subseteq \mathbb{Z}$  such that  $C_A$  has this property while  $A$  contains the sumset of arbitrarily large sets. This theorem also provides a new proof that the disc algebra has no unconditional basis and answers [21, Question 6.1.6].

*Example 7.5.* Neither the span of products  $\{r_i r_j\}$  of two Rademacher functions in the space of continuous functions on  $\{-1, 1\}^\infty$  nor the span of products  $\{s_i s_j\}$  of two Steinhaus functions in the space of continuous functions on  $\mathbb{T}^\infty$  have an unconditional basis.

## 8 Relative Schur multipliers of rank one

Let  $\varrho$  be an *elementary* Schur multiplier on  $S^\infty$ , that is,

$$\varrho = x \otimes y = (x_r y_c)_{(r,c) \in R \times C} :$$

then its norm is  $\sup_{r \in R} |x_r| \sup_{c \in C} |y_c|$ . How is this norm affected if  $\varrho$  is only partially specified, that is, if the action of  $\varrho$  is restricted to matrices with a given support?

**Theorem 8.1.** *Let  $I \subseteq R \times C$  and consider  $(x_r)_{r \in R}$  and  $(y_c)_{c \in C}$ . The relative Schur multiplier on  $S_I^\infty$  given by  $(x_r y_c)_{(r,c) \in I}$  has norm  $\sup_{(r,c) \in I} |x_r y_c|$ .*



each of which has norm 1 on  $S_n^\infty$  by Lemma 8.2. □

*Remark 8.3.* We learnt after submitting this article that Timur Oikhberg proved independently Theorem 8.1: see [24, Theorem 1.1].

*Remark 8.4.* As an illustration, let  $C = R = \{1, \dots, n\}$  and  $I = \{(r, c) : r \geq c\}$ , and let  $a_i$  be an increasing sequence of positive numbers. Take  $x_r = a_r$  and  $y_c = 1/a_c$ . Then the relative Schur multiplier  $(a_r/a_c)_{r \leq c}$  has norm 1. The above proof actually constructs the norm 1 extension  $(\min(a_r/a_c, a_c/a_r))_{(r,c)}$ . If we put  $a_i = e^{x_i}$ , we recover that  $(e^{-|x_r - x_c|})_{(r,c)}$  is positive definite, that is,  $|\cdot|$  is a conditionally negative function on  $\mathbb{R}$ .

## References

- [1] A. B. Aleksandrov and V. V. Peller. **Hankel and Toeplitz-Schur multipliers.** *Math. Ann.*, 324(2):277–327, 2002. (p. 18).
- [2] Erik Bédos. On Følner nets, Szegő’s theorem and other eigenvalue distribution theorems. *Exposition. Math.*, 15(3):193–228, 1997. Erratum on p. 384. (pp. 3, 7 and 9).
- [3] Jean Bourgain.  $\Lambda_p$ -sets in analysis: results, problems and related aspects. In *Handbook of the geometry of Banach spaces, Vol. I*, pages 195–232. North-Holland, Amsterdam, 2001. (p. 15).
- [4] Marek Bożejko. **A new group algebra and lacunary sets in discrete non-commutative groups.** *Studia Math.*, 70(2):165–175, 1981. (p. 15).
- [5] Marek Bożejko and Gero Fendler. Herz-Schur multipliers and completely bounded multipliers of the Fourier algebra of a locally compact group. *Boll. Un. Mat. Ital. A (6)*, 3(2):297–302, 1984. (p. 10).
- [6] Nathaniel P. Brown and Narutaka Ozawa.  *$C^*$ -algebras and finite-dimensional approximations*, volume 88 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008. (pp. 10 and 12).
- [7] Jean De Cannière and Uffe Haagerup. **Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups.** *Amer. J. Math.*, 107(2):455–500, 1985. (pp. 6, 11 and 25).
- [8] T. W. Gamelin. *Uniform algebras and Jensen measures*. Cambridge University Press, 1978. (p. 20).
- [9] Israel Gohberg and Naum Krupnik. *One-dimensional linear singular integral equations. I, II*. Birkhäuser Verlag, 1992. (p. 20).
- [10] I. Ts. Gokhberg and N. Ya. Krupnik. **Norm of the Hilbert transformation in the  $L_p$  space.** *Funct. Anal. Appl.*, 2(2):180–181, 1968. (p. 20).

- [11] Uffe Haagerup and Jon Kraus. **Approximation properties for group  $C^*$ -algebras and group von Neumann algebras.** *Trans. Amer. Math. Soc.*, 344(2):667–699, 1994. (p. 22).
- [12] Asma Harcharras. **Fourier analysis, Schur multipliers on  $S^p$  and non-commutative  $\Lambda(p)$ -sets.** *Studia Math.*, 137(3):203–260, 1999. (pp. 15, 16 and 17).
- [13] B. Hollenbeck, N. J. Kalton, and I. E. Verbitsky. **Best constants for some operators associated with the Fourier and Hilbert transforms.** *Studia Math.*, 157(3):237–278, 2003. (p. 20).
- [14] Brian Hollenbeck and Igor E. Verbitsky. **Best constants for the Riesz projection.** *J. Funct. Anal.*, 175(2):370–392, 2000. (p. 21).
- [15] Marius Junge and Zhong-Jin Ruan. **Approximation properties for non-commutative  $L_p$ -spaces associated with discrete groups.** *Duke Math. J.*, 117(2):313–341, 2003. (pp. 5 and 22).
- [16] Daniel Li. **Complex unconditional metric approximation property for  $C_\Lambda(\mathbb{T})$  spaces.** *Studia Math.*, 121(3):231–247, 1996. (p. 25).
- [17] Daniel Li and Hervé Queffélec. *Introduction à l'étude des espaces de Banach*, volume 12 of *Cours Spécialisés [Specialized Courses]*. Société Mathématique de France, Paris, 2004. Analyse et probabilités. [Analysis and probability theory]. (p. 25).
- [18] Jorge M. López and Kenneth A. Ross. *Sidon sets*. Lecture Notes Pure Appl. Math. 13. Marcel Dekker Inc., New York, 1975. (pp. 17, 18 and 25).
- [19] Michael Marsalli and Graeme West. **Noncommutative  $H^p$  spaces.** *J. Operator Theory*, 40(2):339–355, 1998. (p. 19).
- [20] F. Nazarov, G. Pisier, S. Treil, and A. Volberg. **Sharp estimates in vector Carleson imbedding theorem and for vector paraproducts.** *J. Reine Angew. Math.*, 542:147–171, 2002. (p. 23).
- [21] Stefan Neuwirth. *Multiplicateurs et analyse fonctionnelle*. PhD thesis, Université Paris 6, 1999. <http://tel.archives-ouvertes.fr/tel-00010399>. (p. 25).
- [22] Stefan Neuwirth. **Cycles and 1-unconditional matrices.** *Proc. London Math. Soc. (3)*, 93:761–790, 2006. (pp. 10, 11 and 17).
- [23] Daniel M. Oberlin. **Translation-invariant operators on  $L^p(G)$ ,  $0 < p < 1$ .** *Michigan Math. J.*, 23(2):119–122, 1976. (p. 18).
- [24] Timur Oikhberg. **Restricted Schur multipliers and their applications.** *Proc. Amer. Math. Soc.*, 138(5):1739–1750, 2010. (p. 27).

- [25] Victor Olevskii. **A connection between Fourier and Schur multipliers**. *Integral Equations Operator Theory*, 25(4):496–500, 1996. (p. 19).
- [26] Władysław Orlicz. *Linear functional analysis*, volume 4 of *Series in Real Analysis*. World Scientific Publishing Co. Inc., River Edge, NJ, 1992. Translated from the Chinese edition by Peng Yee Lee, With prefaces by Julian Musielak and Wen Qing Li, With an addendum by Cong Xin Wu. (p. 5).
- [27] Vern Paulsen. *Completely bounded maps and operator algebras*, volume 78 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002. (p. 20).
- [28] Vern I. Paulsen, Stephen C. Power, and Roger R. Smith. **Schur products and matrix completions**. *J. Funct. Anal.*, 85(1):151–178, 1989. (pp. 6 and 11).
- [29] S. K. Pichorides. **On the best values of the constants in the theorems of M. Riesz, Zygmund and Kolmogorov**. *Studia Math.*, 44:165–179. (errata insert), 1972. (p. 20).
- [30] Gilles Pisier. *Non-commutative vector valued  $L_p$ -spaces and completely  $p$ -summing maps*. Number 247 in Astérisque. Société mathématique de France, 1998. (pp. 5, 6, 10, 12 and 18).
- [31] Gilles Pisier. *Similarity problems and completely bounded maps*, volume 1618 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, expanded edition, 2001. Includes the solution to “The Halmos problem”. (pp. 5 and 11).
- [32] Narcisse Randrianantoanina. **Hilbert transform associated with finite maximal subdiagonal algebras**. *J. Austral. Math. Soc. Ser. A*, 65(3):388–404, 1998. (p. 19).
- [33] Éric Ricard. **L’espace  $H^1$  n’a pas de base complètement inconditionnelle**. *C. R. Acad. Sci. Paris Sér. I Math.*, 331(8):625–628, 2000. (p. 25).
- [34] Éric Ricard. *Décompositions de  $H^1$ , multiplicateurs de Schur et espaces d’opérateurs*. PhD thesis, Université Paris 6, 2001. [www.institut.math.jussieu.fr/theses/2001/ricard](http://www.institut.math.jussieu.fr/theses/2001/ricard). (p. 25).
- [35] Walter Rudin. **Trigonometric series with gaps**. *J. Math. Mech.*, 9:203–228, 1960. (pp. 15 and 17).
- [36] Nicholas Varopoulos. **Tensor algebras over discrete spaces**. *J. Functional Analysis*, 3:321–335, 1969. (p. 17).
- [37] László Zsidó. **On spectral subspaces associated to locally compact abelian groups of operators**. *Adv. in Math.*, 36(3):213–276, 1980. (pp. 19 and 20).

Laboratoire de Mathématiques  
Université de Franche-Comté  
16 route de Gray  
25030 Besançon Cedex, France

[stefan.neuwirth@univ-fcomte.fr](mailto:stefan.neuwirth@univ-fcomte.fr)  
[eric.ricard@univ-fcomte.fr](mailto:eric.ricard@univ-fcomte.fr)