



HAL
open science

Nitrogen and carbon isotope values of individual amino acids: a tool to study foraging ecology of penguins in the Southern Ocean

Anne Lorrain, Brittany Graham, Frédéric Ménard, Brian Popp, S. Bouillon, Peter van Breugel, Yves Cherel

► To cite this version:

Anne Lorrain, Brittany Graham, Frédéric Ménard, Brian Popp, S. Bouillon, et al.. Nitrogen and carbon isotope values of individual amino acids: a tool to study foraging ecology of penguins in the Southern Ocean. *Marine Ecology Progress Series*, 2009, 391, pp.293-306. hal-00452366

HAL Id: hal-00452366

<https://hal.science/hal-00452366>

Submitted on 2 Feb 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Nitrogen and carbon isotope values of individual amino acids: a tool to study foraging
2 ecology of penguins in the Southern Ocean

3
4 Anne Lorrain¹, Brittany Graham^{2,3}, Frédéric Ménard⁴, Brian Popp⁵, Steven Bouillon^{6,7,8}, Peter van
5 Breugel⁶, and Yves Cherel⁹

6

7

8 ¹ *IRD/LEMAR, Centre IRD de Brest, BP 70, 29280 Plouzané, France*

9 ² *Department of Oceanography, University of Hawaii, USA*

10 ³ *SINLAB, Canadian Rivers Institute, University of New Brunswick, Fredericton, New
11 Brunswick, Canada E3B 5A3*

12 ⁴ *IRD, UMR 212 EME, Centre de Recherche Halieutique (CRH), Sète Cedex, France*

13 ⁵ *Department of Geology and Geophysics, University of Hawaii, USA*

14 ⁶ *NIOO-KNAW, Yerseke, The Netherlands*

15 ⁷ *Vrije Universiteit Brussel, Dept. of Analytical and Environmental Chemistry, Brussels*

16 ⁸ *Katholieke Universiteit Leuven, Dept. of Earth & Environmental Sciences*

17 ⁹ *CEBC, UPR 1934 du CNRS, BP 14, 79360 Villiers-en-Bois, France*

18

19

20

21 Running header: Compound-specific isotope analysis and penguin isotopic niches

22

23

24

25

26 **Keywords:** $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, compound-specific, isotopic niche, trophic level

27 **ABSTRACT**

28 We determined the $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of individual amino acids (AAs) isolated
29 from chick blood of four penguin species that forage in different oceanic regions (from the
30 subtropics of the Indian Ocean to Antarctica) to test if (1) the $\delta^{15}\text{N}$ values of phenylalanine
31 ($\delta^{15}\text{N}_{\text{phe}}$) revealed different foraging areas among the species, (2) the difference between
32 glutamic acid and phenylalanine $\delta^{15}\text{N}$ values ($\Delta\delta^{15}\text{N}_{\text{glu-phe}}$) accurately predicted trophic levels
33 (TL), and (3) the $\delta^{13}\text{C}$ value of AAs could resolve species foraging locations, as bulk $\delta^{13}\text{C}$
34 values did. The $\delta^{13}\text{C}$ values of all AAs decreased with latitude, were positively correlated with
35 bulk $\delta^{13}\text{C}$ data, and therefore, tracked the isotopic baseline. However, we were not able to
36 discern additional ecological information from these $\delta^{13}\text{C}$ values. In contrast, the $\delta^{15}\text{N}$
37 analysis of individual AAs in blood distinguished the isotopic value of the nitrogen at the base
38 of the food web from the trophic level of the consumer, providing new insight for the study of
39 the trophic ecology of seabirds. The difference in the bulk $\delta^{15}\text{N}$ values of northern and
40 southern rockhopper penguins was due to both a difference in their foraging location
41 ($\neq\delta^{15}\text{N}_{\text{phe}}$) and their trophic levels ($\neq\Delta\delta^{15}\text{N}_{\text{glu-phe}}$). The $\delta^{15}\text{N}_{\text{phe}}$ values of king and Adélie
42 penguins were higher than those from rockhoppers and we hypothesize that this difference
43 reflects foraging on mesopelagic prey and in the highly productive Antarctic shelf waters
44 respectively. The $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ accurately reflected relative penguin's TL but further work is
45 required to determine the trophic enrichment factors for compound-specific isotope analysis.

INTRODUCTION

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70

Determining dietary preference together with foraging habitat of marine predators is challenging because of the extent of their pelagic environment and their long-distant movements. Traditionally, the diet of predators has been determined by stomach content, bulk stable isotope, and fatty acid analyses (Hyslop 1980; Michener & Schell 1994; Iverson et al. 2004). Foraging habitat can be investigated with tagging technologies (Wienecke et al. 2000, Charrassin & Bost 2001, Bost et al., 1997) or by linking a predator's stable isotope compositions to the isotope values of the local environment (Lee et al. 2005; Wallace et al. 2006; Cherel et al. 2006, 2007). The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of phytoplankton at the base of marine food webs can vary greatly due to different factors including phytoplankton community composition, nutrient utilization, differences in nutrient sources (e.g., denitrification vs. N_2 fixation) and the subsequent biological transformations of these nutrients (Altabet 2001; Sigman and Casciotti 2001; Karsh et al. 2003; Montoya 2007; Tamelander et al. 2009). The resulting spatial gradients in phytoplankton or zooplankton $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (e.g., inshore/offshore, pelagic/benthic, latitudinal) have been shown to propagate up to consumers and have served as proxies for foraging habitat (Best and Schell 1996). For example, bulk $\delta^{13}\text{C}$ values have been used to determine the foraging habitats of cetaceans and seabirds (Best & Schell 1996; Cherel et al. 2006, 2007; Quillfeldt et al. 2005) and bulk $\delta^{15}\text{N}$ analyses have been used to delineate temporal changes in the foraging regions of marine mammals (e.g., Burton & Koch 1999; Newsome et al. 2007). However, only a few of these studies directly compare the baseline and predator isotope values (Lee et al. 2005). Instead, the spatial variation in the isotopic baseline is inferred by knowledge of the local oceanography and from previous studies that measured proxies for the isotopic baseline (e.g., POM, zooplankton, etc.) (Cherel & Hobson 2007; Ménard et al. 2007). A spatial knowledge

71 of baseline isotope variations and an understanding of the physiology and ecology of the
72 marine predator are required for robust interpretation of the bulk $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of
73 predators. However, characterizing the isotopic baseline at the scale of ocean basins is
74 logistically challenging (Jennings & Warr 2003) and speculative for historical periods when
75 archived specimens are examined.

76 Recent evidence suggests that compound-specific isotopic analyses (CSIA) of
77 individual amino acids isolated from marine consumers could distinguish the isotopic value of
78 the nitrogen at the base of the food web from the trophic level of the consumer (McClelland &
79 Montoya 2002; Popp et al. 2007; Hannides et al. 2009). Results of the laboratory experiments
80 of McClelland and Montoya (2002) showed that the $\delta^{15}\text{N}$ value of “trophic” AAs (e.g.,
81 glutamic acid) can be enriched by as much as $\sim 7\text{‰}$ in the marine rotifer *Brachionus plicatilis*
82 relative to the $\delta^{15}\text{N}$ value in the alga *Tetraselmis suecica*, whereas other “source” AAs (e.g.,
83 phenylalanine) are little affected by trophic status and retain the $\delta^{15}\text{N}$ values of the
84 phytoplankton at the base of this food web. The implication of these results are that both
85 trophic level and the nitrogen isotopic baseline where predators foraged can be determined by
86 analyzing the $\delta^{15}\text{N}$ values of individual amino acids isolated from a predator’s tissue (see also
87 Schmidt et al. 2003; Hannides et al. 2009). However to date, CSIA of individual AAs has
88 been mainly applied to low trophic levels (McClelland & Montoya 2002; Schmidt et al. 2004;
89 Hannides et al. 2009), with only one vertebrate predator study (tuna, Popp et al. 2007), and no
90 work has yet been conducted on birds or mammals. Carbon CSIA on individual AAs has
91 mainly focused on the metabolic pathways of animals (e.g., O’Brien et al. 2005) but the
92 results of Fantle et al. (1999) on blue crabs suggested that the $\delta^{13}\text{C}$ values of individual amino
93 acids (both essential and non essential AAs) could complement bulk results to decipher a
94 consumer’s food sources.

95 In this paper, we analyzed the $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of individual amino acids isolated
96 from chick blood of four Southern Ocean penguin species: the northern rockhopper (NRP,
97 *Eudyptes chrysocome moseleyi*), southern rockhopper (SRP, *Eudyptes chrysocome*
98 *chrysocome*), king (KP, *Aptenodytes patagonicus*) and Adélie (AP, *Pygoscelis adeliae*)
99 penguins. These Southern Ocean penguins could be ideal species to test the efficacy of amino
100 acid $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ analyses to determine the ecological niches of seabirds as their food habits
101 and foraging habitats are well documented and diverse (details in Cherel & Hobson, 2007).
102 Both bulk stable isotope and stomach content analyses showed different foraging strategies
103 among these species (e.g., fish vs. crustaceans, see Table 1). Furthermore, their foraging
104 habitats differ (see Table 1 and related references) and their breeding colonies are located at
105 sites encompassing a large latitudinal range, from the subtropical Amsterdam Island north of
106 the Subtropical Front, the Crozet Island in the Polar Frontal Zone to Adélie Island, Antarctica
107 (Table 1). These regions exhibit different oceanographic characteristics (temperature,
108 chlorophyll *a* concentrations, sea ice extent) that could lead to spatial variations in the carbon
109 and nitrogen isotopic compositions at the base of the food web (Fig 1; Altabet & François
110 1994; Trull & Armand 2001). In the southwest Indian Ocean, the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of
111 particulate organic matter show an abrupt decrease between 40 and 45°S (François et al.
112 1993; Altabet & François 1994) leading to a north-south gradient across the subtropical
113 frontal zone. This latitudinal gradient can however be complicated by inshore-off shore
114 productivity gradients (Cherel and Hobson, 2007), the influence of blooms and nutrient
115 utilization (Karsh et al. 2003; Tamelander et al. 2009), the mixing of water masses across the
116 frontal zone, and the contribution of sea ice phytoplankton to the food web (Hobson et al.
117 1995; Gibson et al. 1999; Norkko et al. 2007).

118 The present study is the first to analyze carbon and nitrogen isotopes of individual
119 amino acids in seabirds, and also the first to analyze amino acids isolated from whole blood. It

120 should be noted that adult penguins can segregate their diet from the food they feed their
121 chicks (Cherel 2008). This study is limited to the chicks' diet and the adults' foraging areas
122 when they feed their chicks, but does not relate to the adult's diet. Based on our current
123 knowledge on this species, three predictions were tested:

124 1. Northern rockhopper penguin chicks (NRP) have $\delta^{15}\text{N}$ values 2.4‰ higher than
125 those of Southern Rockhopper penguin chicks (SRP, Cherel & Hobson 2007). As NRP forage
126 in the Subtropical Zone where $\delta^{15}\text{N}$ values of particulate organic matter (POM) are higher
127 than at latitudes North of the front (Altabet & François 1994), we expect this bulk isotopic
128 difference to be mainly due to isotopic baseline differences. Since phenylalanine (phe) is a
129 source amino acid and should reflect the isotopic baseline, the $\delta^{15}\text{N}_{\text{phe}}$ values in the blood of
130 NRP should be higher compared to SRP that forage within the Polar Frontal Zone (Cherel &
131 Hobson 2007).

132 2. King penguins feed heavily upon fish relative to SRP, NRP and AP, which mainly
133 prey upon crustaceans. Thus KP should have the highest trophic level (Cherel et al. 2007,
134 2008) and we therefore predict KP to have the greatest difference between source and trophic
135 amino acid $\delta^{15}\text{N}$ values.

136 3. High-latitude oceanic ecosystems (without considering onshore-offshore gradients)
137 typically have much lower POM $\delta^{13}\text{C}$ values than subtropical regions (François et al. 1993;
138 Goericke & Fry 1994) and reflect the $\delta^{13}\text{C}$ values of phytoplankton (Popp et al. 1999). The
139 bulk $\delta^{13}\text{C}$ values of these penguins decreased with increasing latitude, which was attributed to
140 the difference in the $\delta^{13}\text{C}$ values of the baseline in their respective foraging areas (Cherel &
141 Hobson, 2007). If some AAs provide information about the carbon source incorporated in
142 food webs, the $\delta^{13}\text{C}$ values of these AAs should also track spatial variations in the $\delta^{13}\text{C}$ values
143 and we expect the $\delta^{13}\text{C}$ value of some specific AAs to decrease with increasing latitude,
144 similar to the bulk $\delta^{13}\text{C}$ values.

MATERIALS AND METHODS

145

146

147 **Sample collection**

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166 **Sample preparation for compound-specific nitrogen isotope analysis**

167

168

169

170 protocols for muscle samples (e.g., Popp et al. 2007; Hannides et al. 2009). Only an overview
171 of the CSIA method is presented here and we refer the reader to Popp et al. (2007) and
172 Hannides et al. (2009) for specific details on the methods and materials. To hydrolyzed
173 samples, 4-6 mg of dried (whole) blood were transferred to high-temperature reaction vials,
174 ~1 mL 6N HCl added, heated to 150°C for 70 minutes, and cooled. These hydrolysates were
175 evaporated and the residue redissolved in 1 mL 0.01N HCl and the solution filtered (0.2 µm).
176 The solution was further purified using the cation-exchange method of Metges et al. (1996).
177 Prior to the derivatization, samples were re-acidified.

178 Amino acid derivatization included esterification of the carboxyl terminus followed by
179 trifluoroacetylation of the amine group. Samples were esterified using 4:1 isopropanol:acetyl
180 chloride and by heating at 110°C for 60 min. Samples were dried and acylated by the addition
181 of 3:1 methylene chloride: trifluoroacetic anhydride (TFAA) and heating at 100°C for 15
182 minutes. The derivatized samples were further purified using the method of Ueda et al.
183 (1989). Finally, to insure complete derivatization of the samples, the TFAA acylation step
184 was repeated. The resulting TFA derivatives were stored in 3:1 methylene
185 chloride:trifluoroacetic anhydride at 4°C.

186

187 **Compound-specific $\delta^{15}\text{N}$ Stable Isotope Analyses**

188 The $\delta^{15}\text{N}$ values of individual amino acids were analyzed by isotope ratio monitoring
189 gas chromatography-mass spectrometry (irmGCMS) using a ThermoFinnigan Delta-Plus XP
190 mass spectrometer interfaced to a Trace GC gas chromatograph through a GC-C III
191 combustion furnace (980°C), reduction furnace (680°C), and liquid nitrogen cold trap. L-2-
192 aminoadipic acid (AAA), for which the $\delta^{15}\text{N}$ value was known, was co-injected as an internal
193 reference. Samples (1-3 µl), plus the AAA reference, were injected (split/splitless, 5:1 split
194 ratio) onto a 50 m HP Ultra-2 column (0.32 mm i.d., 0.5 µm film thickness) at an injector

195 temperature of 180°C and a constant helium flow rate of 2 mL min⁻¹. The column oven was
196 initially held at 50°C for 2 min, ramped to 190°C at 8°C min⁻¹ and then to 280°C at 10°C
197 min⁻¹, and finally held at 280°C for 10 min. The irm-GCMS method allowed isotopic
198 determination of alanine, glycine, leucine, isoleucine, proline, aspartic acid, glutamic acid,
199 phenylalanine, and histidine. Samples were analyzed at least in triplicate and the measured
200 isotopic ratios were normalized to the $\delta^{15}\text{N}$ value of the amino adipic acid reference peak in
201 each chromatogram. Reproducibility associated with these isotopic measurements averaged
202 0.8‰ and ranged from 0.1 to 1.8‰. All $\delta^{15}\text{N}$ values are reported relative to AIR.

203

204 **Sample preparation for compound-specific carbon isotope analysis**

205 For $\delta^{13}\text{C}$ measurements on individual total hydrolyzable amino acids (THAA), 2.3 to
206 7.5 mg of freeze-dried blood was homogenized and hydrolyzed at 110°C in 1 mL 6M HCl in
207 screw-cap vials with a N₂ headspace. After addition of an internal standard (Norleucine), the
208 hydrolysate was evaporated under a gentle N₂ flow at 60°C. The dried THAA extracts were
209 redissolved in MQ water and stored frozen (-20°C). Prior to analyses on the HPLC-IRMS,
210 samples were centrifuged at 3000 rpm for 10 min.

211

212 **Compound-specific $\delta^{13}\text{C}$ Stable Isotope Analyses**

213 The $\delta^{13}\text{C}$ values of specific amino acids were analyzed using a modified HPLC-IRMS
214 method, based on the protocol suggested by McCullagh et al. (2006). A Surveyor HPLC was
215 coupled to a Finnigan Delta V IRMS via the LC Isolink interface (Thermo Electron, Bremen).
216 Amino acid separation was performed using a Primesep A column (3.2 X 250 mm, particle
217 size 5 μm , pore size 100Å, Sielc Technologies, Prospect Heights, IL, USA) by applying a
218 gradient program with two mobile phases (100% H₂O (Milli-Q) and 0.2 %(v/v) H₂SO₄,
219 respectively), supplied by a pump with high precision proportioning valves to control mobile

220 phase composition. Pure H₂O was used for the first 22 min, after which the mobile phase was
221 switched to linearly increase to 100% 0.2% H₂SO₄ after 75 min. The mobile phase then
222 remained at 0.2% H₂SO₄ for 40 min and switched back to 100% H₂O until the end of the run
223 (138 min). All mobile phase and reagent solutions were ultrasonically degassed under reduced
224 pressure prior to use, and stock solutions were continuously purged with He during analysis.
225 The column flow rate was kept stable at 500 $\mu\text{L min}^{-1}$ at 22°C. All samples were analyzed
226 with 10 μL partial loop injections using a 50 μL injection loop.

227 Separated amino acids eluting from the HPLC are oxidized online with a mixture of
228 0.67 M sodium peroxodisulfate (Merck, Darmstadt) and 1.5 M phosphoric acid (Fluka Sigma
229 Aldrich, Buchs) at 99.9°C. The flow of both reagents is kept at 30 $\mu\text{L min}^{-1}$. The resulting
230 CO₂ is extracted from the liquid in a phase separator with a 1 mL He flow (see Krummen et
231 al. 2004). The He containing the CO₂ from the individual amino acids is dried over a Nafion
232 tube and subsequently transferred to the IRMS through an open split.

233 To calibrate $\delta^{13}\text{C}$ values of amino acids, a mixture of individual amino acid laboratory
234 reference compounds was used. The $\delta^{13}\text{C}$ values of these compounds were determined
235 independently with an EA-IRMS using IAEA-CH-6 and an internal laboratory reference
236 compound (Schimmelmann acetanilide). The $\delta^{13}\text{C}$ value of each of these compounds was
237 previously calibrated using NBS 19 and L-SVEC on the VPDB scale where NBS-19 and L-
238 SVEC are defined as exactly +1.95 and -46.6‰, respectively (Coplen et al., 2006). Individual
239 AA calibration was required because the offset in $\delta^{13}\text{C}$ values between measurements made
240 on the HPLC-IRMS and those obtained on the EA-IRMS were different for some amino acids
241 (corrections ranged between -3.8 ‰ for glycine and +5.7 ‰ for threonine). Repeated analyses
242 of glycine over a range of concentrations (200-1000 ng C) showed excellent reproducibility,
243 with the $\delta^{13}\text{C}$ value averaging $-39.8 \pm 0.15\text{‰}$ (n=15).

244 The Primsep A column is a mixed-mode column, with negatively charged functional
245 groups due to the embedded anionic ion-pairing reagent. Amino acids with more than one
246 charge state within the pH range (e.g., aspartic acid and glutamic acid) have retention times
247 that shift in function of the mobile phase pH, which can result in co-elution of amino acid
248 peaks. In the analytical conditions used here, glutamic acid, cysteine and serine showed co-
249 elution, as well as, isoleucine, norleucine and leucine and were therefore not considered in
250 this study. Six amino acids were analyzed for both $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values (alanine, aspartic
251 acid, histidine, glycine, phenylalanine and proline).

252

253 **Statistical modeling of AA $\delta^{15}\text{N}$ values**

254 Statistical analyses must account for the different sources of variation induced by the
255 sampling strategy, and for the small number of samples analyzed. First, blood samples were
256 collected from several individuals per species (two or three individuals selected at random).
257 Secondly, several replicates were performed on each blood sample (at least three replicates,
258 but some amino acid isotope data were removed because of peak co-elution). Simple averages
259 cannot account for the within-individual variability and for the between-individual variability.
260 On the contrary, linear mixed-effects models (LME model; Pinheiro & Bates 2000) are well
261 suited to deal with unbalanced sampling schemes and they allow different sources of variation
262 to be included. In our case, we coped with replicates per blood sample, and with several
263 individuals per species. Therefore, LME models were fitted to the $\delta^{15}\text{N}$ values of individual
264 amino acids, and data were grouped by individual (measurement replicates) and by species.
265 The individual effect was treated as random variations around a population mean. The species
266 effects represent average characteristics of the populations of the four penguin species (i.e.,
267 the fixed effect in LME terms). These models allowed us to predict population values of AA
268 $\delta^{15}\text{N}$ for each penguin species. These predicted values were then used as the best estimates as

269 they account for the different sources of variation. Parameter estimation used the maximum
270 likelihood method and all computations and tests were performed in S-Plus.

271

272 **Comparison of source AA $\delta^{15}\text{N}$ values and trophic level estimates.**

273 These LME models were applied to predict population values of $\delta^{15}\text{N}_{\text{glu}}$ and $\delta^{15}\text{N}_{\text{phe}}$
274 for each species. Indeed, following Schmidt et al. (2004), we assumed (i) that phenylalanine
275 does not fractionate between trophic levels (i.e., a source amino acid), (ii) that glutamic acid
276 demonstrates a step-wise trophic enrichment (i.e., a trophic amino acid) from one trophic
277 level to the next above the primary producers, and (iii) therefore that $\Delta\delta^{15}\text{N}_{\text{glu-phe}} = \delta^{15}\text{N}_{\text{glu}} -$
278 $\delta^{15}\text{N}_{\text{phe}}$ can be considered as an index of trophic level for each penguin species. We then
279 computed $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ for each replicate and modeled these data with an extra LME model.
280 Finally, population predicted values of $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ from the LME model were used to
281 compare the relative trophic levels of each penguin species.

282 In addition, a trophic level (TL) for each penguin species can be estimated from the
283 equation of Hannides et al. (2009):

$$284 \quad TL_{\text{penguin}} = \left[\frac{(\Delta\delta^{15}\text{N}_{\text{glu-phe}})_{\text{penguin}} - (\Delta\delta^{15}\text{N}_{\text{glu-phe}})_{\text{phytoplankton}}}{TEF} \right] + 1 \quad (1),$$

285 where TEF is the trophic enrichment factor that results from a shift in one trophic level.
286 Equation 1 has three unknown variables: TL_{penguin} , $(\Delta\delta^{15}\text{N}_{\text{glu-phe}})_{\text{phytoplankton}}$, and TEF . The
287 trophic enrichment factor has been determined directly and indirectly to be $\sim 7\%$ for samples
288 of fish and crustaceans muscle tissue and whole organisms (McClelland and Montoya 2002;
289 Schmidt et al. 2004; Popp et al. 2007). In addition, $(\Delta\delta^{15}\text{N}_{\text{glu-phe}})_{\text{phytoplankton}}$ was set to 4% , i.e.
290 the value obtained by McClelland and Montoya (2002) in their lab cultures of the marine alga
291 *Tetraselmis suecica*. Although the TEF and $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ values for phytoplankton have not yet

292 been rigorously tested, they have produced reasonable TL estimates for marine zooplankton
293 (Hannides et al. 2009), krill (Schmidt et al. 2004), and yellowfin tuna (Popp et al. 2007).

294

295

RESULTS

296

297 **Patterns in $\delta^{15}\text{N}$ Amino acid values**

298 Bulk $\delta^{15}\text{N}$ values ranged 3.7‰ among all the penguin blood samples (Table 2; Cherel
299 and Hobson 2007). The $\delta^{15}\text{N}$ values of amino acids (AAs) isolated from penguin chick's
300 blood ranged from -0.2 to +26.0‰ (Table 2; Fig. 2). The trophic AAs (glutamic acid, alanine,
301 aspartic acid, isoleucine, leucine and proline; mean: $17.9 \pm 3.0\text{‰}$) were enriched in ^{15}N
302 relative to the source AAs (glycine, phenylalanine and histidine; mean: $5.1 \pm 3.5\text{‰}$) (Table 2).
303 Aspartic acid (asp) (mean: $14.7 \pm 2.1\text{‰}$) showed the least ^{15}N enrichment of the trophic AAs.
304 Among the source AAs, glycine (gly) (mean: $9.2 \pm 2.8\text{‰}$) was enriched in ^{15}N relative to
305 phenylalanine (phe) (mean: $2.5 \pm 1.2\text{‰}$) and $\delta^{15}\text{N}_{\text{gly}}$ values did not reflect the species-specific
306 patterns observed in $\delta^{15}\text{N}_{\text{phe}}$ values (Table 2; Fig. 2). Except for gly and asp, the patterns
307 observed in amino acid $\delta^{15}\text{N}$ values of penguin chicks followed previous CSIA trends in
308 trophic and source AAs measured in marine invertebrates and fish (McClelland & Montoya
309 2002; Schmidt et al. 2004; Popp et al. 2007; Hannides et al. 2009).

310 The linear mixed-effects models fitted to the $\delta^{15}\text{N}_{\text{phe}}$ and to the $\delta^{15}\text{N}_{\text{glu}}$ data indicated
311 that the species effect was significant ($p=0.022$ and $p=0.002$, respectively). The $\delta^{15}\text{N}_{\text{phe}}$ and
312 $\delta^{15}\text{N}_{\text{glu}}$ differed then by penguin species (Table 2). Adélie penguins (AP) had the highest
313 LME estimated $\delta^{15}\text{N}_{\text{phe}}$ values ($3.5 \pm 0.3\text{‰}$), whereas southern rockhopper penguins had the
314 lowest LME estimated values ($1.1 \pm 0.5\text{‰}$). Fig. 3 displays the LME values with their standard
315 errors (SE) for the four species. Northern rockhopper penguins had LME predicted $\delta^{15}\text{N}_{\text{phe}}$

316 values ($2.1 \pm 0.5\%$) higher than SRP ($1.1 \pm 0.5\%$) while king penguins (KP) had moderate
317 $\delta^{15}\text{N}_{\text{phe}}$ values ($2.6 \pm 0.5\%$).

318

319 **Trophic level of Southern Ocean penguins**

320 Table 4 displays the LME estimates for the index of trophic level $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ for the
321 four species. The species effect was significant ($p = 0.018$). Among penguin species, KP had
322 the highest index of trophic level ($\Delta\delta^{15}\text{N}_{\text{glu-phe}} = 17.2\%$, Table 3), and SRP had the lowest
323 ($\Delta\delta^{15}\text{N}_{\text{glu-phe}} = 14.1\%$). Estimates for NRP and AP were close, with a slightly higher $\Delta\delta^{15}\text{N}_{\text{glu-}}$
324 phe for NRP (15.5 vs 15.0%).

325 Using Eq. 1 with a TEF of 7‰ and a $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ in phytoplankton of 4‰, provided
326 consistent underestimates of TL (2.6, 2.4, 2.9 and 2.6 for NRP, SRP, KP and AP respectively)
327 relative to independent TL estimates based on bulk stable isotope analyses for three of the
328 four species (4.0, 4.5 and 3.9 for SRP, KP and AP, respectively; see Table 4, Cherel et al.
329 2008). To match these TL, a TEF for penguin chick's blood was calculated. This TEF
330 estimation is based on the model estimated differences ($\Delta\delta^{15}\text{N}_{\text{glu-phe}}$, see Table 4) and a
331 $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ of 4‰ for phytoplankton (McClelland & Montoya 2002). Given these
332 assumptions (these will be addressed in the Discussion), the new TEF would be 3.4, 3.8 and
333 3.8‰ for SRP, KP and AP respectively (3.6‰ on average), which is less than the 7‰ cited in
334 previous studies.

335

336 **Patterns in $\delta^{13}\text{C}$ Amino acid values**

337 While bulk $\delta^{13}\text{C}$ values ranged 5.5‰ among the penguin samples (see Table 1 and 3),
338 the $\delta^{13}\text{C}$ values of AAs isolated from penguin chick's blood ranged from -5.0 to -34.0‰
339 (Table 3; Fig. 2, 4). There was no clear pattern of ^{13}C enrichment related to essential
340 (arginine, histidine, lysine, phenylalanine, threonine) and non-essential essential AAs

341 (alanine, aspartic acid, glycine, and proline) (Table 3; Fig. 2). Instead, there were three
342 general $\delta^{13}\text{C}$ groups of amino acids, where 2 of the 3 groups included both essential and non-
343 essential essential AAs: (1) a group of AAs with high ^{13}C enrichment (threonine, glycine, and
344 histidine, mean: $-9.9 \pm 2.3\text{‰}$), (2) an intermediate group of AAs with $\delta^{13}\text{C}$ values similar to
345 bulk $\delta^{13}\text{C}$ values (proline, alanine, phenylalanine, aspartic acid, and lysine, mean: $-22.2 \pm$
346 4.3‰), and (3) a final group of only essential AAs that were very depleted in ^{13}C (arginine,
347 mean: $-31.3 \pm 1.6\text{‰}$). The bimodal pattern observed in the $\delta^{15}\text{N}$ values of source and trophic
348 AAs was not seen in the $\delta^{13}\text{C}$ values. Instead, the $\delta^{13}\text{C}$ values of all AAs decreased with
349 increasing latitude, which mirrored the bulk carbon isotope trend (Fig. 4a). A covariance
350 analysis showed that a model with separate slopes for bulk and all the AAs was justified
351 compared to a model with parallel regressions ($p = 0.006$). Slopes varied between $-0.10 \pm$
352 0.03 (lys) and -0.26 ± 0.03 (pro) with one group of four AAs having parallel slopes with bulk
353 (phe, his, arg and gly, Fig. 4a). In addition, the links between the $\delta^{13}\text{C}$ values of all AAs and
354 bulk were investigated with an extra covariance analysis: the model with separate slopes was
355 significant ($p = 0.002$; Fig 4b). Slopes varied between 0.52 ± 0.18 (thr) and 1.51 ± 0.18 (pro).
356 For six AAs (pro, ala, gly, arg, lys and phe) the correlation between bulk and AA-specific
357 $\delta^{13}\text{C}$ was highly significant ($R^2 > 0.8$ and $p < 0.01$), but there are clearly different patterns in
358 the slope of the relationship (Fig. 4b). Looking at those AAs where there is a good correlation
359 between bulk and AA-specific $\delta^{13}\text{C}$: for all non-essential AAs but asp, the slope is >1 , i.e. the
360 range in $\delta^{13}\text{C}$ -AAs is higher than in the bulk. In contrast, for all the essential ones the slope is
361 < 1 . From all AAs that have good correlation with both bulk and latitude, phenylalanine has
362 the closest values to the bulk.

363

364

365

DISCUSSION

366

367

368 **Penguin $\delta^{15}\text{N}$ values and foraging habitat (hypothesis 1)**

369 Small but significant differences were found in $\delta^{15}\text{N}_{\text{phe}}$ values among penguin species
370 (maximum range 3‰). These results suggest then that phenylalanine $\delta^{15}\text{N}$ values can be used
371 as a source AAs to study the foraging habitat of penguins. Northern Rockhopper penguins'
372 $\delta^{15}\text{N}_{\text{phe}}$ values were higher – even if the difference is relatively small - than SRP $\delta^{15}\text{N}_{\text{phe}}$
373 values ($2.1 \pm 0.5\text{‰}$ vs. $1.1 \pm 0.5\text{‰}$), which confirms the hypothesis of Cherel & Hobson (2007)
374 that the observed difference in their bulk $\delta^{15}\text{N}$ value (2.4‰) relates in part to differences in
375 the isotopic baseline of their foraging regions. The nitrogen isotopic composition of
376 particulate matter is higher in the Subtropical Frontal Zone north of 40-45°S where NRP
377 forage (Table 1, Tremblay et al. 2003) than at latitudes south of 45°S in the Southwest Indian
378 Ocean (from 5 to -2‰, see Altabet & François 1994), i.e., where southern rockhopper
379 penguins forage close to Crozet Island (Table 1). These north-south $\delta^{15}\text{N}$ gradients have also
380 been found in modern sediments collected from the NE Indian Ocean, which demonstrates
381 that these spatial gradients can be robust in the Southern Ocean (Altabet & François 1994).

382 The highest $\delta^{15}\text{N}_{\text{phe}}$ values were observed for king and Adélie penguins that forage at
383 the highest latitudes in the Southern Ocean. Previous tagging and observational data suggests
384 that these penguins forage at the Polar Front (~50°S) and over the Antarctic shelf (~66°S),
385 respectively (Wienecke et al. 2000; Charrassin & Bost 2001). Both of these oceanic regions
386 are south of the Subtropical Front where one would have expected low baseline $\delta^{15}\text{N}$ values (-
387 1 to -2‰; Altabet & François 1994; Lourey et al. 2003). However, several factors can lead to
388 elevated $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values at the base of the food web. The elevated $\delta^{15}\text{N}_{\text{phe}}$ values
389 observed in these penguins could be explained by different processes: a) a local increase in
390 the $\delta^{15}\text{N}$ value of the isotopic baseline (neritic vs. oceanic waters, high nutrient utilization by

391 phytoplankton, or sea ice influence) or b) a difference in the vertical foraging habitat, with
392 penguins foraging on a greater proportion of mesopelagic prey that have elevated $\delta^{15}\text{N}$ values.
393 A local region can have high baseline $\delta^{15}\text{N}$ value because of extensive NO_3^- utilization after a
394 large phytoplankton bloom (Tamelander et al. 2009). From Fig. 1, high chlorophyll a
395 concentrations are located near Crozet Island and Adélie Land. However, the waters close to
396 Crozet Islands are deep and well-mixed and the injection of new NO_3^- to the surface waters
397 will not produce high POM $\delta^{15}\text{N}$ values because the NO_3^- pool size is already large. On the
398 other hand, over the Antarctic shelf, the water column is stratified, and nitrogen delivery to
399 the surface waters, and the subsequent uptake by phytoplankton will lead to high $\delta^{15}\text{N}$ values
400 of the POM. Trull et al. (2008) showed a 2‰ increase in the $\delta^{15}\text{N}$ values of POM on
401 Kerguelen plateau relative to $\delta^{15}\text{N}$ values of POM collected off the plateau. The higher $\delta^{15}\text{N}$
402 values of POM were attributed to an increase in the uptake of NO_3^- by phytoplankton on the
403 Kerguelen plateau. Neritic waters found above plateaus or coastal shelves can then have
404 higher $\delta^{15}\text{N}$ values at the base of the food web compared to offshore oceanic regions.
405 Accordingly, the isotopic baseline of the waters directly surrounding the Antarctic shelf may
406 be higher than offshore waters. Adélie penguins forage within 50 km of their Antarctic
407 colonies (Cherel et al. 2008). Therefore, the relatively high $\delta^{15}\text{N}_{\text{phe}}$ values observed in Adélie
408 penguins could reflect their neritic foraging behavior in waters above the Antarctic shelf.
409 Finally, the high $\delta^{15}\text{N}$ values of Adélie penguins could be explained by feeding in a food web
410 supported in part by sea ice phytoplankton, which has been shown to have elevated $\delta^{13}\text{C}$ and
411 $\delta^{15}\text{N}$ values (Hobson et al., 1995; Norkko et al. 2007).

412 The high $\delta^{15}\text{N}_{\text{phe}}$ values of king penguins that forage in the polar front could be
413 explained by the relative importance of mesopelagic prey in their diet (prey leaving around
414 300-500m). Previous studies have suggested there is a positive depth gradient in the $\delta^{15}\text{N}$
415 value of particulate nitrogen in the open ocean (Saino & Hattori 1980, 1987; Dore et al. 2002,

416 Trull et al. 2008). Saino & Hattori (1980) found an overall increase of 9‰ in POM $\delta^{15}\text{N}$
417 between 0 and 1000m in the NE Indian Ocean and Trull et al. (2008) found +0.5‰ with depth
418 on Kerguelen plateau up to 140m. If this nitrogen is incorporated and conserved in the
419 foodweb, then prey that forage at depth (below 100-200m) should have higher $\delta^{15}\text{N}$ values
420 that similar prey in the surface waters (Rau et al. 1989; Graham et al. 2007). Mintenbeck et al.
421 (2007) showed a significant increase in the $\delta^{15}\text{N}$ values of benthic suspension feeders with
422 water depth on the Weddell Sea shelf (up to 1000 m). Thus, if predators forage on a greater
423 proportion of mesopelagic prey, their $\delta^{15}\text{N}_{\text{phe}}$ values would be higher than those of consumers
424 feeding in the same region, but on a more epipelagic resource. King penguins make deep
425 dives to forage, regularly exceeding 150 m depth (Kooyman et al. 1992), and feeding almost
426 exclusively on mesopelagic fish of the family Myctophidae, while other penguins (including
427 AP, NRP, SRP) dive to shallower depths and mainly prey upon crustaceans (Cherel et al.
428 1993, 2007, Rodary et al. 2000; Tremblay & Cherel 2003, Cherel 2008).

429 The $\delta^{15}\text{N}_{\text{phe}}$ values of SRP and NRP suggest that they do not forage in the same
430 oceanic regions and that the difference in their bulk $\delta^{15}\text{N}$ values is due in part to baseline
431 differences. These results also revealed that northern rockhoppers, king and Adélie penguins
432 have close $\delta^{15}\text{N}$ values, which would have remained unclear without knowledge on their
433 ecology and feeding behavior (i.e., inshore vs. offshore feeding, deep feeding). Phenylalanine
434 can then be used as a source AA that records baseline $\delta^{15}\text{N}$ variations in penguins, but we
435 recommend analyzing a larger number of individuals and interpreting the $\delta^{15}\text{N}_{\text{phe}}$ data with
436 additional ecological information on the studied species.

437

438 **Penguin $\delta^{15}\text{N}$ values and trophic levels (hypothesis 2)**

439 Results of previous stomach content and bulk stable isotope analyses suggest that king
440 penguins have a higher trophic level than southern rockhoppers and Adélie penguins (Cherel

441 et al. 2008). The compound-specific isotope data supports these observations, as the
442 difference between the $\delta^{15}\text{N}$ values of trophic (glutamic acid) and source (phenylalanine)
443 amino acids was greatest in king penguins (17.2‰, Table 4). If bulk isotope $\delta^{15}\text{N}$ values
444 (Table 1) are interpreted only in the context of variations in trophic ecology, NRP (9.2‰) was
445 at a lower trophic level than AP (10.1‰), and SRP (6.8‰) was at the lowest TL. The amino
446 acid $\delta^{15}\text{N}$ data however indicated that NRP had a higher $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ value, or trophic level,
447 than both SRP (15.5 vs. 14.1‰) and AP (15.5 vs. 15.0‰). These conclusions are consistent
448 with stomach content analysis, which indicate that NRP fed on squids and crustaceans,
449 whereas, SRP and AP forage mostly on crustaceans (Table 1). The bulk $\delta^{15}\text{N}$ difference
450 (2.4‰) between NRP and SRP is therefore not only due to baseline difference as discussed
451 previously, but also to a difference in their trophic level.

452 Our study thus suggests that the $\delta^{15}\text{N}$ analyses of individual amino acids, such as
453 glutamic acid and phenylalanine, can provide an opportunity to distinguish the relative
454 influence of baseline variations and trophic level on the bulk $\delta^{15}\text{N}$ values of penguins.
455 However, using equation 1 and a trophic enrichment factor of 7‰ (cf. McClelland &
456 Montoya 2002), the trophic level of penguin chicks ranged from 2.6 to 2.9. A trophic level
457 lower than 3 is not possible for these penguins as they are strictly carnivorous (Cherel et al.
458 1993, 2008). For example, king penguin is a myctophid-eater, and myctophids forage mainly
459 on meso- and macrozooplankton, including some herbivorous, omnivorous and even
460 carnivorous species. Consequently, a TL of king penguins cannot be lower than 4. To match
461 the expected TLs for penguins, and considering that the 4‰ phytoplankton $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ is
462 correct, a new TEF of 3.6‰ (see results) has to be used. Assuming this $\text{TEF}_{\text{blood}}$ of 3.6‰ and
463 a value of 4‰ for the $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ value for phytoplankton, this study produced the first
464 estimate of TL for Northern Rockhopper penguin chicks (4.2) and we find a TL for SRP, KP
465 and AP to be 3.8, 4.6 and 4.0 respectively. However, the $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ value for phytoplankton

466 is not well constrained, and it could also explain why the results from equation 1 consistently
467 under-estimated the TL for these penguins. Although our dataset suggests that the 7‰ trophic
468 enrichment factor is not correct for blood tissue collected from penguins, uncertainty still
469 exists in this emerging ecological method (e.g., TL estimates from Cherel et al. 2008,
470 $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ value for phytoplankton) and trophic enrichment factors should be determined
471 with experimental work conducted on a diverse assemblage of consumers.

472 To conclude, the $\delta^{15}\text{N}$ values of glutamic acid and phenylalanine in penguin blood
473 ($\Delta\delta^{15}\text{N}_{\text{glu-phe}}$) successfully estimated the relative trophic level of the different species of
474 penguins. However before absolute TLs can be calculated, controlled experiments should be
475 performed on seabirds and their diet to better constrain the $\text{TEF}_{\text{blood}}$ and the $\Delta\delta^{15}\text{N}_{\text{glu-phe}}$ value
476 for phytoplankton.

477

478 **Penguin $\delta^{13}\text{C}$ values and foraging habitat (hypothesis 3)**

479 Variations in bulk $\delta^{13}\text{C}$ values have been interpreted as differences in the foraging
480 habitats of the four penguin species and to corresponding spatial differences in the $\delta^{13}\text{C}$ values
481 at the base of the food web (Cherel & Hobson 2007). Strong spatial gradients have been
482 observed in the Southern Ocean, with a ~9‰ decrease in the $\delta^{13}\text{C}$ values of POM from low to
483 high latitudes (François et al. 1993; Popp et al. 1999; Trull et al. 2001). The laboratory and
484 field results of Fantle et al. (1999) showed that the essential amino acids (E AAs) valine,
485 leucine and phenylalanine did not exhibit significant ^{13}C enrichment from the diet and had
486 lower $\delta^{13}\text{C}$ values than the non-essential amino acids (NE AAs). Based upon these
487 observations we predicted that essential AAs would mirror the latitudinal bulk isotopic trends
488 exhibited in Southern Ocean phytoplankton, i.e., the $\delta^{13}\text{C}$ values of E AAs in penguins that
489 forage in lower latitudes would be higher than those that forage at higher latitudes.

490 The $\delta^{13}\text{C}$ values of 6 AAs (phe, lys, arg, gly, pro, ala) had very good correlation with
491 bulk $\delta^{13}\text{C}$ values ($R^2 > 0.8$) and decreased with increasing latitude, suggesting that these AAs
492 track $\delta^{13}\text{C}$ baseline variations. In contrast to what has been found in blue crabs (Fantle et al.
493 1999), the $\delta^{13}\text{C}$ values of E AAs do not segregate relative to NE AAs in penguin's blood.
494 Both E AAs and NE AAs had higher $\delta^{13}\text{C}$ values than bulk $\delta^{13}\text{C}$ values. Interestingly, all E
495 AAs exhibited lower slopes relative to bulk $\delta^{13}\text{C}$ values meaning that the range of variation in
496 these specific AAs $\delta^{13}\text{C}$ values was lower relative to bulk $\delta^{13}\text{C}$ values, while NE AAs (except
497 for aspartic acid) had higher ranges. We are unable to interpret this pattern with our current
498 understanding of carbon isotope fractionation of specific amino acids in seabirds. As such,
499 applying carbon CSIA to determine the foraging ecology and location of marine consumers is
500 not straightforward, and may not even be applicable. Without baseline or prey $\delta^{13}\text{C}$ data, it is
501 not possible to determine if some E AAs fractionate or not relative to their diet. In an
502 experimental study conducted on fish, McCullagh et al. (2008) also found that there was no
503 clear pattern in ^{13}C fractionation relative to essentiality of AAs. Instead these authors found
504 that only phenylalanine showed no isotopic difference between the $\delta^{13}\text{C}$ value of the
505 consumer and its diet. In our study, phenylalanine had $\delta^{13}\text{C}$ values close to the bulk values for
506 all penguin species, which suggest that it may be the most appropriate amino acid to track
507 changes in the baseline $\delta^{13}\text{C}$ values and determine a marine consumer's foraging habitat. If
508 one specific AA had to be chosen for simultaneous C and N isotope analysis, we propose
509 phenylalanine which has the closest values relative to bulk, and is also a source AA for
510 nitrogen.

511

512

513

CONCLUSION

514

515 Few studies have been conducted on carbon and nitrogen isotope analyses of
516 individual amino acids, and none of them examined seabirds. This study shows for the first
517 time that the $\delta^{15}\text{N}$ values of individual amino acids, such as glutamic acid as a trophic AA and
518 phenylalanine as a source AA, can be used to study the foraging ecology of penguins. These
519 results further support the use of compound-specific $\delta^{15}\text{N}$ isotope analysis to determine the
520 foraging areas and trophic levels of marine consumers, from primary consumers to top
521 predators. Previous analyses however focused on muscle and whole body while this study
522 examined blood. Our results suggest that blood can be used to estimate relative trophic levels,
523 but that the trophic enrichment factor reported in previous studies (i.e., 7‰) might not be
524 appropriate to calculate absolute TL in penguins, and, in general, seabirds. Controlled
525 experiments are therefore needed to better constrain the $\text{TEF}_{\text{blood}}$ value for penguins. The $\delta^{13}\text{C}$
526 values of 6 individual AAs tracked $\delta^{13}\text{C}$ isotopic baseline but without additional CSIA data on
527 the diet or base of the food web, we were unable to further interpret the $\delta^{13}\text{C}$ values of
528 specific amino acids isolated from penguin blood. Our study however suggests that glutamic
529 acid ($\delta^{15}\text{N}$) and phenylalanine (for both $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) could be key individual amino acids to
530 study the foraging habitat and behavior of marine consumers.

531

532 **Acknowledgements**

533 The authors thank T. Rust and C. Hannides for their help in $\delta^{15}\text{N}$ analyses and J.
534 Middelburg, and B. Veuger for having made possible $\delta^{13}\text{C}$ analyses on the penguin samples.
535 We also greatly thank S. Sokolov for providing Indian Ocean fronts' positions and Chl a data
536 from Fig. 1, and M. Patriat and P Lopez for map and figure processing. This work was
537 supported financially by the Program REMIGE – ANR Biodiversité 2005-011 and partially
538 by the Netherlands Organisation for Scientific Research for carbon analyses.

539

540 **References**

- 541 Altabet MA, François R (1994) Sedimentary nitrogen isotopic ratio as a recorder for surface
542 ocean nitrate utilization. *Glob Biogeochem Cycles* 8:103-116
- 543 Best PB, Schell DM (1996) Stable isotopes in southern right whale (*Eubalaena australis*)
544 baleen as indicators of seasonal movements, feeding and growth. *Mar Biol* 124:483-494
- 545 Bost CA, Georges JY, Guinet C, Cherel Y, Pütz K, Charrassin JB, Handrich Y, Zorn T, Lage
546 J, Le Maho Y (1997) Foraging habitat and food intake of satellite-tracked king penguins
547 during the austral summer at Crozet Archipelago. *Mar Ecol Prog Ser* 150: 21-33
- 548 Bugoni L, McGill RAR, Furness RW (2008) Effects of preservation methods on stable
549 isotope signatures in bird tissues. *Rapid Commun Mass Spectrom* 22: 2457-2462
- 550 Burton RK, Koch PL (1999) Isotopic tracking of foraging and long-distance migration in
551 northeastern Pacific pinnipeds, *Oecologia* 119: 578–585
- 552 Charrassin JB, Bost CA (2001) Utilisation of oceanic habitat by king penguins over the
553 annual cycle. *Mar Ecol Prog Ser* 221:285-297
- 554 Cherel Y, Verdon C, Ridoux V (1993) Seasonal importance of oceanic myctophids in king
555 penguin diet at Crozet Islands. *Polar Biol* 13:355-357
- 556 Cherel Y, Hobson KA, Hassani S (2005) Isotopic discrimination between food and blood and
557 feathers of captive penguins: implications for dietary studies in the wild. *Physiol Biochem*
558 *Zool* 78:106-115
- 559 Cherel Y, Phillips RA, Hobson KA, McGill R (2006) Stable isotope evidence of diverse
560 species-specific and individual wintering strategies in seabirds. *Biol Lett* 2:301-303
- 561 Cherel Y, Hobson KA (2007) Geographical variation in carbon stable isotope signatures of
562 marine predators: a tool to investigate their foraging areas in the Southern Ocean. *Mar*
563 *Ecol Prog Ser* 329:281-287

564 Cherel Y, Hobson KA, Guinet C, Vanpé C (2007) Stable isotopes document seasonal changes
565 in trophic niches and winter foraging individual specialisation in diving predators from the
566 Southern Ocean. *J Anim Ecol* 76, 826-836

567 Cherel Y (2008) Isotopic niches of emperor and Adélie penguins in Adélie Land, Antarctica.
568 *Mar Biol* 154:813-821

569 Cherel Y, Ducatez S, Fontaine C, Richard P, Guinet C (2008) Stable isotopes reveal the
570 trophic position and mesopelagic fish diet of female southern elephant seals breeding on
571 the Kerguelen Islands. *Mar Ecol Prog Ser* 370:239-247

572 Coplen TB, Brand WA, Gehre M, Gröning M, Meijer HAJ, Toman B, Verkouteren RM
573 (2006): New guidelines for $\delta^{13}\text{C}$ measurements. *Anal Chem* 78: 2439-2441

574 Deniro MJ, Epstein S (1981) Influence of diet on the distribution of nitrogen isotopes in
575 animals. *Geochim Cosmochim Acta* 45:341-351

576 Dore JE, Brum JR, Tupas LM, Karl DM (2002) Seasonal and interannual variability in sources
577 of nitrogen supporting export in the oligotrophic subtropical North Pacific Ocean. *Limnol*
578 *Oceanogr* 47:1595-1607

579 Fantle MS, Dittel AI, Schwalm SM, Epifanio CE, Fogel ML (1999) A food web analysis of
580 the juvenile blue crab, *Callinectes sapidus*, using stable isotopes in whole animals and
581 individual amino acids. *Oecologia* 120:416-426

582 François R, Altabet MA, Goericke R (1993) Changes in the $\delta^{13}\text{C}$ of surface water particulate
583 matter across the Subtropical Convergence in the SW Indian Ocean. *Global Biogeochem*
584 *Cycles* 7:627-644

585 Fry B (1988) Food web structure on Georges Bank from stable C, N, and S isotopic
586 compositions. *Limnol Oceanogr* 33:1182-1190

587 Gibson JAE, Trull TW, Nichols PD, Summons RE, McMinn A (1999) Sedimentation of ^{13}C -
588 rich organic matter from Antarctic sea-ice algae: a potential indicator of past sea-ice
589 extent. *Geology* 27: 331-334

590 Goericke R, Fry B (1994) Variations of marine plankton $\delta^{13}\text{C}$ with latitude, temperature, and
591 dissolved CO_2 in the world ocean. *Global Biogeochem Cycles* 8:85-90

592 Gouretski VV, Koltermann KP (2004) WOCE Global Hydrographic Climatology. A
593 Technical Report 35, Berichte des BSH

594 Graham BS, Grubbs D, Holland K, Popp BN (2007) A rapid ontogenetic shift in the diet of
595 juvenile yellowfin tuna from Hawaii. *Mar Biol* 150:647-658

596 Hannides, CCS, Popp BN, Landry MR, Graham BS (2009) Quantification of zooplankton
597 trophic position in the North Pacific Subtropical Gyre using stable nitrogen isotopes.
598 *Limnol and Oceanogr* 54(1) 50-61

599 Hobson KA, Clark RG (1992) Assessing avian diets using stable isotopes II: Factors
600 influencing diet-tissue fractionation. *The condor* 94: 189-197

601 Hobson KA, Ambrose WG Jr, Renaud PE (1995) Sources of primary production, benthic-
602 pelagic coupling, and trophic relationships within the Northeast Water Polynya: insights
603 from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis. *Mar Ecol Prog Ser* 128: 1-10

604 Hobson KA, Gibbs HL, Gloutney ML (1997) Preservation of blood and tissue samples for
605 stable-carbon and stable nitrogen isotope analysis. *Can J Zool* 75:1720–1723

606 Hyslop EJ (1980) Stomach content analysis. A review of methods and their application. *J Fish*
607 *Biol* 17:411-429

608 Iverson SJ, Field C, Bowen WD, Blanchard W (2004) Quantitative fatty acid signature
609 analysis: a new method of estimating predator diets. *Ecological Monographs* 74:211-235

610 Jennings S, Warr KJ (2003) Environmental correlates of large-scale spatial variation in the
611 $\delta^{15}\text{N}$ of marine animals. *Mar Biol* 142:1131-1140

612 Karsh KL, Trull TW, Lourey MJ, Sigman DM (2003) Relationship of nitrogen isotope
613 fractionation to phytoplankton size and iron availability during the Southern Ocean Iron
614 RElease Experiment (SOIREE). *Limnol and Oceanogr* 48: 1058-1068

615 Kooyman GL, Cherel Y, Le Maho Y, Croxall JP, Thorson PH, Ridoux V, Kooyman CA
616 (1992) Diving behavior and energetics during foraging cycles in King Penguins. *Ecol*
617 *Monogr* 62:143-163

618 Krummen M, Hilkert AW, Juchelka D, Duhr A, Schluter HJ, Pesch R (2004) A new concept
619 for isotope ratio monitoring liquid chromatography/mass spectrometry. *Rapid Commun*
620 *Mass Spectrom* 18: 2260-2266

621 Lee SH, Schell DM, McDonald TL, Richardson WJ (2005) Regional and seasonal feeding by
622 bowhead whales *Balaena mysticetus* as indicated by stable isotope ratios. *Mar Ecol Prog*
623 *Ser* 285:271-285

624 Lourey MJ, Trull TW, Sigman DM (2003) Sensitivity of $\delta^{15}\text{N}$ of nitrate, surface suspended
625 and deep sinking particulate nitrogen to seasonal nitrate depletion in the Southern Ocean.
626 *Global Biogeoche Global Biogeochem Cycles* 17: 1081, doi:10.1029/2002GB001973

627 McClelland JW, Montoya JP (2002) Trophic relationships and the nitrogen isotopic
628 composition of amino acids in plankton. *Ecology* 83:2173-2180

629 McCullagh JSO, Juchelka D, Hedges REM (2006) Analysis of amino acid ^{13}C abundance
630 from human and faunal bone collagen using liquid chromatography isotope ratio mass
631 spectrometry. *Rapid Commun Mass Spectrom* 20: 2761-2768

632 McCullagh JSO, Gaye-Siessegger J, Focken U (2008) Determination of underivatized amino
633 acid $\delta^{13}\text{C}$ by liquid chromatography/isotope ratio mass spectrometry for nutritional
634 studies: the effect of dietary non-essential amino acid profile on the isotopic signature of
635 individual amino acids in fish. *Rapid Commun Mass Spectrom* 22: 1817-1822

636 Ménéard F, Lorrain A, Potier M, Marsac F (2007) Isotopic evidence of distinct feeding
637 ecologies and movement patterns in two migratory predators (yellowfin tuna and
638 swordfish) of the western Indian Ocean. *Mar Biol* 153: 141-152

639 Metges CC, Petzke K, Hennig U (1996) Gas chromatography/combustion/isotope ratio mass
640 spectrometric comparison of N-acetyl- and N-pivaloyl amino acid esters to measure ^{15}N
641 isotopic abundances in physiological samples: a pilot study on amino acid synthesis in the
642 upper gastrointestinal tract of minipigs. *J Mass Spectrom* 31:367-376

643 Michener RH, Schell DM (1994) Stable isotope ratios as tracers in marine aquatic food webs.
644 In: Lajtha, K. and Michener, R.H., Editors, 1994. *Stable isotopes in ecology and*
645 *environmental sciences*, Blackwell Scientific Publications, Oxford, pp. 138–157

646 Mintenbeck K, Jacob U, Knust R, Arntz WE, Brey T (2007) Depth-dependence in stable
647 isotope ratio $\delta^{15}\text{N}$ of benthic POM consumers: The role of particle dynamics and organism
648 trophic guild. *Deep-Sea Res I* 54: 1015-1023

649 Newsome SD, Martinez del Rio C, Bearhop S, Phillips DL (2007) A niche for isotopic
650 ecology. *Frontiers Ecol Environ* 5(8):429-436

651 Norkko A, Trish SF, Cummings VJ, Gibbs MM, Andrew NL, Norkko J, Schwarz AM (2007)
652 Trophic structure of coastal Antarctic food webs associated with changes in sea ice and
653 food supply. *Ecology* 88: 2810–2820

654 O'Brien DM, Boggs CL, Fogel ML (2005) The amino acids used in reproduction by
655 butterflies: a comparative study of dietary sources using compound specific stable isotope
656 analysis. *Physiological and Biochemical Zoology* 78(5):819-827

657 Pinheiro JC, Bates DM (2000) *Mixed-effects models in S and SPLUS*. Springer, New York

658 Popp BN, Trull T, Kenig F, Wakeham SG, Rust TM, Tilbrook B, Griffiths FB, Wright SW,
659 Marchant HJ, Bidigare RR, Laws EA (1999) Controls on the carbon isotopic compositions
660 of Southern Ocean phytoplankton. *Global Biogeochem Cycles* 13: 827–844

661 Popp BN, Graham BS, Olson RJ, Hannides CCS, Lott MJ, Lopez-Ibarra GA, Galván-Magaña
662 F, Fry B (2007) Insight into the trophic ecology of yellowfin tuna, *Thunnus albacares*,
663 from compound-specific nitrogen isotope analysis of proteinaceous amino acids. In
664 Dawson T. and Siegwolf R. (eds) Stable Isotopes as Indicators of Ecological Change,
665 Elsevier Academic Press pp. 173-190

666 Quillfeldt P, McGill RAR, Furness RW (2005) Diet and foraging areas of Southern Ocean
667 seabirds and their prey inferred from stable isotopes: review and case study of Wilson's
668 storm-petrel. Mar Ecol Prog Ser 295:295–304

669 Rau GH, Heyraud M, Cherry RD (1989) $^{15}\text{N}/^{14}\text{N}$ and $^{13}\text{C}/^{12}\text{C}$ in mesopelagic shrimp from the
670 northeast Atlantic Ocean: evidence for differences in diet. Deep Sea Res 36:1103-1110

671 Rodary D, Wienecke BC, Bost CA (2000) Diving behaviour of Adélie penguins (*Pygoscelis*
672 *adeliae*) at Dumont D'Urville, Antarctica: nocturnal patterns of diving and rapid
673 adaptations to changes in sea-ice condition. Polar Biol 23:113-120

674 Saino T, Hattori A (1980) ^{15}N natural abundance in oceanic suspended particulate matter.
675 Nature 28: 752-754

676 Saino T, Hattori A (1987) Geographical variation of the water column distribution of
677 suspended particulate organic nitrogen and its ^{15}N natural abundance in the Pacific and its
678 marginal seas. Deep-Sea Res 34:807-827

679 Schmidt K, McClelland JW, Mente E, Montoya JP, Atkinson A, Voss M (2004) Trophic-level
680 interpretation based on $\delta^{15}\text{N}$ values: implications of tissue-specific fractionation and
681 amino acid composition. Mar Ecol Prog Ser 266:43-58

682 Sears J, Hatch SA, O'Brien DM (2009) Disentangling effects of growth and nutritional status
683 on seabird stable isotope ratios. Oecologia 159:41-48

684 Sokolov S, Rintoul SR (2002) Structure of Southern Ocean fronts at 140°E. J Mar Sys 37:
685 151-184

686 Sokolov S, Rintoul SR (2007) On the relationship between fronts of the Antarctic

687 Circumpolar Current and surface chlorophyll concentrations in the Southern Ocean. J
688 Geophys Res 112: C07030, doi:10.1029/2006JC004072

689 Sokolov S, Rintoul SR (2009) The circulation structure and distribution of the Antarctic
690 Circumpolar Current fronts. Part 1: Mean circumpolar paths. J Geophys Res in press

691 Tamelander T, Kivimä C, Bellerby RGJ, Renaud PE, Kristiansen S (2009) Base-line
692 variations in stable isotope values in an Arctic marine ecosystem: effects of carbon
693 and nitrogen uptake by phytoplankton. Hydrobiol DOI: 10.1007/s10750-009-9780-2.

694 Tremblay Y, Cherel Y (2003) Geographic variation in the foraging behavior, diet and chick
695 growth of rockhopper penguins. Mar Ecol Prog Ser 251: 279-297

696 Trull TW, Armand L (2001) Insights into Southern Ocean carbon export from the $\delta^{13}\text{C}$ of
697 particles and dissolved inorganic carbon during the SOIREE iron release experiment.
698 Deep-Sea Res II 48:2655–2680

699 Trull TW, Daviesa D, Cascottie K (2008) Insights into nutrient assimilation and export in
700 naturally iron-fertilized waters of the Southern Ocean from nitrogen, carbon and oxygen
701 isotopes. Deep-Sea Res II 55: 220-240

702 Vanderklift MA, Ponsard S (2003) Sources of variation in consumer-diet $\delta^{15}\text{N}$ enrichment: a
703 meta-analysis. Oecologia 136:169-182

704 Vander Zanden MJ, Rasmussen JB (2001) Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ trophic fractionation:
705 Implications for aquatic food web studies Limnol Oceanogr 46: 2061-2066

706 Wallace B, Seminoff J, Kilham S, Spotila J, Dutton P (2006) Leatherback turtles as
707 oceanographic indicators: stable isotope analyses reveal a trophic dichotomy between
708 ocean basins. Mar Biol 149:953-960

709 Wienecke BC, Lawless R, Rodary D, Bost CA, Thomson R, Pauly T, Robertson G, Kerry
710 KR, Le Maho Y (2000) Adélie penguin foraging behaviour and krill abundance along the
711 Wilkes and Adélie Land coasts, Antarctica. Deep-Sea Res II 47:2573-2587

Table 1. Foraging characteristics and blood $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of penguin species during the chick-rearing period.

Species	Locations	Foraging areas	Foraging range (km)	Chick diet	$\delta^{13}\text{C}$ (‰)*	$\delta^{15}\text{N}$ (‰)*	References
Northern rockhopper penguin	Amsterdam (37.8°S)	Subtropical Zone	< 10	Squid & crustaceans	-19.5 ± 0.3 (n=10)	9.2 ± 0.3 (n=10)	Tremblay & Cherel (2003)
Southern rockhopper penguin	Crozet (46.42°S)	Polar Frontal Zone	< 10	Crustaceans	-21.2 ± 0.1 (n=10)	6.8 ± 0.3 (n=10)	Tremblay & Cherel (2003)
King penguin	Crozet (46.42°S)	Polar Front (50°S)	340-450	Pelagic fish	-22.6 ± 0.1 (n=10)	10.3 ± 0.2 (n=10)	Cherel et al. (1993) Charrassin & Bost (2001)
Adélie penguin	Adélie Land (66.7°S)	Antarctic Zone	< 50	Crustaceans (fish)	-24.8 ± 0.5 (n=9)	10.1 ± 0.8 (n=9)	Wienecke et al. (2000)

* Bulk values of whole blood of penguin chicks during the austral summer 2001-2002 from Cherel and Hobson (2007).

Table 2. $\delta^{15}\text{N}$ values of the bulk sample and isolated amino acids (AAs) of blood collected from penguin chicks from the southern Indian Ocean.

Source AAs are indicated by bold print. nc means not considered because of peak co-elution. LME estimate for each species (mean \pm SE) were calculated with linear mixed-effect models that accounted for the heterogeneity of the data set (see methods).

		Amino acid $\delta^{15}\text{N}$ values								
	Bulk $\delta^{15}\text{N}$	phe	his	gly	leu	isoleu	glu	pro	asp	ala
Northern rockhopper penguin (NRP), 37,8°S										
NRP 1	9.2	2.5	3.0	6.9	18.5	19.7	19.0	23.9	14.6	20.3
		3.1	2.9	5.6	17.2	17.4	18.3	22.4	14.0	19.8
		1.6	0.9	3.9	16.1	17.7	16.0	20.4	13.8	18.7
		1.7	2.2	4.7	17.2	18.1	16.3	23.6	13.0	19.8
NRP 2	9.2	2.4	1.2	5.7	18.4	19.5	17.5	21.8	13.3	18.5
		1.5	2.9	5.5	18.3	19.4	18.0	23.8	14.1	19.3
		2.6	3.6	7.7	18.2	17.8	16.2	25.4	14.7	20.1
		1.7	5.1	8.8	18.8	20.4	19.9	26.0	15.4	20.7
LME	estimate	2.1 \pm 0.5	2.7 \pm 1.4	6.1 \pm 0.8	17.8 \pm 0.5	18.8 \pm 0.7	17.7 \pm 0.6	23.4 \pm 0.7	14.1 \pm 0.5	19.7 \pm 0.4
Southern rockhopper penguin (SRP). 46.7°S										
SRP 1	6.8	1.1	2.7	9.6	12.8	12.8	14.8	16.4	12.0	16.8
		1.7	1.0	7.9	12.7	13.2	14.0	16.5	12.9	16.3
		1.4	1.3	7.8	13.1	14.4	14.9	17.4	11.9	15.8
		1.1	2.2	7.9	12.7	12.6	15.8	16.0	11.1	16.4
SRP 2	6.8	-0.2	4.7	6.4	13.5	14.4	15.5	19.0	12.0	16.6
		1.8	5.9	8.6	15.0	15.2	16.1	18.0	12.4	17.3
		1.2	7.6	7.3	14.8	14.4	15.5	17.5	12.4	17.3
LME	estimate	1.1 \pm 0.5	3.9 \pm 1.4	7.9 \pm 0.8	13.6 \pm 0.5	13.9 \pm 0.8	15.2 \pm 0.6	17.3 \pm 0.7	12.1 \pm 0.5	16.6 \pm 0.4

King Penguin (KP). 50.0°S										
KP 1	10.4	2.8	nc	13.6	19.8	21.5	20.2	22.4	18.4	22.3
		3.0	nc	14.5	nc	nc	22.0	24.6	19.3	22.9
		4.1	7.7	12.2	20.0	21.8	21.5	23.2	18.0	21.3
		2.5	5.3	11.5	20.1	20.4	21.4	22.1	18.2	21.0
KP 2	10.5	2.5	5.0	nc	19.6	nc	19.7	23.1	nc	nc
		2.0	4.5	nc	20.1	nc	19.5	23.1	18.1	nc
		1.0	3.0	nc	19.9	nc	17.4	21.2	nc	nc
KP 3	10.5	2.7	4.4	nc	19.3	18.6	18.8	20.7	15.9	20.6
		2.3	2.3	11.2	18.9	18.4	18.7	21.2	16.4	21.2
		3.8	5.9	12.8	20.9	nc	20.0	22.4	17.8	22.3
LME	estimate	2.6±0.5	4.9±1.3	12.6±0.9	19.8±0.4	20.0±0.8	19.9±0.6	22.4±0.7	17.7±0.5	21.7±0.4
Adélie Penguin (AP). 66.7°S										
AP 1	9.5	3.1	9.0	8.8	16.7	15.8	18.6	18.6	14.4	21.0
		3.8	5.7	9.7	17.2	15.4	18.3	18.7	13.6	21.8
		2.9	5.7	7.9	16.4	16.2	18.2	18.3	13.8	20.1
AP 2	10.0	1.4	1.7	10.9	16.5	16.0	17.4	17.3	14.0	19.4
		3.7	3.6	13.2	18.2	17.9	20.1	19.5	15.3	21.9
		2.8	3.8	11.5	17.3	17.6	18.2	18.2	15.0	20.3
		2.9	2.7	11.0	17.0	16.3	18.6	18.0	14.8	21.1
AP 3	10.3	4.7	5.7	9.4	17.3	17.0	17.9	19.1	15.5	19.4
		4.3	6.5	11.3	17.8	17.1	19.7	20.9	16.1	20.9
		5.1	5.4	11.5	17.4	16.3	17.8	19.3	14.6	20.1
LME	estimate	3.5±0.5	5.2±0.9	10.5±0.5	17.2±0.3	16.5±0.5	18.5±0.4	18.8±0.5	14.7±0.3	20.6±0.2

Table 3. $\delta^{13}\text{C}$ values of the bulk sample and isolated amino acids (AAs) of blood collected from penguin chicks from the southern Indian Ocean.

Normal font indicates essential AAs and bold print indicates non-essential AAs. nc means not considered because of peak co-elution.

Bulk $\delta^{13}\text{C}$	Amino Acid $\delta^{13}\text{C}$ values. ‰								
	phe	lys	arg	thr	his	gly	pro	asp	ala
Northern rockhopper penguin (NRP). 37.8°S									
-19.4	-20.7	-18.8	-29.8	-8.4	-7.9	-7.0	-13.4	-18.8	-19.8
-19.5	-20.6	-18.4	-29.5	-8.4	-7.9	-7.9	-13.3	-18.0	-18.7
-19.7	-20.8	-18.4	-29.9	-9.1	-6.7	-6.1	-15.7	-18.5	-19.9
Southern rockhopper penguin (SRP). 46.7°S									
-21.3	-21.9	-19.2	-30.4	-10.5	-12.7	-8.2	-16.8	-19.3	-22.8
-21.0	-21.9	-19.0	-30.0	-9.3	-11.0	-9.9	-18.1	-19.7	-22.4
-21.0	-21.8	-19.1	-30.2	-10.4	-12.0	-6.9	-18.2	-19.1	-21.8
King Penguin (KP). 50.0°S									
-22.5	-22.7	-20.1	-31.8	-7.9	-9.2	-10.4	-18.5	-20.9	-24.2
-22.6	-22.6	-19.9	-31.7	nc	-10.0	nc	-19.6	-20.7	-24.7
-22.5	-22.9	-20.4	-31.9	-5.0	-10.0	-11.7	-20.2	-24.8	-21.7
Adélie Penguin (AP). 66.7°S									
-24.5	-25.2	-21.2	-33.4	-11.5	-14.9	-12.2	-22.2	-21.5	-26.3
-24.9	-24.7	-21.2	-33.3	-12.7	-14.3	-12.0	-21.9	-22.0	-26.4
-24.4	-25.2	-21.6	-34.0	-11.6	-10.6	-12.0	-22.0	-21.1	-25.7

Table 4. Trophic positions estimates from the literature (na for non available), linear mixed-effect model predictions of $\delta^{15}\text{N}_{\text{glu}} - \delta^{15}\text{N}_{\text{phe}}$ for the three penguin species, and estimated TEF between source and trophic transfer amino acids for penguin chick's blood using Equation 1 (see material & methods for more details).

Species	Trophic position*	Predicted $\delta^{15}\text{N}_{\text{glu}} - \delta^{15}\text{N}_{\text{phe}}$ (‰)	Estimated TEF (‰)
Northern rockhopper penguin	na	15.5	nd
Southern rockhopper penguin	4.0	14.1	3.4
King penguin	4.5	17.2	3.8
Adélie penguin	3.9	15.0	3.8

*From Cherel et al. (2008)

Figure legends

Fig. 1. Sampling locations of the four penguin species in the Southern Indian Ocean (●): Northern rockhopper penguin (NRP), Southern rockhopper penguin (SRP), King penguin (KP) and Adélie penguin (AP). Mean chlorophyll distribution averaged over the period from October 1997 to October 2002 (mg m^{-3}) in the Southern Ocean overlaid with the Southern Ocean fronts are also indicated. Mean front positions are mapped using SSH (adopted from [Sokolov and Rintoul, 2007; 2009]). The STF position (light brown line) is based on temperature criterion as in [Sokolov and Rintoul, 2002]. The STF is mapped using WOCE global hydrographic climatology [Gouretski and Koltermann, 2004]. The 2000 m bathymetric contour is indicated by light blue line.

Fig. 2. Stable isotope values ($\delta^{15}\text{N}$ to the left and $\delta^{13}\text{C}$ to the right, ‰) of bulk and individual amino acids of the four penguin species: ○. Northern rockhopper penguin (NRP); ● Southern rockhopper penguin (SRP); △ King penguin (KP) and ■ Adélie penguin (AP). Nb: for nitrogen, mean values are predicted values (see Table 2).

Fig. 3. Variations of $\delta^{15}\text{N}$ values for bulk (■, mean \pm SD), phenylalanine (△, Phe) and glutamic acid (▲, Glu) (predicted values \pm SE, see methods) with latitude for four penguin species: Northern rockhopper penguin (NRP), southern rockhopper penguin (SRP), king penguin (KP) and Adélie penguin (AP).

Fig. 4. Variations of bulk and amino acid (AA) $\delta^{13}\text{C}$ values with latitude (a) and of AA $\delta^{13}\text{C}$ values with bulk (b) for four penguin species: Northern rockhopper penguin (NRP), southern rockhopper penguin (SRP), king penguin (KP) and Adélie penguin (AP).




Fig. 1. Sampling locations of the four penguin species in the Southern Indian Ocean (●): Northern rockhopper penguin (NRP), Southern rockhopper penguin (SRP), King penguin (KP) and Adélie penguin (AP). Mean chlorophyll distribution averaged over the period from October 1997 to October 2002 (mg m^{-3}) in the Southern Ocean overlaid with the Southern Ocean fronts are also indicated. Mean front positions are mapped using SSH (adopted from [Sokolov and Rintoul, 2007; 2009]). The STF position (light brown line) is based on temperature criterion as in [Sokolov and Rintoul, 2002]. The STF is mapped using WOCE global hydrographic climatology [Gouretski and Koltermann, 2004]. The 2000 m bathymetric contour is indicated by light blue line.




Fig. 2. Stable isotope values ($\delta^{15}\text{N}$ to the left and $\delta^{13}\text{C}$ to the right, ‰) of bulk and individual amino acids of the four penguin species: ○. Northern rockhopper penguin (NRP); ● Southern rockhopper penguin (SRP); △ King penguin (KP) and ■ Adélie penguin (AP). Nb: for nitrogen, mean values are predicted values (see Table 2).




Fig. 3. Variations of $\delta^{15}\text{N}$ values for bulk (■, mean \pm SD), phenylalanine (Δ , Phe) and glutamic acid (Δ , Glu) (predicted values \pm SE, see methods) with latitude for four penguin species: Northern rockhopper penguin (NRP), southern rockhopper penguin (SRP), king penguin (KP) and Adélie penguin (AP).




Fig. 4. Variations of bulk and amino acid (AA) $\delta^{13}\text{C}$ values with latitude (a) and of AA $\delta^{13}\text{C}$ values with bulk (b) for four penguin species: Northern rockhopper penguin (NRP), southern rockhopper penguin (SRP), king penguin (KP) and Adélie penguin (AP).