



**HAL**  
open science

# On Subordination Principles for Generalized Shannon Sampling Series

Andi Kivinukk, Gert Tamberg

► **To cite this version:**

Andi Kivinukk, Gert Tamberg. On Subordination Principles for Generalized Shannon Sampling Series. SAMPTA'09, May 2009, Marseille, France. pp.General session. hal-00453086

**HAL Id: hal-00453086**

**<https://hal.science/hal-00453086>**

Submitted on 3 Feb 2010

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On Subordination Principles for Generalized Shannon Sampling Series

Andi Kivinukk <sup>(1)</sup> and Gert Tamberg <sup>(2)</sup>

(1) Dept. of Math., Tallinn University, Narva Road 25, 10120 Tallinn, Estonia

(2) Dept. of Math., Tallinn University of Technology, Ehitajate tee 5 19086 Tallinn, Estonia  
andik@tlu.ee, gert.tamberg@mail.ee

## Abstract:

This paper provides some subordination equalities and their applications for the generalized Shannon sampling series.

## 1. Introduction

For the uniformly continuous and bounded functions  $f \in C(\mathbb{R})$  the generalized Shannon sampling series (see [3] and references cited there) are given by ( $t \in \mathbb{R}; W > 0$ )

$$(S_W f)(t) := \sum_{k=-\infty}^{\infty} f\left(\frac{k}{W}\right) s(Wt - k), \quad (1)$$

where the condition for the operator  $S_W : C(\mathbb{R}) \rightarrow C(\mathbb{R})$  to be well-defined is that for the kernel function  $s = s(t)$  we assume

$$\sum_{k=-\infty}^{\infty} |s(u - k)| < \infty \quad (u \in \mathbb{R}).$$

Let be given an even window function  $\lambda \in C_{[-1,1]}$ ,  $\lambda(0) = 1$ ,  $\lambda(u) = 0$  ( $|u| \geq 1$ ), then in our approach the kernel function will be defined by the equality

$$s(t) := s_\lambda(t) := \int_0^1 \lambda(u) \cos(\pi t u) du. \quad (2)$$

Many window functions have been used in applications (see, e.g. [1], [2], [4], [8]), in Signal Analysis in particular. Next window functions are important for our subordination equalities.

1)  $\lambda_{(r)}(u) = 1 - u^r$ ,  $r \geq 1$  defines the Zygmund (or Riesz) kernel, denoted by  $z_r = z_r(t)$ , which special case  $r = 1$ , the Fejér (or Bartlett, see [8]) kernel  $s_F(t) = \frac{1}{2} \text{sinc}^2 \frac{t}{2}$ , is well-known; the special case  $r = 2$  is called also as the Welch [8] kernel;

2)  $\lambda_j(u) := \cos \pi(j + 1/2)u$ ,  $j = 0, 1, 2, \dots$  defines the Rogosinski-type kernel (see [5]) in the form

$$r_j(t) := \frac{(-1)^j (j + 1/2) \cos \pi t}{\pi (j + 1/2)^2 - t^2}; \quad (3)$$

3)  $\lambda_H(u) := \cos^2 \frac{\pi u}{2} = \frac{1}{2}(1 + \cos \pi u)$  defines the Hann kernel (see [6])

$$s_H(t) := \frac{1}{2} \frac{\text{sinc} t}{1 - t^2}. \quad (4)$$

Concerning some direct (Jackson-type) approximation theorems we present certain subordination equalities, which show that the sampling operators, like Rogosinski, Zygmund, and Hann, are in some sense basic.

## 2. Subordination equalities

Subordination equalities state some relations between two sampling operators.

### 2.1 Subordination by the Rogosinski-type sampling series

Let consider the Rogosinski-type sampling operators  $R_{W,j}$  defined by the kernel functions  $r_j$  in (3). These kernel functions are deduced by the window functions  $\lambda_j(u) := \cos \pi(j + 1/2)u$ , ( $j \in \mathbb{N}$ ) and as a family of functions it forms an orthogonal system on  $[0, 1]$ . Therefore, we may represent a quite arbitrary window function  $\lambda$  by its Fourier series. But the Fourier representation allows us to prove for a given kernel function  $s$  the sampling series

$$s(t) = 2 \sum_{j=0}^{\infty} s(j + 1/2) r_j(t).$$

In following  $B_\sigma^p$  stands for the Bernstein class, it consists of those bounded functions  $f \in L^p(\mathbb{R})$  ( $1 \leq p \leq \infty$ ), which can be extended to an entire function  $f(z)$  ( $z \in \mathbb{C}$ ) of exponential type  $\sigma$ . For  $s \in B_\pi^1$  the sampling series above is absolutely convergence and by (1) we get formally the equalities

$$S_W f = 2 \sum_{j=0}^{\infty} s(j + 1/2) R_{W,j} f,$$

$$f - S_W f = 2 \sum_{j=0}^{\infty} s(j + 1/2) (f - R_{W,j} f),$$

calling as the subordination equalities, since the approximation properties of the general sampling operators (1) can be described via the approximation properties of the Rogosinski-type sampling operators  $R_{W,j} : C(\mathbb{R}) \rightarrow C(\mathbb{R})$ . We have proved that [5]

$$\|R_{W,j}\| = \frac{4}{\pi} \sum_{\ell=0}^{2j} \frac{1}{2\ell + 1} = \frac{2}{\pi} \log(j + 1) + O(1),$$

thus the subordination equalities are valid, when

$$\sum_{j=0}^{\infty} |s(j+1/2)| \log(j+1) < \infty.$$

Similar subordination equalities can be deduced for some interpolating sampling series, i.e. for which the equation  $(\tilde{S}_W f)(\frac{k}{W}) = f(\frac{k}{W})$  ( $k \in \mathbb{Z}$ ) is valid. In [7] we have proved that the interpolating sampling operators will be defined by (1) using the kernel  $\tilde{s}(t) := 2s(2t)$ , where the kernel  $s$  is generated by (2) with a window function  $\lambda$  for which  $\lambda(u) + \lambda(1-u) = 1$  ( $u \in [0, 1]$ ).

Let the operator  $S_W^\alpha : C(\mathbb{R}) \rightarrow C(\mathbb{R})$  be defined by the kernel  $s_\alpha := \alpha s(\alpha \cdot) \in B_{\alpha\pi}^1$  ( $0 < \alpha \leq 2$ ), where  $s \in B_\pi^1$ , and the modified Hann operator  $H_{W,j}^\alpha$  is defined by the kernel

$$s_{H,j}^\alpha(t) := \frac{\alpha}{2} \frac{(2j+1)^2}{(2j+1)^2 - (\alpha t)^2} \text{sinc}(\alpha t). \quad (5)$$

Then here we have (see [7], Th. 2.3 and 2.4)

$$S_W^\alpha f = 4 \sum_{j=0}^{\infty} s(2j+1) H_{W,j}^\alpha f,$$

$$f - S_W^\alpha f = 4 \sum_{j=0}^{\infty} s(2j+1) (f - H_{W,j}^\alpha f).$$

## 2.2 Subordination by the Rogosinski-type sampling series: 2D case

The two-dimensional generalized sampling series has the form

$$(S_W f)(x, y) := \sum_{k, l=-\infty}^{\infty} f\left(\frac{k}{W}, \frac{l}{W}\right) s(Wx - k, Wy - l),$$

in particular, the multiplicative Rogosinski-type sampling series we define as

$$(R_{W;i,j} f)(x, y) := \sum_{k, l=-\infty}^{\infty} f\left(\frac{k}{W}, \frac{l}{W}\right) r_i(Wx - k) r_j(Wy - l),$$

where the Rogosinski-type kernel  $r_j$  is defined by (3). Here our subordination equalities read as

$$S_W f = 4 \sum_{i,j=0}^{\infty} s(i+1/2, j+1/2) R_{W;i,j} f,$$

$$f - S_W f = 4 \sum_{i,j=0}^{\infty} s(i+1/2, j+1/2) (f - R_{W;i,j} f),$$

provided

$$\sum_{i,j=1}^{\infty} |s(i+1/2, j+1/2)| \log i \log j < \infty.$$

By given subordination equalities we see that the non-multiplicative sampling series may be studied by the multiplicative Rogosinski-type sampling series.

## 2.3 Subordination by the Zygmund sampling series

The Zygmund sampling operator  $Z_W^r$  will be defined by the window function  $\lambda_{(r)}(u) = 1 - u^r$ ,  $r \geq 1$ . Let us consider the kernel  $s$  in (2), for which the corresponding window function has the power series representation

$$\lambda(u) = 1 - \sum_{j=r}^{\infty} c_j u^j.$$

Then the formal subordination equalities are in the shape

$$S_W f = \sum_{j=r}^{\infty} c_j Z_W^j f,$$

$$f - S_W f = \sum_{j=r}^{\infty} c_j (f - Z_W^j f).$$

Several other subordination equalities and their applications will be presented.

## 3. Acknowledgments

This research was partially supported by the Estonian Sci. Foundation, grants 6943, 7033, and by the Estonian Min. of Educ. and Research, projects SF0132723s06, SF0140011s09.

## References:

- [1] H. H. Albrecht. A family of cosine-sum windows for high resolution measurements. In *IEEE International Conference on Acoustics, Speech and Signal Processing, Salt Lake City, Mai 2001*, pages 3081–3084. Salt Lake City, 2001.
- [2] R. B. Blackman and J. W. Tukey. *The measurement of power spectra*. Wiley-VCH, New York, 1958.
- [3] P. L. Butzer, G. Schmeisser, and R. L. Stens. An introduction to sampling analysis. In F. Marvasti, editor, *Nonuniform Sampling, Theory and Practice*, pages 17–121. Kluwer, New York, 2001.
- [4] F. J. Harris. On the use of windows for harmonic analysis. *Proc. of the IEEE*, 66:51–83, 1978.
- [5] A. Kivinuuk and G. Tamberg. On sampling series based on some combinations of sinc functions. *Proc. of the Estonian Academy of Sciences. Physics Mathematics*, 51:203–220, 2002.
- [6] A. Kivinuuk and G. Tamberg. On sampling operators defined by the Hann window and some of their extensions. *Sampling Theory in Signal and Image Processing*, 2:235–258, 2003.
- [7] A. Kivinuuk and G. Tamberg. Interpolating generalized Shannon sampling operators, their norms and approximation properties. *Sampling Theory in Signal and Image Processing*, 8:77–95, 2009.
- [8] E. H. W. Meijering, W. J. Niessen, and M. A. Viergever. Quantitative evaluation of convolution-based methods for medical image interpolation. *Medical Image Analysis*, 5:111–126, 2001.