



# Quasi-one- and quasi-two-dimensional perfect Bose gas: the second critical density and generalised condensation

Mathieu Beau, Valentin A. Zagrebnov

## ► To cite this version:

Mathieu Beau, Valentin A. Zagrebnov. Quasi-one- and quasi-two-dimensional perfect Bose gas: the second critical density and generalised condensation. 2010. hal-00453337v1

**HAL Id: hal-00453337**

**<https://hal.science/hal-00453337v1>**

Preprint submitted on 4 Feb 2010 (v1), last revised 14 Apr 2010 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Quasi-one- and quasi-two-dimensional perfect Bose-gas: the second critical density and generalised condensation

Mathieu Beau and Valentin A. Zagrebnov

Université de la Méditerranée and Centre de Physique Théorique - UMR 6207 \*,  
Luminy - Case 907, 13288 Marseille, Cedex 09, France

(Dated: February 03, 2010)

In this letter we discuss a relevance of the 3D Perfect Bose gas (PBG) condensation in extremely elongated vessels for the study of anisotropic condensate coherence and the "quasi-condensate". To this end we analyze the case of exponentially anisotropic (*van den Berg*) boxes, when there are *two* critical densities  $\rho_c < \rho_m$  for a generalised Bose-Einstein Condensation (BEC). Here  $\rho_c$  is the standard critical density for the PBG. We consider three examples of anisotropic geometry: slabs, squared beams and "cigars" to demonstrate that the "quasi-condensate" which exists in domain  $\rho_c < \rho < \rho_m$  is in fact the van den Berg-Lewis-Pulé generalised condensation (vdBLP-GC) of the type III with no macroscopic occupation of any mode.

We show that for the slab geometry the second critical density  $\rho_m$  is a threshold between *quasi-two-dimensional* (*quasi-2D*) condensate and the *three dimensional* (*3D*) regime when there is a coexistence of the "quasi-condensate" with the standard one-mode BEC. On the other hand, in the case of squared beams and "cigars" geometries critical density  $\rho_m$  separates *quasi-1D* and *3D* regimes. We calculate the value of difference between  $\rho_c$ ,  $\rho_m$  (and between corresponding critical temperatures  $T_m$ ,  $T_c$ ) to show that observed space anisotropy of the condensate coherence can be described by a critical exponent  $\gamma(T)$  related to the anisotropic ODLRO. We compare our calculations with physical results for extremely elongated traps that manifest "quasi-condensate".

PACS numbers: 05.30.Jp, 03.75.Hh, 67.40.-w

1. One can rigorously show that there is no a *conventional* Bose-Einstein condensation (BEC) in the one- (*1D*) and two-dimensional (*2D*) boson systems or in the three-dimensional squared beams (cylinders) and slabs (films). For interacting Bose-gas it results from the Bogoliubov-Hohenberg theorem [1], [2], based on a non-trivial Bogoliubov inequality, see e.g. [3]. For the perfect Bose-gas this result is much easier, since it follows from the explicit analysis of the occupation number density in one-particle eigenstates, see e.g. [4]. A common point is the Bogoliubov  $1/q^2$ -theorem [1],[5], [6], which implies destruction of the macroscopic occupation of the ground-state by thermal fluctuations.

Renewed interest to eventual possibility of the "condensate" in the quasi-one-, or -two-dimensional (*quasi-1D* or *-2D*) boson gases (i.e., in cigar-shaped systems or slabs) is motivated by recent experimental data indicating the existence of so-called "quasi-condensate" in anisotropic traps [7]-[9] and BKT crossover [10].

The aim of this letter is *twofold*. First we show that a natural modeling of slabs by highly anisotropic *3D*-cuboid implies in the thermodynamic limit the van den Berg-Lewis-Pulé *generalised condensation* (vdBLP-GC) [11] of the Perfect Bose-Gas (PBG) for densities larger than the *first*, i.e. the standard critical  $\rho_c(\beta)$  for the inverse temperature  $\beta = 1/(k_B T)$ . Notice, that a spe-

cial case of this (induced by the geometry) condensation was pointed out for the first time by Casimir [12], although the theoretical concept and the name are due to Girardeau [13]. So, for the PBG the "quasi-condensate" is in fact the vdBLP-GC. Here we generalise these results to the highly anisotropic *3D*-cuboid with anisotropy in *one-dimension*, which is a model for infinite squared beams or cylinders, and "cigar" type traps.

Second, we show that for the slab geometry with *exponential* growing (for  $\alpha > 0$  and  $L \rightarrow \infty$ ) of two edges,  $L_1 = L_2 = Le^{\alpha L}$ ,  $L_3 = L$ , of the anisotropic boxes:  $\Lambda = L_1 \times L_2 \times L_3 \in \mathbb{R}^3$ , there is a *second* critical density  $\rho_m(\beta) := \rho_c(\beta) + 2\alpha/\lambda_\beta^2 \geq \rho_c(\beta)$  such that the vdBLP-GC changes its properties when  $\rho > \rho_m(\beta)$ . This surprising behaviour of the BEC for the PBG was discovered by van den Berg [14], developed in [15], and then in [16],[17] for the spin-wave condensation.

Our original observation concerns the coexistence of two types of the vdBLP-GC for  $\rho > \rho_m(\beta)$  (or for corresponding temperatures  $T < T_m(\rho)$  for a fixed density) and the analysis of the coherence length (ODLRO) in this anisotropic geometry. We extend also our observation to obtain another new result proving the existence of the *second* critical density in the squared beam and in the "cigar" type traps for exponentially weak harmonic potential confinement in *one* direction.

We use these results to calculate the temperature dependence of the vdBLP-GC particle density for the case of two critical densities,  $\rho_m(\beta) < \rho_c(\beta)$  and to apply the recent scaling approach [18] to the ODLRO asymptotics in this case.

---

\*Université de Provence - Aix-Marseille I, Université de la Méditerranée - Aix-Marseille II, Université du Sud - Toulon - Var, FRUMAM (FR 2291)

2. It is known that all kinds of BEC in the PBG are defined by the *limiting spectrum* of the one-particle Hamiltonian  $T_\Lambda^{(N=1)} = -\hbar^2 \Delta / (2m)$ , when cuboid  $\Lambda \uparrow \mathbb{R}^3$ . In this paper we make this operator self-adjoint by fixing the *Dirichlet* boundary conditions on  $\partial\Lambda$ , although our results are valid for all *non-attractive* boundary conditions. Then the spectrum is the set

$$\{\varepsilon_s = \frac{\hbar^2}{2m} \sum_{j=1}^3 (\pi s_j / L_j)^2\}_{s_j \in \mathbb{N}} \quad (1)$$

and  $\{\phi_{s,\Lambda}(x) = \prod_{j=1}^3 \sqrt{2/L_j} \sin(\pi s_j x_j / L_j)\}_{s_j \in \mathbb{N}}$  are the eigenfunctions. Here  $\mathbb{N}$  is the set of the natural numbers and  $s = (s_1, s_2, s_3) \in \mathbb{N}^3$  is the multi-index.

In the grand-canonical ensemble  $(T, V, \mu)$ , here  $V = L_1 L_2 L_3$  is the volume of  $\Lambda$ , the mean occupation number of the state  $\phi_{s,\Lambda}$  is  $N_s(\beta, \mu) = (e^{\beta(\varepsilon_s - \mu)} - 1)^{-1}$ , where  $\mu < \inf_s \varepsilon_{s,\Lambda}$ . Then for the fixed total particle density  $\rho$  the corresponding value of the chemical potential  $\mu_\Lambda(\beta, \rho)$  is a unique solution of the equation  $\rho = \sum_{s \in \mathbb{N}^3} N_s(\beta, \mu) / V =: N_\Lambda(\beta, \mu) / V$ . Independent of the way  $\Lambda \uparrow \mathbb{R}^3$ , one gets the limit  $\rho(\beta, \mu) = \lim_{V \rightarrow \infty} N_\Lambda(\beta, \mu) / V$ , which is the total particle density for  $\mu \leq \lim_{V \rightarrow \infty} \inf_s \varepsilon_s = 0$ . Since  $\rho_c(\beta) := \sup_{\mu \leq 0} \rho(\beta, \mu) = \rho(\beta, \mu = 0) < \infty$ , it is the (*first*) critical density for the 3D PBG:  $\rho_c(\beta) = \zeta(3/2) / \lambda_\beta^3$ . Here  $\zeta(s)$  is the Riemann  $\zeta$ -function and  $\lambda_\beta := \hbar \sqrt{2\pi\beta/m}$  is the de Broglie thermal length.

For  $\Lambda = L e^{\alpha L} \times L e^{\alpha L} \times L$  one gets ([14, 15]) that for any fixed  $s_1, s_2$  and  $\mu \leq 0$  the limit of Darboux-Riemann sums

$$\lim_{L \rightarrow \infty} \sum_{s \neq (s_1, s_2, 1)} \frac{N_s(\beta, \mu)}{V_L} = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \frac{d^3 k}{e^{\beta(\hbar^2 k^2 / 2m - \mu)} - 1}. \quad (2)$$

We denote by  $\mu_L(\beta, \rho) := \varepsilon_{(1,1,1)} - \Delta_L(\beta, \rho)$ , where  $\Delta_L(\beta, \rho) \geq 0$  is a unique the solution of the equation:

$$\rho = \sum_{s=(s_1, s_2, 1)} \frac{N_s(\beta, \mu)}{V_L} + \sum_{s \neq (s_1, s_2, 1)} \frac{N_s(\beta, \mu)}{V_L}. \quad (3)$$

Since by (2):  $\lim_{L \rightarrow \infty} \sum_{s \neq (s_1, s_2, 1)} N_s(\beta, \mu = 0) / V_L = \rho_c(\beta)$ , for  $\rho > \rho_c(\beta)$  the limit  $L \rightarrow \infty$  of the first sum in (3) is equal to

$$\begin{aligned} \lim_{L \rightarrow \infty} \sum_{s=(s_1, s_2, 1)} \frac{N_s(\beta, \mu)}{V_L} &= \\ \lim_{L \rightarrow \infty} \frac{1}{L} \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} \frac{d^2 k}{e^{\beta(\hbar^2 k^2 / 2m + \Delta_L(\beta, \rho))} - 1} &= \\ \lim_{L \rightarrow \infty} -\frac{1}{\lambda_\beta^2 L} \ln[\beta \Delta_L(\beta, \rho)] &= \rho - \rho_c(\beta). \end{aligned} \quad (4)$$

This implies the asymptotics:

$$\Delta_L(\beta, \rho) = \frac{1}{\beta} e^{-\lambda_\beta^2(\rho - \rho_c(\beta))L} + \dots \quad (5)$$




FIG. 1: For the slab geometry, the blue curve  $\rho_c(1/(k_B T))$  is the first critical line for the BEC transition as a function of  $T$ , the red curve  $\rho_m(1/(k_B T)) = \rho_c(1/(k_B T)) + 2\alpha/\lambda_\beta^2$  is the second critical line. Notice that above the red curve there is a *coexistence* between "quasi-condensate" (vdBLP-GC of type III) and the *conventional* condensate in the ground state (vdBLP-GC of type I), between two curve there is only "quasi-condensates" phase and below the blue curve there is no condensate.

Notice that representation of the limit (4) by the integral (see (1)) is valid only when  $\lambda_\beta^2(\rho - \rho_c(\beta)) < 2\alpha$ . For  $\rho$  larger than the *second* critical density:  $\rho_m(\beta) := \rho_c(\beta) + 2\alpha/\lambda_\beta^2$  the correction  $\Delta_L(\beta, \rho)$  must converge to zero faster than  $e^{-2\alpha L}$ . Now to keep the difference  $\rho - \rho_m(\beta) > 0$  we have to return back to the original sum representation (3) and (as for the standard BEC) to take into account the impact of the ground state occupation density *together* with a saturated *non-ground state* (i.e. generalised) condensation  $\rho_m(\beta) - \rho_c(\beta)$  as in (4). For this case the asymptotics of  $\Delta_L(\beta, \rho > \rho_m(\beta))$  is completely different than (5) and it is equal to  $\Delta_L(\beta, \rho) = [\beta(\rho - \rho_m(\beta))V_L]^{-1}$ . Since  $V_L = L^3 e^{2\alpha L}$ , we obtain:

$$\begin{aligned} \lim_{L \rightarrow \infty} \sum_{s=(s_1 > 1, s_2 > 1, 1)} \frac{N_s(\beta, \mu)}{V_L} &= \\ \lim_{L \rightarrow \infty} -\frac{1}{\lambda_\beta^2 L} \ln[\beta \Delta_L(\beta, \rho)] &= 2\alpha/\lambda_\beta^2 = \\ \rho_m(\beta) - \rho_c(\beta), \end{aligned} \quad (6)$$

and the ground-state term gives the *macroscopic* occupation:

$$\rho - \rho_m(\beta) = \lim_{L \rightarrow \infty} \frac{1}{V_L} \frac{1}{e^{\beta(\varepsilon_{(1,1,1)} - \mu_L(\beta, \rho))} - 1}. \quad (7)$$

Notice that for  $\rho_c(\beta) < \rho < \rho_m(\beta)$  we obtain the vdBLP-GC (of the *type III*), i.e. *none* of the single-particle states are *macroscopically* occupied, since by virtue of (1) and (5) for any  $s$  one has:

$$\rho_s(\beta, \rho) := \lim_{L \rightarrow \infty} \frac{1}{V_L} \frac{1}{e^{\beta(\varepsilon_s - \mu_L(\beta, \rho))} - 1} = 0. \quad (8)$$

On the other hand, the asymptotics  $\Delta_L(\beta, \rho > \rho_m(\beta)) = [\beta(\rho - \rho_m(\beta))V_L]^{-1}$  implies

$$\rho_{s \neq (1,1,1)}(\beta, \rho) := \lim_{L \rightarrow \infty} \frac{1}{V_L} \frac{1}{e^{\beta(\varepsilon_s - \mu_L(\beta, \rho))} - 1} = 0, \quad (9)$$

i.e. for  $\rho > \rho_m(\beta)$  there is a *coexistence* of the *saturated* type III vdBLP-GC, with the constant density (6), and the standard BEC (i.e. the *type* I vdBLP-GC) in the single state (7).

3. It is curious to note that neither Casimir shaped boxes [11], nor the van den Berg boxes  $\Lambda = Le^{\alpha L} \times L \times L$ , with one-dimensional anisotropy do not produce the *second* critical density  $\rho_m(\beta) \neq \rho_c(\beta)$ . To model infinite squared beams with BEC transitions at two critical densities we propose the one-particle Hamiltonian:  $T_\Lambda^{(N=1)} = -\hbar^2 \Delta / (2m) + m\omega_1^2 x_1^2 / 2$ , with *harmonic trap* in direction  $x_1$  and, e.g., Dirichlet boundary conditions in directions  $x_2, x_3$ . Then the spectrum is the set

$$\{\epsilon_s := \hbar\omega_1(s_1 + 1/2) + \frac{\hbar^2}{2m} \sum_{j=2}^3 (\pi s_j / L_j)^2\}_{s \in \mathbb{N}}. \quad (10)$$

Here multi-index  $s = (s_1, s_2, s_3) \in (\mathbb{N} \cup \{0\}) \times \mathbb{N}^2$ , and the ground-state energy is  $\epsilon_{(0,1,1)}$ . Then for  $\mu_L(\beta, \varrho) := \epsilon_{(0,1,1)} - \Delta_L(\beta, \varrho)$ , the value of  $\Delta_L(\beta, \varrho) \geq 0$ , is a solution of the equation:

$$\varrho := \sum_{s=(s_1,1,1)} \omega_1 \frac{N_s(\beta, \mu)}{L_2 L_3} + \sum_{s \neq (s_1,1,1)} \omega_1 \frac{N_s(\beta, \mu)}{L_2 L_3}, \quad (11)$$

where  $N_s(\beta, \mu) = (e^{\beta(\epsilon_s - \mu)} - 1)^{-1}$ .

Let  $\omega_1 := \hbar/(mL_1^2)$  and  $L_2 = L_3 = L$ . Here  $L_1$  is the harmonic-trap characteristic size in direction  $x_1$ . Then for any  $s_1 \geq 0$  and  $\mu \leq 0$

$$\begin{aligned} \varrho(\beta, \mu) &:= \lim_{L_1, L \rightarrow \infty} \sum_{s \neq (s_1,1,1)} \omega_1 \frac{N_s(\beta, \mu)}{L_2 L_3} = \\ &= \frac{1}{(2\pi)^2} \int_0^\infty dp \int_{\mathbb{R}^2} \frac{d^2 k}{e^{\beta(\hbar p + \hbar^2 k^2 / 2m - \mu)} - 1}. \end{aligned} \quad (12)$$

Therefore, the *first* critical density is *finite*:  $\varrho_c(\beta) := \sup_{\mu \leq 0} \varrho(\beta, \mu) = \varrho(\beta, \mu = 0) < \infty$ . If  $\varrho > \varrho_c(\beta)$ , then the limit  $L \rightarrow \infty$  of the first sum in (11) is

$$\begin{aligned} \lim_{L_1, L \rightarrow \infty} \sum_{s=(s_1,1,1)} \omega_1 \frac{N_s(\beta, \mu_L)}{L_2 L_3} &= \\ \lim_{L \rightarrow \infty} \frac{1}{L^2} \int_0^\infty \frac{dp}{e^{\beta(\hbar p + \Delta_L(\beta, \varrho))} - 1} &= \\ \lim_{L \rightarrow \infty} -\frac{1}{\hbar\beta L^2} \ln[\beta \Delta_L(\beta, \varrho)] &= \varrho - \varrho_c(\beta). \end{aligned} \quad (13)$$

This means that the asymptotics of  $\Delta_L(\beta, \rho)$  is:

$$\Delta_L(\beta, \varrho) = \frac{1}{\beta} e^{-\hbar\beta(\varrho - \varrho_c(\beta))L^2} + \dots \quad (14)$$

Let  $L_1 := Le^{\gamma L^2}$ , for  $\gamma > 0$ . Then, similar to our arguments in 2., the representation of the limit (13) by the integral is valid for  $\hbar\beta(\varrho - \varrho_c(\beta)) < 2\gamma$ . For  $\varrho$  larger than the *second* critical density:  $\varrho_m(\beta) := \varrho_c(\beta) + 2\gamma/(\hbar\beta)$

the chemical potential correction (14) must converge to zero faster than  $e^{-2\gamma L^2}$ . By the same line of reasoning as in 2., to keep the difference  $\varrho - \varrho_m(\beta) > 0$  we have to use the original sum representation (11) and to take into account the input due to the ground state occupation density *together* with a saturated *non-ground state* condensation  $\varrho_m(\beta) - \varrho_c(\beta)$  (13). The asymptotics of  $\Delta_L(\beta, \varrho > \varrho_m(\beta))$  is then equal to  $\Delta_L(\beta, \varrho) = [\beta m(\varrho - \varrho_m(\beta))L^4 e^{2\gamma L^2} / \hbar]^{-1}$ . Hence,

$$\begin{aligned} \lim_{L \rightarrow \infty} \sum_{s=(s_1>0,1,1)} \frac{\hbar}{m} \frac{N_s(\beta, \mu_L)}{L^4 e^{2\gamma L^2}} &= \\ \lim_{L \rightarrow \infty} -\frac{1}{\hbar\beta L^2} \ln[\beta \Delta_L(\beta, \varrho)] &= \frac{2\gamma}{\hbar\beta} = \varrho_m(\beta) - \varrho_c(\beta), \end{aligned} \quad (15)$$

and the ground-state term gives the *macroscopic* occupation:

$$\varrho - \varrho_m(\beta) = \lim_{L \rightarrow \infty} \frac{\hbar}{mL^4 e^{2\gamma L^2}} \frac{1}{e^{\beta(\epsilon_{(0,1,1)} - \mu_L(\beta, \varrho))} - 1}. \quad (16)$$

With this choice of boundary conditions and the one-dimensional anisotropic trap our model of the infinite squared beams manifests the BEC with two critical densities. Again for  $\varrho_c(\beta) < \varrho < \varrho_m(\beta)$  we obtain the *type* III vdBLP-GC, i.e., *none* of the single-particle states are *macroscopically* occupied:

$$\varrho_s(\beta, \varrho) := \lim_{L \rightarrow \infty} \frac{\hbar}{mL^4 e^{2\gamma L^2}} \frac{1}{e^{\beta(\epsilon_s - \mu_L(\beta, \varrho))} - 1} = 0. \quad (17)$$

When  $\varrho_m(\beta) < \varrho$  there is a coexistence of the *type* III vdBLP-GC, with the constant density (20), and the standard *type* I vdBLP-GC in the single state (16), since

$$\varrho_{s \neq (0,1,1)}(\beta, \varrho) := \lim_{L \rightarrow \infty} \frac{\hbar}{mL^4 e^{2\gamma L^2}} \frac{1}{e^{\beta(\epsilon_s - \mu_L(\beta, \varrho))} - 1} = 0. \quad (18)$$

Finally, it is instructive to study a "cigar"-type geometry ensured by the anisotropic harmonic trap:

$$T_\Lambda^{(N=1)} = -\hbar^2 \Delta / (2m) + \sum_{1 \leq j \leq 3} m\omega_j^2 x_j^2 / 2. \quad (19)$$

with  $\omega_1 = \hbar/(mL_1^2), \omega_2 = \omega_3 = \hbar/(mL^2)$ . Here  $L_1, L_2 = L_3 = L$  are the *characteristic* sizes of the trap in three directions and  $\eta_s = \sum_{1 \leq j \leq 3} \hbar\omega_j(s_j + 1/2)$  is the corresponding one-particle spectrum. Then the same reasoning as in (12),(13), yields for  $\mu_L(\beta, n) := \eta_{(0,0,0)} - \Delta_L(\beta, n)$  and auxiliary dimensionality factor  $\kappa > 0$ :

$$\begin{aligned} \lim_{L_1, L \rightarrow \infty} \sum_{s=(s_1,0,0)} \kappa^3 \omega_1 \omega_2 \omega_3 N_s(\beta, \mu_L) &= \\ \lim_{L \rightarrow \infty} -\frac{\kappa^3 \hbar}{\beta(mL^2)^2} \ln[\beta \Delta_L(\beta, n)] &= n - n_c(\beta). \end{aligned} \quad (20)$$

Here the *finite* critical density  $n_c(\beta) := n(\beta, \mu = 0)$  is defined similarly to (12), where the particle density is

$$n(\beta, \mu) := \lim_{L_1, L \rightarrow \infty} \sum_{s \neq (s_1, 0, 0)} \kappa^3 \omega_1 \omega_2 \omega_3 N_s(\beta, \mu) = \int_{\mathbb{R}^3_+} \frac{\kappa^3 d\omega_1 d\omega_2 d\omega_3}{e^{\beta[(\omega_1 + \omega_2 + \omega_3) - \mu]} - 1}. \quad (21)$$

Equation (20) implies for  $\Delta_L(\beta, n)$  the same asymptotics as in (13):

$$\Delta_L(\beta, n) = \frac{1}{\beta} e^{-\beta(n - n_c(\beta))m^2 L^4 / (\hbar \kappa^3)} + \dots \quad (22)$$

If we choose  $L_1 := L e^{\hat{\gamma} L^4}$ , for  $\hat{\gamma} > 0$ , then the *second* critical density  $n_m(\beta) := n_c(\beta) + (\hat{\gamma} \hbar \kappa^3) / (\beta m^2)$ . For  $n_c(\beta) < n < n_m(\beta)$  we obtain the *type III* vdBLP-GC, i.e., *none* of the single-particle states are *macroscopically* occupied:

$$n_s(\beta, \rho) := \lim_{L \rightarrow \infty} \frac{\kappa^3 \omega_1 \omega_2 \omega_3}{e^{\beta(\eta_s - \mu_L(\beta, n))} - 1} = 0. \quad (23)$$

Although for  $n_m(\beta) < n$  there is a coexistence of the *type III* vdBLP-GC, with the constant density  $n_m(\beta) - n_c(\beta)$ , and the standard *type I* vdBLP-GC in the ground-state:

$$n - n_m(\beta) = \lim_{L \rightarrow \infty} \frac{\kappa^3 \omega_1 \omega_2 \omega_3}{e^{\beta(\eta_{(0,0,0)} - \mu_L(\beta, n))} - 1}. \quad (24)$$

4. In experiments with BEC, it is important to know the critical temperatures associated with corresponding critical densities. The *first* critical temperatures:  $T_c(\rho)$ ,  $\tilde{T}_c(\rho)$  or  $\hat{T}_c(\rho)$  are well-known. For a given density  $\rho$  they verify the identities:

$$\rho = \rho_c(\beta_c(\rho)), \quad \varrho = \varrho_c(\tilde{\beta}_c(\varrho)), \quad n = n_c(\hat{\beta}_c(n)), \quad (25)$$

respectively for our models of slabs, squared beams or "cigars". Since definition of the critical densities yield the representations:  $\rho_c(\beta) =: T^{3/2} I_{sl}$ ,  $\varrho_c(\beta) =: T^2 I_{bl}$ ,  $n_c(\beta) =: T^3 I_{cg}$ , the expressions for the *second* critical densities one gets the following relations between the *first* and the *second* critical temperatures:

$$\begin{aligned} T_m^{3/2}(\rho) + \tau^{1/2} T_m(\rho) &= T_c^{3/2}(\rho) \quad (\text{slab}), \\ \tilde{T}_m^2(\varrho) + \tilde{\tau} \tilde{T}_m(\varrho) &= \tilde{T}_c^2(\varrho) \quad (\text{beam}), \\ \hat{T}_m^3(n) + \hat{\tau}^2 \hat{T}_m(n) &= \hat{T}_c^3(n) \quad (\text{cigar}). \end{aligned}$$

Here  $\tau = [\alpha m k_B / (\pi \hbar^2 I_{sl})]^2$ ,  $\tilde{\tau} = 2\gamma k_B / (\hbar I_{bl})$  and  $\hat{\tau} = [(\hat{\gamma} \hbar \kappa^3 k_B) / (m^2 I_{cg})]^{1/2}$  are "effective" temperatures related to the corresponding geometrical shapes. Notice that the *second* critical temperature modifies the usual law for the condensate fractions temperature dependence, since now the total condensate density is  $\rho - \rho_c(\beta) := \rho_0(\beta) = \rho_{0c}(\beta) + \rho_{0m}(\beta)$ . Here  $\rho_{0m}(\beta) := (\rho - \rho_m(\beta)) \theta(\rho - \rho_m(\beta))$ .




FIG. 2: The first (blue fit), The second (pink fit) and the total (green fit) condensate fractions as a function of the temperature for  $^{87}\text{Rb}$  atoms in the slab geometry with  $T_{c1} = 10^{-7} \text{ K}$  and  $\tau = 4.43 \times 10^{-7}$ .

For example, in the case of the slab geometry the *type III* vdBLP-GC (i.e. the "quasi-condensate")  $\rho_{0c}(\beta)$  behaves for a given  $\rho$  like (see Fig.2.)

$$\frac{\rho_{0c}(\beta)}{\rho} = \begin{cases} 1 - (T/T_c)^{3/2}, & T_m \leq T \leq T_c, \\ \sqrt{\tau} T/T_c^{3/2}, & T \leq T_m. \end{cases} \quad (26)$$

Similarly, for the *type I* vdBLP-GC in the ground state  $\rho_{0m}(\beta)$  (i.e. the conventional BEC) we obtain:

$$\frac{\rho_{0m}(\beta)}{\rho} = \begin{cases} 0, & T_m \leq T \leq T_c, \\ 1 - (T/T_c)^{3/2} (1 + \sqrt{\tau/T}), & T \leq T_m, \end{cases} \quad (27)$$

see Fig.2. The total condensate density  $\rho_0(\beta) := \rho_{0c}(\beta) + \rho_{0m}(\beta)$  is the result of *coexistence* of both of them: it gives the standard PBG expression  $\rho_0(\beta)/\rho = 1 - (T/T_c)^{3/2}$ .

For the "cigars" geometry case the temperature dependence of the "quasi-condensate"  $\rho_{0c}(\beta)$  is

$$\frac{n_{0c}(\beta)}{n} = \begin{cases} 1 - (T/\hat{T}_c)^3, & \hat{T}_m \leq T \leq \hat{T}_c, \\ \hat{\tau}^2 T/\hat{T}_c^3, & T \leq \hat{T}_m. \end{cases} \quad (28)$$

The corresponding ground state conventional BEC behaves as

$$\frac{n_{0m}(\beta)}{n} = \begin{cases} 0, & \hat{T}_m \leq T \leq \hat{T}_c, \\ 1 - (T/\hat{T}_c)^3 (1 + \hat{\tau}^2/T^2), & T \leq \hat{T}_m, \end{cases} \quad (29)$$

and again for the two coexisting condensates one gets  $n - n_c(\beta) := n_0(\beta) = n_{0c}(\beta) + n_{0m}(\beta) = (1 - (T/T_c)^{3/2})n$ .

Notice that for a given density the difference between two critical temperatures for the slab geometry can be calculated explicitly:

$$(T_c - T_m)/T_c = g(\rho_\alpha/\rho), \quad (30)$$

where  $\rho_\alpha := 8\alpha^3/\zeta(3/2)^2$  and  $g(x)$  is an explicit algebraic function. For illustration consider a *quasi-2D* PBG model of  $^{87}\text{Rb}$  atoms in trap with characteristic sizes  $L_1 = L_2 = 100 \mu\text{m}$ ,  $L = 1 \mu\text{m}$  and with typical



critical temperature  $T_c = 10^{-7}K$ . The *anisotropy* parameter is  $\alpha = (1/L)\ln(L_1/L) = 4,6 \cdot 10^6 m^{-1}$ . Then for  $\tau = 4,4 \cdot 10^{-7}K$  we find  $T_m = 3,7 \cdot 10^{-8}K$  and  $(T_c - T_m)/T_c = 0,63$ .

5. Another physical observable to characterise this second critical temperature is the condensate coherence length or the global spacial particle density distribution.

The usual criterion is the ODLRO, which is going back to Penrose and Onsager [19]. For a fixed particle density  $\rho$  it is defined by the kernel:

$$K(x, y) := \lim_{L \rightarrow \infty} K_\Lambda(x, y) = \lim_{L \rightarrow \infty} \sum_s \frac{\bar{\phi}_{s,\Lambda}(x) \phi_{s,\Lambda}(y)}{e^{\beta(\varepsilon_s - \mu_L(\beta, \rho))} - 1}. \quad (31)$$

The limiting diagonal function  $\rho(x) := K(x, x)$  is *local*  $x$ -independent particle density.

To detect a trace of the geometry (or the second critical temperature) impact on the spatial density distribution we follow a recent scaling approach to the generalised BEC developed in [18] (see also [11],[15]) and introduce a *scaled global* particle density:

$$\xi_L(u) := \sum_s \frac{|\phi_{s,\Lambda}(L_1 u_1, L_2 u_2, L_3 u_3)|^2}{e^{\beta(\varepsilon_s - \mu)} - 1}, \quad (32)$$

with the scaled distances  $\{u_j = x_j/L_j \in [0, 1]\}_{j=1,2,3}$ .

For a given  $\rho$  the scaled density (32) in the slab geometry is

$$\xi_{\rho,L}^{sl}(u) := \sum_s \frac{1}{e^{\beta(\varepsilon_s - \mu_L(\beta, \rho))} - 1} \prod_{j=1}^{d=3} \frac{2}{L_j} [\sin(\pi s_j u_j)]^2. \quad (33)$$

Since  $2[\sin(\pi s_j u_j)]^2 = 1 - \cos\{(2\pi s_j/L_j)u_j L_j\}$  and  $\lim_{L \rightarrow \infty} \mu_L(\beta, \rho < \rho_c(\beta)) < 0$ , by the Riemann-Lebesgue lemma we obtain that  $\lim_{L \rightarrow \infty} \xi_{\rho,L}^{sl}(u) = \rho$  for any  $u \in (0, 1)^3$ . If  $\rho > \rho_c(\beta)$ , one has to proceed as in (3)-(5). Then for any  $u \in (0, 1)^3$ :

$$\begin{aligned} & \lim_{L \rightarrow \infty} \sum_{s=(s_1, s_2, 1)} \frac{1}{e^{\beta(\varepsilon_s - \mu_L(\beta, \rho))} - 1} \prod_{j=1}^{d=3} \frac{2}{L_j} [\sin(\pi s_j u_j)]^2 \\ &= \lim_{L \rightarrow \infty} \frac{2[\sin(\pi u_3)]^2}{(2\pi)^2 L} \int_{\mathbb{R}^2} \frac{\prod_{j=1}^2 (1 - \cos(2k_j u_j L_j) d^2 k)}{e^{\beta(\hbar^2 k^2/2m + \Delta_L(\beta, \rho))} - 1} \\ &= (\rho - \rho_c(\beta)) 2[\sin(\pi u_3)]^2, \end{aligned} \quad (34)$$

$$\begin{aligned} & \lim_{L \rightarrow \infty} \sum_{s \neq (s_1, s_2, 1)} \frac{1}{e^{\beta(\varepsilon_s - \mu_L(\beta, \rho))} - 1} \prod_{j=1}^{d=3} \frac{2}{L_j} [\sin(\pi s_j u_j)]^2 \\ &= \rho_c(\beta). \end{aligned} \quad (35)$$

Then the limit of (33) is equal to

$$\xi_\rho^{sl}(u) = (\rho - \rho_c(\beta)) 2[\sin(\pi u_3)]^2 + \rho_c(\beta). \quad (36)$$

It manifests a *space anisotropy* of the type III vdBLP-GC for  $\rho_c(\beta) < \rho < \rho_m(\beta)$  in direction  $u_3$ .

For  $\rho > \rho_m(\beta)$  one has to use representation (3) and asymptotics (6),(7). Then following the arguments developed above we obtain

$$\begin{aligned} \xi_\rho^{sl}(u) &= (\rho - \rho_m(\beta)) \prod_{j=1}^3 2[\sin(\pi u_j)]^2 + \\ &(\rho_m(\beta) - \rho_c(\beta)) 2[\sin(\pi u_3)]^2 + \rho_c(\beta). \end{aligned} \quad (37)$$

So, the anisotropy of the space particle distribution is still in direction  $u_3$  due to the type III vdBLP-GC.

It is instructive to compare this anisotropy with a *coherence length* analysis within the scaling approach [18] to the BEC space distribution. To this end let us center the box  $\Lambda$  at the origin of coordinates:  $x_j = \tilde{x}_j + L_j/2$  and  $y_j = \tilde{y}_j + L_j/2$ . Then the ODLRO kernel (31) is:

$$K_\Lambda(\tilde{x}, \tilde{y}) = \sum_{l=1}^{\infty} e^{l\beta\mu_L(\beta, \rho)} R_l^{(2)} R_l^{(1)}, \quad (38)$$

where after the shift of coordinates and using (1) we put

$$\begin{aligned} R_l^{(2)}(\tilde{x}^{(2)}, \tilde{y}^{(2)}) &= \\ \sum_{s=(s_1, s_2)} e^{-l\beta\varepsilon_{s_1, s_2}} \bar{\phi}_{s_1, s_2, \Lambda}(\tilde{x}_1, \tilde{x}_2) \phi_{s_1, s_2, \Lambda}(\tilde{y}_1, \tilde{y}_2) \end{aligned} \quad (39)$$

$$\begin{aligned} R_s^{(1)}(\tilde{x}_3, \tilde{y}_3) &= \sum_{s=(s_3)} e^{-l\beta\varepsilon_{s_3}} \sqrt{\frac{2}{L_3}} \sin\left(\frac{\pi s_3}{L_3}(\tilde{x}_3 + \frac{L_3}{2})\right) \\ &\times \sqrt{\frac{2}{L_3}} \sin\left(\frac{\pi s_3}{L_3}(\tilde{y}_3 + \frac{L_3}{2})\right). \end{aligned} \quad (40)$$

Similar to (3), for  $\rho_c(\beta) < \rho < \rho_m(\beta)$  we must split the sum over  $s = (s_1, s_2, s_3)$  in (38) into two parts. Since by the generalized Weyl theorem one gets:

$$\lim_{L \rightarrow \infty} R_l^{(2)}(\tilde{x}^{(2)}, \tilde{y}^{(2)}) = \frac{1}{l\lambda_\beta^2} e^{-\pi\|\tilde{x}^{(2)} - \tilde{y}^{(2)}\|^2/l\lambda_\beta^2},$$

by (38) we obtain for the first part the representation:

$$\begin{aligned} & \lim_{L \rightarrow \infty} \sum_{l=1}^{\infty} e^{l\beta\mu_L(\beta, \rho)} \sum_{s=(s_1, s_2, 1)} e^{-l\beta\varepsilon_{s_1, s_2, 1}} \times \\ & \times \bar{\phi}_{s_1, s_2, 1\Lambda}(\tilde{x}) \phi_{s_1, s_2, 1\Lambda}(\tilde{y}) = \\ & \lim_{L \rightarrow \infty} \sum_{l=1}^{\infty} e^{-l\beta\Delta_L(\beta, \rho)} \frac{1}{l\lambda_\beta^2} e^{-\pi\|\tilde{x}^{(2)} - \tilde{y}^{(2)}\|^2/l\lambda_\beta^2} \times \\ & \times \frac{2}{L} \sin\left(\frac{\pi}{L}(\tilde{x}_3 + \frac{L}{2})\right) \sin\left(\frac{\pi}{L}(\tilde{y}_3 + \frac{L}{2})\right). \end{aligned} \quad (41)$$

For the second part we apply the Weyl theorem for the 3-dimensional Green function:

$$\begin{aligned} & \lim_{L \rightarrow \infty} \sum_{l=1}^{\infty} e^{l\beta\mu_L(\beta, \rho)} \sum_{s \neq (s_1, s_2, 1)} e^{-l\beta\varepsilon_s} \times \\ & \times \bar{\phi}_{s, \Lambda}(\tilde{x}) \phi_{s, \Lambda}(\tilde{y}) = \sum_{l=1}^{\infty} \frac{1}{l\lambda_\beta^3} e^{-\pi\|\tilde{x} - \tilde{y}\|^2/l\lambda_\beta^2}. \end{aligned} \quad (42)$$




FIG. 3: Exponent  $\gamma(T)$  for evolution of the coherence length for the quasi-condensate with temperature corresponding to  $^{87}\text{Rb}$  atoms in the slab geometry with  $T_{c1} = 10^{-7}\text{K}$  and  $\tau = 4.43 \times 10^{-7}\text{K}$

If in (41) we change  $l \rightarrow l \Delta_L(\beta, \rho)$ , then it gets the form of the integral Darboux-Riemann sum, where  $\|\tilde{x}^{(2)} - \tilde{y}^{(2)}\|^2$  is scaled as  $\|\tilde{x}^{(2)} - \tilde{y}^{(2)}\|^2 \Delta_L(\beta, \rho)$ . Therefore, the coherence length  $L_{ch}$  in direction perpendicular to  $x_3$  is  $L_{ch}(\beta, \rho)/L := 1/\sqrt{\Delta_L(\beta, \rho)}$ . A similar argument is valid for  $\rho > \rho_m(\beta)$  with obvious modifications due to BEC for  $s = (1, 1, 1)$  (7) and to another asymptotics (6) for  $\Delta_L(\beta, \rho)$ . To compare the coherence length with the scale  $L_{1,2} = Le^{\alpha L}$ , let us define the critical exponent  $\gamma(T, \rho)$  such that  $\lim_{L \rightarrow \infty} (L_{ch}(\beta, \rho)/L)(L_1/L)^{-\gamma(T, \rho)} = 1$ . Then we get:

$$\begin{aligned} \gamma(T, \rho) &= \lambda_\beta^2 (\rho - \rho_c(\beta))/2\alpha, \quad \rho_c(\beta) < \rho < \rho_m(\beta) \\ &= \lambda_\beta^2 (\rho_m(\beta) - \rho_c(\beta))/2\alpha, \quad \rho_m(\beta) \leq \rho. \end{aligned} \quad (43)$$

For a fixed density, taking into account (26) we find the temperature dependence of the exponent  $\gamma(T) := \gamma(T, \rho)$ , see Fig.3:

$$\begin{aligned} \gamma(T) &= \sqrt{T/\tau} \{(T_c/T)^{3/2} - 1\}, \quad T_m < T < T_c, \\ &= 1, \quad T \leq T_m. \end{aligned} \quad (44)$$

Notice that in the both cases the ODLRO kernel is anisotropic due to impact of the type III condensation (41) in the states  $s = (s_1, s_2, 1)$ , whereas the other states give a symmetric part of correlations (42), which includes a constant term  $\rho_c(\beta)$ .

Numerically, for  $L_1 = L_2 = 100\mu\text{m}$ ,  $L_3 = 1\mu\text{m}$  and  $T_m < T = 0.75T_c$  the coherence length of the condensate is equal to  $2.8\mu\text{m} \ll 100\mu\text{m}$ . This decreasing of the coherence length is experimentally observed in [7]. 6. In conclusion we add several remarks about a possible impact of particle interaction. Since the "quasi-condensate" is observed in extremely anisotropic traps [7]-[9], we think that the geometry of the vessels is predominant. So, the study of the PBG is able to catch the phenomenon and so is relevant. Next, in this letter did not enter into details of the phase-fluctuations [8], [7], although we suppose that for the vdBLP-GC it can be

studied by switching different Bogoliubov quasi-average sources in condensed modes. Finally, since a repulsive interaction is able to transform the conventional one-mode BEC (type I) into the vdBLP-GC of type III, [20], [21], it is important to combine this interaction with the results already obtained for interacting gases in [7]-[9].

The pioneer calculations of a crossover in a trapped 1D PBG are due to [22]. It is similar to the vdBLP-GC in our exact calculations for the "cigars" geometry and it apparently persists for a *weakly* interacting Bose-gas as argued in [9]. Although the ultimate aim is to understand the relevance of these quasi-1D calculations for the Lieb-Liniger exact analysis of a strongly interacting gas [23].

We are going to return to these questions in our next papers.

- 
- [1] N.N.Bogoliubov, Phys.Abhandl.Sowijetunion **6**, 1-110 (1962); *ibid* **6**, 113-229 (1962).
  - [2] P.C.Hohenberg, Phys.Rev. **158**, 383-386 (1967).
  - [3] M.Bouziane and Ph.A.Martin, J.Math.Phys. **17**, 1848-1851 (1976).
  - [4] R.M.Ziff, G.E.Uhlenbeck and M.Kac, Phys.Reports **32**, 169-248 (1977).
  - [5] N.N.Bogoliubov, Selected Works, vol.II : Quantum Statistical Mechanics (Gordon and Breach, N.Y.,1991).
  - [6] N.N.Bogoliubov, Collection of Scientific Works in 12 vols.: Statistical Mechanics, vol.6, Part II (Nauka, Moscow, 2006).
  - [7] F. Gerbier, J. H. Thywissen, S. Richard, M. Hugbart, P. Bouyer, and A. Aspect, Phys. Rev. A **67** 051602(R) (2003)
  - [8] D. S. Petrov, G. V. Shlyapnikov, et J. T. M. Walraven, Phys. Rev.Lett. **87**, 050404 (2001).
  - [9] D. S. Petrov, G. V. Shlyapnikov, et J. T. M. Walraven, Phys. Rev.Lett. **85**, 3745 (2000).
  - [10] Z.Hadzibabic, P. Krüger, M. Cheneau, B. Battelier, and J. Dalibard, Nature **441**, 1118 (2006).
  - [11] M.van den Berg, J.T.Lewis and J.Pulé, Helv.Phys.Acta **59**, 1273-1288 (1986).
  - [12] H.B.G.Casimir, *On Bose-Einstein condensation*, in Fond.Probl.in Stat.Mech.v.III, 188-196, ed.E.G.D.Cohen (North-Holland Publ.Comp., Amsterdam, 1968).
  - [13] M.Girardeau, J.Math.Phys. **1**, 516-523 (1960)
  - [14] M. van den Berg, J.Stat. Phys. **31**, 623-637 (1983).
  - [15] M.van den Berg, J.T.Lewis and M.Lunn, Helv.Phys.Acta **59**, 1289-1310 (1986).
  - [16] J.Gough and J.Pulé, Helv.Phys.Acta **66**, 17 (1993).
  - [17] A.Patrick, J.Stat.Phys. **75**, 253 (1994).
  - [18] M.Beau, J.Phys.A **42**, 235204 (2009).
  - [19] O.Penrose and L.Onsager, Phys.Rev. **104**, 576 (1956).
  - [20] T.Michoel and A.Verbeure, J.Math.Phys.**40**, 1268 (1999).
  - [21] J.-B.Bru and V.A.Zagrebnov, Physica A**268**, 309 (1999).
  - [22] W.Ketterle and N.J.van Druten, Phys.Rev.A**54**, 656 (1996).
  - [23] E.H.Lieb and W.Liniger, Phys.Rev.**130**, 1605 (1963); E.H.Lieb, Phys.Rev.**130**, 1616 (1963).